Main Manuscript for

Electrical control of the transduction channels' gating force

in mechanosensory hair cells.

SHORT TITLE: Control of the channel-gating force in hair cells

Achille Joliot[1,2], Laure Stickel[1,2] and Pascal Martin[1,2,*]

[1] Physique des Cellules et Cancer, Institut Curie, PSL Research University, CNRS
UMR168, F-75248 Paris, France.
[2] Sorbonne Université, F-75252 Paris, France.
10
[*] Correspondence: pascal.martin@curie.fr

Keywords: auditory system, bioelectricity, gating compliance, mechanosensitive ion channels,
vestibular system.

The main manuscript includes:
Main Text
Figures 1 to 6
19
20

ABSTRACT

The inner ear's hair cells rely on mechanosensitive ion channels to convert vibrations of their hair bundles into electrical signals. The mechanical correlates of channel gating—the gating force and the gating swing—are fundamental determinants of hair-cell mechanosensitivity but are still poorly understood. Here we show that varying the electrical potential across the sensory hair-cell epithelium continuously modulates the gating force, by up to $\pm 100\%$. Our observations also revealed an abrupt transition between states of weak and strong gating force at a threshold potential, so that strong gating forces associated to high mechanosensitivity are observed only when the calcium influx through the channels is large enough, but not too large. Gating-force changes, remarkably enough, were explained by the modularity of the gating swing, ranging from values comparable to the channel pore size to nearly tenfold larger. Gating-swing control is expected to underly the hair cell's ability to tune its mechanosensitivity to minute sound stimuli.

TEASER: Mechanosensitivity of ion-channel gating in hair cells can be tuned by the electrical potential across the sensory epithelium.

INTRODUCTION

Mechanoreception by the inner ear's sensory hair cells begins with the deflection of their hair bundle—a cohesive tuft of cylindrical stereocilia arranged in a staircase pattern (1–3). A deflection toward the taller row of stereocilia increases mechanical tension in oblique tip links, which connect the tip of each stereocilium to the flank of a taller neighbor. The increased tip-link tension opens mechanosensitive ion channels and evokes an influx of cations into the hair cell, primarily K^+ ions but also Ca^{2+} ions.

Our understanding of hair-cell mechanosensitivity is informed by the gating-spring model of mechanoelectrical transduction (4, 5). Within this framework, the transduction channels are gated directly by force, which is transmitted to the channels via elastic elements—the gating springs—that operate in series with the tip links. The product of the gating-spring stiffness and the gating swing defines the gating force, a fundamental determinant of hair-cell mechanosensitivity: its magnitude sets the maximal slope of the sigmoidal relation between the transduction current and the deflection of the hair bundle (fig. S1, A and B). The greater the gating force, the larger the transduction current evoked by a small stimulus, making the hair cell more sensitive to a given deflection of its hair bundle.

The gating-spring model imposes a reciprocal relationship between tip-link tension and channel gating. If tensioning the tip links evokes channel opening, then channel opening must reduce tip-link tension by an amount given precisely by the gating force. Feedback between tip-link tension and channel gating produces gating compliance, the effective softening of the hair bundle upon channel gating (6, 7). For large positive or negative bundle displacements, at which the transduction channels are nearly all open or closed, respectively, the force-displacement relation is linear. At intermediate displacements that elicit channel gating, however, the slope of the force-displacement relation is reduced: the hair bundle is effectively softer. Gating compliance is associated with a shift between the two linear limbs of the force-displacement relation; this shift provides an estimate of the combined gating force, denoted F_G in the following, produced by the ensemble of transduction channels that operate in parallel within the hair bundle (fig. S1B). Furthermore, upon an oscillatory movement of the hair bundle, opening and closing of the transduction channels also increase friction on the hair bundle, a phenomenon called gating friction (8). The experimental demonstration of gating compliance and gating friction at the cellular scale of a micron-sized hair bundle provides strong support for the gating-spring model. Because these phenomena reflect the mechanical correlate of

channel gating, they can also be used to infer fundamental properties of channel gating at the
molecular scale such as the gating force (6, 7, 9).

Biophysical estimates of the gating force are variable, even with hair cells from the same
animal species and organ. With hair cells in excised preparations of the mammalian cochlea,
there is some evidence of gating compliance (10–12) but gating forces can also be so weak that
the force-displacement relation remains linear (13, 14). In these experiments, however, the
endocochlear potential that normally boosts transduction currents in vivo is absent and the hair
cells are fully immersed in a standard extracellular fluid called perilymph. This condition
disrupts the two ionic compartments that hair cells experience in vivo, with endolymph and
perilymph bathing the cells' apical and basal aspects, respectively. In the frog saccule, gating
forces can lead to a moderate (about 40%) decrease in hair-bundle stiffness (6) or be strong
enough to result in negative stiffness, which can lead to spontaneous oscillations of the hair
bundle and, in turn, frequency-selective amplification of periodic stimuli (9, 15–17).
Remarkably enough, strong gating forces and negative stiffness were reported only under ex-
vivo conditions that closely mimic natural ones, with reconstituted endolymph-perilymph
compartments.

The reason for the apparent variability of the gating force is unclear. Variability may result
from hair-cell damage inflicted by artificial ex-vivo conditions or instead reflect an intrinsic
ability of the hair cell to control the magnitude of its gating force under variable conditions.
These observations call for a comprehensive and systematic investigation of the gating force
under a range of physiological conditions.

In this work, we report that the electrical potential across the sensory epithelium that houses
the hair cells can serve as a control parameter of the gating force. Our results indicate that the
gating force varies primarily because the gating swing varies. Our work further reveals a
reversible, but hysteretic, transition between a state of strong gating force, for which the hair
bundles oscillate spontaneously, and a state of weak gating force, for which hair-bundle
mechanics is linear and the hair bundles are quiescent. We finally provide evidence that these
effects are mediated, at least in part, by electrical control of the influx of Ca^{2+} through the
transduction channels.

RESULTS

We took advantage of an excised preparation of the frog saccule that mimics *ex vivo* the two-
compartment ionic conditions that the sensory epithelium experiences *in vivo*. Under these
conditions, the hair cells routinely displayed spontaneous oscillations of their hair bundles, as
recognized before (18–20). We aimed here at evaluating the effects of a transepithelial potential
on hair-bundle mechanics. To this end, we used a pair of electrodes and a current generator to
impose a transepithelial current, I, within a range of $\pm 10\ \mu A$ (fig. S2A). Upon application of a
current step, the transepithelial potential, U, reached a steady-state value in about $600\ \mu s$
(fig. S2B). As a rule of thumb, a positive current of $+1\ \mu A$ evoked a positive potential in the
endolymphatic compartment of $+10\ mV$ with respect to the perilymphatic compartment;
accordingly, the transepithelial potential thus varied within a range of $\pm 100\ mV$. We emphasize
for the following that a positive endolymphatic potential is expected to increase the influx of
K^+ and Ca^{2+} ions through the transduction channels, whereas a negative potential does the
opposite. All numerical results are given below as the average and standard deviations (SD)
over an ensemble of n independent measurements.

The potential modulates strongly the gating force but weakly the hair-bundle stiffness

To determine the effects of an endolymphatic potential on the mechanical properties of a hair
bundle, we measured the bundle's force-displacement relation in response to a series of step
displacements applied to the base of a stimulus fiber (Fig. 1). Under control conditions, for
which no transepithelial current was applied ($I = 0$), the force-displacement relation showed a
prominent region of reduced slope, or gating compliance, which is expected for hair bundles
that oscillate spontaneously (9, 16). A fit of the relation to Equation 1 (Methods) yielded a set-
point deflection $X_0 = -2 \pm 7$ nm, a gating force $F_G = 17 \pm 4$ pN and a linear stiffness $K =$
$0.85 \pm 0.2\ pN \cdot nm^{-1}$ (mean $\pm$ SD; $n = 29$ in 16 frogs). We recall that the set-point deflection
can be interpreted as the deflection that leads to a channels' open probability of ½ (16). For
each hair bundle, the procedure was repeated in the presence of a static transepithelial current
($I \neq 0$).

We start by presenting the effects of a transepithelial current on a force-displacement relation
in four representative examples (Fig. 1); the effects on the corresponding spontaneous hair-
bundle oscillations are shown in Supplementary Materials (fig. S3, A to D). Applying a positive
current, here $I = +3\ \mu A$, corresponding to a positive endolymphatic potential $U \simeq +30\ mV$,

evoked an increase of the set-point deflection: the transduction channels closed (Fig. 1A). In

addition, we observed a small (-5%) decrease of the linear bundle's stiffness and, remarkably,

a large decrease (-38%) of the gating force. In another cell, a larger current $I = +6\ \mu A$ resulted

in a similar change in stiffness (-6%) but evoked a larger decrease of the gating force (-50%;

Fig. 1B). Conversely, applying a negative current of moderate magnitude, here $I = -3\ \mu A$,

corresponding to a negative endolymphatic potential $U \simeq -30$ mV, elicited a decrease of the

set-point deflection and thus opening of the transduction channels, a small increase ($+10\%$) of

the linear stiffness, and a large increase ($+89\%$) of the gating force (Fig. 1C). Furthermore,

applying a large transepithelial current, here $I = -6\ \mu A$, led to an unexpected finding: the force-

displacement relation became linear (Fig. 1D). This is not because the integrity of the hair cell

was impaired by application of a large transepithelial current, for all the effects were fully

reversible.

Performing an ensemble of $n = 41$ pairs of measurements in 29 hair bundles from 16 frogs

confirmed the effects illustrated by the four representative examples. We found a strong positive

correlation between the change, $\Delta X_0 = X_0(I) - X_0(I = 0)$, of the set-point deflection and the

applied transepithelial current (Fig. 2A). This correlation came as no surprise: the

endolymphatic potential resulting from the transepithelial current should influence the

magnitude of the Ca^{2+} influx through the transduction channels and calcium is known to

promote channel closure (4, 16, 21–23). The change in linear stiffness, $\Delta K = K(I) - K(I =$

$0)$, was negatively correlated to the transepithelial current, but only weakly (Fig. 2B). Instead,

for currents $I > -4\ \mu A$, the change, $\Delta F_G = F_G(I) - F_G(I = 0)$, of the hair-bundle's gating

force displayed a strong negative correlation with the current (Fig. 2C). The relative variation

of the gating force was correspondingly much larger than that of the linear stiffness (Fig. 2D).

A positive transepithelial current evoked a reduction of the gating force by as much as -80%,

whereas the stiffness decreased at most by -10% over the same range of transepithelial

currents. Conversely, the gating force could nearly double in the presence of a negative

transepithelial current, corresponding to a relative increase of $+100\%$, whereas stiffness did

not increase beyond $+25\%$. These correlations between mechanical properties of the hair

bundle and the transepithelial current were associated to systematic changes of the frequency,

amplitude and waveform of spontaneous hair-bundle oscillations (fig. S3, E to G).

For all the seven hair bundles for which the transepithelial current was more negative than

$-5\ \mu A$, corresponding to an endolymphatic potential $U \leq -50$ mV, the gating force did not

follow the trend observed at more positive potentials. Instead of increasing further or saturating

at more negative potentials, the gating force dropped to values so low that the force-

displacement relation was linear, thus with no detectable signature of gating compliance

(Figs. 1D and 2, C and D). At these endolymphatic potentials, no spontaneous oscillation was

observed; the hair bundles were quiescent (fig. S3D). Our study thus revealed the existence of

a critical value of the transepithelial current, $I = I_C \simeq -5\,\mu A$, and of the endolymphatic

potential, $U = U_C \simeq -50\,\text{mV}$, that delimited two functional states of the hair bundle: a state of

strong gating force for which gating compliance was large enough to yield spontaneous

oscillations of the hair bundle and a state of weak gating force for which hair-bundle mechanics

was linear and the hair bundle was quiescent.

The gating force varies because the gating swing varies

Within the framework of the gating-spring model (fig. S1A), the gating force for the whole hair

bundle, $F_G = K_{GS}D$, is the product of the combined stiffness of the gating springs, K_{GS}, and of

the gating swing, D, of a transduction channel. Remarkably enough, the relative effect of an

endolymphatic potential on the gating force was much larger than that on the hair-bundle

stiffness (Fig. 2D): the gating force could vary by as much as about $\pm 100\%$, whereas stiffness

in the same hair bundles varied only from -10% to $+25\%$. Because the gating springs are

expected to contribute a significant fraction of the total hair-bundle stiffness (24), K, we

reasoned that the change in gating force resulted mainly from a change in the magnitude of the

gating swing. We could confirm this inference by measuring the hair-bundle stiffness, $K_{SP} =$

$0.17 \pm 0.03\,\text{pN} \cdot \text{nm}^{-1}$ $(n = 14) \simeq 0.2\,K$, after disruption of the tip links. By subtracting the

average value of K_{SP} to individual measurements of K, we could then estimate, for each hair

bundle, the gating-spring stiffness $K_{GS} = K - K_{SP}$, and in turn the gating swing D. After

projection along the oblique axis of the tip links using the projection factor $\gamma = 0.14$ (25), we

found a gating swing $d = \gamma D = 3.7 \pm 1.4\,\text{nm}$ $(n = 29)$ under control conditions (Fig. 3A).

Replicating the behavior of the gating force, the gating-swing change displayed a strong

negative correlation with the transepithelial current for currents $I \geq I_C \simeq -5\,\mu A$ (Pearson

correlation $r = -0.86$; p-value $= 1 \cdot 10^{-10}$) and dropped to undetectable levels for $I \leq I_C$. For

$I \geq I_C$, the gating swing d could reach a value of up to about 8 nm in the presence of a negative

endolymphatic potential or, conversely, be as small as 2 nm upon application of a potential of

opposite polarity (Fig. 3B).

The transition between states of strong and weak gating force shows hysteresis

To further explore the transition between states of strong and weak gating force (Figs. 1D and 2C), we applied ramps of transepithelial current (Fig. 4A) while the hair bundle was subjected to triangular mechanical stimulation (Fig. 4B). This stimulation protocol allowed to monitor, continuously for the same hair bundle, how the mechanical properties varied as a function of the transepithelial current. The ramps were slow enough that the transepithelial current varied by only $\sim 0.2\ \mu A$ over the time required to measure a force-displacement cycle (see Methods). Because this variation was too small to elicit a significant variation of the gating force (Fig. 2D), the mechanical measurements were quasi static. The mechanical stimulus elicited a force-displacement cycle with clockwise circulation, corresponding to energy dissipation (8). The cycle displayed a region of lower slope—gating compliance—on the positive and negative half cycles (Fig. 4C), where the channels opened and closed, respectively. In addition, the height of the force-displacement cycle was larger within the region of channel gating. This property was previously shown to result from the contribution of the transduction channels' gating force to hair-bundle friction, a phenomenon called gating friction, and the displacement at which the cycle had a maximal height (abscissa of disks in Fig. 4C) was identified as the set-point deflection X_0 (8). In the following, the maximal half-height of the cycle, ϕ_{MAX}, was used as a reporter for the magnitude of the gating force.

As the transepithelial current decreased from zero to negative values, the region of gating compliance and gating friction in the force-displacement cycle shifted toward more negative displacements, corresponding to a decrease of the set-point deflection and thus opening of the transduction channels (Fig. 4C, movie S1). Following the point of inversion symmetry in a force-displacement cycle resulted in a nearly linear trajectory of slope $0.4 \pm 0.2\ \mathrm{pN \cdot nm^{-1}}$ ($n = 8$; Fig. 4D). In addition, the height of the force-displacement cycle, and thus the maximal friction force ϕ_{MAX} increased before saturating (Fig. 4E). When the absolute magnitude of the negative current became large enough, corresponding to $I \leq I_C = -3.7 \pm 1\ \mu A$ and an endolymphatic potential $U \leq U_C \simeq -37 \pm 10\ \mathrm{mV}$ ($n = 8$), the friction force ϕ_{MAX} dropped (Fig. 4, E and F, and movie S1) until the nonlinearity vanished (Fig. 4C, red). These observations on single hair cells (solid line in Fig. 4E) mirrored the measured changes of the gating force upon application of a static transepithelial current in an ensemble of cells (Fig. 2C). We note that gating friction generally depends on the activation time of the transduction channels, not just on the gating force (8). Detailed analysis of the force-displacement cycles

suggests that the channel-activation time had increased by about 50%, on average, when the
friction force ϕ_{MAX} reached a saturating value upon application of a negative transepithelial
current (fig. S4). The observed variation of hair-bundle friction (Fig. 4, E and F) thus resulted
from modulation of both the channel-activation time and the gating force (Fig. 2C).

The polarity of the current ramp was then reversed. As the negative current increased, the
set-point deflection increased and the point of inversion symmetry in the force-displacement
cycle showed a nearly linear trajectory of slope 0.6 ± 0.2 pN $\cdot$ nm^{-1} ($n = 8$; Fig. 4, C and D).
In addition, the signatures of gating compliance and gating friction, associated to nonlinear
force-displacement relations, were retrieved. The nonlinearity, however, reappeared at a
negative current with a smaller absolute value than that necessary to conceal the signatures of
channel-gating forces on the descending ramp. Over an ensemble of $n = 8$ cells, we measured
a shift in the transepithelial current of 2.3 ± 0.8 µA, corresponding to a 20-mV difference of
endolymphatic potential. Thus, the transition between states of strong and weak gating force
displayed hysteresis (Fig. 4E).

The transition kinetics between the two states occurs on a sub-second timescale
To characterize the kinetics of the transition between states of strong and weak gating force, we
applied large negative steps of transepithelial current, here $I = -8$ µA (Fig. 5), while
continuously measuring a hair bundle's force-displacement cycle in response to triangular
mechanical stimulation. As expected from the effects of current ramps (Fig. 4), the force-
displacement cycle shifted in the negative direction along the displacement axis and the force-
displacement cycle lost the signatures of gating compliance and gating friction (Fig. 5A,
movie S2). The time courses of the average hair-bundle position on the cycle, denoted $\bar{X}$ in the
following, and of the maximal friction force ϕ_{MAX} were quantified by measuring the
$10 - 90\%$ decay and rise times following the onset and offset of the current step, respectively.
Over an ensemble of $n = 4$ hair cells, the characteristic time, $\tau_X^{ON} = 500 \pm 50$ ms, of negative
hair-bundle motion evoked by the current step was fivefold longer than that, $\tau_X^{OFF} = 100 \pm$
30 ms, necessary to return to the initial position after the current was turned off (Fig. 5B). The
friction force ϕ_{MAX} showed an opposite behavior: its decay time, $\tau_\Phi^{ON} = 180 \pm 40$ ms, after
the onset of the current step was twofold shorter than its rise time, $\tau_\Phi^{OFF} = 400 \pm 100$ ms,
during recovery (Fig. 5B). Note that the time required to measure a force-displacement cycle

was here of only 20 ms. This duration was about ten times shorter than $\tau_\Phi^{OFF/ON}$ and thus short

enough to ensure a quasistatic assessment of hair-bundle mechanics (Methods).

There was thus a separation of timescales between the changes of $\bar{X}$ and ϕ_{MAX}. At the onset

of the current step, gating compliance vanished before the negative shift had completed:

$\tau_\Phi^{ON}/\tau_X^{ON} = 0.37 \pm 0.17$ (Fig. 5, B and C). In contrast, after the external transepithelial current

was turned off, the mechanical nonlinearity recovered more slowly than the average position of

the hair bundle: $\tau_\Phi^{OFF}/\tau_X^{OFF} = 4.2 \pm 2$ (Fig. 5, B and C).

Reducing the endolymphatic Ca²⁺ concentration replicates the effects of a negative

endolymphatic potential

A negative endolymphatic potential may have mediated its effects on the gating force by

reducing the magnitude of the Ca^{2+} influx through the transduction channels. To test this

hypothesis, we used iontophoresis of a Ca^{2+} chelator to reduce the endolymphatic Ca^{2+}

concentration near a hair bundle and, in turn, the Ca^{2+} influx through the channels. In response

to descending and ascending ramps of iontophoretic current (Fig. 6A), or equivalently of Ca^{2+}

concentration, we again observed an increase in hair-bundle friction followed by a reversible

but hysteretic transition between states of strong and weak gating force (Fig. 6, B and C, and

movie S3). Repeating the experiments over an ensemble of $n = 8$ cells confirmed the

robustness of these effects (Fig. 6D) and how they mirrored those observed upon modulation

of the endolymphatic potential (Fig. 4). In addition, the force-displacement relation of a hair

bundle could show strong gating compliance under control conditions but reversibly become

linear upon step application of a Ca^{2+} chelator, with no noticeable change in the linear stiffness

of the hair bundle (Fig. 6E). The same behavior was observed in $n = 7$ hair cells. Thus, enough

calcium in endolymph was required for the hair bundle to evince strong gating forces.

DISCUSSION

In this work, we showed that varying the endolymphatic potential modulates the gating force

continuously and reversibly, thus serving as a control parameter for the mechanosensitivity of

the hair cell to minute deflections of its hair bundle. This bioelectric effect is large; the gating

force could nearly double upon application of negative potentials and, conversely, could be

reduced by up to 80% of the control value at positive potentials (Fig. 2D). A major finding of

our study is that the gating force varied because the gating swing varied, with only minor effects

on the gating-spring stiffness (Figs. 2D and 3). In addition, our observations revealed an

unexpected transition between two functional states of the hair bundle—one with low
mechanosensitivity and one with high mechanosensitivity: a quiescent state of weak gating
force with gating swings of about one nanometer or less, and an oscillatory state of strong gating
force with gating swings up to ten times larger, respectively. The hair cell switched from a state
of strong gating force to a state of weak gating force when the endolymphatic potential became
more negative than a critical value $U = U_C \simeq -50$ mV (Figs. 1D, 2C, and 4, C and E). At such
negative potentials, the electromotive force driving the influx of cations through the
transduction channels is expected to be greatly reduced. Because reducing the Ca^{2+}
concentration near a hair bundle evoked a similar transition (Fig. 6), the strong gating forces
and large gating swings associated to high mechanosensitivity likely required a resting Ca^{2+}
influx through the transduction channels that was large enough. Altogether, our findings
demonstrate the modularity of the molecular determinants of hair-bundle mechanosensitivity—
specifically, the transduction channels' gating force and gating swing—in response to
variations in physiological control parameters.

Physiological role of the endolymphatic potential

Like other epithelia, the sensory epithelia that house the hair cells in the vertebrate inner ear are
polar tissues that experience a transepithelial potential under natural conditions (26). This
potential generally lies within a range $1 - 10$ mV, except in the mammalian cochlea where the
endolymphatic potential can reach $+100$ mV. Remarkably enough, this high value of the
endolymphatic potential is associated with a Ca^{2+} concentration in endolymph, about 20 μM,
which is about one of order of magnitude lower than in hearing organs from non-mammalian
species or than in vestibular organs. Our results on frog saccular hair cells (Figs. 2 and 6)
suggest that the combination of a large endolymphatic potential and a low Ca^{2+} concentration
in the cochlea might maintain a Ca^{2+} influx within the optimal range to allow for large gating
forces while boosting the magnitude of transduction currents. In support of this inference, a
mechanical nonlinearity attributed to hair-bundle mechanics has been measured in an ex-vivo
preparation of the Gerbil cochlea, but only under conditions that replicate a normal
endolymphatic potential and Ca^{2+} concentration (27). This nonlinearity disappears when no
endolymphatic potential is applied, which makes sense within the framework of our results: the
potential is then effectively very negative with respect to natural conditions—by 100 mV,
perhaps enforcing a similar state of weak gating force as we report here in frog upon application
of a potential that is negative enough (Fig. 1D, 2C and 4). Note that immersing the sensory

epithelium in artificial perilymph, as is routinely performed in the field, is also expected to lead

to weak gating forces but here as a result of a too high Ca^{2+} influx. In the cochlea, the Ca^{2+}

concentration in perilymph is indeed nearly two orders of magnitude higher than in endolymph.

Our results on frog saccular hair cells thus help putting ex-vivo measurements on mammalian

hair cells into prospective. In addition, our observations also imply that a spatial gradient in the

endolymphatic potential, as observed in vivo in the mammalian cochlea (28), might provide a

mechanism to tune the gating force and thus mechanosensitivity across hair cells.

In our experiments, control conditions approach natural conditions in the frog saccule: the

top and bottom compartments have ionic compositions that mimic those of endolymph and

perilymph, respectively, and there is no transepithelial potential. Also, our approach is based

on force measurements, which in contrast to standard electrophysiological approaches does not

disrupt the integrity of the hair cell or their internal ionic concentrations. Under these nearly

natural conditions, the gating force is strong, as a matter of fact strong enough to evoke negative

stiffness and promote spontaneous oscillations of the hair bundle (9). We applied

endolymphatic potentials within a range of ± 100 mV. While this range departs from natural

conditions, pushing the hair cells into uncharted territory disclosed gating-force control by the

endolymphatic potential and also indicated that the hair cell operates near an optimum of

mechanosensitivity under natural conditions.

It is worth noting that the potential difference between endolymph and the hair-cell interior

likely constitutes the relevant control parameter, not the endolymph potential per se. From this

perspective, modifying the endolymphatic potential may serve as a proxy for evaluating the

effects of varying the hair cell's transmembrane potential—at least insofar as

mechanosensitivity of the transduction channels is concerned. The efferent nervous system is

known to provide descending feedback from the brain to tune sensitivity of the auditory system

(29). Activation of the efferent system leads to hyperpolarization of the hair cell (30–32).

Assuming that hyperpolarization of the hair cell has similar effects on transduction channels'

operation as raising the endolymph potential to more positive values, our work opens the

possibility that efferent activation might affect the transduction channels' gating force and

gating swing.

Effects of Ca^{2+} on hair-bundle mechanics

Varying the endolymphatic potential afforded a means to reversibly and robustly control the

electric field in the membrane near the transduction channels and in turn the electromotive force

that controls the magnitude of transduction currents, including the Ca^{2+} influx into the hair cell.

Reducing the Ca^{2+} concentration near a hair bundle resulted in similar effects on gating

compliance and gating friction than applying negative endolymphatic potentials (Fig. 6). This

finding indicates that negative endolymphatic potentials modulated the gating force, at least in

part, by reducing the Ca^{2+} influx through the transduction channels. We cannot exclude,

however, that the endolymphatic potential also directly affected channel gating by mediating

an electric force on the transduction-channel protein complex. Depolarization of the hair cell

has been previously shown to evoke fast movements of the hair bundle, even with channels

blocked and thus with no change in the Ca^{2+} influx though the channels, which may be

indicative of a rearrangement of the channel protein complex (33, 34).

Calcium is also known to affect the channel activation time, the operating point of the

transducer and the kinetics of the active adaptation process that drives active hair-bundle

motility (4, 16, 18, 21–23, 35, 36). Moreover, there is some evidence that the extracellular

calcium concentration can modulate the stiffness of the hair bundle (23, 24, 37). In our

experiments, negative endolymphatic potentials produced effects on spontaneous hair-bundle

oscillations (fig. S3) that were qualitatively similar as those reported earlier in response to

depolarization of the hair cell (33) or to a decrease of the Ca^{2+} concentration in endolymph (18).

Conversely, positive endolymphatic potentials produced similar effects as hyperpolarization of

the hair cell or increasing the endolymphatic Ca^{2+} concentration.

Previous models had postulated stiffer gating springs at lower Ca^{2+} (18, 38), in apparent

contradiction with recent in-vitro measurements of PCDH15 elasticity (39). Our observations

(Figs. 2, 3 and 6) suggest instead that the effect of Ca^{2+} on gating-spring stiffness is small

compared to that on the gating swing, at least under our experimental conditions.

Implications for the mechanism of channel gating

As recognized earlier (9), we observed that the gating swing (Fig. 3) can be nearly one order of

magnitude larger than the expected diameter of the transduction channels' pore, which has been

estimated at about 1.2 nm (2). Hair cells thus appear to have implemented a mechanism to

magnify the conformational change associated to opening and closing of the transduction

channel's pore, a feature also relevant to other mechanosensitive channels (40). In hair cells,

the ion channels that mediate mechanoelectrical transduction are part of a molecular complex

that includes TMC1—the pore-forming subunit (41), TMIE, CIB2, LHFPL5 and the lower end

of the tip link, PCDH15 (42, 43). In addition, the complex may comprise two copies of each

protein, forming a dimeric assembly with an axis of twofold symmetry (44, 45) and comprising

a total of at least 10 proteins. Interactions between the multiple components of the transduction

apparatus might lead to a collective rearrangement upon opening of the channel's pore,

effectively producing a large gating swing. Alternatively, channel gating has been proposed to

elicit a membrane-mediated interaction between two channel complexes, each attached to one

strand of the PCDH15 dimer, leading to large movement within the membrane that magnifies

the apparent gating swing (46).

Whatever the molecular substrate for the gating swing, our work indicates that the

transepithelial potential and the extracellular calcium concentration can modulate the

magnitude of the movement associated to channel gating (Fig. 3). In addition, under conditions

for which the calcium influx through the channels becomes too low, we found that the gating

swing drops to undetectable levels. This observation suggests that the channel-protein complex

must be primed by enough intracellular calcium to elicit a large gating swing, perhaps via Ca^{2+}-

mediated conformational change in CIB2—the calcium- and integrin-binding protein (47, 48)

or by affecting the properties of the lipid membrane that houses the transduction channels (49).

Hysteresis upon a continuous back-and-forth variation of the endolymphatic potential (Fig. 4)

or of the extracellular concentration (Fig. 6), and the relatively short timescale of the transition

between states of strong and weak gating force (Fig. 5) provide further support for the bistability

of the channel-protein complex.

To cope with the physical requirements of hearing, hair cells have evolved the ability to

respond to sound stimuli that vary over timescales as short as a few tens of microseconds, detect

deflections of their hair bundle as small as one nanometer or less, and mediate sharp frequency

selectivity over three orders of magnitude of sound frequencies (50). These unique capabilities

rely on a transduction-channel protein complex with a highly intricate molecular composition

and architecture (43, 45). Inasmuch that the molecular underpinnings of mechanoelectrical

transduction by hair cells remains similar across species and organs, our force measurements

on hair bundles from frog saccular hair cells demonstrate that the effective conformational

change associated to channel gating—the gating swing—can be modulated over a broad range

by varying physiological control parameters. The modularity of the gating swing provides a

mechanism to tune passive and active mechanosensitivity of the hair cell while maintaining the

molecular composition of the transduction channels. Elucidating the molecular rearrangement

associated to channel gating and how it can be modulated remain major challenges for future

studies.

METHODS

Experimental preparation

All experimental procedures were approved by the Ethics committee on animal experimentation of the Institut Curie; they complied with the European and French National Regulation for the Protection of Vertebrate Animals used for Experimental and other Scientific Purposes (Directive 2010/63; French Decree 2013–118). Details of the experimental preparation have been published elsewhere (18, 23). Briefly, an excised preparation of the frog's saccule (Rivan 92 (51)) was glued with tissue-compatible surgical glue (Histoacryl, B. Braun) over a ~1 mm hole in a plastic disk (fig. S2A). The preparation was then secured in a two-compartment chamber, which exposed the apical and basal aspects of the sensory tissue to fluids with different ionic compositions to mimic physiological conditions. The basal bodies of the hair cells were bathed in artificial perilymph, a standard saline solution containing (in mM): 110 NaCl, 2 KCl, 1.5 CaCl$_2$, 3 D-glucose and 5 Hepes. Hair bundles instead projected into artificial endolymph with (in mM): 2 NaCl, 118 KCl, 0.25 CaCl$_2$, 3 D-glucose, and 5 Hepes. Each solution had a pH of ~ 7.3 and an osmotic strength of ~ 225 mmol $\cdot$ kg^{-1} . To disconnect the hair bundles from the overlying otolithic membrane, the apical surface of the preparation was exposed for 20 min to endolymph supplemented with 67 μg $\cdot$ mL^{-1} of the protease subtilisin (type XXIV, Sigma). The otolithic membrane was then peeled off to obtain access to individual hair bundles. Experiments were performed at a temperature of 20°C.

Microscopic apparatus and mechanical stimulation

The preparation was viewed through a $\times 60$ water-immersion objective of an upright microscope (BX51WI, Olympus). The mechanical properties of single hair bundles were characterized using published methods (8, 23). In short, the tip of a stimulus fiber was affixed to the kinociliary bulb of an individual hair bundle and imaged at a magnification of $\times 944$ onto a displacement monitor comprising a dual photodiode. Stimulus fibers were fabricated from borosilicate capillaries (TW120-3, World Precision Instrument) and coated with a thin layer of gold-palladium to enhance contrast (Hummer 6.1, Anatech ltd.). The fiber was secured by its base to a stack-type piezoelectric actuator (PA-8/14, Piezosystem Jena) driven by a custom-made power supply (Elbatech). Its stiffness, k_F, was estimated from the analysis of thermal fluctuations at the fiber's tip.

In a first stimulation protocol, a series of step displacements, Δ, was applied to the fiber's

base. The standard protocol consisted in the application of 7 to 9 successive pairs of step

displacements in opposite directions with a magnitude growing with a constant increment,

within the range $20 - 40$ nm. The force, $F = k_F (\Delta - X)$, and the bundle's displacement, X,

were estimated $5 - 9$ ms after the stimulus onset. This time was long enough for the open

probability of the transduction channels to have reached a new steady state and for the frictional

contribution to the applied force to have vanished, but short enough to limit the effect of

adaptation. The resulting force-displacement relation, $F(X)$, was fitted by the equation (5, 16):

$$F = KX - F_G/[1 + \exp(-(X - X_0)/\delta)] + F_\emptyset, \tag{1}$$

from which we estimated the bundle's linear stiffness, K, the gating force, F_G, for the whole

hair bundle, and the set-point deflection, X_0. The parameter $F_\emptyset$ ensured that $F = 0$ at $X = 0$.

Because we did not use displacement-clamp feedback to stabilize positions of negative stiffness

(9), the characteristic displacement, δ, could not be exploited. In this protocol, both photodiode

and command signals were acquired and generated at a sampling rate of 2.5 kHz and the half-

power frequency of the lowpass filter was fixed at 1 kHz. A force-displacement relationship

was obtained by averaging 7 to 15 repetitions of the same stimulation protocol, which

collectively took 10 to 20 seconds.

In a second stimulation protocol, we applied a symmetric triangular waveform of motion to

the fiber's base, at a frequency of either 40 Hz or 100 Hz and with a peak-to-peak amplitude

within the range $600 - 1000$ nm. The force applied to the hair bundle was estimated using a

detailed description of the fiber's vibrational modes (8). This protocol resulted in force-

displacement cycles with counterclockwise circulation. The half-height at any position, X, in

the cycle provides an estimate of the average friction force $\phi(X) = [F^+(X) - F^-(X)]/2$

impeding the hair bundle motion at this position, where $F^\pm(X)$ represents the force in the

positive and negative half cycle of stimulation, respectively. The friction force displayed a

maximum, $\phi_{\text{MAX}} = \phi(X_0)$, at a displacement that was identified earlier as the set-point

deflection X_0 in Equation 1 (8). The force-displacement cycle also displayed a point of

inversion symmetry of coordinates $(X_0, F_0 = [F^+(X_0) + F^-(X_0)]/2)$. The shift, ΔX, between

the positions where the hair-bundle speed reached a maximal value, $|dX/dt| = V_{MAX}$, in the

positive and negative half cycle of stimulation provided an estimate of the activation time of

the transduction channels, $\tau = \Delta X/(2V_{MAX})$ (8). In this protocol, the force-displacement cycle

was averaged over ten periods of the triangular stimulus at 40 Hz or, when a fast assessment of

hair-bundle mechanics became necessary (as for the data shown in Fig. 5), over only two

periods of a 100-Hz stimulus. Both photodiode and command signals were acquired and

generated at sampling rate of 25 kHz. The half-power frequency of the filter was fixed at

12.5 kHz for the photodiode signals and 0.5 kHz for the command signals.

Control of the transepithelial current and potential

Using a current source (Linear Stimulator Isolator A395, World Precision Instruments)

connected to a pair of silver chloride electrodes (fig. S2A), we applied steps of transepithelial

current with a magnitude that was varied within a range of ± 10 μA. In some preparations, we

measured the transepithelial potential, $U(t)$, evoked by a step of transepithelial current, $I_{STEP} =$

1 μA (fig. S2B), using an additional pair of electrodes connected to the headstage of a

differential amplifier (voltage gain $\times$ 50; Model 3000, Systems Inc). The time course of the

transepithelial potential was well described by an exponential relation $U(t) = U_{MAX} \exp(-t/$

$\tau)$, with a characteristic timescale $\tau = 230 \pm 50$ μs (mean $\pm$ SD, $n =$13). Over an ensemble

of 13 saccules, we found a transepithelial resistance $R = U_{MAX}/I_{STEP} = 10 \pm 2$ kΩ and

capacitance $C = 23 \pm 4$ nF. Thus, over the range of transepithelial currents that we explored,

the potential into the artificial endolymph that bathed the hair bundles—the endolymphatic

potential— varied within a range of ± 100 mV with respect to the potential in the perilymphatic

compartment.

In other experiments, we applied successive ramps of descending and ascending

transepithelial currents at an absolute rate within the range of $0.5 - 0.9$ μA $\cdot$ s^{-1}. These rates

were low enough to ensure that the variation of the transepithelial current remained small during

the time required for probing hair-bundle mechanics, allowing these mechanical measurements

to remain quasistatic. The minimal current was maintained for $1 - 2.5$ s before the ascending

ramp started.

Iontophoresis of a calcium chelator

We used iontophoresis of a calcium chelator to decrease the extracellular calcium concentration

in the vicinity of individual hair bundles (13, 23). Coarse microelectrodes were fabricated from

borosilicate capillaries (TW120F-3, World Precision Instrument) with a pipette puller (P97,

Sutter Instrument); their resistance ranged from 0.8 to 1.5 MΩ when filled with 3-M KCl and

immersed in the same solution. The electrode's tip was positioned at ~3 μm from the hair

bundle.

In a first series of experiments, the electrodes were filled with 400-µM pentasodium

triphosphate. Triphosphate allowed to decrease the calcium concentration while preserving the

integrity of the tip links (23). A holding current of $+10$ nA was continuously applied to

counteract the diffusive release of the calcium chelator from the electrode, to which we added

either a step or a succession of descending and ascending ramps of iontophoretic current. The

steps had a magnitude of varying between -80 nA and -60 nA, while the ramps reached a

minimal value between -90 nA and -50 nA. The current varied at the same rate on the

descending and ascending ramps, within a range $5 - 8$ nA $\cdot$ s^{-1}, and the minimal current was

maintained for $1 - 2$ s before the ascending ramp started.

In a second series of experiments, we aimed at disrupting the tip links to evaluate their

contribution to hair-bundle stiffness. In this case, the electrode was filled with EDTA at a

concentration of 100 mM. We applied current steps of -100 nA on top of a resting current of

$+10$ nA. In three cells, reducing the amplitude of the step to about -35 nA allowed to maintain

the integrity of the tip links, adding data to those obtained with iontophoresis of triphosphate to

evaluate the effects of reducing the calcium concentration on a force-displacement relation.

Signal generation and acquisition

All signals were generated and acquired under the control of a computer running a user interface

programmed with the LabVIEW software (version 2011, National Instruments). Command

signals were produced by a 16-bit interface card (PCI-6733, National Instruments). A second

16-bit interface card (PCI-6250, National Instruments) conducted signal acquisition. Sampling

rates for signal generation and acquisition varied within the range $2.5 - 25$ kHz. All signals

were conditioned with an eight-pole Bessel antialiasing filter (Model 3384, Krohn-Hite

Corporation) at a lowpass cut-off frequency set at half the sampling frequency unless otherwise

indicated.

Analysis of spontaneous hair-bundle oscillations

Details of the analysis of the spontaneous oscillations have been published elsewhere (20).

Briefly, the displacement, $X(t)$, of an oscillating hair bundle was monitored over a duration of

$10 - 30$ s. The relation between its power spectrum, $C(f)$, and frequency, f, was first

smoothed by a moving average with a frequency window of 1 Hz and then fitted by the sum of

two Lorentzians

$$\tilde{C}(f) = \frac{A}{(f_0/2Q)^2 + (f - f_0)^2} + \frac{A}{(f_0/2Q)^2 + (f + f_0)^2}. \tag{2}$$

The fit provided values for the quality factor Q, the characteristic frequency f_0, and parameter

A, which is related to the root-mean-square magnitude $X_{RMS} = \langle X(t) - \langle X(t)\rangle^2\rangle =$

$\int_{-\infty}^{+\infty} \tilde{C}(f)df = \sqrt{4\pi AQ/f_0}$ of the movement. In cases where it was bimodal, the probability

density of hair bundle positions was fitted by the sum of two Gaussian functions. The fractional

area under the Gaussian that was centered at a positive position with respect to the mean bundle

position provided a functional estimate of the open probability P_O^{SS} of transduction channels at

steady state.

REFERENCE

1.   A. J. Hudspeth, How the ear's works work. *Nature* **341**, 397–404 (1989).

2.   R. Fettiplace, K. X. Kim, The physiology of mechanoelectrical transduction channels in
hearing. *Physiol. Rev.* **94**, 951–86 (2014).

3.   P. Martin, G. A. Manley, "Auditory processing by the cochlea" in *Principles of Neural*
Science, McGraw-Hill Education / Medical; 6th edition, (2021), pp. 594–624.

4.   D. P. Corey, A. J. Hudspeth, Kinetics of the receptor current in bullfrog saccular hair cells.
J. Neurosci. **3**, 962-976. (1983).

5.   V. S. Markin, A. J. Hudspeth, Gating-spring models of mechanoelectrical transduction by
hair cells of the internal ear. *Annu. Rev. Biophys. Biomol. Struct.* **24**, 59–83 (1995).

6.   J. Howard, A. J. Hudspeth, Compliance of the hair bundle associated with gating of
mechanoelectrical transduction channels in the bullfrog's saccular hair cell. *Neuron* **1**,
189-199. (1988).

7.   A. J. Ricci, A. C. Crawford, R. Fettiplace, Mechanisms of active hair bundle motion in
auditory hair cells. *J. Neurosci.* **22**, 44–52 (2002).

8.   V. Bormuth, J. Barral, J.-F. Joanny, F. Julicher, P. Martin, Transduction channels' gating
can control friction on vibrating hair-cell bundles in the ear. *Proc. Natl. Acad. Sci. USA*
111, 7185–7190 (2014).

9.   P. Martin, A. D. Mehta, A. J. Hudspeth, Negative hair-bundle stiffness betrays a
mechanism for mechanical amplification by the hair cell. *Proc. Natl. Acad. Sci. USA* **97**,
12026–12031 (2000).

10.   I. J. Russell, M. Kossl, G. P. Richardson, Nonlinear mechanical responses of mouse
cochlear hair bundles. *Proc. R. Soc. Lond. B* **250**, 217–27. (1992).

11.   D. K. Chan, A. J. Hudspeth, Ca2+ current–driven nonlinear amplification by the
mammalian cochlea in vitro. *Nat. Neurosci.* **8**, 149–155 (2005).

12.   H. J. Kennedy, A. C. Crawford, R. Fettiplace, Force generation by mammalian hair
bundles supports a role in cochlear amplification. *Nature* **433**, 880–883 (2005).

13.   M. Tobin, A. Chaiyasitdhi, V. Michel, N. Michalski, P. Martin, Stiffness and tension
gradients of the hair cell's tip-link complex in the mammalian cochlea. *eLife* **8**, e43473
(2019).

14.   M. Beurg, E. T. Schwalbach, R. Fettiplace, LHFPL5 is a key element in force transmission
from the tip link to the hair cell mechanotransducer channel. *Proc. Natl. Acad. Sci. USA*
121, e2318270121 (2024).

15. P. Martin, A. J. Hudspeth, Active hair-bundle movements can amplify a hair cell's
response to oscillatory mechanical stimuli. *Proc. Natl. Acad. Sci. USA* **96**, 14306–14311
(1999).

16. P. Martin, A. J. Hudspeth, Mechanical frequency tuning by sensory hair cells, the
receptors and amplifiers of the inner ear. *Annu. Rev. Condens. Matter Phys.* **12**, 29–49
(2021).

17. P. Martin, A. J. Hudspeth, F. Julicher, Comparison of a hair bundle's spontaneous
oscillations with its response to mechanical stimulation reveals the underlying active
process. *Proc. Natl. Acad. Sci. USA* **98**, 14380–14385 (2001).

18. P. Martin, D. Bozovic, Y. Choe, A. J. Hudspeth, Spontaneous oscillation by hair bundles
of the bullfrog's sacculus. *J. Neurosci.* **23**, 4533–4548 (2003).

19. D. Ramunno-Johnson, C. E. Strimbu, L. Fredrickson, K. Arisaka, D. Bozovic, Distribution
of frequencies of spontaneous oscillations in hair cells of the bullfrog sacculus. *Biophys.*
J. **96**, 1159–68 (2009).

20. J. Barral, F. Jülicher, P. Martin, Friction from transduction channels' gating affects
spontaneous hair-bundle oscillations. *Biophys. J.* **114**, 425–436 (2018).

21. N. Hacohen, J. A. Assad, W. J. Smith, D. P. Corey, Regulation of tension on hair-cell
transduction channels: displacement and calcium dependence. *J. Neurosci.* **9**, 3988-3997.
(1989).

22. A. J. Ricci, Y. C. Wu, R. Fettiplace, The endogenous calcium buffer and the time course
of transducer adaptation in auditory hair cells. *J. Neurosci.* **18**, 8261–77 (1998).

23. J.-Y. Tinevez, F. Jülicher, P. Martin, Unifying the various incarnations of active hair-
bundle motility by the vertebrate hair cell. *Biophys. J.* **93**, 4053–4067 (2007).

24. R. E. Marquis, A. J. Hudspeth, Effects of extracellular Ca2+ concentration on hair-bundle
stiffness and gating-spring integrity in hair cells. *Proc. Natl. Acad. Sci. USA* **94**, 11923–
11928 (1997).

25. R. A. Jacobs, A. J. Hudspeth, Ultrastructural correlates of mechanoelectrical transduction
in hair cells of the bullfrog's internal ear. *Cold Spring Harb. Symp. Quant. Biol.* **55**, 547–
561 (1990).

26. C. Köppl, V. Wilms, I. J. Russell, H. G. Nothwang, Evolution of endolymph secretion and
endolymphatic potential generation in the vertebrate inner ear. *Brain. Behav. Evol.* **92**, 1–
31 (2018).

27. D. K. Chan, A. J. Hudspeth, Ca2+ current–driven nonlinear amplification by the
mammalian cochlea in vitro. *Nat. Neurosci.* **8**, 149–155 (2005).

28. Y. Li, H. Liu, X. Zhao, D. Z. He, Endolymphatic Potential Measured From Developing
and Adult Mouse Inner Ear. *Front. Cell. Neurosci.* **14**, 584928 (2020).

29.　P. A. Fuchs, A. M. Lauer, Efferent Inhibition of the Cochlea. *Cold Spring Harb. Perspect.*
Med. **9**, a033530 (2019).

30.　M. Castellano-Muñoz, S. H. Israel, A. J. Hudspeth, Efferent Control of the Electrical and
Mechanical Properties of Hair Cells in the Bullfrog's Sacculus. *PLoS ONE* **5**, e13777
(2010).

31.　J. J. Art, R. Fettiplace, Efferent desensitization of auditory nerve fibre responses in the
cochlea of the turtle Pseudemys scripta elegans. *J. Physiol.* **356**, 507–523 (1984).

32.　C.-H. J. Lin, D. Bozovic, Effects of Efferent Activity on Hair Bundle Mechanics. *J.*
Neurosci. **40**, 2390–2402 (2020).

33.　S. W. Meenderink, P. M. Quinones, D. Bozovic, Voltage-mediated control of spontaneous
bundle oscillations in saccular hair cells. *J. Neurosci.* **35**, 14457–66 (2015).

34.　E. L. Cheung, D. P. Corey, Ca2+ changes the force sensitivity of the hair-cell transduction
channel. *Biophys J* **90**, 124–39 (2006).

35.　Y. C. Wu, A. J. Ricci, R. Fettiplace, Two components of transducer adaptation in auditory
hair cells. *J Neurophysiol* **82**, 2171–81 (1999).

36.　A. J. Ricci, H. J. Kennedy, A. C. Crawford, R. Fettiplace, The transduction channel filter
in auditory hair cells. *J. Neurosci.* **25**, 7831–9 (2005).

37.　M. Beurg, J.-H. Nam, A. Crawford, R. Fettiplace, The actions of calcium on hair bundle
mechanics in mammalian cochlear hair cells. *Biophys. J.* **94**, 2639–2653 (2008).

38.　Y. Roongthumskul, L. Fredrickson-Hemsing, A. Kao, D. Bozovic, Multiple-timescale
dynamics underlying spontaneous oscillations of saccular hair bundles. *Biophys. J.* **101**,
603–610 (2011).

39.　Villasante, C. M., Deng, X., Cohen, J. E., Hudspeth, A.J., Nanomechanics of wild-type
and mutant dimers of the inner-ear tip-link protein protocadherin 15. *Proc. Natl. Acad.*
Sci. USA **121**, e2404829121 (2024).

40.　Y. R. Guo, R. MacKinnon, Structure-based membrane dome mechanism for Piezo
mechanosensitivity. *eLife* **6**, e33660 (2017).

41.　B. Pan, *et al.*, TMC1 forms the pore of mechanosensory transduction channels in
vertebrate inner ear hair cells. *Neuron* **99**, 736-753.e6 (2018).

42.　W. Zheng, J. R. Holt, The mechanosensory transduction machinery in inner ear hair cells.
Annu. Rev. Biophys. **50**, 31–51 (2021).

43.　X. Qiu, U. Müller, Sensing sound: Cellular specializations and molecular force sensors.
Neuron **110**, 3667–3687 (2022).

44.　J. Ge, *et al.*, Structure of mouse protocadherin 15 of the stereocilia tip link in complex
with LHFPL5. *eLife* **7**, e38770 (2018).

45. H. Jeong, *et al.*, Structures of the TMC-1 complex illuminate mechanosensory
transduction. *Nature* **610**, 796–803 (2022).

46. F. Gianoli, T. Risler, A. S. Kozlov, Lipid bilayer mediates ion-channel cooperativity in a
model of hair-cell mechanotransduction. *Proc. Natl. Acad. Sci. USA* **114**, E11010–E11019
(2017).

47. R. Vallone, G. Dal Cortivo, M. D'Onofrio, D. Dell'Orco, Preferential binding of Mg2+
over Ca2+ to CIB2 triggers an allosteric switch impaired in Usher syndrome type 1J.
Front. Mol. Neurosci. **11**, 274 (2018).

48. S. Wu, *et al.*, Mechano-electrical transduction components TMC1-CIB2 undergo a Ca2+-
induced conformational change linked to hearing loss. *Dev. Cell* S1534580725000048
(2025). https://doi.org/10.1016/j.devcel.2025.01.004.

49. S. S. George, C. R. Steele, A. J. Ricci, Rat Auditory Inner Hair Cell Mechanotransduction
and Stereociliary Membrane Diffusivity Are Similarly Modulated by Calcium. *iScience*
23, 101773 (2020).

50. A. J. Hudspeth, P. Martin, The Critical Thing about the Ear's Sensory Hair Cells. *J.
Neurosci.* **44**, e1583242024 (2024).

51. A. Neveu, Suitability of European green frogs for intensive culture: Comparison between
different phenotypes of the esculenta hybridogenetic complex. *Aquaculture* **295**, 30–37
(2009).

Acknowledgements

Funding:

Fondation pour l'Audition grant FPA RD 2020-7 (PM)

Labex Cell(n)Scale grant ANR-11-LABX-0038 (PM)

Labex Cell(n)Scale grant ANR-10-IDEX-001-02 (PM)

Université Paris Sciences et Lettres and doctoral school "Physique en Île-de-France" grant (AJ)

Author contributions:

Competing interest:

The authors declare that they have no conflict of interest.

Data and materials availability: All data needed to evaluate the conclusions in the paper are present in the paper and/or the Supplementary Materials. Datasets for all the figures in the Main Text and Supplementary materials can be found on Zenodo (https://doi.org/10.5281/zenodo.14452874). Additional data related to this paper may be requested from the authors.

     MAIN-FIGURE LEGENDS

Fig. 1 : Representative examples of mechanical changes evoked by a transepithelial current. **(A-D)** Force-displacement relations, $F(X)$, for four different hair bundles under control conditions ($I = 0$ µA; black) and in the presence of a transepithelial current (magenta) of amplitude $I = +3$ µA (A), $I = +6$ µA (B), $I = -3$ µA (C) and $I = -6$ µA (D). Each relation was fitted by Equation 1 (solid lines; parameter values in table S1). Insets: the fitted curves are replotted with respect to the origin $(X_0, F(X_0))$, where X_0 is the set-point deflection. Under each condition, the hair bundle's gating force, F_G, was given by the vertical distance between the two symbols in the inset ($I = 0$: triangles; $I \neq 0$: disks). In (A), (B) and (D), the open black symbols correspond to a measurement after restoring control conditions to assess reversibility; in (C), the experiment was repeated 4 times.

Fig. 2 : Changes in set-point deflection, stiffness and gating force upon application of a
transepithelial current. (A) The change, ΔX_0, of the set-point deflection showed a strong
positive correlation to the transepithelial current, I (Pearson correlation r $= 0.87$; p-value: $4 \cdot$
10^{-11}; $n = 34$). (B) The change, ΔK, of the linear stiffness was negatively correlated to the
current, but only weakly (Pearson correlation r $= -0.57$; p-value: 10^{-4}; $n = 41$). (C) For
I $> -4\ \mu$A, the change, ΔF_G, of the bundle's gating force showed a strong negative correlation
to the current (Pearson correlation r $= -0.86$; p-value: $8 \cdot 10^{-11}$; $n = 34$). (D) Relative
change of the hair-bundle's gating force, $\Delta F_G / F_G(0)$ (solid disks) and linear stiffness, $\Delta K / K(0)$
(open disks) as a function of the transepithelial current, I. Under control conditions ($I = 0$), the
gating force was $F_G = 17 \pm 4\ $pN (mean $\pm$ SD, $n = 29$). In (C) and (D), the red and blue
dashed lines mark, respectively, a reduction and an increase of the gating force by 100% with
respect to the mean value under control conditions. For $I < -5\ \mu$A, all the force displacement
relations were linear: the gating force had dropped to undetectable levels and the set-point
deflection could no longer be measured from the mechanics.
713

Fig. 3: Electrical control of the gating swing. The gating swing, d, **(A)** and its relative change,

$\Delta d/d_0$, **(B)** are shown as a function of the transepithelial current, I, where $\Delta d = d(I) -$

$d(I = 0)$ and $d_0 = d(I = 0)$. For $I \geq -5$ μA, the relative change displayed a strong negative

correlation to the current (Pearson correlation $r = -0.86$; p-value: $1 \cdot 10^{-10}$). Same dataset as

that used in Fig. 2. After each measurement of a bundle's force-displacement relation, the

gating swing was estimated as $d = \gamma F_G/(K - K_{SP})$, where F_G and K are respectively the gating

force and linear stiffness resulting from a fit of the force-displacement relation with Equation 1

and K_{SP} is the mean stiffness measured over an ensemble of hair bundles after disruption of

their tip links. The projection factor $\gamma = 0.14$ was used to estimate the gating swing along the

oblique axis of the tip link.

Fig. 4 : **Transition between states of strong and weak gating force.** **(A)** A descending ramp of a negative transepithelial current was followed by an ascending ramp of current at the same absolute rate of $0.9 \, \mu A \cdot s^{-1}$. The colored square symbols indicate the time points of the analysis shown in (C). **(B)** The hair bundle was continuously moved back and forth (bottom) by imposing a periodic triangular movement to the stimulus fiber's base, here with a peak-to-peak amplitude of 600 nm and at a frequency of 40 Hz (top). The movements were averaged over 10 stimulus cycles. **(C)** Force-displacement cycles for the values of the transepithelial current and at times indicated at the top of each sub-panel (see colored square symbols in (A)). **(D)** Trajectory of the point of inversion symmetry (X_0, F_0) in a force-displacement cycle, marked by disks in (C). Linear fits to these trajectories: $F_0 = 0.41 \cdot X_0 + 9$ ($R^2 = 0.92$) (magenta) and $F_0 = 0.78 \cdot X_0 + 24$ ($R^2 = 0.98$) (orange), respectively. **(E)** Maximal friction force, ϕ_{MAX}, as a function of the transepithelial current, I. **(F)** For all the 8 different hair bundles that followed the same protocol, the maximal friction force, ϕ_{MAX}, first increased as the

transepithelial current decreased, before dropping to a low value as the hair bundle transitioned

from a state of strong gating force to a state of weak gating force at more negative values of the

current than a threshold $I = I_{DROP} = -3.7 \pm 1\ \mu A$ (mean $\pm$ SD, $n = 8$). Colored disks mark,

for each hair-bundle, the value of ϕ_{MAX} at the start of the experiment ($I = 0$, black), before

($I = I_{DROP} + 1\ \mu A$, magenta) and after ($I = -9\ \mu A$, red) the drop of ϕ_{MAX}, and at the end of

the experiment ($I = 0\ \mu A$, white). Error bars: mean $\pm$ SD. Statistical significance was assessed

using a paired t-test with p-values from left to right: $2 \cdot 10^{-2}$, $8 \cdot 10^{-4}$, 10^{-1}. ns > 0.05, * p-

value ≤ 0.05, *** p-value ≤ 0.001. In (D-E), black solid and dashed lines correspond to the

descending and ascending ramps of transepithelial currents, respectively. Arrows indicate the

direction of circulation along the curve from start (black disk) to finish (white disk).

Fig. 5 : Transition kinetics between states of strong and weak gating force. **(A)** Force-displacement cycle here averaged over 50 cycles of stimulation, under control conditions ($I = 0$; black solid line), 400 ms after the onset of a step of transepithelial current ($I = -8\ \mu$A; magenta) and 600 ms after termination of the current step ($I = 0$; black dashed line). **(B)** Time course of the maximal friction force, ϕ_{MAX}, (blue line) and of the average displacement, $\bar{X}$, (dashed magenta line) in a force-displacement cycle. The current step started at $t = 1$ s and ended at $t = 2$ s. **(C)** For all the 4 different hair bundles that followed the same protocol, time, τ_{10-90}, required for the maximal friction force, ϕ_{MAX} (blue disks), and the average displacement, $\bar{X}$ (magenta disks), to transition from 10% to 90% of the total change after the onset (ON) and the termination (OFF) of a step of transepithelial current ($I = -8\ \mu$A). Error bars indicate the mean $\pm$ SD. Statistical significance was assessed using a paired t-test with p-values from left to right: $2 \cdot 10^{-2}$, $3 \cdot 10^{-2}$; * p-value ≤ 0.05. The base of the stimulus fiber followed a periodic triangular waveform of motion, with a peak-to-peak amplitude of 600 nm and at a frequency of 100 Hz. For the data shown in (B) and (C), the force-displacement cycles were averaged over only two cycles of stimulation.

Fig. 6 : Replicating the effects of negative endolymphatic potentials with iontophoresis of a Ca²⁺ chelator. **(A)** We used iontophoresis of a calcium chelator to decrease the extracellular Ca^{2+} concentration in the vicinity of a hair bundle. A descending ramp of a negative iontophoretic current, I_{IONTO}, which was carried in part by a Ca^{2+} chelator, was followed by an ascending ramp of current at the same absolute rate of $6.6 \text{ nA} \cdot \text{s}^{-1}$. The colored square symbols indicate the time points of the analysis shown in (B). **(B)** Each sub-panel shows the bundle's force-displacement cycle in response to a triangular mechanical stimulus for the value of the iontophoretic current, I_{IONTO}, and at the time, t, indicated at the top. **(C)** Maximal friction force, ϕ_{MAX}, as a function of the iontophoretic current, I_{IONTO}, which varied in time from start (black disk) to finish (white disk). Arrows indicate the direction of circulation along the curve. Solid and dashed lines correspond to the descending and ascending ramps of iontophoretic currents, respectively. **(D)** For all the 8 different hair bundles that followed the same protocol, the maximal friction force, ϕ_{MAX}, first increased as the extracellular Ca^{2+} concentration

decreased, before dropping to a low value as the hair bundle transitioned from a state of strong

gating force to a state of weak gating force at more negative values of the iontophoretic current

than a threshold $I_{IONTO} = I_{DROP} = -40 \pm 10$ nA (mean $\pm$ SD, $n = 8$). Colored disks mark, for

each hair-bundle, the value of ϕ_{MAX} at the start of the experiment ($I_{IONTO} = +10$ nA, black),

before ($I_{IONTO} = I_{DROP} + 5$ nA, magenta) and after ($I_{IONTO} = -50$ nA, red) the drop of ϕ_{MAX},

and at the end of the experiment ($I_{IONTO} = +10$ nA, white). Error bars: mean $\pm$ SD. Statistical

significance was assessed using a paired t-test with p-values from left to right: $1.5 \cdot 10^{-2}$, $5 \cdot$

10^{-3}, $6 \cdot 10^{-1}$; ns > 0.05; * p-value ≤ 0.05, ** p-value ≤ 0.01. (E) The force-displacement

relation of a different hair bundle displays a region of gating compliance under control

conditions (black triangles) but was linear in the presence of a steady iontophoretic current of

-60 nA (red circle). This effect was reversible (grey triangles). Each force-displacement

relation was fitted by Equation 1 (solid lines; parameter values in table S1). In (B-D), a periodic

triangular waveform of motion was applied to the base of the stimulus fiber, with a peak-to-

peak amplitude of 600 nm and at a frequency of 40 Hz. In (E), force-displacement relations

were instead measured in response to step displacements at the base of the stimulus fiber (see

Methods).

<p style="text-align:center">Supplementary Materials for</p>

Electrical control of the transduction channels' gating force in sensory hair cells

<p style="text-align:center">Achille Joliot[1,2] , Laure Stickel[1,2] and Pascal Martin[1,2,*]</p>

[1] Physique des Cellules et Cancer, Institut Curie, PSL Research University, CNRS UMR168, F-75248 Paris, France.

[2] Sorbonne Université, F-75252 Paris, France.

[*] Correspondence: pascal.martin@curie.fr

The supplementary materials include:

Table S1

Figures S1 to S4

Legends for Movies S1 to S3

Other supporting materials for this manuscript include the following:

Movies S1 to S3

822   **Supplementary Table**

823   **Table S1** Parameter values for fits of force-displacement relations with Equation 1.

Variable	Definition	Fig. 1A		Fig. 1B		Fig. 1C		Fig. 1D		Fig. 5E	
I (μA)	Transepithelial current	0	+3	0	+6	0	−3	0	−6	n/a	n/a
I_{IONTO} (nA)	Iontophoretic current,	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	+10	−60
X_0 (nm)	Set-point deflection	−3	1	1	+6	−0.2	−12.7	−2	n/a	−14	n/a
K (pN $\cdot$ nm^{-1})	Stiffness	0.96	0.91	0.83	0.78	0.6	0.66	0.82	0.96	0.75	0.80
F_G (pN)	Gating force	21	13	18	9	19	36	22	n/a	22	n/a
δ (nm)	Characteristic length	4	2	2	5	8.9	13	6	n/a	7	n/a
$F_\emptyset$ (pN)	Force constant	14	5	7	0.6	10	25	11	n/a	18	n/a
R^2	Coefficient of determination	0.999	0.994	0.986	0.986	0.986	0.986	0.986	0.986	0.979	0.993

Supplementary Figures

Fig. S1 : Gating-spring model of mechanoelectrical transduction.   **(A)** The concept of
gating force. When the transduction channels switch between their open and closed state,
tension in the gating springs is reduced (black arrows); this mechanical correlate of channel
gating is called the gating force, $F_G = K_{GS} \cdot D$ (red arrow), where K_{GS} is the combined stiffness
of all the gating springs operating in parallel in the hair bundle and D is the gating swing of
each channel.   **(B)** Electrical and mechanical responses to hair-bundle displacement. Top:
sigmoidal relationship of channel open probability to displacement in the case of weak (dashed
magenta line) and strong (solid blue line) gating force. The larger the gating force, the steeper
the sigmoid and the more sensitive the hair cell is to displacement of its hair bundle. Bottom:
The force–displacement relationship displays a shallow nonlinearity when sensitivity is low
(dashed magenta curve) and a strong nonlinearity when sensitivity is high (smooth blue curve).
The gating force is given by the vertical shift between the two linear limbs of the force-
displacement relation; it is here shown in the case of high sensitivity as the distance between
the two disks.

Fig. S2 : Two-compartment ex-vivo preparation of the frog saccule. **(A)** Schematic of the
setup. The tissue was glued over a 1-mm hole in a plastic disk. Its apical and basal aspects were
bathed in artificial endolymph and perilymph, respectively. A transepithelial current, I, was
produced by a pair of silver chloride electrodes connected to a current generator, resulting in an
endolymphatic potential, U, with respect to the potential in perilymph. The sign convention is
such that positive currents lead to a positive value of the endolymphatic potential. **(B)** A +1-
μA step of transepithelial current (top) evoked an endolymphatic potential $U \simeq +10$ mV at
steady state (bottom, grey), corresponding to a transepithelial resistance of 10 kΩ. An
exponential fit (black line), $U(t) = A \exp(-t/\tau)$, to the response yielded the amplitude $A =$
10.2 mV and the characteristic time $\tau = 210$ μs.

Fig. S3 : Effects of a transepithelial current on spontaneous hair-bundle oscillations.

Under control conditions ($I = 0$), over an ensemble of $n = 10$ oscillatory hair bundles, the

oscillations had a characteristic frequency $f_0 = 22 \pm 14$ Hz, a root-mean-squared (RMS)

amplitude $X_{RMS} = 13 \pm 4$ nm and their bimodal position histograms were associated with a

steady-state open probability for the transduction channels $P_o^{SS} = 0.4 \pm 0.1$, thus close to $1/2$.

These characteristics are remarkably similar to those reported earlier with hair cells from a

different frog species (18, 20). **(A-D)** Representative examples of spontaneous hair-bundle

oscillations under control conditions ($I = 0$; top) and in the presence of a transepithelial current

(bottom) $I = +3$ µA (A), $I = +6$ µA (B), $I = -3$ µA (C), and $I = -6$ µA (D). **(E)** The change,

Δf_0, of the oscillation frequency displayed a strong positive correlation to the transepithelial

current (Pearson correlation $r = +0.9$; p-value $= 10^{-4}$; $n = 10$). **(F)** The change, ΔX_{RMS}, of

the RMS amplitude of spontaneous oscillations displayed a strong negative correlation to the

transepithelial current (Pearson correlation $r = -0.9$; p-value $= 1 \cdot 10^{-3}$; $n = 10$). **(G)** The

change, ΔP_o^{SS}, of the open probability at steady state displayed a strong negative correlation to

the transepithelial current (Pearson correlation r $= -0.9$, p-value $= 1 \cdot 10^{-4}$, $n = 10$). In (A),

the transepithelial current evoked an increase of the oscillation frequency from 17 Hz to 39 Hz,

a decrease of the RMS amplitude of oscillation from 7 nm to 5 nm and a decrease of the steady-

state open probability from 0.5 to 0.3. Conversely in (B), the frequency of oscillation decreased

from 49 Hz to 25 Hz, the RMS amplitude of oscillation increased from 11 nm to 15 nm and the

steady-state open probability increased from 0.4 to 0.6. In the presence of a large positive

current ($I = +6$ µA; panel B), the hair bundle only produced irregular spiky movements in the

positive direction: the transduction channels were closed most of the time and evinced brief

openings. Instead, a large negative current ($I = -6$ µA; panel D) completely abolished

spontaneous oscillations. Red lines in (E-G) correspond to proportional fits of respective slopes

$7\,\text{Hz} \cdot \mu\text{A}^{-1}$ ($R^2 = 0.86$; panel E), $-1.1\,\text{nm} \cdot \mu\text{A}^{-1}$ ($R^2 = 0.6$; panel F), and $-0.08\,\mu\text{A}^{-1}$ ($R^2 =$

0.83; panel G).

Fig. S4 : Effect of a transepithelial current on the channel-activation time. Same data as in

Fig. 4A-E, to which this figure is associated. The time course of successive ramps of descending

and ascending transepithelial current can be found in Fig. 4A. **(A)** Fiber-base displacement, Δ

(*top*), and hair-bundle displacement, X (*bottom*), as a function of time for one stimulus cycle at

time points and values of the transepithelial current indicated at the top of each time trace. **(B)**

Hair-bundle speed, $|dX/dt|$, on the positive (*solid line*) and negative (*dashed line*) half cycles

of mechanical stimulation as a function of displacement, X. The distance between the two

peaks, ΔX, defines the gating shift. **(C-D)** Maximum speed, V_{MAX} (C), and gating shift, ΔX

(D), as a function of the transepithelial current, I, which varied in time from start (black disk)

to finish (white disk). Arrows indicate the direction of circulation along the curve. Solid and

dashed lines correspond to the descending and ascending ramps of transepithelial currents,

respectively. In panel (D), a black square indicates the current at which the signatures of gating

compliance and gating friction vanished on the descending ramp; bellow this current, the gating

shift is no longer defined. Conversely, a white square indicates recovery of the signatures of

gating compliance and gating friction on the ascending ramp. **(E)** For the same ensemble of

hair bundles as in Fig. 4, the distributions of the estimated activation time of the transduction

channel, $\tau = \Delta X/(2\,V_{MAX})$, are shown at the start of the experiment ($I = 0$, black), before the

drop of the maximal friction force (magenta, see Fig. 4F), and at the end of the experiment ($I =$

0, white). Error bars: mean $\pm$ SD. Statistical significance was assessed using a paired t-test

with, from p-values from left to right: $0.0014, 0.021$ *p-value $\leq$ 0.05, **p-value $\leq$ 0.01.

Legends for supplementary movies

Movie S1 (separate file): Transition between states of strong and weak gating force. Movie associated to Fig. 3A-E. The force-displacement cycle in response to triangular mechanical stimulation of the hair bundle (top) is shown over time as the transepithelial current follows successive descending and ascending ramps (bottom). The friction force, $\phi(X)$, at each displacement X of the hair bundle is color coded as indicated on the right; it is given by the half-height of the force-displacement cycle. To obtain smooth videos, the force-displacement cycles were here averaged over 15 cycles, instead of 10 cycles in the data shown in Fig. 3.

Movie S2 (separate file): Kinetics of the transition between states of strong and weak gating force. Movie associated to Fig. 4A-B. The force-displacement cycle in response to triangular mechanical stimulation of the hair bundle (top) is shown over time before, during and after application of a negative step of transepithelial current (bottom). The friction force, $\phi(X)$, at each displacement X of the hair bundle is color coded as indicated on the right; it is given by the half-height of the force-displacement cycle. The force-displacement cycles were here averaged over 5 cycles.

Movie S3 Transition between states of strong and weak gating force upon a decrease of extracellular calcium concentration. Movie associated to Fig. 5A-D. The force-displacement cycle in response to triangular mechanical stimulation of the hair bundle (top) is shown over time as an iontophoretic current trough a pipette containing a calcium chelator and positioned near a hair bundle follows successive descending and ascending ramps (bottom). The friction force, $\phi(X)$, at each displacement X of the hair bundle is color coded as indicated on the right; it is given by the half-height of the force-displacement cycle. To obtain smooth videos, the force-displacement cycles were here averaged over 15 cycles, instead of 10 cycles for the data shown in Fig. 5.