Thyroid hormones act as a timer for the postnatal maturation of parvalbumin neurons in mouse neocortex

Juan Ren[1, 3], Suzy Markossian[1], Romain Guyot[1], Denise Aubert[1], Jacques Brocard[2], Jiemin Wong[3], Frédéric Flamant[1,*], Sabine Richard[1]

[1] Ecole Normale Supérieure de Lyon, INRAE, CNRS, Institut de Génomique Fonctionnelle de Lyon, 69364 Lyon, France

[2] Universite Claude Bernard Lyon 1, CNRS UAR3444, Inserm US8, ENS de Lyon, SFR Biosciences, Lyon 69007, France

[3] Shanghai Key Laboratory of Regulatory Biology, Institute of Biomedical Sciences and School of Life Sciences, East China Normal University, 200241, Shanghai, China

[*] Corresponding author. Frederic.flamant@ens-lyon.fr Tel: 00 (33) 4 26 73 13 32

Significance statement

Thyroid hormones are crucial for neurodevelopment, but the underlying mechanisms are still
unclear. In the neocortex, mutations in thyroid hormone nuclear receptors affect the maturation of
GABAergic inhibitory interneurons, and in particular a subset of these neurons that express
parvalbumin. We found that, during a defined postnatal development stage, thyroid hormones are
essential for the maturation of parvalbumin neurons. They notably promote the formation of the
perineuronal net, a specialized extracellular matrix that contributes to the maturation of neocortical
networks by stabilizing synapses around parvalbumin neurons. Experimental evidence strongly
suggests that thyroid hormones act as a timer to define the temporal boundaries of the well-known
critical period of heightened plasticity, during which neuronal networks are shaped by experience.

Abstract

Thyroid hormones (THs) are required for proper maturation of parvalbumin-expressing GABAergic
interneurons (PV neurons) in the mouse neocortex. However, the timeline of this TH action remains
to be elucidated. Here, we used several mouse models expressing dominant negative mutations of
the TH nuclear receptor α1 (TRα1), to better define the time window during which TH promotes the
postnatal maturation of PV neurons. Histological observations revealed that TH action during the first
three postnatal weeks was necessary to initiate the expression of parvalbumin and the elaboration of
a specialized extracellular matrix called perineuronal net. By contrast, after the third postnatal week,
TH action on PV neuron maturation appeared to be somewhat dispensable. Transcriptome analysis
of neocortical GABAergic neurons identified a small set of putative target genes for TRα1. These
genes are at the origin of the postnatal remodeling of the repertoire of ions channels and the
elaboration of perineuronal nets. These data suggest that THs act as a timer to define the temporal
boundaries of the critical period of heightened plasticity, which plays a fundamental role in the
development of neuronal circuits.

Keywords

Introduction

PV neurons are GABAergic inhibitory interneurons that play a central role in the organization of
neuronal circuits in the neocortex (1). They are fast-spiking and synchronize network oscillations,
which are thought to play an important role in information processing and memory formation. In
mice, PV neurons begin to express parvalbumin during the first postnatal weeks (2, 3). During this
time, the electrical activity of these neurons progressively increases (4) and synaptogenesis is very
active. This opens the so-called critical period of heightened synaptic plasticity (5) during which
sensory input and experience shape the neuronal networks of the neocortex. The critical period ends
when excessive circuit rewiring becomes actively dampened (6). Notably, the elaboration of the
perineuronal net (PNN) around PV neurons, a densely packed extracellular matrix enriched in
proteoglycans stabilizes synapses and restrict plasticity (7). Defects in PV neuron maturation during
the critical period cause a persistent imbalance between neuronal excitation and inhibition, which is
suspected to be at the origin of several pathological conditions, including attention deficit
hyperactivity disorder, autism, and epilepsy (8).

THs (including T3 or 3,3',5-triiodo-L-thyronine, and the less active precursor T4 or thyroxine) exert a
broad influence on brain development. They directly act on gene transcription by binding to the TRα1
and TRβ1/2 nuclear receptors, which are encoded by the *Thra* and *Thrb* genes in mice. Unlike
TRβ1/2, TRα1 is ubiquitous from early stages of development. In both humans and mice, TRα1
mutations have dramatic and irreversible consequences on neurodevelopment, and these
consequences are similar to those of TH deficiency (9-11). Although all types of neural cells are
directly or indirectly dependent on THs for proper terminal differentiation, previous studies have
revealed that in different brain regions, the cell-autonomous influence of THs is limited to a fraction
of neural cells, and eventually amplified by TH-dependent release of neurotrophic factors (12).

In the postnatal mouse neocortex, PV neurons display a high sensitivity to TRα1 mutation (13) and TH
deficiency (14, 15). We have previously found that expressing TRα1^{L400R}, a dominant-negative TRα1
receptor, only in the GABAergic lineage from an early neurodevelopmental stage, is sufficient to
dramatically inhibit the expression of parvalbumin and the elaboration of PNNs (16). By contrast,
other subtypes of neocortical GABAergic interneurons, expressing somatostatin or calretinin, are
much less affected. We have also recently observed that the postnatal expression of TRα1^{L400R} in PV
neurons has no visible effects on their maturation and has only moderate consequences on their
function (Ren et al., 2024). This suggests that TH signaling stimulates the maturation of PV neurons
during a specific time window and then becomes less important to maintain their function.
Accordingly, the appearance of hypothyroidism in the adult stage (14, 17) does not prevent the
expression of parvalbumin. Therefore, it seems that THs exert a selective action on PV neuron

maturation during a specific time window. However, the boundaries of this time window remain
elusive. Moreover, whether blocking TH signaling during development only delays or permanently
inhibits PV neuron maturation remains an open question.

To gain more insight into the timeline of TH action on PV neuron maturation, we generated novel
mouse models to restrict the expression of dominant-negative TRα1 receptors in GABAergic neurons
during specific time windows. Histological observations of parvalbumin and PNNs in the
somatosensory cortex allowed us to pin point a key time window for TH action around the second
postnatal week. A genome-wide analysis of GABAergic neuron transcriptome in the whole neocortex
also allowed us to identify a set of genes whose regulation is expected to be involved in TH-mediated
maturation of PV neurons.

Results

TRα1^{E395fs401X} expression in GABAergic neurons from an early developmental stage permanently
inhibits PV neuron maturation

In our previous work, we used Cre/loxP recombination to restrict the expression of the dominant-
negative TRα1^{L400R} mutant receptor in the GABAergic lineage from embryonic day 12.5 (16). As most
of the mice had frequent epileptic seizures and died before postnatal day 21 (PND21), we could not
address a possible recovery of PV neuron maturation during adulthood. We thus used here a similar
model, based on the _Thra^{Slox}_ allele, encoding TRα1^{E395fs401X}, a receptor with moderate dominant-
negative activity (18). The floxed _ROSA-tdTomato^{lox}_ reporter transgene was also introduced to trace
the cells belonging to the GABAergic lineage. One third of _Thra$^{Slox/+}$ ROSA-tdTomato$^{lox/+}$ Gad2-Cre_ mice
survived to adulthood. We therefore compared these mutant mice and _Thra$^{+/+}$ ROSA-tdTomato$^{lox/+}$_
Gad2-Cre littermates (controls), to histologically study whether TRα1 mutation only delays or
permanently inhibits PV neuron maturation.

At PND21, while the red fluorescence did not indicate any cell loss in the GABAergic lineage
(Supplementary Fig. 1), the density of parvalbumin-positive (PV+) cells was drastically lower in
mutants than in controls (Fig. 1a, b). PNNs were labeled using Wisteria Floribunda Agglutinin (WFA),
which binds N-acetyl-galactosamine-β1 residues of PNN glycoproteins. The density of WFA-positive
(WFA+) cells was also significantly lower in mutants (Fig. 1a, c). Interestingly, the two phenotypes
were accompanied by a clearly lower ratio of PV+ cells that were surrounded by PNNs in mutants,
compared to controls (Fig. 1a, d). At adult stage, the same analysis showed signs of some recovery of
PV neuron density in mutants (Fig. 1e, f), while the PNN defect persisted (Fig. 1e, g and h). Overall,

TRα1^{E395fs401X} appears to have a similar negative influence to TRα1^{L400R} and permanently impairs the

maturation of PV neurons.

Postnatal expression of TRα1^{L400R} in GABAergic neurons is sufficient to inhibit PV neuron

maturation

We have recently observed that the postnatal expression of TRα1^{L400R} in PV neurons has little effect

on their maturation (Ren et al., 2024). Although adult *Thra$^{AMI/+}$ ROSA-tdTomato$^{lox/+}$ Pvalb-Cre* mice

displayed minor histological, electrophysiological and behavioral defects, they did not spontaneously

present epileptic seizures or lethality (19). This suggests that TH signaling intensively stimulates the

maturation of PV neurons during a specific time window, and then becomes secondary to maintain

their function. To better evaluate the temporal restriction of TH signaling for PV neurons maturation,

we used *Thra$^{AMI/+}$ ROSA-tdTomato$^{lox/+}$ Gad2-CreER^{T2}* mice (mutants) and *Thra$^{+/+}$ ROSA-tdTomato$^{lox/+}$*

Gad2-CreER^{T2} or *Gad2-CreER^{T2}* negative mice (controls), where Cre/loxP recombination can be

induced at a chosen time by tamoxifen treatment.

A first batch of these mice were treated with tamoxifen at PND7 and histological analysis was

performed at PND14. This analysis revealed that expressing TRα1^{L400R} during the second postnatal

week was sufficient to compromise the onset of parvalbumin expression (Fig. 2a, b). But the

amplitude of the defect in PV expression in mutant mice was not as high as that in our previous work,

where TH signaling was blocked from embryonic day 12.5 to PND14 in *Thra$^{AMI/+}$ ROSA-tdTomato$^{lox/+}$*

Gad2-Cre mice (16). Besides, we did not observe any clear change in either the density of WFA+ cells

(Fig. 2a, c) or the ratio of PV+ cells that were surrounded by PNNs (Fig. 2a, d). This may be due to the

fact that the elaboration of PNNs mainly takes place after PND14, so that the density of WFA+ cells

was low, with high interindividual variability. By contrast, a tamoxifen treatment performed during

the fourth postnatal week (from PND21 to PND25), did not cause any significant change in either PV+

cell density or WFA+ cell density at adult stage (Fig. 2e-h). Therefore, THs exert their influence during

a postnatal time window, which includes the second postnatal week and precedes weaning, since

weaning usually occurs after the end of the third postnatal week in mice.

Consequences of TRα1^{L400R} expression on gene expression in neocortical GABAergic neurons two

weeks after birth

We set out to analyze the influence of TRα1^{L400R} expression on the nuclear transcriptome of

GABAergic neurons at PND14, a developmental stage which appears to be quite central in the

timeline of TH action in PV neurons. To do this, we used the *ROSA-GSL10gfp^{lox}* reporter transgene,

which labels cell nuclei, to extract nuclear RNA from neocortical GABAergic neurons, in which PV

neurons represent around 50% of the population (20). RNA-seq analysis was performed for $Thra^{AMI/+}$

$ROSA\text{-}GSL10gfp^{lox/+}$ $Gad2\text{-}Cre$ mice and $Thra^{+/+}$ $ROSA\text{-}GSL10gfp^{lox/+}$ $Gad2\text{-}Cre$ control littermates at

PND14. Differential analysis (Deseq2 adjusted p-value > 0.05; fold-change >2.0; basemean > 30)

identified 275 down-regulated genes and 265 up-regulated genes in mutant mice. This large gene set

contains TRα1 target genes whose transcription is normally up-regulated by liganded TRα1. Notably,

among these were genes (*Hr, Klf9, Dbp, Shh, Gabrd*) that are known to be TRα1 target genes in other

cell types (21). However, many of the observed changes in mRNA level probably represent indirect

consequences of the long-term influence of TRα1^{L400R} expression on the cell physiology.

To distinguish direct TRα1 target genes from other genes that are downstream in the cascade, we

carried out a series of additional studies. First, we looked for genes which are down-regulated by

hypothyroidism and up-regulated at PND14 after a short TH treatment. To do this, we used a

propylthiouracyl (PTU) treatment to induce hypothyroidism in gestating females and their offspring.

Half of the hypothyroid pups with a $Thra^{+/+}$ $ROSA\text{-}GSL10gfp^{lox/+}$ $Gad2\text{-}Cre$ genotype then received THs

at PND13. The nuclear RNA transcriptome of GABAergic neurons in the neocortex from hypothyroid

pups, and from pups treated with THs for 24 hours, was analyzed as above. In this setting we

expected a selective recovery of mRNA level for TRα1 target genes.

A total of 1306 genes were sensitive to either TRα1^{L400R} expression, hypothyroidism, or TH

stimulation (Fig. 3a, b). Among them, a large set of genes were only deregulated in hypothyroid mice.

This suggests an important influence of hypothyroidism on the microenvironment (notably glial cells)

of GABAergic neurons, with indirect consequences on gene expression in GABAergic neurons. The

genes whose expression in GABAergic neurons was sensitive to both TRα1^{L400R} expression and

hypothyroidism rather reflect the cell-autonomous influence of TH signaling in these neurons.

Importantly, among a total of 388 of such genes was the *Pvalb* gene, which encodes parvalbumin.

Recovery after TH treatment (TH response reached a two-fold threshold) was achieved for only a

small subset of these genes (61 genes up-regulated after TH treatment and 5 genes down-regulated

after TH treatment). However, clustering analysis revealed a clear trend toward recovery, which was

more visible for genes up-regulated by THs (Fig. 3c) than for genes down-regulated by THs (Fig. 3d).

The 5 genes which were down-regulated after TH treatment could reflect a poorly understood

negative regulation of transcription by the liganded TRα1 (Fig. 3b). In the end, only 61 genes behaved

as *bona fide* TRα1 target genes, with a robust negative response to both TRα1^{L400R} expression and

hypothyroidism, and a clear induction after the short TH treatment of hypothyroid mice (Fig. 3a).

Gene Ontology analysis indicated that many of the genes deregulated in hypothyroid and mutant

mice encode components of the extracellular matrix (GO:0031012; enrichment 4.4, p-value 3.10^{-13}).

This includes genes encoding collagens or proteins binding to the glycosaminoglycans of PNNs

(GO:0005539; enrichment 4.2, p-value 5.10^{-6}). This is consistent with the hypothesis that TH/TRα1

signaling stimulates the elaboration of PNNs around PV neurons. The expression of two of the

downregulated genes in this family (*Sema7a,* encoding semaphorin 7A and *Hapnl4,* encoding

hyaluronan and proteoglycan link protein 4) was restored after a short treatment with THs,

suggesting that they are TRα1 target genes.

Another prominent characteristic of deregulated genes in hypothyroid and mutant mice was the

presence of genes that encode voltage-gated potassium channels (GO:0005249; enrichment 6.1, p-

value 10^{-6}). As these genes appeared to be insensitive to acute TH treatment, their expression is

unlikely to be under the direct control of TRα1. This observation suggests that hypothyroidism or

long-term expression of TRα1^{L400R} in GABAergic neurons damage the postnatal remodeling of ions

channels, which is a key event for proper maturation of PV neurons in the neocortex (4). To further

evaluate this possibility, we extracted the genes that mostly encode ions channels and whose

expression is modified in this process of postnatal maturation as defined by Okaty and colleagues (4).

This analysis (Fig. 3e) revealed that this postnatal transition is impaired and that the immature status

of PV neurons persists in hypothyroid and mutant mice.

Discussion

In the present study, we extend our previous observations showing that a cell-autonomous action of

TH/TRα1 in GABAergic neurons is required for the maturation of neocortical PV neurons in mice. We

identified a time window that spans from birth to the end of the third postnatal week, during which

the action of THs is critical for the proper maturation of PV neurons. The analysis of gene expression

also pinpoints a set of genes that are directly regulated by TH/TRα1 in GABAergic neurons. Some of

these genes encode ion channels and PNN components, reflecting the involvement of THs in

essential steps of PV neuron maturation.

Using genetically modified mouse models expressing TRα1^{L400R}, which exerts dominant negative

activity over TRα1, we have gathered evidence suggesting that the time window during which THs

promote PV neuron maturation is in the early postnatal weeks, notably during the second postnatal

week. First, blocking TH/TRα1 signaling in GABAergic neurons after PND7 is sufficient to alter the

expression of parvalbumin, as evidenced at PND14. On the contrary, blocking TH/TRα1 signaling in

GABAergic neurons after PND21 does not significantly alter PV neuron maturation, as assessed in

adulthood. PV neurons also show near-normal maturation when the blockade of TH/TRα1 signaling is

restricted to PV neurons and begins only after the onset of the expression of *Pvalb*, the gene that

encodes parvalbumin (19). Furthermore, a previous whole cortex RNA-seq analysis showed that gene

expression changes rapidly in wild-type mice during the second postnatal week and that this

evolution is stopped when TH/TRα1 signaling is blocked in GABAergic neurons (16). Finally, our novel

transcriptome analysis, which was carried out with sorted GABAergic neuron nuclei, indicates that

the postnatal maturation process of PV neurons is blunted by TRα1^{L400R} expression. In particular,

blocking TH/TRα1 signaling appears to compromise the expression of a number of genes involved in

a well-documented transition, during which the repertoire of ion channels is extensively modified

(Okaty et al., 2009). It is remarkable that the second postnatal week, during which THs seem to play a

crucial role in PV neuron maturation, coincides with a peak in circulating levels of THs (22). This

strongly suggests that THs represent a temporal cue to synchronize the postnatal maturation of PV

neurons.

The striking influence exerted by THs on cortical PV neuron maturation in the first postnatal weeks

does not exclude the possibility that THs also play a role in these neurons earlier during

development, or after weaning. First, we cannot fully rule out that TRα1 has an earlier function in PV

neurons, as recent observations suggest that it could act prenatally to promote the tangential

migration of GABAergic neuron progenitors (23). However, there is no indication that this effect is

exerted in a cell-autonomous manner. Second, the hypothesis that THs act mainly during the second

postnatal week is at odds with observations that were previously reported in mice expressing

TRα1^{R384C} in all tissues, for which a complete recovery in PV neuron density was observed in adults

(13). By contrast, in the present study we have observed little recovery in PV neuron maturation in

adult mice expressing TRα1^{E395fs401} in GABAergic neurons. This discrepancy is probably explained by

the residual capacity of TRα1^{R384C} to bind T3 and activate transcription (24), which is not the case for

TRα1^{E395fs401} (16). Therefore, the late onset of TRβ1 expression in PV neurons (25) does not appear to

be sufficient to counteract the influence of the dominant negative TRα1 mutated receptor.

The mode of TH action in PV neurons during the second postnatal week is clarified by RNA-seq data.

In the present study, we first found that PV neurons were much more sensitive to variations in TH

signaling at PND14 than at the adult stage, where the same method only showed minor changes in

gene expression (19). Another finding of the present study was that one prominent outcome of the

late maturation process of PV neurons, the elaboration of PNNs, appears to strongly rely on TH

signaling. Indeed, genes that were consistently deregulated when TH/TRα1 signaling was blocked,

encode extracellular matrix proteins, specific components of PNNs, or proteins interacting with PNN

glycosaminoglycans. We used rapid recovery of gene expression after TH treatment in hypothyroid
mice, as an indication of direct transcriptional control of gene expression by TRα1. This allowed us to
identify a small subset of genes whose deregulation is more likely to be the initial set of events
preventing PV neuron maturation. Among these are genes encoding transcription factors or cofactors
(*Dbp, Hr, Klf9*), that were already reported to be regulated by TRα1 in other contexts (21) and that
could act as upstream regulators of other deregulated genes. However, the presence in this subset of
putative TRα1 target genes of genes encoding components of PNNs suggests a more direct link
between TH and PNN content. Thus, it appears that liganded TRα1 might notably activate the
transcription of genes encoding collagens (*Col24a1* and *Col9a2* genes), ephrin A4 (*Epha4*), laminin
alpha 3 subunit (*Lama3*), semaphorin 7A (*Sema7a*) and hyaluronan and proteoglycan link protein 4
(*Hapln4* gene). This last protein notably represents an essential component of PNNs (26-28).

The novel data presented in this paper provide strong support for the hypothesis that TH is a major
temporal signal for the opening and closing of the critical period of heightened plasticity, as
previously suggested (29). The critical period occurs at slightly different times in different neocortical
areas, as does the onset of TH signaling (30). It is characterized by a series of cellular events that play
a crucial role in the elaboration of functional neuronal circuits (5). The beginning of parvalbumin
synthesis marks the opening of the critical period, while the elaboration of PNNs around PV neurons
stabilizes synapses and marks its closure. Thus, our findings suggest that both the opening and the
closure of the critical period are directly stimulated by liganded TRα1. Therefore, THs appear to act in
the postnatal neocortex as it does during amphibian metamorphosis, as an external cue that
represents an important temporal signal. During brain development, the influence of THs is also
found in glial cells, as TH signaling is particularly required for timely differentiation of
oligodendrocytes and myelin formation (12, 31). Of note, oligodendrocyte maturation has also been
shown to play a role in PV neuron maturation and in the closure of the critical period (32).

Patients with early TH deficiency caused by congenital hypothyroidism (33) or carrying TRα1
mutations (11) are currently treated with TH replacement therapy. Treatment can have a positive
influence on cognitive development only if it is initiated soon after birth. This is an indication that the
TH action on the maturation of the human neocortex is also restricted to an early temporal window.
An alteration in the timing in the critical period is also suspected to cause autism spectrum disorders
(8). In that regard, it is noticeable that some of the rare patients in whom *THRA* mutations have been
discovered were first diagnosed for autism spectrum disorders (34, 35). Within the framework of our
hypothesis, irreversible cognitive and neurological disorders that are associated with TH related

diseases are related to ASD symptoms and might also be regarded as consequences of a disruption of
the critical period of heightened plasticity.

Study limitations

The complete characterization of TRα1 target genes in PV neurons would require the definition of
chromatin occupancy by the receptor, as it informs of its presence on regulatory sequences. This is
not currently feasible with the small number of cells that are collected when sorting PV neurons from
mouse neocortex. Also, the histological methods that we have used only indirectly inform of the
timing of the critical period of heightened plasticity, which can be defined in a number of different
ways (36, 37).

Methods

Mouse models and treatments

Experiments involving the use of live animals for the current project were approved by local ethics
committees (C2EA015 and C2EA017) and subsequently authorized by the French Ministry of
Research (Projects #33279-2021082516194165 and #25496-2020032811174506). Mice were bred
and maintained at Plateau de Biologie Expérimentale de la Souris (SFR BioSciences Gerland - Lyon
Sud, France).

The *Thra^{AMI}* allele allows the expression of the dominant-negative TRα1^{L400R} after the Cre/loxP-
mediated deletion of a cassette with polyadenylation signals (38). *Thra^{Slox}* is identical, except for the
presence of a C-terminus frameshift mutation which leads to the expression of the dominant-
negative TRα1^{E395fs401X}. The *ROSA-tdTomato^{lox}* reporter transgene (also known as Ai9, MGI Cat#
4436851, RRID: MGI: 4436851) drives the Cre-dependent expression of a red fluorescent protein,
mainly localized in cytoplasm (39). The *ROSA-GSL10gfp^{lox}* transgene encodes the GS-EGFP-L10a
protein, which is a green fluorescent protein fused to the N-terminus part of the large subunit
ribosomal protein L10a, mainly localized in cell nuclei (19). All transgenes were in C57Bl6/J genetic
background. *Gad2-Cre* and *Gad2-CreER^{T2}* are knock-in alleles in which the Cre recombinase reading
frame is inserted into the *Gad2* gene in order to restrict the expression of the recombinase in
GABAergic neurons (40) . Recombination was induced in mice carrying the *Gad2-CreER^{T2}* transgene
by intraperitoneal injection of tamoxifen (T5648, Sigma, dissolved in sterile sunflower oil (10 mg/ml).
PND7 mice received a single 50 µL subcutaneous injection whereas 3-week-old mice received 5 daily
intraperitoneal injections (daily dose: 7.5 µL/g).

Gestating mice were fed for 2 weeks with iodine deficient food supplemented with 0.15% propyl-thio

uracyl (Envigo ref TD.95125) to cause deep hypothyroidism. TH levels were restored by a single

intraperitoneal injection in half of the pups (10 µg T4 + 1 µg T3 dissolved in 50 µL of phosphate buffer

saline, all chemicals from Sigma Aldrich France). The other PTU treated animals were injected with 50

uL of phosphate buffer saline.

RNA-seq analysis from sorted nuclei

Nuclei were isolated from individual cortex frozen in liquid nitrogen and nuclear RNA was extracted

as previously described (19). Nuclear RNA was extracted from sorted nuclei (RNeasy Micro Kit,

Qiagen ref 74004) and quantified using Tapestation4150 (Agilent). 1 ng RNA was reverse-transcribed

using the SMART-Seq v4 Low Input RNA kit (Takara). cDNA was quantified and qualified using Qubit

(Invitrogen) and Tapestation4150 (Agilent). Libraries were then prepared from 1 ng cDNA using the

Nextera XT DNA Library Kit (Illumina). Libraries were sequenced (> 2.10^7 reads/library) on a

Nextseq500 DNA sequencer (Illumina). Reads were aligned on the mouse genome (mm10 GRCm38

release) with Bowtie2 (Galaxy Version 2.4.2 + galaxy 0) (41). Count table was prepared using htseq-

count (Galaxy Version 0.9.1 + galaxy 1, mode union, feature type: gene) (42). Differential gene

expression analysis was performed with DEseq2 (Galaxy Version 2.1.8.3) (43) using the following

thresholds: False Discovery Rate < 0.05; p-adjusted value < 0.05; maximum adjusted basecount > 30;

fold-change > 2.0. Hierarchical clustering was performed using Euclidian distance and complete

distance with Clustvis (44). Gene Ontology analysis was performed with Gorilla (https://cbl-

gorilla.cs.technion.ac.il/).

Brain slice collection

Each mouse was given a lethal intraperitoneal injection (6 mL/kg) of a mixture of ketamine (33

mg/mL) and xylazine (6.7 mg/mL). The thorax was opened and each mouse was perfused with 3.7%

paraformaldehyde in 0.1 M phosphate buffer at room temperature. Each brain was dissected out,

immersed in fixative at 4°C for 3 h and then in phosphate buffered saline (PBS) at 4°C until sectioning.

Coronal sections (50 µm) were cut with the aid of a vibrating microtome (Integraslice 7550 SPDS,

Campden Instruments, Loughborough, UK), in PBS at room temperature. Brain sections were stored

at -20°C in cryoprotectant (30% ethylene glycol and 20% glycerol in 10 mM low-salt PBS).

Brain slice histology

Immunohistochemistry was performed on free-floating brain sections. For PV labeling, a mouse anti-

PV primary antibody (PARV19, Sigma P3088, 1:2000), and a secondary antibody made in donkey

(anti-mouse DyLight 633, ThermoFisher Scientific, 1:1000) were used. For the concurrent PNN

labeling, a biotinylated Wisteria floribunda lectin (WFL/WFA, Vector Laboratories B-1355, 20 µg/mL),

and Streptavidin coupled to DyLight 488 (Vector Laboratories, SA-5488, 1:1000) were used. Non-

specific binding sites were blocked by incubating sections for 1 h in PBS with 1% BSA and 0.2% Triton

X-100. Brain sections were incubated overnight at 4°C with PV antibody and biotinylated WFA, which

were diluted in PBS with 1% BSA, 0.2% Triton X-100 and 1% dimethyl sulfoxide. Sections were

washed in PBS and further incubated for 15 min at room temperature with DAPI (4',6-diamidino-2-

phenylindole, 1:5000, Sigma). Incubation with the secondary antibody lasted for 3 h at room

temperature. Sections were mounted in Fluoroshield™ (Sigma), coverslipped and imaged using an

inverted confocal microscope (Zeiss LSM 780 and Zeiss LSM 800).

Image analysis

Image analysis was performed using ImageJ. Final images resulted from the z projection of 3 optical

sections (Maximum intensity tool) that were 2 µm apart (the resulting images thus reflected the

fluorescence collected over a thickness of 4 µm). The densities of Tomato+, PV+, WFA+ cells, and

their co-localizations were measured in the whole area of one image taken in the somatosensory

cortex area. Three to four images in both hemispheres were analyzed for each mouse. Except for Fig.

2a-d, for which the counting was performed manually, the Cellpose wrapper for Fiji (45) was used to

count Tomato+, PV+ and WFA+ cells. Numbers of mono-, double- or triple-labelled cells per slice

were measured automatically. The code of the original macro may be found here:

https://github.com/jbrocardplatim/MacroPV.

Quantitative data were analyzed using two-tailed unpaired T-test. Data are expressed as mean ±

SEM. Data analyses were performed with Excel 2016 and GraphPad Prism software (v.8.). The level of

significance was set at $p < 0.05$.

Figure legends:

Fig. 1 | Persistent defect in PV neuron maturation in mice expressing TRα1^{E395fs401X}. Histological

analysis of *Thra$^{+/+}$ ROSA-tdTomato$^{lox/+}$ Gad2-Cre* littermates (controls) and *Thra$^{Slox/+}$ ROSA-*

tdTomato$^{lox/+}$ Gad2-Cre mice (mutants) that express TRα1^{E395fs401X} in the GABAergic neurons and their

progenitors from embryonic day 12.5 to PND21 (**a-d**, controls: n = 6, mutants: n = 6) and to adult

stage (**e-h**, controls: n = 6, mutants: n = 5). **a, e** Immunohistochemistry of parvalbumin and WFA-

labeled PNNs in PV neurons of somatosensory cortex at striatum level. **b, f** The density of PV+ cells was lower in mutants at both ages. **c, g** The density of WFA+ cells was lower in mutants at both ages. **d, h** The proportion of PV neurons surrounded by PNNs was lower in mutants at both ages. Data are presented as mean ± SEM and analyzed using an unpaired two-tailed T-test (*$p < 0.05$, **$p < 0.01$, ***$p < 0.001$).

Fig. 2 | A time window for TH action on PV neuron maturation. Histological analysis of *Thra$^{+/+}$ ROSA-tdTomato$^{lox/+}$ Gad2-CreER^{T2}* or *Gad2-CreER^{T2}* negative littermates (controls) and *Thra$^{AMI/+}$ ROSA-tdTomato$^{lox/+}$ Gad2-CreER^{T2}* mice (mutants) that express TRα1^{L400R} in the GABAergic neurons from PND7 to PND14 (**a-d**, controls: n = 8, mutants: n = 6) and from PND21 to adult (**e-h**, controls: n = 11, mutants: n = 7). **a, e** Immunohistochemistry of parvalbumin and WFA-labeled PNNs in PV neurons of somatosensory cortex at striatum level. **b, f** The density of PV+ cells was lower in mutants in former but not in latter. **c, g** The density of WFA+ cells had no clear difference in both cases. **d, h** The proportion of PV neurons surrounded by PNNs had no clear difference in both cases. Data are presented as mean ± SEM and analyzed using unpaired two-tailed T-test (*$p < 0.05$, **$p < 0.01$, ***$p < 0.001$).

Fig. 3 | RNA-seq analysis of gene expression in sorted GABAergic neuron nuclei. Nuclei were sorted at PND14 from the neocortex of *Thra$^{+/+}$ ROSA-GSL10gfp^{lox} Gad2-Cre* mice (Ctrl), *Thra$^{AMI/+}$ ROSA-GSL10gfp^{lox} Gad2-Cre* mice expressing TRα1^{L400R} in GABAergic neurons (Mut), hypothyroid *Thra$^{+/+}$ ROSA-GSL10gfp^{lox} Gad2-Cre* mice (Hypoth.), or hypothyroid *Thra$^{+/+}$ ROSA-GSL10gfp^{lox} Gad2-Cre* mice that received a single TH injection at PND13 (+TH). **a** Venn diagram of genes that were either downregulated by TRα1^{L400R}, downregulated by hypothyroidism or upregulated after TH treatment. **b** Venn diagram of genes that were either upregulated by TRα1^{L400R}, upregulated by hypothyroidism or downregulated after TH treatment (thresholds: Fold change > 2 or < 0.5, maximal adjusted base counts > 30, adjusted p-value > 0.05). **c** Heatmap showing the genes that were downregulated by both TRα1^{L400R} and hypothyroidism. Note the variable response to TH. **d** Heatmap showing the genes that were upregulated by both TRα1^{L400R} and hypothyroidism. **e** Heatmap of markers of PV neuron maturation. The marks on the right side are for genes whose expression normally increases (red) or decreases (green) between PND7 and PND21. The alterations of gene expression in mutant mice reveals a blockade of the PV neuron maturation program, caused either by TRα1^{L400R} expression or by hypothyroidism. A 24h TH treatment two weeks after birth is not sufficient to restore the expression of these marker genes.

Acknowledgements:

We acknowledge the contribution of SFR Biosciences (Université Claude Bernard Lyon 1, CNRS UAR3444, Inserm US8, ENS de Lyon): Nadine Aguilera and the Plateau de Biologie Expérimentale de la Souris (ANIRA-PBES) for mouse breeding; Sébastien Dussurgey from ANIRA-Cytométrie for nuclei sorting; ANIRA-AGC for mouse genotyping; LyMIC-PLATIM for imaging. We thank Benjamin Gillet and Sandrine Hughes of the deep sequencing facility (PSI IGFL, Lyon). We also thank Elodie Martel for carrying out histology experiments as part of her internships and Isabelle Dusart (Institut de Biologie Paris Seine, France) for helpful discussions. We acknowledge China Scholarship Council for a scholarship to Juan REN to do her PhD project in ENS de Lyon. Research fundings were from European Union's Horizon 2020 research and innovation program, under grant agreement No. 825753 (ERGO), Agence Nationale de la Recherche (Thyromut2 program; ANR-15-CE14-0011-01).

Declaration of interest

No conflict of interest

References

1. S. Hijazi, A. B. Smit, R. E. van Kesteren, Fast-spiking parvalbumin-positive interneurons in brain physiology and Alzheimer's disease. *Mol Psychiatry* **28**, 4954-4967 (2023).
2. J. A. del Rio, L. de Lecea, I. Ferrer, E. Soriano, The development of parvalbumin-immunoreactivity in the neocortex of the mouse. *Brain Res Dev Brain Res* **81**, 247-259 (1994).
3. Z. J. Huang *et al.*, BDNF regulates the maturation of inhibition and the critical period of plasticity in mouse visual cortex. *Cell* **98**, 739-755 (1999).
4. B. W. Okaty, M. N. Miller, K. Sugino, C. M. Hempel, S. B. Nelson, Transcriptional and electrophysiological maturation of neocortical fast-spiking GABAergic interneurons. *J Neurosci* **29**, 7040-7052 (2009).
5. T. K. Hensch, Critical period plasticity in local cortical circuits. *Nat Rev Neurosci* **6**, 877-888 (2005).
6. A. E. Takesian, T. K. Hensch, Balancing plasticity/stability across brain development. *Prog Brain Res* **207**, 3-34 (2013).
7. J. C. Wingert, B. A. Sorg, Impact of Perineuronal Nets on Electrophysiology of Parvalbumin Interneurons, Principal Neurons, and Brain Oscillations: A Review. *Frontiers in synaptic neuroscience* **13**, 673210 (2021).
8. V. S. Sohal, J. L. R. Rubenstein, Excitation-inhibition balance as a framework for investigating mechanisms in neuropsychiatric disorders. *Mol Psychiatry* **24**, 1248-1257 (2019).
9. F. Flamant, K. Gauthier, Thyroid hormone receptors: the challenge of elucidating isotype-specific functions and cell-specific response. *Biochim Biophys Acta* **1830**, 3900-3907 (2013).
10. P. Berbel, D. Navarro, G. C. Roman, An evo-devo approach to thyroid hormones in cerebral and cerebellar cortical development: etiological implications for autism. *Frontiers in endocrinology* **5**, 146 (2014).

11. A. L. M. van Gucht *et al.*, Resistance to Thyroid Hormone due to Heterozygous Mutations in
Thyroid Hormone Receptor Alpha. *Current topics in developmental biology* **125**, 337-355
(2017).
12. F. Picou, T. Fauquier, F. Chatonnet, S. Richard, F. Flamant, Deciphering direct and indirect
influence of thyroid hormone with mouse genetics. *Mol Endocrinol* **28**, 429-441 (2014).
13. K. Wallis *et al.*, Locomotor deficiencies and aberrant development of subtype-specific
GABAergic interneurons caused by an unliganded thyroid hormone receptor alpha1. *J*
Neurosci **28**, 1904-1915 (2008).
14. K. Uchida *et al.*, Thyroid hormone insufficiency alters the expression of psychiatric disorder-
related molecules in the hypothyroid mouse brain during the early postnatal period.
Scientific reports **11**, 6723 (2021).
15. S. Mayerl, H. Heuer, Thyroid hormone transporter Mct8/Oatp1c1 deficiency compromises
proper oligodendrocyte maturation in the mouse CNS. *Neurobiol Dis* **184**, 106195 (2023).
16. S. Richard *et al.*, A Pivotal Genetic Program Controlled by Thyroid Hormone during the
Maturation of GABAergic Neurons. *iScience* **23**, 100899 (2020).
17. M. E. Gilbert *et al.*, Thyroid hormone insufficiency during brain development reduces
parvalbumin immunoreactivity and inhibitory function in the hippocampus.
Endocrinology **148**, 92-102 (2007).
18. S. Markossian *et al.*, CRISPR/Cas9 Editing of the Mouse Thra Gene Produces Models with
Variable Resistance to Thyroid Hormone. *Thyroid* **28**, 139-150 (2018).
19. J. Ren *et al.*, Thyroid hormones maintain parvalbumin neuron functions in mouse neocortex.
bioRxiv 10.1101/2024.07.16.603713, 2024.2007.2016.603713 (2024).
20. B. Tasic *et al.*, Adult mouse cortical cell taxonomy revealed by single cell transcriptomics. *Nat*
Neurosci **19**, 335-346 (2016).
21. Y. Zekri, R. Guyot, F. Flamant, An Atlas of Thyroid Hormone Receptors' Target Genes in
Mouse Tissues. *International journal of molecular sciences* **23** (2022).
22. S. Friedrichsen *et al.*, Regulation of iodothyronine deiodinases in the Pax8-/- mouse model of
congenital hypothyroidism. *Endocrinology* **144**, 777-784 (2003).
23. E. da Cunha Menezes *et al.*, Effects of gestational hypothyroidism on mouse brain
development: Gabaergic systems and oxidative stress. *Dev Biol* **515**, 112-120 (2024).
24. A. Tinnikov *et al.*, Retardation of post-natal development caused by a negatively acting
thyroid hormone receptor alpha1. *Embo J* **21**, 5079-5087 (2002).
25. D. R. Hochbaum *et al.*, Thyroid hormone remodels cortex to coordinate body-wide
metabolism and exploration. *Cell* **187**, 5679-5697 e5623 (2024).
26. K. Nojima, H. Miyazaki, T. Hori, L. Vargova, T. Oohashi, Assessment of Possible Contributions
of Hyaluronan and Proteoglycan Binding Link Protein 4 to Differential Perineuronal Net
Formation at the Calyx of Held. *Front Cell Dev Biol* **9**, 730550 (2021).
27. M. Edamatsu *et al.*, Hapln4/Bral2 is a selective regulator for formation and transmission of
GABAergic synapses between Purkinje and deep cerebellar nuclei neurons. *J Neurochem* **147**,
748-763 (2018).
28. J. Popelar *et al.*, The absence of brain-specific link protein Bral2 in perineuronal nets hampers
auditory temporal resolution and neural adaptation in mice. *Physiological research /*
Academia Scientiarum Bohemoslovaca **66**, 867-880 (2017).
29. G. Batista, T. K. Hensch, Critical Period Regulation by Thyroid Hormones: Potential
Mechanisms and Sex-Specific Aspects. *Front Mol Neurosci* **12**, 77 (2019).
30. L. Quignodon *et al.*, Thyroid hormone signaling is highly heterogeneous during pre- and
postnatal brain development. *J Mol Endocrinol* **33**, 467-476 (2004).
31. B. Durand, M. Raff, A cell-intrinsic timer that operates during oligodendrocyte development.
Bioessays **22**, 64-71 (2000).
32. B. T. Kalish *et al.*, Single-nucleus RNA sequencing of mouse auditory cortex reveals critical
period triggers and brakes. *Proc Natl Acad Sci U S A* **117**, 11744-11752 (2020).

33.   G. Van Vliet, J. Deladoey, Diagnosis, treatment and outcome of congenital hypothyroidism.
Endocr Dev **26**, 50-59 (2014).
34.   R. K. Yuen *et al.*, Whole-genome sequencing of quartet families with autism spectrum
disorder. *Nat Med* **21**, 185-191 (2015).
35.   M. K. Kalikiri, M. P. Mamidala, A. N. Rao, V. Rajesh, Analysis and functional characterization
of sequence variations in ligand binding domain of thyroid hormone receptors in autism
spectrum disorder (ASD) patients. *Autism research : official journal of the International*
Society for Autism Research 10.1002/aur.1838 (2017).
36.   K. B. Quast *et al.*, Rapid synaptic and gamma rhythm signature of mouse critical period
plasticity. *Proc Natl Acad Sci U S A* **120**, e2123182120 (2023).
37.   R. K. Reh *et al.*, Critical period regulation across multiple timescales. *Proc Natl Acad Sci U S A*
117, 23242-23251 (2020).
38.   L. Quignodon, S. Vincent, H. Winter, J. Samarut, F. Flamant, A point mutation in the
activation function 2 domain of thyroid hormone receptor alpha1 expressed after CRE-
mediated recombination partially recapitulates hypothyroidism. *Mol Endocrinol* **21**, 2350-
2360 (2007).
39.   L. Madisen *et al.*, A robust and high-throughput Cre reporting and characterization system
for the whole mouse brain. *Nat Neurosci* **13**, 133-140 (2010).
40.   H. Taniguchi *et al.*, A resource of Cre driver lines for genetic targeting of GABAergic neurons
in cerebral cortex. *Neuron* **71**, 995-1013 (2011).
41.   B. Langmead, C. Trapnell, M. Pop, S. L. Salzberg, Ultrafast and memory-efficient alignment of
short DNA sequences to the human genome. *Genome Biol* **10**, R25 (2009).
42.   S. Anders, P. T. Pyl, W. Huber, HTSeq--a Python framework to work with high-throughput
sequencing data. *Bioinformatics* **31**, 166-169 (2015).
43.   M. I. Love, W. Huber, S. Anders, Moderated estimation of fold change and dispersion for
RNA-seq data with DESeq2. *Genome Biol* **15**, 550 (2014).
44.   T. Metsalu, J. Vilo, ClustVis: a web tool for visualizing clustering of multivariate data using
Principal Component Analysis and heatmap. *Nucleic Acids Res* **43**, W566-570 (2015).
45.   C. Stringer, T. Wang, M. Michaelos, M. Pachitariu, Cellpose: a generalist algorithm for cellular
segmentation. *Nat Methods* **18**, 100-106 (2021).