Thyroid hormones act as a timer for the postnatal maturation of parvalbumin
neurons in the neocortex

Juan Ren[1,3], Suzy Markossian[1], Romain Guyot[1], Denise Aubert[1], Jacques Brocard[2], Jiemin Wong[3],
Frédéric Flamant[1,*], Sabine Richard[1]

[1] Ecole Normale Supérieure de Lyon, INRAE, CNRS, Institut de Génomique Fonctionnelle de Lyon,
69364 Lyon, France

[2] Université Claude Bernard Lyon 1, CNRS UAR3444, Inserm US8, ENS de Lyon, SFR Biosciences, Lyon
69007, France

[3] Shanghai Key Laboratory of Regulatory Biology, Institute of Biomedical Sciences and School of Life
Sciences, East China Normal University, 200241, Shanghai, China

[*] Corresponding author. Frederic.flamant@ens-lyon.fr Tel: 00 (33) 4 26 73 13 32

Abbreviated title: Thyroid hormone action in postnatal cortex

Declaration of conflict of interest: none.

Abstract

Thyroid hormones (THs) are required for proper maturation of parvalbumin-expressing GABAergic

interneurons in the mouse neocortex, which function is essential to maintain a proper balance

between neuronal excitation and inhibition. However, the timeline of this TH action has yet to be

elucidated. Here, we used several mouse models expressing dominant negative mutations of the TH

nuclear receptor α1 (TRα1), to better define the time window during which THs promote the

maturation of parvalbumin neurons. The action of THs during the first three postnatal weeks appears

to be necessary to initiate the elaboration of a specialized extracellular matrix called the

perineuronal net. Transcriptome analysis identified a small set of putative target genes for TRα1,

which are at the origin of the postnatal remodeling of the repertoire of ions channels and the

elaboration of perineuronal nets. These data suggest that THs act as a timer to define the temporal

boundaries of the critical period of heightened cortical plasticity, which plays a fundamental role in

the development of neuronal circuits.

Significance statement

Thyroid hormones exert a major influence on late neurodevelopment, acting directly on gene
expression by binding to nuclear receptors present in all cell types. This study reports the discovery
of a direct link between the transcriptional regulation exerted by the thyroid hormones in GABAergic
neurons and the timing of the critical period of heightened plasticity. This provides a general
framework to understand the pathology associated with altered thyroid hormone signaling.

Keywords

Thyroid hormones / TRα1 / PV neurons / GABAergic neurons / neocortex

Introduction

Parvalbumin-expressing neurons (PV neurons) are GABAergic inhibitory interneurons that play a
central role in the organization of neuronal circuits in the neocortex (Hijazi, Smit et al., 2023). They
are fast-spiking and synchronize network oscillations, which are thought to play an important role in
information processing and memory formation. In mice, PV neurons begin to express parvalbumin
during the first postnatal weeks (del Rio, de Lecea et al., 1994, Huang, Kirkwood et al., 1999). During
this time, the electrical activity of these neurons progressively increases (Okaty, Miller et al., 2009)
and synaptogenesis is very active. This opens the so-called critical period of heightened synaptic
plasticity (Hensch, 2005) during which sensory input and experience shape neuronal networks in the
neocortex. The critical period ends when circuit rewiring becomes actively dampened (Takesian &
Hensch, 2013). Notably, the elaboration of the perineuronal net (PNN) around PV neurons, a densely
packed extracellular matrix enriched in proteoglycans, stabilizes synapses and restricts plasticity
(Wingert & Sorg, 2021). Defects in PV neuron maturation during the critical period cause a persistent
imbalance between neuronal excitation and inhibition, which is suspected to be at the origin of
several pathological conditions, including attention deficit hyperactivity disorder, autism, and
epilepsy (Sohal & Rubenstein, 2019).

THs (including T3 or 3,3',5-triiodo-L-thyronine, and the less active precursor T4 or thyroxine) exert a
broad influence on brain development, and they are critically required for proper maturation of PV
neurons (Bernal, Morte et al., 2022, Ren & Flamant, 2023, Richard, Ren et al., 2023). THs directly act
on gene transcription by binding to the TRα1 and TRβ1/2 nuclear receptors, which are encoded by
the *Thra* and *Thrb* genes in mice. In both humans and mice, TRα1 mutations have dramatic and
irreversible consequences on neurodevelopment, and these consequences are similar to those of TH

deficiency, with most human presenting with mild to severe intellectual disability (Berbel, Navarro et
al., 2014, van Gucht, Moran et al., 2017).

In the postnatal mouse neocortex, PV neurons display a high sensitivity to TRα1 mutation (Wallis,
Sjogren et al., 2008) and TH deficiency (Mayerl & Heuer, 2023, Uchida, Hasuoka et al., 2021). We
have previously found that expressing TRα1^{L400R}, a dominant-negative TRα1 receptor, only in the
GABAergic lineage from an early neurodevelopmental stage, was sufficient to dramatically inhibit the
expression of parvalbumin and the elaboration of PNNs (Richard, Guyot et al., 2020). By contrast, we
have recently observed that expressing TRα1^{L400R} specifically in PV neurons had no visible effects on
their maturation and had only moderate consequences on their function (Okaty et al., 2009). In the
latter study, the specific blockade of TH signaling in PV neurons was driven by the expression of the
Pvalb gene, which starts very gradually and with considerable temporal variation between PV
neurons, ranging from a few days up to several weeks after birth (de Lecea, del Rio et al., 1995).
Taken together, these results suggest that TH signaling stimulates the maturation of PV neurons
during a specific time window and then becomes less important to maintain their function.
Accordingly, the appearance of hypothyroidism in adult stages does not prevent the expression of
parvalbumin (Gilbert, Sui et al., 2007, Uchida et al., 2021). However, the boundaries of the time
window during which THs act on PV neuron maturation remain elusive. Moreover, whether blocking
TH signaling during development only delays or permanently inhibits PV neuron maturation remains
an open question.

To gain more insight into the timeline of TH action on PV neuron maturation, we generated novel
mouse models to restrict the expression of dominant-negative TRα1 receptors in GABAergic neurons
during specific time windows. Histological observations of parvalbumin and PNNs in the
somatosensory cortex allowed us to pinpoint a key time window for TH action in PV neurons, around
the second postnatal week. Moreover, we evidenced a lifelong persistence of defects in PV neuron
maturation when TH signaling was blocked in GABAergic neurons from early stages of development.
A genome-wide analysis of GABAergic neuron transcriptome in the whole neocortex two weeks after
birth also allowed us to identify a set of genes whose regulation is expected to be involved in TH-
mediated maturation of PV neurons. Several of these genes are notably known to be linked to the
closure of the critical period. As a whole, our results show that THs, via their action on PV neurons,
play a key role in cortical circuit maturation in the mouse brain.

Results

$TR\alpha1^{E395fs401X}$ expression in GABAergic neurons from an early developmental stage permanently

inhibits PV neuron maturation

In our previous work, we used Cre/loxP recombination to restrict the expression of the dominant-

negative $TR\alpha1^{L400R}$ mutant receptor in the GABAergic lineage from embryonic day 12.5 (Richard et al.,

2020). As most of the mice had frequent epileptic seizures and died before postnatal day 21 (PND21),

we could not address a possible recovery of PV neuron maturation during adulthood. We thus used

here a similar model, based on the $Thra^{Slox}$ allele, encoding $TR\alpha1^{E395fs401X}$, a receptor with moderate

dominant-negative activity (Markossian, Guyot et al., 2018). The floxed $ROSA$-$tdTomato^{lox}$ reporter

transgene was also introduced to trace the cells belonging to the GABAergic lineage. One third of

$Thra^{Slox/+}$ $ROSA$-$tdTomato^{lox/+}$ $Gad2$-Cre mice survived to adulthood. We therefore compared these

mutant mice and $Thra^{+/+}$ $ROSA$-$tdTomato^{lox/+}$ $Gad2$-Cre littermates (controls), to histologically study

whether TRα1 mutation only delays or permanently inhibits PV neuron maturation.

At PND21, while the red fluorescence did not indicate any cell loss in the GABAergic lineage (Suppl.

Fig. 1), the density of parvalbumin-positive (PV+) cells was drastically lower in mutants than in

controls (Fig. 1a, b). PNNs were labeled using Wisteria Floribunda Agglutinin (WFA), which binds N-

acetyl-galactosamine-β1 residues of PNN glycoproteins. The density of WFA-positive (WFA+) cells

was also significantly lower in mutants (Fig. 1a, c). Interestingly, the two phenotypes were

accompanied by a clearly lower ratio of PV+ cells that were surrounded by PNNs in mutants,

compared to controls (Fig. 1a, d). At adult stage, the same analysis showed signs of some recovery of

PV neuron density in mutants (Fig. 1e, f), while the PNN defect persisted (Fig. 1e, g and h). Overall,

$TR\alpha1^{E395fs401X}$ appears to have a similar negative influence to $TR\alpha1^{L400R}$ and permanently impairs the

maturation of PV neurons.

Postnatal expression of $TR\alpha1^{L400R}$ in GABAergic neurons is sufficient to inhibit PV neuron

maturation

We have recently observed that the postnatal expression of $TR\alpha1^{L400R}$ in PV neurons has little effect

on their maturation (submitted data). Although adult $Thra^{AMI/+}$ $ROSA$-$tdTomato^{lox/+}$ $Pvalb$-Cre mice

displayed minor histological, electrophysiological and behavioral defects, they did not spontaneously

present epileptic seizures or lethality (Ren, Markossian et al., 2024). This suggests that TH signaling

intensively stimulates the maturation of PV neurons during a specific time window, and then

becomes secondary to maintain their function. To better evaluate the temporal restriction of TH

signaling for PV neurons maturation, we used $Thra^{AMI/+}$ $ROSA$-$tdTomato^{lox/+}$ $Gad2$-$CreER^{T2}$ mice

(mutants) and *Thra*$^{+/+}$ *ROSA-tdTomato*$^{lox/+}$ *Gad2-CreER*^{T2} or *Gad2-CreER*^{T2} negative mice (controls),

where Cre/loxP recombination can be induced at a chosen time by tamoxifen treatment.

A first batch of these mice were treated with tamoxifen at PND7 and histological analysis was

performed at PND14. This analysis revealed that expressing TRα1^{L400R} during the second postnatal

week was sufficient to compromise the onset of parvalbumin expression (Fig. 2a, b). But the

amplitude of the defect in PV expression in mutant mice was not as high as that in our previous work,

where TH signaling was blocked from embryonic day 12.5 to PND14 in *Thra*$^{AMI/+}$ *ROSA-tdTomato*$^{lox/+}$

Gad2-Cre mice (Richard et al., 2020). Besides, we did not observe any clear change in either the

density of WFA+ cells (Fig. 2a, c) or the ratio of PV+ cells that were surrounded by PNNs (Fig. 2a, d).

This may be due to the fact that the elaboration of PNNs mainly takes place after PND14, so that the

density of WFA+ cells was low, with high interindividual variability. By contrast, a tamoxifen

treatment performed during the fourth postnatal week (from PND21 to PND25), did not cause any

significant change in either PV+ cell density or WFA+ cell density at adult stage (Fig. 2e-h). Therefore,

THs exert their influence during a postnatal time window, which includes the second postnatal week

and precedes weaning, since weaning usually occurs after the end of the third postnatal week in

mice.

Consequences of TRα1^{L400R} *expression on gene expression in neocortical GABAergic neurons two weeks after birth*

We set out to analyze the influence of TRα1^{L400R} expression on the nuclear transcriptome of

GABAergic neurons at PND14, a developmental stage which appears to be quite central in the

timeline of TH action in PV neurons. To do this, we used the *ROSA-GSL10gfp*^{lox} reporter transgene,

which labels cell nuclei, to extract nuclear RNA from neocortical GABAergic neurons, in which PV

neurons represent around 50% of the population (Tasic, Menon et al., 2016). RNA-seq analysis was

performed for *Thra*$^{AMI/+}$ *ROSA-GSL10gfp*$^{lox/+}$ *Gad2-Cre* mice and *Thra*$^{+/+}$ *ROSA-GSL10gfp*$^{lox/+}$ *Gad2-Cre*

control littermates at PND14 Differential analysis (Deseq2 adjusted p-value > 0.05; fold-change > 2.0;

basemean > 30) identified 275 down-regulated genes and 265 up-regulated genes in mutant mice

(Suppl. Table 1). This large gene set contains TRα1 target genes whose transcription is normally up-

regulated by liganded TRα1. Notably, among these were genes (*Hr, Klf9, Dbp, Shh, Gabrd*) that are

known to be TRα1 target genes in other cell types (Zekri, Guyot et al., 2022). However, many of the

observed changes in mRNA level probably represent indirect consequences of the long-term

influence of TRα1^{L400R} expression on the cell physiology.

To distinguish direct TRα1 target genes from other genes that are downstream in the cascade, we

carried out a series of additional studies. First, we looked for genes which are down-regulated by

hypothyroidism and up-regulated by a short TH treatment at PND14. To do this, we used a

propylthiouracyl (PTU) treatment to induce hypothyroidism in gestating females and their offspring.

Half of the hypothyroid pups with a *Thra$^{+/+}$ROSA-GSL10gfp$^{lox/+}$ Gad2-Cre* genotype then received THs

at PND14. The nuclear RNA transcriptome of GABAergic neurons in the neocortex from hypothyroid

pups, and from pups treated with THs for 24 hours, was analyzed at PND14 as above. In this setting

we expected a selective recovery of mRNA level for TRα1 target genes.

A total of 1306 genes were sensitive to either TRα1^{L400R} expression, hypothyroidism, or TH

stimulation (Fig. 3a, b and Suppl. Table 1). Among them, a large set of genes were only deregulated in

hypothyroid mice. This suggests an important influence of hypothyroidism on the microenvironment

(notably glial cells) of GABAergic neurons, with indirect consequences on gene expression in

GABAergic neurons. The genes whose expression in GABAergic neurons was sensitive to both

TRα1^{L400R} expression and hypothyroidism rather reflect the cell-autonomous influence of TH signaling

in these neurons. Importantly, among a total of 388 of such genes was the *Pvalb* gene, which

encodes parvalbumin. Recovery after TH treatment (TH response reached a two-fold threshold) was

achieved for only a small subset of these genes (61 genes up-regulated after TH treatment and 5

genes down-regulated after TH treatment). However, clustering analysis revealed a clear trend

toward recovery, which was more visible for genes up-regulated by THs (Fig. 3c) than for genes

down-regulated by THs (Fig. 3d). The five genes which were down-regulated after TH treatment

could reflect a poorly understood negative regulation of transcription by the liganded TRα1 (Fig. 3b).

In the end, only 61 genes behaved as *bona fide* TRα1 target genes, with a robust negative response

to both TRα1^{L400R} expression and hypothyroidism, and a clear induction after the short TH treatment

of hypothyroid mice (Fig. 3a).

Gene Ontology analysis indicated that many of the genes deregulated in hypothyroid and mutant

mice encode components of the extracellular matrix (GO:0031012; enrichment 4.4, p-value 3.10^{-13}).

This includes genes encoding collagens or proteins binding to the glycosaminoglycans of PNNs

(GO:0005539; enrichment 4.2, p-value 5.10^{-6}). This is consistent with the hypothesis that TH/TRα1

signaling stimulates the elaboration of PNNs around PV neurons. The expression of two of the

downregulated genes in this family (*Sema7a* encoding semaphorin 7A; *Hapnl4* encoding hyaluronan

and proteoglycan link protein 4) was restored after a short treatment with THs, suggesting that they

are TRα1 target genes.

Another prominent characteristic of deregulated genes in hypothyroid and mutant mice was the
presence of genes that encode voltage-gated potassium channels (GO:0005249; enrichment 6.1, p-
value 10^{-6}). As these genes appeared to be sensitive to acute TH treatment, their expression is
unlikely to be under the direct control of TRα1. This observation suggests that hypothyroidism or
long-term expression of TRα1^{L400R} in GABAergic neurons damage the postnatal remodeling of ions
channels, which is a key event for proper maturation of PV neurons in the neocortex (Okaty et al.,
2009). To further evaluate this possibility, we extracted the genes that mostly encode ions channels
and whose expression is modified in this process of postnatal maturation, as defined by Okaty and
colleagues (Okaty et al., 2009). This analysis (Fig. 3e) revealed that this postnatal transition is
impaired and that the immature status of PV neurons persists in hypothyroid and mutant mice.

Discussion

In the present study, we extend our previous observations showing that a cell-autonomous action of
TH/TRα1 in GABAergic neurons is required for the maturation of neocortical PV neurons in mice. We
identified a time window that spans from birth to the end of the third postnatal week, during which
the action of THs is critical for the proper maturation of PV neurons. We have also found that the
defects induced in PV neuron maturation when TH signaling is blocked in GABAergic neurons from
early stages of development persist up to adulthood. Finally, analysis of gene expression at the end
of the second week of life pinpointed a set of genes that are directly regulated by TH/TRα1 in
GABAergic neurons. Some of these genes encode ion channels and PNN components, reflecting the
involvement of THs in essential steps of PV neuron maturation in the developing cortex.

Using genetically modified mouse models expressing TRα1^{L400R}, which exerts dominant negative
activity over TRα1, we have gathered evidence suggesting that the time window during which THs
promote PV neuron maturation is in the early postnatal weeks, notably during the second postnatal
week. First, blocking TH/TRα1 signaling in GABAergic neurons after PND7 is sufficient to alter the
expression of parvalbumin, as evidenced at PND14. On the contrary, blocking TH/TRα1 signaling in
GABAergic neurons after PND21 does not significantly alter PV neuron maturation, as assessed in
adulthood. PV neurons also show near-normal maturation when the blockade of TH/TRα1 signaling is
restricted to PV neurons and begins only after the onset of the expression of *Pvalb*, the gene that
encodes PV (Ren et al., 2024). Furthermore, a previous whole cortex RNA-seq analysis showed that
gene expression changes rapidly in wild-type mice during the second postnatal week and that this
evolution is stopped when TH/TRα1 signaling is blocked in GABAergic neurons (Richard et al., 2020).
Finally, our novel transcriptome analysis, which was carried out with sorted GABAergic neuron nuclei,
indicates that the postnatal maturation process of PV neurons is blunted by TRα1^{L400R} expression. In

particular, blocking TH/TRα1 signaling appears to compromise the expression of a number of genes
involved in a well-documented transition, during which the repertoire of ion channels is extensively
modified (Okaty et al., 2009). It is remarkable that the second postnatal week, during which THs
seem to play a crucial role in PV neuron maturation, coincides with a peak in circulating levels of THs
(Friedrichsen, Christ et al., 2003). This strongly suggests that THs represent a temporal cue to
synchronize the postnatal maturation of PV neurons.

The striking influence exerted by THs on cortical PV neuron maturation in the first postnatal weeks
does not exclude the possibility that THs also play a role in these neurons earlier during
development, or after weaning. First, we cannot fully rule out that TRα1 has an earlier function in PV
neurons, as recent observations suggest that it could act prenatally to promote the tangential
migration of GABAergic neuron progenitors (da Cunha Menezes, de Abreu et al., 2024). However,
there is no indication that this effect is exerted in a cell-autonomous manner. Second, a potential
role of THs on PV neuron maturation after weaning is supported by observations that were
previously reported in mice expressing TRα1^{R384C} in all tissues, for which a complete recovery in PV
neuron density was observed in adults (Wallis et al., 2008). By contrast, in the present study we have
observed little recovery in PV neuron maturation in adult mice expressing TRα1^{E395fs401} in GABAergic
neurons. This discrepancy is probably explained by the residual capacity of TRα1^{R384C} to bind T3 and
activate transcription (Tinnikov, Nordstrom et al., 2002), which is not the case for TRα1^{E395fs401}
(Richard et al., 2020). Therefore, the late onset of TRβ1 expression in PV neurons (Hochbaum,
Hulshof et al., 2024) did not appear to be sufficient to counteract the influence of the dominant
negative TRα1^{E395fs401} mutated receptor. Thus, up to now it is only for the first post-natal weeks that
there is evidence for a direct, cell-autonomous, action of THs in GABAergic neurons, impacting PV
neuron maturation.

The mode of action of THs in PV neurons during the second postnatal week is clarified by RNA-seq
data. In the present study, we first found that PV neurons were much more sensitive to variations in
TH signaling at PND14 than in adulthood, where the same method only showed minor changes in
gene expression (Ren et al., 2024). Another finding of the present study was that genes that were
consistently deregulated by blocking TH/TRα1 signaling in GABAergic neurons encode extracellular
matrix proteins, specific PNN components or proteins interacting with PNN glycosaminoglycans. We
used rapid recovery of gene expression after TH treatment in hypothyroid mice as an indication of
direct transcriptional control of gene expression by TRα1. This allowed us to identify a small subset of
genes whose deregulation is more likely to be the initial set of events preventing PV neuron
maturation. Among these were genes encoding transcription factors or cofactors (*Dbp, Hr, Klf9*), that

were already reported to be regulated by TRα1 in other contexts (Zekri et al., 2022) and that could

act as upstream regulators of other deregulated genes. Moreover, in this subset of putative TRα1

target genes, we found again genes encoding PNN components, suggesting a direct regulation of

PNNs by THs. Thus, it appears that liganded TRα1 might notably activate the transcription of genes

encoding collagens (*Col24a1* and *Col9a2* genes), ephrin A4 (*Epha4*), laminin alpha 3 subunit (*Lama3*),

semaphorin 7A (*Sema7a*) and hyaluronan and proteoglycan link protein 4 (*Hapln4*). This later protein

notably represents an essential component of the PNN (Edamatsu, Miyano et al., 2018, Nojima,

Miyazaki et al., 2021, Popelar, Diaz Gomez et al., 2017). Thus, a major finding of the present study is

that THs directly regulate the elaboration of PNNs, which is a prominent outcome of the late

maturation process of PV neurons.

The novel data presented in this paper provide strong support for the hypothesis that TH is a major

temporal signal for the opening and closing of the critical period of heightened plasticity, as

previously suggested (Batista & Hensch, 2019). The critical period occurs at slightly different times in

different neocortical areas, as does the onset of TH signaling (Quignodon, Legrand et al., 2004). It is

characterized by a series of cellular events that play a crucial role in the elaboration of functional

neuronal circuits (Hensch, 2005). The beginning of parvalbumin synthesis marks the opening of the

critical period, while the elaboration of PNNs around PV neurons stabilizes synapses and decides its

closure. The results presented above show that TH/TRα1 signaling directly regulate the onset of

parvalbumin expression, as well as the synthesis of several PNN components. Thus, our findings

suggest that both the opening and the closure of the critical period are directly stimulated by

liganded TRα1. Therefore, THs appear to act in the postnatal neocortex as it does during amphibian

metamorphosis, as an external cue that represents an important temporal signal. During brain

development, the influence of THs is also found in glial cells, as TH signaling is particularly required

for timely differentiation of oligodendrocytes and myelin formation (Durand & Raff, 2000, Picou,

Fauquier et al., 2014). Of note, oligodendrocyte maturation has also been shown to play a role in PV

neuron maturation and in the closure of the critical period (Kalish, Barkat et al., 2020).

Patients with early TH deficiency caused by congenital hypothyroidism (Van Vliet & Deladoey, 2014)

or carrying TRα1 mutations (van Gucht et al., 2017) are currently treated with TH replacement

therapy. Treatment can have a positive influence on cognitive development only if it is initiated soon

after birth. This is an indication that the action of THs on the maturation of the human neocortex is

restricted to an early temporal window, as we have found in mice. An alteration in the timing of the

critical period is also suspected to cause autism spectrum disorders (Sohal & Rubenstein, 2019). In

that regard, it is noticeable that some of the rare patients in whom *THRA* mutations have been

discovered were first diagnosed for autism spectrum disorders (Kalikiri, Mamidala et al., 2017, Yuen,

Thiruvahindrapuram et al., 2015). Within the framework of our hypothesis, the irreversible cognitive

and neurological defects that are seen both in patients with TH-related diseases and in patients with

autism spectrum disorders might be similarly regarded as consequences of a disruption of the critical

period of heightened cortical plasticity. We have pinpointed THs as essential players in defining the

critical period. Delving deeper into the detailed molecular events occurring in the neocortex during

the critical period will be key to finding therapeutic approaches to these debilitating pathologies.

Study limitations

The complete characterization of TRα1 target genes in PV neurons would require the definition of

chromatin occupancy by the receptor, as it informs of its presence on regulatory sequences. This is

not currently feasible with the small number of cells that are collected when sorting GABAergic

neurons from mouse neocortex. Also, the histological methods that we have used only indirectly

inform of the timing of the critical period of heightened plasticity, which can be defined in a number

of different ways (Quast, Reh et al., 2023, Reh, Dias et al., 2020).

Material and methods

Mouse models and treatments

Experiments involving the use of live animals for the current project were approved by local ethics

committees (C2EA015 and C2EA017) and subsequently authorized by the French Ministry of

Research (Projects #33279-2021082516194165 and #25496-2020032811174506). Mice were bred

and maintained at Plateau de Biologie Expérimentale de la Souris (SFR BioSciences Gerland - Lyon

Sud, France).

The $Thra^{AMI}$ allele allows the expression of the dominant-negative TRα1^{L400R} after the Cre/loxP-

mediated deletion of a cassette with polyadenylation signals (Quignodon, Vincent et al., 2007).

$Thra^{Slox}$ is identical, except for the presence of a C-terminus frameshift mutation which leads to the

expression of the dominant-negative TRα1^{E395fs401X}. The $ROSA$-$tdTomato^{lox}$ reporter transgene (also

known as Ai9, MGI Cat# 4436851, RRID: MGI: 4436851) drives the Cre-dependent expression of a red

fluorescent protein, mainly localized in cytoplasm (Madisen, Zwingman et al., 2010). The $ROSA$-

$GSL10gfp^{lox}$ transgene encodes the GS-EGFP-L10a protein, which is a green fluorescent protein fused

to the N-terminus part of the large subunit ribosomal protein L10a, mainly localized in cell nuclei

(Wu, Dellinger et al., 2025). All transgenes were in C57Bl6/J genetic background. *Gad2-Cre* and *Gad2-*

CreER^{T2} are knock-in alleles in which the Cre recombinase reading frame is inserted into the *Gad2*

gene in order to restrict the expression of the recombinase in GABAergic neurons (Taniguchi, He et

al., 2011). Recombination was induced in mice carrying the *Gad2-CreER^{T2}* transgene by

intraperitoneal injection of tamoxifen (T5648, Sigma, dissolved in sterile sunflower oil (10 mg/ml).

PND7 mice received a single 50 µL subcutaneous injection whereas 3-week-old mice received 5 daily

intraperitoneal injections (daily dose: 7.5 µL/g).

Gestating mice were fed for 2 weeks with iodine deficient food supplemented with 0.15% propyl-thio

uracyl (Envigo ref TD.95125) to cause deep hypothyroidism. TH levels were restored by a single

intraperitoneal injection in half of the pups (10 µg T4 + 1 µg T3 dissolved in 50 µL of phosphate buffer

saline, all chemicals from Sigma Aldrich France). The other PTU treated mice were injected with 50 uL

of phosphate buffer saline.

RNA-seq analysis from sorted nuclei

Nuclei were isolated from individual cortex frozen in liquid nitrogen and nuclear RNA was extracted

as previously described (Wu et al., 2025). Nuclear RNA was extracted from sorted nuclei (RNeasy

Micro Kit, Qiagen ref 74004) and quantified using Tapestation4150 (Agilent). 1 ng RNA was reverse-

transcribed using the SMART-Seq v4 Low Input RNA kit (Takara). cDNA was quantified and qualified

using Qubit (Invitrogen) and Tapestation4150 (Agilent). Libraries were then prepared from 1 ng cDNA

using the Nextera XT DNA Library Kit (Illumina). Libraries were sequenced (> 2.10^7 reads/library) on a

Nextseq500 DNA sequencer (Illumina). Reads were aligned on the mouse genome (mm10 GRCm38

release) with Bowtie2 (Galaxy Version 2.4.2 + galaxy 0) (Langmead, Trapnell et al., 2009). Count table

(Suppl. Table 1) was prepared using htseq-count (Galaxy Version 0.9.1 + galaxy 1, mode union,

feature type: gene) (Anders, Pyl et al., 2015). Differential gene expression analysis was performed

with DEseq2 (Galaxy Version 2.1.8.3) (Love, Huber et al., 2014) (Suppl. Table 1) using the following

thresholds: False Discovery Rate < 0.05; p-adjusted value < 0.05; maximum adjusted basecount > 30;

fold-change > 2.0. Hierarchical clustering was performed using Euclidian distance and complete

distance with Clustvis (Metsalu & Vilo, 2015). Gene Ontology analysis was performed with Gorilla

(https://cbl-gorilla.cs.technion.ac.il/). Raw RNA-seq data are available at NCBI Gene Expression

Omnibus (GSE290072 and GSE290207).

Brain slice collection

Each mouse was given a lethal intraperitoneal injection (6 mL/kg) of a mixture of ketamine (33

mg/mL) and xylazine (6.7 mg/mL). The thorax was opened and each mouse was perfused with 3.7%

paraformaldehyde in 0.1 M phosphate buffer at room temperature. Each brain was dissected out,

immersed in fixative at 4°C for 3 h and then in phosphate buffered saline (PBS) at 4°C until sectioning.

Coronal sections (50 μm) were cut with the aid of a vibrating microtome (Integraslice 7550 SPDS,

Campden Instruments, Loughborough, UK), in PBS at room temperature. Brain sections were stored

at -20°C in cryoprotectant (30% ethylene glycol and 20% glycerol in 10 mM low-salt PBS).

Brain slice histology

Immunohistochemistry was performed on free-floating brain sections. For PV labeling, a mouse anti-

PV primary antibody (PARV19, Sigma P3088, 1:2000), and a secondary antibody made in donkey

(anti-mouse DyLight 633, ThermoFisher Scientific, 1:1000) were used. For the concurrent PNN

labeling, a biotinylated Wisteria floribunda lectin (WFL/WFA, Vector Laboratories B-1355, 20 μg/mL),

and Streptavidin coupled to DyLight 488 (Vector Laboratories, SA-5488, 1:1000) were used. Non-

specific binding sites were blocked by incubating sections for 1 h in PBS with 1% BSA and 0.2% Triton

X100. Brain sections were incubated overnight at 4°C with PV antibody and biotinylated WFA, which

were diluted in PBS with 1% BSA, 0.2% Triton X-100 and 1% dimethyl sulfoxide. Sections were

washed in PBS and further incubated for 15 min at room temperature with DAPI (4',6-diamidino-2-

phenylindole, 1:5000, Sigma). Incubation with the secondary antibody lasted for 3 h at room

temperature. Sections were mounted in Fluoroshield™ (Sigma), coverslipped and imaged using an

inverted confocal microscope (Zeiss LSM 780 and Zeiss LSM 800).

Image analysis

Image analysis was performed using ImageJ. Final images resulted from the z projection of 3 optical

sections (Maximum intensity tool) that were 2 μm apart (the resulting images thus reflected the

fluorescence collected over a thickness of 4 μm). PV+ WFA+ Tomato+ cells were detected from final

images using the Cellpose wrapper for Fiji (Stringer, Wang et al., 2021). Numbers of mono-, double-

or triple-labelled cells per slice were measured automatically from 3-4 slices per mouse. The code of

the original macro may be found here: https://github.com/jbrocardplatim/MacroPV

The densities of Tomato+, PV+, WFA+ cells, and their co-localizations were counted in the whole area

of one image taken in the somatosensory cortex area. 3-4 images in each hemisphere were analyzed

for each mouse. Quantitative data were analyzed using two-tailed unpaired T-test. Data are

expressed as mean ± SEM. Data analyses were performed with Excel 2016 and GraphPad Prism

software (v.8.). The level of significance was set at $p < 0.05$.

References

Anders S, Pyl PT, Huber W (2015) HTSeq--a Python framework to work with high-throughput
sequencing data. *Bioinformatics* 31: 166-9
Batista G, Hensch TK (2019) Critical Period Regulation by Thyroid Hormones: Potential Mechanisms
and Sex-Specific Aspects. *Front Mol Neurosci* 12: 77
Berbel P, Navarro D, Roman GC (2014) An evo-devo approach to thyroid hormones in cerebral and
cerebellar cortical development: etiological implications for autism. *Front Endocrinol (Lausanne)* 5:
146
Bernal J, Morte B, Diez D (2022) Thyroid hormone regulators in human cerebral cortex development.
J Endocrinol 255: R27-R36
da Cunha Menezes E, de Abreu FF, Davis JB, Maurer SV, Roshko VC, Richardson A, Dowell J, Cassella
SN, Stevens HE (2024) Effects of gestational hypothyroidism on mouse brain development: Gabaergic
systems and oxidative stress. *Dev Biol* 515: 112-120
de Lecea L, del Rio JA, Soriano E (1995) Developmental expression of parvalbumin mRNA in the
cerebral cortex and hippocampus of the rat. *Brain Res Mol Brain Res* 32: 1-13
del Rio JA, de Lecea L, Ferrer I, Soriano E (1994) The development of parvalbumin-immunoreactivity
in the neocortex of the mouse. *Brain Res Dev Brain Res* 81: 247-59
Durand B, Raff M (2000) A cell-intrinsic timer that operates during oligodendrocyte development.
Bioessays 22: 64-71
Edamatsu M, Miyano R, Fujikawa A, Fujii F, Hori T, Sakaba T, Oohashi T (2018) Hapln4/Bral2 is a
selective regulator for formation and transmission of GABAergic synapses between Purkinje and
deep cerebellar nuclei neurons. *J Neurochem* 147: 748-763
Friedrichsen S, Christ S, Heuer H, Schafer MK, Mansouri A, Bauer K, Visser TJ (2003) Regulation of
iodothyronine deiodinases in the Pax8-/- mouse model of congenital hypothyroidism. *Endocrinology*
144: 777-84
Gilbert ME, Sui L, Walker MJ, Anderson W, Thomas S, Smoller SN, Schon JP, Phani S, Goodman JH
(2007) Thyroid hormone insufficiency during brain development reduces parvalbumin
immunoreactivity and inhibitory function in the hippocampus. *Endocrinology* 148: 92-102
Hensch TK (2005) Critical period plasticity in local cortical circuits. *Nat Rev Neurosci* 6: 877-88
Hijazi S, Smit AB, van Kesteren RE (2023) Fast-spiking parvalbumin-positive interneurons in brain
physiology and Alzheimer's disease. *Mol Psychiatry* 28: 4954-4967

Hochbaum DR, Hulshof L, Urke A, Wang W, Dubinsky AC, Farnsworth HC, Hakim R, Lin S, Kleinberg G,
Robertson K, Park C, Solberg A, Yang Y, Baynard C, Nadaf NM, Beron CC, Girasole AE, Chantranupong
L, Cortopassi MD, Prouty S et al. (2024) Thyroid hormone remodels cortex to coordinate body-wide
metabolism and exploration. *Cell* 187: 5679-5697 e23

Huang ZJ, Kirkwood A, Pizzorusso T, Porciatti V, Morales B, Bear MF, Maffei L, Tonegawa S (1999)
BDNF regulates the maturation of inhibition and the critical period of plasticity in mouse visual
cortex. *Cell* 98: 739-55

Kalikiri MK, Mamidala MP, Rao AN, Rajesh V (2017) Analysis and functional characterization of
sequence variations in ligand binding domain of thyroid hormone receptors in autism spectrum
disorder (ASD) patients. *Autism research : official journal of the International Society for Autism*
Research

Kalish BT, Barkat TR, Diel EE, Zhang EJ, Greenberg ME, Hensch TK (2020) Single-nucleus RNA
sequencing of mouse auditory cortex reveals critical period triggers and brakes. *Proc Natl Acad Sci U*
S A 117: 11744-11752

Langmead B, Trapnell C, Pop M, Salzberg SL (2009) Ultrafast and memory-efficient alignment of short
DNA sequences to the human genome. *Genome biology* 10: R25

Love MI, Huber W, Anders S (2014) Moderated estimation of fold change and dispersion for RNA-seq
data with DESeq2. *Genome biology* 15: 550

Madisen L, Zwingman TA, Sunkin SM, Oh SW, Zariwala HA, Gu H, Ng LL, Palmiter RD, Hawrylycz MJ,
Jones AR, Lein ES, Zeng H (2010) A robust and high-throughput Cre reporting and characterization
system for the whole mouse brain. *Nature neuroscience* 13: 133-40

Markossian S, Guyot R, Richard S, Teixeira M, Aguilera N, Bouchet M, Plateroti M, Guan W, Gauthier
K, Aubert D, Flamant F (2018) CRISPR/Cas9 Editing of the Mouse Thra Gene Produces Models with
Variable Resistance to Thyroid Hormone. *Thyroid* 28: 139-150

Mayerl S, Heuer H (2023) Thyroid hormone transporter Mct8/Oatp1c1 deficiency compromises
proper oligodendrocyte maturation in the mouse CNS. *Neurobiol Dis* 184: 106195

Metsalu T, Vilo J (2015) ClustVis: a web tool for visualizing clustering of multivariate data using
Principal Component Analysis and heatmap. *Nucleic Acids Res* 43: W566-70

Nojima K, Miyazaki H, Hori T, Vargova L, Oohashi T (2021) Assessment of Possible Contributions of
Hyaluronan and Proteoglycan Binding Link Protein 4 to Differential Perineuronal Net Formation at
the Calyx of Held. *Front Cell Dev Biol* 9: 730550

Okaty BW, Miller MN, Sugino K, Hempel CM, Nelson SB (2009) Transcriptional and
electrophysiological maturation of neocortical fast-spiking GABAergic interneurons. *The Journal of*
neuroscience : the official journal of the Society for Neuroscience 29: 7040-52

Picou F, Fauquier T, Chatonnet F, Richard S, Flamant F (2014) Deciphering direct and indirect
influence of thyroid hormone with mouse genetics. *Mol Endocrinol* 28: 429-41

Popelar J, Diaz Gomez M, Lindovsky J, Rybalko N, Burianova J, Oohashi T, Syka J (2017) The absence
of brain-specific link protein Bral2 in perineuronal nets hampers auditory temporal resolution and
neural adaptation in mice. *Physiol Res* 66: 867-880

Quast KB, Reh RK, Caiati MD, Kopell N, McCarthy MM, Hensch TK (2023) Rapid synaptic and gamma
rhythm signature of mouse critical period plasticity. *Proc Natl Acad Sci U S A* 120: e2123182120

Quignodon L, Legrand C, Allioli N, Guadano-Ferraz A, Bernal J, Samarut J, Flamant F (2004) Thyroid
hormone signaling is highly heterogeneous during pre- and postnatal brain development. *J Mol*
Endocrinol 33: 467-76

Quignodon L, Vincent S, Winter H, Samarut J, Flamant F (2007) A point mutation in the activation
function 2 domain of thyroid hormone receptor alpha1 expressed after CRE-mediated recombination
partially recapitulates hypothyroidism. *Mol Endocrinol* 21: 2350-60

Reh RK, Dias BG, Nelson CA, 3rd, Kaufer D, Werker JF, Kolb B, Levine JD, Hensch TK (2020) Critical
period regulation across multiple timescales. *Proc Natl Acad Sci U S A* 117: 23242-23251

Ren J, Flamant F (2023) Thyroid hormone as a temporal switch in mouse development. *Eur Thyroid J*
12

Ren J, Markossian S, Guyot R, Aubert D, Li D, Cauli B, Riet F, Wong J, Flamant F, Richard S (2024)
Thyroid hormones maintain parvalbumin neuron functions in mouse neocortex. *bioRxiv*:
2024.07.16.603713

Richard S, Guyot R, Rey-Millet M, Prieux M, Markossian S, Aubert D, Flamant F (2020) A Pivotal
Genetic Program Controlled by Thyroid Hormone during the Maturation of GABAergic Neurons.
iScience 23: 100899

Richard S, Ren J, Flamant F (2023) Thyroid hormone action during GABAergic neuron maturation: The
quest for mechanisms. *Front Endocrinol (Lausanne)* 14: 1256877

Sohal VS, Rubenstein JLR (2019) Excitation-inhibition balance as a framework for investigating
mechanisms in neuropsychiatric disorders. *Mol Psychiatry* 24: 1248-1257

Stringer C, Wang T, Michaelos M, Pachitariu M (2021) Cellpose: a generalist algorithm for cellular
segmentation. *Nat Methods* 18: 100-106

Takesian AE, Hensch TK (2013) Balancing plasticity/stability across brain development. *Prog Brain Res*
207: 3-34

Taniguchi H, He M, Wu P, Kim S, Paik R, Sugino K, Kvitsiani D, Fu Y, Lu J, Lin Y, Miyoshi G, Shima Y,
Fishell G, Nelson SB, Huang ZJ (2011) A resource of Cre driver lines for genetic targeting of GABAergic
neurons in cerebral cortex. *Neuron* 71: 995-1013

Tasic B, Menon V, Nguyen TN, Kim TK, Jarsky T, Yao Z, Levi B, Gray LT, Sorensen SA, Dolbeare T,
Bertagnolli D, Goldy J, Shapovalova N, Parry S, Lee C, Smith K, Bernard A, Madisen L, Sunkin SM,
Hawrylycz M et al. (2016) Adult mouse cortical cell taxonomy revealed by single cell transcriptomics.
Nature neuroscience 19: 335-46

Tinnikov A, Nordstrom K, Thoren P, Kindblom JM, Malin S, Rozell B, Adams M, Rajanayagam O,
Pettersson S, Ohlsson C, Chatterjee K, Vennstrom B (2002) Retardation of post-natal development
caused by a negatively acting thyroid hormone receptor alpha1. *Embo J* 21: 5079-87

Uchida K, Hasuoka K, Fuse T, Kobayashi K, Moriya T, Suzuki M, Katayama N, Itoi K (2021) Thyroid
hormone insufficiency alters the expression of psychiatric disorder-related molecules in the
hypothyroid mouse brain during the early postnatal period. *Sci Rep* 11: 6723

van Gucht ALM, Moran C, Meima ME, Visser WE, Chatterjee K, Visser TJ, Peeters RP (2017)
Resistance to Thyroid Hormone due to Heterozygous Mutations in Thyroid Hormone Receptor Alpha.
Current topics in developmental biology 125: 337-355

Van Vliet G, Deladoey J (2014) Diagnosis, treatment and outcome of congenital hypothyroidism.
Endocr Dev 26: 50-9

Wallis K, Sjogren M, van Hogerlinden M, Silberberg G, Fisahn A, Nordstrom K, Larsson L, Westerblad
H, Morreale de Escobar G, Shupliakov O, Vennstrom B (2008) Locomotor deficiencies and aberrant
development of subtype-specific GABAergic interneurons caused by an unliganded thyroid hormone
receptor alpha1. *The Journal of neuroscience : the official journal of the Society for Neuroscience* 28:
1904-15

Wingert JC, Sorg BA (2021) Impact of Perineuronal Nets on Electrophysiology of Parvalbumin
Interneurons, Principal Neurons, and Brain Oscillations: A Review. *Front Synaptic Neurosci* 13:
673210

Wu S, Dellinger J, Markossian S, Dusabyinema Y, Guyot R, Hughes S, Aubert D, Fackeure M, Gauthier
K, Gillet B, Jiang W, Flamant F (2025) An Atlas of Thyroid Hormone Responsive Genes in Adult Mouse
Hypothalamus. *Endocrinology* 166

Yuen RK, Thiruvahindrapuram B, Merico D, Walker S, Tammimies K, Hoang N, Chrysler C,
Nalpathamkalam T, Pellecchia G, Liu Y, Gazzellone MJ, D'Abate L, Deneault E, Howe JL, Liu RS,
Thompson A, Zarrei M, Uddin M, Marshall CR, Ring RH et al. (2015) Whole-genome sequencing of
quartet families with autism spectrum disorder. *Nature medicine* 21: 185-91

Zekri Y, Guyot R, Flamant F (2022) An Atlas of Thyroid Hormone Receptors' Target Genes in Mouse
Tissues. *Int J Mol Sci* 23

Figure legends:

Fig. 1 | Persistent defect in PV neuron maturation in mice expressing TRα1^{E395fs401X}. Histological

analysis of *Thra^{+/+} ROSA-tdTomato^{lox/+} Gad2-Cre* littermates (controls) and *Thra^{Slox/+} ROSA-*

tdTomato^{lox/+} Gad2-Cre mice (mutants) that express TRα1^{E395fs401X} in the GABAergic neurons and their

progenitors from embryonic day 12.5 to PND21 (**a-d**, controls: n = 6, mutants: n = 6) and to adult

stage (**e-h**, controls: n = 6, mutants: n = 5). **a, e** Immunohistochemistry of parvalbumin and WFA-

labeled PNNs in PV neurons of somatosensory cortex at striatum level. **b, f** The density of PV+ cells

was lower in mutants at both ages. **c, g** The density of WFA+ cells was lower in mutants at both ages.

d, h The proportion of PV neurons surrounded by PNNs was lower in mutants at both ages. Data are

presented as mean ± SEM and analyzed using an unpaired two-tailed T-test (*$p < 0.05$, **$p < 0.01$,

***$p < 0.001$). Also see Suppl. Fig. 1 for related analyses.

Fig. 2 | A time window for TH action on PV neuron maturation. Histological analysis of *Thra^{+/+} ROSA-*

tdTomato^{lox/+} Gad2-CreER^{T2} or *Gad2-CreER^{T2}* negative littermates (controls) and *Thra^{AMI/+} ROSA-*

tdTomato^{lox/+} Gad2-CreER^{T2} mice (mutants) that express TRα1^{L400R} in the GABAergic lineage from

PND7 to PND14 (**a-d**, controls: n = 8, mutants: n = 6) and from PND21 to adult (**e-h**, controls: n = 11,

mutants: n = 7). **a, e** Immunohistochemistry of parvalbumin and WFA-labeled PNNs in PV neurons of

somatosensory cortex at striatum level. **b, f** The density of PV+ cells was lower in mutants in former

but not in latter. **c, g** The density of WFA+ cells had no clear difference in both cases. **d, h** The

proportion of PV neurons surrounded by PNNs had no clear difference in both cases. Data are

presented as mean ± SEM and analyzed using unpaired two-tailed T-test (*$p < 0.05$, **$p < 0.01$, ***p

< 0.001).

Fig. 3 | RNA-seq analysis of gene expression in sorted GABAergic neuron nuclei. Nuclei were sorted

at PND14 from the neocortex of *Thra^{+/+} ROSA-GSL10gfp^{lox} Gad2-Cre* mice (Ctrl), *Thra^{AMI/+} ROSA-*

GSL10gfp^{lox} Gad2-Cre mice expressing TRα1^{L400R} in GABAergic neurons (Mut), hypothyroid *Thra^{+/+}*

ROSA-GSL10gfp^{lox} Gad2-Cre mice (Hypoth.), or hypothyroid *Thra^{+/+} ROSA-GSL10gfp^{lox} Gad2-Cre* mice

that received a single TH injection at PND13 (+TH). **a** Venn diagram of genes that were either

downregulated by TRα1^{L400R}, downregulated by hypothyroidism or induced after TH treatment. **b**

Venn diagram of genes that were either upregulated by TRα1^{L400R}, upregulated by hypothyroidism or

repressed after TH treatment (thresholds: Fold change > 2 or < 0.5, maximal adjusted base counts >

30, adjusted p-value > 0.05). **c** Heatmap showing the genes that were downregulated by both

TRα1^{L400R} and hypothyroidism. Note the variable response to TH. **d** Heatmap showing the genes that

were upregulated by both TRα1^{L400R} and hypothyroidism. **e** Heatmap of markers of PV neuron

maturation. The marks on the right side are for genes whose expression normally increases (red) or

decreases (green) between PND7 and PND21. The alterations of gene expression in mutant mice

reveals a blockade of the PV neuron maturation program, caused either by TRα1^{L400R} expression or by

hypothyroidism. A 24h TH treatment two weeks after birth is not sufficient to restore the expression

of these marker genes. Also see Suppl. Table 1 for related datasets.

Acknowledgements:

We acknowledge the contribution of SFR Biosciences (Université Claude Bernard Lyon 1, CNRS

UAR3444, Inserm US8, ENS de Lyon): Nadine Aguilera and the Plateau de Biologie Expérimentale de

la Souris (ANIRA-PBES) for mouse breeding; ANIRA-CYTOMETRIE and particularly Sébastien

Dussurgey for nuclei sorting; ANIRA-AGC for mouse genotyping. We thank Benjamin Gillet and

Sandrine Hughes of the deep sequencing facility (PSI IGFL, Lyon). We also thank Elodie Martel for

carrying out histology experiments as part of her internships and Isabelle Dusart (Institut de Biologie

Paris Seine, France) for helpful discussions. We acknowledge China Scholarship Council for a

scholarship to Juan REN to do her PhD project in ENS de Lyon. Research fundings were from

European Union's Horizon 2020 research and innovation program, under grant agreement No.

825753 (ERGO), Agence Nationale de la Recherche (Thyromut2 program; ANR-15-CE14-0011-01).

Disclosure and competing interests statement

The authors have nothing to declare.