Induction of hemodynamic traveling waves by glial-related vasomotion in a

rat model of neuroinflammation: implications for functional neuroimaging

Mickaël Pereira[1], Marine Droguerre[2], Marco Valdebenito[3], Louis Vidal[1], Guillaume Marcy[4], Sarah Benkeder[5],

Jean-Christophe Comte[1], Olivier Pascual[5], Luc Zimmer[1, 3, 6], Benjamin Vidal[1,3*]

[1] Université Claude Bernard Lyon 1, Lyon Neuroscience Research Center, Lyon, France
[2] Theranexus, Lyon, France
[3] CERMEP-Imaging Platform, Lyon, France
[4] Université Claude Bernard Lyon1, Bioinformatic Platform of the Labex Cortex, Lyon, France
[5] Institut MeLiS, INSERM U1314 CNRS UMR5284, Lyon, France
[6] Hospices Civils de Lyon, Lyon, France
*Corresponding author: benjamin.vidal@cermep.fr

Keywords: functional ultrasound imaging, vasomotion, neurovascular coupling, functional connectivity, glia,

neuroinflammation

Abstract

Cerebral hemodynamics are crucial for brain homeostasis and serve as a key proxy for brain activity. Although this process involves coordinated interaction between vessels, neurons and glial cells, its dysregulation in neuroinflammation is not well understood.

We used in vivo mesoscopic functional ultrasound imaging to monitor cerebral blood volume changes during neuroinflammation in rats injected with lipopolysaccharide (LPS) in the visual cortex, under resting-state or visual stimulation, combined to advanced ex vivo techniques for glial cell reactivity analysis.

Cortical neuroinflammation induced large oscillatory hemodynamic traveling waves in the frequency band of vasomotion (~0.1 Hz) in both anesthetized and awake rats. Vasomotor waves traveled through large distances between adjacent penetrating vessels, spanning the entire cortex thickness, and even extending to subcortical areas. Moreover, vasomotion amplitude correlated with microglial morphology changes and was significantly reduced by astrocytic toxins, suggesting that both microglia and astrocytes are involved in the enhancement of vasomotion during neuroinflammation. Notably, functional connectivity was increased under this oscillatory state and functional hyperemia was exacerbated.

These findings reveal new spatiotemporal properties and cellular mechanisms of cerebral vasomotion, and suggest that this is a major component of brain hemodynamics in pathological states. Moreover, reactive microglia and astrocytes are participating to increased vasomotion during neuroinflammation. These results call for a reassessment of vasomotion and traveling waves as primary phenomena when imaging brain hemodynamic activity, particularly in conditions associated with neuroinflammation.

Introduction

Cerebral hemodynamics constitute a fundamental component of the central nervous system (CNS) by providing the necessary oxygen and nutrients to support its substantial energy consumption. Cerebral blood flow is dynamically adapted in response to changes in neuronal activity, a process known as neurovascular coupling. This fundamental process underlies functional neuroimaging techniques that measure hemodynamic parameters to infer changes in brain activity, such as functional magnetic resonance imaging (fMRI), intrinsic optical imaging, laser speckle contrast imaging, near-infrared spectroscopy, and more recently functional ultrasound imaging (fUSi)[1–6]. It is therefore of major importance to understand the processes governing cerebral hemodynamics to gain deeper insights into brain functions, as well as to accurately interpret functional neuroimaging findings, in both physiological and pathological states.

Despite many efforts and advances, the cellular mechanisms that govern hemodynamic fluctuations in resting-state as well as functional hyperemia (i.e, the increase of cerebral blood flow in response to an increased neuronal activity) remain elusive and controversial. An emerging hypothesis is that vasomotion, i.e., the rhythmic fluctuations of vessel diameter (~0.1 Hz) produced by the autonomous contraction of smooth muscle cells, is naturally synchronized to neuronal activity to meet the energetic demands under physiological conditions[7]. This model, although debated[8], provides a possible cellular basis for functional connectivity. In parallel, other studies have suggested that glial cells are equally important for regulating hemodynamic fluctuations. Astrocytes, in particular, are strategically positioned to mediate interactions between neurons and blood vessels as their end-feet envelop most of the cerebral vasculature. Astrocytic control of cerebral blood flow during functional hyperemia has been extensively studied, and involves the Ca^{2+}-dependent release of vasoactive metabolites of arachidonic acid, as well as potassium efflux from the endfeet[9–16]. They also appear to be involved in the regulation of cerebral hemodynamics during resting-state, as they were recently suggested to exert a negative control on the amplitude of vasomotion at 0.1 Hz in arterioles[17]. This brings vasomotion at the center of a hypothetic cellular model for explaining spontaneous hemodynamic fluctuations and resting-state connectivity, with SMCs setting the tempo around 0.1 Hz, astrocytes calibrating the gain, and neurons serving as the conductor of this orchestrated response. Other glial actors such as microglia, the key regulators of inflammatory processes in the brain, may also be involved as they can

dynamically interact with the neurovascular unit and have been shown to modulate functional
hyperemia[18].

Fewer studies have focused on the cellular mechanisms involved in the control of
cerebral blood flow in pathological conditions[9,19]. This is important because all cellular
players of the neurovascular unit, and in particular microglia and astrocytes, undergo major
morphological, functional and molecular changes during inflammatory processes that can
occur in many neuropsychiatric conditions. Such changes could have major impact on the
regulation of the cerebral hemodynamics, which must be taken into account for the
interpretation of neuroimaging studies and for understanding the alterations of brain
homeostasis that can occur during neuroinflammatory processes .

Using functional ultrasound imaging (fUSi), we aimed to extensively characterize the
cerebral hemodynamics changes that occur in a neuroinflammatory state and the influence of
reactive glial cells in these alterations. This recent technique can map cerebral blood volume
(CBV) changes at a mesoscopic scale with higher sensitivity than fMRI[20] even in deep areas,
unlike optical techniques, and can be used in freely moving rodents[21-23]. CBV changes were
studied during LPS-induced neuroinflammation through injection in the primary visual cortex,
during resting-state as well as visual stimulation. In parallel, the impact of the inflammatory
challenge on the reactivity of glial cells was assessed by advanced morphological analyzes
and single-nuclei transcriptomics. We show that cortical neuroinflammation can drastically
enhance hemodynamic oscillations at 0.1 Hz, the typical rhythm of vasomotion. Adding to the
recent descriptions of vasomotion as long-wavelength traveling waves along cortical pial
vessels[24], the vasomotor waves traveled across large areas between neighboring penetrating
vessels and even reached subcortical vessels. We also highlight that vasomotion amplitude
may be locally driven by the presence of reactive glial cells, as suggested by a correlation
between microglial morphological changes and hemodynamic oscillations at 0.1 Hz, and the
significant inhibition of inflammatory vasomotion by astrocytic toxins. Finally, enhanced
vasomotion was associated with an increase in cortical connectivity, extending previous
findings that vasomotor fluctuations are a major component of functional connectivity, and
with an increased hyperemic response during visual stimulation, suggesting that the two
phenomena share common but non-competitive pathways under specific states. These
findings demonstrate that vasomotion and traveling waves around 0.1 Hz are fundamental
players of cerebral hemodynamic fluctuations, especially during neuroinflammation, and
emphasize the importance of reactive glial cells in the control of spontaneous changes of

cerebral blood volume in pathological states, usually interpreted only in terms of neuronal
activity in functional neuroimaging studies.

Materials and methods

Animals

All animal experimentation and housing were carried out in accordance with the guidelines of
the European Union (Directive 2010/63/EU) and approved by the French Ministry of
Research (authorization reference: APAFIS-36043). The animals were kept under controlled
environmental conditions (12/12 h light-dark cycle, food and water ab libitum, controlled
temperature) with at least one week of acclimatation before any experiment. All efforts were
made to ensure the animal welfare. A total of 62 Wistar male rats (from Janvier Labs, Le
Genest-St-Isle, France) were used in this study.

For the main experiment, 40 rats were used. The rats first underwent a skull thinning surgery
the day before beginning the imaging protocol (6 animals were excluded after the surgery due
to the presence of hematomas). The day of the first fUSi session, the rats received
intracerebral bilateral injections of the following agents: NaCl 0.9% (VEH, n = 6),
lipopolysaccharide (LPS, n = 6), L-alpha amino acid (AAA, n = 6), fluorocitrate (FC, n = 5), a
combination of LPS and AAA (LPS+AAA, n = 5), or a combination of LPS and FC
(LPS+FC, n = 5), 1 h before the fUSi acquisition under anesthesia (except one rat in the
LPS+AAA group that died before starting the imaging session). The rats were then used for a
second fUSi session under anesthesia, 48 h after the intracerebral injections. The brains of all
rats imaged twice were used for immunofluorescence staining (n = 6 for VEH, LPS and AAA;
n = 5 for FC, LPS+AAA and LPS+FC) for assessment of different neuropathological features
and verification of astrocyte loss in the groups that received the gliotoxins (leading to the
exclusion of one rat from the LPS+AAA group because no astrocytic lesion could be detected
for this animal).

For further evaluation of the temporal effects of LPS, 12 additional rats underwent thinned-
skull surgery (24 h before imaging and/or intracerebral injection) and were used for one
anesthetized fUSi session, either without injection (n = 3), 24 h (n = 3) or 7 days (n = 3) after
LPS injection. For awake fUSi, 4 other rats underwent skull thinning (24 h before imaging
and/or intracerebral injection) and were imaged in freely moving protocol before LPS

injection (n = 3), 1 h (n = 1) and/or 48 h (n = 3) after bilateral LPS injections. For single-

nuclei transcriptomic analysis, 6 other rats were bilaterally injected with VEH or LPS and

euthanized before sampling the brains 48 h (n = 2 VEH, n = 2 LPS) or 7 days (n = 2, LPS)

after injection.

Surgical procedures

Skull Thinning. A thinned-skull window was performed before fUSi to enable sufficient

penetration of acoustic waves, as previously described in the rat[25,26]. For this purpose, rats

were anesthetized with isoflurane (induction 4%, then 2% in a 50/50 mix of air and O_2 during

the surgery) and received a subcutaneous injection of buprenorphine (0.05 mg/kg; Buprecare,

Axience). The top of the head was shaved and cleaned with povidone-iodine (Vetedine,

Vetoquinol), then 30 min after buprenorphine pretreatment, the scalp was incised. The skin

was carefully pulled to visualize the lateral sides of the skull from bregma to lambda

anatomical landmarks. The skull thinning was performed from -3.00 mm to -8.00 mm (AP)

and +5.00 to 5.00 mm (L) from bregma using a drill at low speed with a micro drill steel burr

(Harvard Apparatus, 75-1887). To avoid overheating during drilling sessions, cold saline

solution was regularly added on the skull. When the skull was thin enough to be flexible, the

scalp was sutured, povidone-iodine and lidocaine 5% (EMLA 5%, Aspen) were locally

applied, and the rats were placed under heating light to recover from anesthesia.

Intracerebral injections. After 24 h of recovery, rats were anesthetized with isoflurane (4%

for induction, then lowered to 2% in a 50/50 mix of air and O_2 during the surgery) and placed

in a stereotaxic frame with a heating blanket. The wound was cleaned and lidocaine 2.5% was

applied before removing sutures and exposing the thinned-skull window. Animals received

bilateral injections of 4 µl of solution (LPS, FC, AAA, VEH, LPS+AAA or LPS+FC) at a rate

of 0.4 µl/min into the primary visual cortex (AP: -6; L: ±4.0; D: 1.6 - in mm from bregma). At

the end of each injection, the needle was left in place for 5 min before being slowly removed.

LPS (1.25 µg/µl; L2880), FC (0.25 nmol/µl; F9634) and AAA (12.5 µg/µl; A7275) were

purchased from Sigma-Aldrich. For the preparation of FC solution, DL-fluorocitric acid

barium salt was dissolved in 1 mL of HCl 0.1 M. Two drops of Na_2SO_4 were then added to

precipitate Ba^{2+}. After adding 2 mL of Na_2HPO_4, the solution was centrifuged at 1000 g for 5

min, and 0.31 mL of supernatant was diluted in 0.9% NaCl to the final concentration (while

adjusting the pH to 7.4).

Functional Ultrasound Imaging

Rats were imaged using a preclinical ultrasound imaging system (Iconeus V1, Paris, France). Doppler vascular images were obtained using the Ultrafast Compound Doppler Imaging technique[27]. Each frame corresponded to a Compound Plane Wave frame[28] resulting from the coherent summation of backscattered echoes obtained after successive tilted plane waves emissions. Each stack of 200 compounded frames acquired at 500 Hz frame rate was processed with a dedicated spatiotemporal filter using Singular Value Decomposition[29] to extract the blood volume signal from the tissue signal, to obtain the Power Doppler images. These final images have a temporal resolution of 0.4 s, a spatial resolution of 100 µm and are directly proportional to the cerebral blood volume.

Anesthetized imaging protocol. At the end of the intracerebral injection procedure, the animals received a subcutaneous injection of medetomidine (0.05 mg/kg; Domitor, Vetoquinol), and were moved in another stereotaxic frame under the ultrasound imaging probe before performing the acute fUSi acquisition, with isoflurane being still delivered at 2% through a mask to maintain anesthesia. A subcutaneous catheter was placed and the medetomidine infusion (0.1 mg/kg/h) was started, while progressively decreasing isoflurane to 0.4%, to follow the consensus protocol for studying functional connectivity in the rat[30]. Respiratory rate and body temperature were continuously monitored (TCMT, Minerve, France) and the animal temperature was maintained using a heating blanket. The thinned skull was exposed and covered with ultrasound gel. After reaching the final concentration of isoflurane, a period of at least 10 min was added before starting the imaging experiments to ensure physiological stability of the animals. Then the acute session of fUSi was performed, that included a resting-state acquisition followed by a visual stimulation acquisition. At the end of this imaging session, the scalp was sutured, and betadine and lidocaine were applied on the wound. Finally, rats received a subcutaneous injection of atipamezole (5 mg/kg; Antisedan, Vetoquinol) and were placed under heating light to recover from anesthesia. 48 h later, the second, delayed fUSi session was performed. Animals were anesthetized first with isoflurane under the same conditions and duration as for the stereotaxic injections performed 2 days before, and then were placed in a stereotaxic frame under the ultrasound imaging probe for the second session under the same anesthesia protocol (combination of isoflurane 0.4% and subcutaneous medetomidine, bolus 0.05 mg/kg followed by infusion at 0.1 mg/kg/h), that also included a

resting-state acquisition followed by a visual stimulation. At the end of this second session, an ultrasound localization microscopy (ULM) acquisition was also performed.

Resting-state protocol. For resting-state fUSi, the probe was placed above the injection site (AP: -6; L: ±4.0; D: 1.6 - in mm from bregma) to follow the CBV changes in the primary visual cortex during a 20-minute period, with a sampling rate of 0.4 s.

Visual stimulation protocol. For fUSi acquisition with visual stimuli, the probe was kept in the same coronal plane. The visual stimuli protocol was performed using a screen placed in front of the animal, as previously described[31]. Briefly, the acquisition began with an initial 30-second rest period in the dark, followed by 4 blocks of stimuli, all separated by a 45-second rest period in the dark. Each visual stimuli block consisted of a 30-second period with black and white flickering at 3 Hz, as this frequency was shown to elicit the best response in the visual cortex in a previous fUSi study in the rat[32]. The stimulation was performed using a homemade Python script.

Ultrasound localization microscopy protocol. At the end of the last imaging session, an ULM acquisition was performed for each animal. An intravenous catheter was set into the caudal vein before starting the acquisition. Two boluses of 300 µL of a microbubble ultrasound contrast agent (Sonovue, Bracco Imaging, France) were slowly injected at the start of acquisition and at 2.5 min of acquisition, for a total acquisition time of 5 min. This sequence enables to visualize the vasculature and to measure the blood flow velocity with very high spatial resolution (10 µm)[33].

Kinetics of LPS effect. To further evaluate the temporal effects of LPS, an additional group of animals underwent a resting-state acquisition after 0, 1 or 7 days following the intracerebral injection of LPS using the same imaging, anesthesia and surgery conditions as stated above. The impact of skull thinning on our results was also assessed with three rats without injection of LPS, scanned in fUSi under the same conditions.

Freely moving awake imaging protocol. A resting-state fUSi protocol was also performed in awake freely-moving rats. These experiments included a first phase of handling/habituation of 5 days, followed by a surgery to implant the head plate (designed by Iconeus, Paris, France). After a minimal period of 24 h of recovery, a 3-days habituation period was performed, during which the rats were free to move between their home cage and the cage used for imaging experiments. 24 h later, a first baseline fUSi session was performed. Then, the bilateral stereotaxic injections of LPS or VEH were performed as stated previously in primary visual

261 cortex (AP: -6; L: ±4.0; D: 1.6 - in mm from bregma), followed by the second resting-state

262 fUSi in awake condition, 1 h after the intracerebral injections. Animals were left in their initial

263 cage for 2 days before their last resting-state imaging session.

Single-nucleus transcriptomics

Single-nuclei isolation. Dissected frozen cortex was resuspended into 600µl of cold

homogenization buffer that consisted of 10mM Tris Hcl pH 7.5, 146mM NaCl, 1mM $CaCl_2$,

21mM $MgCl_2$, 0.03% Tween20, 0.01% BSA, 10% EZ buffer (Sigma) and 0.2U/µl Protector

RNase Inhibitor (Roche). Tissues were then transferred into 2mL dounce (Kimble) and

homogenized using 10 strokes of the loose pestle followed by 8 strokes of the tight pestle to

release nuclei, on ice. Homogenate was then strained through a 70µm cell strainer

(Pluriselect) and centrifuged at 500g for 5 minutes to pellet nuclei. After removing

supernatant, nuclei were washed in 1mL resuspension buffer containing 10mM Tris Hcl pH

7.5, 10mM NaCl, 3mM $MgCl_2$, 1% BSA and 0.2U/µl Protector RNase Inhibitor and

centrifuged at 500g for 5 minutes. Nuclei were then fixed with the Evercode Nuclei Fixation

kit (#ECF2003) according to manufacturer instructions (Parse Biosciences) and FACS-sorted

based on DAPI intensity to remove debris (BD ARIA). Nuclei were then frozen down.

Single-nucleus RNA sequencing. Nuclei were thawed and loaded in round 1, 2 and 3 plates for

combinatorial barcoding according to manufacturer instructions (#CW02030, Parse

Biosciences). Six sub-libraries of 12,500 nuclei were prepared to target 75,000 nuclei and

12,000 nuclei per condition. Libraries were sequenced using the Novaseq 6000 platform

(Illumina) to target 50,000 reads per nucleus. Parse demultiplexing pipeline v1.1.2 was used

to align reads on the rat reference genome (Rattus_norvegicus.mRatBN7.2) and to produce

the count matrix.

Immunofluorescence

Rats were deeply anesthetized and transcardially perfused with 4% paraformaldehyde in

phosphate-buffered saline (PBS). Brain tissues were post-fixed in 4% paraformaldehyde for

24 h, then immersed in 20% sucrose for 48 h. Fixed brain sections were cut at 40 µm using a

cryostat (Leica Biosystems – CM3050S). After washing in PBS, the sections were

permeabilized in PBST-BSA (0.5% Triton X + 5% BSA in PBS) for 90 min. Primary

antibodies, diluted in blocking solution, were applied overnight at 4°C. After three washing in

PBS, appropriate secondary antibodies coupled to fluorochromes were diluted in PBST-BSA

and applied for 1 h at RT. Slides were rinsed and coverslipped with DAPI fluoromount

mounting media. The antibodies list used is provided in the Supplementary Table 1.

Data analysis

fUSi. All data processing steps were performed as previously described[34,35]. Briefly, a custom

template was created by a first automatic registration step and by computing the mean of all

fUSi acquisitions gathered during resting-state and visual stimulation. This template enabled

to manually refine the registration of all images to each other.

Resting-state. For the analysis of CBV changes and static functional connectivity, the

registered images were band-pass filtered between 0.0008 Hz and 0.2000 Hz. Then, a 2D slice

corresponding to the imaging plane (−6 mm from bregma) was extracted from the SIGMA

atlas[36] and manually co-registered on the fUSi template in the ITK-Snap software. The Power

Doppler signal was automatically extracted from each region of interest - namely, the primary

and secondary visual cortices (PrimVisCtx and SecVisCtx), the retrosplenial dysgranular

cortex (RetroDysCtx), the parts A and B of the retrosplenial granular cortex (RetroGrCtx$_A$ and

RetroGrCtx$_B$) and the superficial part of superior colliculus (SuperiorColl) for each individual

scan using a Matlab script. For the functional connectivity analysis, the temporal correlations

were computed using the Pearson correlation coefficient for each pair of regions. Fisher Z

transformation was applied to the correlation coefficients before calculating the differences

between the correlation matrices in each condition relative to the vehicle.

For the analysis of the amplitude of specific frequency bands (also known as fALFF for

fractional amplitude of low-frequency fluctuations), no frequency filtering was done before

applying the slice of atlas on the registered images for extraction of the fALFF in the different

ROIs. fALFF was obtained using a custom Matlab script adapted from the REST toolbox[37].

Briefly, the power over a frequency band of interest (0.08 to 0.15 Hz) was calculated for each

pixel and divided by the power of the entire frequency range.

For the principal component analysis, the images were band-pass filtered between 0.0008 Hz

and 0.2000 Hz, centered by subtracting the global mean, and normalized prior to reshaping

each acquisition into two dimensions. Following the approach of Cabral et al.,, we calculated

and then averaged the covariance matrices of all acquisitions for each group to capture

common variance patterns[38]. The eigenvectors were derived from this mean covariance matrix

and sorted in descending order of their corresponding eigenvalues. The first principal

components' spatial maps were visualized by reshaping each eigenvector into the original

image format (only the components with eigenvalues higher than 0.1 were retained, with up to

10 components for simplification). The temporal patterns associated with these components

were obtained by projecting the time series data onto the eigenvectors. The Fourier transform

was applied to these temporal patterns to calculate their amplitude spectra, which were then

averaged across acquisitions. For visualization, some acquisitions were reconstructed using

the first 10 principal components, summing the product of each principal component and its

corresponding temporal pattern. The scree plot, showing the proportion of total variance

explained by each principal component, was achieved by computing the total variance across

all components using the global sum of eigenvalues and calculating the cumulative sum of the

individual variances explained by the components as a proportion of the total variance, in

descending order of explained variance.

For the awake dataset, the same preprocessing and analysis (functional connectivity matrices,

fALFF and PCA) were performed on selected continuous periods of 120s that were devoid of

movement artifacts for each scan (estimated using a threshold defined on the time-courses of

the Power Doppler signal, and further confirmed by the absence of obvious artifacts in the

spatial maps of the principal components).

For the analysis of traveling waves, the images were filtered between 0.08 Hz and 0.15 Hz to

focus on the frequency of oscillations. We used the AquA software[39] to automatically extract

static and traveling events. This software detects signal peaks propagating across space

without predefined assumptions about event locations or timings. Originally developed for

fluorescence imaging and astrocytic calcium waves, AquA is explicitly designed for working

with various biological contexts and spatiotemporal scales. After normalization to correct for

the baseline differences in signal intensity and estimate the noise variance, the groups of

active pixels were found using an intensity threshold of 20, a smoothing parameter of 0.5 and

a minimum size criterion of 15 pixels. Significant pixels (z-score above 5) were retained, and

events were detected by identifying local maxima (seeds) in these pixels, extending them in

space and time to estimate their propagation. Events were further refined using a z-score

threshold of 5, based on peak size and noise level. We classified events as dynamic/traveling

if their propagation distance exceeded 5 pixels, and as static otherwise. We averaged the

different features of interest across all dynamic events for each scan for group comparisons

(percentage of dynamic events relative to the total number of events, amplitude of peaks, and

propagation, corresponding to the distance crossed from the beginning to the end of the

event). To explore traveling wave trajectories and locations in the different groups, we

clustered the propagation maps generated by AquA (total of 17377 events with an area > 5 pixels). The individual maps were first binarized, reshaped and concatenated in a two-dimensional array, followed by singular value decomposition (SVD) to retain the first 3000 components. We then performed k-means clustering to identify 4 main clusters corresponding to different spatial locations, regardless of the direction of the waves. These clusters were further subclustered based on horizontal and vertical propagation directions, obtained by averaging the delay values of the original propagation maps in these directions and concatenation of the directional vectors into a two-dimensional array, followed by k-means clustering. This process allowed us to identify 6 distinct spatiotemporal patterns in our dataset, in addition to the events that were static or followed an undefined propagation pattern.

The dynamic seed-based connectivity results were computed on 0.08-0.15 Hz band-pass filtered representative individual scans, with a window size of 15 successive images and a step of 1 image, by measuring the Pearson correlation coefficients between the average time course in the right visual cortex as a seed and all pixels of the image. The time course of the average correlation coefficients in the contralateral hemisphere was displayed as a heatmap with superimposed average CBV changes occurring in both hemispheres.

Visual stimulation. For visual stimulation analysis, the average CBV changes during all stimulations were extracted in the regions of interest and the four time-courses corresponding to each stimulation block were averaged for all individuals. The mean CBV changes values during stimulation as compared to the baseline were estimated for each individual before group comparisons.

Ultrasound localization microscopy. The speed maps were reconstructed using a dedicated Iconeus code, based on the tracking of individuals microbubbles in the ULM data. An additional spatial filter was applied to remove outliers, which decreased the spatial resolution to 10 x 10 x 400 μm. The analysis was restricted to the primary visual cortex area. The pial vessels were manually drawn on each individual image on the absolute speed map, and the penetrating vessels were further differentiated as arterioles or venules depending on the vertical direction of the flow indicated in the directional map. The absolute speed values were averaged in these different vascular compartments for each individual scan and compared between groups.

Immunohistochemistry. Slides were scanned with ZEISS Axioscan 7 Microscope Slide Scanner at the East Lyon Quantitative imaging center of University Lyon1 (CIQLE). For quantification of Iba1, GFAP, and NDRG2 markers, the mean intensity was measured via ImageJ in four to eight areas in the primary visual cortex for each animal (after normalizing the intensity to that of the whole image for NDRG2). For NeuN counting, the neuron number was automatically determined using Spot detector tool of the Icy software.

Microglia morphological analysis. A total of 250 to 400 microglia were cropped individually randomly in four to eight areas in the primary visual cortex for each group. The crops were then binarized and their morphology automatically analyzed with MorphoCellSorter[40] using Matlab R2018. Briefly, 20 non dimensional morphological indexes are measured. Then, a principal component analysis (PCA) is performed to filter the parameters and select only the most discriminant ones. The selected parameters and their weight on the first principal component are subsequently used to form a derivative of an Andrews plot, which allow the two-dimensional representation of high dimensional data. In this representation, each curve represents one microglia, and cells having similar morphological characteristics follow similar trajectories. It is used to establish a ranking of the cells according to their morphological features by using the Andrews score, corresponding to the value of the Andrews plots at the time point where the curves are the most dispersed (i.e. when the variance is maximal).

Single-nucleus RNA-seq data analysis. The gene expression matrix from Parse pipeline was used for downstream analysis using the software R (version 4.4.0) and the R toolkit Seurat (version 5.1.0). Nuclei were excluded from downstream analysis when they had more than 1% mitochondrial genes, fewer than 100 unique genes, more than 15,000 unique molecular identifiers (UMIs) and detected as doublets using AUC cell type marker scores based on AUCell R package. A total of 64,471 nuclei were selected for the three groups and 6 animals studied. Gene expression was normalized using the standard Seurat workflow and the 2000 most variable genes were identified and used for principal component analysis (PCA) at 50 dimensions. The top most significant principal components (PCs) were selected for generating the UMAP, based on the ElbowPlot method in Seurat. Clustering of cells was obtained following Seurat graph-based clustering approach with the default Louvain algorithm for community detection. We then performed differential expression analysis using the FindMarkers function of Seurat and annotated clusters based on expression of marker genes. We thus manually annotated the major classes of cells: Neurons, Microglia, Astrocytes, Oligodendrocytes (OLs) and Vascular cells.

421   *Gene set enrichment.* Overrepresentation analyses of biological processes (BPs) were

422   performed using the R package "Cluster Profiler"[41]. DEGs were extracted using the following

423   criteria: Logfc.threshold = 0.25, min.pct = 0.1.

Statistical analysis

All statistical analyses were performed using GraphPad Prism 8.0.1. For all experiments, a

level of $p < 0.05$ was accepted as evidence for a statistically significant effect (*$p < 0.05$; **p

< 0.01; ***$p < 0.001$). For the correlation and the fALFF analysis, the different groups were

compared using two-way ANOVAs followed by Sidak's or Dunnett's multiple comparisons

test depending on the number of groups. For the comparisons of the ROIs time-courses during

stimulation, two-way repeated-measures ANOVAs were used followed by Sidak's multiple

comparisons tests ($p < 0.05$). For ULM analysis, the difference of perfusion were assessed

using one-way ANOVAs followed by Dunnett's multiple comparisons test or unpaired t-tests

depending on the number of groups. For the traveling waves analysis, the general features of

dynamic events were calculated in AquA software, averaged for each individual, and

compared using one-way ANOVA followed by Dunnett's test. For comparison of the

occurrence of different wave patterns among groups, the percentages of the different patterns

were compared between each group and the vehicle using two-way ANOVAs for each time-

point, and between each time-point for each group using mixed-effect analysis. The p-value of

the interaction between the condition factor (group or time-point) and the patterns factor was

used to identify the significant changes of propagation patterns.

Results

Inflammation induced local hemodynamic oscillations at 0.1 Hz and increased functional connectivity

The effects of bilateral intracortical injection of LPS in the visual cortex were monitored in a

coronal plane centered around the injected area using fUSi, first during resting-state at

different times post-injection (Fig. 1A). The spontaneous CBV changes were extracted from

different ROIs (Fig. 1B). No obvious modification of CBV changes was observed 1h or 48h

after vehicle injection (exemplified in Fig. 1C). However, the administration of LPS induced a

high-amplitude low frequency oscillatory pattern that appeared 48-h post-injection (Fig. 1D;

sky-blue curves, as compared to the dark-blue curves at 1h). From the frequency spectrum,

these waves are occurring between 0.08 and 0.15 Hz, on average centered around 0.125 Hz
(Fig. 1E-F), the typical frequency band of vasomotion. To quantify this phenomenon, the
frequency amplitudes between 0.08 and 0.15 Hz were computed for each individual scan and
each group in the different ROIs. Whereas no significant difference were found between
groups at 1h ($F(1,60) = 1.241$; $p = 0.2697$ - Fig. 1G), the amplitude of oscillations between
0.08-0.15 Hz was significantly higher in LPS compared to vehicle at 48h post-injection
($F(1,60) = 21.71$, $p < 0.0001$ - Fig. 1H) in the primary visual cortex ($p = 0.0331$), and even
more in the retrosplenial dysgranular cortex ($p = 0.001$). Indeed, when looking at the
amplitude maps, oscillations occurred not only near the injection site but also spanned
neighboring cortical areas, as shown in Fig. 1I. Functional connectivity variations between
LPS and vehicle were also assessed for each time point (Fig. 1J). No difference was found
during the first imaging session, 1h after injection (left diagonal). However, 48 h later (right
diagonal), functional connectivity after LPS injection was significantly different ($F (1,150) =$
27.51, $p < 0.0001$), with an increase between the secondary visual and the retrosplenial
dysgranular cortices ($p = 0.0374$). Hence, the increase of functional connectivity was
observed between highly oscillating cortical areas and neighboring areas.
468
LPS-induced hemodynamic waves are traveling through the cortex and subcortical
regions.

Next, we examined the dynamic changes occurring during inflammatory hemodynamic
oscillations at 48 h after LPS injection. When measuring the dynamic functional connectivity
between the left and right primary visual cortices, we observed an alternation between periods
of hyper- and hypo connectivity towards extreme values (Fig. 2A) that were not observed at 1
h post-injection (Fig. 2B). A closer look at the individual maps of relative changes of CBV
revealed that this was associated with the presence of hemodynamic traveling waves with
variable spatial trajectories (See also Supplementary Movie 1). For illustration of this
phenomenon, three representative time periods from Fig. 2A are displayed in terms of relative
CBV maps (Fig. 2C), reflecting high correlation (time-period 1), no correlation (time-period
2) or anticorrelation (time-period 3) in one typical animal. Period 1 was associated with a fast
bilateral cortical wave with successive decrease and increase of CBV that also involved
subcortical regions. Period 2 involved unilateral waves with a high amplitude event that
started from the left cortex to propagate to the left dorsal and then ventral hippocampus, while
period 3 ended with a wave that traveled from the left hippocampus to the right retrosplenial

and then right visual cortex, leading to the anticorrelation. We then used a peak detection software dedicated to the analysis of traveling events in stacks of 2D images to automatically identify CBV increases in our data and understand the trajectories of traveling waves (see Methods). The delay maps of all detected events (that is, local peaks of CBV) were classified in terms of spatial location and horizontal or vertical directionality by k-means clustering, enabling to reveal six different patterns of propagation (Fig. 2D). The traveling waves mainly involved adjacent penetrating vessels in the cortex, either from the retrosplenial cortex to the visual cortex (medio-temporal) or in the reverse direction (temporo-medial), in one hemisphere or bilaterally, or sometimes from one hemisphere to another (trans-hemispheric). In some cases, waves traveled from the cortex to the hippocampus (cortico-hippocampal) or reversely (hippocampo-cortical), or from the cortex to other subcortical regions (cortico-subcortical). This analysis also enabled to globally quantify the amount of traveling waves and their properties. Interestingly, dynamic traveling events were already observed in the vehicle group or at only 1h. However, a significant increase in the relative occurrence of such events as compared to the total number of events was found in the LPS group 48 h post-injection, as compared to the vehicle (+25.5%, p = 0.0066; Fig. 2E). The peak amplitude (+2.2%, p = 0.0269, Fig. 2F) and the propagation distance (+2.6 mm, p=0.0313, Fig. 2G) of the traveling waves were also significantly increased at 48 h post-injection of LPS.

As we individually classified each event, we studied the frequency of occurrence of events depending on their pattern of propagation (Fig. 2H). The medio-temporal or temporo-medial waves, either in one hemisphere or bilaterally, were found in LPS and VEH groups at both time points. However, their occurrence was always higher at 48 h as compared to 1 h for all groups, leading to significant differences in terms of global patterns representation. The distribution of the patterns was also significantly different between LPS and VEH at 48 h, with an increase in the relative proportion of cortical events and the appearance of propagations between cortical and subcortical areas in both directions (VEH: 4.2%, 4.1%, 0.1% and 91.6% of medio-temporal, temporo-medial, transhemispheric and unclassified/static events, respectively; LPS: 5.3%, 4.5%, 1.4%, 1.2%, 0.7%, 1.5% and 85.4% of medio-temporal, temporo-medial, transhemispheric, cortico-subcortical, cortico-hippocampal, hippocampo-cortical and unclassified/static events, respectively; $F_{(6,70)}$ = 4.2, p = 0.0012, interaction between patterns factor and group factor).

To summarize, the hemodynamic oscillations around 0.1 Hz displayed propagational properties, reflected by an increase of occurrence of traveling waves in the LPS group that

518 were also of higher amplitude and longer distance of propagation. They were mainly located

519 in the cortex, not only in pial vessels but largely involving vertical penetrating vessels. They

520 also occasionally involved the hippocampus or other subcortical regions, but only in the LPS

521 group.

Reconfiguration of resting hemodynamic activity towards oscillatory modes following

inflammation

To further characterize the contribution of these oscillations on the global hemodynamic

fluctuations, a principal component analysis (PCA) was performed on the mean covariance

matrix for each group (see Methods). Compared to the VEH group (Fig. 3A-B) and the

immediate effects after 1 h (Fig. 3C), the injection of LPS alone produced at 48 h spatial maps

of principal components with highly fragmented cortical regions, characterized by small,

complex patches with correlated or anticorrelated activity, indicating a fine-grained functional

subdivision of the cortex (Fig. 3D). The frequency spectra of the temporal signature of the

components for this specific condition showed frequency peaks between 0.08 and 0.15 Hz

(Fig. 3D), as observed previously with the raw Power Doppler signal in these regions.

Strikingly, at 48 h post-injection, the first eight components alone were sufficient to explain

more than 99.5% of the variance for the LPS group, as opposed to the VEH group or to the

early time-point in which more than 1000 components were required (Fig. 3E), highlighting

the importance of this phenomenon. In other words, the resting state activity under

inflammation could be fully described by these large-amplitude hemodynamic oscillations

occurring at a handful of distinct spatial nodes with their own frequency spectra.

Hemodynamic waves in neuroinflammatory conditions are related to the reactivity of

glial cells

As LPS-induced inflammation is known to heavily modify the morphology and function of

microglia and astrocytes (that become "reactive" in such conditions), we decided to

investigate the relationship between glial cells and inflammatory vasomotion. For studying

microglial reactivity, we used a new data-driven approach that relies on principal component

analysis of multiple morphological features of segmented microglial cells[40], visualized by Iba-

1 immunofluorescence on cerebral slices of the same animals that were used for fUS imaging.

For the astrocytic contribution, we chose a different strategy since astrocytes' morphology is

very complex and usual astrocytic protein markers are not able to label the numerous finest

processes that would be needed for assessing the overall cell morphology. Instead, we used

additional experimental groups that relied on different intracerebral injections of gliotoxins
that are known to induce specific lesions of astrocytes through different mechanisms, namely
alpha-amino-adipate (AAA, which acts by blocking the glutamine synthase[42]) and
fluorocitrate (FC, which blocks the Krebs cycle after selective uptake in astrocytes[43,44]). These
gliotoxins were injected alone or in addition to LPS before fUS imaging to evaluate the
influence of astrocytes on the hemodynamic oscillations, before sacrificing the animals.

Using relevant immunostainings, we first verified the specificity of astrocytic lesions and
absence of obvious toxicity on other cell types in those groups, as shown in Supplementary
Fig. 1A. The staining intensity of GFAP (an astrocytic protein upregulated in reactive
astrocytes) was more pronounced in the LPS group compared to the vehicle injection,
although non significantly ($p = 0.1740$). Conversely, the groups that received the gliotoxin
FC alone or in addition to LPS displayed significantly reduced GFAP staining (FC: $p = 0.0050$; LPS+FC: $p = 0.0006$), whereas the gliotoxin AAA alone did not inhibit GFAP
0.0050; LPS+FC: $p = 0.0006$), whereas the gliotoxin AAA alone did not inhibit GFAP
expression significantly ($p = 0.826$), but did when combined with LPS (LPS+AAA: $p < 0.0001$). Although GFAP is often used as an astrocytic marker, it is not recommended alone to
0.0001). Although GFAP is often used as an astrocytic marker, it is not recommended alone to
assess overall astrocytes density, as it is overexpressed in reactive state, and only few
astrocytes express GFAP in cortical grey matter. For these reasons, we also performed
NDRG2 (a marker of cortical astrocytes[45]) labelling. All groups receiving either only
gliotoxins or a gliotoxin in co-injection with LPS showed significant decrease of NDRG2
intensity (Supplementary Fig. S1B - AAA, $p = 0.0113$; FC, $p = 0.0002$; LPS+AAA, $p = 0.0226$; LPS+FC, $p = 0.0018$) in comparison to the vehicle group. These observations
0.0226; LPS+FC, $p = 0.0018$) in comparison to the vehicle group. These observations
confirmed an astrocytic lesion at the injection site by AAA or FC. In contrast, no significant
modification of NDRG2 level was observed after LPS administration alone, indicating that
LPS did not induce an astrocyte loss ($p > 0.9999$). Furthermore, no significant difference in
neuronal density (NeuN+ cells counting) was observed between groups (Supplementary Fig.
S1B; LPS, $p = 0.0569$; AAA, $p = 0.5120$; FC, $p = 0.9986$; LPS+AAA, $p = 0.9464$; LPS+FC, $p = 0.1213$); suggesting that the injections did not induce neurotoxicity in our models. Finally,
= 0.1213); suggesting that the injections did not induce neurotoxicity in our models. Finally,
we also analyzed the expression of the microglial marker Iba-1, that remained similar to
vehicle in most groups (Supplementary Fig. S1B ; LPS, $p = 0.9996$; AAA, $p = 0.3923$;
LPS+FC, $p = 0.6743$) but was slightly decreased in the FC and LPS+AAA groups (FC, $p = 0.0445$; LPS+AAA, $p = 0.0283$), suggesting that global density of microglia was unchanged
0.0445; LPS+AAA, $p = 0.0283$), suggesting that global density of microglia was unchanged
by LPS (consistently with an expression of Iba-1 in both activated and quiescent microglial
cells[46]) but were occasionally slightly lowered by the astrocytic toxins.

Then we focused on the morphology of microglial Iba-1+ cells. As expected, the extensive

morphology analysis showed a switch from a ramified to an ameboid morphology 48h

following LPS injection (Fig. 4A, segmented cells in red, as compared to the vehicle in blue).

Numerous parameters (see [40] for the definition of all parameters) such as the perimeter area

ratio (p = 0.0018), the ramification index (p = 0.0065), the processes/cell areas ratio (p =

0.0010), the skeleton/processes ratio (p = 0.0104), the polarization index (p = 0.0001), the

lacunarity (p = 0.0247), the density (p = 0.0162), the branching index (p = 0.0203), the inertia

(p = 0.0107), the processes/soma areas ratio (p = 0.0162), the convexity (p = 0.0324), the

convex hull radii ratio (p = 0.0076), the endpoints/branch points ratio (p = 0.0078) and the

fractal dimension (p = 0.0241) were significantly decreased 48h after LPS injection as

compared to vehicle injection, while the circularity (p = 0.0006), the roundness factor (p =

0.0004), the convex hull circularity (p = 0.0004), and the solidity (p = 0.0006) were increased

(Fig. 4B and Supplementary Fig. S1C). The features span ratio and linearity showed no

significant differences. The other additional groups that received astrocytic toxins were also

assessed for microglial morphology (Fig. 4A-B). No significant morphological differences

with control condition were observed 48h after the injection of each astrocytic toxin alone.

When gliotoxins were co-injected with LPS, morphological differences were still observed

compared to vehicle, although they were slightly less important than in the LPS alone group.

Indeed, the perimeter area ratio (LPS+AAA: p = 0.0332; LPS+FC: p = 0.0141), the

ramification index (LPS+AAA: p = 0.0488; LPS+FC: p = 0.0282), the processes/cell areas

ratio (LPS+AAA: p = 0.0340; LPS+FC: p = 0.0073), the skeleton/processes ratio (LPS+AAA:

p = 0.0448; LPS+FC: p = 0.0275), the polarization index (LPS+AAA: p = 0.0119; LPS+FC: p

= 0.0041), the circularity (LPS+AAA: p = 0.0265 ; LPS+FC: p = 0.0079), the roundness

factor (LPS+AAA: p = 0.0282; LPS+FC: p = 0.0059), and the solidity (LPS+AAA: p =

0.0169; LPS+FC: p = 0.0056) were modified in both the LPS+AAA and LPS+FC groups.

Other features were significantly different only for the LPS+FC group, such as convex hull

circularity (p = 0.0216), branching index (p = 0.0454), endpoints/branch points ratio (p =

0.0129) and processes/soma areas ratio (p = 0.0379) (Fig. 4B and Supplementary Fig. S1C).

Finally, Andrew's score ranking, which integrates multiple parameters that characterized best

our dataset, was used to classify each microglial cell from the different animals in terms of

reactivity (a negative score depicts a round-shape morphology, typical of the reactive state,

whereas a positive score depicts a well-ramified, basal morphology – see Fig. 4C for the

relative frequency of microglia in each group as a function of their Andrew's score). The

frequency of reactive microglia was much higher in the LPS alone as compared to vehicle and

to any other group. A slight increase of microglial reactivity was found in the FC group, while a moderate increase was found for the LPS+gliotoxins groups, as shown by the mean of Andrew's scores between the groups (Fig. 4D ; $F_{(5,20)}$ = 5.377, p = 0.0027).

As fUS imaging was performed for all animals of these different groups, the microglial morphological changes could be correlated with the amplitude of hemodynamic oscillations for all animals that received the different injections. Interestingly, the Andrew's score of overall cell morphology was negatively correlated with the increase of the frequency amplitude of 0.08-0.15 Hz range in the same animals (Fig. 4E; rho: -0.445; p = 0.023). In other words, as negative values of Andrew's score are reflective of round-shaped morphology, the rats that displayed the most reactive microglial cells showed higher amplitudes of vasomotion. Significant correlations were also observed for many individual morphological parameters in addition to the global Andrew's score (results not shown). Moreover, focusing on individual groups, the injection of gliotoxins alone did not significantly impact the amplitude of vasomotion compared to the control animals in any region at both time points (Fig. 4F-G), but the LPS-induced increase of amplitude at 48h was fully prevented by AAA and reduced by FC co-injection (Primary visual cortex: LPS+AAA, p = 0.9976; LPS+FC, p = 0.2099; Retrosplenial dysgranular cortex: LPS+AAA, p = 0.9597; LPS+FC, p = 0.0541). This suggests that the inflammatory vasomotion in the LPS group alone also depends on the local presence of astrocytes. Accordingly, no more increase of resting-state functional connectivity was found at 48h post-injection of LPS when it was co-injected with AAA or FC, while none of these molecules alone produced any effect on functional connectivity at 1h or 48h post-injection (Fig. 4H). When comparing the contributions of hemodynamic oscillations to the overall signal changes in LPS with the other groups that received the astrocytic toxins using PCA, here again only the variance from the LPS-48h group was fully described by less than 10 components, whereas FC or AAA alone or combined with LPS required about 1000 components to explain the same amount of variance, similarly to the VEH group (Fig. 4I). In parallel, the spatial maps of the components from the different gliotoxins groups (see Supplementary Fig. S2) displayed broader, more globally coherent activity patterns than LPS alone, with entire cortical areas showing correlation or anticorrelation with subcortical structures, and occasionally, inter-hemispheric anticorrelation (i.e. component 3 for the FC group at 1 h and 48 h – Supplementary Fig. S2C-D). Interestingly, the injection of gliotoxins alone or with LPS tended to produce acute subcortical effects, particularly in the hippocampus, that are visible on the spatial maps of the components at 1 h post-injection (i.e.

components 3 for AAA, 2 for FC, 1 for LPS+AAA and 2 for LPS+FC – Supplementary Fig.
S2A, C, E and G, respectively). In terms of frequency properties of the components, only
certain components of the LPS+FC group still showed frequency peaks around 0.1 Hz at 48 h.
We also noticed that the gliotoxins tended to produce irregular very low frequency
fluctuations at the acute phase, 1 h post-injection, as showed in the temporal signature of
some components in Figure S2 (i.e., components 3 to 5 of AAA, 2 to 3 of FC, and 1 and 2 of
LPS+FC). Taken together, these PCA results show that LPS oscillations at 48h were
respectively attenuated or fully blocked by FC and AAA, whereas these toxins alone produced
distinct hemodynamic effects by themselves that were most obvious at 1h.

When analyzing the properties of hemodynamic changes as traveling waves, it also appeared
that LPS combined with AAA or FC was no more inducing any significant change in the
proportion (LPS+AAA, p = 0.9833, LPS+FC, p = 0.2480), peak amplitude (LPS+AAA, p =
0.9987, LPS+FC, p = 0.3411) or propagation (LPS+AAA, p > 0.9999, LPS+FC, p = 0.7374)
of such waves (Fig. 4J), with no effect of the gliotoxins alone. The combination of AAA or FC
with LPS was also found to restore the normal profile of traveling waves spatial patterns (Fig.
4K), that is, mostly medio-temporal or temporo-medial waves. We still observed an overall
difference of waves repartition with less unclassified events for all groups at 48h compared to
1h, and FC alone was significantly different from VEH at 48h due to the appearance of some
waves involving the subcortical structures or transhemispheric propagation (5.6%, 3.7%,
0.3%, 0.06%, 0.06%, 0.06% and 90.2% of medio-temporal, temporo-medial,
transhemispheric, cortico-subcortical, cortico-hippocampal, hippocampo-cortical and
unclassified/static events, respectively; $F_{(6,63)}$ = 2.624, p = 0.0247).

To summarize, the injection of toxins that induce lesions of astrocytes, in combination with
LPS, was followed by an almost complete normalization of the inflammatory vasomotion, as
shown by the different parameters that we assessed (amplitude of oscillations, functional
connectivity, traveling waves proportion and trajectories, and variance explained by principal
components). Moreover, when taking into account all animals from the different groups, a
correlation was found between the microglial morphology changes and the amplitude of
vasomotion. Altogether, this suggest that microglia and astrocytes both participate (and most
probably, interact together) in this hemodynamic phenomenon.

Kinetics of hemodynamic waves induction after LPS exposure.

To understand the kinetics of the oscillatory effect resulting from LPS injection, the amplitude

of 0.08-0.15 Hz frequencies was evaluated at 1- and 7-days post injection in addition to the 1h

(0 day) and 48h (2-days) groups (Fig. (5A). The animals without injection (wI; only with a

skull thinning) did not differ from animals measured 1 h after LPS injection (primary visual

cortex, p = 0.6835; secondary visual cortex, p = 0.8632; retrosplenial dysgranular cortex, p =

0.8976, retrosplenial granular cortex part A, p = 0.9617; retrosplenial granular cortex part B, p

> 0.9999; superior colliculus, p = 0.9952). As expected, 48 h after injection, the amplitude of

the frequency range of interest was increased in the primary visual cortex (1 h, 0.057 ± 0.002;

48 h, 0.078 ± 0.002; p = 0.0001), the secondary visual cortex (1 h, 0.056 ± 0.001 ; 48 h, 0.070

± 0.007, p = 0.0320) and retrosplenial dysgranular cortex (1 h, 0.057 ± 0.003; 48 h, 0.083 ±

0.024; p < 0.0001), compared to 1 h post-injection. The increase was already present in the

primary and secondary visual cortices, 24 h after the intracerebral injection of LPS (primary

visual cortex: 24 h, 0.086 ± 0.026, p< 0.0001; secondary visual cortex: 24h, 0.079 ± 0.024, p

= 0.0025), and was no longer different from baseline state 7 days after injection (primary

visual cortex: 7 d, 0.061 ± 0.003, p = 0.9509; secondary visual cortex: 7 d, 0.059 ± 0.0004 , p

= 0.9917 ; retrosplenial dysgranular cortex: 7 d, 0.061 ± 0.002, p = 0.9627). Therefore, the

appearance of hemodynamic oscillations was present in the earlier phase of

neuroinflammation and was no more observed after one week.

To gain an in-depth overview of the temporality of the intracerebral LPS model at the cellular

level, we performed large-scale single-nuclei profiling from the microdissected visual cortex

sampled at 2 days or 7 days following LPS injection, or at 2 days following vehicle injection.

Following quality controls, we obtained the transcriptome of 64 471 nuclei by snRNA-seq

using PARSE Biosciences split-seq protocol allowing multiplexing. Two brain samples from

different individuals were used for each condition and these duplicates were analyzed

independently for each condition (Supplementary Fig. S3). Clustering revealed 5 distinct

clusters which corresponded to the brain major cell classes (Suppl Fig. S3A). Cell cluster

identification was based on the detection of landmark cell type markers, such as *Rbfox3* for

neurons, *Aqp4* for astrocytes, *Sox10* for oligodendroglial lineage cells (OLs), *Ptprc* for

immune cells and *Cdh5* for vascular cells. The three conditions grouped well together at the

general population level (Suppl Fig. S3B) with overall similar proportions of cells but

respectively lower and higher relative frequencies of neurons and immune cells at 7d (Suppl

Fig. S3C). However, *de novo* clustering of only astrocytes resulted in two different clusters of

reactive and homeostatic astrocytes, exclusively enriched in LPS astrocytes at 48h *versus*

control VEH astrocytes and LPS astrocytes at 7d, respectively (Fig. 5B,C). Reactive

astrocytes over-expressed *Gfap* and *Nes* (Fig. 5C) and genes enriched in wound healing,

gliogenesis, glial cell differentiation and response to axon injury biological processes

determined using gene ontology (GO) terms enrichment analysis (Suppl. Fig S3D), as

compared to vehicle. These enrichments were specific to the reactive astrocytes at 48h and

were no more found at 7d, consistently with the UMAP showing that the LPS-7d astrocytes

cluster was located much closer to the control cluster, suggesting a peak of reactivity around

two days and a return to a state closer to basal astrocytes (Fig. 5B). The direct comparison of

astrocytes at 48h post-LPS versus 7 days showed the significant enrichment of the same

biological processes (wound healing, gliogenesis, glial cell differentiation and response to

axon injury biological processes) as with vehicle injection. The expression of the 2 markers

that best described this transient activation at 48h (*Gfap* and *Nes*) went back to normal at 7

days (Fig. 5C). Correlation heatmap between samples confirmed that the whole transcriptional

signature of astrocytes from the VEH group correlates more with LPS-7d astrocytes than

LPS-48h astrocytes (Fig. 5D). We observed a similar kinetic pattern in Aif1-expressing

(Iba1□) microglia, although reactive microglia persisted at LPS-7d (Suppl. Fig. S3E).

However, further investigation is needed due to the low number of identified microglial nuclei

(as compared to the global number of immune cells, with extensive recruitment of peripheral

macrophages at LPS-48h).

Sustained neurovascular coupling in presence of oscillations.

To gain further insights into the functional properties of hemodynamic changes under

inflammation, we also evaluated the functional hyperemia changes during visual stimuli (Fig.

6A), which were performed immediately after the resting-state session at both time points

(Fig. 1A). As expected in this coronal plane, typical increases of CBV were found in the

primary visual cortex and the superior colliculus during the light flickering (Fig. 6B-C). At

both 1 h (Fig. 6D-F) and 48 h post-injection (Fig. 6G-I), the response to visual stimuli showed

globally no difference between VEH and LPS in the different visual areas. Diving into

temporal properties, the time-course of the response at 1h after injection was virtually

unchanged compared to vehicle (Fig; 6E), but the oscillations that we observed during

resting-state persisted during the visual stimulation at 48h post-injection (Fig. 6H).

Interestingly, despite these oscillations, the mean amplitude of the response and the rising and

decaying slopes of the curve appeared similar to the vehicle injection, suggesting a sustained

neurovascular coupling in this particular state. In fact, in the LPS group specifically, the

amplitude of the visual response tended to be higher in animals that displayed the strongest

basal oscillations, (as shown in Fig. 6J and in Supplementary Fig. S4A), and there was a

significant positive correlation between the fluctuations at 0.08-0.15 Hz during resting-state

and the mean amplitude of the functional hyperemia (rho = 0.6084, p = 0.0399; Fig. 6K). This

suggests that inflammatory vasomotion shares some common but non-competitive molecular

mechanisms with functional hyperemia.

We also studied the functional hyperemia in the groups that received AAA and FC

(Supplementary Fig. S4B-E). The gliotoxins alone had some effects at 1h: AAA led to a

modest increase of the visual response in the primary visual cortex at 1h (some time points

were significantly higher compared to vehicle from 21 to 30.2 s after the beginning of the

stimulation period; Supplementary Fig. S4B), and a global higher response in the superior

colliculus (VEH, 4.47% ± 1.85; AAA, 9.88% ± 6.02, p = 0.0012; Supplementary Fig. S4D).

For the FC group, the response shape in primary visual cortex was dramatically changed 1 h

after the injection, from the usual "plateau"-shape to a dome shape with a much higher

amplitude of the response (from 21.2 to 33.2 s after the beginning of the stimulation period;

Suppl Fig. S4B). This modification of the response shape was also present after the co-

injection of LPS+FC. No obvious effects of gliotoxins, either alone or combined with LPS,

was observed when compared to VEH at 48H (Suppl Fig. S4C, E). A slight decrease of visual

responses was observed in the primary visual cortex after co-injection of LPS+AAA (from

23.8 to 25 s after the beginning of the stimulation period; Suppl Fig. S4C). To summarize,

both gliotoxins acutely impacted the response to visual stimuli in rats, but with different

features, notably a strong perturbation of the temporal response of visual cortex following FC

injection. In agreement with the previous observations in resting-state conditions, the co-

injection of the gliotoxins with LPS prevented the appearance of the vascular oscillations

during the visual stimulation 48 h post-injection (as seen in Suppl Fig. S4C,E).

Hemodynamic waves are not associated to a change in blood flow velocity in cortical

vessels.

As the hemodynamic oscillations may be related to a global change in vessel tone, we also

evaluated quantitative hemodynamic parameters to estimate the level of cerebral perfusion in

the vessels using Ultrasound Localization Microscopy (ULM) sequences performed in the

same animals (Fig. 7), 48 h after intracerebral injections, immediately after the second fUS

imaging session (Fig. 1A). In addition to the density of bubbles (Fig. 7A), their absolute

velocity (Fig. 7B) and their direction (Fig. 7C) can be estimated, enabling to identify different

vascular compartments (pial vessels, arterioles and venules) in the cortex (Fig. 7D). The

microbubbles density and velocity were globally unchanged in the different compartments

following LPS injection, although the density tended to be slightly higher in arterioles (Fig.

7E-F). This enables to conclude that the appearance of vascular oscillatory waves was not due

to a significant change of the cerebral blood flow absolute velocity or microbubbles density,

suggesting that the vascular tone was unchanged compared to the control group.

The gliotoxins' influence on cerebral perfusion were also assessed, as displayed in Suppl. Fig

S5. Both AAA and FC, either alone or coinjected with LPS, had minimal impact on the

cerebral perfusion at 48h post-injection, with the exception of a significant increase in the

penetrating arterioles from the LPS+AAA group (VEH: 15.5 ± 2.1; LPS+AAA: 21.8 ± 4.0, p

= 0.0074).

Confirmation of the LPS-related hemodynamic oscillations in awake rats.

As anesthesia could potentially interact with LPS effects on hemodynamic activity, additional

rats were imaged in awake freely moving condition using skull implants for probe fitting after

progressive habituation (Fig. 8A). For each acquisition, a continuous temporal window of 120

sec was selected based on the absence of any movement artefact (Fig. 8B). Like in

anesthetized animals, awake rats showed an increased functional connectivity 48 h after LPS

injection between different visual areas (Fig. 8C - primary and secondary visual cortices: Δz-

score = 1.126, p = 0.0083; primary visual and retrosplenial dysgranular cortices: Δz-score =

0.8781, p = 0.0448; secondary visual and retrosplenial dysgranular cortices: Δz-score =

0.9915, p = 0.0215). As previously, the PCA showed a reconfiguration after 48 h post-LPS

with a very low number of components that could explain the signal variance, as opposed to

the baseline measurements or after 1 h (Fig. 8D). Furthermore, the frequency amplitude

related to the oscillations were also significantly increased after 48 h compared to the baseline

or 1 h post injection in the same regions (Fig. 8E), namely in the primary visual cortex

(Baseline: 0.064; 1 h: 0.062; 48 h: 0.080, p = 0.0015), secondary visual cortex (Baseline:

0.061; 1 h: 0.061; 48 h: 0.078, p = 0.0006) and retrosplenial dysgranular cortex (Baseline:

0.064; 1 h, 0.061; 48 h, 0.076, p = 0.0088). An example of long-distance traveling waves in an

awake rat can be visualized in Supplementary Movie 2.

Discussion

Cerebral hemodynamics have been extensively used as a correlate of brain function in animals

and humans. Although the interpretation of functional neuroimaging data focuses mainly on

neuronal activity and the influence of non-neuronal phenomena on blood flow could be

considered a drawback of such techniques[47], we argue that with careful interpretation,

cerebrovascular dynamics have the advantage to provide an integrative picture of the

interactions between the different components of the brain. Indeed, neuronal, glial and

vascular cells all critically interact to regulate the CBF fluctuations, both at rest and during

functional hyperemia. In this regard, understanding how inflammation locally shapes cerebral

hemodynamics is essential as this phenomenon is observed in many brain pathologies and

drastically modifies the functional properties of the different components of the

neurogliovascular unit (such as the induction of microglial and astrocytic reactivity). In

addition to providing new elements for functional neuroimaging interpretation in patients, this

may also lead to the identification of biomarkers or therapeutic perspectives.

Our main finding relates to the reorganization of cerebral hemodynamic activity towards an

oscillatory state characterized by quasi-sinusoidal, high-amplitude oscillations at ∼0.1 Hz, that

locally propagate through the vascular tree as traveling waves under inflammation, as early as

24 h post-injection of LPS. This oscillatory mode is transient and is no longer detectable after

seven days. Other studies have reported such sinusoidal hemodynamic oscillations at similar

frequencies occurring sporadically in animals or humans under anesthesia[48,49] or in the awake

state[50], but this has not yet been reported under the controlled induction of inflammation. The

observed frequency is typical of two distinct phenomena, namely vasomotion and "Mayer

waves". The latter is due to regular fluctuations in blood pressure that can occur at around 0.1

Hz, which could obviously affect CBV changes[51]. However, as noted by other authors, such

waves should be observed in all vessels simultaneously, whereas we and others have found

these oscillations to be localized in discrete regions of the vasculature, propagating through

the cortex - and in our case, occasionally between the cortex and subcortical regions - at a

speed between 0.1 and 0.8 mm/s (previous reports[49,50] and our own findings). Therefore, the

most plausible phenomenon is vasomotion[52], i.e. an autonomous and regular change in vessel

diameter at about 0.1 Hz, of myogenic origin (through intracellular calcium fluctuations in

smooth muscle cells[53]), distinct from the pulsatility due to the heartbeat. This is compatible

with traveling waves, as vasomotion can propagate along the vessel walls thanks to gap-

junctions and shear stress mechanisms[54]. A novel finding here is that vasomotor waves may

also occur in subcortical areas and can travel long distances, for instance from the ventral

hippocampus to the cortex in the contralateral hemisphere, which can be easily detected using

fUSi as compared to previous studies describing hemodynamic oscillations using optical

techniques. Recently, Broggini et al. also reported long-wavelength traveling waves in the cortical arterioles of awake mice due to vasomotion. Our findings align with their observation that such waves can span the entire cortical mantle[24]. Additionally, we reveal that the waves along the horizontal plane can propagate not only along the pial vessels, but also through adjacent penetrating cortical vessels. This results in hemodynamic changes that can span the entire thickness of cortex while traveling through distinct areas, not just the superficial layers. Importantly, we show here that fUS is sensitive enough to track individual traveling waves, which calls for further mesoscopic studies of vasomotion in physiological or pathological conditions with this imaging modality.

Although vasomotion can occur in peripheral as well as cerebral vessels, it has previously been shown that cerebral vasomotion is further entrained by changes in neuronal activity in the gamma-band, and this vaso-neuronal coupling would be the main driver of resting-state connectivity[7]. Here, we confirm under pathological conditions that vasomotion can enhance homotopic functional connectivity. Whether this phenomenon was entrained by or independent from neuronal activity remains to be investigated. As the oscillatory phenomenon did not reduce the amplitude of response during functional hyperemia, it seems unlikely that local neuronal activity would be strongly disrupted to the point of generating such sinusoidal hemodynamic waves by itself, because a high basal neuronal activity in the absence of stimulation should result in a smaller difference of activity during the stimulation. Nevertheless, it is puzzling that the amplitude of functional hyperemia tended to be higher in the presence of strong vasomotion, and it suggests that the two phenomena can co-exist while engaging some common cellular and molecular mechanisms. The most plausible hypothesis to us is that vasomotion frequency is indeed controlled by internal vascular mechanisms and external neuronal activity, and vasomotion amplitude is further influenced by other players which are also involved in neurovascular coupling under functional hyperemia. Our data suggest that the missing piece of this puzzle is glial cells, particularly when they become reactive. In the same way, another study showed that non-reactive astrocytes are able to limit the amplitude of vasomotion by releasing prostaglandins during the constriction phase, which stimulate TRPV4 channels, increase Ca^{2+} in the terminal foot and activate COX-1[17]. As LPS has been shown to decrease the expression of COX-1 in reactive astrocytes[55], this could increase the amplitude of vasomotion in our study by limiting negative feedback exerted by prostaglandins. In addition, there may be other mechanisms that reactive astrocytes could use to directly enhance vasomotion rather than reduce it. Our findings also suggest that microglia

might be directly involved, although this remains speculative, and the possible molecular

mechanisms are currently unclear.

Other interesting points should be discussed in terms of cellular changes in our model.

Transcriptomics revealed a higher astrocytic reactivity at 48 h than at 7 days, with the

identification of two astrocytic subclusters corresponding to a more basal state, characterized

by moderate expression of *Gfap* and very low expression of *Nes*, including astrocytes from

the VEH group and the LPS group at 7 days post-injection, and a strongly reactive state found

only in the 48 h post-LPS group, with high expression of *Gfap* and *Nes*. This unbiased

approach is recommended to take into account the complexity and heterogeneity of glial cells

reactivity among different pathologies, as opposed to measuring a single marker[56–58].

Interestingly, nestin and GFAP are both intermediate filament proteins that are consistent with

the known hypertrophy and morphological alterations occurring in reactive astrocytes, but

Nestin increase has previously been reported in stroke and not following a peripheral injection

of LPS in mice[59], which indicates that the effects of LPS can vary greatly depending on the

administration method. *Nestin* has also been described as absent from normal astrocytes and

transiently expressed in reactive state, with a return to baseline levels at 7 days post-stroke[59],

which is similar to our own findings with LPS intracerebral injection in the rat cortex. Future

works with optimized preparation protocols for enrichment of gliovascular unit should enable

to better dissect the link between hemodynamic oscillations and astrocytic reactivity. An

additional observation is that the morphology of microglial cells was slightly less impacted in

the groups that received a co-injection of LPS and astrocytic toxins, suggesting that although

microglia are thought to be the primary target of LPS effect through activation of the TLR4

pathway (with subsequent release of cytokines that render astrocytes reactive), astrocytes may

further increase the reactivity of microglia through a bidirectional interaction during

inflammation[60,61]. This also illustrates the difficulty to isolate the specific contributions of

microglia *versus* astrocytes under neuroinflammatory conditions.

Regarding the effects of astrocytic toxins alone, we found that both tended to acutely increase

functional hyperemia in the rat, with different effects that could be related to their different

mechanisms of action. Surprisingly, AAA did not change the response in the visual cortex

where it was injected, but the amplitude of response in the superior colliculus was

significantly increased. The reason for this is unclear, although it might be possible that local

acute perturbations of astrocytic activity in the visual cortex, and secondary neuronal activity,

could alter inputs from the visual cortex to the superior colliculus[62]. The effect of FC was

much more pronounced and resulted in a change of the shape of the response, with a sustained
increase in CBV rather than a plateau at the end of the stimulation period. This is in line with
a previous study showing that astrocytes are particularly important for the late component of
functional hyperemia, observed during sustained sensory stimulation, such as in our own
conditions[63]. Indeed, chemogenetic activation of astrocytes enhances late hyperemia in awake
mice. We also found that FC induced such an effect, although we would expect it to reduce
the response, since the toxin is supposed to reduce the calcium concentration in astrocytes and
inhibit their activity[64]. This difference may be due to the use of anesthesia in our visual
stimulation experiments, or to an acute compensatory hyperexcitability of astrocytes *in vivo* as
opposed to the inhibitory effect of FC *in vitro*.

In terms of implications for functional imaging, the present study sheds new light on the
complexity of cerebral hemodynamics and the involvement of glial cells in the regulation of
cerebral blood flow. Such findings are important for understanding and interpreting the data,
particularly in studies of patients with brain diseases. While future work will be needed to
firmly demonstrate the causative influence of reactive glial cells, the present results suggest
that vasomotion may serve as a hemodynamic biomarker for glial cell reactivity. The fact that
sinusoidal cerebral vasomotion has also been sporadically reported in patients (without
insights into the cellular mechanisms) demonstrates the clinical relevance of the
phenomenon[49,50]. In practice, these results demonstrate that vasomotion and consequently
hemodynamic waves can represent a huge part of the data variance (up to 99% in the LPS
group). Whether frequencies above 0.08 Hz should be filtered out in fMRI resting-state
analysis has been vastly debated[8,65]; here we argue that biologically relevant phenomenon are
occurring at such frequencies, providing insights into the glial components as well as our
basic understanding of functional connectivity and hyperemia. Therefore, implementing
separate analyzes of this frequency band, or other approaches for characterization of traveling
waves[66] may increase the interpretability of future functional neuroimaging studies.

Finally, although vasomotion is still a poorly understood phenomenon, it has been suggested
to facilitate oxygen exchange between vessels and the tissues[52]. As hypoxia is frequently
observed in inflammatory conditions[67] and LPS can activate the transcription factor hypoxia-
inducible factor 1 (HIF-1)[68], the present effect could compensate neuronal hypoxia in
pathological conditions. Importantly, it has been shown that an "artificial" increase in cerebral
vasomotion by functional hyperemia improves the clearance of waste products along the
vessels[69] and several studies have emphasized that this should be a major driving force for

improving exchanges between the cerebrospinal (CSF) and interstitial (ISF) fluids, according

to the glymphatic system hypothesis[70] as well as the intra-mural peri-arterial drainage

hypothesis[71,72]. Although this remains speculative at this early stage, it is conceivable that in

inflammatory conditions, enhancement of vasomotion may improve CSF/ISF exchange to

enhance waste removal, or even facilitate the communication between the CSF/ISF and the

meningeal lymphatic system[73]. This may explain the reported paradoxical improvement in

soluble amyloid beta-peptide following intracerebral injection of LPS in the mice[74]. These

different possibilities suggest new therapeutic opportunities by modulating the cerebral

vasomotion in a pathological context such as neurodegenerative disorders, and deserve further

investigation.

Data availability

956 Data will be made available upon request to the authors, without undue reservation.

Acknowledgments

958 This work was funded by the ANR (LabCom "Neuroimaging for Drug Discovery") and the Auvergne-

959 Rhône-Alpes Region (Pack Ambition Recherche "Brain Imaging for Drug Discovery"). We thank the

960 CIQLE facility and Bruno Chapuis for the assistance with the slide scanner.

Author contributions

MP, MD, LZ and BV conceived the project. MP, MV and BV performed the imaging experiments and

surgeries. MP, LV and BV analyzed the fUS and ULM data. MP performed the immunofluorescence

staining and analysis. GM performed the single-nuclei RNA-seq experiment and analysis. SB, JCC

and OP designed and performed the analysis of microglia morphology. MD, LZ and BV supervised the

project. MP and BV wrote the first draft. MP, MD, LZ and BV reviewed the article. All authors

approved the final manuscript.

Declaration of interests

969 MD is full-time employee of Theranexus. BV was full-time employee of Theranexus during a part of

970 the project. The other authors declare no competing interests.

Figure titles and legends

Figure 1: fUSi reveals spontaneous hemodynamic oscillations in anesthetized rats following the

intracerebral injection of LPS. (A) Experimental timeline following the intracortical injection

procedure. Each imaging session (1h and 48h post-injection) included a first resting-state acquisition

followed by a second acquisition under visual stimulation. The second session was followed by an

Ultrasound Localization Microscopy acquisition and animals were euthanized immediately after for

brain sampling. (B) Mean of all registered fUSi images with overlay of the selected anatomical regions

of interest from the SIGMA atlas. (C) Representative time curves of spontaneous CBV changes in the

primary visual cortex of rats, 1h (dark blue) or 48h (light blue) following saline injection. (D) Same as

(C) following LPS injection. The mean amplitude spectra ($\pm$ SEM) after injection of LPS (red) or

vehicle (blue) are shown for 1h (E) and 48h (F) post-injection. (G) Frequency amplitude in 0.08-0.15

Hz range at 1 h post-injection. (H) Frequency amplitude in 0.08-0.15 Hz range at 48 h post-injection.

Two-way ANOVA followed by Sidak's multiple comparisons test. Each bar is the mean $\pm$ SEM. (I)

Mean spatial maps of frequency amplitude in 0.08-0.15 Hz range in the different groups, with a

threshold of fALFF>0.075. (J) Functional connectivity changes after 1 and 48 hours. Matrices of the

correlation coefficient variations of LPS compared to vehicle are shown (bottom-left for 1 h post-

injection, top-right for 48 h post-injection). For statistical comparisons, a Fisher Z-transformation was

applied to the correlation coefficients before two-way ANOVA followed by Sidak's multiple

comparisons test (*$p < 0.05$; **$p < 0.01$; ***$p < 0.001$, ****$p < 0.001$).

Figure 2: Spatial and temporal dynamics of CBV oscillations. (A) Dynamic functional connectivity

analysis in a representative individual, 48h after LPS injection. The same individual is shown in (B),

only 1h after LPS injection. The CBV changes over time in the left and right visual cortex are shown

in grey and black curves, respectively. The dynamic connectivity between the left and right visual

cortex is shown in the colored heatmap, using a sliding window of 6 seconds and a step of 0.4 second.

The hemodynamic oscillations are associated with small epochs of hyperconnectivity or

anticorrelation between cortical hemispheres. Corresponding changes in CBV, pixel-by-pixel, are

mapped during such periods in (C), showing that such changes in dynamic functional connectivity

reflect different patterns of hemodynamic traveling waves that can occur almost simultaneously across

the bilateral cortex (Period 1 of hyperconnectivity), unilaterally (Period 2 of low connectivity), or

sequentially between the two sides (Period 3 of negative connectivity). (D) Different patterns of

traveling waves identified across all acquisitions after clustering of peak events, showing that waves

could travel along different directions across the cortex and also propagate from subcortical areas to

the cortex, and reversely. Examples of delay maps and corresponding maps of CBV changes across time are shown for each pattern. (E) Percentage of dynamic traveling events identified as compared to the total number of events including static and localized increases of CBV, showing that traveling waves were observed more often in the LPS group 48h after injection. Each bar is the mean across individuals ± SEM. (F) Amplitude of the CBV increases for the traveling events, showing that the traveling waves were of higher amplitude in the LPS group 48h after injection. Each individual point is the mean for all dynamic events detected over the acquisition. (G) Propagation distance for the traveling waves, showing that the waves traveled on longer distances in the LPS-48h group. Each individual point is the mean for all dynamic events detected over the acquisition. Unpaired t-tests in (E), (F), (G). (H) Pie charts showing the proportion of each traveling pattern as compared to the total number of detected events. To better visualize the relative occurrence of each pattern, the proportions are also shown with inlays excluding the unclassified events. Two-way ANOVAs for between-groups comparisons and mixed-effect analyzes for between-time points comparisons.

Figure 3: Principal component analysis reveals oscillatory modes following LPS exposure. Spatial, temporal and spectral signature of the principal components in the vehicle group, 1h (A) and 48h (B) after injection, and in the LPS group, 1h (C) or 48h (D) after injection. Only the components with eigenvalues >0.1 are shown. The spatial and spectral signatures are obtained from the group averaging; the temporal profile is shown for representative individuals. Resonant modes can be visualized in the LPS 48h group with peaks in the frequency spectrum between 0.08 Hz and 0.15 Hz, cutting the cortex into distinct subterritories. (E) Scree plot showing the percentage of explained variance for each component. The ten first components in the LPS 48h group can explain more than 99% of the variance, contrary to the other groups.

Figure 4: Relationship between reactive glial cells and CBV oscillations. (A) Representative illustrations of segmented microglia for each group (vehicle, LPS, and/or gliotoxins injections). (B) Main parameters that best describe the microglia morphology in the dataset in the different groups. Two-way ANOVA followed by Dunnett's multiple comparisons test. Each bar is the mean ± SEM. (C) Distribution graph of Andrew's scores of microglial cells (that estimates the reactivity of microglia based on their morphology) in the different groups. (D) Bar plot of the global Andrew's score for each group. Two-way ANOVA followed by Dunnett's multiple comparisons test. Each bar is the mean ± SEM. (E) Scatter plot and linear regression showing the relationship between CBV oscillations in 0.08-0.15 Hz range and global Andrew's score in all individuals. (F) Frequency amplitude of CBV changes in 0.08-0.15 Hz range at 1 h post-injection for all experimental groups. (G) Frequency amplitude in 0.08-0.15 Hz range at 48 h post-injection for all experimental groups. Two-way ANOVA followed by Dunnett's multiple comparisons test. Each bar is the mean ± SEM. (H) Functional

connectivity changes after 1 and 48 hours. Matrices of the correlation coefficient variations of each group (that received gliotoxin alone or in combination with LPS) compared to vehicle are shown (bottom-left for 1 h post-injection, top-right for 48 h post-injection). For statistical comparisons, a Fisher Z-transformation was applied to the correlation coefficients before two-way ANOVA followed by Dunnett's multiple comparisons test (all non-significant). (I) Scree plot showing the percentage of explained variance for each component after PCA for all groups. The ten first components in the LPS 48h group can explain more than 99% of the variance, contrary to all the other groups that received gliotoxins alone or coinjected with LPS. (J) Percentage, amplitude and propagation of traveling waves for all groups at 1h and 48h post-injection. Each bar is the mean across individuals ± SEM. One-way ANOVAs followed by Dunnett's tests. (H) Bar charts showing for all groups the proportion of each traveling pattern as compared to the total number of detected events. To better visualize the relative occurrence of each pattern, the proportions are also shown excluding the unclassified events. Two-way ANOVAs for between-groups comparisons and mixed-effect analyzes for between-time points comparisons.

Figure 5: Kinetics of LPS effects on the hemodynamic oscillations and astrocytes reactivity. (A) Frequency amplitude in 0.08-0.15 Hz range in animals without any injection (wI), or at 1h, 24h, 48h (2d) or seven days (7d) after injection . Two-way ANOVA followed by Dunnett's multiple comparisons test. Each bar is the mean ± SEM. The oscillations were already present at 24h post-injection but disappeared after 7 days. (B) UMAP plot showing the distribution of astrocytes in the three conditions: 48h post-injection of VEH in green or LPS in red, and 7 days post-injection of LPS in yellow. Note that at 7 days post-injection of LPS, the molecular phenotype of astrocytes is much closer to the VEH group than at 48h post-injection. (C) Density plot showing the selective increase of GFAP and Nestin expression in the cluster that corresponds to LPS at 48h post-injection, but not in the cluster that corresponds to VEH and LPS-7d groups. (D) Correlation matrix showing the global correlation of gene expression between the different conditions, with higher correlation between LPS-7d and VEH than LPS-48h and VEH. LPS-48h was also markedly different from LPS-7d in terms of global gene expression.

Figure 6: Inflammatory spontaneous oscillations of CBV are associated with increased functional hyperemia during visual stimulations in anesthetized rats. (A) Schematic representation of the experimental setup during fUSi with visual stimulation. (B) Representative time curve of CBV changes in the primary visual cortex of rats following visual stimuli. (C) Mean map of relative CBV changes pixel-by-pixel in the VEH group 48 h post injection. The CBV changes during stimulation after 1h post-injection of LPS in shown as mean map of relative changes in (D), mean time curve in the primary visual cortex in (E) and global changes in the different regions of interest (mean ± SEM; Two-way ANOVA followed by Sidak's multiple comparisons correction) in (F). The same results are shown in (G), (H) and (I) for 48h postinjection of LPS. (J) Individual mean time curves (averaged

across the 4 blocks) for all animals in the LPS group, 48h after injection. (K) Scatter plot and linear regression showing the relationship between amplitude of oscillations during resting-state and the mean CBV changes during visual stimulation in the LPS group at 1h and 48h post-injection.

Figure 7: Measurement of cerebral perfusion at high resolution using ultrasound localization microscopy following LPS injection. (A) Representative image of microbubbles density. (B) Representative image of the microbubbles absolute velocity in the same individual. (C) Representative image of the microbubbles directional velocity along the vertical axis in the same individual (red = downward movement, blue = upward movement). (D) Magnification of the cortical area in (C), showing the identification of the pial vessels, penetrating arterioles and veinules based on the direction of the microbubbles. (E) Absolute velocity of microbubbles reflecting the cerebral perfusion in the different vessel compartments. Each bar is the mean ± SEM. Unpaired t-tests (non-significant). (F) Density of microbubbles in the different vessel compartments. Each bar is the mean ± SEM. Unpaired t-tests (non-significant).

Figure 8: Resting-state fUSi in freely-moving animals revealing hemodynamic oscillations in awake rats after LPS exposure. (A) Illustration showing the probe, probe holder and implant enabling fUSi in freely-moving rats. (B) Representative time-course of the Power Doppler signal in one animal over the 1200s period, enabling to visualize alternance between active periods leading to movement artifacts and quiet state periods, with analysis performed on the consecutive minutes devoid of movement artifact (as shown in the inlay). (C) Functional connectivity matrices 1h or 48h after LPS injection in awake rats. Two-way ANOVA followed by Dunnett's test. Similarly to anesthetized rats, a significant increase of cortical connectivity was observed at 48h. (D) Scree plot showing the percentage of explained variance for each component obtained after principal component analysis of the different groups (Baseline = without injection, or 1h/48h following LPS injection). Similarly to anesthetized rats, LPS was followed by a global reconfiguration toward oscillatory modes, with less than 10 components that were able to explain almost all the variance in the 48h group only. (E) Frequency amplitude in 0.08-0.15 Hz in awake animals, showing a significant increase of amplitude 48h after LPS injection. Two-way ANOVA followed by Dunnett's multiple comparisons test. Each bar is the mean ± SEM.

Supplemental information titles and legends

Figure S1: Validation of gliotoxins effects and additional characterization of glial reactivity. (A) Representative fluorescence microscopy images of perilesional site showing GFAP, Iba1, NDRG2 and NeuN expression in the different groups. (B) Iba, GFAP and NDRG2 expressions and NeuN-positive cells counting for each group. Two-way ANOVA followed by Dunnett's multiple comparisons test.

Each bar is the mean ± SEM. (C) Microglial morphology parameters. Bar plots of the parameters in

the different groups. Each bar is the mean ± SEM. All tests are two-way ANOVAs followed by

Dunnett's multiple comparisons test.

Figure S2: Principal component analysis in the gliotoxins groups. Spatial, temporal and spectral

signature of the principal components in the AAA group, 1h (A) and 48h (B) after injection, in the FC

group, 1h (C) and 48h (D) after injection, in the LPS+AAA group, 1h (E) and 48h (F) after injection,

and in the LPS+FC group, 1h (G) and 48h (H) after injection. Only the components with eigenvalues

>0.1 are shown. The spatial and spectral signatures are obtained from the group averaging; the

temporal profile is shown for representative individuals.

Figure S3: Single-nucleus RNA sequencing. (A) UMAP plot showing the spatial distribution of the

major cell classes within the dataset comprising of 70k nuclei (top left) and the expression of class

gene markers used to identify the clusters. (B) Spatial distribution of each of the three conditions

within the UMAP plot. (C) Barplots of cell classes distribution within each condition (VEH 48h, LPS

48h and LPS 7d). Note the increase of immune cells over the time after LPS. (D) Barplots showing the

enrichment of biological processes (BPs) in astrocytes after 48h post LPS injection versus the control.

(E) Barplots of microglia distribution within each condition (VEH 48h, LPS 48h and LPS 7d),

showing a maximal proportion of reactive microglia in the LPS 48h group.

1125

Figure S4: Additional observations regarding functional hyperemia during visual stimulations in

anesthetized rats. (A) Whole individual time curves for all animals in the LPS group, 1h and 48h

after injection. (B) Mean time curves and spatial maps of CBV changes during stimulation in the

gliotoxins groups, 1h after injection. For time curves, significant differences with controls are shown

by horizontal bars. Two-way repeated-measures ANOVAs followed by Sidak's multiple comparisons

tests (p < 0.05, mean ± SEM). (C) Mean time curves and spatial maps of CBV changes during

stimulation in the gliotoxins groups, 48h after injection. For time curves, significant differences with

controls are shown by horizontal bars. Two-way repeated-measures ANOVAs followed by Sidak's

multiple comparisons tests (p < 0.05, mean ± SEM). (D) Global CBV changes during stimulation in

the different regions of interest for all experimental groups (mean ± SEM; Two-way ANOVA followed

by Dunnett's multiple comparisons correction) at 1h post-injection. (E) Same as (D) but at 48h post-

injection.

Figure S5: Ultrasound localization microscopy results for all experimental groups. (A) Density of

microbubbles in the different vessel compartments. Each bar is the mean ± SEM. One-way ANOVAs

followed by Dunnett's test. (B) Absolute velocity of microbubbles reflecting the cerebral perfusion in

the different vessel compartments. Each bar is the mean ± SEM. One-way ANOVAs followed by

Dunnett's test.

Table S1: Complete list of antibodies used in the present study for the immunofluorescence stainings.

Video S1: Representative mapping of CBV changes pixel-by-pixel across 150 consecutives images,
during resting-state in an anesthetized animal, 48h after LPS injection. The raw images are displayed
on the left, and the reconstructed images following the first 10 group-derived principal components are
displayed on the right.

Video S2: Representative mapping of CBV changes pixel-by-pixel across 150 consecutives images,
during resting-state in an awake animal, 48h after LPS injection. The raw images are displayed on the
left, and the reconstructed images following the first 10 group-derived principal components are
displayed on the right.

1152

REFERENCES:

1.   Howarth, C., Mishra, A. & Hall, C. N. More than just summed neuronal activity: how multiple
cell types shape the BOLD response. *Philos. Trans. R. Soc. B Biol. Sci.* **376**, 20190630 (2021).

2.   Mishra, A., Hall, C. N., Howarth, C. & Freeman, R. D. Key relationships between non-invasive
functional neuroimaging and the underlying neuronal activity. *Philos. Trans. R. Soc. B Biol. Sci.* **376**,
20190622 (2021).

3.   Villringer, A. Understanding Functional Neuroimaging Methods Based on Neurovascular
Coupling. in *Optical Imaging of Brain Function and Metabolism 2* (eds. Villringer, A. & Dirnagl, U.) vol.
413 177–193 (Springer US, Boston, MA, 1997).

4.   Masamoto, K. & Vazquez, A. Optical imaging and modulation of neurovascular responses. *J.*
Cereb. Blood Flow Metab. **38**, 2057–2072 (2018).

5.   Liao, L.-D. *et al.* Neurovascular coupling: in vivo optical techniques for functional brain
imaging. *Biomed. Eng. OnLine* **12**, 38 (2013).

6.   Deffieux, T., Demené, C. & Tanter, M. Functional Ultrasound Imaging: A New Imaging Modality
for Neuroscience. *Neuroscience* **474**, 110–121 (2021).

7.   Mateo, C., Knutsen, P. M., Tsai, P. S., Shih, A. Y. & Kleinfeld, D. Entrainment of Arteriole
Vasomotor Fluctuations by Neural Activity Is a Basis of Blood-Oxygenation-Level-Dependent "Resting-
State" Connectivity. *Neuron* **96**, 936-948.e3 (2017).

8.   Lambers H, Wachsmuth L, Lippe C, Faber C. The impact of vasomotion on analysis of rodent
fMRI data. *Front Neurosci.* **17,** 1064000 (2023).

9.   Mishra, A., Gordon, G.R., MacVicar, B.A., Newman, E.A. Astrocyte Regulation of Cerebral
Blood Flow in Health and Disease. *Cold Spring Harb Perspect Biol.* **16**, 4 (2024).

10.   Nippert, A. R., Biesecker, K. R. & Newman, E. A. Mechanisms Mediating Functional Hyperemia
in the Brain. *The Neuroscientist* **24**, 73–83 (2018).

11.   Howarth, C. The contribution of astrocytes to the regulation of cerebral blood flow. *Front.*
Neurosci. **8**, (2014).

12.   Petzold, G. C. & Murthy, V. N. Role of Astrocytes in Neurovascular Coupling. *Neuron* **71**, 782–
797 (2011).

13.   Bazargani, N. & Attwell, D. Astrocyte calcium signaling: the third wave. *Nat. Neurosci.* **19**,
182–189 (2016).

14.   Lia, A., Di Spiezio, A., Speggiorin, M. & Zonta, M. Two decades of astrocytes in neurovascular
coupling. *Front. Netw. Physiol.* **3**, 1162757 (2023).

15.   MacVicar, B. A. & Newman, E. A. Astrocyte Regulation of Blood Flow in the Brain. *Cold Spring*
Harb. Perspect. Biol. **7**, a020388 (2015).

16.   Filosa, J. A. *et al.* Local potassium signaling couples neuronal activity to vasodilation in the
brain. *Nat. Neurosci.* **9**, 1397–1403 (2006).

17.   Haidey, J. N. *et al.* Astrocytes regulate ultra-slow arteriole oscillations via stretch-mediated
TRPV4-COX-1 feedback. *Cell Rep.* **36**, 109405 (2021).

18.   Császár E, et al. Microglia modulate blood flow, neurovascular coupling, and hypoperfusion
via purinergic actions. *J Exp Med* **219**, e20211071 (2022).

19.   Verkhratsky, A., Butt, A., Li, B., Illes, P., Zorec, R., Semyanov, A., Tang, Y., Sofroniew, M.V.
Astrocytes in human central nervous system diseases: a frontier for new therapies. *Signal Transduct*
Target Ther. **8**, 396 (2023).

20.   Boido D, et al. Mesoscopic and microscopic imaging of sensory responses in the same animal.
Nat Commun. **10**, 1110 (2019).

21.   Urban, A. *et al.* Real-time imaging of brain activity in freely moving rats using functional
ultrasound. *Nat. Methods* **12**, 873–878 (2015).

22.   El Hady, A. *et al.* Chronic brain functional ultrasound imaging in freely moving rodents
performing cognitive tasks. *J. Neurosci. Methods* **403**, 110033 (2024).

23.   Tiran, E. *et al.* Transcranial Functional Ultrasound Imaging in Freely Moving Awake Mice and
Anesthetized Young Rats without Contrast Agent. *Ultrasound Med. Biol.* **43**, 1679–1689 (2017).

24.   Broggini, T., Duckworth, J., Ji, X., Liu, R., Xia, X., Mächler, P., Shaked, I., Munting, L.P., Iyengar,
S., Kotlikoff, M., van Veluw, S.J., Vergassola, M., Mishne, G., Kleinfeld, D. Long-wavelength traveling
waves of vasomotion modulate the perfusion of cortex. *Neuron.* **24**, 324-6 (2024).

25.   Osmanski, B.-F., Pezet, S., Ricobaraza, A., Lenkei, Z. & Tanter, M. Functional ultrasound
imaging of intrinsic connectivity in the living rat brain with high spatiotemporal resolution. *Nat.*
Commun. **5**, 5023 (2014).

26.   Urban, A. *et al.* Chronic assessment of cerebral hemodynamics during rat forepaw electrical
stimulation using functional ultrasound imaging. *NeuroImage* **101**, 138–149 (2014).

27.   Bercoff, J. *et al.* Ultrafast compound doppler imaging: providing full blood flow
characterization. *IEEE Trans. Ultrason. Ferroelectr. Freq. Control* **58**, 134–147 (2011).

28.   Montaldo, G., Tanter, M., Bercoff, J., Benech, N. & Fink, M. Coherent plane-wave
compounding for very high frame rate ultrasonography and transient elastography. *IEEE Trans.*
Ultrason. Ferroelectr. Freq. Control **56**, 489–506 (2009).

29.   Demene, C. *et al.* Spatiotemporal Clutter Filtering of Ultrafast Ultrasound Data Highly
Increases Doppler and fUltrasound Sensitivity. *IEEE Trans. Med. Imaging* **34**, 2271–2285 (2015).

30.   Grandjean, J., *et al.* A consensus protocol for functional connectivity analysis in the rat brain.
Nat Neurosci. **4**, 673-681 (2023).

31.   Droguerre, M., Vidal, B., Valdebenito, M., Mouthon, F., Zimmer, L., Charvériat, M. Impaired
Local and Long-Range Brain Connectivity and Visual Response in a Genetic Rat Model of Hyperactivity
Revealed by Functional Ultrasound. *Front Neurosci.* **16**, 865140 (2022).

32.   Gesnik, M. *et al.* 3D functional ultrasound imaging of the cerebral visual system in rodents.
NeuroImage **149**, 267–274 (2017).

33.   Errico, C. *et al.* Ultrafast ultrasound localization microscopy for deep super-resolution vascular
imaging. *Nature* **527**, 499–502 (2015).

34.   Vidal, B. *et al.* Pharmaco-fUS in cognitive impairment: Lessons from a preclinical model. *J.*
Psychopharmacol. (Oxf.) **36**, 1273–1279 (2022).

35.   Vidal, B., Droguerre, M., Venet, L., Valdebenito, M., Mouthon, F., Zimmer, L., Charvériat, M.
Inter-subject registration and application of the SIGMA rat brain atlas for regional labeling in
functional ultrasound imaging. *J Neurosci Methods.* **355**, 109139 (2021).

36.   Barrière, D. A. *et al.* The SIGMA rat brain templates and atlases for multimodal MRI data
analysis and visualization. *Nat. Commun.* **10**, 5699 (2019).

37.   Song, X.-W. *et al.* REST: A Toolkit for Resting-State Functional Magnetic Resonance Imaging
Data Processing. *PLoS ONE* **6**, e25031 (2011).

38.   Cabral, J., Fernandes, F. F. & Shemesh, N. Intrinsic macroscale oscillatory modes driving long
range functional connectivity in female rat brains detected by ultrafast fMRI. *Nat. Commun.* **14**, 375
(2023).

39.   Wang, Y. *et al.* Accurate quantification of astrocyte and neurotransmitter fluorescence
dynamics for single-cell and population-level physiology. *Nat. Neurosci.* **22**, 1936–1944 (2019).

40.   Benkeder, S., Dinh, S.M., Marchal, P., De Gea, P., Thoby-Brisson, M., Hubert, V., Hristovska, I.,
Pitollat, G., Combet, K., Cardoit, L., Pillot, B., Léon, C., Wiart, W., Marty, S., Honnorat, J., Pascual, O.,
Comte, J.C. MorphoCellSorter: An Andrews plot-based sorting approach to rank microglia according
to their morphological features. *eLife.* **13**, RP101630 (2024).

41.   Yu, G., Wang, L.G., Han, Y., He, Q.Y. clusterProfiler: an R package for comparing biological
themes among gene clusters. *OMICS.* **5**, 284-7 (2012).

42.   Khurgel, M., Koo, A. C. & Ivy, G. O. Selective ablation of astrocytes by intracerebral injections
of ?-aminoadipate. *Glia* **16**, 351–358 (1996).

43.   Fonnum, F., Johnsen, A. & Hassel, B. Use of fluorocitrate and fluoroacetate in the study of
brain metabolism. *Glia* **21**, 106–113 (1997).

44.   Hassel, B., Paulsen, R. E., Johnsen, A. & Fonnum, F. Selective inhibition of glial cell metabolism
in vivo by fluorocitrate. *Brain Res.* **576**, 120–124 (1992).

45.   Zhang, Z. *et al.* The Appropriate Marker for Astrocytes: Comparing the Distribution and
Expression of Three Astrocytic Markers in Different Mouse Cerebral Regions. *BioMed Res. Int.* **2019**,
1–15 (2019).

46.   Boche, D., Perry, V. H. & Nicoll, J. A. R. Review: Activation patterns of microglia and their
identification in the human brain. *Neuropathol. Appl. Neurobiol.* **39**, 3–18 (2013).

47.   Figley, C. R. & Stroman, P. W. The role(s) of astrocytes and astrocyte activity in
neurometabolism, neurovascular coupling, and the production of functional neuroimaging signals:
Astrocyte control of fMRI and PET signals. *Eur. J. Neurosci.* **33**, 577–588 (2011).

48.   Rivadulla, C., de Labra, C., Grieve, K.L., Cudeiro, J. Vasomotion and neurovascular coupling in
the visual thalamus in vivo. *PLoS One.* **6**, e28746 (2011).

49.   Noordmans, H.J., van Blooijs, D., Siero, J.C.W., Zwanenburg, J.J.M., Klaessens, J.H.G.M.,
Ramsey, N.F. Detailed view on slow sinusoidal, hemodynamic oscillations on the human brain cortex
by Fourier transforming oxy/deoxy hyperspectral images. *Hum Brain Mapp.* **39**, 3558-3573 (2018).

50.   Rayshubskiy, A. *et al.* Direct, intraoperative observation of ~0.1 Hz hemodynamic oscillations
in awake human cortex: Implications for fMRI. *NeuroImage* **87**, 323–331 (2014).

51.   Julien, C. The enigma of Mayer waves: Facts and models. *Cardiovasc. Res.* **70**, 12–21 (2006).

52.   Aalkjær, C., Boedtkjer, D. & Matchkov, V. Vasomotion – what is currently thought? *Acta*
Physiol. **202**, 253–269 (2011).

53.   Li, J. *et al.* Ca2+ oscillation in vascular smooth muscle cells control myogenic spontaneous
vasomotion and counteract post-ischemic no-reflow. *Commun. Biol.* **7**, 332 (2024).

54.   Koenigsberger, M., Sauser, R., Bény, J.-L. & Meister, J.-J. Effects of Arterial Wall Stress on
Vasomotion. *Biophys. J.* **91**, 1663–1674 (2006).

55.   Font-Nieves., M, Sans-Fons, M.G., Gorina, R., Bonfill-Teixidor, E., Salas-Pérdomo, A., Márquez-
Kisinousky, L., Santalucia, T., Planas, A.M. Induction of COX-2 enzyme and down-regulation of COX-1
expression by lipopolysaccharide (LPS) control prostaglandin E2 production in astrocytes. *J Biol Chem.*
9:6454-68 (2012).

56.   Hasel, P., Aisenberg, W. H., Bennett, F. C. & Liddelow, S. A. Molecular and metabolic
heterogeneity of astrocytes and microglia. *Cell Metab.* **35**, 555–570 (2023).

57.   Sun, J. *et al.* Heterogeneity and Molecular Markers for CNS Glial Cells Revealed by Single-Cell
Transcriptomics. *Cell. Mol. Neurobiol.* **42**, 2629–2642 (2022).

58.   Escartin, C. *et al.* Reactive astrocyte nomenclature, definitions, and future directions. *Nat.*
Neurosci. **24**, 312–325 (2021).

59.   Zamanian, J.L., Xu, L., Foo, L.C., Nouri, N., Zhou, L., Giffard, R.G., Barres, B.A. Genomic analysis
of reactive astrogliosis. J Neurosci. 18,6391-410 (2012).

60.   Rueda-Carrasco, J. *et al.* SFRP1 modulates astrocyte-to-microglia crosstalk in acute and
chronic neuroinflammation. *EMBO Rep.* **22**, e51696 (2021).

61.   Pascual, O., Ben Achour, S., Rostaing, P., Triller, A. & Bessis, A. Microglia activation triggers
astrocyte-mediated modulation of excitatory neurotransmission. *Proc. Natl. Acad. Sci.* **109**, (2012).

62.   Zhao, X., Liu, M. & Cang, J. Visual Cortex Modulates the Magnitude but Not the Selectivity of
Looming-Evoked Responses in the Superior Colliculus of Awake Mice. *Neuron* **84**, 202–213 (2014).

63.   Institoris, A. *et al.* Astrocytes amplify neurovascular coupling to sustained activation of
neocortex in awake mice. *Nat. Commun.* **13**, 7872 (2022).

64.   Padmashri, R., Suresh, A., Boska, M. D. & Dunaevsky, A. Motor-Skill Learning Is Dependent on
Astrocytic Activity. *Neural Plast.* **2015**, 1–11 (2015).

65.   Davey, C.E, Grayden, D.B., Egan, G.F., Johnston, L.A. Filtering induces correlation in fMRI
resting state data. *Neuroimage.* **64**:728-40 (2013).

66.   Bolt, T., Nomi, J.S., Bzdok, D., Salas, J.A., Chang, C., Thomas Yeo, B.T., Uddin, L.Q., Keilholz, S.D.
A parsimonious description of global functional brain organization in three spatiotemporal patterns.
Nat Neurosci. **8**:1093-1103 (2022).

67.   Kölliker-Frers, R. *et al.* Neuroinflammation: An Integrating Overview of Reactive-
Neuroimmune Cell Interactions in Health and Disease. *Mediators Inflamm.* **2021**, 1–20 (2021).

68.   Frede, S., Stockmann, C., Freitag, P., Fandrey, J. Bacterial lipopolysaccharide induces HIF-1
activation in human monocytes via p44/42 MAPK and NF-kappaB. *Biochem J.* **3**,517-27 (2006).

69.   Van Veluw, S. J. *et al.* Vasomotion as a Driving Force for Paravascular Clearance in the Awake
Mouse Brain. *Neuron* **105**, 549-561.e5 (2020).

70.   Hablitz, L. M. & Nedergaard, M. The Glymphatic System: A Novel Component of Fundamental
Neurobiology. *J. Neurosci.* **41**, 7698–7711 (2021).

71.   Diem, A. K., Carare, R. O., Weller, R. O. & Bressloff, N. W. A control mechanism for intra-mural
peri-arterial drainage via astrocytes: How neuronal activity could improve waste clearance from the
brain. *PLOS ONE* **13**, e0205276 (2018).

72.   Aldea, R., Weller, R. O., Wilcock, D. M., Carare, R. O. & Richardson, G. Cerebrovascular
Smooth Muscle Cells as the Drivers of Intramural Periarterial Drainage of the Brain. *Front. Aging
Neurosci.* **11**, 1 (2019).

73.   Da Mesquita, S., Fu, Z. & Kipnis, J. The Meningeal Lymphatic System: A New Player in
Neurophysiology. *Neuron* **100**, 375–388 (2018).

74.   DiCarlo, G. Intrahippocampal LPS injections reduce Aβ load in APP+PS1 transgenic mice.
Neurobiol. Aging **22**, 1007–1012 (2001).

1321

1322

1323

1324

A

Frequency amplitude (0.08-0.15Hz)

Legend:
- wl
- 1 h
- 24 h
- 48 h
- 7 d

X-axis: PrimVisCtx, SecVisCtx, RetroDysCtx, RetroGrCtx$_A$, RetroGrCtx$_B$, SuperiorColl

PrimVisCtx: ***, ****
SecVisCtx: *, **
RetroDysCtx: ****

B

Reactive astrocytes (LPS 48h)

Transient reactivity

Control homeostatic astrocytes (VEH 48h)

Sub-reactive astrocytes (LPS 7d)

C

GFAP   NES

D

VEH, LPS-7d, LPS-48h

Pearson coef.
-1 -0.5 0 0.5 1

A

B   C   D

Pial vessel   Arteriole

Veinule

E

Pial vessels

Perfusion (mm.s⁻¹)

VEH   LPS

Penetrating vessels

Perfusion (mm.s⁻¹)

VEH   LPS

Penetrating arterioles

Perfusion (mm.s⁻¹)

VEH   LPS

Penetrating veins

Perfusion (mm.s⁻¹)

VEH   LPS

F

Pial vessels

Bubble density

VEH   LPS

Penetrating vessels

Bubble density

VEH   LPS

Penetrating arterioles

Bubble density

VEH   LPS

Penetrating veins

Bubble density

VEH   LPS