# Single neuron diversity supports area functional specialization along the visual cortical pathways

## Authors

Feyerabend M[1-3].*, Pommer S[4-6].*, Jimenez-Sosa MS[1-3]., Rachel J[4], Sunstrum J[1], Preuss F[4], Mestern S[1], Hinkel R[7-9], Mietsch M[7], Viyajraghavan S[1-3], Everling S[1], Treue S[10], Arnsten AFT[11], Lewis DA[12], Wolf F[6,13-16], Murray J[17-20], Steve McCarroll[21-22], Lyle Muller[23-26], Krienen F[27], Datta D[11,28], Wang XJ[29], Tripathy SJ[30-33], Gonzalez-Burgos G[12], Inoue W[1]**, Neef A[5,6,13,14,34,35]**, Staiger JF[4]**, Martinez-Trujillo J[1-3]**

*First authors contributed equally to the study

**Senior Authors contributed equally to the study

1. Department of Physiology and Pharmacology, Schulich School of Medicine and Dentistry, Western University, London, Ontario, Canada
2. Department of Psychiatry, Schulich School of Medicine and Dentistry, Western University, London, Ontario, Canada
3. Department of Clinical Neurological Sciences, Schulich School of Medicine and Dentistry, Western University, London, Ontario, Canada
4. Institute for Neuroanatomy, University Medical Center Göttingen, Georg-August University, Göttingen, Germany
5. Max Planck Institute for Multidisciplinary Sciences, Göttingen, Germany
6. Göttingen Campus Institute for Dynamics of Biological Networks, Göttingen, Germany
7. Laboratory Animal Science Unit, German Primate Center, Leibniz Institute for Primate Research, Göttingen, Germany.
8. German Center for Cardiovascular Research (DZHK), Partner Site Lower Saxony, Göttingen, Germany
9. Institute of Molecular and Translational Therapeutic Strategies, Hannover Medical School, Hannover, Germany
10. Cognitive Neuroscience Laboratory, German Primate Center, Leibniz Institute for Primate Research, Göttingen, Germany.
11. Departments of Neuroscience, Yale Medical School, New Haven, CT USA
12. Department of Psychiatry, University of Pittsburgh, Pittsburgh, PA, USA
13. Max Planck Institute for Dynamics and Self-Organization, Göttingen, Germany
14. Bernstein Center for Computational Neuroscience, Göttingen, Germany
15. Faculty of Physics, Georg-August-Universität Göttingen, Göttingen, Germany
16. Cluster of Excellence "Multiscale Bioimaging: from Molecular Machines to Networks of Excitable Cells", Georg-August University, Göttingen, Germany
17. Department of Psychiatry, Yale School of Medicine, New Haven, CT, USA
18. Manifest Technologies, New Haven, CT, USA
19. Department of Physics, Yale University, New Haven, CT, USA
20. Department of Psychological and Brain Sciences, Dartmouth College, Hanover, NH, USA
21. Stanley Center for Psychiatric Research, Broad Institute of MIT and Harvard, Cambridge, MA, USA
22. Department of Genetics, Harvard Medical School, Boston, MA, USA
23. Department of Mathematics, Western University, London, Ontario, Canada
24. Western Institute for Neuroscience, Western University, London, Ontario, Canada
25. Western Academy for Advanced Research, Western University, London, Ontario, Canada
26. Fields Lab for Network Science, Fields Institute, Toronto, Ontario, Canada.
27. Princeton Neuroscience Institute, Princeton University, Princeton, NJ, USA.
28. Department of Psychiatry, Yale University School of Medicine, New Haven, Connecticut.
29. Center for Neural Science, New York University, New York, USA.
30. Institute of Medical Science, University of Toronto, Toronto, Ontario, Canada
31. Krembil Centre for Neuroinformatics, Centre for Addiction and Mental Health, Toronto, Ontario, Canada
32. Department of Psychiatry, University of Toronto, Toronto, Ontario, Canada
33. Department of Physiology, University of Toronto, Toronto, Ontario, Canada
34. Institute for Nonlinear Dynamics, Georg-August-University Göttingen, Göttingen, Germany
35. Center for Biostructural Imaging of Neurodegeneration, Göttingen, Germany

# Abstract

Humans and other primates have specialized visual pathways composed of interconnected cortical areas. The input area V1 contains neurons that encode basic visual features, whereas downstream in the lateral prefrontal cortex (LPFC) neurons acquire tuning for novel complex feature associations. It has been assumed that each cortical area is composed of repeatable neuronal subtypes, and variations in synaptic strength and connectivity patterns underlie functional specialization. Here we test the hypothesis that diversity in the intrinsic make-up of single neurons contributes to area specialization along the visual pathways. We measured morphological and electrophysiological properties of single neurons in areas V1 and LPFC of marmosets. Excitatory neurons in LPFC were larger, less excitable, and fired broader spikes than V1 neurons. Some inhibitory fast spiking interneurons in the LPFC had longer axons and fired spikes with longer latencies and a more depolarized action potential trough than in V1. Intrinsic bursting was found in subpopulations of both excitatory and inhibitory LPFC but not V1 neurons. The latter may favour temporal summation of spikes and therefore enhanced synaptic plasticity in LPFC relative to V1. Our results show that specialization within the primate visual system permeates the most basic processing level, the single neuron.

# Introduction

Human and other primates have hierarchically organized cortical visual processing streams composed of serially connected areas, starting in the primary visual area V1 and extending downstream to the lateral prefrontal cortex (LPFC, Felleman and Essen, 1991). The diversity of neuronal response patterns, selectivity to different visual stimuli, and involvement in diverse perceptual and cognitive functions across areas has been extensively documented in the macaque monkey (Merigan and Maunsell, 1993; Nassi and Callaway, 2009; Martinez-Trujillo, 2022). This inter-area functional diversity has been implicitly assumed to be due to variations in expression levels of receptors subtypes, synaptic strength, inputs, and connectivity within an area's microcircuit (Carandini and Heeger, 2011; Godlove et al., 2014; Ardid et al., 2015; Jiang et al., 2015). It has been further proposed that neurons across cortical areas of mammals follow the principle of serial homology, i.e., single neuron classes are small quantitative variations of a common theme (Harris and Shepherd, 2015). Here we test the alternative hypothesis that single neurons in different visual areas of primates show qualitative and quantitative variations in their intrinsic morphological and functional properties that contribute to inter-area specialization.

Previous studies have shown that neurons in macaque area V1 show little changes in their tuning for basic visual features during training, even when subjects' show a strong improvement of behavioral discrimination thresholds for the same features (Ghose et al., 2002). On the other hand, neurons in the lateral intraparietal area (LIP) downstream from V1 acquire selectivity for categories during training (Freedman and Assad, 2006). In the LPFC, downstream from LIP, neurons flexibly become selective for new associations of visual features during intervals of minutes or hours (Rouzitalab et al., 2023; Abbass et al., 2024). At the molecular level, a recent study in macaques reported that the ratio of NMDA to AMPA receptors in synapses is larger in the LPFC than in area V1 suggesting that LPFC circuits support learning via synaptic plasticity to a larger degree than V1 circuits (Yang et al., 2018). Thus, the primate visual pathways extend along a cortical *stability – plasticity* axis: Neurons in early areas are *stably* tuned to basic visual features and neurons in far downstream areas *flexibly* 'learn' novel feature combinations encountered in new situations or environments. One issue that remains poorly investigated is whether variations in cells' intrinsic properties and morphology could shape single neuron computations and synaptic plasticity, ultimately contributing to inter-area specialization along this axis.

Reports of intrinsic electrophysiological properties and morphology of neurons in primates are scarce relative to other species such as the mouse. Previous studies showed that excitatory pyramidal neurons in area V1 and the LPFC of macaques diverge in their morphology (Amatrudo et al., 2012; Medalla and Luebke, 2015; Gilman et al., 2017), whereas pyramidal cells in mouse areas V1 and dorsal PFC do not have such prominent differences (Gilman et al., 2017). Moreover, a study in macaques has revealed differences in the response patterns of pyramidal cells between the lateral intraparietal (LIP) area and the LPFC (González-Burgos et al., 2019). It is possible that not only excitatory but also inhibitory neurons have diversified their intrinsic properties to contribute to the functional specialization of cortical areas in the primate visual pathways. However, we have not found any study systematically comparing intrinsic properties of inhibitory interneurons between different areas of the primate visual processing hierarchy.

We investigate this issue in the common marmoset (*Callithrix jacchus*), a non-human primate
(NHP) model that has become popular amongst neuroscientists due to their faster reproductive
cycle, amenability to transgenic manipulations and similarity of visual processing with humans
(Mitchell and Leopold, 2015; Okano et al., 2016). We compare intrinsic electrophysiological
properties and morphology of single neurons between areas V1 and the LPFC of marmosets.
We use methods, protocols and tools similar to those the Allen Institute of Brain Science (AIBS)
used in the mouse and human cell type database (Gouwens et al., 2019),  i.e. whole-cell patch
clamp recordings in acute slices and morphological reconstructions of single neurons. We built
a "Primate Cell Type Database" resource (https://primatedatabase.com). It contains 374
biophysical characterizations of intrinsic membrane properties as well as morphological
reconstructions of neurons from several areas (predominantly V1 and LPFC). We found that
intrinsic properties and morphology of excitatory and inhibitory neurons quantitatively and
qualitatively varied between V1 and the LPFC. Thus, area functional specialization along the
primate visual cortical pathways occurs at the most basic level of processing, the single neuron.

# Materials & Methods

## Statement on animal research

Research on NHPs is an ethically sensitive but irreplaceable part of neuroscience research. The
authors of this study are committed to pursue the best scientific result with the least possible
harm to the animals. Data was collected from 42 (male: 31, female: 11) common marmosets
(*Callithrix jacchus*, 26 at Western University + 16 at University of Göttingen) with a median age
of 4.43 years (IQR: 4.81, min: 1.52, max 18.06). Animal research from the Canadian group was
conducted in accordance with the Canadian Council of Animal Care policy on the care and use
of laboratory animals. The experiments were approved by the Animal Care Committee of the
University of Western Ontario. Animals from the German research group were fostered and kept
at the German Primate Center, Göttingen, Germany. Husbandry and experiments were
conducted in compliance with the Directive 2021/63/EU of the European Union and the German
Animal Welfare Act and therefore meet the regulations of the European Animal Research
Association. The animals were sacrificed as part of a broad study at the German Primate Center
approved by the Lower Saxony State Office for Consumer Protection and Food Safety (LAVES;
reference number 33.19-42502-04-20/3458). Experimental procedures were in accordance with
regulations of the German Animal Welfare Act. All animals were under constant veterinary
supervision.

## Acute marmoset ex vivo brain slice preparation

Marmosets were anaesthetized with Ketamine (20 mg/kg, intramuscular) and isoflurane (2–5
%) and then euthanized by trans-cardial perfusion (Canada) or pentobarbital (Germany). In the
latter case, ice-cold solution was poured over the head before opening the skull to slow down
cellular processes and decrease brain deterioration during organ removal. Afterwards, the brain
was immediately rinsed with pre-chilled (2–4 °C) N-methyl-D-glucamine (NMDG) substituted
artificial cerebrospinal fluid (aCSF) containing 92 mM NMDG, 2.5 mM KCl, 1.2 mM
$NaH_2PO_4·H_2O$, 30 mM $NaHCO_3$, 20 mM 4-(2-hydroxyetyl)-1-piperazineethansulfonic acid
(HEPES), 25 mM glucose, 5 mM Na-ascorbate, 2 mM thiourea, 3 mM Na-pyruvate, 10 mM
$MgSO_4·7H_2O$ and 0.5 mM $CaCl_2·2H_2O$. The pH was adjusted to 7.3–7.4 with concentrated
hydrochloric acid (37 %) and the osmolality was set to 300–305 mOsm/kg. The solution was

oxygenized with 95 % $O_2$ and 5 % $CO_2$ (carbogen) for 15 minutes prior to animal surgery. The brain was transferred into a container containing the same NMDG aCSF and quickly transported from the animal facility to the laboratory room (10–20 min) (Ting et al., 2018; Gouwens et al., 2019).

The hemispheres were separated and trimmed to blocks of prefrontal and visual cortex. Slices of 300 μm were cut with a vibratome VT1200 S (Leica) using the same ice-cold NMDG aCSF as above. Subsequently, the slices were transferred to a recovery chamber filled with NMDG aCSF at 32 °C and incubated for 12 minutes before stored in a holding chamber at room temperature. This chamber was filled with HEPES aCSF containing 92 mM NaCl, 2.5 mM KCl, 1.2 mM $NaH_2PO_4 \cdot H_2O$, 30 mM $NaHCO_3$, 20 mM HEPES, 25 mM glucose, 5 mM Na-ascorbate, 2 mM thiourea, 3 mM Na-pyruvate, 2 mM $MgSO_4 \cdot 7H_2O$ and 2 mM $CaCl_2 \cdot 2H_2O$. The slices were stored for up to 10 hours with minimal submersion and transferred to fresh HEPES aCSF after 6 hours. HEPES provided additional support to pH buffering and reduced slice deterioration over extended time periods. All solutions were continuous oxygenized with carbogen gas and set to 300–310 mOsm/kg.

All solutions for ex vivo brain slice preparation and electrophysiological recordings were adopted from the technical white paper from the AIBS Cell Type Data base. The latter facilitates the comparison between our marmoset dataset with mouse and human datasets from AIBS Cell Type Data base (Gouwens et al., 2019; Berg et al., 2021).

## Patch clamp electrophysiology

Individual brain slices were transferred into the recording chamber and continuously perfused with ~32 °C carbogen-saturated aCSF containing 126 mM NaCl, 2.5 mM KCl, 1.25 mM $NaH_2PO_4 \cdot H_2O$, 26 mM $NaHCO_3$, 20 mM HEPES, 12.5 mM glucose, 1 mM $MgSO_4 \cdot 7H_2O$, 2 mM $CaCl_2 \cdot 2H_2O$ and a mix of synaptic blockers (2 mM kynurenic acid and picrotoxin 0.1 mM). Thick-walled borosilicate glass pipettes were manufactured on the day of recording and filled with intracellular solution containing 126 mM K-gluconate, 4 mM KCl, 0.3 mM EGTA, 10 mM HEPES, 4 mM $K_2$-ATP, 0.3 mM Na-GTP, 10 mM $Na_2$-Phosphocreatinine. The pH was adjusted to 7.3–7.4 with 1 M KOH and the osmolality was set to 290 mOsm/kg with sucrose when necessary. The internal solution was supplemented with 0.5% biocytin to fill the recorded neurons for histology. Pipette resistance in the bath was 3-7 MΩ. Recording equipment varied across experimenters: data were acquired with either a Multiclamp 700B and a Digidata (Axon instruments/ molecular devices), a SEC-10LX or SEC-05X (npi electronics) digitized with a 1401-3A (CED) or an EPC 10 USB (HEKA). After formation of a stable seal above 1 GΩ and break-in, access resistance was determined in voltage clamp (VC) mode. Then, the recording mode was switched to current clamp (CC) and bridge balance and pipette neutralization were applied. When necessary, a holding current was applied and continuously readjusted throughout to maintain a membrane potential close to the initial value after break-in. Cells that required more than 20 pA holding current to keep the membrane potential at -60 mV in the initial whole-cell configuration were generally not recorded from. Recording protocols and procedures were standardized across laboratories and acquisition equipment. Neurons were subjected to hyper- and depolarizing 1 s square pulse current injections. Furthermore, 275 (73.5%) cells were recorded with an additional 3 ms short square pulse protocol.

## Histology

After recording, slices were fixed with 4% m/v PFA + 15% v/v picric acid (1.3% saturated in water) at 4 °C for a period of 24–36 hours and then washed in PBS. Then, slices were permeabilized by incubation in PBS + 2% Triton-X for 2 x 15 min. Biocytin filling of patched cells was made visible by a 4-hour incubation of streptavidin conjugated with Alexa Flour 647. After a subsequent washing step, slices were incubated in PBS with DAPI (1:4000), before mounted on specimen slides with aqua-polymount.

Imaging was done using a SP8 confocal microscope (Leica, Germany) or a LSM 880 with Airyscan (Zeiss, Germany) to assess the cell location, quality of the filling and dendritic type. Dendritic type was determined based on spininess and somatodendritic configuration of the cell similar to Gouwens et al. 2018. Cells with sufficiently retained dendritic and axonal arborization were used for another acquisition with 40x or 63x magnification (z-step size 0.5–0.8 µm). These image stacks were then used to reconstruct the neuronal morphology using Neurolucida (MBF Bioscience). Automated quantifications of neurites were obtained with Neurolucida explorer (MBF Bioscience, USA).

## Analysis of electrophysiological features

Raw recordings were converted into the neurodata without borders (nwb) format (version 2.4.0; (Rübel et al., 2022)) using the MATNWB repository (https://neurodatawithoutborders.github.io/matnwb/). Additional information such as subject data (age, sex, etc.), acquisition parameters (access resistance, temperature, etc.) or histological specifications (layer, putative superclass, etc.) were added if available. A custom MATLAB analysis pipeline was used for quality control and extraction of electrophysiological features. The code is publicly available on GitHub (GitHub - mfeyerab/MATFX). Recordings from cells were not considered for further analysis if one of the following criteria was met: (1) an initial membrane potential above -55 mV or (2) input resistance or action potential (AP) waveform features could not be determined. In addition, recording quality was assessed in a sweep-wise fashion. For this purpose, a time window of 250 ms for each pre and post stimulus period was selected. A sweep was discarded if: (1) the membrane potential in either segment exceeded a root mean square error of 0.65, (2) the membrane potential difference between pre and post stimulus exceeds 2.5 mV, (3) the membrane potential in the pre-stimulus segment is not more depolarized than -52.5 mV, (4) the membrane potential of the pre-stimulus deviates more than 6 mV from the median pre-stimulus membrane potential of all passing sweeps, (5) flagged by manual curation based on evaluation of the pre-stimulus test pulse (see **Table 1** and **STab 1 for a list of key features, and STab 2** for description of all electrophysiological features).

## UMAP and cross-species classification

Uniform Manifold Approximation Projection, UMAP (McInnes et al., 2018) space was created by combining three data sets each obtained from a different species. Mouse and human data were available at the AIBS Cell Type Database. For the classification analyses separate training and test data sets for each species were chosen. Cells with a missing value in more than 15% of features were excluded from the subsequent analysis yielding a total number of 2421 cells (1757 mouse, 356 human, and 308 marmoset). Initially, 29 electrophysiological features (see supplement for more detail) were reduced to 16 components by probabilistic PCA explaining more than 95% of the total variance. Then data was transformed into UMAP space (McInnes et al., 2018) using custom MATLAB scripts in two instances: (1) in a space with two components

for visualization of electrophysiological diversity in the different data sets (number of nearest
neighbours = 60, minimum distance = 0.2) and (2) in a space with three components for
supervised classification across species (number of nearest neighbours = 30, minimum distance
= 0.3).

For classification, three labels were chosen: Class 1 (C1), which contains fast-spiking
interneurons (FSI); Class 2 (C2), which contains the non-fast spiking population of interneurons
and Class 3 (C3), which contains excitatory cells (EXC). Ground truth data (**SFig. 1B**) was
obtained by Cre reporter lines and electrophysiological types (e-types, determined by
unsupervised clustering by Gouwens et al. 2019), to exclude contamination by off-target
expression (Hu et al., 2013). Cells which expressed Cre with the parvalbumin (PV) promoter
and an e-type ranging from Inh_8 to Inh_13 were considered fast spiking interneurons (FSI).
Cells which had a spiny dendritic type were considered EXC. Cells that expressed Cre with a
somatostatin (SST) promoter and an e-type ranging from Inh_4 to Inh_7 were considered SST
cells, whereas cells that expressed Cre with a serotonin receptor 3a (5-HT3A) or vasoactive
(poly)peptide (VIP) promoter and had either e-type 2 or 3 were considered VIP cells. Lastly,
cells that had the e-type "Inh_1", which was marked by low spike-frequency adaptation and
commonly high latency of the rheobase spikes and expressed Cre with either a 5-Htr3a or a
neuron-derived-neurotrophic-factor (Ndnf) promoter were considered neurogliaform (NGF) cells.
C2 consisted of the SST, VIP and NGF subtype, which do not show the distinctive fast spiking
phenotype of PV cells. Cell type composition of training data was informed by histological data
of the primate neocortex, consequently the ratio of SST, VIP and NGF cells within the training
data was fixed to 2:2:1. A random forest classifier for cross species cell labeling was trained
(Seiffert et al., 2010) with uniform priors with a subset of the 3 component UMAP data
consisting of 1174 mouse cells with marker labels. Cell composition of test data was matched
across species and it was biased towards excitatory cells (~65 %). Marmoset and human test
data set for cross-species validation was determined by unambiguous morpho-
electrophysiological identity (see examples in **Fig 2** and **SFig 1**).

## Statistical analysis

All hypothesis testing regarding differences between areas was done with 2-sided rank sum
tests in MATLAB and effect sizes were reported by rank bi-serial correlation ($r_{rb}$). All $r_{rb}$ values
and their 95% confidence intervals were calculated with the measures-of-effect-size-toolbox
(Hentschke and Stüttgen, 2011). Reported p-values were corrected for false discoveries rates
via the Benjamini-Hochberg procedure. Correlations reported were either tested by Spearman
(correlation coefficient: $\rho$) or Pearson (correlation coefficient: r).

The residual values of rheobase were obtained from a linear regression model with rheobase as
dependent variable and the ratio between the difference in the action potential threshold and the
resting membrane potential (Dvoltage), and the input resistance (R) as predictor variable. This
model (rheobase = B0 + B1*(Dvoltage/R) + residuals) captures the passive/ohmic aspect of the
rheobase.

## Effect of different acquisition systems on AP waveform

Our data were recorded at two different sites with different acquisition systems. Therefore, we
checked whether this had an effect on electrophysiological measurements. Data recorded with
the HEKA amplifier show significantly higher AP amplitudes than those recorded with npi (**SFig
2A**), likely because differences in pipette capacitance compensation between amplifiers.

Uncompensated pipette capacitance, together with the access resistance, create a passive low-
pass filter which artificially shorten and widen the AP. This effect was stronger in neurons with
narrow spike waveforms. Selection of AP waveform features for the UMAP procedure was
informed by these observations. AP width was the only wave form parameter chosen as input,
since it showed no significant effect between the amplifier systems (**SFig 2B**). Ultimately,
neurons in the UMAP projection did not suggest clustering by acquisition equipment (**SFig 2C**)
and we conclude that our intrinsic property profiles are mainly shaped by biological factors such
as cell identity.

# Results

We recorded from a total of 463 neurons in slices obtained from 42 marmosets. 374 neurons
(80.8%) passed our quality control (QC pipeline) to enter the primate cell type database (see
**Fig 1**). This database contains 107 V1 cells, 256 LPFC cells (focused on Brodmann area (BA) 8
and medial part of BA 46) and 11 cells from cingulate cortex (not included in this study). We
targeted both pyramidal and non-pyramidal cells focusing on supragranular layers. A total of
144 neurons were examined histologically, and spine density was evaluated to classify the cells
into aspiny, sparsely spiny or spiny. The latter could further inform whether a neuron was
inhibitory or excitatory (Gouwens et al., 2019). 29% of the cells evaluated were either aspiny or
sparsely spiny (for clarity purposes we will refer to the latter two categories as aspiny), and 71%
were considered spiny. The marmoset dataset contains a wide spectrum of morphologically
different cell types (**Fig. 2A, B**). Of the 144 neurons evaluated morphologically, a total of 32
cells were reconstructed. We also used two other data sets from mouse and human,
respectively, which are publicly available from the Allen Institute for Brain Science's (AIBS) Cell
Type Database (Gouwens et al., 2019). After applying the same QC pipeline as in the marmoset
data, we included 1907 of 1920 AIBS mouse cells (99.32%) and 403 of 413 human cells
(97.58%) in our analyses **(SFig 1A)**.

### Comparing profiles of intrinsic properties across species

We combined our marmoset data with the two AIBS datasets in a multivariate map of
electrophysiological features to determine whether the cell type profiles of intrinsic membrane
properties of marmoset, mouse and human cortex neurons overlap under varying factors such
as species or cortical area (**Fig 2C**). Importantly, our data were acquired with protocols and
solutions identical to those used in the AIBS studies, thus minimizing the influence of
methodological factors on potential inter-species differences (see Methods). Intrinsic properties
of all cells were quantified in the same analysis pipeline by 29 electrophysiological features
covering the subthreshold domain, action potential (AP) shape, and AP (spike) train pattern
diversity (see **STab 1** for a list and respective quantification of features). Subsequently, we used
the dimensionality reduction procedure "Uniform Manifold Approximation and Projection"
(UMAP) for visualizing intrinsic properties profiles (**Fig 2C**). While the data showed a general
overlap across species, marmoset and human cells were concentrated in hot zones marked by
large sag: a delayed deflection in the membrane potential response to hyperpolarizing current
injections (**Fig. 2C** upper right panel, mouse data indicated by gray filled circles in **Fig. 2C** upper
left panel).
327
328  To better understand the distribution of different cell types on the UMAP projection, we utilized
329  available cell type labels from 1460 mouse cells (76.6 % of analyzed mouse cells). Neurons
330  showing the "spiny" dendritic type were considered to be excitatory and 4 major interneuron

subtypes (PV/SST/VIP/NGF) were identified by a combination of e-type (Gouwens et al., 2019) and labeling by one of five Cre-lines (PV-Cre/SST-Cre/VIP-Cre/5HT3A-Cre/NDNF-Cre; for more detail see **SFig 1**). High density regions of the corresponding cell identity are indicated by color shading in **Fig 2D** (see also **SFig 1**). One of the most prominent differences between GABAergic and EXC cells is the speed of repolarization after the peak of an AP, with inhibitory cells having a narrower AP. Indeed, the first UMAP component is strongly influenced by AP width (r = 0.745, p < 0.001, see gradient in **Fig. 2C** lower panel) with FSI cells clustering to the left of the map. This electrophysiological profile is strongly associated with PV fast spiking interneurons, which was in line with the corresponding cell type cluster (see orange background colors of **Fig. 2D**). Although different cell types tend to localize to a certain region in the UMAP projection, they rarely separated into well-defined clusters but rather occupied a certain segment on a continuous spectrum of intrinsic properties. VIP and SST cells, indicated by purple and gold, showed a high degree of overlap.

The dendritic type of marmoset neurons (spiny vs. aspiny) showed a good correspondence with mouse cell types, with occasional intermixing between aspiny and spiny cells (see **Fig 2D** and **SFig 3B** for dendritic type of marmoset cells). The only notable shift of intrinsic property profiles could be observed for neurogliaform (NGF) cells, which in mouse were associated with low spike frequency adaptation and commonly high spike latencies (**Fig. 2D** cyan marker and background). Interestingly, all 6 morphologically identified NGF cells in marmoset showed strong spike frequency adaptation and short spike latencies (see **SFig 4**). Except for NGF cells, these results suggest that overall feature profiles of distinct cell types overlap across the different species.

## Classification of marmoset cell types based on intrinsic properties

We merged the four interneuron subpopulations of AIBS mouse data (i.e., PV, SST, VIP, NGF) into two group labels, labelling the PV cells were labeled as class 1 (C1), while the SST, VIP, and NGF were grouped as class 2 (C2). The label EXC was preserved, making class 3 (C3). We projected the intrinsic electrophysiological properties onto a 3D UMAP (**Fig 3**). We trained a random forest classifier (Seiffert et al., 2010) to predict the class label (C1, C2, C3) from the UMAP components using a sample of 1174 mouse neurons. The procedure was repeated 500 times with random training sets of fixed numbers of cells (**Fig. 3A**). We finally used a sample of 150 cells (not included in the training set) to assess the classifier performance in the mouse data. Median test performance in the mouse across the different classes was 91.3% (**Fig. 3B**).

We used the mouse classifier to predict the different cell classes (C1, C2, C3) in the marmoset. We assessed the classifiers' performance using a test set of 48 marmoset cells not included in any of the previous analyses but with available morphological information which was used to provide "ground truth" labels, as follows: fast spiking neurons (FSI) with congruent (i.e., basket cell) morphology were determined to be C1 cells, whereas other aspiny neurons that could be identified as different morphological types (such as bi-tufted cells, NGFs, etc.) were determined to be C2 cells (see **Fig 2A,B**). C3 cells were drawn from the pool of all spiny cells. Marmoset cells were classified with a median accuracy of 83.3% (**Fig 3B**). Confusion matrices provided insight into classification accuracy by class (**Fig 3C**). Compared to mouse, erroneous classification of C1 and C3 as C2 cells was more common (positive predictive values: mouse = 78.7% vs marmoset = 55.7%). Marmoset C1 and C3 cells were identified with high specificity (positive predictive value: C1 = 94.3% and C3 = 95.8%).

We next examined the correspondence between the categories identified by the classifier and dendritic type, as an additional evaluation independent of intrinsic membrane properties (**Fig

**3E-F**). Final class labels were determined by the most frequent prediction throughout different
rounds of classification (repetitions) with different training subsamples (**Fig 3E**). Composition of
dendritic type by putative class (**Fig 3F**) is in line with positive predictive values of the test data
sets. In marmoset, 99% of putative C3 cells were spiny. The fraction of aspiny cells in the
putative C2 cells (68.1%) matches very well with the share of ground truth interneurons (i.e., C1
and C2) in the totals of the test data (68.8%). In summary, these findings indicate that the cross-
species classification leads to reliable labels for both C1 and C3, but less reliable labels for C2.

We repeated the same procedure in a sample of human neurons obtained from the AIBS
database and the results were qualitatively similar for the different classes. However, there was
a slight decrease in performance relative to the marmoset (**SFig 3D**).

## Comparison of intrinsic properties between V1 and LPFC neurons

We used the classifier to identify C1 and C3 cells and then compared the intrinsic properties of
these classes between V1 and LPFC. First, we focused on C3, which includes EXC cells. A
summary of comparisons in key features is depicted in **Table 1**. We focus on supragranular
(layers II-III) neurons because these neurons process visual information from upstream areas
(Roussy et al., 2022a, 2022b), and encode visual long term memories (Corrigan et al., 2022;
Rouzitalab et al., 2023; Abbass et al., 2024), so they were intentionally targeted in our
recordings (**Fig. 1D**).

We found that C3 cells from LPFC had a lower input resistance ($p < 1e-4$, effect size $r_{rb} = 0.62$),
higher rheobase ($p < 1e-4$, $r_{rb} = -0.63$), and less rectification ($p < 1e-4$, $r_{rb} = -0.64$) than V1 cells
(**Fig 4A, B, Tab1**). These results suggest that LPFC EXC cells are less excitable than V1 cells.
To investigate the factors that may underlie the differences between areas, we examine how the
size of C3 dendritic trees (quantified by total length) relates to the cell's input resistance (**Fig
4C**). Indeed, both parameters showed a correlation of rho = -0.929 (p-value < 0.001), meaning
that cells with shorter dendritic trees have a higher input resistance. V1 C3 cells had
systematically smaller dendritic trees than LPFC C3 cells. V1 cells also fired significantly
narrower action potentials with a robust effect size ($p < 1e-4$, $r_{rb} = -0.42$; **SFig 5**). Interestingly,
firing rate adaptation, the increase in inter-spike interval duration after stimulation onset, was
larger in LPFC than in V1 ($p = 0.0032$, $r_{rb} = -3.14$). A list of comparisons for several intrinsic
features appears in **Tab1**. A total of 14 out of 19 features showed significant differences
between areas.

We further compared intrinsic properties of C1 fast spiking inhibitory interneurons between the
two areas (see **STab 1**). Out of 19 analyzed features, 4 showed significant differences indicating
that intrinsic features in interneurons were more similar between areas. We found that AP width
was smaller in V1 than LPFC ($p = 0.0369$; $r_{rb} = -0.51$). The latency at the herosweep (see
**methods** and **STab 2**) was longer in LPFC than in V1 ($p = 0.0369$, $r_{rb} = -0.58$). The difference in
trough was also significant with a more negative value in V1 than in LPFC ($p = 0.0369$, $r_{rb} = -
0.54$). The rheobase showed a large difference and effect size between areas (see **STab1**), but
it did not reach statistical significance.

## Burst firing in C1 and C3 cells

Bursts are short, high frequency trains of action potentials or spikes that could elicit a response
in the postsynaptic cell with higher probability than isolated spikes (Zeldenrust et al. 2018). We
computed burst indices in all the recorded neurons (see methods). Both C1 and C3 neurons

showed an increased burst index in LPFC relative to V1 (**Fig 4D;** C1: p = 0.0369, $r_{rb}$ = -0.51; C3:
p < 1e-04, $r_{rb}$ = -0.39). Some C3 neurons in both areas fired duplets of action potentials at the
beginning of stimulation at medium to high intensities (**Fig 4E**). Violin plots show a distribution
that was best fitted by the sum of 2 Gaussians rather than by a single Gaussian function (LPFC
data, Akaike Information Criteria, AIC 1 Gaussian = -140; AIC 2 Gaussians = -181), indicating
bimodality. Consequently, we divided all C3 cells into two subpopulations according to the
presence of duplets, or two-spikes bursts (see example traces in **Fig. 4E**). After isolating C3
bursting cells we found that: 1) bursting cells were more frequent in LPFC than in V1 (54% vs
34%, **Fig 4E**), and 2) bursting cells in LPFC have a higher burst index than bursting cells in V1
(p < 1e-4, $r_{rb}$ = -0.50, **Fig 4F**). In summary, C3 LPFC neurons show an increase in bursting
relative to V1.

Interestingly, some of the C1 neurons in LPFC also showed a significant increase in burst index
(**Fig 4D**), which was associated with a bursting fast-spiking phenotype (purple traces in **Fig 5A).**
Unlike C3 neurons, C1 cells responded with bursts of multiple action potentials at rheobase
stimulation, which was often accompanied by other features of intrinsic bursting at higher
stimulation intensities such as sustained depolarization and varying AP wave forms. These
bursting C1 neurons were exclusive to LPFC (5 out of 19, i.e., 26.3 %, **Fig 5B**) and appeared to
be basket cells (BCs, see **Fig 5C**). With increasing stimulation, the gap in inter-spike intervals
between the initial burst and post-burst narrowed, making it harder to distinguish bursting vs
non-bursting fast spiking cells. At medium stimulus intensity, such as at the hero sweep,
bursting C1 cells (purple color) had a higher burst index relative to other C1 LPFC cells (pink
color, **Fig 5D**). LPFC cells we considered as not intrinsically bursting (pink color), show a higher
burst index than their V1 counterparts (blue color).

Intrinsic bursting cells typically have a prolonged active depolarization mechanism that
facilitates the generation of action potentials. To explore this issue, we next examine the
relationship between burst index and variables potentially linked to neuronal excitability. We first
developed a linear regression model with rheobase as dependent variable and the ratio
between the difference in the action potential threshold and the resting membrane potential
(Dvoltage), and the input resistance (R) as predictor variable. This model (rheobase = B0 +
B1*(Dvoltage/R) + residuals) captures the passive/ohmic aspect of the rheobase and provides a
reasonable goodness of fit (adj. R2 = 0.73, p-value = 2.75e-09; model parameters [B0 = -3.654,
p = 0.82597], [B1 = -826.86, p = 3.0106e-9*]). The model equation was used to obtain predicted
values and residuals.

Negative residuals of this model reflect active depolarization mechanisms. We found a
significant negative correlation between these residuals and burst index ( r = -0.642, p < 0.0001,
see **Fig 5E**). These results suggest that an active depolarization mechanism underlies bursting
in C1 cells. Burst index also correlated significantly with other variables, i.e., latency of the first
action potential and amplitude of the afterhyperpolarization (latency p = 0.017, rho = 0.441; AHP
amp , p = 0.009, rho = 0.474; see **Fig 5F**). Cells with higher burst index showed longer latencies
to first spike at rheobase and weaker afterhyperpolarization, consistent with a depolarizing
mechanism that activates slowly and persist beyond the first action potential within a burst.

**Differences in morphology between V1 and LPFC neurons**
We investigated morphological features of C1 and C3 cells in LPFC and V1. We reconstructed 4
V1 and 11 LPFC pyramidal cells (PC). Two representative examples of PCs from V1 and LPFC
are shown in **Fig 6A**. We focused on dendrites from PCs because their long axons are more

likely to be truncated following slice preparation (Bruno et al., 2009; Oberlaender et al., 2011; Mohan et al., 2015; Goriounova et al., 2018). The total dendritic length of V1 PCs was half of their LPFC counterparts (median$_{LPFC}$ = 5760 μm, median$_{V1}$ = 3119 μm, p = 0.0352, r$_{rb}$ = -0.63). Quantitative analysis showed that both apical dendrite length and complexity was significantly higher for PCs in LPFC than in V1 (apical length: p = 0.0352, r$_{rb}$ = -0.63, apical complexity: p = 0.0352, r$_{rb}$ = -0.63, **Fig 6C**). The result followed the same trend for basal dendrites with effect sizes of 0.42 and 0.52, although it did not reach statistical significance at 0.05.

We also reconstructed 9 C1 (3 V1, 6 LPFC) fast-spiking basket cells (BC), all in layer III. Two example cells from V1 and LPFC are shown in **Fig. 6B**. In contrast to PCs, interneurons have a more localized axonal arborization, less likely to be affected by the thickness of the slice preparation. The axonal length of LPFC fast spiking BCs doubled the one of V1 neurons (median$_{LPFC}$ = 19.743 mm, median$_{V1}$ = 9.172 mm). There was over 5-fold increase in axon complexity in LPFC relative to V1 (median$_{LPFC}$ = 52,0e6, median$_{V1}$ = 9,2e6, **Fig 6D**). A larger sample of reconstructed cells would be needed to more systematically assess differences between areas.

In summary, we found that PCs in LPFC have larger dendritic complexity and length than in V1. On the other hand, a small sample of reconstructed fast-spiking BCs in LPFC show a more extensive and complex axon than in V1. These findings suggest that morphological changes have accompanied changes in intrinsic electrophysiological properties observed between V1 and LPFC neurons.

# Discussion

We compared electrophysiological and morphological features of single neurons in two different areas of the common marmoset visual cortical processing streams: V1, the input and earliest sensory area, and the LPFC, located far downstream at the end of the visual processing stream (Felleman and Essen, 1991). We conducted intracellular recordings using whole-cell patch-clamp in brain slices and anatomical reconstructions of neurons in both areas. To identify cell types, we built a random forest classifier using anatomical and transgenic labels, with electrophysiological features as predictors. We used the classifier to identify three classes of cells: C1 (aspiny fast-spiking interneurons), C2 (aspiny non-fast spiking interneurons), and C3 (spiny excitatory cells). We focused on C1 and C3 groups and compared neuronal intrinsic features between areas V1 and LPFC. C3 cells in V1 were more excitable, smaller, and fired narrower spikes than in LPFC. C1 neurons in the LPFC had longer latencies and a more depolarized action potential trough than V1 cells. In both C3 and C1 groups, intrinsic bursting was more commonly found in LPFC relative to V1 neurons.

**A tool for classifying neuron types using intrinsic properties and morphology**

Neuronal diversity has been a puzzle for neuroscientists since the times of Santiago Ramon y Cajal, who described different cell morphologies across brain regions and species and suggested a relationship between the structure/morphology and function of a neuron (Szentágothai, 1990). Indeed, morphology and intracellular electrophysiological tools have been used to classify neurons (Ascoli et al., 2008). Progress in transcriptomics has advanced this field considerably (Saunders et al., 2018; Krienen et al., 2020; Yuste et al., 2020). However, studies of cell types have been mainly conducted in mice, where transgenic tools for single neuron classification are available and one can use large number of animals. In primates such

as the macaque monkey some studies have examined electrophysiological and morphological
features of pyramidal and non-pyramidal neurons (Gonzalez-Burgos et al., 2000; Zaitsev et al.,
2005, 2012; Povysheva et al., 2007, 2008; Amatrudo et al., 2012; Luebke et al., 2015; Medalla
and Luebke, 2015; Gilman et al., 2017; Medalla et al., 2017). However, in general they had low
sample sizes and/or were limited to one cell type and/or brain area. More recent studies in
resected tissue from human patients with epilepsy or brain tumors have also documented
intrinsic features of neurons and their divergence from those of in the mouse brain (Kalmbach et
al., 2018; Berg et al., 2021). However, they have also been restricted to one brain region, the
temporal lobe, and changes due to the underlying pathology may have influenced their results.

Over the last decade, the common marmoset has emerged as a model in neuroscience
research (Cyranoski, 2014; Mitchell et al., 2014; Mitchell and Leopold, 2015; Miller et al., 2016;
Servick, 2018) opening a door to explore cell types in primate brains. In the current study we
took advantage of the marmoset model to explore cell type diversity in areas V1 and LPFC of
the primate cortical visual pathways. We used the same protocols as the AIBS to minimize
technical and methodological confounds that could hinder comparisons of neuronal intrinsic
properties and morphology between datasets (e.g., our marmoset data vs. AIBS mouse data).
Our classification using transgenic mouse labels, as well as anatomical features (e.g., spiny and
aspiny) resulted in three main classes of neurons (C1, C2, and C3) that include known cell
types (Group et al., 2008; Tremblay et al., 2016; Zhou et al., 2020). C3 were mainly excitatory
pyramidal cells, C1 fast spiking interneurons, and C2, more variable in their electrophysiological
features, was mainly comprised of VIP and SST positive neurons in the mouse (Group et al.,
2008; Defelipe et al., 2013; Tremblay et al., 2016; Merino et al., 2019; Zhou et al., 2020). This
latter group likely included CB and CR positive interneurons in primates, thought to be
homologues of SST and VIP positive cells in the mouse (Condé et al., 2004; Zaitsev et al.,
2009, 2012; Torres-Gomez et al., 2020).

Our classification tool could be used by studies in humans and non-human primates that
investigate the properties of different neocortical cell types when transgenic labels are not
available. One important contribution of our study is highlighting homologies and differences
between broad categories of cell types across species, which is critical to the use of animal
models in research (**Fig. 2C, D)**. The differences between areas observed here agrees with the
finding of transcriptomics studies in marmosets (Krienen et al., 2020). One general limitation of
the existing maps is that the samples are biased to few brain areas (e.g., V1 in mice and
Temporal cortex in humans (Gouwens et al., 2019), V1 and LPFC in marmosets (this study),
which makes it difficult to perform wide inter-area comparisons across species. Future work is
needed to fill this gap and provide a comprehensive map of functional and anatomical properties
of neurons across different cortical areas of primates.

**Differences in intrinsic properties and morphology between cell types in V1 and LPFC**

A main goal of this study was to identify distinctive features of cell types in areas V1 and the
LPFC of the common marmoset. V1 and LPFC are in 'opposite' extremes (beginning and end)
of the visual processing cortical hierarchy. In the initial stages of visual processing V1 neurons
must fast and reliably process incoming information from the thalamus. We found that C3
excitatory cells in V1 showed high excitability and narrow action potentials, which may
contribute to the elevated firing rates and short spike latencies relative to downstream areas
reported by extracellular studies in macaques (Schmolesky et al., 1998; Reynolds et al., 1999).
V1 also has the highest neuronal densities along the visual cortical pathways. The small C3

neuronal sizes may allow 'packing' more cells within the area. Indeed, excitatory V1 neurons
receive highly spatially-specific LGN inputs corresponding to a small region of the retina,
resulting in a high-resolution map of the visual field (Sincich and Horton, 2005). We also found
that fast spiking interneurons (C1) in V1 show sustained high firing rates with low adaption in
response to square current pulses. Additionally, V1 excitatory cells (C3) had high excitability,
small dendritic trees and less intrinsic bursting. These features make V1 neurons suitable for
linearly or quasi-linearly encoding important spatiotemporal details of visual inputs.

LPFC pyramidal neurons have large dendritic arborizations and many spines that can
integrate inputs from a variety of sources (Young et al., 2014; Preuss and Wise, 2022).
Pyramidal cells in the LPFC are known to fire with lower rates than in early sensory areas
(Lennert and Martinez-Trujillo, 2013). Moreover, neurons output (i.e., firing rates) in LPFC has a
higher correlation with performance than in upstream areas such as MT (Mendoza-Halliday et
al., 2014) and presumably V1. Thus, LPFC signals seem to be more directly linked to our
perceptions and actions than V1 signals. We found that excitatory cells (C3) in LPFC fired fewer
spikes with the same current stimulation levels as V1. The lower excitability of LPFC pyramidal
neurons is suited for selecting strong inputs corresponding to stimuli that are salient and/or
behaviorally relevant and for producing sparse codes that may underlie our perceptual
capabilities and sense of awareness (Panagiotaropoulos, 2024). However, the same structural
and functional complexity may make LPFC pyramidal cells vulnerable to new categories of
disease that would affect the highly developed cognitive skills of primates such as
Schizophrenia (Glantz and Lewis, 2000; Lewis and González-Burgos, 2008; Arnsten, 2011) and
Alzheimer disease (Arnsten et al., 2012).

Most interneurons receive their main inputs from local pyramidal cells. Pyramidal
neurons and fast spiking interneurons intrinsic properties in V1 and LPFC may have diversified
to perform computations according to the area's input-output landscape. Indeed, certain types of
interneurons are more abundant in LPFC than in early sensory areas (Condé et al., 2004;
Torres-Gomez et al., 2020). One issue that remains unclear is when and how such
diversification occurs. For the case of LPFC it is possible that the protracted development of
primate species provides a prolonged window to adjust mRNA and protein expression (Bakken
et al., 2016) based on local feedback (i.e., levels of firing rate) and therefore produce variations
of what has been considered canonical cell types of the neocortex. In species with shorter
protracted periods such as mice, neuronal types may appear more serially homologous (Harris
and Shepherd, 2015; Gilman et al., 2017) perhaps due to a mode limited time window for
epigenetics factors to transform the neuronal diversity landscape. This issue may be highly
relevant for the study of mental disease (Lewis and Sweet, 2009) and deserves further
investigation.

**Intrinsic bursting as a distinctive feature of LPFC neurons**

A burst is a short, high frequency trains of spikes with a higher probability than single spikes to
elicit a postsynaptic spike via temporal summation of excitatory postsynaptic potentials
(Zeldenrust et al., 2018). Bursts could arise by intrinsic mechanisms or by network dynamics
(Zeldenrust et al., 2018). Intrinsic bursters usually need a slow depolarization mechanism on top
of which action potentials are generated. Intrinsic currents such as T-type Calcium have been
isolated in bursters (Jahnsen and Llinás, 1984; Williams et al., 1997). The intrinsic bursting
found in our LPFC cells may be linked to differential expression of T-type currents. In some
LPFC fast-spiking C1 interneurons, depolarizing current injections elicited depolarizing events

with slow kinetics that led to long latencies of action potentials evoked at low rheobase
suggesting that at least a group of fast spiking interneurons in LPFC, likely basket cells, may
express these channels (**Fig. 4A**).

Bursting has been associated to a variety of functions such as increase in the efficiency of
synaptic transmission (Lisman, 1997; Csicsvari et al., 1998), and modulation of synaptic
plasticity by increasing temporal summation of EPSPs and Long-Term Potentiation (LTP) in
postsynaptic terminals (Thomas et al., 1998). The increase in bursting in LPFC relative to V1
may be linked to the ability of LPFC neurons to flexibly become tuned to novel objects
(Freedman et al., 2002; Fitzgerald et al., 2012; Martinez-Trujillo, 2022). One interpretation of our
results is that neurons in early areas such as V1 consistently process basic visual features (high
stability - low plasticity), which is necessary for encoding the same sensory information (e.g.,
color, orientation) in a stable manner, regardless of whether it belongs to familiar or novel
objects. On the other hand, neurons in the LPFC rapidly learn new associations of these basic
features (low stability - high plasticity), which is necessary when encountering novel complex
objects. Previous studies have reported a gradient in the ratio of NMDA- to AMPA-receptors
(Yang et al., 2018; Froudist-Walsh et al., 2023) along the hierarchy of visual processing with
LPFC having a high amount of NMDA glutamate receptors compared to V1. Combined with
intrinsic bursting, NMDA gradients can lead to the enhanced ability of LPFC to build selectivity
for new categories or rules (Freedman et al., 2001; Rouzitalab et al., 2023; Abbass et al., 2024).
Our results can be interpreted within the framework of the feature integration theory of
perception and attention (Treisman and Gelade, 1980; Treisman, 1982; Humphreys, 2015), in
which essential features pre-attentively processed in early visual areas such as V1 and are
latter assembled by the mechanisms of attention in the LPFC (Martinez-Trujillo, 2022).

**Serial homology and neuronal diversity in the primate visual system**
It has been proposed that in the mammalian neocortex, cell types follow the principle of *serial*
*homology*: cell types across brain areas are variations on a common theme organized in the
same basic pattern. Differences between serially homologous structures are typically
quantitative rather than qualitative (Harris and Shepherd, 2015). However, much of the data
supporting this hypothesis have been collected in a single species, the mouse, and in sensory
areas (e.g., barrel cortex or V1, Magrou et al., 2024). Mice do not show the stark differences in
the morphology of pyramidal cells between V1 and PFC (Gilman et al., 2016) and perhaps the
principle of serial homology applies to a larger degree than in primates. Previous studies
observed a strong difference in structural complexity, from branching patterns (Gilman et al.,
2016) to increases in the number of dendritic spines (Arnsten et al., 2012; Yang et al., 2013)
between V1 and LPFC neurons. Moreover, we see qualitative differences in firing patterns such
as the emergence of intrinsic bursting in LPFC. Thus, primate brains with expanded neocortices
seem to have diversified single neuron structure and function to cope with the demands of their
ecological niches achieving more brain computational power and behavioral complexity and
making them some of the most adaptable species in the planet.
639
Inhibitory interneurons, have been classically considered less diverse across areas and species
(Tremblay et al., 2016). However, most of what we know about interneuron diversity is from
studies in mice. Interestingly, a recent study has shown substantial variation in the expression of
different types of mRNAs across various species (mice, ferrets, marmosets and human,
(Krienen et al., 2020)). In agreement with our results, several transcripts were differentially
expressed in primate interneurons depending on the area (Krienen et al., 2020), supporting the
notion of existing qualitative differences in interneurons subtypes across areas in primates.
Thus, if there is serial homology across different areas of the primate visual system, it may be

restricted to some cell types. Neuronal diversity across brain areas may be a hallmark of
primate evolution and may arise during long-protracted periods of brain development (Otani et
al., 2016). How this diversification of cell types has impacted normal brain function and
vulnerability to new categories of brain diseases affecting humans is an open question.
652

**Conclusion**

654 Our study revealed several differences in intrinsic properties of pyramidal cells and fast spiking
655 inhibitory interneurons between areas V1 and the LPFC. An important contribution of our study
656 is to show that area specialization in the primate visual system permeates the most basic level
657 of signal processing, the single neuron. The data corresponding to the manuscript are available
658 as a resource ([www.primatedatabase.com](www.primatedatabase.com)).

**Acknowledgements**

We would like to thank all veterinarians, veterinary technicians and experimenters taking care of
the animals used in the study. At Western University: Kristy Gibbs, Rhonda Kersten, Cheryl
Vander Tuin, Jarrod Dowell, Diego Buitrago Piza, Tsz Wai Bentley Lo. In Goettingen: Corinna
Boike, Diana Kaltenborn, Verena Arndt, Diana Kues, Martina Bleyer, Daniel Aschoff.

In addition, we would like to thank Irma Meteluch, Patricia Sprysch, Pavel Truschow for
excellent technical assistance around tissue processing and imaging as well as numerous
student assistants for manual reconstruction of neuronal morphology, in particular Rakshit
Sharma, Abdirahman Mohamed Salat, Katharina Bodes, Ima Mansori, Nicolas Zdun, Marvin
Luca Schmidt, Sabrina Schild, Sophia Heidenreich, Leander Isaak Matthes, Serena Lam, Luca
Auge, Josi Dittmann, Omar Aljameh, Amirhossein Haddadbakhodaei.

Furthermore, we would like to thank other members of the NeuroNex grant for giving their time
and feedback on discussion of the project, namely John Meng, Xiao Xing Wang, Takeaki
Miyamae, Benjamin Corrigan.

Last, but not least, we want to show our gratitude to Eric Kuebler, Kartik Pradeepan and William
Assis, who have created the website featured in this publication together with coauthor Sam
Mestern.

This work was primarily supported by the internation research grant NeuroNex funded by the
Canadian Institutes of Health Research (CIHR reference: FL6GV84CKN57, MF, MSJS, SM, SV,
SE, SJT, WI, JMT) , the Deutsche Forschungsgemeinschaft (DFG, Projektnummer 436260547,
SP, JR, FP, ST, AN, JFS) and the National Science Foundation (NSF, grant number: 2015276,
AFTA, DAL, GGB). JMT and WI were also supported by the Natural Sciences and Engineering
Research Council of Canada (NSERC) and BrainsCAN grants.

**Author Contribution Statement**

M Feyerabend: data curation (lead), formal analysis (lead), investigation (lead in Canada), visualization (equal), writing – original draft (equal) , writing – review & editing (support)

S Pommer: data curation (support), formal analysis (support), investigation (lead in Germany), visualization (equal), writing – original draft (equal) , writing – review & editing (support)

M Jimenez-Sosa: data curation (support), investigation (equal)

J Rachel: investigation (equal)

J Sunstrum: investigation (equal)

F Preuss: investigation (equal)

S Mestern: investigation (equal)

R Hinkel: resources (lead)

M Mietsch: resources (support)

S Viyajraghavan: resources (support)

S Everling: resources (support), funding acquisition (support)

S Treue: resources (support), funding acquisition (lead),

AFT Arnsten: funding acquisition (lead), project administration (lead), conceptualization (support)

DA Lewis: funding acquisition (support), writing – review & editing (support)

F Wolf, , S McCarroll: funding acquisition (support)

L Muller: funding acquisition (support), project administration (support)

F Krienen, J Murray: funding acquisition (support), conceptualization (support)

SJ Tripathy, G Gonzalez-Burgos: funding acquisition (support), conceptualization (support), writing – review & editing (support)

W Inoue, JF Staiger, A Neef : funding acquisition (support),  resources (support), supervision (equal), writing – review & editing (support)

J Martinez-Trujillo: funding acquisition (lead),  writing – original draft (equal) , writing – review & editing (lead), supervision (equal)

# References

1. Abbass M, Corrigan B, Johnston R, Gulli R, Sachs A, Lau JC, Martinez-Trujillo J (2024) Neural Ensembles in the Lateral Prefrontal Cortex Temporally Multiplex Task Features During Virtual Navigation. bioRxiv:2024.01.10.574378.
2. Amatrudo JM, Weaver CM, Crimins JL, Hof PR, Rosene DL, Luebke JI (2012) Influence of highly distinctive structural properties on the excitability of pyramidal neurons in monkey visual and prefrontal cortices. J Neurosci 32:13644–13660 Available at: http://www.jneurosci.org/cgi/doi/10.1523/JNEUROSCI.2581-12.2012.
3. Ardid S, Vinck M, Kaping D, Marquez S, Everling S, Womelsdorf T (2015) Mapping of Functionally Characterized Cell Classes onto Canonical Circuit Operations in Primate Prefrontal Cortex. J Neurosci 35:2975–2991.
4. Arnsten AFT (2011) Prefrontal cortical network connections: key site of vulnerability in stress and schizophrenia. Int J Dev Neurosci 29:215–223.
5. Arnsten AFT, Wang MJ, Paspalas CD (2012) Neuromodulation of thought: flexibilities and vulnerabilities in prefrontal cortical network synapses. Neuron 76:223–239 Available at: https://linkinghub.elsevier.com/retrieve/pii/S0896627312008045.
6. Ascoli et al. (2008) Petilla terminology: nomenclature of features of GABAergic interneurons of the cerebral cortex. Nat Rev Neurosci 9:557–568 Available at: http://www.nature.com/articles/nrn2402.
7. Bakken TE et al. (2016) Comprehensive transcriptional map of primate brain development. Nature 535:367–375.
8. Berg J et al. (2021) Human neocortical expansion involves glutamatergic neuron diversification. Nature 598:151–158.
9. Bruno RM, Hahn TTG, Wallace DJ, Kock CPJ de, Sakmann B (2009) Sensory Experience Alters Specific Branches of Individual Corticocortical Axons during Development. J Neurosci 29:3172–3181.
10. Carandini M, Heeger DJ (2011) Normalization as a canonical neural computation. Nat Rev Neurosci 13:51–62 Available at: http://www.nature.com/articles/nrn3136.
11. Condé F coise, Lund JS, Jacobowitz DM, Baimbridge KG, Lewis DA (2004) Local circuit neurons immunoreactive for calretinin, calbindin D-28k or parvalbumin in monkey prefronatal cortex: Distribution and morphology. J Comp Neurol 341:95–116 Available at: http://doi.wiley.com/10.1002/cne.903410109.
12. Corrigan BW, Gulli RA, Doucet G, Roussy M, Luna R, Pradeepan KS, Sachs AJ, Martinez-Trujillo JC (2022) Distinct neural codes in primate hippocampus and lateral prefrontal cortex during associative learning in virtual environments. Neuron 110:2155-2169.e4.
13. Csicsvari J, Hirase H, Czurko A, Buzsáki G (1998) Reliability and State Dependence of Pyramidal Cell–Interneuron Synapses in the Hippocampus an Ensemble Approach in the Behaving Rat. Neuron 21:179–189.
14. Cyranoski D (2014) Marmosets are stars of Japan's ambitious brain project. Nature 514:151–152 Available at: http://www.nature.com/doifinder/10.1038/514151a.
15. Defelipe J et al. (2013) New insights into the classification and nomenclature of cortical GABAergic interneurons. Nat Rev Neurosci 14:202–216 Available at: http://www.nature.com/doifinder/10.1038/nrn3444.
16. Felleman DJ, Essen DCV (1991) Distributed hierarchical processing in the primate cerebral cortex. Cereb Cortex 1:1–47 Available at:

http://eutils.ncbi.nlm.nih.gov/entrez/eutils/elink.fcgi?dbfrom=pubmed&id=1822724&retmode=ref&cmd=prlinks.

17. Fitzgerald JK, Swaminathan SK, Freedman DJ (2012) Visual categorization and the parietal cortex. Front Integr Neurosci 6:18.

18. Freedman DJ, Assad JA (2006) Experience-dependent representation of visual categories in parietal cortex. Nature 443:85–88 Available at: http://www.nature.com/doifinder/10.1038/nature05078.

19. Freedman DJ, Riesenhuber M, Poggio T, Miller EK (2001) Categorical Representation of Visual Stimuli in the Primate Prefrontal Cortex. Science 291:312–316 Available at: http://www.sciencemag.org/cgi/doi/10.1126/science.291.5502.312.

20. Freedman DJ, Riesenhuber M, Poggio T, Miller EK (2002) Visual Categorization and the Primate Prefrontal Cortex: Neurophysiology and Behavior. J Neurophysiol 88:929–941.

21. Ghose GM, Yang T, Maunsell JHR (2002) Physiological Correlates of Perceptual Learning in Monkey V1 and V2. J Neurophysiol 87:1867–1888.

22. Gilman JP, Medalla M, Luebke JI (2017) Area-Specific Features of Pyramidal Neurons-a Comparative Study in Mouse and Rhesus Monkey. Cereb Cortex 27:2078–2094 Available at: http://cercor.oxfordjournals.org/cgi/doi/10.1093/cercor/bhw062.

23. Glantz LA, Lewis DA (2000) Decreased Dendritic Spine Density on Prefrontal Cortical Pyramidal Neurons in Schizophrenia. Arch Gen Psychiat 57:65–73 Available at: http://archpsyc.jamanetwork.com/article.aspx?doi=10.1001/archpsyc.57.1.65.

24. Godlove DC, Maier A, Woodman GF, Schall JD (2014) Microcircuitry of agranular frontal cortex: testing the generality of the canonical cortical microcircuit. J Neurosci 34:5355–5369 Available at: http://www.jneurosci.org/cgi/doi/10.1523/JNEUROSCI.5127-13.2014.

25. Gonzalez-Burgos G, Barrionuevo G, Lewis DA (2000) Horizontal synaptic connections in monkey prefrontal cortex: an in vitro electrophysiological study. Cereb Cortex 10:82–92 Available at: http://eutils.ncbi.nlm.nih.gov/entrez/eutils/elink.fcgi?dbfrom=pubmed&id=10639398&retmode=ref&cmd=prlinks.

26. González-Burgos G, Miyamae T, Krimer Y, Gulchina Y, Pafundo DE, Krimer O, Bazmi H, Arion D, Enwright JF, Fish KN, Lewis DA (2019) Distinct Properties of Layer 3 Pyramidal Neurons from Prefrontal and Parietal Areas of the Monkey Neocortex. J Neurosci 39:7277–7290 Available at: http://www.jneurosci.org/lookup/doi/10.1523/JNEUROSCI.1210-19.2019.

27. Goriounova NA, Heyer DB, Wilbers R, Verhoog MB, Giugliano M, Verbist C, Obermayer J, Kerkhofs A, Smeding H, Verberne M, Idema S, Baayen JC, Pieneman AW, Kock CP de, Klein M, Mansvelder HD (2018) Large and fast human pyramidal neurons associate with intelligence. eLife 7:e41714.

28. Gouwens NW et al. (2019) Classification of electrophysiological and morphological neuron types in the mouse visual cortex. Nat Neurosci 22:1182–1195 Available at: http://www.nature.com/articles/s41593-019-0417-0.

29. Harris KD, Shepherd GMG (2015) The neocortical circuit: themes and variations. Nat Neurosci 18:170–181.

30. Hentschke H, Stüttgen MC (2011) Computation of measures of effect size for neuroscience data sets. Eur J Neurosci 34:1887–1894.

31. Hu H, Cavendish JZ, Agmon A (2013) Not all that glitters is gold: off-target recombination in the somatostatin–IRES-Cre mouse line labels a subset of fast-spiking interneurons. Front Neural Circuits 7:195.

32. Humphreys GW (2015) Feature confirmation in object perception: Feature integration theory 26 years on from the Treisman Bartlett lecture. Q J Exp Psychol (2006) 69:1910–1940.

33. Jahnsen H, Llinás R (1984) Ionic basis for the electro‑responsiveness and oscillatory
properties of guinea‑pig thalamic neurones in vitro. J Physiol 349:227–247.
34. Jiang X, Shen S, Cadwell CR, Berens P, Sinz F, Ecker AS, Patel S, Tolias AS (2015)
Principles of connectivity among morphologically defined cell types in adult neocortex.
Science 350:aac9462–aac9462 Available at:
http://www.sciencemag.org/cgi/doi/10.1126/science.aac9462.
35. Kalmbach BE et al. (2018) h-Channels Contribute to Divergent Intrinsic Membrane
Properties of Supragranular Pyramidal Neurons in Human versus Mouse Cerebral
Cortex. Neuron 100:1194-1208.e5 Available at:
https://linkinghub.elsevier.com/retrieve/pii/S0896627318309000.
36. Krienen FM et al. (2020) Innovations present in the primate interneuron repertoire.
Nature 586:262–269 Available at: http://www.nature.com/articles/s41586-020-2781-z.
37. Lennert T, Martinez-Trujillo JC (2013) Prefrontal neurons of opposite spatial preference
display distinct target selection dynamics. J Neurosci 33:9520–9529 Available at:
http://www.jneurosci.org/cgi/doi/10.1523/JNEUROSCI.5156-12.2013.
38. Lewis DA, González-Burgos G (2008) Neuroplasticity of neocortical circuits in
schizophrenia. Neuropsychopharmacol 33:141–165 Available at:
http://www.nature.com/articles/1301563.
39. Lewis DA, Sweet RA (2009) Schizophrenia from a neural circuitry perspective:
advancing toward rational pharmacological therapies. J Clin Investig 119:706–716.
40. Lisman JE (1997) Bursts as a unit of neural information: making unreliable synapses
reliable. Trends Neurosci 20:38–43.
41. Luebke JI, Medalla M, Amatrudo JM, Weaver CM, Crimins JL, Hunt B, Hof PR, Peters A
(2015) Age-related changes to layer 3 pyramidal cells in the rhesus monkey visual
cortex. Cereb Cortex 25:1454–1468 Available at:
https://academic.oup.com/cercor/article-lookup/doi/10.1093/cercor/bht336.
42. Magrou L, Joyce MKP, Froudist-Walsh S, Datta D, Wang X-J, Martinez-Trujillo J,
Arnsten AFT (2024) The meso-connectomes of mouse, marmoset, and macaque:
network organization and the emergence of higher cognition. Cereb Cortex 34:bhae174.
43. Martinez-Trujillo J (2022) Visual Attention in the Prefrontal Cortex. Annu Rev Vis Sci
8:407–425.
44. McInnes L, Healy J, Saul N, Großberger L (2018) UMAP: Uniform Manifold
Approximation and Projection. J Open Source Softw 3:861.
45. Medalla M, Gilman JP, Wang J-Y, Luebke JI (2017) Strength and Diversity of Inhibitory
Signaling Differentiates Primate Anterior Cingulate from Lateral Prefrontal Cortex. J
Neurosci 37:4717–4734 Available at:
http://www.jneurosci.org/lookup/doi/10.1523/JNEUROSCI.3757-16.2017.
46. Medalla M, Luebke JI (2015) Diversity of glutamatergic synaptic strength in lateral
prefrontal versus primary visual cortices in the rhesus monkey. J Neurosci 35:112–127
Available at: http://www.jneurosci.org/cgi/doi/10.1523/JNEUROSCI.3426-14.2015.
47. Mendoza-Halliday D, Torres S, Martinez-Trujillo JC (2014) Sharp emergence of feature-
selective sustained activity along the dorsal visual pathway. Nat Neurosci 17:1255–1262
Available at: http://www.nature.com/doifinder/10.1038/nn.3785.
48. Merigan WH, Maunsell JHR (1993) How Parallel are the Primate Visual Pathways?
Annu Rev Neurosci 16:369–402.
49. Merino RM, Leon-Pinzon C, Stühmer W, Möck M, Staiger JF, Wolf F, Neef A (2019)
Background correlations selectively boost the gamma-sensitivity of cortical GABAergic
neurons. Biorxiv 12:2019.12.19.882639 Available at:
http://biorxiv.org/lookup/doi/10.1101/2019.12.19.882639.

50. Miller CT, Freiwald WA, Leopold DA, Mitchell JF, Silva AC, Wang X (2016) Marmosets: A Neuroscientific Model of Human Social Behavior. Neuron 90:219–233 Available at: http://linkinghub.elsevier.com/retrieve/pii/S0896627316300071.

51. Mitchell JF, Leopold DA (2015) The marmoset monkey as a model for visual neuroscience. Neurosci Res 93:20–46 Available at: http://linkinghub.elsevier.com/retrieve/pii/S0168010215000267.

52. Mitchell JF, Reynolds JH, Miller CT (2014) Active Vision in Marmosets: A Model System for Visual Neuroscience. J Neurosci 34:1183–1194.

53. Mohan H, Verhoog MB, Doreswamy KK, Eyal G, Aardse R, Lodder BN, Goriounova NA, Asamoah B, Brakspear ABCB, Groot C, Sluis S van der, Testa-Silva G, Obermayer J, Boudewijns ZSRM, Narayanan RT, Baayen JC, Segev I, Mansvelder HD, Kock CPJ de (2015) Dendritic and Axonal Architecture of Individual Pyramidal Neurons across Layers of Adult Human Neocortex. Cereb Cortex (N York, NY) 25:4839–4853.

54. Nassi JJ, Callaway EM (2009) Parallel processing strategies of the primate visual system. Nat Rev Neurosci 10:360–372.

55. Oberlaender M, Boudewijns ZSRM, Kleele T, Mansvelder HD, Sakmann B, Kock CPJ de (2011) Three-dimensional axon morphologies of individual layer 5 neurons indicate cell type-specific intracortical pathways for whisker motion and touch. Proc Natl Acad Sci 108:4188–4193.

56. Okano H, Sasaki E, Yamamori T, Iriki A, Shimogori T, Yamaguchi Y, Kasai K, Miyawaki A (2016) Brain/MINDS: A Japanese National Brain Project for Marmoset Neuroscience. Neuron 92:582–590.

57. Otani T, Marchetto MC, Gage FH, Simons BD, Livesey FJ (2016) 2D and 3D Stem Cell Models of Primate Cortical Development Identify Species-Specific Differences in Progenitor Behavior Contributing to Brain Size. Cell Stem Cell 18:467–480.

58. Panagiotaropoulos TI (2024) An integrative view of the role of prefrontal cortex in consciousness. Neuron 112:1626–1641.

59. Povysheva NV, Zaitsev AV, Kröner S, Krimer OA, Rotaru DC, Gonzalez-Burgos G, Lewis DA, Krimer LS (2007) Electrophysiological differences between neurogliaform cells from monkey and rat prefrontal cortex. J Neurophysiol 97:1030–1039 Available at: http://www.physiology.org/doi/10.1152/jn.00794.2006.

60. Povysheva NV, Zaitsev AV, Rotaru DC, Gonzalez-Burgos G, Lewis DA, Krimer LS (2008) Parvalbumin-positive basket interneurons in monkey and rat prefrontal cortex. J Neurophysiol 100:2348–2360 Available at: https://www.physiology.org/doi/10.1152/jn.90396.2008.

61. Preuss TM, Wise SP (2022) Evolution of prefrontal cortex. Neuropsychopharmacology 47:3–19.

62. Reynolds JH, Chelazzi L, Desimone R (1999) Competitive Mechanisms Subserve Attention in Macaque Areas V2 and V4. J Neurosci 19:1736–1753 Available at: http://www.jneurosci.org/lookup/doi/10.1523/JNEUROSCI.19-05-01736.1999.

63. Roussy M, Busch A, Luna R, Leavitt ML, Mofrad MH, Gulli RA, Corrigan B, Mináč J, Sachs AJ, Palaniyappan L, Muller L, Martinez-Trujillo JC (2022a) Neural sequences in primate prefrontal cortex encode working memory in naturalistic environments. Biorxiv:2022.08.18.504406.

64. Roussy M, Corrigan B, Luna R, Gulli RA, Sachs AJ, Palaniyappan L, Martinez-Trujillo JC (2022b) Stable working memory and perceptual representations in macaque lateral prefrontal cortex during naturalistic vision. J Neurosci 42:JN-RM-0597-22.

65. Rouzitalab A, Boulay CB, Park J, Martinez-Trujillo JC, Sachs AJ (2023) Ensembles code for associative learning in the primate lateral prefrontal cortex. Cell Rep 42:112449.

66. Rübel O, Tritt A, Ly R, Dichter BK, Ghosh S, Niu L, Baker P, Soltesz I, Ng L, Svoboda K, Frank L, Bouchard KE (2022) The Neurodata Without Borders ecosystem for neurophysiological data science. eLife 11:e78362.

67. Saunders A, Macosko EZ, Wysoker A, Goldman M, Krienen FM, Rivera H de, Bien E, Baum M, Bortolin L, Wang S, Goeva A, Nemesh J, Kamitaki N, Brumbaugh S, Kulp D, McCarroll SA (2018) Molecular Diversity and Specializations among the Cells of the Adult Mouse Brain. Cell 174:1015-1030.e16 Available at: https://linkinghub.elsevier.com/retrieve/pii/S0092867418309553.

68. Schmolesky MT, Wang Y, Hanes DP, Thompson KG, Leutgeb S, Schall JD, Leventhal AG (1998) Signal Timing Across the Macaque Visual System. J Neurophysiol 79:3272–3278.

69. Seiffert C, Khoshgoftaar TM, Hulse JV, Napolitano A (2010) RUSBoost: A Hybrid Approach to Alleviating Class Imbalance. IEEE Trans Syst, Man, CybernPart A: Syst Hum 40:185–197.

70. Servick K (2018) Why are U.S. neuroscientists clamoring for marmosets? Science Available at: https://www.sciencemag.org/news/2018/10/why-are-us-neuroscientists-are-clamoring-marmosets.

71. Sincich LC, Horton JC (2005) THE CIRCUITRY OF V1 AND V2: Integration of Color, Form, and Motion. Annu Rev Neurosci 28:303–326.

72. Szentágothai J (1990) Cajal on the cerebral cortex (An annotated translation of the complete writings). By J. de Felipe and E. G. Jones. Oxford University Press, Oxford. 654 pp. $65.00. Neuroscience 36:569–570.

73. Thomas MJ, Watabe AM, Moody TD, Makhinson M, O'Dell TJ (1998) Postsynaptic complex spike bursting enables the induction of LTP by theta frequency synaptic stimulation. J Neurosci 18:7118–7126 Available at: /pmc/articles/PMC6793261/.

74. Ting JT, Kalmbach B, Chong P, Frates R de, Keene CD, Gwinn RP, Cobbs C, Ko AL, Ojemann JG, Ellenbogen RG, Koch C, Lein E (2018) A robust ex vivo experimental platform for molecular-genetic dissection of adult human neocortical cell types and circuits. Sci Rep-uk 8:8407–8413 Available at: http://www.nature.com/articles/s41598-018-26803-9.

75. Torres-Gomez S, Blonde JD, Mendoza-Halliday D, Kuebler E, Everest M, Wang X-J, Inoue W, Poulter MO, Martinez-Trujillo J (2020) Changes in the Proportion of Inhibitory Interneuron Types from Sensory to Executive Areas of the Primate Neocortex: Implications for the Origins of Working Memory Representations. Cereb Cortex 9:557 Available at: https://academic.oup.com/cercor/advance-article/doi/10.1093/cercor/bhaa056/5811846.

76. Treisman A (1982) Perceptual grouping and attention in visual search for features and for objects. J Exp Psychology Hum Percept Perform 8:194–214.

77. Treisman AM, Gelade G (1980) A feature-integration theory of attention. Cogn Psychol 12:97–136.

78. Tremblay R, Lee S, Rudy B (2016) GABAergic Interneurons in the Neocortex: From Cellular Properties to Circuits. Neuron 91:260–292.

79. Williams SR, Tóth TI, Turner JP, Hughes SW, Crunelli V (1997) The 'window' component of the low threshold Ca2+ current produces input signal amplification and bistability in cat and rat thalamocortical neurones. J Physiol 505:689–705.

80. Yang S-T, Wang M, Paspalas CD, Crimins JL, Altman MT, Mazer JA, Arnsten AFT (2018) Core Differences in Synaptic Signaling Between Primary Visual and Dorsolateral Prefrontal Cortex. Cereb Cortex 28:1458–1471 Available at: https://academic.oup.com/cercor/article/28/4/1458/4812491.

81. Yang Y, Paspalas CD, Jin LE, Picciotto MR, Arnsten AFT, Wang M (2013) Nicotinic $\alpha$7 receptors enhance NMDA cognitive circuits in dorsolateral prefrontal cortex.

Proc National Acad Sci 110:12078–12083 Available at:
http://www.pnas.org/cgi/doi/10.1073/pnas.1307849110.

82. Young ME, Ohm DT, Dumitriu D, Rapp PR, Morrison JH (2014) Differential effects of aging on dendritic spines in visual cortex and prefrontal cortex of the rhesus monkey. Neuroscience 274:33–43.

83. Yuste R et al. (2020) A community-based transcriptomics classification and nomenclature of neocortical cell types. Nat Neurosci 23:1456–1468 Available at: http://www.nature.com/articles/s41593-020-0685-8.

84. Zaitsev AV, Gonzalez-Burgos G, Povysheva NV, Kröner S, Lewis DA, Krimer LS (2005) Localization of calcium-binding proteins in physiologically and morphologically characterized interneurons of monkey dorsolateral prefrontal cortex. Cereb Cortex 15:1178–1186 Available at: http://academic.oup.com/cercor/article/15/8/1178/304661/Localization-of-Calciumbinding-Proteins-in.

85. Zaitsev AV, Povysheva NV, Gonzalez-Burgos G, Lewis DA (2012) Electrophysiological classes of layer 2/3 pyramidal cells in monkey prefrontal cortex. J Neurophysiol 108:595–609 Available at: https://www.physiology.org/doi/10.1152/jn.00859.2011.

86. Zaitsev AV, Povysheva NV, González-Burgos G, Rotaru D, Fish KN, Krimer LS, Lewis DA (2009) Interneuron diversity in layers 2-3 of monkey prefrontal cortex. Cereb Cortex 19:1597–1615 Available at: https://academic.oup.com/cercor/article-lookup/doi/10.1093/cercor/bhn198.

87. Zeldenrust F, Wadman WJ, Englitz B (2018) Neural Coding With Bursts-Current State and Future Perspectives. Front Comput Neurosc 12:48 Available at: http://eutils.ncbi.nlm.nih.gov/entrez/eutils/elink.fcgi?dbfrom=pubmed&id=30034330&retmode=ref&cmd=prlinks.

88. Zhou X, Mansori I, Fischer T, Witte M, Staiger JF (2020) Characterizing the morphology of somatostatin-expressing interneurons and their synaptic innervation pattern in the barrel cortex of the GFP-expressing inhibitory neurons mouse. J Comp Neurol 528:244–260 Available at: https://onlinelibrary.wiley.com/doi/abs/10.1002/cne.24756.

992 # Tables

993 *Table 1: Electrophysiological characteristics of class 3 neurons in LPFC and V1.*

| Feature | LPFC n = 104 | V1 n = 56 | adj. p-value | z-val. | Effect size [CI] |
|---|---|---|---|---|---|
| Vrest, mV | -71.61 | -71.68 | 0.6721 | 0.47 | -0.04 [-0.19; 0.11] |
| $R_{inHD}$, MΩ | 212.6 | 461.4 | <1e-4* | -7.80 | 0.62 [0.53; 0.70] |
| rheobase, pA | 90 | 30 | <1e-4* | 7.95 | -0.63 [-0.71;-0.54] |
| tau, ms | 26.74 | 30.28 | 0.2640 | -1.18 | 0.09 [-0,06; 0.25] |
| sag ratio | 1.13 | 1.21 | 0.0012* | -3.45 | 0.27 [0.11;0.43] |
| inst. rectification | 0.92 | 0.75 | <1e-4* | 8.08 | -0.64 [-0.73; -0.54] |
| AP width | 0.35 | 0.27 | <1e-4* | 5.28 | -0.42 [-0.54; -0.29] |
| amp slow trough | -19.65 | -21.65 | 0.0127* | 2.60 | -0.21 [-0.36; -0.05] |
| rate of rheo, Hz | 1.00 | 1.00 | 0.1274 | -1.64 | 0.13 [-0.03; 0.30] |
| med. inst. rate, Hz | 22.2 | 19.3 | 0.0116* | -2.65 | 0.22 [0.07; 0.37] |
| IQR inst. rate, Hz | 7.30 | 8.10 | 0.2234 | 1.32 | -0.11 [-0.27; 0.05] |
| adaRat Blast/B1 | 0.225 | 0.048 | 0.0032* | -3.14 | 0.25 [0.11; 0.39] |
| fIslope, Hz/pA | 0.048 | 0.123 | <1e-4* | -5.07 | 0.41 [0.26; 0.53] |
| cvISI | 0.354 | 0.249 | 2.25e-4* | 3.90 | -0.34 [-0.49; -0.17] |
| rate of hero, Hz | 8 | 13 | 0.0116* | -2.68 | 0.21 [0.06; 0.36] |
| latency at hero, ms | 14.35 | 16.41 | 0.9544 | 0.06 | 0.00 [-0.17; 0.16] |
| adaptation index | 0.091 | 0.048 | <1e-4* | 4.51 | -0.40 [-0.55; -0.25] |
| difference in trough | 4.608 | 6.760 | 0.0053* | -2.96 | 0.24 [0.09; 0.39] |
| burst index | 0.888 | 0.782 | <1e-4* | 4.48 | -0.39 [-0.52; -0.24] |

995 **Table 1: Comparison between V1 and LPFC in key electrophysiological features of C3 Cells**
996 The measure of central tendency of the distribution median. Negative effect size measures show
997 smaller values for V1 compared to PFC. p Values are corrected with Benjamini-Hochberg
998 procedure.

1000

1001

1002 *Table 2: Comparison between V1 and LPFC in key morphological features of class 1 and 3 neurons.*

| Feature | LPFC n = 11/5 | V1 n = 4/3 | adj. p-value | z-val. | Effect size [CI] |
|---|---|---|---|---|---|
| # primary dendrites (C3) | 6 | 6 | 0.7341 | -0.71 | -0.11 [-0.66; 0.53] |
| apical length (C3) | 2.8321e+03 | 972.1027 | 0.0352* | -2.07 | -0.63 [-0.78; -0.28] |
| apical complexity (C3) | 5.4943e+05 | 7.3401e+04 | 0.0352* | -2.23 | -0.63 [-0.71;-0.54] |
| basal length (C3) | 2.6794e+03 | 1.8978e+03 | 0.0835 | -1.83 | -0.52 [-0.77; -0.14] |
| basal complexity (C3) | 4.5996e+04 | 3.0933e+04 | 0.1653 | -2.07 | -0.63 [-0.78; -0.35] |
| # primary dendrites (C1) | 5 | 8 | 0.2976 | - | 0.57 [0; 0.91] |
| tot. dendritic length (C1) | 3.9230e+03 | 1.00 | 0.7857 | - | 0.17[-0.54; 0.88] |
| dendritic complexity (C1) | 5.8131e+04 | 7.4153e+04 | 0.7143 | - | 0.28 [-0.52; 0.88] |
| total axonal length (C1) | 1.9743e+04 | 9.1722e+03 | 0.2976 | - | -0.73 [-0.89; -0.40] |
| axonal complexity (C1) | 52045624 | 9163046 | 0.2976 | - | -0.62 [-0.89; 0.06] |

1003

1004 **Table 2.** Values are median of electrophysiological characteristics in class 1 cells of LPFC and V1.
1005 Effect sizes were calculated as rank-biserial correlation. Negative values indicate larger LPFC
1006 values.

1007

# Figures

**Figure 1: Tissue acquisition and characterization of neuronal morphology and electrophysiology in multiple areas of the marmoset neocortex.**
*A: Acute slices of LPFC and V1 were obtained from 42 marmosets with a median age of 4.4 years. Neurons were characterized by electrophysiology (B) and morphology (C). B: Cells (grayscale picture) were subjected to 1 sec long square pulse current injection with a 20 pA increment from -110 pA till suprathreshold saturation. Example traces show hyperpolarizing, subthreshold, rheobase and suprathreshold injections. Data was acquired across different laboratories and recording systems (right pie chart). C, D: Anatomical evaluation of all filled neurons consisted of determination of broad class (spiny, putative Excitatory, pEXC versus aspiny, putative Inhibitory, pINH) via somatodendritic configuration, spininess (C) and cortical home layer of the soma (D). Left image in C shows a biocytin filled pyramidal neuron in layer III of LPFC; nuclei are stained with DAPI (blue); scale bar equals 150 µm. Small images show example spiny (top) and aspiny (bottom) somatodendritic morphology; scale bars equal 20 µm. Pie-chart shows the distribution of morphologically identified cells. In D the image shows NeuN staining with layer delineation in LPFC and V1; scale bars equal 100 µm. Right pie-charts represent the distribution of cells by layers for LPFC and V1. Cells recorded in both areas were predominantly located in supragranular (I-III) layers.*

1026

**Figure 2: Neuronal diversity within marmoset data and co-projection with known cell types.**
*A, B: Examples of different marmoset cells separated by cortical area (A: V1, B: LPFC). Cells 1 and 4 are regular-spiking pyramidal cells. Cell 2 is a fast-spiking basket cell. Cell 3 is a vertically oriented basket cell. The cell was recorded in the dorsal medial V1 with expanded layers IV–VI. Cell 5 is a neurogliaform cell. Cell 6 is a double bouquet cell and cell 7 is a regular spiking spiny non-pyramidal cell (Below each cell is the corresponding subthreshold, rheobase and hero sweep. Scale bars equal 100 µm. C: Panel of 2D UMAP projections visualizing distribution of cells by species and key electrophysiological features. D: 2D UMAP projection visualizing distribution of marmoset cells color-coded by dendritic type, referenced with known cell type in mouse. Colored background delineates areas with high density for the respective cell type. Location of examples from subfigures A and B are indicated by their respective color (downward facing triangle: V1, circle: PFC). The Logos indicate the origin of the data. Allen Institute in blue/black. Our NEURONEX consortium in black + country flags.*

1038

**Figure 3: Objective classification of primate cells.**
**A:** Panel of pie charts showing cell type composition of training and test data. Three different classes were used: Class 1 (dark blue; C1), Class 2 (ocher, C2), merged from SST (yellow), VIP (purple) and NGF (turquoise) cells and Class 3 (dark green, C3). Mouse and marmoset test data had identical proportion of classes. For marmoset test data we chose electrophysiology with congruent morphology for C1 and C2, respectively. Marmoset C3 cells were drawn from spiny cells. The procedure was repeated 500 times with randomly drawn training and test sets. **B:** Performance of classifier across repetitions. Horizontal boxplots show distribution of performances. Median test performances by species were 91.3% for mouse and 83.3 % for marmoset. **C:** Confusion matrices for mouse and marmoset test data showing all classification totals. The positive predictive value (PPV) for C1 and C3 cells is over 90% in the marmoset. FDR = False Discovery Rate, TPR = True Positive Rate, FNR = False Negative Rate. **D:** Example UMAP visualization of one model prediction referenced with dendritic type of marmoset cells. Shaded areas indicate high density of the respective class. **E:** Colormap visualizing prediction outcome of marmoset cells for all 500 repetitions. Cells are grouped by dendritic type and then sorted by the first UMAP component. Color scale indicates class prediction same as in D. **F:** Bar charts showing proportion of dendritic type per majority classifier prediction.

1053

**Figure 4: Input resistance and bursting across areas**

*A, D: Violin plots of class 3 (V1, turquois; PFC, orange) and class 1 (V1: blue, LPFC: pink) for input resistance (A)*
*and burst index (D). Black contours signify example cells (triangle: V1, circle: LPFC) shown in B (for A) and E (for D).*
*Asterisk in D labels bursting class 3 neuron. B: Above: Dendritic morphology of two example of class 3 cells (LPFC:*
*orange, circle in A; V1: turquois, triangle in A) together with 4 subthreshold traces. Below: UI curve showing how input*
*resistance of cells above was determined as the slope of linear fit of three lowest subthreshold stimulation. C: Total*
*dendritic length and input resistance strongly correlate in class 3 cells. E: Example of bursting (V1, dark green; LPFC,*
*brown) and non-bursting (V1, turquois; LPFC, orange) class 3 cells: Sweeps above the dashed line are from*
*rheobase and show the first 0.5 s, instead of the first 0.25 s. Bursting is marked by doublets at medium to high*
*stimulation, but not on rheobase level. F: Comparison of burst index of class 3 cells split by burst/non-burst*
*distinction.*

1065

**Figure 5: A bursting phenotype in C1 fast spiking basket cells**
*A: Example of various firing patterns of class 1 cells: neurons were divided into groups depending on bursting or lack*
*thereof and cortical area: Bursting is marked by multiple spikes at rheobase level. B: Number of burst-type fast*
*spiking basket cells among class 1 neurons in LPFC and V1. Bursting was observed in 28 % of LPFC class 1 cells*
*(purple vs pink) and not found in V1 (blue). C: Example morphology of bursting and non-bursting class 1 cells in*
*LPFC. The dendritic tree is shown in purple (bursting) pink (non-bursting) while the axon is shown in dark blue. The*
*asterisk corresponds to the example traces in A. Both groups are associated with basket cells. D: Burst index across*
*stimulus intensity by subpopulation (burst-LPFC/non-burst-LPFC/V1). Line indicates median, shaded area indicates*
*the IQR. E: Burst index negatively correlates with residual rheobase after regression with input resistance and*
*difference between AP threshold and resting membrane potential. Markers are blue triangles for V1 cells, pink circles*
*for non-bursting LPFC and purple stars for bursting LPFC cells. F: Burst index correlates with logarithm of latency*
*and depolarized trough of the rheobase AP. G: Box charts comparing both logarithm of latency (left y-axis) and AP*
*trough (right y-axis) by cortical area.*

1079

**Figure 6: Comparison of morphology of pyramidal cells and fast spiking basket cells across areas.**
**A, B:** *Examples of reconstructed marmoset neurons for area (left: V1, right: LPFC) and cell type of interest (A: pyramidal cells, B: fast spiking interneurons). Dendrites of pyramidal cells are colored in a lighter shade of orange for LPFC and in turquoise for V1, while the axons are in dark green. Dendrites of fast spiking basket cells were colored in pink for LPFC and light blue for V1, while axons are in dark blue. Below each neuron is the corresponding hero, rheobase and subthreshold response to the square-pulse stimulus. **C, D:** Parallel plots of key morphological quantifications for pyramidal neurons (C) and fast spiking basket cells (D). Transparent areas represent the range of the morphological feature for PFC and V1 respectively, while lines show the average for each brain region.*