Zero valent sulfur is the product of assimilatory sulfite reductase and a substrate of cysteine synthase

Qun Cao, [1,2,3†] Xuanyu Liu,[1†] Qingda Wang,[1] Yongzhen Xia,[1] Luying Xun,[1,4]

Huaiwei Liu[1,*]

[1] State Key Laboratory of Microbial Technology, Shandong University,

Qingdao, 266200, People's Re-public of China.

[2] Department of Neurosurgery, Qilu Hospital, Cheeloo College of Medicine and

Institute of Brain and Brain-Inspired Science, Shandong University, Jinan, 250012,

China.

[3] Jinan Microecological Biomedicine Shandong Laboratory and Shandong Key

Laboratory of Brain Health and Function Remodeling, Jinan, 250017, China.

[4] School of Molecular Biosciences, Washington State University, Pullman,

WA, 991647520, USA.

[†] These authors contribute equally to this study.

[*] Correspondence: liuhuaiwei@sdu.edu.cn

Abstract

The long process of the assimilatory sulfate-reduction consists of sulfate reduction

to H_2S and then H_2S assimilation for cysteine synthesis. However, recent studies

indicate that both H_2S and zero valent sulfur (S^0) are produced during dissimilatory

sulfate reduction process, which leads to the question whether S^0 also is involved in

assimilatory sulfate-reduction. Herein, phylogenetic analysis grouped assimilatory

sulfite reductases into five clusters and we re-examined four representatives from two

major clusters. These enzymes produced both S^0 and H_2S from sulfite. However,

S^0 was more like a direct product, while H_2S was just a derivative of S^0 via chemical

reduction. Phylogenetic analysis of cysteine synthases grouped them into six clusters,

and we re-examined six representatives from two major clusters. These enzymes used

both glutathione persulfide (GSSH, a common intracellular compound containing a

S^0 atom) and H_2S as substrates to synthesize cysteine, but preferred to use GSSH.

These findings indicated that S^0, other than H_2S, is the key intermediate during the

process of inorganic sulfur converting to organic sulfur.

Keywords

Assimilatory sulfate-reduction; hydrogen sulfide; zero valent sulfur; sulfite

reductase; cysteine synthase

Importance

Sulfate reduction is an important link in the global sulfur cycle. Assimilatory sulfate-

reduction consists of sulfate-reduction to H_2S and then H_2S assimilation for cysteine

synthesis. This concept was founded in 1950s and has not been challenged until today.

In this study, we re-examined four representative assimilatory sulfite reductases

(aSiRs) and found that these enzymes directly reduced sulfite to zero valent sulfur

(S^0). A small portion of produced S^0 was reduced to H_2S via aSiR-independent

chemical reactions. Further, we re-examined six representative cysteine synthases and

found that these enzymes used S^0 derivative compounds (thiosulfate and glutathione

persulfide) as substrates to synthesize cysteine. Thus, S^0 is the key intermediate of

assimilatory sulfate reduction process, other than H_2S.

Introduction

Sulfate reduction is a critical step of the sulfur biogeochemical cycle in which microorganisms play important roles. Depending on the fate of reduced sulfur atom, sulfate reduction process can be categorized into two types, dissimilatory sulfate reduction (DSR) and assimilatory sulfate reduction (ASR) (1, 2). DSR mainly happens in anaerobic environments where sulfate-reducing microorganisms (SRM) use sulfate as a terminal electron acceptor in the process of organic matter degradation. The canonical DSR process is thought to reduce sulfate to hydrogen sulfide (H_2S) as the end product. However, zero valent sulfur (S^0) has been widely detected in anaerobic conditions along with H_2S (2). Recent studies indicated that dissimilatory sulfite reductases (dSiRs), specifically, DsrAB and DsrC generated S^0 as an intermediate product (3-5).

Different from DSR, ASR is active in various environments where microorganisms use sulfate as a sulfur donor to synthesize organic compounds including cysteine, methionine, and glutathione. Sulfate is activated by an ATP sulfurylase (ATPS) to generate adenosine 5'-phosphosulfate (APS), which is directly reduced by an APS reductase (APSR) to generate sulfite. Alternatively, APS can be further phosphorylated to 3'-phosphoadenosine 5'-phosphosulfate (PAPS) by the APS kinase (APSK), which is then reduced to sulfite by a PAPS reductase (PAPSR) (6). Assimilatory sulfite reductases (aSiRs, some literatures used the name aSir, Asr, SiR and other names) finally reduced sulfite to H_2S, which then reacts with O-acetyl-serine (OAS) to produce cysteine (6-9). Hence, H_2S has been regarded as the key knot connecting sulfate-reduction and cysteine synthesis for over 60 years (10).

Most of early studies that focused on ASR ignored the presence and involvement of S^0, mainly due to lack of related knowledge of S^0 and its

detection methods (11-15). It is possible that S^0 also is present in ASR process

because, first, recent studies indicated that S^0 is widely present in all types of cells. It

is not stable and can easily react with H_2S, small cellular thiols, such as glutathione

(GSH), and sulfite to form hydrogen polysulfides (HS_nH, $n \geq 2$), glutathione

polysulfides (GS_nH, $n \geq 2$), and thiosulfate ($S_2O_3^{2-}$) (16-20). Second, phylogenetic

sequence analysis indicates that aSiRs diverged from the same ancestral gene as dSiRs

(21-23). Therefore, aSiRs may share similar catalytic mechanisms as dSiRs and also

product S^0, but no related study has been reported as so far.

In recent years, a number of H_2S producing enzymes have been redefined as S^0

producing enzymes thanks to the development of S^0 analysis methods. For instance,

3-mercaptopyruvate sulfurtransferase (3-MST), which was reported to cleave the C–

S bond of 3-mercaptopyruvate to produce H_2S. In 2015, Kimura *et al*., discovered

that the released sulfur atom actually existed as HS_nH (24). Cystathionine β-

synthetase (CBS) and cystathionine γ-lyase (CSE) both produced S^0 from cystine (25,

26). Methanethiol oxidase (MTO) catalyzed methanethiol degradation to produce S^0

other than previously reported H_2S (27). From sulfur cycle viewpoint, these findings

indicate that when organic sulfur is converted to inorganic sulfur by biological

systems, S^0 is the direct product.

Previous studies indicated that H_2S is the main substrate of cysteine synthases (28),

but a few studies found that some microbial cysteine synthases also can use

thiosulfate as an alternative substrate to produce cysteine, which is less efficient

probably due to thiosulfate is weaker than H_2S in aspect of nucleophilicity and

thiosulfate forms a disulfide bond intermediate with OAS before generates cysteine

(29, 30). In our previous study, we discovered that the *Schizosaccharomyces pombe*

cysteine synthase Cys11 uses glutathione persulfide (GSSH) as a substrate to

produce cysteine (31). Moreover, for Cys11, the K_m value of GSSH is ~400 μM, 2-

fold lower than that of H_2S (~800 μM). Knocking out the sulfide:quinone

oxidoreductase (Sqr), which converts H_2S to S^0 in mitochondria, leads to cysteine-

deficient phenotypes. Therefore, we proposed that S^0 might be the *defacto* substrate

of Cys11 inside of *S. pombe* cells. Whether Cys11 is an exception or it represents a

common characteristic of microbial cysteine synthases needs further investigation.

In this study, we performed phylogenetic analysis of sulfite reductases and

cysteine synthases. Representative enzymes were purified and their activities were

examined. We found that all examined assimilatory sulfite reductases (aSiRs)

produced more S^0 than H_2S from sulfite. H_2S was a derivative product of S^0.

Microorganisms overexpressing aSiRs accumulated more S^0 than H_2S from sulfite

reduction. All representative cysteine synthases can use GSSH as a substrate to

synthesize cysteine and most of them preferred GSSH to H_2S. These findings

suggest that as in the H_2S oxidation pathway, S^0 also is the intermediate in the ASR

pathway. S^0 is more like the key knot connecting sulfate reduction and cysteine

synthesis other than H_2S.

Results

Methylene blue method cannot distinguish between hydrogen sulfide and hydrogen

persulfide

Methylene blue method was commonly used in early studies of sulfite reductases.

The mechanism is that two molecules of *N,N*-dimethyl-p-phenylenediamine

(DMPD) and one H_2S molecule are oxidized by ferric ion (Fe^{3+}) to form a

methylene blue dye, which has a maximal absorbance at around 670 nm (A_{670}) (Fig.

1). The A_{670} value is linearly correlated with amount of H_2S in the presence of

excessive Fe^{3+}/DMDP (32, 33). We found that Fe^{3+}/DMPD not only reacted with

H_2S, but also with hydrogen persulfide (HSSH). Furthermore, the latter generated

higher A_{670} values than the former at the same doses (Fig. 1A). LC-ESI-MS

analysis indicated that when H_2S reacted with Fe^{3+}/DMDP, not only methylene blue

compound (**a**) was produced, another compound whose structure is similar as **a** but

contains an extra sulfur atom (**b**) was also produced (Fig. S1). We proposed that **a**

and **b** were generated from reactions 2 and 3, respectively (Fig. 1B). When HSSH

reacted with Fe^{3+}/DMDP, **a** and **b** also were produced. The detailed mechanism

was not investigated, but could be because HSSH is inherently unstable and easily

breaks into H_2S and S^0 in aqueous solution, or HSSH could react with Fe^{3+}/DMDP

directly (reactions 4 and 5).

Elemental sulfur (S_8) did not generate any methylene blue dye, suggesting that

only nucleophilic polysulfide (polysulfide with terminal hydrogen atoms) can react

with Fe^{3+}/DMDP to produce S-containing heterocyclic dyes (Fig. 1B).

Representative aSiRs produced more S^0 than H_2S from sulfite in vitro

A total of 457,826 protein sequences are denoted as sulfite reductase in

GenBank (update to August 29, 2024). These sequences were downloaded and

subjected to redundancy analysis. After removing redundant sequences, 4,378

sulfite reductases were left. Their characteristic sequences were identified (PF01077

and PF03460) and using the presence or absence of the identified characteristic

sequence as a screening criterion, 289 sulfite reductases were finally extracted out.

Origins of these enzymes encompass bacteria, archaea, fungi, and plants.

Phylogenetic analysis of these enzymes indicated that 289 enzymes were

categorized into five major clusters with clusters 3 and 4 were two largest ones (Fig.

2 and Table S1).

We chose four aSiRs as representatives for further examination: *E. coli* CysIJ

(AAC75805.1), *S. cerevisiae* MET5&MET10 (NP_ 116579.1), and *S. pombe* Sir1

(CAB11176.1). They all belong to cluster 3; *Ruegeria pomeroyi* DSS-3 SPO2634

(AAV95879.1) belongs to cluster 4. Among them, CysIJ contains two subunits (Cys

I and Cys J) and has been intensively studied as a model aSiR in 1960s (10).

Others are from widely used model microorganisms. *S. cerevisiae* MET5&MET10

also contains two subunits, while *S. pombe* Sir1 and *R. pomeroyi* DSS-3 SPO2634

are single subunit enzymes. These subunits/enzymes were fused with an N-

terminal His-Tag, expressed in *E. coli* BL21(DE3), and purified using a nickel

column (Fig. S2). During cell disruption and protein purification processes, no

dithiothreitol (DTT) or other reductants was added. Freshly purified enzymes were

immediately subjected to activity analysis.

We mixed purified CysI and CysJ to obtain the CysIJ complex. Likewise, the

MET5&MET10 complex was obtained. CysIJ, MET5&MET10, and Sir1 were

NADPH dependent enzymes, we mixed them with NADPH, hemin, and sulfite. Since

SPO2634 was a ferredoxin dependent enzyme, we mixed it with ferredoxin (Fd),

ferredoxin-NADP+ oxidoreductase (FNR), NADPH, and sulfite. Fd and FNR

encoding genes were amplified from genomic DNA of *R. pomeroyi* DSS-3. We

expressed them in *E. coli* BL21(DE3) and purified these two proteins for this

experiment. To analyze the produced sulfur compound, we used a LC-MS dependent

method instead of the methylene blue method. The enzymatic products were

derivatized with monobromobimane (mBBr) and analyzed using LC-ESI-MS. For the

reaction catalyzed by CysIJ, H_2S derivatives mB-SH and mB-S-mB were detected.

Signal intensity of the former was 1.9×10^6 and the latter was 4.6×10^5 (Fig. 3A, Fig.

S3A and S3B). Surprisingly, derivative of HSSH (mB-SSH) also was detected and its

signal intensity was 3.6×10^6 (Fig. 3A, Fig. S3C). Previous studies indicated that when

derivatizing H_2S and HSSH with mBBr, dominant products were mB-S-mB and mB-

SS-mB, respectively (34). Here, we detected two derivatives of H_2S, mB-S-mB and

mB-SH (partial derivatization product), and one derivative of HSSH, mB-SSH

(partial derivatization product). The most probable reason is aSiRs and sulfite also

react with mBBr, which disturbs the reaction between H_2S/HSSH and mBBr, and

leads to partial derivatization. Considering mB-SH and mB-S-mB have similar

fluorescence and structure, we summed their signal intensities (roughly representing

H_2S amount) and compared it with signal intensity of mB-SSH (roughly representing

HSSH amount). The latter was higher than that of the former. Similarly, signal

intensity of H_2S derivatives (mB-SH + mB-S-mB) from MET5&MET10 was 2.4×10^6

; whereas signal intensity of HSSH derivative (mB-SSH) was 3.2×10^6 (Fig. 3B).

Signal intensities of H_2S derivatives from SPO2634 and Sir1 also were lower than that

of HSSH derivative (Fig. 3C and 3D).

A previous report indicated that hemin itself can catalyze sulfite reduction (35). The

authors used methyl viologen (MeV) as electron source. By monitoring the decrease

in absorbance at 600 nm from the MeV radical cation, they calculated the reaction

kinetics and concluded that free siroheme was a more efficient catalyst than dSiRs.

No product analysis was performed in that study but the authors proposed that

trithiolate was a product. Here, for CysIJ, MET5&MET10, and Sir1 catalyzing

reactions, hemin was added as a cofactor. To find out whether hemin itself can reduce

sulfite to H_2S or HSSH, we mixed hemin with sulfite. The reaction conditions were

the same except that no sulfite enzyme was added. LC-ESI-MS analysis demonstrated

that no H_2S or HSSH was detected.

To further confirm the S^0 products of these enzymes, we analyzed the products

with the trifluoromethanesulfonate derivatization-based method, which can

discriminate polysulfides with different length of chains (H_2S_n, n=2-8) (36). The

reaction mixtures were reacted with methyl trifluoromethanesulfonate to convert

hydrogen polysulfide (H_2S_n) to dimethylpolysulfide. Results showed that HSSH

(H_2S_2) was the main product of these enzymes while HSSSSH (H_2S_4) was the minor

one (Fig. 3E). We then quantified the produced total sulfane sulfur (representing

amount of S^0 atoms) with a cyanide method and compared it to the amount of

produced H_2S. Results showed that CysIJ, MET5&MET10, and SPO2634 produced

80~130 μM S^0, Sir was less efficient than them and produced about 50 μM S^0 from

150 μM sulfite. In comparison, CysIJ produced about 15 μM H_2S and the other three

aSiRs produced only 1~5 μM H_2S (Fig. 3F and 3G). These results indicated that all of

the four aSiRs produced more S^0 than H_2S.

For the presence of HS_nH in the products, a possible explanation is that

aSiRs firstly produce H_2S from sulfite, and HS_nH are generated from autonomous

oxidation of H_2S (by dissolved oxygen). However, according to our experience, the

process of H_2S autonomous oxidation is slow, which usually takes hours without

the help of related enzymes or strong oxidants (such as Fe^{3+} and H_2O_2). For

confirmation, we mixed H_2S with reaction buffer containing no aSiR. After 10 min

incubation at 25°C, no HS_nH was produced, suggesting that HS_nH was not

produced from autonomous oxidation of H_2S. As the control, we also mixed HSSH

with reaction buffer containing no aSiR and other reagents. Other conditions were the

same as that of enzymatic reaction. H_2S production was detected and its amount was

about 1/10 of HSSH. Considering that HSSH is easily reduced to H_2S when reductant

is present, we also mixed HSSH, reaction buffer, and NADPH. In this case, H_2S

production was significantly increased to about 1/2 of HSSH (Fig. S4).

Together, the above results suggested that under the tested conditions, S^0 was the

direct product of all four aSiRs. As showing in equation 1, aSiR gets four electrons

from NADPH and passes them to sulfite. Sulfite is reduced to S^0. The produced S^0

atoms either attach to the enzyme and exist as protein-S_nH, or exist as free S_n (equation

2). In either case, S_n can get more electrons and become water soluble HS_nH (equation

3). When there is abundant NADPH present, HS_nH is easily reduced to H_2S (equation

4). Since the H_2S generation reaction is aSiR independent, H_2S is more like a

derivative product of ASR process, not the direct product.

$SO_3^{2-} + 3\ NADPH + 3\ H^+ \longrightarrow S^0 + 3H_2O + 2\ NADP^+$ (equation 1)

$n\ S^0 \longrightarrow S_n$ (equation 2

$S_n + NADPH + H^+ \longrightarrow HS_nH + 3H_2O$ (equation 3)

$HS_nH + NADPH + H^+ \longrightarrow HS_{n-1}H + H_2S$ (equation 4)

Representative aSiRs produced more S^0 than H_2S in vivo.

Since *E. coli* MG1655 contains no H_2S oxidation enzyme, testing the products of

aSiRs in it is convenient. We expressed the four aSiRs (using the pBBR1MCS5

plasmid as a vector) in *E. coli* MG1655. *E. coli* MG1655 with the pBBR1MCS5

empty plasmid was used as a control. The cell suspension (10 ml, $OD_{600}=3$) was

mixed with 0.5 mM sulfite in PBS buffer (pH 7.4, 50 mM) and the mixture was

incubated at 37□ with shaking. After 1 h incubation, consumed sulfite and

generated products were quantified. We observed that compared with the control, all

aSiR expression strains consumed more sulfite. More importantly, they all produced

more total S^0 than H_2S from the consumed sulfite (Fig. 3H).

We also constructed a *S. cerevisiae* S288C strain (that contains no *sqr*) with

MET5&MET10 overexpression and a *R. pomeroyi* DSS-3 Δ*sqr*Δ*fccAB* strain

with SPO2634 overexpression (Since *sqr* and *fccAB* encode H_2S oxidation enzymes,

deleting these two genes prevent the oxidation of H_2S to S^0). Cell suspensions of

these strains were mixed with sulfite, and the products were analyzed. Similarly,

they also produced more S^0 than H_2S (Fig. 3I and 3J). These results demonstrated that

the tested aSiRs tended to produce S^0 other than H_2S *in vivo*.

Representative cysteine synthases utilized S^0 to produce cysteine

A total of 499,263 protein sequences are denoted as cysteine synthase in

GenBank (update to August 29, 2024). The similar screening process was

performed and 735 cysteine synthases were finally extracted out. Origins of these

enzymes encompass bacteria, archaea, fungi, and plants. Phylogenetic analysis of

these enzymes indicated that they were categorized into six major clusters with

clusters 4 and 5 were two largest ones (Fig. 4, Table S2).

We selected 5 microbial cysteine synthases for further study: CysK and CysM from

E. coli MG1655 (EcCysK and EcCysM), CysK from *R. pomeroyi* DSS-3

(RpCysK), SiRe_ 1641 from *Sulfolobus islandicus* REY15A, a model archaeon

that belongs to Crenarchaeota. They all belong to cluster 4. CysM from *R. pomeroyi*

DSS-3 (RpCysM) belongs to cluster 5. We expressed their encoding genes in *E. coli*

BL21(DE3). These enzymes (fused with a His-tag at N-terminus) were purified

using a nickel column (Fig. S5). Freshly purified enzymes were subjected to

activity analysis immediately.

A catalytic mechanism of cysteine synthase has been reported. OAS first reacted

with PLP to form the α-aminoacrylate intermediate containing a C=C bond,

which was electrophilic. Subsequently, nucleophilic H_2S attacked the C=C bond of α-

aminoacrylate to generate cysteine (Fig. 5) (37). We mixed purified cysteine

synthases with H_2S/GSSH/$Na_2S_2O_3$, OAS, and pyridoxal 5-phosphate (PLP) in PBS

buffer (50 mM, pH 7.4) and incubated the mixture at 30°C for 5 min. LC-ESI-

MS method was used to analyze the products.

From the catalyzing mechanism perspective, GSSH and $Na_2S_2O_3$ react with OAS

will generate GS-cysteine and sulfo-cysteine intermediates, respectively (equations 5-

6). We detected these two intermediates from reactions catalyzed by all five

enzymes (Fig. S6-S11). After treating the intermediates with tris-(2-carboxyethyl)-

phosphine (TCEP), L-cysteine was detected (equations 7-8). These enzymes also used

H_2S as substrate to produce L-cysteine (equation 9).

GSSH + OAS ——> GS-cysteine + Acetate   (equation 5)

GS-cysteine + 2 e ——> GSH + Cysteine   (equation 6)

$H_2S_2O_3$ + OAS ——> HSO_3-cysteine + Acetate   (equation 7)

HSO_3-cysteine + 2 e ——> H_2SO_3 + Cysteine   (equation 8)

H_2S + OAS ——> Cysteine + Acetate   (equation 9)

We then assayed the kinetics of all five cysteine synthases (Fig. 6, Fig. S12-

S14). The obtained data were compared with the reported data of *S. pombe* cysteine

synthase Cys11 (31). For all 6 enzymes, the K_m values for GSSH were between

202~426 μM, K_m values for H_2S were between 157~3192 μM, and K_m values for

$Na_2S_2O_3$ were between 408~3774 μM. The V_{max} values for GSSH were between

0.125~0.280 mM·min^{-1}, V_{max} values for H_2S were between 0.139~0.581 mM·min^{-1},

and V_{max} values for $Na_2S_2O_3$ were between 0.021~0.173 mM·min^{-1}. The K_{cat} values

for GSSH were between 2.083~9.778 s^{-1}, K_{cat} values for H$_2$S were between

3.127~13.833 s^{-1}, and K_{cat} values for Na$_2$S$_2$O$_3$ were between 0.495~2.454 s^{-1}. These

results indicated that among GSSH, H$_2$S, and Na$_2$S$_2$O$_3$, Na$_2$S$_2$O$_3$ was the most

unpreferred substrate. Most cysteine synthases showed higher affinity to GSSH

than to H$_2$S.

Discussion

Inorganic sulfur (sulfate) becomes organic sulfur via the assimilatory sulfate

reduction and cysteine synthesis pathways. The concept that H$_2$S is a key

intermediate of this process is founded in 1960s when S^0 analysis methods were

short and knowledge of biological S^0 were scarce (10). In this study, we re-

examined the product of aSiRs with modern methods that can distinguish S^0

containing compounds from H$_2$S. We found that all tested aSiRs produced both S^0

and H$_2$S from sulfite. S^0 was more like the direct product of aSiRs while H$_2$S

was its derivative. We also re-examined model cysteine synthases and discovered

that all tested cysteine synthases can use S^0 (S$_2$O$_3$$^{2-}$ and GSSH) as alternative

substrates to produce cysteine.

From chemistry viewpoint, biological S^0 compounds are better intermediates than

H$_2$S. First, the pK_a1 of H$_2$S is 6.8 and gasification temperature is 34.3°C, so it easily

releases out of cells under normal physiological condition. In comparison, the pK_a of

GSSH is 5~6, and therefore, it mainly exist in deprotonated form inside cells

(38). Thiosulfate is even more stable than GSSH. Second, H$_2$S is toxic to the

respiration system, its intracellular concentration is limited to 10~30 nM level (33),

which was far below the K_m value of cysteine synthases. In contrast, intracellular

S^0 chemicals are high to 100~400 μM (39, 40), which was near the K_m values of

cysteine synthases. Third, biological S^0 compounds have distinctive activities. For

instance, GSSH is more nucleophilic than GSH because of the α-effect and hence

displays high reactivity toward electrophiles such as reactive oxygen species

(ROS) and toxic metal cations (41). In consistent, we found that cysteine synthases

preferred GSSH the most. It is noteworthy that GSSH leads to the production of GS-

cysteine intermediate which contains a disulfide bond. We used TCEP to cut this

disulfide bond to release cysteine *in vitro*. It is still unknown how this disulfide bond is

cleaved in the cell, small reducing molecules like GSH might be responsible for this

reaction. In addition, we also found that microorganisms accumulated high

concentration of intracellular S^0 during the fast growth phase (cysteine synthesis

process is active in fast growth phase), but reduced its level quickly after that (40).

Combining the above evidence, we proposed that S^0 is the *de facto* substrate of

cysteine synthases *in vivo*. S^0 probably is a key knot connecting sulfate reduction and

cysteine synthesis in microorganisms. H_2S is more like an efflux of intracellular S^0

(Fig. 7).

Recent studies indicated that S^0 is an intermediate product in DSR process (4). A

proposed mechanism is, first, DsrAB complex reduces sulfite to S^0, which then is

transferred to DsrC in the form of DsrC-SSH or DsrC trisulfide (3, 5). Second, DsrC-

SSH or DsrC trisulfide receives electrons from the Dsr membrane complex and

reduce the bound S^0 to H_2S. Therefore, DsrC is an important S^0 carrier in this process.

Some bacteria contain only DsrABC and no Dsr membrane complex. These bacteria

may only produce S^0 during sulfate reduction and DsrC may transfer its S^0 to other

acceptors in the cell. Using the *Archaeoglobus fulgidus* DsrC as seed and 35%

homology as criteria, we searched for the DsrC homologs in genomes of *E. coli*, *S.*

cerevisiae, *S. pombe*, and *R. pomeroyi*, no DsrC homologue was found. Probably

because DsrC mainly exists in anaerobic bacteria but the four strains are all

aerobic/facultative bacteria. No evidence indicates that DsrC is involved in ASR

process as so far. In conclusion, we found that like in DSR process, S^0 presents in

ASR process as a direct product of aSiRs, probably because both aSiRs and dSiRs

diverge from a common ancestor.

Materials and methods

Strains and cultivation conditions

R. pomeroyi DSS-3 is a gift from Prof. Yuzhong Zhang of Ocean University of

China. *R. pomeroyi* DSS-3 derivative *ΔsqrΔfccAB* were constructed in a previous

work(29). *S. islandicus* REY15A is a gift from Prof. Yulong Shen of Shandong

University. *S. pombe* YHL6381 is a gift from Prof. Huang Ying. *E. coli* strains

used for plasmid construction and protein expression, and plasmids constructed in

this study are all listed in Table S4. *E. coli* strains were cultured in LB medium.

R. pomeroyi DSS-3 strains were cultured in 1/2 YTSS medium. *S. cerevisiae* S288C

strains were cultured in YPD or SD medium. *S. pombe* YHL6381 strains were

cultured in YES or SD medium. *S. islandicus* REY15A were grown in MTSV

medium [mineral salt supplemented with 0.2% w/v (weight/volume) tryptone (T),

0.2% (w/v) sucrose (S), mixed solution of vitamins (V)] or MTSVU medium [MTSV

supplemented with 0.001% (w/v) uracil (U)]. For cultivation, *E. coli* strains were

cultured at 37□ with shaking (220 rpm). *R. pomeroyi* DSS-3, *S. cerevisiae* S288C

and *S. pombe* YHL6381 strains were cultivated at 30□ with shaking (220 rpm). *S.*

islandicus REY15A strains were cultivated at 75□ with shaking (220 rpm).

Reagents

Sodium hydrosulfide (NaHS), sodium sulfide (Na_2S), cysteine, reduced glutathione

(GSH), sulfite and thiosulfate were purchased from Sigma-Aldrich (Saint Louis,

MO). S_8 were purchased from TCI Company (Shanghai, China). HSSH was

prepared following the protocol of Xin *et al* (34). GSSH was prepared following

the protocol of Luebke *et al* (42). H_2S_n was prepared by a modifying protocol of Xin

et al. Briefly, 13 mg sulfur powder and 28 mg sodium sulfide were added to 5 ml of

anoxic distilled water under argon gas, and the bottle was sealed. The bottle was

incubated at 37^oC for 24 hours. The pH was adjusted to 9.3 with 6 M HCl, and the

solution was diluted to 10 ml with anoxic distilled water. The stock polysulfide

concentration was determined with cyanide method and calibrated by using

thiosulfate as the standard.

Trifluoromethanesulfonate derivatization and following HPLC analysis was

performed following the protocol of a previous report (36).

Methylene blue method

The methylene blue method was performed as reported previously (32). Two

reagents were required, 0.02 M *N,N*-dimethyl-p-phenylenediamine sulfate in 7.2 N

hydrochloric acid (DPD reagent), and 0.03 M ferric chloride in 1.2 N hydrochloric

acid (ferric chloride agent). Assays are performed in 1.5-mL microfuge tubes.

First, 250 μl deionized water was added into a 1.5-ml tube, followed by the addition

of 250 μl NaHS, HSSH, or S_8 solution (in 50 μM, 100 μM, 200 μM, or 500 μM

concentration). Second, 20 μl DPD reagent was added, rapidly followed by

addition of 20 μl of ferric chloride reagent. The mixture was shaken vigorously and

placed in the dark for 30 minutes. After being centrifuged at $17,000 \times g$ for 2

minutes. Supernatant (200 µl) was pipetted out and the absorbance at 670 nm was

measured. The compounds in supernatant were analyzed using LC-ESI-MS.

Phylogenetic analysis

The protein sequences of sulfite reductase and cysteine synthase were

downloaded from NCBI database (update to 29 August 2024). Redundant sequences

were removed using the CD-hit tool with conventional criteria (sequence identity

threshold ≥45%, word length =2). The characteristic sequences of the conserved

regions were analyzed in the InterPro database. The characteristic sequences of

sulfite reductases were PF01077 and PF03460, and the characteristic sequences of

cysteine synthases was PF00291. The hmmsearch command in Hmmer tool was

used to identify all the featured sequences. Phylogenetic trees were constructed by a

neighbor-joining method using ETE3 with a partial deletion, p-distance

distribution, and bootstrap at 1,000 repeats. Multiple sequence alignment was

performed using MUSCLE.

Protein expression and purification

Cysteine synthases and sulfite reductases encoding genes were amplified from

genomic DNA of *E. coli*, *R. pomeroyi* DSS-3, *S. cerevisiae* S288C, *S. pombe*

YHL6381 and *S. islandicus* REY15A genomic DNA. Ferredoxin (Fd, SPO2377),

and ferredoxin-NADP reductase (FNR, SPO2637) encoding genes were amplified

from genomic DNA of *R. pomeroyi* DSS-3. The expression plasmids (see Table S4)

were constructed using primers listed in Table S5. These plasmids were

introduced into *E. coli* BL21(DE3) and the strains were incubated in LB

medium containing kanamycin (50 µg/ml). When OD_{600} reached 0.6, 0.4 mM

isopropyl β-D-1-thiogalactopyranoside (IPTG) was added and the temperature was

decreased to 25°C. The cultivation was further continued overnight. Cells were

harvested by centrifugation and then re-suspended in binding buffer (200 mM NaCl,

50 mM NaH$_2$PO$_4$, 20 mM Imidazol, pH 8.0). For protein purification, cell

disruption was performed using a Pressure Cell Homogeniser (SPCH-18) at 4°C.

Cell lysate was centrifuged to remove the debris and cysteine synthetase/sulfite

reductase was purified using the nickel-nitrilotriacetic acid (Ni-NTA) agarose resin

(Invitrogen, Waltham, MA, USA). Purification was conducted by following the

manufacturer's instructions. The eluted protein was loaded onto the PD-10 desalting

column (GE) for buffer exchange to sodium phosphate buffer (20 mM, pH 7.6).

Purity of the proteins was examined via SDS-PAGE.

Sulfite reductase activity assay in vitro

The assays were performed as previously reported (12). Briefly, for CysIJ,

Met5&Met10, and Sir1 activity assays, 0.3 ml reaction buffer (50 mM K$_3$PO$_4$, pH

7.7, containing 0.1 mM Na$_2$EDTA) contained 2 μg/mL purified sulfite reductase, 200

μM sulfite, 100 μM hemin and 2 mM NADPH. The reaction was conducted at

25°C for 10 min. For SPO2634 activity assay, 0.3 ml Tris-HCl buffer (pH 7.5,

containing 0.1 M NaCl) contained 2 μg/ml of purified SPO2634, purified ferredoxin

(Fd, SPO2377), and purified ferredoxin-NADP reductase (FNR, SPO2637), 200

μM sulfite and 2 mM NADPH. The reaction was conducted at 25°C for 10 min.

Produced H$_2$S and HSSH were derivatized by mBBr following a previously reported

protocol (34), and derivatized products were analyzed by LC-ESI-MS.

As control experiments, 500 μM HSSH was added into 0.3 ml reaction buffer (50

mM K$_3$PO$_4$, pH 7.7, containing 0.1 mM Na$_2$EDTA) with or without 2 mM NADPH.

After 10 min incubation at 25°C, produced H_2S and remaining HSSH were derivatized

by mBBr and quantified using RP-HPLC.

LC-ESI-MS Analysis

The quadrupole time-of-flight, high-resolution mass spectrometer (Ultimate 3000,

Burker impact HD, Thermo Fisher, Waltham, MA, USA) was used. The column

was InertSustain C18 5 μm (4.6 mm × 250 mm, GL Sciences, Shanghai, China).

Injection volume of sample was 10 μl. The source temperature was set at 200°C

and the ion spray voltage was at 4.5 kv. Nitrogen was used as the nebulizer and

drying gas. The mobile phase was a mixture of pure water and methanol. A linear

gradient of solvent A (0.25% acetic acid in ddH_2O) and solvent B (100% methanol)

from 7.5% to 100% to 7.5% (solvent B) in 31 min was used for elution, the

column temperature was maintained at 38°C. Data were acquired in full MS

(100–1500 m/z) mode. Full mass resolution (40000@ m/z 1222) at full sensitivity.

Sulfite reductase activity assay in vivo

E. coli MG1655-CysJI/Met5&Met10/Sir1 overexpression strains were constructed

with pBBR1MCS5 plasmids (Table S4). The strains were cultured in LB medium

at 37°C for overnight. The overnight culture (1 ml) was transferred into 100 ml fresh

LB medium and cultured to OD_{600}=0.4, and then 0.4 mM IPTG was added and the

cultivation continued at 37°C for 3.5 h. Cells were collected by centrifugation

(4,500×g, 10 min) and re-suspended in PBS buffer (pH 7.4, 50 mM) to make a

OD_{600}=3.0 cell suspension. Sulfite (0.5 mM) was added into 10 ml cell suspension

in a 50-ml scale tube. The tube was then incubated at 37□ for 1 h with shaking

(200 rpm).

R. pomeroyi-SPO2634 overexpression strains were constructed with *R.*

pomeroyi DSS-3 Δ*sqr*Δ*fccAB* strain and pBBR1-SPO2634 plasmid (Table S4).

Strains were cultured in 1/2 YTSS medium at 30□ for overnight. The overnight

culture (1 ml) was transferred into 100 ml of fresh medium and cultured to OD_{600}=

0.7, and then 0.1 mM sulfite was added and the cultivation continued until OD_{600}

reached 2.0. Cells were collected by centrifugation (4,500×g, 10 min) and re-

suspended in Tris-HCl buffer (pH 8.4, 50 mM, 20 mM $MgCl_2$, 20g/L NaCl) to

make a OD_{600}=5.0 cell suspension. Sulfite (0.5 mM) was added into 10 ml cell

suspension in a 50-ml scale tube. The tube was then incubated at 30□ for 1 hwith

shaking (200 rpm).

S. cerevisiae S288C-MET5&MET10 overexpression strain was constructed with

pJFE3 vector (Table S1). The strain was cultured in SD-URA medium at 30□

for overnight. The overnight culture (1 ml) was transferred into 100 ml of fresh

medium and cultured to OD_{600}= 0.6, and then 0.1 mM sulfite was added and the

cultivation continued until OD_{600} reached 1.0. Cells were collected by

centrifugation (4,500×g, 10 min) and re-suspended in PBS buffer (pH 7.4, 50 mM)

to make a OD_{600}=10.0 cell suspension. Sulfite (0.5 mM) was added into 10 ml cell

suspension in a 50-ml scale tube. The tube was then incubated at 30□ for 1 h with

shaking (200 rpm).

After the incubation, cell suspension was centrifugated (4,500×g, 10 min) and 100

μl supernatant was taken after centrifugation. H_2S, total S^0, and sulfite

concentrations in the supernatant were quantified with following methods.

H_2S and sulfite quantification

For quantification of H_2S and remaining sulfite in obtained supernatant and in aSiRs

reaction system, the supernatant and reaction mixture were passed through a 3K

Amicon® Ultra centrifugation filter (Sigma-Aldrich, China). The passed-through

solution was collected and subjected to mBBr derivatization following protocols

reported previously (43-44). Briefly, 30 µl of passed-through solution was added to a

PCR tube containing 70 µl of 100mM Tris-HCl buffer (pH 9.5, 0.1mM DTPA),

followed by addition of 50 µl of 10 mM mBBr (dissolved in deoxygenated

acetonitrile). The reaction was stopped by adding 50 µl of 200 mM 5-sulfosalicylic

acid after a 30-minute incubation. All derivatized samples are removed from the

hypoxic chamber and stored at 4°C until analyzed by RP-HPLC. Pre-made standard

curves of H_2S and sulfite were used for calculation.

Total S^0 quantification

Total S^0 quantification was performed with the cyanide method (45). First, 0.55 mL 1%

boric acid was added into a 1.5-mL tube, heated with boiling water for 1 min and then

cooled to room temperature, followed by the addition of 0.25 mL sample (the same

volume of deionized water was added to the blank control group). Second, 0.2 mL of

0.1 M potassium cyanide was added, heated with boiling water for 1 min, and then

cooled to room temperature. Finally, 0.1 mL $Fe(NO_3)_3$ solution was added and

centrifuged at 13000 rpm for 2 min. The supernatant was taken and the absorbance at

460 nm was measured. According to the pre-made standard curve, the detection result

is converted to the total S^0 concentration.

RP-HPLC Analysis

A C18 reverse phase HPLC column (VP-ODS, 150 3 4 mm, Shimadzu) was pre-

equilibrated with 80% Solvent A (10% methanol and 0.25% acetic acid) and 20%

Solvent B (90% methanol and 0.25% acetic acid). The column was eluted with the

following gradients of Solvent B: 20% from 0 to 10 min; 20%–40% from 10 to 25

min; 40%–90% from 25 to 30 min; 90%–100% from 30 to 32 min; 100% from 32 to

35 min; 100 to 20% from 35 to 37 min; and 20% from 37 to 40 min. The flow rate

was 0.75 ml min21. And the bimane adducts were detected by a fluorescence detector

(RF-10AXL, Shimadzu) with excitation at 340 nm and emission at 450 nm.

Cysteine synthase activity assay

Cysteine synthases were expressed in *E. coli* BL21(DE3) (Table S4) and purified with

Ni-NTA agarose resin. Activity assay was carried out as described previously.

Briefly, 0.3 ml reaction buffer (50 mM PBS buffer, pH 7.4) contained 1 μg/ml

purified enzyme, 100 μM pyridoxal phosphate, 1.0 mM–2.0 mM O-acetyl-L-serine

(OAS), 1.0–5.0 mM NaSH, GSSH, or $Na_2S_2O_3$. The reaction was conducted at 30°C

for 5 min and the products were derivatized with mBBr and analyzed with LC-ESI-

MS. For kinetics assay, the same reaction system contained varied concentrations of

NaSH, GSSH or $Na_2S_2O_3$ (see maintain text). The reaction was conducted at 30°C

for 5 min. For the reaction using NaSH as a substrate, the produced cysteine was

labeled with mBBr and quantified by HPLC. For the reaction using GSSH or

thiosulfate as a substrate, after 5 min reaction, the product was mixed with 0.5 mM

TCEP, and the released cysteine was labeled with mBBr and quantified by HPLC.

All kinetic parameters were determined using the Michaelis-Menten equation (non-

linear regression) embedded in Graphpad prism software.

Acknowledgements

This work was supported by the National Key R&D Program of China (2022YFC3401301) and the National Natural Science Foundation of China (92351302).

Declaration of competing interest

The authors declare no conflict of interest regarding the publication of this article.

References

1. Jorgensen BB, Findlay AJ, Pellerin A. 2019. The biogeochemical sulfur cycle of marine sediments. Front Microbiol 10:849.

2. Wu B, Liu F, Fang W, Yang T, Chen GH, He Z, Wang S. 2021. Microbial sulfur metabolism and environmental implications. Sci Total Environ 778:146085.

3. Oliveira TF, Vonrhein C, Matias PM, Venceslau SS, Pereira IAC, Archer M. 2008. The crystal structure of *Desulfovibrio vulgaris* dissimilatory sulfite reductase bound to DsrC provides novel insights into the mechanism of sulfate respiration. J Biol Chem 283: 34141-34149.

4. Wang S, Lu Q, Liang Z, Yu X, Lin M, Mai B, Qiu R, Shu W, He Z, Wall JD. 2023. Generation of zero-valent sulfur from dissimilatory sulfate reduction in sulfate-reducing microorganisms. Proc Natl Acad Sci USA 120(20):e2074242176.

5. Santos AA, Venceslau SS, Grein F, Leavitt WD, Dahl C, Johnston DT, Pereira IAC. 2015. A protein trisulfide couples dissimilatory sulfate reduction to energy conservation. Science 350(6267):1541-1545.

6. Jespersen M, Wagner T. 2023. Assimilatory sulfate reduction in the marine methanogen *Methanothermococcus thermolithotrophicus*. Nat Microbiol 8:1227-1239.

7. Takahashi H, Kopriva S, Giordano M, Saito K, Hell R. 2011. Sulfur
assimilation in photosynthetic organisms: molecular functions and regulations
of transporters and assimilatory enzymes. Annu Rev Plant Biol 62:157-184.

8. Demin KA, Prazdnova EV, Minkina TM, Gorovtsov AV, Reguera G. 2024.
Sulfate-reducing bacteria unearthed: ecological functions of the diverse
prokaryotic group in terrestrial environments. Appl Environ Microbiol
90:e139023.

9. Ghazi Esfahani B, Walia N, Neselu K, Garg Y, Aragon M, Askenasy I, Wei HA,
Mendez JH, Stroupe ME. 2025. Structure of dimerized assimilatory NADPH-
dependent sulfite reductase reveals the minimal interface for diflavin reductase
binding. Nat Commun 16: 2955.

10. Pasternak CA, Ellis RJ, Jones-Mortimer MC, Crichton CE. 1965. The control of
sulphate reduction in bacteria. Biochem J 1:270-275.

11. Kline BC, Schoenhard DE. 1970. Biochemical characterization of sulfur
assimilation by *Salmonella pullorum*. J Bacteriol 1:142-150.

12. Ostrowski J, Barber MJ, Rueger DC, Miller BE, Siegel LM, Kredich NM.
1989. Characterization of the flavoprotein moieties of nadph-sulfite reductase
from *Salmonella typhimurium* and *Escherichia coli*: Physico-chemical and
catalytic properties, amino acid sequence deduced from dna sequence of *cysJ*,
and comparison with nadph-cytochrome P-450 reductase. J Biol Chem
264:15796-15808.

13. Lewis MS, Patricia SD, Henry K. 1974. Reduced nicotinamide adenine-
dinucleotide phosphate-sulfite reductase of enterobacteria .3. *Escherichia-coli*
hemoflavoprotein - catalytic parameters and sequence of electron flow. Journal of
Biological Chemistry Volume 249:1572-1586.

14. Covès J, Nivière V, Eschenbrenner M, Fontecave M. 1993. NADPH-sulfite

reductase from Escherichia coli. A flavin reductase participating in the

generation of the free radical of ribonucleotide reductase. J Biol Chem

268:18604-18609.

15. Siegel LM, Leinweber FJ, Monty KJ. 1965. Characterization of the Sulfite and

Hydroxylamine Reductases of *Neurospora crassa*. J Biol Chem 240:2705-2711.

16. Ogata S, Matsunaga T, Jung M, BarayeuU, Morita M, Akaike T. 2023. Persulfide

biosynthesis conserved evolutionarily in all organisms. Antioxid Redox Signal

39:983-999.

17. Lau N, Pluth MD. 2019. Reactive sulfur species (RSS): persulfides,

polysulfides, potential, and problems. Curr Opin Chem Biol 49:1-8.

18. Filipovic MR, Zivanovic J, Alvarez B, Banerjee R. 2018. Chemical biology

of H_2S signaling through persulfidation. Chem Rev 118:1253-1337.

19. Sawa T, Takata T, Matsunaga T, Ihara H, Motohashi H, Akaike T. 2021.

Chemical biology of reactive sulfur species: hydrolysis-driven equilibrium of

polysulfides as a determinant of physiological functions. Antioxid Redox

Signal 36:327-336.

20. Olson KR. 2019. Hydrogen sulfide, reactive sulfur species and coping with

reactive oxygen species. Free Radic Biol Med 140:74-83.

21. Molitor M, Dahl C, Molitor I, Schafer U, Speich N, Huber R, Deutzmann R,

Truper HG. 1998. A dissimilatory sirohaem-sulfite-reductase-type protein from the

hyperthermophilic archaeon Pyrobaculum islandicum. Microbiology (Reading)

144 (Pt 2): 529-541.

22. Dhillon A, Goswami S, Riley M, Teske A, Sogin M. 2005. Domain evolution and

functional diversification of sulfite reductases. Astrobiology 5: 18-29.

23. Crane BR, Siegel LM, Getzoff ED. 1995. Sulfite reductase structure at 1.6 A:

evolution and catalysis for reduction of inorganic anions. Science 270: 59-67.

24. Kimura Y, Toyofuku Y, Koike S, Shibuya N, NagaharaN, Lefer D,

OgasawaraY, Kimura H. 2015. Identification of H_2S_3 and H_2S produced by 3-

mercaptopyruvate sulfurtransferase in the brain. Sci Rep 5:14774.

25. Mishanina TV, Libiad M, Banerjee R. 2015. Biogenesis of reactive sulfur

species for signaling by hydrogen sulfide oxidation pathways. Nat Chem Biol

11:457-464.

26. Kimura H. 2021. Hydrogen Sulfide (H_2S) and Polysulfide (H_2S_n) Signaling:

The First 25 Years. Biomolecules 11:896.

27. Cao Q, Liu X, Wang Q, Liu Z, Xia Y, Xun L, Liu H. 2024.

Rhodobacteraceae methanethiol oxidases catalyze methanethiol degradation to

produce sulfane sulfur other than hydrogen sulfide. mBio 15:e02907-23.

28. Becker MA, Kredich NM, Tomkins GM. 1969. The purification and

characterization of O-acetylserine sulfhydrylase-A from *Salmonella*

typhimurium. J Biol Chem 244:2418-2427.

29. Nakamura T, Iwahashi H, Eguchi Y. 1984. Enzymatic proof for the identity of

the S-sulfocysteine synthase and cysteine synthase B of *Salmonella*

typhimurium. J Bacteriol 158:1122-1127.

30. Hensel G, Truper HG. 1983. O-Acetylserine sulfhydrylase and S-sulfocysteine

synthase activities of *Rhodospirillum tenue*. Arch Microbiol 134:227-232.

31. Zhang X, Xin Y, Chen Z, Xia Y, Xun L, Liu H. 2021. Sulfide-quinone

oxidoreductase is required for cysteine synthesis and indispensable to

mitochondrial health. Redox Biol 47:102169.

32. Siegel LM. 1965. A direct microdetermination for sulfide. Anal Biochem 11:126-
132.

33. Kubán V, Dasgupta PK, Marx JN. 1992. Nitroprusside and methylene blue
methods for silicone membrane differentiated flow injection determination of
sulfide in water and wastewater. Analytical chemistry (Washington) 64:36-43.

34. Xin Y, Liu H, Cui F, Liu H, Xun L. 2016. Recombinant *Escherichia coli* with
sulfide:quinone oxidoreductase and persulfide dioxygenase rapidly oxidises
sulfide to sulfite and thiosulfate via a new pathway. Environ Microbiol 18:5123-
5136.

35. Soriano A, Cowan JA. 1995. Sulfite reductase-active-site residues are
noncatalytic-comparison of reaction energetics for enzyme-catalyzed and
siroheme-catalyzed reduction of inorganic substrates. J Am Chem Soc 117:4724-
4725.

36. Kamyshny AJ, Goifman A, Gun J, Rizkov D, Lev O. 2004. Equilibrium
Distribution of Polysulfide Ions in Aqueous Solutions at 25°C: A New Approach
for the Study of Polysulfides' Equilibria. Environ Sci Technol 38:6633-6644.

37. Claus MT, Zocher GE, Maier TH, Schulz GE. 2005. Structure of the O-
acetylserine sulfhydrylase isoenzyme CysM from *Escherichia coli*. Biochemistry
44:8620-8626.

38. Benchoam D, Semelak JA, Cuevasanta E, Mastrogiovanni M, Grassano JS, Ferrer-
Sueta G, Zeida A, Trujillo M, Möller MN, Estrin DA, Alvarez B. 2020. Acidity and
nucleophilic reactivity of glutathione persulfide. Journal of Biological Chemistry
295:15466-15481.

39. Kimura H. 2015. Signaling molecules: hydrogen sulfide and polysulfide.
Antioxid Redox Signal 22:362-376.

40. Ran M, Wang T, Shao M, Chen Z, Liu H, Xia Y, Xun L. 2019. Sensitive

method for reliable quantification of sulfane sulfur in biological samples. Anal

Chem 91:11981-11986.

41. Bianco CL, Chavez TA, Sosa V, Saund SS, Nguyen QNN, Tantillo DJ,

Ichimura AS, Toscano JP, Fukuto JM. 2016. The chemical biology of the

persulfide (RSSH)/perthiyl (RSS·) redox couple and possible role in biological

redox signaling. Free Radic Biol Med 101:20-31.

42. Luebke JL, Shen J, Bruce KE, Kehl-Fie TE, Peng H, Skaar EP, Giedroc DP.

2014. The CsoR-like sulfurtransferase repressor (CstR) is a persulfide sensor in

Staphylococcus aureus. Mol Microbiol 94:1343-1360.

43. Ohta, S. 2015. Molecular hydrogen as a novel antioxidant: overview of the

advantages of hydrogen for medical applications. Methods Enzymol 555: 289-317.

44. Smith DA, Sessions AL, Dawson KS, Dalleska N, Orphan VJ. 2017. Rapid

quantification and isotopic analysis of dissolved sulfur species. Rapid Commun

Mass Spectrom 31: 791-803.

45. Bartlett JK, Skoog DA. 1954. Colorimetric determination of elemental sulfur

in hydrocarbons. Anal Chem 26:1008-1011.

Figure legends
685

FIG 1. Application of methylene blue method in H_2S and S^0 quantification.

(A) HSSH and H_2S can react with methylene blue reagents to generate

thiaazaheterocyclic dyes but S_8 cannot. Data are from three replicated experiments

(n=3) and presented as average $\pm$ s.d; (B) Mechanisms of how HSSH and H_2S

react with methylene blue reagents. Reactions 1 and 2 are from reference (24).

Other reactions are proposed by authors.

692
693

694 **FIG 2. Phylogenetic analysis of 289 sulfite reductases.** Only part origins of the

695 enzymes were shown. Details of these enzymes and their origins are shown in

696 Table S1.

697
698

FIG 3. Analysis of the products generated from sulfite reductase-catalyzed

reactions. Purified CysIJ (A), MET5&10 (B), and SPO2634 (C) were mixed with

NADPH andsulfite; purified Sir1 (D) was mixed with ferredoxin, ferredoxin-

NADP+ oxidoreductase, NADPH, and sulfite. The reactions were conducted at 25°C

for 10 min. Products were derivatized by mBBr. After mBBr derivatization, H_2S

became mB-SH and mB-S-mB, and HSSH became mB-SSH. The derivatives were

detected by LC-ESI-MS. MS signal intensities of H_2S (mB-SH + mB-S-mB) and

HSSH (mB-SSH) were shown. MS spectra of these chemicals were shown in Fig.

S3; (E) The enzyme-produced H_2S_n was derivatized by methyl

trifluoromethanesulfonate and analyzed using HPLC with UV detection. H_2S_n

mixture was used as the standard; (F) The enzyme-produced H_2S_n was quantified

using cyanide method and shown as total S^0; (G) The enzyme-produced H_2S was

quantified by HPLC; (H) Sulfite consumption, S^0 production, and H_2S production

from *E. coli* MG1655 strains overexpressing sulfite reductases; (I) Sulfite

consumption, S^0 production, and H_2S production from *S. cerevisiae* S288C

overexpressing MET5&10; (J) Sulfite consumption, S^0 production, and H_2S

production from *R. pomeroyi* DSS-3$\Delta sqr\Delta fccAB$ overexpressing SPO2634. Controls

are strains without sulfite reductase overexpression. For F–J, data are from three

replicated experiments (n=3) and presented as average ± s.d. T-tests were performed

to calculate the p-values. Asterisks indicate statistically significant difference (* $p <$

0.05, ** $p < 0.01$, ***$p < 0.001$, ****$p < 0.0001$).

FIG 4. Phylogenetic analysis of 735 cysteine synthases. Only part origins of the

enzymes were shown. Details of these enzymes and their origins are shown in

Table S2.

FIG 5. Pathway of cysteine synthesis using different sulfur donors.

FIG 6. Kinetic assays of cysteine synthases with different substrates. Details of

the data are shown in Table S3 and Fig. S12–S14.

FIG 7. Process of the assimilatory sulfate reduction. Zero valent sulfur (S^0) is the

key intermediate linking sulfate reduction and cysteine synthesis and H_2S is a side-

product.

733
734

Candidatus	Planctomycetota
Streptomyces	Gammaproteobacteria
Pseudomonas	Glycine
Nocardia	Bacillus
Amycolatopsis	Fusarium
Symbiodinium	Francisella
Colletotrichum	Streptococcus
Ceratobasidium	Mycolicibacterium
Clostridium	Aspergillus
Corynebacterium	Acidobacteriota
Burkholderia	Chlorella
Citrus	Leptospira
Chloroflexota	Mycobacteroides
Rhizoctonia	Mesorhizobium
Staphylococcus	Flavobacterium
bacterium	Nonomuraea
Aerococcus	Micromonospora
Xenorhabdus	Bradyrhizobium
Parcubacteria	Sphingobacterium
Paraburkholderia	Coemansia
Vibrio	Actinomycetota
Paenibacillus	Cafeteria
Arthrobacter	Rhodobacteraceae
Pseudovibrio	Ancylostoma
Scenedesmus	uncultured