A cysteine-rich domain of the Cuf1 transcription factor is required for high copper stress

sensing and fungal virulence

Corinna Probst [1], Catherine A. Denning-Jannace [4], Lukas M. du Plooy [2], Charles Giamberardino [1], Yohannes

Asfaw [5] Katherine J. Franz [4], J. Andrew Alspaugh [1,2,3*]

Duke University School of Medicine, Durham, NC, USA: Departments of Medicine [1], Molecular Genetics

and Microbiology [2], and Cell Biology [3]

Duke University, Durham, NC, USA: Department of Chemistry [4]

Duke University, Durham, NC: Division of Laboratory Animal Resources [5]

Running title**: High copper sensing in the human fungal pathogen *Cryptococcus neoformans***

Keywords**: cell wall, nutritional immunity, metal homeostasis, fungal pathogenesis**

(*) Corresponding Author: J. Andrew Alspaugh, andrew.alspaugh@duke.edu

Abstract

The ability to sense, import but also detoxify copper (Cu) has been shown to be crucial for microbial pathogens to survive within the host. Previous studies conducted with the opportunistic human fungal pathogen *Cryptococcus neoformans* (*Cn*) have revealed two extreme Cu environments encountered during infection: A high Cu environment within the lung and a low Cu environment within the brain. However, how *Cn* senses these different host Cu microenvironments, and the consequences of a blunted Cu stress adaption for pathogenesis, are not well understood.

In contrast to other fungi, *Cn* has a single transcription factor, Cuf1, to regulate adaptive responses to both high- and low-Cu stress. Sequence analysis of *Cn* Cuf1 identified three conserved cysteine (Cys)-rich regions that may play a role in Cu sensing. We mutated the 1st Cys-rich region within the *CUF1* gene to investigate its role for *Cn* high Cu stress sensing. Subsequent analysis of Cuf1 transcriptional activity and target gene promoter binding demonstrated that the 1st Cys-rich region is required for Cuf1 transcriptional activity in high Cu stress. We performed an inhalational murine infection to analyze the effects of a blunted high Cu stress response on pathogenesis. No significant differences in lung fungal burden were observed based on variable Cuf1 activity. However, strains with defective high Cu stress regulation induced a markedly altered immune response in mice. Based on these findings, we hypothesize that Cuf1-driven high Cu responses are not required for initial survival but instead modulate immune recognition and inflammation within the mouse lung.

Importance

Copper is an essential micronutrient required for survival in all kingdoms of life as it is used as a

catalytic cofactor for many essential processes in the cell. In turn, this reactivity of copper ions makes

elevated levels of free copper toxic for the cell. This dual nature of copper-essential for life but toxic at

elevated levels- is used by our innate immune system in a process called nutritional immunity to combat

and kill invading pathogens. In this work we explore how the fungal human pathogen *Cryptococcus*

neoformans senses high copper stress, a copper microenvironment encountered within the host lung.

We identified a specific cysteine-rich region within the copper responsive transcription factor Cuf1 to be

essential for high copper stress sensing. Mutation of this region led to an impaired high copper stress

adaptation, which did not affect fitness of the yeast but did impact immune recognition and

inflammation inside the host lung.

Introduction

Copper (Cu) is an essential trace element in both microbes and mammalian cells, serving as a

catalytic cofactor for a diverse range of cellular processes including iron uptake and distribution;

mitochondrial cytochrome oxidase activity; Cu, Zinc (Zn) superoxide dismutase catalysis; and melanin

pigment production [1, 2]. Although Cu is required for life, elevated Cu levels are toxic due to its ability

to induce uncontrolled redox cycling, causing the formation of toxic reactive oxygen species (ROS) and

oxidative damage of DNA, proteins, and lipids [3, 4]. Free Cu ions were also shown to induce protein

aggregation, triggering protein miss-folding and precipitation [5, 6]. Furthermore, Cu has been shown to

irreversibly displace iron (Fe) from Fe-S-cluster containing proteins, rendering these proteins non

functional [7, 8]. Therefore, organisms have evolved processes to sense and tightly regulate cellular Cu

availability, to maintain but not exceed the required cellular Cu quota [2, 4, 9-11].

Pathogenic microbes encounter unique extremes of Cu stress. Infected hosts often employ Cu-

sequestering strategies known as "nutritional immunity" to limit Cu availability for invading pathogens

[12-17]. In other sites, such as the macrophage phagolysosome, the host ATP7A Cu pump actively

imports Cu to toxic concentrations to kill engulfed microorganisms [18] and, more recently, Mohamed

and colleagues demonstrated that brain resident microglia provide Cu to a phagocytosed fungal

pathogen in the central nervous system (CNS) [13].

The basidiomycete *Cryptococcus neoformans* (*Cn*) is an opportunistic fungal pathogen that causes

life-threatening disease in immune compromised individuals [19, 20]. *Cn* is an environmental fungus

frequently isolated from degrading plants, and soil contaminated with bird guano. Infectious spores or

desiccated yeast cells are inhaled by mammalian hosts from the environment, resulting in an initial

pulmonary infection [20]. In patients with weakened immune systems, including those with late-stage

HIV infection, *Cn* disseminates through the bloodstream, crosses the blood-brain barrier, and colonizes

the brain, where it causes lethal infection [19, 20]. Previous studies have revealed that *Cn* encounters

two extreme Cu environments during infection. During the initial infection within the lung, *Cn* senses a

high Cu environment. In contrast, once the microorganism reaches the CNS, it senses low Cu availability

[21, 22]. Thus, to maintain its proper cellular Cu quota, *Cn* needs to switch from Cu-detoxification during

pulmonary infection to Cu-acquisition upon entering the CNS.

Several adaptive strategies for *Cn* Cu homeostasis have been described. To combat high Cu stress, *Cn*

induces the Cu-detoxifying metallothionein genes *MT1* and *MT2*, the ROS detoxifying superoxide

dismutase (*SOD*) 1 gene, as well as *ATM1,* a gene encoding a mitochondrial ABC transporter required to

provide a sulfuric precursor for cytosolic Fe-S cluster biosynthesis [23]. These genes were previously

shown to be required for *Cn* survival inside macrophages or in a mouse model of *cryptococcosis* [21, 24-

26]. To combat the Cu-restrictive CNS environment, *Cn* induces its two high affinity copper importer

genes *CTR1* and *CTR4*, as well as the *CBI1* gene, encoding a Cu-binding protein required for efficient Cu

acquisition [22, 27, 28]. Additionally, Smith and colleagues described a transcription factor-mediated

switch between the Cu/Zn-dependent Sod1 and a Cu-independent/ manganese (Mn)-dependent Sod2 to

prioritize Cu towards other required Cu-dependent pathways when labile Cu is limited [11]. Even though

these adaptive strategies have been described, it is still unclear how they function together to facilitate

adaption against the encountered host Cu microenvironment and how failure of adaption to high or low

Cu stress effects *Cn* survival and pathogenesis.

Interestingly, most fungi employ two distinct transcription factors to control responses to high and

low Cu stress [2]. For example, the *Saccharomyces cerevisiae* (*Sc*) Ace1 transcription factor is activated in

a high Cu environment to regulate gene expression to protect the yeast from toxic Cu levels [29, 30],

while the *Sc* Mac1 transcription factor is activated by low Cu environments to favor Cu acquisition [31].

In contrast, *Cn* maintains a single Cu-responsive transcription factor Cuf1, that controls both high and

low Cu stress responsive genes [23]. Deletion of *CUF1* renders *Cn* less virulent in an intravenous mouse

model of *cryptococcosis* [32], further underscoring the importance of Cu homeostasis in *Cn* pathogenesis

and suggests the potential of Cuf1 as a fungal-specific intervention target to disrupt pathogen-driven

adaptions in the host nutritional environment during infection. However, the actual mechanism of gene

regulation by Cuf1 in response to various Cu stresses is not well understood and has yet to be

elucidated.

Sequence comparison between *Cn*Cuf1 and other fungal Cu-responsive transcription factors

identified the highest homology within the first 39 amino acids (aa) of the N-terminal domain [23]. This

region includes a Zn-binding motif and the highly conserved positively charged 'KGRP' motif that is

crucial for binding the minor groove of the DNA binding sites [29, 31, 33-36]. Mutational studies in

*Cn*Cuf1 have demonstrated that the KGRP motif is required for regulation of both high and low Cu

responses, consistent with the role of Cuf1 in transcription under both conditions [23]. Following the

minor groove binding region, the similarity between Cuf1 and other Cu-responsive transcription factors

decreases considerably, except for three cysteine (Cys)-rich regions: The first Cys-rich region is similar to

the Cys-rich region found in the high Cu-responsive transcription factor *Sc*Ace1. The second and third

Cys-rich regions are not found in *Sc*Ace1 but are conserved in the low Cu-responsive transcription factor

*Sc*Mac1 [23]. A role in Cu-binding and regulation of transcriptional activity has been established for the

Cys-rich regions in *Sc*Ace1 and *Sc*Mac1. In both cases Cu binding to the Cys-rich region of these

transcription factors acts as an ON-OFF switch either activating (Ace1-like transcription factors) or

deactivating (Mac1-like transcription factors) transcriptional activity [29, 33, 34, 37]. However, in

contrast to Ace1- and Mac1-like transcription factors, *Cn*Cuf1 binds and regulates genes in response to

low and high Cu stress, and additionally is bound constitutively to a rather unexpected subset of genes,

regardless of the encountered Cu stress [23]. These findings point towards a more complex regulation of

Cuf1 dependent gene regulation. Specifically, a single transcription factor must be activated by a specific

Cu stress condition to bind to and direct gene expression of its target genes, but also has to be activated

by the opposed Cu stress condition where it also has to bind to and direct gene expression of a

completely different set of target genes, while still being bound constitutively at a small subset of

promoters independent of the present Cu stress. Consequently, the requirements of Cuf1 transcriptional

activity at high and low Cu stress suggests a more multi-faceted model of gene regulation than the

ON/OFF model previously described for Ace1- or Mac1-like transcription factors.

Here we present an initial study of the role of the Ace1-like Cys site for high Cu stress sensing and

gene regulation by *Cn*Cuf1. We analyzed the consequential role of a blunted high Cu stress response for

cupro-enzyme regulation *in vitro* and its effects on pathogenesis *in vivo* within the mouse lung

environment using an inhalational mouse model of *cryptococcosis*.

RESULTS

Mutation of the Ace1-like Cys site in the Cuf1 transcription factor renders the fungus susceptible to

high Cu stress.

Cuf1 is the central Cu-responsive transcription factor known to regulate high and low copper genes in

Cn. Previous research conducted on other fungal transcription factors implicates conserved Cys-rich

motifs to bind Cu and regulate transcriptional activity in response to cellular Cu levels [29, 33, 34, 37]

(Fig1 A). The first Cys-rich region encoded by *CUF1* is formed by an 11 aa motif, containing 4 cysteines,

which are highly conserved among Cuf1 and Ace1-like transcription factors [23]. To elucidate how Cuf1

senses high Cu and is guided to its target genes in response to high Cu stress, we mutated all 4

conserved Cys located in the Ace1-like Cys rich region to alanine (Cuf1-FLAG ΔAce1, Fig 1A). Alleles

encoding FLAG-tagged versions of the Cuf1 wild type (WT) protein and the Cuf1 ΔAce1 mutant were

incorporated into the *cuf1Δ* mutant strain at the same genomic site (safe haven 1), both expressed by

the *CUF1* promoter (Fig 1A-B).

To explore the effects of the Ace1-like Cys site mutation on Cuf1-FLAG protein levels at high and low

Cu stress, we conditioned the *CUF1*-FLAG WT and two independent *CUF1-FLAG* ΔAce1 strains for 3h in

either SC medium + 1 mM $CuSO_4$ (= high Cu) or SC medium + 1 mM of the extracellular Cu chelator

bathocuproinedisulfonic acid (BCS,= low Cu) at 30°C. Western blot analysis using a FLAG-HRP conjugate

demonstrated no differences in Cuf1-FLAG protein levels in response of high or low Cu stress between

the WT or the Ace1- like Cys site mutant proteins, indicating that the Ace1-like site mutation does not

affect Cuf1 protein levels (Fig 1C).

To determine the physiological effects of the Ace1-like Cys-rich region on regulation of Cu

homeostasis during Cu stress, we assessed the ability of each allele to complement the *cuf1Δ* growth

phenotype in high and low Cu stress environments. In all stress conditions, the *CUF1-FLAG* WT allele

fully restored growth of the *cuf1Δ* mutant strain, demonstrating that the FLAG-tag did not compromise

Cuf1 protein function. Similarly, *CUF1-FLAG* ΔAce1 complemented the *cuf1Δ* growth defect in Cu-limiting

conditions, however it failed to restore growth in a high Cu environment (Fig1D). This finding suggests a

specific role for the Ace1- like Cys rich region for regulating transcriptional activity in response to high Cu

stress.

The Ace1-like Cys site is required for Cuf1 to specifically bind to and regulate high Cu responsive genes

To test whether the growth phenotype observed under high Cu for both the *cuf1Δ* strain and

the *cuf1Δ* strain expressing the *CUF1*-FLAG ΔAce1 allele were due to a similar failure to activate Cuf1

target genes in response to high Cu stress, we assessed transcript levels for established high and low Cu

regulated genes in each strain. The *MT1* gene encoding the Cu detoxifying metallothionein 1 protein and

the *ATM1* gene, encoding for a mitochondrial inner membrane ABC transporter involved in cytosolic Fe-

S assembly, were used to assess transcriptional responses to high Cu stress [21, 24]; and the *CTR4* gene,

encoding a high affinity Cu importer, and the *CBI1* gene, encoding the copper binding and release

protein 1, were used to assess responses to low Cu stress [22, 23, 27]. The WT and the *cuf1Δ* mutant

strain, as well as the *cuf1Δ* mutant strain expressing either *CUF1-FLAG* WT or the *CUF1-FLAG-*ΔAce1

mutant allele, were incubated for 3h in SC medium supplemented with either 1 mM $CuSO_4$ (=high Cu) or

1 mM BCS (=low Cu) at 30°C prior to RNA extraction. Transcript levels of the Cu-response genes for each

strain in each condition were determined by quantitative real-time PCR (qRT-PCR). Consistent with prior

results, the *MT1* and *ATM1* genes were induced in the WT strain under high Cu conditions, and the *CTR4*

and *CBI1* genes were induced in low Cu conditions (Fig 2 A-B) [23]. In contrast, the transcript levels of

both high Cu and low Cu genes were not induced in either condition in the *cuf1Δ* mutant. The *CUF1-*

FLAG WT strain fully restored gene expression in both conditions. However, consistent with the

observed growth phenotype on high Cu media, mutation of the Ace1-like Cys site was associated with

defective transcriptional activation of high Cu-regulated genes (*MT1* and *ATM1*, Fig 2 A), while fully

restoring WT transcript levels of genes involved in low Cu stress response (*CTR4* and *CBI1,* Fig 2 B)*.*

The Cuf1 transcription factor was previously demonstrated to directly bind the promoters of

these high and low Cu reporter genes [23]. We used chromatin immunoprecipitation followed by qRT-

PCR analysis (ChIP-PCR) to determine whether the observed differential transcript levels in each strain

were associated with changes in Cuf1 promoter binding. We specifically chose oligonucleotides

corresponding to the *CTR4* and *MT1* promoters as indicators of low- and high-Cu stress gene promoter

binding, respectively. To do so, we incubated the strains in the identical high- and low-Cu stress

conditions as above prior to protein-DNA crosslinking and chromatin immunoprecipitation. Enrichment

of the *MT1* or *CTR4* promoter fragments in each sample was calculated relative to the WT strain and was

further normalized against the *TUB1* tubulin gene as a non Cuf1 target gene. Consistent with previous

studies, we observed a significant enrichment of the *MT1* promoter, but not the *CTR4* promoter, in the

Cuf1-FLAG WT sample conditioned in high Cu stress and enrichment of the *CTR4* promoter sequence

during low Cu stress [23](Fig 2C-D). In contrast, no enrichment of the *MT1* promoter was observed by

ChIP-PCR in the *cuf1Δ + CUF1-FLAG* ΔAce1 strain conditioned in high Cu stress, whereas the *CTR4*

promoter was enriched in this strain to similar levels as in the Cuf1-FLAG WT allele expressing strain (Fig

2C-D). Taken together, these data show that the Ace1-like Cys-rich region is required for Cuf1 binding to

and regulation of high Cu-responsive promoters while it does not impact binding or regulation of low

Cu-responsive promoters.

Unlike other Cu-responsive transcription factors, Cuf1 does not only bind and regulate specific

gene promoters at either low and high Cu conditions. It also constitutively binds the promoters of a

group of genes regardless of changes in Cu concentration [23]. We therefore performed transcript and

Cuf1-promoter binding analysis for a subset of these genes (*CDA2*, *CFO1*) in both Cu stress conditions

(Supp. Fig 1). The Chitin deacetylase 2 (Cda2) was previously described to be involved in chitosan

synthesis within the lung environment [38]. *CDA2* was identified as a Cuf1-regulated gene, with Cuf1

binding of its promoter at high and low Cu levels and a modest but statistically significant transcript

induction only during high Cu stress [23]. The ferroxidase gene (*CFO1*), which is involved in high affinity

iron uptake [39], was likewise identified to have constitutive promoter binding by Cuf1. However, in

contrast to *CDA2*, *CFO1* is induced by low Cu stress [23]. In line with previous published RNAseq data, we

observed a decrease in *CDA2* transcript in the *cuf1Δ* mutant strain in presence of high Cu stress which

was restored by the expression of the WT *CUF1-FLAG* allele, but not the *CUF1-FLAG-ΔAce1* mutant allele

(Supp Fig 1A), indicating a role for the Ace1-like site in Cuf1-dependent regulation of the *CDA2* gene

during high Cu stress. However, in contrast to *MT1* or *CTR4,* no Cuf1-associated enrichment of the *CDA2*

promoter fragment was found by ChIP-PCR at either stress condition (Supp. Fig 2B-C).

In contrast to *CDA2*, which seemed to be only regulated by Cuf1 during high Cu stress, we

observed a ~55% reduction of the *CFO1* transcript level in the *cuf1Δ* background in both Cu stress

conditions, suggesting *CFO1* expression to be constitutively driven by Cuf1 (Supp. Fig 1A). *CFO1*

transcript levels were mostly restored in the *cuf1Δ* mutant by the expression of either the WT *CUF1-*

FLAG allele or the *CUF1-FLAG* ΔAce1 mutant allele (Supp Fig 1A). Similar to *CDA2* promoter binding*,* no

enrichment of the *CFO1* promoter fragment was found in cells conditioned with high Cu stress

conditions in either strains (Supp. Fig 2B). However, we did observe a modest enrichment of the *CFO1*

promoter at low Cu stress, independent from the absence or presence of the Ace1-like Cys site (Supp.

Fig1C). This finding suggests that, in contrast to Cuf1-target genes whose promoters are bound

exclusively by Cuf1 during high Cu stress, the Ace1-like Cys site is not absolutely required to support

CFO1 expression in the presence of high Cu stress.

Together, these transcript and promoter binding data demonstrate that the Ace1-like Cys site is

required for the Cuf1 transcription factor to transcriptionally activate high Cu-responsive genes but not

low Cu-responsive genes. This activity likely explains the defective cellular adaptation to high Cu stress

for strains with mutations of this site.

The Ace1-like Cys site is required for proper regulation of ROS detoxification and melanization in high

Cu environments.

Cn encounters both low- and high-Cu microenvironments in the infected host [22]. We therefore

assessed the role of Cuf1 and the Ace1-like Cys region in the activity of cupro-enzymes required for *Cn*

stress resistance and survival within these microenvironments in the human host.

Laccase 1 and melanin production:

The *Cn* Lac1 protein is a Cu-containing diphenol oxidase/laccase involved in formation of

melanin, a cell wall-associated pigment that protects the fungal from host-derived oxidative and

nitrosative stress as well as from antifungal drugs [40]. The process of melanization is tightly linked to Cu

homeostasis and Cuf1 in multiple ways. First, melanin production requires cellular Cu as Lac1 oxidase

activity is dependent on Cu as a catalytic cofactor [41]. Second, *LAC1* gene expression is controlled in a

Cuf1-dependent manner [23, 28, 42, 43](Fig 3A). To analyze the effects of the Ace1-like Cys site

mutation on melanin production, we performed a combination of an *in vitro* pigmentation assay using L-

DOPA media plates supplemented with BCS or $CuSO_4$ to create a Cu limiting or Cu excess environment as

well as *LAC1* transcript analysis in L-DOPA medium in the presence of Cu limitation or excess (Fig 3B-D).

To test for melanin production, we incubated the WT strain, the *cuf1Δ* mutant, and the *cuf1Δ*

strain expressing either *CUF1-FLAG* WT or the ΔAce1-like Cys site mutant allele on L-DOPA medium

supplemented with either 10 µM BCS (to slightly restrict Cu) or 10 µM $CuSO_4$ (to provide excess Cu). As

controls, we included the completely melanin-defective *lac1Δ* mutant strain and the *cbi1Δ ctr4Δ* double

mutant strain that is impaired in copper import [27, 41]. For each strain, relative *LAC1* transcript levels

were measured after 3h of incubation in liquid L-DOPA medium similarly supplemented with either 10

μM BCS or 10 μM CuSO$_4$.

The WT strain was strongly melanized by 72h, and the addition of excess Cu to the medium

resulted in a higher degree of pigmentation (Fig 3B). In contrast, the *cuf1Δ* mutant displayed decreased

melanin in Cu-limited conditions, similar to that of the *cbi1Δ ctr4Δ* Cu import mutant (Fig 3B). In these

conditions, *LAC1* gene expression in the *cuf*1Δ mutant is similar to WT, suggesting that its decreased

melanin formation is due to failed induction of Cu import genes, such as *CTR4*, resulting in reduced

levels of intracellular Cu (Fig 3 C-D). Consistent with this hypothesis, we observed full restoration of

melanization in the Cu-import mutant, when Cu is supplemented to the L-DOPA medium, restoring Lac1

Cu loading and thus function (Fig 3B).

In contrast to the Cu import mutant, melanin formation was not restored in the *cuf1Δ* mutant

when the medium was supplemented with Cu. On the contrary, the *cuf1Δ* strain demonstrated an even

more severe melanin-deficient phenotype, similar to that of the *lac1Δ* mutant (Fig 3 B). In this condition,

we observed a marked relative decrease in *LAC1* transcript levels in the *cuf1Δ* mutant strain, consistent

with the role of Cuf1 supporting *LAC1* expression when Cu is in excess. (Fig 3C).

Reintroduction of the *CUF1-FLAG* WT allele fully restored WT levels of melanin production in

both high and low Cu environments, as well as WT levels of *LAC1* and *CTR4* expression (Fig 3B-D). In

contrast, the *cuf1Δ* strain expressing the ΔAce1-like Cys site mutant allele only restored melanization

when Cu was limited but not when Cu was in excess (Fig 3B). In this strain, *CTR4* expression was fully

restored in Cu-limiting conditions, but *LAC1* expression was not restored in Cu excess (Fig 3C-D). Taken

together, these data demonstrate a major role for Cuf1 in controlling melanin formation, by ensuring

adequate levels of Cu as catalytical cofactor for Lac1 in low Cu conditions, and by supporting *LAC1* gene

expression in high Cu conditions. Additionally, the Cuf1 Ace1-like Cys site is required for ensuring proper

LAC1 expression and melanization when Cu is in excess but is not required for melanization during Cu

limitation.

<u>Superoxide dismutase and ROS detoxification:</u>

The Cu,Zn-superoxide dismutase Sod1 helps to protect the fungal cell from ROS generated by

host immune cells during infection [26]. Previous studies demonstrated that Cuf1 binds to the *SOD1*

promoter in high Cu stress and induces *SOD1* gene expression. In contrast, during prolonged Cu

starvation, *SOD1* expression is repressed to divert limited Cu to other cell processes. In this condition,

Cuf1 binds to the *SOD2* promoter and induces the expression of a cytosolic retained variant of the Mn-

dependent mitochondrial superoxide dismutase 2 (Sod2) to uncouple ROS detoxification from a Cu-

requiring enzyme when cellular Cu is scarce [11]. Therefore, deletion of *CUF1* renders the cells

vulnerable to ROS stress not only during high Cu stress due to failure of up-regulation of *SOD1* but also

during low Cu stress due to inability to switch to a Cu-independent ROS detoxification pathway (Fig 4A).

To test effects of the Ace1-like Cys site mutation on cellular ROS detoxification, we incubated

the WT and *cuf1Δ* strains as well as strains expressing *CUF1-FLAG* WT or the ΔAce1-like Cys mutant allele

in the presence of increasing concentrations of the superoxide-generating compound menadione by disk

diffusion (Fig 4B). Consistent with previous studies, the *cuf1Δ* mutant showed an increased susceptibility

to menadione, measured by zone size of growth inhibition, in high or low Cu conditions (Fig 4C) [11].

Both Cu-stress phenotypes were fully restored in the *cuf1Δ* strain expressing the *CUF1-FLAG* WT allele.

In contrast, the strain expressing the *CUF1-FLAG* ΔAce1-like Cys site mutant allele only restored the ROS

resistance during low Cu stress but not high Cu stress (Fig 4C).

To fully analyze the role of the Ace1-like Cys site in regulating Sod-driven ROS detoxification, we

measured Sod activity in whole cell extracts using an in-gel Sod activity assay [10] from cells conditioned

in either low or high Cu stress. The Sod activity gels demonstrated three activity bands from cell extracts

from the WT strain incubated in low Cu conditions. The lowest band likely represents Sod1, while the

middle band likely represents Sod2 activity given their absence in the respective *sod1Δ* and *sod2Δ*

mutants (Fig 4D). The highest of the three bands was absent in the *cuf1Δ* and *sod2Δ* mutant strains (Fig

4D, yellow hash tag), and likely represents the cytosolic Sod2 variant that is specifically enriched in Cu

limitation in a Cuf1-dependent manner [11]. This Sod activity is restored in the *cuf1Δ* mutant by

introduction of either the *CUF1-FLAG* WT or the ΔAce1-like Cys site mutant allele, indicating that Cuf1-

dependent formation of cytosolic Sod2 during Cu-limiting conditions is not dependent on the Cuf1 Ace1-

like Cys site (Fig 4C-D).

The patterns of SOD activity varied when the strains were conditioned in high Cu stress, with

two predominant bands of similar intensity in WT cells. Both bands were significantly reduced in

intensity in the *cuf1Δ* mutant strain. Moreover, in the *sod1Δ* strain, these two bands were only faintly

present, and the upper of these two was either not present or altered in its molecular size (Fig 4D,

asterix), suggesting that this upper band represents Sod1 activity. While expression of *CUF1-FLAG* WT

restored the Sod activity bands to WT-like levels, we observed identical patterns of reduced Sod activity

in the *cuf1Δ* mutant as well as the *cuf1Δ* mutant expressing the *CUF1-FLAG* ΔAce1-Cys mutant allele,

indicating that the first Cys-rich site is required for Cuf1-depenedent induction of Sod1 protein activity in

response to high Cu stress (Fig 4D). Consistent with this finding, the Cuf1 Ace1-like Cys site is required

for supporting Cuf1-dependent *SOD1* transcript induction in high Cu conditions (Fig 4E). Together, these

phenotypic data, enzyme activity assays, and transcript assessments demonstrate the importance of the

Ace1-like Cys site for Cuf1-dependent regulation of ROS detoxification in a high Cu stress environment.

Consequences of a blunted Cuf1 high Cu stress response on fitness and immune response inside the

host lung environment

A previous study conducted by Waterman *et al.* established a role for Cuf1 for virulence in an

intravenous model of *Cn* infection, demonstrating a decreased fungal burden of the *cuf1Δ* mutant

compared to WT in the mouse brain but not in mouse lung [32]. More recent studies have suggested

that these different anatomic sites represent different Cu environments encountered by *Cn* during the

course of infection [21, 22, 27]. Using high and low Cu promoter driven Luciferase strains, one study

found an up-regulation of genes typically induced during Cu deficiency in the brain, while the lung

environment induced genes required for Cu detoxification [21]. This finding was further underscored by

assessing the effects of Cu homeostasis mutant strains on virulence and survival at the different sites of

infection: Strains lacking components of the Cu import machinery are less fit within the mouse brain

environment, while strains lacking components required for Cu detoxification are less fit in the mouse

lung environment [21, 22, 27]*.*

Given the severe *in vitro* phenotypes found for the *cuf1Δ* mutant and the ΔAce1-like Cys site

mutant strain, we hypothesized that these strains would display altered physiology within the high Cu

environment of the lung. We inoculated female A/J mice by inhalation with the WT strain, the *cuf1Δ*

mutant, or the *cuf1Δ* mutant reconstituted with either the WT *CUF1-FLAG* allele or the *CUF1-FLAG*

ΔAce1 mutant allele. Lungs were harvested at 14 days of infection, a time at which lung infection is well

established but before animals succumb to the effects of a CNS infection, to assess lung fungal burden

as well as patterns of inflammation by histology (Fig 5). Consistent with prior studies, we did not observe

significant difference in lung fungal burden for any of the analyzed strains (Fig 5A) [32]. However, we did

note a significant increase in lung wet weight in animals infected with the WT or *CUF1-FLAG* WT strain

(Fig 5B). Additionally, we observed macroscopic differences between WT and *CUF1-FLAG* WT-infected

lungs compared to *cuf1Δ*- and the *CUF1-FLAG* ΔAce1-like Cys site mutant-infected lungs: WT-infected

lungs showed more frequent and larger foci of tissue abnormality compared to lungs infected with

either the *cuf1Δ* mutant or the ΔAce1-like Cys site mutant strain (Fig 5C, foci are indicated with white

arrows).

Histopathology analysis of fixed lung tissue confirmed altered immune reactions between these

two groups. Lungs infected with the WT strain displayed expected patterns of diffuse inflammation

surrounding single and grouped encapsulated yeast cells (Fig 5D) [44]. In the regions of most intense

inflammation, the inflammatory cell infiltration often obscured the clear definition of the underlying

alveolar structures. Titanized and non-titan yeast cells were observed throughout the lung tissue, both

in areas of inflammation as well as in surrounding less inflamed small airways and alveoli. In contrast

lungs infected with the *cuf1Δ* mutant strain displayed inflammation mostly confined to well

circumscribed foci, similar to the early granulomas described previously in studies of *Cn* strains with

slow growth phenotypes due to defects in mitochondrial function [44]. In these *cuf1Δ*-infected lungs, we

observed a sharp distinction between normal-appearing lung tissue and the foci of inflammation. Most

encapsulated yeast cells were present in these inflammatory foci with fewer fungal cells than WT

observed in the surrounding normal lung. Few titan cells were observed.

Lungs infected with *CUF1-FLAG* WT strain demonstrated very similar patterns of inflammation as

observed in WT-infected lungs. In contrast, the *CUF1-FLAG* ΔAce1-like Cys site mutant strain generated

an inflammatory response that had mixed features to both WT and *cuf1Δ* infections. Similar to *cuf1Δ*

infections, there was often a sharp distinction between inflamed and normal lung tissue with most yeast

cells observed in regions with the highest degree of inflammation. However, regions of inflammation

were less frequently circular, and there was more variability in yeast cell size than in the *cuf1Δ*

infections. Based on these findings, we hypothesize that Cuf1-regulated high Cu responses are not

required to colonize the mouse lung but modulate immune recognition and inflammation at the site of

infection.

Discussion

*Cn*Cuf1 is unique among fungal Cu-responsive transcription factors as it regulates a diverse

spectrum of Cu responses, combining the role of a high Cu-responsive transcription factor with the role

of a low Cu-responsive transcription factor [23]. Accordingly, it shares conserved motifs with both high

and low Cu-responsive transcription factors previously identified in ascomycete yeast Cu transcription

factors [23]. However, how these factors work together to regulate this rather complex Cuf1-driven Cu

response has not yet been elucidated. An interspecies cross-complementation study conducted with the

Aspergillus fumigatus (*Af*) AceA high Cu-responsive transcription factor demonstrated that AceA

homologs from ascomycete yeasts did not functionally complement the high Cu growth defect observed

in the *Af ΔaceA* mutant, suggesting that, even though the N-terminal DNA binding and the Cu sensing

motifs are highly conserved among these high Cu-responsive transcription factors, the motifs might

function differently in *Aspergillus* species than in their yeast counterparts [45]. This finding suggests that

despite using similar motifs/domains, different fungal clades may have evolved slightly different

molecular mechanisms to sense and/or transcriptionally regulate Cu stress responses. *Cn* Cuf1 is another

example of divergence of fungal Cu-responsive transcription factors, having evolved into a dual Cu-stress

responsive transcription factor by combining motifs from high- and low-Cu responsive transcription

factors with potentially yet unknown regulatory mechanisms.

The transition metal Cu has been established as an important virulence factor in all major

human fungal pathogens as it is used as a cofactor for several enzymes required for stress adaptation

within the host [10, 11, 40, 46, 47]. Consistently, Cu import genes and the Cu-responsive transcription

factors sensing and maintaining the cellular Cu quota were demonstrated to play a role in fungal

pathogenesis [22, 27, 32, 48-51]. For instance in *Candida albicans* (*Ca*), researchers demonstrated that

the low Cu-responsive *Ca*Mac1 transcription factor is required for host invasion, adhesion to host cells,

as well as antifungal tolerance [49]. Similarly, deletion of the low Cu-responsive MacA transcription

factor in *Af* attenuates pathogenicity in a neutropenic immunocompromised mice model [50].

Nutritional immunity responses for other transition metals mostly involve sequestering the

micronutrient from the site of infection [15, 17]. In contrast, Cu nutritional immunity responses appear

to be more dynamically regulated and are highly tissue dependent. While Cu might be restricted at one

site or time point of an infection, elevated and toxic Cu levels are present at other sites or time points to

act as an antimicrobial agent to kill the invading pathogen [10, 11, 13, 16, 21, 22, 48, 52]. Therefore, high

Cu stress responses are as important as Cu scavenging responses for pathogenesis. However, they must

be balanced to ensure proper adaptation to the Cu environment established at the particular infection

microniche.

Analysis of this dynamic shift of Cu nutritional immunity in the host and the fungal adaptive

stress response has been challenging to study in *Cn* since deletion of the *CUF1* gene ultimately blunts

both high and low Cu stress responses [23]. This makes deciphering the role of high and low Cu stress

responses during infection difficult as these stresses can have compensatory effects. For instance,

deletion of the *CTR4* Cu import gene was shown to be beneficial for *Cn* survival during pulmonary

infection, preventing an excessive influx of Cu and thus reducing the toxic effect of the high Cu lung

environment. In contrast, the same gene deletion resulted in reduced fungal virulence when the strain

was directly injected into the CNS, a site where Cu is limited [22]. Our mutational study aimed to identify

amino acid motifs specifically required for transcriptional activity in high Cu environments to uncoupled

Cuf1-driven high Cu stress responses from low Cu stress responses. In this work using a combination of

transcriptional analysis, Cu stress, and enzyme activity assays, we have established an essential role of

the Ace1-like Cys site for Cuf1 transcriptional activity during high Cu stress; mutation of this site does

not affect low Cu stress adaption. Using this site mutant, we were able to show the consequences of a

blunted high Cu stress adaptation on Cu-dependent virulence factors *in vitro* and on fungal growth and

immune recognition *in vivo.*

Recent work conducted with *Af*AceA demonstrated that mutation of its conserved Cys-rich

region attenuates virulence of the fungus similar to a Δ*aceA* mutant [45], highlighting the importance of

its Cys-rich region for activating Cu stress responses within the mouse lung environment. Similar to a

Δ*aceA* mutant in *Af*, deletion of the *CUF1* gene results in an attenuation of virulence [32]. However,

unlike the *Af* Δ*aceA* , a *CUF1* deletion did not affect *Cn* growth and survival inside the lung environment

[32, 45]. This observation stands in contrast with the severe *in vitro* phenotypes depicted by the *cuf1Δ*

mutant or the *CUF1-FLAG* ΔAce1 site mutant strains during high Cu stress. The observed *in vivo* growth

phenotype of the *cuf1Δ* and the *CUF1-FLAG* ΔAce1 mutant is also very different from the strongly

attenuated phenotype observed for a *Af* Δ*aceA* mutant. These very distinct outcomes of a blunted high

Cu stress response observed in the pathogenesis of two fungal pathogens in the same host environment

suggest different host-pathogen dynamics within the lung Cu environment, which are likely dependent

on the type of pathogen and its lifestyle inside the host tissue. Another possibility is that *Cn* might not

be as dependent on Cuf1 as *Af* on AceA for Cu detoxification *in vivo*. For instance, *Cn* could have evolved

alternate compensatory mechanisms to survive in the host lung environment which are independent

from Cuf1. To address these questions and to better understand the dynamics of Cu nutritional immune

responses within the lung environment, a more thorough assessment is needed at several different

stages of infection.

Although we did not observe significant differences in fungal burden between our WT and

mutant strains at two weeks of infection, we did observe macroscopic and microscopic differences

within the lungs of WT infected mice compared to mice infected with the *cuf1Δ* mutant or *CUF1-FLAG*

ΔAce1 site mutant. This observation suggests a role for Cuf1-driven high Cu stress responses for

modulating inflammatory responses within the lung environment, by mediating adaptive responses at

the fungal cell surface. This fungal feature is composed of ordered layers of polysaccharides and

mannosylated proteins and is the site of recognition and interaction with the host immune system:

Changes in carbohydrate composition as well as masking or unmasking of certain cell surface

components can modulate and define the pattern of inflammation caused by the pathogen [53-55]. For

instance, chitin has been associated with non-protective immune responses, and the degree of

unmasking coupled with the amount of chitosan within the *Cn* cell wall, modulates the severity of

immune recognition and inflammation within the host tissue [55, 56].

Previous work has suggested that the fungal cell wall plays a role in regulating intracellular Cu

levels during Cu stress [28]. In turn, Cuf1 regulates multiple genes associated with the fungal cell wall in

response to Cu stress [23, 28]. One example of these Cu-regulated cell wall genes is the chitin

deacetylase *CDA2,* an enzyme involved in chitosan synthesis. Our study suggests that *CDA2* is up-

regulated in response to high Cu stress and that this transcriptional activation is blunted in the *cuf1*Δ

null and ΔAce1-site mutant. Another study showed that cell wall architecture and cell wall chitosan

levels are significantly altered in a Cu homeostasis mutant strain during Cu limitation [28]. These findings

document a role for Cuf1 in modulating the fungal cell wall and in particular cell wall chitosan levels as

part of its Cu homeostasis response towards Cu stress. In this way, strains lacking chitosan in their cell

wall were shown to be more sensitive to high Cu stress [28]. We therefore propose a model in which

the observed differences in inflammatory patterns between WT-infected groups and the *cuf1*Δ mutant-

infected groups are caused by Cuf1-driven cell surface changes in response to the lung Cu environment.

We hypothesize that either loss of *CUF1* or simply a blunted high Cu-response prevents Cu-stress-driven

cell wall remodeling. This in turn, triggers a more focal, granuloma-like inflammatory response with

fungal containment within these inflammatory foci. Further investigations will more precisely define the

effects of Cu and Cuf1 on the fungal cell surface and their subsequent effects on fungal recognition and

inflammation at different stages and sites of infection.

Material and Methods

Ethics statement

All animal experiments conducted in this study were approved by the Duke University Institutional Animal Care and Use Committee (IACUC) (protocol #A102-20-05). Mice were handled according to IACUC guidelines.

Strains, media and growth conditions

Cryptococcus neoformans strains used in this study are shown in **Table S1.** Strains generated in this study were created using the *C. neoformans var. grubii H99* background. Strains were created using biolistic transformation as previously described [57]. Yeast extract (2%)-peptone (1%)-dextrose (2%) (YPD) medium supplemented with 2% agar and 200 μg ml-1 of hygromycin B (HYG) was used for colony selection after biolistic transformation. Transformants were screened by PCR for intended mutations and gene expression was confirmed by western blot. Plasmids and oligonucleotides used in this study for strain generation are listed in **Tables S2-S3.**

Strains were cultivated in synthetic complete (SC) medium (MP Biomedicals) at 30°C. Media was supplemented with indicated concentrations of $CuSO_4$ or the Cu^+ chelator bathocuproine disulfonate (BCS) to induce Cu sufficiency or deficiency, respectively.

Growth phenotype analysis was performed as previously described [28, 58]. Melanization assays were performed on L-3,4-dihydroxyphenylalanine (L-DOPA) media (7.6 mM L-asparagine monohydrate, 5.6 mM glucose, 22 mM KH_2PO_4, 1 mM MgSO4 heptahydrate, 0.5 mM L-DOPA, 0.3 μM thiamine-HCl, 20 nM biotin, pH 5.6) as previously described [28, 58]. L-DOPA plates were incubated at 30°C for 3 days.

Trichloroacetic acid (TCA)-based protein precipitation and western blotting

Cn cultures grown overnight for 18h in synthetic complete (SC) medium (MP Biomedicals) were

diluted to OD_{600} 0.5 and incubated for 2 h at 30°C in SC. Cu stress was then induced by the addition of

either 1 mM BCS or 1 mM $CuSO_4$, and cultures were incubated for 3 h at 30°C. TCA-precipitated protein

extracts were generated as previously described [59]. Precipitated proteins were solubilized in alkylating

buffer (0.1 M Tris pH 8.3, 1 mM EDTA, 1% SDS) for 10 min at 65°C. Crude TCA extracts were used in all

experiments. Prior to analysis, indicated amounts of the crude TCA extract were treated with 1x SDS-

loading buffer (Biorad) and boiled for 10 min at 95°C. Protein samples were run on a 7.5% TGX Mini

polyacrylamide gel (Biorad) using SDS running buffer (Biorad) for 35 to 40 min at 200 V. Then, proteins

were blotted on nitrocellulose using the Transblot Turbo system (Biorad) at the high molecular weight

blotting setting. Cuf1-FLAG protein levels were analyzed using a monoclonal Anti-FLAG M2-Peroxidase

antibody (1:5000 dilution) (Sigma, αFLAG-HRP) using a rabbit anti-β-tubulin at 1:500 dilution in 5% milk

+ TBS-T and 0.02% sodium azide (abcam, ab15568) combined with an anti-rabbit-HRP in a dilution of

1:4000 (GE Healthcare) as loading control. Blots were blocked with 5% Milk in TBS-T and washed 3x for

15 min with TBS-T after each antibody incubation. Primary antibodies were incubated overnight at 4°C,

the FLAG-HRP conjugate as well as secondary antibodies were incubated for 1 h at RT. Proteins were

detected by chemiluminescence using the SuperSignal West Pico substrate (Thermo Fisher Scientific).

RNA isolation and qRT-PCR

For transcript analysis during high versus low Cu stress, overnight cultures (~18h) incubated in

synthetic complete (SC) medium (MP Biomedicals) were diluted to OD_{600} 0.5, grown for 2 h at 30°C in SC

medium. Cu stress was then induced by the addition of either 1 mM BCS or 1mM $CuSO_4$ and cultures

were grown for 3 h at 30°C.

For transcript analysis in melanin inducing conditions, overnight cultures incubated in SC

medium were diluted to OD_{600} 0.5 and incubated for 3h at 30°C in L-DOPA medium supplemented with

either 10 µM BCS or 10 µM $CuSO_4$.

RNA extraction and cDNA analysis was performed as described previously [28]. For transcript

analysis via qRT-PCR, cDNA was diluted 1:5 in RNase-free water and mixed with the PowerUP SYBR

Green Master mix (Applied Biosystems) per protocol instruction and analyzed on a QuantStudio 6 Flex

(Applied Biosystems). The oligonucleotides used for transcript analysis are listed in **Table S3.** Cycle

threshold (C_T) values were calculated using the QuantStudio 6 Flex and gene expression values were

normalized to *GAPDH*. Expression fold changes were calculated using the $\Delta\Delta C_T$ method. For all transcript

analysis a minimum of 4 independent biological replicates were analyzed.

Chromatin immunoprecipitation followed by quantitative PCR analysis (ChIP-PCR)

Overnight cultures (~18h) incubated in SC medium were diluted to OD_{600} 0.2, and incubated for

2 h at 30°C in SC medium. Cu stress was then induced by the addition of either 1 mM BCS or 1mM $CuSO_4$

and cultures were grown for 2.5 h at 30°C. Crosslinking of the samples was performed in a 1%

formaldehyde solution for 30 min at RT. The downstream processing and ChIP were performed as

previously described [23] with following adaptions: Instead of sonication 5 times, with 30s ON followed

by 1 min OFF, samples were processed using a Bioruptor UCD200 (Diagenode) via 24 cycles of 15 s "ON "

followed by 15 s "OFF" at the high level to shear chromatin. Immune precipitated DNA fragments were

purified with a PCR clean up kit (Macherey Nagel) following manufacturer's instructions. qPCR was

performed with the PowerUP SYBR Green Master mix (Applied Biosystems) per protocol instructions

and analyzed on a QuantStudio 6 Flex (Applied Biosystems) using oligonucleotides listed in **Table S3**. For

quantification, the C_T from each input was subtracted from the C_T from the target gene obtained for the

Cuf1–FLAG strain ($\Delta C_{T\ target}$) or the negative control (WT strain, no FLAG-tagged Cuf1, $\Delta C_{T\ control}$). Then,

the $\Delta\, C_{T\ control}$ was subtracted from the $\Delta C_{T\ target}$ to give a $\Delta\Delta\, C_T$ value. For each gene promoter, Cuf1

enrichment was calculated as the inverse log of $\Delta\Delta\, C_T$ base 2 ($2^{\Delta\Delta Ct}$). Cuf1-FLAG enrichment at the

tubulin promoter was used for normalization.

Disc diffusion assay

To measure menadione-induced ROS stress sensitivity, SC overnight cultures (~18h) of indicated

Cn strains were diluted 1:500 in SC medium, and 500 μL cell suspension spread onto SC plates or SC

plates supplemented with indicated amounts of BCS or $CuSO_4$. Inoculated plates were dried for 45 min

at RT. Filter discs were spotted with 15 μL of the indicated concentrations of menadione (dissolved in

ethanol) and set in quadrants around the plates. Plates were incubated at 30 °C for 3 to 4 days.

Protein extraction and Sod activity assay

For Sod activity analysis, *Cn* overnight cultures (~18h) incubated in synthetic SC medium were diluted to

OD_{600} 0.1 in SC medium or SC medium supplemented with either 250 μM BCS or 50 μM or 100 μM

$CuSO_4$. Cultures were incubated for 24 h at 30°C. For protein extraction, cells were pelleted (1 min,

11000xg, 4°C), washed 1x with cold Lysis buffer (10 mM sodium phosphate, 50 mM NaCl, 0.1% triton X-

100, 10% glycerol pH 7.8 made in chelex-treated H_2O) and resuspended in 350 μL of cold lysis buffer + 1x

Protease Inhibitor cocktail (ROCHE) and PMSF. Cells were lysed with a Bead Ruptor 12 (Omni

international) using 6 cycles of 60s "ON" and 90s "OFF", 5.5 speed at 4°C. Cell lysates were clarified for

15 min at 21000xg at 4°C and the clarified lysate was transferred into a 1.5 mL reaction tube. Total

protein levels were measured using a BCA kit (Thermo Fisher Scientific) according to manufacturer's

instruction. Lysates were diluted to a concentration of 75 μg protein in 25 μL native sample buffer (Bio-

Rad, catalog #1610738) and lysis buffer to a final volume of 50 μL. Diluted lysates at a volume of 20 μL

were loaded onto a 10% mini-PROTEAN TGX precast protein gel (Bio-Rad, catalog #4561033).

Electrophoresis was performed in 1X tris glycine running buffer at 200 V and 4°C for approximately 1 h.

Gels were then incubated at RT in the dark for 1 h in 50 mL NBT staining solution (50 mM potassium

phosphate, 30 µM riboflavin, 0.48 mM nitro blue tetrazolium chloride, pH 7.8) while shaking at 60 rpm.

NBT staining solution was decanted and gels were soaked in diH$_2$O in the light for approximately 30 min

before imaging on an iBright 1500 FL Imager (Thermo Fisher Scientific).

Infection experiments

We used the murine inhalation model of cryptococcosis [59] to assess fungal survival and host

response after lung infection. For each strain, 5 female A/J mice were used. Mice were anesthetized by

isoflurane inhalation and intranasally inoculated with 1 x 10^5 fungal cells of PBS-washed overnight

cultures (~18h) of indicated strains in SC. Mice were sacrificed at 14 days post infection and the mouse

lung harvested, weighed, and homogenized in cold PBS. Colony forming units (CFU) were calculated by

quantitative culture and are represented as CFU/lung. For Histopathology at 14 days post infection, mice

were sacrificed using a lethal dose of pentobarbital sodium, and lungs were fixed via intratracheal

instillation of neutral buffered formalin under gravity flow [60]. Fixed lungs were submitted to the

Immunohistopathology Core at Duke University for paraffin embedding and staining with hematoxylin

and eosin. A pathologist assessed the distribution of the yeast cells and the immune response in a

blinded analysis.

Statistical analysis

All data error bars represent standard errors of the means (SEM) of results from a number of

biological replicates (N), as indicated in figure legends. Prior to statistical analysis, data from all

experiments was log$_2$ transformed for comparison of proportions. Statistical analysis was conducted

using the GraphPad Prism software v9. The statistical tests used in each experiment and their results

(i.e., p values) are indicated in figure legends. Asterisks in figures correspond to statistical significance as

follows: ****, $P < 0.0001$; ***, $P = 0.0001$ to $P < 0.001$; **, $P = 0.001$ to $P < 0.01$; *, $P = 0.01$ to $P < 0.05$;

ns (not significant), $P > 0.05$.

Acknowledgements

577 We thank the Duke Immunohistopathology Core for preparation and staining of the infected lung tissue.

578 We thank Dr. Connie Nichols for the careful reading of the manuscript. C.D.J. and K.J.F. acknowledge

579 support by the National Institute of General Medical Sciences of the National Institutes of Health (NIH)

580 under award number R01GM145035. This work was further supported by an NIH grant R01GM041840

581 (J.A.A. and C.P.) and by Duke University School of Medicine Bridge Funding 4532750.

FIGURE LEGENDS

Figure 1: The Cuf1-FLAG ΔAce1 mutant protein is produced to WT levels but fails to restore growth

during high Cu stress.

(A-B) Schematic representation of the *Cn*Cuf1 N- and C-terminal domains and the Cuf1-FLAG WT and

ΔAce1 mutant proteins. The first and last residues of each domain are indicated. The N-terminal domain

is shown in grey; the C-terminal domain is shown in pale yellow. The N-terminal DNA binding motif,

which contains the conserved KGRP DNA binding motif, is highlighted in orange. The Ace1-like cysteine-

rich region is indicated via blue box, and the Mac1-like regions are indicated by an orange box. The

sequence of the identified cysteine-rich regions is indicated for each region as well as (B) the cysteine-

to-alanine mutations performed to generate the ΔAce1 mutant protein. **(C)** Cuf1-FLAG WT and ΔAce1

mutant protein expression assessed by western blot analysis. The *CUF1-FLAG* WT or the *CUF1-FLAG*

ΔAce1 mutant strains were incubated in SC medium supplemented with either 1mM $CuSO_4$ or 1 mM BCS

at a starting OD_{600} of 0.5 for 3 h at 30°C. A TCA based protein extraction was performed. 25 µL of crude

TCA extract was analyzed per lane with the α FLAG-HRP conjugate. The α β-Tubulin (anti-β-tubulin)

antibody was used as a loading control. **(D)** Growth analysis in the presence of high and low Cu stress.

Five-fold serial dilutions of cell suspensions for each strain were spotted onto SC medium plates. To

induce low Cu stress, the SC medium was supplemented with 0.5 or 1 mM BCS. To induce varying

degrees of high Cu stress, the SC medium was supplemented with either 0.02, 0.05, 0.1, 0.25 or 1 mM

$CuSO_4$. Plates were incubated at 30°C for 3 d. Two independent *CUF1-FLAG ΔAce1* mutant strains were

analyzed. As Cu stress controls, the Cu import mutant *ctr1Δ ctr4Δ* and the Cu detoxification mutant

mt1Δ mt2Δ were included in the growth analysis.

Figure 2:   Mutation of the Ace1-like Cys-rich region renders Cuf1 unable to bind to and activate high

Cu regulated genes.

(A-B) Transcript abundance of the high Cu-regulated *MT1* and *CTR4* genes (A) and the low Cu-regulated

CTR4 and *CBI1* genes (B). Indicated strains were diluted to OD_{600} of 0.5 and incubated for 2 h at 30°C in

SC medium. Cu stress was induced by the addition of either 1 mM BCS or 1mM $CuSO_4$ and cultures were

incubated for 3 h at 30°C. Transcript abundance for each indicated gene was assessed by quantitative

real-time PCR (qPCR), and the $\log_2$ fold change of transcripts was calculated using the $\Delta\Delta\ C_T$ method in

high Cu stress versus low Cu stress (A) or low Cu stress versus high Cu stress (B). Transcript levels were

normalized to *GAPDH*. Shown is the mean +/- SEM of 4 biological replicates. Statistical analysis was

performed using one-way ANOVA. **(C-D)** Cuf1-FLAG promoter binding at high Cu stress (A) or low Cu

stress (B). Indicated strains were diluted to OD_{600} of 0.25 and incubated for 2 h at 30°C in SC medium. Cu

stress was induced by the addition of either 1 mM BCS or 1mM $CuSO_4$, and cultures were incubated for

2.5 h at 30°C prior to crosslinking. Promoter binding to the target promoters was analyzed via chromatin

immunoprecipitation followed by quantitative PCR (ChIP-PCR). Fold enrichment was calculated by

comparison of target promoter abundance in the FLAG-tagged strains to the promoter abundance in the

non-FLAG tagged, WT strain. The mean +/- SEM of 3 biological replicates is shown. A two-tailed t-test

was performed as statistical analysis.

Figure 3: Cuf1 Ace1-like Cys rich region is essential for *LAC1* regulation in response to excess copper.

(A) Schematic summary of the role of copper and Cuf1 for Lac1 activity and melanin production during

both Cu limitation and Cu excess. In the absence of Cuf1-induction of Cu-import genes during Cu

limitation, melanin production by the Cu-dependent laccase Lac1 is impaired. Cuf1 is also required for

expression of the *LAC1* gene during Cu excess. Images were designed using Biorender. **(B)** Melanization

during Cu limitation (10 µM BCS) and in presence of exogenous copper (10 µM $CuSO_4$). Indicated strains

were grown for 18h in SC at 30°C, diluted to OD_{600} of 2.5 and spotted onto L-DOPA agar plates with

indicated supplementation. L-DOPA plates were incubated for 3 d at 30°C. Melanin production is

demonstrated by colony pigmentation **(C-D)** *LAC1* (C) and *CTR4* (D) transcript abundance in melanin-

inducing conditions. Indicated strains were diluted to OD_{600} of 0.5 in liquid L-DOPA medium

supplemented with either 10 µM BCS or 10 µM $CuSO_4$ and grown for 3 h at 30°C. Relative transcript

abundance for *LAC1* (C) or *CTR4* (D) was assessed by quantitative RT-PCR and normalized to WT

transcript levels via the $\Delta\Delta C_T$ method. The data is presented as the mean +/- SEM of the Log_2 fold change

of 3 biological replicates. Statistical analysis was performed using a one-way ANOVA.

Figure 4: Cuf1 Ace1-like Cys rich region is required for superoxide dismutase (Sod)-dependent ROS detoxification during high Cu stress.

(A) Schematic summary of the role of copper and Cuf1 for Sod protein regulation during Cu limitation and excess. Cu, Zn-containing Sod1 is the major cytosolic Sod in Cu sufficiency and excess. The Mn-dependent mitochondrial Sod2 assumes the majority of cytosolic Sod function in Cu limitation. Images were designed using Biorender. **(B-C)** Disc diffusion assay to test cellular superoxide sensitivities. Indicated strains were inoculated on SC agar plates supplemented with indicated amounts of BCS or $CuSO_4$. Cells were challenged with increasing concentrations of the superoxide generator menadione on filter discs as outlined (B) for 3 days at 30 °C (C). **(D)** In-gel SOD activity assays from whole cell extracts of indicated strains. For Cu stress induction, strains were incubated in SC medium supplemented with indicated amounts of BCS or $CuSO_4$ for 24 h at 30°C. Whole cell extracts were generated, separated by native PAGE, and stained for SOD activity using nitroblue tetrazolium (NBT) staining solution. **(E)** Transcript abundance of the *SOD1* gene during high Cu and low Cu stress. Indicated strains were diluted to OD_{600} 0.5, and incubated for 2 h at 30°C in SC medium. Cu stress was induced by the addition of either 1 mM BCS or 1mM $CuSO_4$ and cultures were incubated for 3 h at 30°C. Relative transcript abundance for *SOD1* was assessed by quantitative RT-PCR and normalized to WT transcript levels using the $\Delta\Delta C_T$ method. The data is presented as the mean +/- SEM of the Log_2 fold change of 3 biological replicates and statistical analysis was performed using a one-way ANOVA.

Figure 5: Effects of Cuf1 high Cu stress regulation on survival and immune recognition in the infected lung.

(A-C) Lung fungal burden 14 days postinfection. Female A/J mice (5 mice per strain) were infected by

inhalation with 10^5 cells of the indicated strains. Mice were sacrificed 14 d postinfection and their lungs

harvested. Lung lysates were assessed by quantitative culture (A), lung wet weight (B) and macroscopic

lung abnormalities (C). (A-B) Data were plotted, and an unpaired t-test was performed using GraphPad

Prism (B only). (C) Foci are indicated with white arrows. **(D)** Histopathology analysis of infected murine

lungs 14 days post infection. Mice were infected as described above. Mice were sacrificed and lungs

were fixed via intratracheal instillation of neutral buffered formalin under gravity flow. Hematoxylin and

eosin staining were used to visualize microscopic lung pathology. Yeast enriched foci of inflammation

are highlighted by a black arrowhead. The length of the black bar represents 250 μm.

SUPPLEMENTAL

Supplemental Figure S1:

(A) *CDA2* and *CFO1* transcript abundance at high end low Cu stress. Indicated strains were diluted to

OD_{600} of 0.5 and, incubated for 2 h at 30°C in SC medium. Cu stress was induced by the addition of either

1 mM BCS or 1mM $CuSO_4$ and cultures were incubated for 3 h at 30°C. Relative transcript abundance

was assessed by quantitative RT-PCR and normalized to WT transcript levels via the $\Delta\Delta C_T$ method. The

data is presented as the mean +/- SEM of the Log_2 fold change of 4 biological replicates. A one-way

ANOVA was performed. **(B-C)** Cuf1-FLAG promoter binding at high Cu stress (B) or low Cu stress (C).

Indicated strains were diluted to OD_{600} of 0.25 and incubated for 2 h at 30°C in SC medium. Cu stress was

induced by the addition of either 1 mM BCS or 1mM $CuSO_4$, and cultures were incubated for 2.5 h at

30°C prior to crosslinking. Promoter binding was analyzed via chromatin immunoprecipitation followed

by quantitative PCR (ChIP-PCR). Fold enrichment was calculated by comparison of promoter abundance

in the Cuf1-FLAG strain to the promoter abundance in the WT strain. The presented data show the mean

+/- SEM of 3 biological replicates.

Supplemental Table S1: Strains used in this study.

Supplemental Table S2: Plasmids used in this study.

Supplemental Table S3: Oligonucleotides used in this study.

References

1.   Maret W. The Metals in the Biological Periodic System of the Elements: Concepts and

Conjectures. *International journal of molecular sciences*. 2016;17(1). Epub 2016/01/08. doi:

10.3390/ijms17010066.

2.   Smith AD, Logeman BL, Thiele DJ. Copper Acquisition and Utilization in Fungi. *Annual review of*

microbiology. 2017;71:597-623. Epub 2017/09/10. doi: 10.1146/annurev-micro-030117-020444.

3.   Halliwell B, Gutteridge JM. The importance of free radicals and catalytic metal ions in human

diseases. *Molecular aspects of medicine*. 1985;8(2):89-193. Epub 1985/01/01. doi: 10.1016/0098-

2997(85)90001-9.

4.   Giachino A, Waldron KJ. Copper tolerance in bacteria requires the activation of multiple

accessory pathways. 2020;114(3):377-90. doi: https://doi.org/10.1111/mmi.14522.

5.   Robison ATR, Sturrock GR, Zaengle-Barone JM, Wiebelhaus N, Dharani A, Williams IG, *et al.*

Analysis of copper-induced protein precipitation across the *E. coli* proteome. *Metallomics : integrated*

biometal science. 2023;15(1). Epub 2022/12/23. doi: 10.1093/mtomcs/mfac098.

6.   Zuily L, Lahrach N, Fassler R, Genest O, Faller P, Sénèque O, *et al.* Copper Induces Protein

Aggregation, a Toxic Process Compensated by Molecular Chaperones. *mBio*. 2022;13(2):e0325121. Epub

2022/03/16. doi: 10.1128/mbio.03251-21.

7.   Macomber L, Imlay JA. The iron-sulfur clusters of dehydratases are primary intracellular targets

of copper toxicity. *Proceedings of the National Academy of Sciences of the United States of America*.

2009;106(20):8344-9. Epub 2009/05/07. doi: 10.1073/pnas.0812808106.

8.   Imlay JA. The mismetallation of enzymes during oxidative stress. *The Journal of biological*

chemistry. 2014;289(41):28121-8. Epub 2014/08/28. doi: 10.1074/jbc.R114.588814.

9. Merchant SS, Schmollinger S, Strenkert D, Moseley JL, Blaby-Haas CE. From economy to luxury:

Copper homeostasis in *Chlamydomonas* and other algae. *Biochimica et biophysica acta Molecular cell*

research. 2020;1867(11):118822. Epub 2020/08/18. doi: 10.1016/j.bbamcr.2020.118822.

10. Li CX, Gleason JE, Zhang SX, Bruno VM, Cormack BP, Culotta VC. *Candida albicans* adapts to host

copper during infection by swapping metal cofactors for superoxide dismutase. *Proceedings of the*

National Academy of Sciences of the United States of America. 2015;112(38):E5336-42. Epub

2015/09/10. doi: 10.1073/pnas.1513447112.

11. Smith AD, Garcia-Santamarina S, Ralle M, Loiselle DR, Haystead TA, Thiele DJ. Transcription

factor-driven alternative localization of *Cryptococcus neoformans* superoxide dismutase. *The Journal of*

biological chemistry. 2021;296:100391. Epub 2021/02/11. doi: 10.1016/j.jbc.2021.100391.

12. Ray SC, Rappleye CA. Mac1-Dependent Copper Sensing Promotes *Histoplasma* Adaptation to the

Phagosome during Adaptive Immunity. 2022;13(2):e03773-21. doi: doi:10.1128/mbio.03773-21.

13. Mohamed SH, Fu MS, Hain S, Alselami A, Vanhoffelen E, Li Y, *et al.* Microglia are not protective

against *cryptococcal* meningitis. *Nature communications*. 2023;14(1):7202. Epub 2023/11/08. doi:

10.1038/s41467-023-43061-0.

14. García-Santamarina S, Thiele DJ. Copper at the Fungal Pathogen-Host Axis. *The Journal of*

biological chemistry. 2015;290(31):18945-53. Epub 2015/06/10. doi: 10.1074/jbc.R115.649129.

15. Murdoch CC, Skaar EP. Nutritional immunity: the battle for nutrient metals at the host–

pathogen interface. *Nature Reviews Microbiology*. 2022;20(11):657-70. doi: 10.1038/s41579-022-00745-

6.

16. Besold Angelique N, Gilston Benjamin A, Radin Jana N, Ramsoomair C, Culbertson Edward M, Li

Cissy X, *et al.* Role of Calprotectin in Withholding Zinc and Copper from *Candida albicans*. *Infection and*

Immunity. 2018;86(2):10.1128/iai.00779-17. doi: 10.1128/iai.00779-17.

17.   Garg R, David MS, Yang S, Culotta VC. Metals at the Host–Fungal Pathogen Battleground.

2024;78(Volume 78, 2024):23-38. doi: https://doi.org/10.1146/annurev-micro-041222-023745.

18.   White C, Lee J, Kambe T, Fritsche K, Petris MJ. A role for the ATP7A copper-transporting ATPase

in macrophage bactericidal activity. *The Journal of biological chemistry*. 2009;284(49):33949-56. Epub

2009/10/08. doi: 10.1074/jbc.M109.070201.

19.   Rajasingham R, Smith RM, Park BJ, Jarvis JN, Govender NP, Chiller TM, *et al.* Global burden of

disease of HIV-associated *cryptococcal* meningitis: an updated analysis. *The Lancet Infectious diseases*.

2017;17(8):873-81. Epub 2017/05/10. doi: 10.1016/s1473-3099(17)30243-8.

20.   Heitman J, Kozel TR, Kwon-Chung KJ, Perfect JR, Casadevall A. *Cryptococcus* : from human

pathogen   to   model   yeast.   *ASM   Press*;   2011.   Available   from:

http://www.asmscience.org/content/book/10.1128/9781555816858.

21.   Ding C, Festa RA, Chen YL, Espart A, Palacios Ò, Espín J, *et al. Cryptococcus neoformans* copper

detoxification machinery is critical for fungal virulence. *Cell host & microbe*. 2013;13(3):265-76. Epub

2013/03/19. doi: 10.1016/j.chom.2013.02.002.

22.   Sun TS, Ju X, Gao HL, Wang T, Thiele DJ, Li JY, *et al.* Reciprocal functions of *Cryptococcus*

neoformans copper homeostasis machinery during pulmonary infection and meningoencephalitis.

Nature communications. 2014;5:5550. Epub 2014/11/25. doi: 10.1038/ncomms6550.

23.   Garcia-Santamarina S, Festa RA, Smith AD, Yu CH, Probst C, Ding C, *et al.* Genome-wide analysis

of the regulation of Cu metabolism in *Cryptococcus neoformans*. *Molecular microbiology*.

2018;108(5):473-94. Epub 2018/04/03. doi: 10.1111/mmi.13960.

24.   Garcia-Santamarina S, Uzarska MA, Festa RA, Lill R, Thiele DJ. *Cryptococcus neoformans* Iron-

Sulfur Protein Biogenesis Machinery Is a Novel Layer of Protection against Cu Stress. *mBio*. 2017;8(5).

Epub 2017/11/02. doi: 10.1128/mBio.01742-17.

25.   Do E, Park S, Li MH, Wang JM, Ding C, Kronstad JW, *et al*. The mitochondrial ABC transporter

Atm1 plays a role in iron metabolism and virulence in the human fungal pathogen *Cryptococcus*

neoformans. *Medical mycology*. 2018;56(4):458-68. Epub 2018/02/09. doi: 10.1093/mmy/myx073.

26.   Cox GM, Harrison TS, McDade HC, Taborda CP, Heinrich G, Casadevall A, *et al.* Superoxide

dismutase influences the virulence of *Cryptococcus neoformans* by affecting growth within

macrophages. *Infect Immun*. 2003;71(1):173-80. Epub 2002/12/24. doi: 10.1128/iai.71.1.173-180.2003.

27.   Garcia-Santamarina S, Probst C, Festa RA, Ding C, Smith AD, Conklin SE, *et al.* A lytic

polysaccharide monooxygenase-like protein functions in fungal copper import and meningitis. *Nature*

Chemical Biology. 2020;16(3):337-44. doi: 10.1038/s41589-019-0437-9.

28.   Probst C, Garcia-Santamarina S, Brooks JT, Van Der Kloet I, Baars O, Ralle M, *et al.* Interactions

between copper homeostasis and the fungal cell wall affect copper stress resistance. *PLoS pathogens*.

2022;18(6):e1010195. Epub 2022/06/24. doi: 10.1371/journal.ppat.1010195.

29.   Thiele DJ. *ACE1* regulates expression of the *Saccharomyces cerevisiae* metallothionein gene.

Molecular and cellular biology. 1988;8(7):2745-52. Epub 1988/07/01. doi: 10.1128/mcb.8.7.2745-

2752.1988.

30.   Culotta VC, Hsu T, Hu S, Fürst P, Hamer D. Copper and the *ACE1* regulatory protein reversibly

induce yeast metallothionein gene transcription in a mouse extract. *Proceedings of the National*

Academy of Sciences of the United States of America. 1989;86(21):8377-81. Epub 1989/11/01. doi:

10.1073/pnas.86.21.8377.

31.   Jungmann J, Reins HA, Lee J, Romeo A, Hassett R, Kosman D, *et al. MAC1*, a nuclear regulatory

protein related to Cu-dependent transcription factors is involved in Cu/Fe utilization and stress

resistance in yeast. *The EMBO journal*. 1993;12(13):5051-6. Epub 1993/12/15. doi: 10.1002/j.1460-

2075.1993.tb06198.x.

32.   Waterman SR, Hacham M, Hu G, Zhu X, Park YD, Shin S, *et al.* Role of a CUF1/CTR4 copper

regulatory axis in the virulence of *Cryptococcus neoformans*. *The Journal of clinical investigation*.

2007;117(3):794-802. Epub 2007/02/10. doi: 10.1172/jci30006.

33.   Szczypka MS, Thiele DJ. A cysteine-rich nuclear protein activates yeast metallothionein gene

transcription. *Molecular and cellular biology*. 1989;9(2):421-9. Epub 1989/02/01. doi:

10.1128/mcb.9.2.421-429.1989.

34.   Graden JA, Posewitz MC, Simon JR, George GN, Pickering IJ, Winge DR. Presence of a copper(I)-

thiolate regulatory domain in the copper-activated transcription factor Amt1. *Biochemistry*.

1996;35(46):14583-9. Epub 1996/11/19. doi: 10.1021/bi961642v.

35.   Koch KA, Thiele DJ. Autoactivation by a *Candida glabrata* copper metalloregulatory transcription

factor requires critical minor groove interactions. *Molecular and cellular biology*. 1996;16(2):724-34.

Epub 1996/02/01. doi: 10.1128/mcb.16.2.724.

36.   Serpe M, Joshi A, Kosman DJ. Structure-function analysis of the protein-binding domains of

Mac1p, a copper-dependent transcriptional activator of copper uptake in *Saccharomyces cerevisiae*. *The*

Journal of biological chemistry. 1999;274(41):29211-9. Epub 1999/10/03. doi:

10.1074/jbc.274.41.29211.

37.   Fürst P, Hu S, Hackett R, Hamer D. Copper activates metallothionein gene transcription by

altering the conformation of a specific DNA binding protein. Cell. 1988;55(4):705-17. Epub 1988/11/18.

doi: 10.1016/0092-8674(88)90229-2.

38.   Upadhya R, Lam WC, Hole CR, Parchment D, Lee CK, Specht CA, *et al. Cryptococcus neoformans*

Cda1 and Cda2 coordinate deacetylation of chitin during infection to control fungal virulence. *Cell*

surface. 2021;7:100066. Epub 2021/10/30. doi: 10.1016/j.tcsw.2021.100066.

39.   Kim J, Cho Y-J, Do E, Choi J, Hu G, Cadieux B, *et al.* A defect in iron uptake enhances the

susceptibility of *Cryptococcus neoformans* to azole antifungal drugs. *Fungal Genetics and Biology*.

2012;49(11):955-66. doi: https://doi.org/10.1016/j.fgb.2012.08.006.

40.   Camacho E, Vij R, Chrissian C, Prados-Rosales R, Gil D, O'Meally RN, *et al.* The structural unit of

melanin in the cell wall of the fungal pathogen *Cryptococcus neoformans*. *The Journal of biological*

chemistry. 2019;294(27):10471-89. Epub 2019/05/24. doi: 10.1074/jbc.RA119.008684.

41.   Zhu X, Williamson PR. Role of laccase in the biology and virulence of *Cryptococcus neoformans*.

FEMS Yeast Research. 2004;5(1):1-10. doi: 10.1016/j.femsyr.2004.04.004 %J FEMS Yeast Research.

42.   Jiang N, Sun N, Xiao D, Pan J, Wang Y, Zhu X. A copper-responsive factor gene *CUF1* is required

for copper induction of laccase in *Cryptococcus neoformans*. *FEMS microbiology letters*. 2009;296(1):84-

90. Epub 2009/05/23. doi: 10.1111/j.1574-6968.2009.01619.x.

43.   Xue P, Hu G, Jung WH, Kronstad JW. Metals and the cell surface of *Cryptococcus neoformans*.

Current Opinion in Microbiology. 2023;74:102331. doi: https://doi.org/10.1016/j.mib.2023.102331.

44.   Telzrow CL, Esher Righi S, Castro-Lopez N, Campuzano A, Brooks JT, Carney JM, *et al.* An

Immunogenic and Slow-Growing *Cryptococcal* Strain Induces a Chronic Granulomatous Infection in

Murine Lungs. *Infect Immun*. 2022;90(6):e0058021. Epub 2022/05/20. doi: 10.1128/iai.00580-21.

45.   Cai Z, Du W, Zhang Z, Guan L, Zeng Q, Chai Y, *et al.* The transcription factor AceA is involved not

only in Cu but also in Zn detoxification through regulating transporters CrpA and ZrcA.

2018;20(10):e12864. doi: https://doi.org/10.1111/cmi.12864.

46.   Upadhyay S, Torres G, Lin X. Laccases involved in 1,8-dihydroxynaphthalene melanin

biosynthesis in *Aspergillus fumigatus* are regulated by developmental factors and copper homeostasis.

Eukaryotic cell. 2013;12(12):1641-52. Epub 2013/10/15. doi: 10.1128/ec.00217-13.

47.   Saikia S, Oliveira D, Hu G, Kronstad J. Role of ferric reductases in iron acquisition and virulence in

the fungal pathogen *Cryptococcus neoformans*. *Infect Immun*. 2014;82(2):839-50. Epub 2014/01/31. doi:

10.1128/iai.01357-13.

48.   Ray Stephanie C, Rappleye Chad A. Mac1-Dependent Copper Sensing Promotes *Histoplasma*

Adaptation to the Phagosome during Adaptive Immunity. *mBio*. 2022;13(2):e03773-21. doi:

10.1128/mbio.03773-21.

49.   Khemiri I, Tebbji F, Sellam A. Transcriptome Analysis Uncovers a Link Between Copper

Metabolism, and Both Fungal Fitness and Antifungal Sensitivity in the Opportunistic Yeast *Candida*

albicans. 2020;11. doi: 10.3389/fmicb.2020.00935.

50.   Cai Z, Du W, Zeng Q, Long N, Dai C, Lu L. Cu-sensing transcription factor Mac1 coordinates with

the Ctr transporter family to regulate Cu acquisition and virulence in *Aspergillus fumigatus*. *Fungal*

Genetics and Biology. 2017;107:31-43. doi: https://doi.org/10.1016/j.fgb.2017.08.003.

51.   Ballou ER, Wilson D. The roles of zinc and copper sensing in fungal pathogenesis. *Current*

Opinion in Microbiology. 2016;32:128-34. doi: https://doi.org/10.1016/j.mib.2016.05.013.

52.   Moraes D, Tristão GB, Rappleye CA, Ray SC, Ribeiro-Dias F, Gomes RS, *et al.* The influence of a

copper efflux pump in *Histoplasma capsulatum* virulence. *The FEBS journal*. 2024;291(4):744-60. Epub

2023/11/11. doi: 10.1111/febs.16999.

53.   Ost KS, Esher SK, Leopold Wager CM, Walker L, Wagener J, Munro C, *et al.* Rim Pathway-

Mediated Alterations in the Fungal Cell Wall Influence Immune Recognition and Inflammation. *mBio*.

2017;8(1). doi: https://doi.org/10.1128/mbio.02290-16.

54.   Esher SK, Ost KS, Kohlbrenner MA, Pianalto KM, Telzrow CL, Campuzano A, *et al.* Defects in

intracellular trafficking of fungal cell wall synthases lead to aberrant host immune recognition. *PLoS*

pathogens. 2018;14(6):e1007126. Epub 2018/06/05. doi: 10.1371/journal.ppat.1007126.

55. Hole CR, Lam WC, Upadhya R, Lodge JK. *Cryptococcus neoformans* Chitin Synthase 3 Plays a

Critical Role in Dampening Host Inflammatory Responses. 2020;11(1):10.1128/mbio.03373-19. doi:

doi:10.1128/mbio.03373-19.

56. Upadhya R, Lam WC, Hole CR, Vasselli JG, Lodge JK. Cell wall composition in *Cryptococcus*

neoformans is media dependent and alters host response, inducing protective immunity. *Frontiers in*

fungal biology. 2023;4. Epub 2023/08/04. doi: 10.3389/ffunb.2023.1183291.

57. Toffaletti DL, Rude TH, Johnston SA, Durack DT, Perfect JR. Gene transfer in *Cryptococcus*

neoformans by use of biolistic delivery of DNA. *Journal of Bacteriology*. 1993;175(5):1405-11. doi:

10.1128/jb.175.5.1405-1411.1993.

58. Upadhya R, Probst C, Alspaugh JA, Lodge JK. Measuring Stress Phenotypes in *Cryptococcus*

neoformans. *Methods in molecular biology*. 2024;2775:277-303. Epub 2024/05/17. doi: 10.1007/978-1-

0716-3722-7_19. PubMed PMID: 38758325.

59. Probst C, Hallas-Møller M, Ipsen J, Brooks JT, Andersen K, Haon M, *et al.* A fungal lytic

polysaccharide monooxygenase is required for cell wall integrity, thermotolerance, and virulence of the

fungal human pathogen *Cryptococcus neoformans*. *PLoS pathogens*. 2023;19(4):e1010946. Epub

2023/04/26. doi: 10.1371/journal.ppat.1010946.

60. Hsia CC, Hyde DM, Ochs M, Weibel ER. An official research policy statement of the American

Thoracic Society/European Respiratory Society: standards for quantitative assessment of lung structure.

American journal of respiratory and critical care medicine. 2010;181(4):394-418. Epub 2010/02/05. doi:

10.1164/rccm.200809-1522ST.

A

B

C

D

Figure 1

Figure 2

Figure 3

A

Cu Limitation

Cu Sufficiency

Cu Excess

B

[mM] Menadione in EtOH

C

cuf1Δ:: CUF1-FLAG

WT | *cuf1Δ* | WT | ΔAce1 #1 | ΔAce1 #2

100 μM BCS

-

75 μM CuSO₄

100 μM CuSO₄

D

cuf1Δ :: CUF1-FLAG

ΔAce1

WT | *cuf1Δ* | WT | #1 | #2 | *sod1Δ* | *sod2Δ*

250 μM BCS

50 μM CuSO₄

100 μM CuSO₄

(*) Sod1 protein activity
(#) cytosolic Sod2 protein activity

E

$SOD1$ Log₂ fold change WT=1

SC + 1 mM CuSO₄
SC + 1 mM BCS

WT | *cuf1Δ* | WT | #1 | #2 | WT | *cuf1Δ* | WT | #1 | #2
ΔAce1 | ΔAce1
CUF1-FLAG | *CUF1-FLAG*

Figure 4

Figure 5