Identification of novel plasma proteomic biomarkers of Dupuytren Disease

Full title: Identification of novel plasma proteomic biomarkers of Dupuytren

Disease

Short title: Dupuytren plasma proteomic biomarkers

Blake Hummer[1¶], Paola Sebastiani[2,3&], Anastasia Leshchyk[4&], Anastasia Gurinovich[2,3&], Cecilie

Bager[5&], Morten Karsdal[5&], Signe Nielsen[5&], Charles Eaton[1¶*]

[1] Dupuytren Research Group, West Palm Beach, Florida, United States of America

[2] Institute for Clinical Research and Health Policy Studies, Tufts Medical Center, Boston,

Massachusetts, United States of America

[3] Department of Medicine, Tufts University, Boston, Massachusetts, United States of America

[4] Department of Medicine, Computational Biomedicine Section, Chobanian & Avedisian School

of Medicine, Boston University, Boston, Massachusetts, United States of America

[5] Nordic Bioscience, Herlev, Denmark

* Corresponding Author

Email: c.eaton.md@dupuytrens.org (CE)

[¶] These authors contributed equally to this work.

[&] These authors also contributed equally to this work.

Abstract

Dupuytren Disease (DD) is a chronic progressive disease that can cause disabling hand

deformities. The most common treatments have either high complication rates or high early

recurrence rates. Dupuytren lacks a staging biomarker profile to inform the development of

preventive therapeutics to improve long-term outcomes. This multi-omic study aimed to create a

DD blood proteomic biomarker profile by comparing DD plasma with that of a healthy control

group. We measured circulating collagen metabolism peptides and found normal Collagen I

synthesis but impaired Collagen I degradation in DD. We measured 6995 serum protein

aptamers and identified 68 proteins that showed statistically significant differences compared

with the control group. We developed two Diagnostic Proteomic Risk Scores (DPRS) based on

hypothesis-free and hypothesis-based analyses. In independent data, our hypothesis-free and

hypothesis-based DPRS distinguished Dupuytren from control subjects with accuracies of

76.5% and 70.6%, respectively. Our hypothesis-based DPRS also distinguished DD subjects

with different disease progression rates by age at their first corrective procedure (p=0.0018).

This pilot study is the first to provide evidence to suggest that Collagen I accumulation in DD

results from impaired degradation rather than increased collagen synthesis. It also describes

novel DPRS that have potential use as diagnostic and staging biomarker panels for Dupuytren

disease.

Introduction

Dupuytren disease (DD) is a progressive fibrotic condition that often results in finger deformities

called Dupuytren contracture (DC). DD affects an estimated 10 million Americans. DC is

associated with loss of productivity, higher healthcare costs, and a greater prevalence of benign

diseases [1].

The greatest need in DD care is preventing DC progression and recurrence. As with other

fibrotic diseases, developing preventive treatments requires a biomarker profile for staging and

measuring therapeutic response. The authors have not found published reports with statistically

significant evidence of preventing DC recurrence or a biomarker tool to develop such

treatments. This pilot study aimed to develop preliminary data on a diagnostic blood biomarker

profile of DD.

DC is a common outcome of DD. Of those diagnosed with DD at age 65, one in four men and

one in eight women will require at least one corrective procedure over their lifetime [2]. The

most common current treatments are palliative, often resulting in partial or temporary

improvement [3]. Surgery for DC has a greater risk of major complications than other common

elective hand operations [4-7], and recurrent DC is the most common diagnosis leading to

elective finger amputation [8].

Genetic factors account for about 80% of the risk of developing DD [9]. Family history, age, and

gender are strongly associated. A genetic risk score (GRS) based on 11 SNPs correlates with

DD diagnosis [10], and a weighted GRS based on 26 SNPs correlates with post-procedure

recurrence [11].

DD affects the palm skin, subcutaneous tissues, and fascia [12]. Early disease is cellular, with

fibroblasts, myofibroblasts, and perivascular clusters of myofibroblasts, pericytes, and

mononuclear immune cells [13]. Early changes include local variations of immune [13],

hematopoietic, and mesenchymal cell markers [12], cytokines, signaling factors [14], and

extracellular matrix components [15, 16]. Progression transforms tissues into nearly acellular,

densely packed collagen.

Physical changes are the basis for DD diagnosis, staging, and treatment, but physical findings

do not predict progression [17]. There are no standard noninvasive tests of DD biological

activity. Biopsy is problematic because it can trigger disease activity and progression. Blood

remains the most promising source for a disease activity profile.

Materials and methods

Study Approval

The Institutional Review Board Ascension Via Christi Hospitals, Wichita, Inc., approved this

study and the associated informed consent form. Written informed consent was obtained from

all participants before their participation. This written informed consent included a description of

the use of the photographs/ images to be reviewed for study entry. The records of these signed

informed consents are held on file by the investigators in accordance with PLOS policies. All

samples remained blinded throughout the processing and data collection in the laboratories. All

methods were performed in accordance with relevant guidelines, regulations, and the

Declaration of Helsinki.

Subjects

We selected disease cohort subjects from the International Dupuytren Database (IDDB), a

patient-reported registry developed by the Dupuytren Research Group (DRG) in collaboration

with the National Databank of Rheumatic Diseases (NDB). Control subjects were enrolled by

word of mouth and from the NDB database (https://dupuytrens.org/enroll-in-the- iddb-2/). Study

candidate subjects were identified on December 3, 2019, using the entry criteria described

below. We excluded minors from the study. Our study involved male and female subjects. Sex

was considered a biological variable. While we controlled for sex differences, we did not stratify

by sex due to limited sample sizes. The authors had no access to information that could identify

individual participants during or after data collection. All subjects met the entry criteria, and each

subject submitted a signed physical Informed Consent Form by postal mail. Recruitment started

on December 13, 2019, and ended on May 3, 2021.

92   Entry criteria

The DD cohort met criteria based on clinical evidence of DD with strong traditional predictors of

recurrent deformity after a corrective procedure: age of disease onset younger than 50; two or

more first-degree relatives with DD; current DD nodules; three or more separate locations of

DD, or two palm locations and Ledderhose disease (a related disorder). Control cohort subjects

were healthy individuals with no clinical evidence of DD, no personal history of DD or

Ledderhose disease, and no family history of DD.

We confirmed study eligibility by requiring candidates to upload recent images of the palmar and

dorsal sides of each hand. The Principal Investigator/senior author confirmed the entry criteria

and reviewed images to confirm clinical evidence of active DD in the disease cohort and the

absence of these findings in the control cohort.

103   Blood Sample Collection

The study used a mobile phlebotomy service (Phlebotek Solutions Corporation, Fort

Lauderdale, FL) to collect blood specimens. Once confirmed that the subject met all study entry

criteria, they were contacted by DRG and later by Phlebotek to schedule their blood draw. The

phlebotomist visited the subjects in their homes and collected two tubes of blood. The last 16

blood draws included a third 2.5ml PAXgene tube collection for additional genomic testing.

Processing of the EDTA tubes for proteomic testing was standardized. Immediately following

collection, blood samples were centrifuged for 15 minutes at 2000 g to generate plasma. The

plasma was harvested from the top of the container and pooled before aliquoting into storage

cryotubes. The last 500 µL of plasma was discarded to avoid contamination with platelets.

Samples were immediately packaged for shipping on dry ice.

Number of subjects

One hundred subjects were planned (50 DD/50 Control). This number was a practical

consideration, given the limited number of potential participants enrolled in the International

Dupuytren Data Bank who met all disease cohort entry criteria. Blood collection launched in

February 2020, just as the COVID pandemic began. Due to COVID-related issues, we amended

the protocol to enroll 50 subjects. Forty-seven subjects were enrolled;46 subjects had samples

drawn. One sample was disqualified due to failed shipping refrigeration. Mass spectrometry

(Bioproximity, Manassas, VA) was performed on samples from the remaining 45 subjects (25

DD, 20 Control). Twenty-seven subjects (13 DD, 14 control) had second blood samples drawn 6

to 14 months after their first draw for mass spectrometry testing by a second lab (MacCoss Lab,

Seattle, WA). Additional proteomic testing was conducted on remaining aliquots: 1) n= 27

redrawn samples had ELISA tests for collagen metabolites (Nordic Bioscience, Herlev, DK); 2)

n= 72 from both initial and redraw specimens had Aptamer-based testing (SomaLogic, Boulder,

CO). S1 Fig outlines the distribution of samples and laboratories.

Blinding

129 All subjects were assigned and identified by participant numbers. Investigators were blinded to

130 the sample group allocation. All samples remained blinded throughout the processing and data

131 collection in the laboratories.

Assays

133 No laboratory protocols were conducted in-house.

Nordic Bioscience collagen markers

Circulating collagen synthesis and degradation markers were measured using proprietary

assays [18]. Synthesis biomarkers included collagens I (PRO-C1), II (PRO-C2), III (PRO-C3), IV

(PRO-C4), V (PRO-C5), VI (PRO-C6), VII (PRO-C7), and IX (PRO-C9). Circulating degradation

markers were measured for collagens I (C1M), II (CALC2, C2M), III (C3M, CTX-III), IV (C4M), VI

(C6M), and VII (C7M). All biomarkers were assessed by competitive ELISAs and validated to

measure in human plasma samples. The inter- and intra-assay coefficients of variation were

>15% and 10%, respectively, for all assays.

Mass spectrometry

A novel untargeted proteomics assay using large-scale liquid chromatography-tandem mass

spectrometry (LC-MS/MS) with data-independent acquisition (DIA) was used to collect

proteomic data. Each sample (10 µl plasma), following tryptic digestion, was diluted 50:50 by

the addition of 4% sodium dodecyl sulfate (SDS) buffer to a final concentration of ~2% SDS.

Approximately 200 µg of protein was added to 96-well plates, and the remainder of the sample

preparation was performed using the single-pot, solid-phase enhanced sample preparation for

proteomics methodology (SP3) [19, 20]. The entire process of protein capture, reduction and

alkylation, protein cleanup, digestion, and peptide cleanup was performed using a KingFisher

magnetic bead robot [20]. Peptides were reconstituted in HPLC starting aqueous buffer (5%

acetonitrile and 0.1% formic acid) with analysis performed on the high-resolution quadrupole-

orbitrap hybrid mass spectrometer (Exploris 480, Orbitrap Fusion Lumos, or Orbitrap Fusion

Eclipse).

Data was acquired in a single, 2-hour LC-MS/MS run (90 min HPLC gradient) on a Thermo

Quadrupole-Orbitrap mass spectrometer. Crucially, the DIA method comprehensively sampled

all peptides between 400 and 1000 m/z using a repeated cycle of 75 MS/MS scans with an 8

m/z-wide isolation width. Using a multiplexing scheme based on staggering the isolation window

in alternating cycles of MS/MS scans, the precursor selectivity of the resulting data was 4 m/z

after computational demultiplexing [21]. From the resulting data, fragment ion chromatograms

were extracted for any peptide precursor between 400 and 1000 m/z. To determine which

peptides were confidently detected in each acquisition, the software tool EncyclopeDIA [22] was

used to query the data using a reference chromatogram library generated from a prior analysis

of the tissue. Chromatogram-based relative quantitation was then performed for each peptide

detected. All measurements were made relative to a common reference. In the case of plasma,

a common reference was a pooled standard from 400 age-matched samples from subjects who

were both healthy and spanning the phenotypes being studied, processed in each sample

batch. Within each 96-well plate, one copy of the common reference standard was placed in

each row for a total of 8 copies per plate [23]. Before quantification, it was confirmed that

peptides were stable during the time scale of the data collection, had a suitable linear response,

and had suitable intra- and inter-data precision.

SomaScan

SomaLogic, Inc. (Boulder, CO, USA) performed proteomic analysis of the plasma samples using

the SomaScan platform, which quantifies the presence of 6995 human proteins (secreted

proteins, extracellular domains, and intracellular proteins). These proteins are found in diverse

biological groups such as receptors, kinases, cytokines, proteolytic enzymes and inhibitors,

growth factors, protease inhibitors, hormones, and structural proteins [24]. Most of these

proteins are involved in pathways related to signal transduction, stress response, immune

processes, phosphorylation, proteolysis, cell adhesion, cell differentiation, and intracellular

transport [25].

Proteins were measured using the Slow Off-Rate Modified Aptamer (SOMAmer)-based capture

array, which uses chemically modified single-stranded DNA sequences capable of uniquely

recognizing individual proteins via one or more binding sites. SOMAmers transform a protein

signal to a nucleotide signal that is quantified using relative fluorescence on microarrays. Each

SOMAmer array underwent validation for specificity, upper and lower limits of detection, and

both intra- and inter-assay variability. Plasma dilutions (0.005%, 0.05%, and 20%) were applied

to capture low-, medium-, and high-abundance proteins. Positive and negative controls were

also positioned on the array to confirm if the experimental procedure was performed correctly.

After the hybridization step, the microarrays were washed and scanned using a laser scanner

that excites the fluorescence of the fluorochrome used in the labeling step. The amount of the

emitted signal is directly proportional to the amount of dye on the microarray. The scanner

measured this quantity and created a digital image map of the position of each signal based on

the location of its origin. Image analysis software generated a text file from this map describing

the pixel intensity of the spot and the background. This information was processed, and a final

value was generated that summarized the level of each detectable protein on the microarray.

Quality control analysis of the sample and SOMAmer levels involved SOMAmer control arrays

on the microarray and calibration samples. Hybridization controls measured the sample-by-

sample variation in hybridization, while the median signal over all SOMAmer arrays measured

the technical variability. The SomaScan measures were reported as relative fluorescence units

(RFU) in a summary ADAT file.

Analysis and statistics

Nordic collagen marker analysis

203   Nordic collagen data turnover rates were calculated by performing ratios of the propeptide over

204   the mature peptide (e.g., PRO-C1/ C1M). This ratio is an estimate of the formation/degradation

205   of each collagen type and can be interpreted as a net gain of collagen/loss of collagen

206   remodeling [18]. All comparisons were done using Student's T-test.

SomaScan analysis

A series of quality control and data cleaning steps was performed on the SomaScan proteomics

data. All non-human proteins were excluded from the analyses. Samples that did not pass the

default normalization acceptance criteria for all row scale factors were removed. All aptamers

that did not pass the default quality control criteria across all plates/sets were filtered out. The

data had no missing values, and the protein values were log-transformed. No further

normalization was performed.

We identified differentially expressed aptamers using linear regression of each log-transformed

protein as the outcome and disease status as the main predictor. We adjusted each analysis by

age and sex. We analyzed all repeated measurements of each individual and employed

generalized estimating equations to account for technical and biological variability using the gee

package in R. Due to the high number of aptamers tested in the SomaScan assay (6995), we

were at a statistical disadvantage when applying multiple corrections. Therefore, we calculated

adjusted p-values using Benjamini-Hochberg methods. Our SomaScan analyses report only

proteins with statistically significant differences from the control at an FDR-adjusted p-value <

0.25.

Mass spectrometry analysis

224   For the MS protein data, all zeroes were replaced with NAs across samples, and protein values

225   were log-transformed. The MS dataset was subset based on proteins that overlapped with the

226   signature proteins associated with DD previously identified using the SomaScan proteomics

227   data. We used a generalized estimation equation model to account for the repeated blood

228   draws, adjusted by age and gender.

Proteomic Risk Score calculations

To prepare the data for the Diagnostic Proteomic Risk Score (DPRS) calculation, we first

removed all highly correlated proteins (Pearson correlation > 0.8). 3369 out of 6995 aptamers

were highly correlated and were excluded from the DPRS calculation. The DPRS calculation

contained the following two steps. First, we fit a logistic regression for each protein, with disease

status (i.e., case or control) as the outcome variable, adjusted for age and gender. We used a

generalized estimation equation model for repeated measurements of two independent blood

draws. We included the same subject's repeated measurements and used the generalized

estimation equations model with the gee R package [26] to estimate the regression coefficients.

Next, the DPRS was computed using summary statistics from each regression model. Only the

proteins with statistically significant differences from control at an FDR-adjusted p-value < 0.25

were included in the score calculation. The DPRS was calculated by summing the beta

estimated effect for all significant proteins, each multiplied by normalized protein abundance

and divided by its standard error. Finally, we fit a logistic regression to examine the association

of the computed DPRS with DD onset. We split the dataset into training (75%) and testing sets

utilizing the caret R package for a leave-one-out cross-validation approach for model fitting and

assessment (25%) sets [27]. We used R version 4.2.1 for all analyses.

Pathway analysis

STRING (12.0) is a web tool that provides and illustrates protein-protein interactions and

networks [28]. STRING integrates evidence from public databases, including genomic context,

high-throughput and biochemical interaction experiments, known protein complexes,

computational predictions, and text mining from scientific literature. Given a list of gene names,

STRING constructs protein-protein interaction networks in which each gene is a node, with

edges representing predicted functional associations.

Networks were generated using the whole genome as the statistical background, and interaction

sources included text mining, experiments, databases, co-expression, neighborhood, gene

fusions, and co-occurrence. Interaction scores were computed by combining the probabilities

across all evidence channels and correcting for the probability of observing an interaction by

chance. All nodes and interactions were computed with a required medium confidence

interaction score (0.400) and an FDR<=0.05.

Results

Plasma collagen metabolites in DD

Collagen synthesis and degradation processes release protein fragments specific to the

process, collagen type, and enzyme involved. The ratio of synthesis to degradation-related

fragments correlates with net accumulation or loss of that collagen type. We evaluated plasma

markers of collagen synthesis and degradation using a competitive ELISA (Nordic Biosciences;

Herlev, Denmark) and calculated collagen synthesis-to-degradation ratios [18].

We measured markers of types I, II, III, IV, V, VI, VII, and IX collagen metabolism in blood from a

cohort of healthy controls (n = 14) and subjects with DD (n = 13). We found significantly lower

levels of markers of Collagen I degradation (C1M; Fig 1A, p = 0.004), Collagen V synthesis

(Pro-C5; Fig 1B, p = 0.046), and Collagen VII synthesis (Pro-C7; Fig 1C, p = 0.048) in the DD

cohort compared to the control group. The remaining collagen biomarkers did not show

significant differences between the disease and control groups (Table 1).

Fig 1. Differential production of Dupuytren vs. control collagen metabolism markers. A)

Collagen I degradation marker C1M. B) Collagen I synthesis to degradation marker ratio Pro-

C1/C1M. C) Collagen V synthesis marker Pro-C5. D) Collagen VII synthesis marker Pro-C7. * p-

value <0.05, **p-value <0.005.

Table 1. Collagen epitope markers and findings.

Assay	D/C	Description	Represents	p-value	SE	FC (D/C)
PRO_C1_HP	ns	Internal epitope in the N-terminal propeptide of type I collagen	Col I Synthesis	0.9437	C 10.99 D 11.89	0.93
C1M_HP	Down**	MMP-generated fragment of the a1 chain of type I collagen	Col I Degradation	0.0036	C 5.25 D 4.30	0.73
PRO-C2_HP	ns	N-terminal fragment of type IIB procollagen	Col II Synthesis	0.7808	C 1.45 D 3.11	1.01
CALC2	ND	C-terminal propeptide of type II collagen	Col II Synthesis	NA	NA	NA
C2M_HP	ns	MMP-generated fragment of type II collagen	Col II Degradation	0.1607	C 1.52 D 1.16	1.29
PRO-	ns	N-terminal propeptide of type III	Col III	0.9951	C 1.04	0.93

C3_roHP		collagen	Synthesis		D 1.24	
C3M_HP	ns	MMP-9 generated fragments of type III collagen	Col III Degradation	0.1615	C 16.96 D 13.61	0.86
CTX-III_HP	ns	MMP-generated fragment of crosslinked type III collagen	Col III Degradation	0.4887	C 40.88 D 53.26	0.91
PRO-C4_HP	ns	Internal epitope in the 7S domain of type IV collagen	Col IV Synthesis	0.0739	C 195.98 D 183.12	0.84
C4M_HP	ns	MMP-generated fragment of the a1 chain of type IV collagen	Col IV Degradation	0.1014	C 8.89 D 7.42	0.84
PRO-C5_HP	Down*	C-terminal propeptide of type V collagen	Col V Synthesis	0.0458	C 33.17 D 35.88	0.82
PRO-C6_roHP	ns	C-terminal propeptide of type VI collagen alpha-3 chain	Col VI Synthesis	0.7222	C 0.27 D 0.27	0.92
C6M_HP	ns	MMP2-generated fragment of type VI collagen	Col VI Degradation	0.1097	C 22.31 D 23.41	0.83
PRO_C7 HP	Down*	BMP-I mediated neo-epitope in the NC-2 propeptide of type VII collagen	Col VII Synthesis	0.0480	C 2.49 D 1.08	0.72
C7M	ND	Neo-epitope of MMP13 degradation of type VII collagen	Col VII Degradation	NA	NA	NA
PRO-C9	ns	C-terminus of NC-I domain of the a1 chain of type IX collagen	Col IX Synthesis	0.9658	C 0.80 D 0.65	0.92

Table 1. Synthesis and degradation markers of Collagen I, II, III, IV, V, VII, and IX.

Compared to controls, there were significant reductions in markers of Collagen I degradation,

Collagen V synthesis, and Collagen VII synthesis. D/C: ratio of Dupuytren to control values. FC:

fold change. **: <0.005, *: <0.05. ns: not significant. ND: undetectable in both disease and

control cohorts.

We calculated collagen synthesis-to-degradation ratios for collagen types I, II, III, IV, and V. The

Collagen I synthesis-to-degradation ratio, Pro-C1/C1M, was significantly greater in the DD

cohort than in the control group (Fig 1D, p = 0.024). The remaining ratios were not significant

(Col II, III, IV, VI) or could not be calculated because the degradation marker was either

unavailable (Col V, IX) or not detectable in both disease and control cohorts (Col VII).

Differentially expressed blood proteins in DD

We used two orthogonal, unbiased proteomic methods: the aptamer-based SomaScan

(SomaLogic; Boulder, CO) and liquid chromatography-tandem mass spectrometry (LC-MS/MS)

(MacCoss Lab, University of Washington, WA, USA).

We used sixty-nine samples for SomaScan testing after quality control (36 DD and 33 controls).

These represent 42 subjects, 27 of whom had second samples collected 6 to 14 months after

the first (S1 Fig). The SomaScan panel measures 6995 protein binding sites applying to 5444

proteins.

Hypothesis-free approach

The 6995 aptamer measurements identified 54 proteins with statistically significant differences

between the DD and control groups, with 24 overexpressed and 30 underexpressed after

multiple-comparison adjustment (S1 Table). Fig 2 is a heatmap of these 54 protein expressions.

Fig 2. Heatmap of the 54 significant proteins in the hypothesis-free analysis. This heatmap

shows natural log-transformed and standardized relative fluorescence units (RFU) for proteins

(rows) across samples (columns). Red colors represent higher-than-average RFU values, and

blue represents lower-than-average RFU values. Rows and/or columns are organized using

hierarchical clustering, and cohorts are indicated by the top bar [purple: Dupuytren; green:

Control]. We performed pathway analysis of these 54 proteins to identify network relationships

using STRING [28]. We found significant protein-protein interactions (PPI) in 29, with multiple

enriched categories (Fig 3A, p-value = 4.55E-04). S2 Table summarizes categories of enriched

gene groups, and Fig S2 provides a visual cross-reference of functional categories and

individual genes. The dominant categories were coagulation and complement cascades, the

extracellular matrix, enzyme inhibition, cell membrane structures, phosphorylation, and signaling

pathways.

Fig 3. Hypothesis-Free DPRS and Protein-Protein Interactions. A) Network expression of 54

differentially expressed proteins. Of the 54 proteins, 30 had strong pathway connections in the

network: ACAN, AKR1A1, C5, C6, CALCB, CASP3, CGA, CGB3, CPB1, CRKL, DSG2, EIF4H,

FKBP5, G3BP2, GP1BB, GUSB, HGS, HSP90AA1, KNG1, MAP3K11, MBNL1, MBNL2, NRG4,

OSCAR, PCSK9, POSTN, PRSS1, SERPINC1, SOCS3, SYK, TF, USP8, and YWHAB. B) The

Hypothesis-free Diagnostic Proteomic Risk Score has an AUC-ROC value of 0.8. C)

Hypothesis-free Diagnostic Proteomic Risk Scores (HF-DPRS) of the control vs Dupuytren

cohorts. p-value = 5.28E-06.

Using SomaScan results, we aimed to develop a hypothesis-free indicator of DD diagnosis, the

Diagnostic Proteomic Risk Score (DPRS), to estimate an individual's disease risk based on

plasma protein abundance.

Hypothesis-free DPRS

We developed a hypothesis-free DPRS estimation based on two of the 54 proteins. These

proteins, OSCAR (overexpressed) and SPART (underexpressed), had a low relative correlation

(Pearson's correlation coefficient = -0.19) (Table 2), suggesting minimal interaction. Using this

hypothesis-free DPRS as a diagnostic measure for DD achieved 79% accuracy in the training

set (AUC 0.8) and 76.5% in the test set (Fig 3B), indicating predictive value for discriminating

between disease and control. Fig 3C shows a box plot of control and DD DPRS. The DPRS for

controls and DD differed significantly by the Wilcoxon test (p-value = 2.5E-06).

Table 2. Hypothesis-free Diagnostic Proteomic Risk Score.

Protein	Eff	SE	p-value	FDR
Spartin	-11.62125	2.97479	0.000094	0.18
Oscar	13.99037	3.596067	0.0001	0.18

333   **Table 2. Proteins of the Hypothesis-free DPRS**. Eff = log (odds ratio comparing Dup to

334   control) for a change of 1 in the log (protein data). SE: Standard error. FDR: False Discovery

335   Rate (Adjusted p-value).

Hypothesis-based approach

We also used a hypothesis-based approach, identifying 328 relevant proteins from a literature

search and available on the SomaScan platform (S3 Table). Twenty-three proteins in this subset

showed statistically significant differences between DD and control, with nine overexpressed

and 14 underexpressed after multiple-comparison adjustment (S4 Table).

We constructed PPI networks and performed enrichment analysis of these 23 proteins using

STRING. We found significantly enriched PPI in 21 proteins with multiple enriched categories

(Fig 4A, p-value = 1.35E-07). We found more enrichment categories and greater significance in

the hypothesis-based than the hypothesis-free analysis, which may reflect a higher-yield group,

selection bias, or both. S5 Table lists enriched gene categories, and Fig S3 provides a visual

cross-reference between functional categories and individual genes.

Fig 4. Hypothesis-based DPRS and Protein-protein interactions. A) Network expression of

23 differentially expressed proteins. Of the 23 proteins, 20 had strong pathway connections:

DAB2, SMAD1, USP8, PLAT, SFRP4, ACAN, DDR2, HPX, KNG1, LCN2, SERPINH1, TF,

MAP2K1, PRKCA, STAT1, YWHAZ, POSTN, COL1A1, CASP3, and STAT3. B) The Hypothesis-

based Diagnostic Proteomic Risk Score has an AUC-ROC value of 0.9. AUC-ROC (Area Under

the Curve - Receiver Operating Characteristics) is an index of predictive accuracy. C)

Hypothesis-free Diagnostic Proteomic Risk Scores (HF-DPRS) of the control vs Dupuytren

cohorts. (p-value = 7.79E-09).

Hypothesis-based DPRS

We developed a hypothesis-based DPRS using 11 of 23 DE proteins identified in the

hypothesis-based analysis. Six genes were underexpressed (*SMAD1, SH3BP2, LCN2, USP8,*

CSNK1G2, SERPINH1) and five overexpressed (*CSMD1, POSTN, ACAN, AOC3, DDR2*). Six

proteins were present in hypothesis-based and hypothesis-free DPRS analyses (Table 3). Our

hypothesis-based DPRS achieved a training accuracy of 78.85% (AUC 0.9) and a test accuracy

of 70.59% in distinguishing Dupuytren from control cohorts (Fig 4B). Fig 4C shows that DPRS

distributions of Control and DD groups differ significantly (Wilcoxon test, p-value = 2.2E-09).

Table 3. Hypothesis-based Diagnostic Proteomic Risk Score.

Protein	Eff	SE	p-value	FDR
SMAD1*	-6.3649	1.877804	0.0007	0.12
CSMD1	5.397535	1.907423	0.00466	0.18
SH3BP2	-4.59151	1.582341	0.00371	0.18
POSTN*	4.305245	1.502627	0.00417	0.18
ACAN*	5.589134	1.75746	0.00147	0.12

LCN2	-4.73895	1.734879	0.0063	0.19
AOC3*	7.050175	2.62144	0.00716	0.2
DDR2	1.63418	0.595212	0.00604	0.19
USP8*	-3.30095	1.041269	0.00152	0.12
CSNK1G2*	-10.3331	3.593633	0.00404	0.18
SERPINH1	-2.6813	0.799538	0.0008	0.12

365 **Table 3. Proteins of the Hypothesis-based DPRS**. Eff = log (odds ratio comparing Dup to

366 control) for a change of 1 in the log (protein data). SE: Standard error. FDR: False Discovery

367 Rate (Adjusted p-value). * Six also had significant differential expression in the 54 proteins

368 identified in the hypothesis-free analysis.

Evaluation of DRPS for prognosis

We explored the potential of each DPRS to profile DD prognosis. We used age at the time of

the first DC procedure as an index of biological severity. We segregated our DD samples into

subjects younger (11 specimens) or older (16 specimens) than age 50 at the time of their first

DC procedure. We excluded from this analysis subjects younger than 50 at the time of their

blood draw. A one-way ANOVA revealed a statistically significant difference in the means of the

hypothesis-based DPRS across groups with different rates of pretreatment contracture

progression (Fig S4; p-value = 0.0018). Conversely, we found no associations in the DPRS

generated through the hypothesis-free approach.

Mass Spectrometry

379   The LC-MS dataset contained 48 samples: 27 DD and 21 controls. This assay identified 561

380   peptides. No peptides had statistically significant differences in abundance across groups (FDR

381   > 0.25).

Mass-Spectrometry-SomaScan concordance

To further validate our SomaScan findings, we evaluated the MS results for concordance. Of the

54 significant proteins identified in the SomaScan analysis, six were also detected in the Mass

Spectrometry findings: AOC3, POSTN, SERPINC1, TF, PCSK9, and KNG1. We ran the same

analyses on these six proteins using Mass Spectrometry data. AOC3, POSTN, and SERPINC1

had adjusted p-values (FDR) <0.25 in both Mass Spec and SomaScan analyses (Table 4).

Table 4. Mass Spectrometry (MS) – SomaScan (SS) concordance.

Protein	Est MS	p-val MS	FDR MS	Est SS	p-val SS	FDR SS
AOC3	0.214854	0.076413	0.232677	0.116624	0.000845	0.217972
POSTN	0.410212	0.098842	0.232677	0.182892	0.000312	0.211479
SERPINC1	0.071331	0.116338	0.232677	0.091474	0.001500	0.225251
TF	0.035200	0.567393	0.829196	0.076715	0.000025	0.172072
PCSK9	0.165811	0.755318	0.829196	0.185675	0.001371	0.222267
KNG1	0.010065	0.829196	0.829196	0.065561	0.000490	0.217972

Table 4. Proteins identified in both MS and SS analyses. Six proteins identified by MS were

also in the 54 proteins found in the hypothesis-free SS analysis: AOC3, POSTN, SERPINC1,

TF, PCSK9, and KNG1. Of these, AOC3, POSTN, and SERPINC1 had False Discovery Rate

(FDR, Adjusted p-value) <0.25 in both Mass Spec and SomaScan analyses. Est MS: Natural log

fold change in protein abundance between Dupuytren disease cases and controls, measured by

Mass Spectrometry, estimated using a generalized estimating equation (GEE) model adjusted

for age and gender. p-val MS: P-value for the Est MS, calculated from the robust standard error

of the GEE model. FDR MS: False discovery rate (FDR)–adjusted p-value for Mass

Spectrometry results, computed using the Benjamini–Hochberg procedure. Est SS: Natural log

fold change in protein abundance between Dupuytren disease cases and controls, measured by

SomaScan, estimated using a generalized estimating equation (GEE) model adjusted for age

and gender. p-val SS: P-value for the Est SS, calculated from the robust standard error of the

GEE model. FDR SS: False discovery rate (FDR)–adjusted p-value for SomaScan results,

computed using the Benjamini–Hochberg procedure.

Discussion

This study aimed to identify candidate biomarkers that correlate with DD diagnosis as a first

step toward a diagnostic DD blood test.

Dynamic biomarker panels are essential for developing preventive therapeutics for chronic

diseases such as DD. The relevance of this study is that DD lacks such biomarkers. Blood is the

most promising unexplored source of proteomic biomarkers for trend analysis.

There are no broad surveys of circulating proteins in DD. In addition, blood findings may not

parallel tissue findings because of rapid degradation, intracellular or membrane-bound

locations, post-translational variants, protein binding, and other mechanisms. Because of this,

new and diverse discovery research is necessary to profile DD-associated blood protein

biomarkers.

There are two common strategies for biomarker discovery: hypothesis-free and hypothesis-

based. The hypothesis-free approach (also called unbiased or non-hypothesis-based) screens

many markers and is the standard in systems biology and discovery research. The hypothesis-

based strategy evaluates selected markers based on published information and is the most

common model of DD research. We used both methods in this study.

Plasma collagen metabolites in DD

Collagen accumulation is the hallmark of DD. DD-affected tissues have unique proportions of

collagen types I [29], III [30], IV [16], and V [29]. The most prominent tissue changes in the

progression from preclinical to end-stage disease are increases in total Collagen I and the

proportion of Collagen III [29].

Our hypothesis-based approach compared markers of collagen synthesis and degradation

between the DD and control groups. Collagen turnover releases circulating peptides unique to

the synthesis and degradation of each collagen type. In the absence of disease, synthesis and

degradation markers are tightly coupled. Plasma synthesis-to-degradation marker ratios can

indicate net collagen gain, loss, or homeostasis [18]. We used ELISA to measure plasma levels

of 16 collagen metabolism markers across 8 collagen types. We summarize our findings in

Table 1.

The DD cohort had normal Pro-C1 levels, a marker of Collagen I synthesis, but showed a

statistically significant decrease in C1M, a marker of Collagen I degradation (Fig 1A), resulting

in an elevated synthesis-to-degradation marker ratio in DD compared to controls (Fig 1D).

These findings suggest DD collagen accumulation follows impaired Collagen I degradation

rather than excess collagen synthesis. Both degradation-resistant collagen variants and

reduced collagenase activity may contribute to reduced collagen degradation.

Dupuytren collagen has two features that increase resistance to enzymatic degradation. Typical

Collagen I fibrils are heterotrimers of two Collagen I α1 strands and one Collagen I α2 strand.

DD-affected tissues have an increased proportion of α1(I) homotrimeric Collagen I [31]. DD

collagen also has more crosslinks stabilized by hydroxylation and glycosylation [32]. Both

homotrimer Collagen I [33] and glycosylation-stabilized crosslinks [34] increase resistance to

enzymatic cleavage.

DD-related dysregulation of collagenases and their inhibitors impairs collagen degradation by

several mechanisms. TIMP1 (Tissue Inhibitor of Metalloproteinase 1) inhibits multiple MMPs

and is elevated in the circulation of DD patients [35]. MMP1, MMP8, MMP13, and MMP14 gene

polymorphisms are associated with Dupuytren disease [36, 37], including MMP14 partial loss-

of-function variants [37]. MMP14 is necessary for MMP2 activation, and both enzymes facilitate

myofibroblast contraction [38].

Two clinical findings support a relationship between impaired collagen degradation and DD. The

first is the rapid onset of DD during treatment with a metalloproteinase inhibitor [39]. The second

is abnormally slow matrix remodeling in DD. Enzymatic cleavage of collagen fibrils is necessary

for matrix shortening [40], and impaired cleavage would slow this process. Slow matrix

remodeling could explain why DD-related tissue shortening progresses more slowly than wound

healing despite similar biology, and why DD surgery is associated with an increased risk of

delayed wound healing [41]. The role of impaired collagen degradation in pathogenesis and

intervention is unclear, and simply normalizing collagen degradation might have the undesirable

effect of accelerating tissue contraction.

The hallmark of fibrotic diseases is the accumulation of collagen. Increased collagen synthesis

contributes to collagen accumulation in pulmonary fibrosis [42], liver fibrosis [43], primary biliary

cirrhosis [44], renal fibrosis [45], and cardiovascular disease [46]. Our finding of normal levels of

circulating Collagen I synthesis biomarkers in DD suggests that the mechanism of collagen

accumulation in DD is not due to the increased collagen synthesis associated with these

diseases. Although often used interchangeably with increased collagen synthesis, neither

collagen accumulation nor upregulated collagen gene expression is, in itself, evidence of

increased collagen synthesis. Cell culture studies show increased *COL1A1* and *COL1A2* mRNA

expression in DD-derived fibroblasts [47, 48], but these RNA findings are not associated with

increased collagen I protein production [48]. Our findings of normal Collagen I synthesis and

impaired degradation inform future drug development research beyond the traditional focus of

reducing collagen synthesis [49] to normalizing collagen degradation.

Collagen II markers were normal. We evaluated Collagen II markers because of the correlation

between DD and psoriasis [50] and the finding of abnormal Collagen II in psoriasis [51]. We

found no significant differences between groups in the Collagen II synthesis marker PRO-

C2_HP, the degradation marker C2M_HP, or their ratios. The CALC2 synthesis marker was

undetectable in both cohorts, preventing the calculation of CALC2 synthesis-to-degradation

ratios.

Collagen III synthesis markers were normal, consistent with a prior report of normal circulating

Collagen III synthesis marker amino-terminal propeptide of type III procollagen (PIIINP) in DD

subjects [52]. Given the accumulation of Collagen III in DD, we expected to find elevated

Collagen III synthesis-to-degradation marker ratios. Instead, we found normal Collagen III

synthesis-to-degradation ratios. This finding may reflect that other proteases also degrade

Collagen III, and we did not measure all potential collagen degradation fragments. To

investigate further, we assayed a different Col III degradation marker, CTX-III_HP, on the

remaining aliquots. Only 17 (10 DD, 7 Controls) of the initial 27 (13 DD, 14 Controls) leftover

aliquots had sufficient residual volumes for this assay. Although this small subgroup's CTX-III

violin plot appearance suggested reduced degradation in DD compared to controls (Fig S5), this

difference was not statistically significant.

We found significantly lower Pro-C5 levels, a marker of Collagen V synthesis, in the DD cohort

than in controls (Fig 1B). This finding is unexplained. An increased percentage of Collagen V is

reported in DD tissues [29], suggesting a high Collagen V synthesis-to-degradation ratio.

However, we could not calculate this ratio because a Collagen V degradation marker was

unavailable.

The DD cohort also had significantly lower levels of the Collagen VII synthesis marker Pro-C7

than controls (Fig 1C). We could not calculate Collagen VII synthesis-to-degradation ratios

because Collagen VII degradation markers were undetectable in both cohorts. Although there

are no reports of Collagen VII in DD, its role in matrix mechanobiology and elevated tissue

levels in systemic sclerosis [53] make it a potential marker for collagen metabolism

abnormalities in DD.

We found no significant differences in the remaining collagen markers. We measured Collagen

IV markers because Collagen IV is elevated in DD tissues [16]. We assayed Collagen VI

markers and ratios because prior pathway analysis suggests a role for Collagen VI in DD and

other fibrotic diseases [54]. We selected Collagen IX markers based on unpublished data

suggesting a role in DD. None of these markers or ratios showed significant differences

between disease and control cohorts.

Plasma proteins and interactions in DD

In addition to the Nordic collagen fragment analysis, we analyzed whole proteins using the

SomaScan aptamer-based platform. The SomaScan panel measures 6995 protein-binding sites

associated with 5444 proteins. S6 Table lists genes and gene aliases of proteins mentioned in

this manuscript.

Our SomaScan hypothesis-free analysis identified 54 distinct proteins with plasma levels that

were significantly different from those of controls. Twenty-four proteins showed increased levels,

and 30 showed lower DD levels than controls (S1 Table). Our findings are consistent with

existing literature.

The body of DD research publications mentions only seven of these 54 differentially expressed

proteins: Aggrecan (*ACAN*), Amine Oxidase Copper Containing 3 (*AOC3*), Caspase-3 (*CASP3*),

Periostin (*POSTN*), SMAD Family Member 1 (*SMAD1*), Spartin (*SPART*), and Ubiquitin

carboxyl-terminal hydrolase 8 (*USP8*). Our differential expression findings concur with reports of

ACAN [55], *AOC3 [13]*, and *POSTN [56]* overexpression in tissue and *SMAD1 [57]*, *SPART [58]*,

and *USP8 [58]* underexpression in cell culture. CASP3 levels were low in our findings, but are

upregulated in a transcriptomic profiling report [55]. This finding is unexplained. *ACAN* is also

near the variant rs6496519, which is associated with DD [59]. S1 Table summarizes the

relevance of these 54 proteins and their parent genes to DD.

We constructed PPI networks and enrichment analysis of these 54 plasma proteins, revealing

significantly enriched protein-protein interactions in 32, with multiple enriched protein groups

and categories (Fig S2, S2 Table). These categories align with known Dupuytren cellular

biology. Abnormalities in the KEGG category has04610 (complement and coagulation

cascades) were consistent with DD pathway meta-analysis findings [60]. Complement

fragments induce adhesion molecules that act on endothelial cells, consistent with DD-related

perivascular clusters of immune cells and small vessel occlusion [13]. Complement fragments

also activate mast cells that release TNF, a known DD pathway component [61], and histamine,

a driver of itching, which is common in early-stage DD [62]. Cell membrane and extracellular

matrix signaling receptor binding of growth factors, cytokines, and other signaling factors play

pivotal roles in DD pathobiology [14]. Protein phosphorylation processes are consistent with

increased phosphorylated STAT1, STAT3, SMAD2, SMAD3, and ERK1/2 in DD tissues [57, 63,

64]. The enriched Reactome HSA-162582 (Signal Transduction) pathway is noteworthy, as it

includes signaling by TGF-Beta, WNT, and NOTCH, all of which are prominent in DD biology

[65].

We based our hypothesis-free DPRS on high levels of Osteoclast-associated immunoglobulin-

like receptor (*OSCAR*) and low levels of Spartin (*SPART*) (Table 2). Our finding of elevated

OSCAR levels is interesting because Collagen I is an OSCAR ligand. OSCAR also stimulates T-

cells to release TNF [66], a component of DD biology [61]. Our finding of decreased *SPART*

levels is consistent with low *SPART* expression in DD fibroblasts [58]. SPART is also associated

with Epidermal Growth Factor Receptor (EGFR) degradation and transport to the cell

membrane [67], consistent with increased ratios of intracellular to cell membrane EGFR levels

observed in DD [68].

In addition to the hypothesis-free analysis, we conducted a hypothesis-based analysis using a

subset of SomaScan proteins selected from an independent literature review (S3 Table). We

identified 23 proteins in this subset with significantly different plasma levels compared to

controls. Nine proteins had higher levels in DD than in controls, and fourteen had lower levels.

Nine proteins were present in both hypothesis-free and hypothesis-based groups. Our findings

are consistent with existing literature. S4 Table summarizes the potential relationships among

these proteins, their parent genes, and DD.

With our hypothesis-free analysis, we performed a STRING PPI analysis of the hypothesis-

based findings to identify network relationships of the 23 proteins. We found significantly

enriched protein-protein interactions in 21 with multiple enriched categories (p-value = 1.35E-

07, S5 Table). This analysis implicated additional enrichment categories beyond those from the

hypothesis-free analysis. Myofibroblast differentiation, proliferation, motility, WNT signaling, and

interleukin pathways are unsurprising, given their known roles in DD biology [10, 69]. Cancer

pathway associations are also unsurprising, given the correlations between DC and carcinomas

and sarcomas [70, 71]. Infection-related pathways were unexpected and were exclusively

associated with viral and parasitic infection categories. This finding might be due to a hidden

variable of tissue repair biology, which overlaps interleukin-driven parasitic immune defense

biology and interleukin-related DD biology.

This hypothesis-based analysis generated data for a second Diagnostic Proteomic Risk Score

(DPRS) calculated from 11 proteins (Table 3). Fig 4A shows the AUC-ROC of this panel (AUC =

0.9), and Fig 4B shows the DPRS based on this panel, distinguishing DD from control cohorts.

We evaluated the potential of each DPRS for staging. Recurrence after a procedure is the

benchmark of Dupuytren's biological severity. Traditionally, surgeons use clinical factors such

as early age of onset, affected family members, and disease locations other than the palms to

predict post-procedure recurrence. However, the study's inclusion criteria included these

factors, so we could not use them to subset our Dupuytren cohort. Instead, we used age at the

time of the first DC procedure. Recurrence is more common in patients who are younger at their

first DC procedure [72-76]. We evaluated each DPRS to distinguish Dupuytren subjects who

had their first corrective procedure before age 50 from those who had their first correction at age

50 or older. The hypothesis-based DPRS distinguished DD subjects who underwent their first

DC procedure before age 50 from those who underwent their first DC procedure after age 50

(p=0.0018), suggesting potential use as a staging and predictive tool. An unexpected and

unexplained finding is that DPRS scores were higher in DD than in controls, but hypothesis-

based DPRS scores were higher in DD patients who had their first procedure after age 50 than

in those who were younger at the time of their first corrective procedure.

Although Mass Spectrometry did not reveal statistically significant differences in abundance

across cohorts, there was statistically significant concordance between Mass Spec and

SomaScan findings. The Mass Spec data contained 6 of the 54 proteins with statistically

significant differences between disease and control cohorts using the hypothesis-free

SomaScan data analysis: Amine Oxidase Copper Containing 3 *AOC3*, Periostin *POSTN*, Serpin

Family C Member 1 *SERPINC1*, Transferrin *TF*, Proprotein Convertase Subtilisin/Kexin Type 9

PCSK9, and Kininogen 1 *KNG1*. Table 4 compares the results of the SomaScan and Mass

Spectrometry data analysis. A pathway analysis of these six proteins identified two highly

significant Reactome categories (S7 Table). In addition to protein phosphorylation noted above,

Reactome HSA-381426 involving Insulin-like Growth Factor *IGF* and Insulin-like Growth Factor

Binding Proteins *IGFBP*s is relevant because of the role of IGF-II and IGFBP-6 in regulating

Dupuytren fibroblast proliferation and contractility [77].

This study has limitations. Statistical disadvantages included the small sample sizes across all

assays, and the large number of aptamers in the SomaScan assay. Although our findings align

with current research on Dupuytren proteomics, there are differences. Elevated serum TIMP1

[35] and TNF [69] have been reported in DD, but we found no statistically significant DE

between disease and control groups. This discrepancy may reflect that SomaScan assays

target specific protein conformations, whereas ELISA measures all conformations.

Conclusions

We used ELISA, aptamer, and mass spectrometry assays to identify differential plasma

proteomic signatures between DD and control cohorts. We found significant DD-related

proteomic abnormalities in both circulating collagen metabolism peptide fragments and whole

proteins. This study is the first to provide evidence suggesting net collagen accumulation in DD

results from normal collagen synthesis and impaired collagen degradation, challenging the

model of excess collagen synthesis and distinguishing DD from other organ fibroses. We

independently analyzed proteins from the same cohorts. Our hypothesis-free analysis of 5444

proteins revealed 54 with statistically significant differences between the DD and control

cohorts. Our independent hypothesis-based analysis of 328 proteins selected from the literature

review identified 23 with statistically significant differences. We developed two Dupuytren

Diagnostic Proteomic Risk Scores, one based on hypothesis-free and the other based on

hypothesis-driven analyses. We used pathway analyses to identify enrichments in the DD

protein-protein network, confirming known DD interactions and revealing previously unreported

DD pathway categories. Finally, we showed our hypothesis-based DPRS correlated with the

subject's age at the time of the first DD procedure, an index of biological severity. These findings

suggest that a validated, hypothesis-based DPRS could guide management and the

development of DD therapeutics.

We recommend a larger study to validate our findings on the metabolites of Collagen I, III, V,

and VII. We also recommend a validating study of 68 unique proteins identified in the protein

aptamer analysis (54 in the hypothesis-free analysis and 23 in the hypothesis-based analysis,

with 9 in both groups). Testing only the subset of the 68 most pertinent proteins would increase

statistical power. Individual ELISA assays would be cost-prohibitive. An alternative approach

would be to develop a custom multiplex panel for the 13 proteins in our two DPRS [14] to

validate our DPRS and to assess the correlation with our hypothesis-based DPRS on age at the

first DD procedure. Our findings strongly support the need to fund and conduct more extensive

validating studies of circulating collagen metabolites and proteins in DD. This biomarker

research is fundamental to developing DD diagnostic tests, staging DD, and informing the

development of therapeutic drugs for DD.

Acknowledgments

Gail Solomon assisted with manuscript writing.

Databank and study support from FORWARD, the National Databank for Rheumatic Diseases,

is gratefully acknowledged.

Author Contributions

BH: Formal Analysis, Data Curation, Methodology, Software, Validation, Visualization, Writing

original draft, review, and editing. PS: Formal Analysis, Data Curation, Methodology, Software,

Validation, Visualization, Writing original draft, review, and editing. AL: Formal Analysis,

Software, Validation, Visualization. AG: Formal Analysis, Software, Validation, Visualization. CB:

Formal Analysis. MK: Formal Analysis, Methodology. SN: Formal Analysis. CE:

Conceptualization, Data Curation, Investigation, Methodology, Visualization, Writing original

draft, review, and editing.

Supporting information

S1 Fig. Distribution of samples to laboratories. Forty-five subjects provided initial samples.

Seventeen of these subjects had second samples drawn at least 6 months after the first. Mass

spec lab 1 performed analysis on the initial 45 samples. Mass spec lab 2 and collagen

metabolite analyses were performed on redraw specimens. Aptamer-based analysis was

performed on both initial and redraw specimens. Three of the original forty-five samples for

Aptamer analysis failed quality control, leaving 23 DD and 19 Controls available for analysis.

S2 Fig. Protein-protein network enrichment categories in the hypothesis-free analysis.

Thirty-two differentially expressed proteins showed significant network enrichment (p.adj < 0.05)

in the following biological categories. C&C: coagulation and complement cascades, ECM:

extracellular matrix, ENZ: Enzyme inhibition, MBR: cell membrane structures, PPL:

phosphorylation, SGN: other signaling pathways.

S3 Fig. Protein-protein network enrichment categories in the hypothesis-based analysis.

Twenty-one differentially expressed proteins showed significant network enrichment (p.adj <

0.05) in the following biological categories. CAN: Cancer, C&C: coagulation and complement

cascades, ECM: extracellular matrix, INL: interleukin interactions, INF: infection, MBR: cell

membrane structures, MFB: myofibroblast differentiation, proliferation, and motility, PPL:

phosphorylation, SGN: other signaling pathways, WNT: WNT pathways. Hypothesis-based

selection bias from a preselected list of candidate proteins may influence the enrichment results.

S4 Fig. Hypothesis-based DPRS correlates with age at the time of the first Dupuytren

procedure. This 11-protein Hypothesis-Based Dupuytren Proteomic Risk Score (HB-DPRS)

distinguished DD subjects with different disease progression rates based on whether the subject

was younger than 50 vs. 50 or older during their first corrective procedure (p=0.0018).

S5 Fig. Differential expression of Dupuytren vs. control Collagen III degradation marker

CTX-III. Although these violin plots visually suggest reduced Collagen III degradation marker

CTX-III values in DD compared to controls, this difference was not significant (p-value =

0.4887). These data were from 17 of the 27 samples used for the other collagen metabolism

markers, due to insufficient volumes in 10 samples. Because small cohorts magnify the effects

of outliers, we recommend repeating these assays on larger cohorts.

S1 Table. 54 Differentially expressed genes in the SomaScan Hypothesis-free analysis. Of

the 6995 aptamers in the Hypothesis-free analysis, 54 proteins showed statistically significant

differences between the DD and control groups, with 24 overexpressed and 30 underexpressed

after adjusting for multiple comparisons. Nine of these proteins were identified in both

hypothesis-free and hypothesis-based analyses, as indicated by *. Plain italics indicate an

indirect DD relationship, if any, to the protein; bold italics indicate a direct relationship between

published DD findings and the protein [10, 13, 55, 57, 58, 61, 66, 68, 71, 78, 82, 84-126]. In the

Notes column, "Referred to" means the referenced publication referred to the gene by an alias

name. Exp: DD expression compared to controls. p-val: p-value of expression difference. FDR:

False discovery rate (adjusted p-value. Nodes: number of pathway analysis connections

(nodes) of this protein to others in this group with a medium confidence interaction score (0.400)

and an FDR<=0.05. Notes: potential relationship to DD biology. DOI: DOIs of publications

referenced in Notes. Cat: categories of possible relationships to DD: 1. Apoptosis and

senescence (3 genes); 2. Bioinformatics (9 genes); 3. Clinical and demographic (6 genes); 4.

Extracellular matrix (6 genes); 5. Fibroblast and myofibroblast cytoskeleton, membrane, and

motility (10 genes); 6. Fibroblast and myofibroblast differentiation and transcriptome (7 genes);

7. Vascular and perivascular (9 genes); Unclear relationship (16 genes).

S2 Table. Functionally enriched pathways derived from the 54 differentially expressed

proteins in the hypothesis-free analysis. ID: Enriched category name. STR: Strength of

protein-protein interaction (PPI). FDR: False Discovery Rate (Adjusted p-value). Overall PPI

enrichment p-value: 0.000455. Figure 4 summarizes enriched pathways for individual genes.

S3 Table. 328 protein-coding genes in the SomaLogic hypothesis-based analysis. We

combined three literature-based search strategies to identify candidate Dupuytren-related genes

for hypothesis-based analytics. We first collected 2547 full-text publications from 1980 to 2023

that included at least three instances of the search terms "Dupuytren*" and/or "palmar fibroma*".

In the first strategy, we searched each publication for 4527 potential Dupuytren-related genes

and their 33907 gene name aliases. This search identified 326 unique proteins or parent genes

mentioned at least three times in at least one publication. The second method identified 117

selected collagen metabolism-related proteins or their parent genes that appeared at least once

in at least one of these publications. The third method identified potentially DD-related single-

nucleotide polymorphisms (SNPs) reported in these publications and compiled a list of 369

genes adjacent to these SNPs. We merged these three lists, resulting in 546 unique protein-

coding genes, 328 of which matched proteins on the SomaScan panel, which contains 6995

protein-binding sites. Proteins expressed by these 328 genes were the targets of our

hypothesis-based analysis.

S4 Table. 23 Differentially expressed genes in the Hypothesis-based analysis. Of the 328

aptamers in the Hypothesis-based analysis, 23 had statistically significant differences between

DD and control, with nine overexpressed and fourteen underexpressed after adjusting for

multiple comparisons. Nine of these genes appeared in hypothesis-free and hypothesis-based

analyses, as indicated by *. Plain italics indicate an indirect DD relationship, if any, to the

protein; bold italics indicate a direct relationship between published DD findings and the protein

[10, 13, 31, 33, 37, 55, 57, 58, 59, 64, 78, 79, 84, 87, 88, 92, 93, 94, 108, 109, 117-119, 125-

143]. In the Notes column, "Referred to" means the referenced publication referred to the gene

by an alias name. Exp: DD expression compared to controls. FDR: False Discovery Rate

(Adjusted p-value) of expression difference. Nodes: number of pathway analysis connections

(nodes) of this protein to others in this group with a medium confidence interaction score (0.400)

and an FDR<=0.05. Notes: potential relationship to DD biology. DOI: DOIs of publications

referenced in Notes. Cat: categories of possible relationships to DD: 1. Apoptosis and

senescence (2 genes); 2. Bioinformatics (7 genes); 3. Clinical and demographic (3 genes); 4.

Extracellular matrix (9 genes); 5. Fibroblast and myofibroblast cytoskeleton, membrane, and

motility (6 genes); 6. Fibroblast and myofibroblast differentiation and transcriptome (5 genes); 7.

Vascular and perivascular (4 genes).

S5 Table. Functionally enriched pathways derived from the 23 differentially expressed

proteins in the hypothesis-based analysis. ID: Enriched category name. STR: Strength of

protein-protein interaction (PPI). FDR: False Discovery Rate (Adjusted p-value). Overall PPI

enrichment p-value: 1.35E-07. Hypothesis-based candidate protein preselection may influence

enrichment results. Figure 6 summarizes enriched pathways for individual genes.

S6 Table. Genes referenced in the manuscript and their aliases. This table lists all gene

names referenced in the manuscript, their UniProt identifiers, protein names, and gene aliases.

S7 Table. Enriched pathways in Mass Spec - SomaScan concordant genes. Six proteins

identified by Mass Spectrometry had statistically significant differences between Dupuytren and

control cohorts in the SomaScan analysis, representing genes AOC3, POSTN, SERPINC1, TF,

PCSK9, and KNG1. Within this network, AOC3, POSTN, and PCSK9 had strong pathway

connections, and PCSK9, SERPINC1, TF, and KNG1 had functionally enriched pathways. PPI

enrichment p-value: 0.00987. STR: Strength of association. FDR: False Discovery Rate

(adjusted p-value).

Data availability

737   All the data and analytic scripts necessary to replicate the findings are available in the GitHub

738   repository Dup_Blood_Proteomic_Data at

739   https://github.com/Dupuytren/Dup_Blood_Proteomic_Data.

Competing interest statement

741 The authors declare the following competing interests: Morten Karsdal, Cecilie Bager, and Signe

742 Holm Nielsen are employees of Nordic Bioscience and hold stocks in Nordic Bioscience. No

743 other competing interests have been declared. This does not alter the authors' adherence to

744 PLOS ONE policies on sharing data and materials.

References

1.   Macaulay D, Ivanova J, Birnbaum H, Sorg R, Skodny P. Direct and indirect costs

associated with Dupuytren's contracture. J Med Econ. 2012;15(4):664–71. Epub 2012/03/01.

doi: 10.3111/13696998.2012.670678. PubMed PMID: 22369346.

2.   Broekstra DC, Kuo RYL, Burn E, Prieto-Alhambra D, Furniss D. Dupuytren Disease:

Prevalence, Incidence, and Lifetime Risk of Surgical Intervention. A Population-Based Cohort

Analysis. Plast Reconstr Surg. 2023;151(3):581–91. Epub 20221122. doi:

10.1097/PRS.0000000000009919. PubMed PMID: 36730480; PubMed Central PMCID:

PMCPMC9944385.

3.   Mendelaar NHA, Poelstra R, van Nieuwenhoven CA, Slijper HP, Feitz R, Hovius SER, et

al. Outcome of Recurrent Surgery in Dupuytren's Disease: Comparison with Initial Treatment.

Plast Reconstr Surg. 2019;144(5):828e–35e. Epub 2019/08/10. doi:

10.1097/PRS.0000000000006150. PubMed PMID: 31397794.

4.   Goodman AD, Gil JA, Starr AM, Akelman E, Weiss A-PC. Incidence of Serious

Complications in Hand Surgery: A 10-Year Review. The Journal of Hand Surgery.

2016;41(9):S27–S8. doi: 10.1016/j.jhsa.2016.07.047.

5.   Schindelar L, McEntee R, D'Amore T, Beredjiklian P, Lutsky K. Unplanned Return to the

Operating Room in Upper-Extremity Surgery: Incidence and Reason for Return. J Hand Surg

Am. 2021;46(8):715 e1– e12. Epub 20210514. doi: 10.1016/j.jhsa.2021.01.019. PubMed PMID:

33994259.

6.   Denkler K. Surgical complications associated with fasciectomy for dupuytren's disease: a

20-year review of the English literature. Eplasty. 2010;Jan 27(10):e15. PubMed Central PMCID:

PMCPMC2828055.

7.   Alser O, Craig RS, Lane JCE, Prats-Uribe A, Robinson DE, Rees JL, et al. Serious

complications and risk of re-operation after Dupuytren's disease surgery: a population-based

cohort study of 121,488 patients in England. Sci Rep. 2020;10(1):16520. Epub 2020/10/07. doi:

10.1038/s41598-020-73595-y. PubMed PMID: 33020582; PubMed Central PMCID:

PMCPMC7536429.

8.   Degreef I, De Smet L. Dupuytren's disease: a predominant reason for elective finger

amputation in adults. Acta Chir Belg. 2009;109(4):494–7. Epub 2009/10/07. doi:

10.1080/00015458.2009.11680467. PubMed PMID: 19803262.

9.   Larsen S, Krogsgaard DG, Aagaard Larsen L, Iachina M, Skytthe A, Frederiksen H.

Genetic and environmental influences in Dupuytren's disease: a study of 30,330 Danish twin

pairs. J Hand Surg Eur Vol. 2015;40(2):171–6. Epub 2014/05/20. doi:

10.1177/1753193414535720. PubMed PMID: 24835475; PubMed Central PMCID:

PMCPMC4810018.

10.   Dolmans GH, Werker PM, Hennies HC, Furniss D, Festen EA, Franke L, et al. Wnt

signaling and Dupuytren's disease. N Engl J Med. 2011;365(4):307–17. Epub 2011/07/08. doi:

10.1056/NEJMoa1101029. PubMed PMID: 21732829.

11.   Riesmeijer SA, Manley OWG, Ng M, Nolte IM, Broekstra DC, Werker PMN, et al. A

Weighted Genetic Risk Score Predicts Surgical Recurrence Independent of High-Risk Clinical

Features in Dupuytren's Disease. Plast Reconstr Surg. 2019;143(2):512–8. Epub 2019/01/29.

doi: 10.1097/PRS.0000000000005208. PubMed PMID: 30688894; PubMed Central PMCID:

PMCPMC6358194.

12.   Iqbal SA, Hindocha S, Farhatullah S, Paus R, Bayat A. Dupuytren's Disease Shows

Populations of Hematopoietic and Mesenchymal Stem-Like Cells Involving Perinodular Fat and

Skin in Addition to Diseased Fascia: Implications for Pathogenesis and Therapy. Dupuytren's

Disease and Related Hyperproliferative Disorders2012. p. 167–74.

13.   Mayerl C, Del Frari B, Parson W, Boeck G, Piza-Katzer H, Wick G, et al.

Characterisation of the inflammatory response in Dupuytren's disease. J Plast Surg Hand Surg.

2016;50(3):171–9. Epub 2016/02/09. doi: 10.3109/2000656X.2016.1140054. PubMed PMID:

26852784.

14.   Gonga-Cave BC, Pena Diaz AM, O'Gorman DB. Biomimetic analyses of interactions

between macrophages and palmar fascia myofibroblasts derived from Dupuytren's disease

reveal distinct inflammatory cytokine responses. Wound Repair Regen. 2021;29(4):627–36.

Epub 20210611. doi: 10.1111/wrr.12928. PubMed PMID: 34212454.

15.   Halliday NL, Rayan GM, Zardi L, Tomasek JJ. Distribution of ED-A and ED-B containing

fibronectin isoforms in Dupuytren's disease. J Hand Surg Am. 1994;19(3):428–34. Epub

1994/05/01. doi: 10.1016/0363-5023(94)90057-4. PubMed PMID: 8056970.

16.   Magro G, Fraggetta F, Travali S, Lanzafame S. Immunohistochemical expression and

distribution of alpha2beta1, alpha6beta1, alpha5beta1 integrins and their extracellular ligands,

type IV collagen, laminin and fibronectin in palmar fibromatosis. Gen Diagn Pathol.

1997;143(4):203–8. Epub 1998/03/07. PubMed PMID: 9489951.

17.   Broekstra DC, Lanting R, Werker PMN, van den Heuvel ER. Disease Course of Primary

Dupuytren Disease: 5-Year Results of a Prospective Cohort Study. Plast Reconstr Surg.

2022;149(6):1371–8. Epub 20220411. doi: 10.1097/PRS.0000000000009115. PubMed PMID:

35404338.

18.   Karsdal MA, Leeming DJ, Henriksen K, Bay-Jensen AC, Nielsen SH, CL B. Biochemistry

of Collagens, Laminins and Elastin. Boston: Elsevier; 2019.

19.   Hughes CS, Moggridge S, Muller T, Sorensen PH, Morin GB, Krijgsveld J. Single-pot,

solid-phase-enhanced sample preparation for proteomics experiments. Nat Protoc.

2019;14(1):68–85. Epub 2018/11/23. doi: 10.1038/s41596-018-0082-x. PubMed PMID:

30464214.

20.   Leutert M, Rodriguez-Mias RA, Fukuda NK, Villen J. R2-P2 rapid-robotic

phosphoproteomics enables multidimensional cell signaling studies. Mol Syst Biol.

2019;15(12):e9021. doi: 10.15252/msb.20199021. PubMed PMID: 31885202; PubMed Central

PMCID: PMCPMC6920700.

21.   Amodei D, Egertson J, MacLean BX, Johnson R, Merrihew GE, Keller A, et al. Improving

Precursor Selectivity in Data-Independent Acquisition Using Overlapping Windows. J Am Soc

Mass Spectrom. 2019;30(4):669–84. Epub 20190122. doi: 10.1007/s13361-018-2122-8.

PubMed PMID: 30671891; PubMed Central PMCID: PMCPMC6445824.

22.   Searle BC, Pino LK, Egertson JD, Ting YS, Lawrence RT, MacLean BX, et al.

Chromatogram libraries improve peptide detection and quantification by data independent

acquisition mass spectrometry. Nat Commun. 2018;9(1):5128. Epub 2018/12/05. doi:

10.1038/s41467-018-07454-w. PubMed PMID: 30510204; PubMed Central PMCID:

PMCPMC6277451.

23.   Hoofnagle AN, Becker JO, Oda MN, Cavigiolio G, Mayer P, Vaisar T. Multiple-reaction

monitoring-mass spectrometric assays can accurately measure the relative protein abundance

in complex mixtures. Clin Chem. 2012;58(4):777–81. Epub 20120203. doi:

10.1373/clinchem.2011.173856. PubMed PMID: 22307200; PubMed Central PMCID:

PMCPMC3665768.

24.   Gold L, Ayers D, Bertino J, Bock C, Bock A, Brody EN, et al. Aptamer-based multiplexed

proteomic technology for biomarker discovery. PLoS One. 2010;5(12):e15004. Epub 20101207.

doi: 10.1371/journal.pone.0015004. PubMed PMID: 21165148; PubMed Central PMCID:

PMCPMC3000457.

25.   Berrone E, Chiorino G, Guana F, Benedetti V, Palmitessa C, Gallo M, et al. SOMAscan

Proteomics Identifies Novel Plasma Proteins in Amyotrophic Lateral Sclerosis Patients. Int J Mol

Sci. 2023;24(3). Epub 20230118. doi: 10.3390/ijms24031899. PubMed PMID: 36768220;

PubMed Central PMCID: PMCPMC9916400.

26.   Network TCRA. gee: Generalized Estimation Equation Solver 2023 [01/06/2023].

Available from: https://cran.r-project.org/package=gee.

27.   Kuhn M. Building Predictive Models in R Using the caret Package. Journal of Statistical

Software [Internet]. 2008; 28(5).

28.   Szklarczyk D, Kirsch R, Koutrouli M, Nastou K, Mehryary F, Hachilif R, et al. The

STRING database in 2023: protein-protein association networks and functional enrichment

analyses for any sequenced genome of interest. Nucleic Acids Res. 2023;51(D1):D638–D46.

doi: 10.1093/nar/gkac1000. PubMed PMID: 36370105; PubMed Central PMCID:

PMCPMC9825434.

29.   Bailey AJ. Collagen. In: McFarlane R, McGrouther D, Flint M, editors. Dupuytren's

Disease Biology and Treatment. Hand and Upper Limb Series. 5. Edinburg: Churchill

Livingstone; 1990.

30.   Bailey AJ, Sims TJ, Gabbiani G, Bazin S, LeLous M. Collagen of Dupuytren's disease.

Clin Sci Mol Med. 1977;53(5):499–502. Epub 1977/11/01. doi: 10.1042/cs0530499. PubMed

PMID: 589933.

31.   Williamson K, Cooper G, Lee KJ, Beamish EL, Simpson D, Carter A, et al. Active

synthesis of type I collagen homotrimer in Dupuytren's fibrosis is unaffected by anti-TNF-α

treatment. bioRxiv preprint 2023. doi: 10.1101/2020.07.13.195107.

32.   Bazin S, Le Lous M, Duance VC, Sims TJ, Bailey AJ, Gabbiani G, et al. Biochemistry

and histology of the connective tissue of Dupuytren's disease lesions. Eur J Clin Invest.

1980;10(1):9–16. Epub 1980/02/01. doi: 10.1111/j.1365-2362.1980.tb00003.x. PubMed PMID:

6768572.

33.   Han S, Makareeva E, Kuznetsova NV, DeRidder AM, Sutter MB, Losert W, et al.

Molecular mechanism of type I collagen homotrimer resistance to mammalian collagenases. J

Biol Chem. 2010;285(29):22276–81. Epub 2010/05/14. doi: 10.1074/jbc.M110.102079. PubMed

PMID: 20463013; PubMed Central PMCID: PMCPMC2903388.

34.   Snedeker JG, Gautieri A. The role of collagen crosslinks in ageing and diabetes - the

good, the bad, and the ugly. Muscles Ligaments Tendons J. 2014;4(3):303–8. Epub 2014/12/10.

PubMed PMID: 25489547; PubMed Central PMCID: PMCPMC4241420.

35.   Ulrich D, Hrynyschyn K, Pallua N. Matrix metalloproteinases and tissue inhibitors of

metalloproteinases in sera and tissue of patients with Dupuytren's disease. Plast Reconstr Surg.

2003;112(5):1279–86. Epub 2003/09/25. doi: 10.1097/01.PRS.0000081462.40448.49. PubMed

PMID: 14504511.

36.   Rodrigues MP, Tissi LH, Oliveira VM, Wistuba G, Araujo FB, Mattar-Junior R, et al.

MMP-1, MMP-8, and MMP-13 Gene Polymorphisms and Haplotype Is a Risk Factor for

Dupuytren Contracture: A Case-Control Study. Hand (N Y). 2024:15589447241242818. Epub

20240425. doi: 10.1177/15589447241242818. PubMed PMID: 38660990.

37.   Itoh Y, Ng M, Wiberg A, Inoue K, Hirata N, Paiva KBS, et al. A common SNP risk variant

MT1-MMP causative for Dupuytren's disease has a specific defect in collagenolytic activity.

Matrix Biol. 2021;97:20–39. Epub 2021/02/17. doi: 10.1016/j.matbio.2021.02.003. PubMed

PMID: 33592276.

38.   Wilkinson JM, Davidson RK, Swingler TE, Jones ER, Corps AN, Johnston P, et al. MMP-

14 and MMP-2 are key metalloproteases in Dupuytren's disease fibroblast-mediated

contraction. Biochim Biophys Acta. 2012;1822(6):897–905. Epub 2012/02/22. doi:

10.1016/j.bbadis.2012.02.001. PubMed PMID: 22342364.

39.   Krzeski P, Buckland-Wright C, Balint G, Cline GA, Stoner K, Lyon R, et al. Development

of musculoskeletal toxicity without clear benefit after administration of PG-116800, a matrix

metalloproteinase inhibitor, to patients with knee osteoarthritis: a randomized, 12-month,

double-blind, placebo-controlled study. Arthritis Res Ther. 2007;9(5):R109. Epub 2007/10/26.

doi: 10.1186/ar2315. PubMed PMID: 17958901; PubMed Central PMCID: PMCPMC2212568.

40.   Castella LF, Buscemi L, Godbout C, Meister JJ, Hinz B. A new lock-step mechanism of

matrix remodelling based on subcellular contractile events. J Cell Sci. 2010;123(Pt 10):1751–

60. Epub 2010/04/30. doi: 10.1242/jcs.066795. PubMed PMID: 20427321.

41.   Bismil Q, Bismil M, Bismil A, Neathey J, Gadd J, Roberts S, et al. The development of

one-stop wide-awake dupuytren's fasciectomy service: a retrospective review. JRSM Short Rep.

2012;3(7):48. Epub 2012/08/22. doi: 10.1258/shorts.2012.012050. PubMed PMID: 22908029;

PubMed Central PMCID: PMCPMC3422854.

42.   Organ LA, Duggan AR, Oballa E, Taggart SC, Simpson JK, Kang'ombe AR, et al.

Biomarkers of collagen synthesis predict progression in the PROFILE idiopathic pulmonary

fibrosis cohort. Respir Res. 2019;20(1):148. Epub 20190712. doi: 10.1186/s12931-019-1118-7.

PubMed PMID: 31299951; PubMed Central PMCID: PMCPMC6624898.

43.   Liu T, Wang X, Karsdal MA, Leeming DJ, Genovese F. Molecular serum markers of liver

fibrosis. Biomark Insights. 2012;7:105–17. Epub 20120723. doi: 10.4137/BMI.S10009. PubMed

PMID: 22872786; PubMed Central PMCID: PMCPMC3412619.

44.   Vesterhus M, Nielsen MJ, Hov JR, Saffioti F, Manon-Jensen T, Leeming DJ, et al.

Comprehensive assessment of ECM turnover using serum biomarkers establishes PBC as a

high-turnover autoimmune liver disease. JHEP Rep. 2021;3(1):100178. Epub 20200903. doi:

10.1016/j.jhepr.2020.100178. PubMed PMID: 33225252; PubMed Central PMCID:

PMCPMC7666353.

45.   Han DC, Isono M, Hoffman BB, Ziyadeh FN. High glucose stimulates proliferation and

collagen type I synthesis in renal cortical fibroblasts: mediation by autocrine activation of TGF-

beta. J Am Soc Nephrol. 1999;10(9):1891–9. doi: 10.1681/ASN.V1091891. PubMed PMID:

10477140.

46.   Genovese F, Goncalves I, Holm Nielsen S, Karsdal MA, Edsfeldt A, Nilsson J, et al.

Plasma levels of PRO-C3, a type III collagen synthesis marker, are associated with arterial

stiffness and increased risk of cardiovascular death. Atherosclerosis. 2024;388:117420. Epub

20231210. doi: 10.1016/j.atherosclerosis.2023.117420. PubMed PMID: 38128431.

47.   Satish L, Gallo PH, Baratz ME, Johnson S, Kathju S. Reversal of TGF-beta1 stimulation

of alpha-smooth muscle actin and extracellular matrix components by cyclic AMP in

Dupuytren's-derived fibroblasts. BMC Musculoskelet Disord. 2011;12:113. Epub 2011/05/27.

doi: 10.1186/1471-2474-12-113. PubMed PMID: 21612641; PubMed Central PMCID:

PMCPMC3125251.

48.   van Beuge MM, Ten Dam EJ, Werker PM, Bank RA. Matrix and cell phenotype

differences in Dupuytren's disease. Fibrogenesis Tissue Repair. 2016;9:9. Epub 2016/07/02.

doi: 10.1186/s13069-016-0046-0. PubMed PMID: 27366208; PubMed Central PMCID:

PMCPMC4928329.

49.   Lambi AG, Popoff SN, Benhaim P, Barbe MF. Pharmacotherapies in Dupuytren Disease:

Current and Novel Strategies. J Hand Surg Am. 2023. Epub 20230317. doi:

10.1016/j.jhsa.2023.02.003. PubMed PMID: 36935324.

50.   Patel M, Freeman NR, Dhaliwal S, Wright N, Daoud Y, Ryan C, et al. The prevalence of

Dupuytren contractures in patients with psoriasis. Clin Exp Dermatol. 2014;39(8):894–9. Epub

2014/09/10. doi: 10.1111/ced.12424. PubMed PMID: 25196833.

51.   Holm Nielsen S, Magee C, Groen SS, Sinkeviciute D, Bay-Jensen AC, Karsdal MA, et al.

Differentiating patients with psoriasis from psoriatic arthritis using collagen biomarkers. Clin Exp

Rheumatol. 2022. Epub 20220728. doi: 10.55563/clinexprheumatol/jmt9jv. PubMed PMID:

35916294.

52.   Gassier N, Brenner P, Berger A, Delbruck A. Biochemical Parameters for the Diagnosis

of Dupuytren's Disease. In: Berger A, Delbruck A, Brenner P, Hinzmann B, editors. Dupuytren's

Disease Pathobiochemistry and Clinical Management. Berlin: Springer-Verlag; 1994. p. 94–8.

53.   Sand JMB, Lamy P, Juhl P, Siebuhr AS, Iversen LV, Nawrocki A, et al. Development of a

Neo-Epitope Specific Assay for Serological Assessment of Type VII Collagen Turnover and Its

Relevance in Fibroproliferative Disorders. Assay Drug Dev Technol. 2018;16(2):123–31. Epub

20180301. doi: 10.1089/adt.2017.820. PubMed PMID: 29493258.

54.   Williams LM, McCann FE, Cabrita MA, Layton T, Cribbs A, Knezevic B, et al. Identifying

collagen VI as a target of fibrotic diseases regulated by CREBBP/EP300. Proc Natl Acad Sci U

S A. 2020;117(34):20753–63. Epub 2020/08/08. doi: 10.1073/pnas.2004281117. PubMed PMID:

32759223; PubMed Central PMCID: PMCPMC7456151.

55.   Rehman S, Salway F, Stanley JK, Ollier WE, Day P, Bayat A. Molecular phenotypic

descriptors of Dupuytren's disease defined using informatics analysis of the transcriptome. J

Hand Surg Am. 2008;33(3):359–72. Epub 2008/03/18. doi: 10.1016/j.jhsa.2007.11.010. PubMed

PMID: 18343292.

56.   Vi L, Feng L, Zhu RD, Wu Y, Satish L, Gan BS, et al. Periostin differentially induces

proliferation, contraction and apoptosis of primary Dupuytren's disease and adjacent palmar

fascia cells. Exp Cell Res. 2009;315(20):3574–86. Epub 2009/07/22. doi:

10.1016/j.yexcr.2009.07.015. PubMed PMID: 19619531; PubMed Central PMCID:

PMCPMC5017872.

57.   Krause C, Kloen P, Ten Dijke P. Elevated transforming growth factor beta and mitogen-

activated protein kinase pathways mediate fibrotic traits of Dupuytren's disease fibroblasts.

Fibrogenesis Tissue Repair. 2011;4(1):14. Epub 2011/06/30. doi: 10.1186/1755-1536-4-14.

PubMed PMID: 21711521; PubMed Central PMCID: PMCPMC3148569.

58.   Forrester HB, Temple-Smith P, Ham S, de Kretser D, Southwick G, Sprung CN.

Genome-wide analysis using exon arrays demonstrates an important role for expression of

extra-cellular matrix, fibrotic control and tissue remodelling genes in Dupuytren's disease. PLoS

One. 2013;8(3):e59056. Epub 2013/04/05. doi: 10.1371/journal.pone.0059056. PubMed PMID:

23554969; PubMed Central PMCID: PMCPMC3595223.

59.   Ng M, Thakkar D, Southam L, Werker P, Ophoff R, Becker K, et al. A Genome-wide

Association Study of Dupuytren Disease Reveals 17 Additional Variants Implicated in Fibrosis.

Am J Hum Genet. 2017;101(3):417–27. Epub 2017/09/09. doi: 10.1016/j.ajhg.2017.08.006.

PubMed PMID: 28886342; PubMed Central PMCID: PMCPMC5591021.

60.   Park TH, Kim D, Lee YS, Kim SY. A meta-analysis to identify novel diagnostic and

therapeutic targets for Dupuytren's disease. Wound Repair Regen. 2020;28(2):202–10. Epub

2019/11/07. doi: 10.1111/wrr.12774. PubMed PMID: 31688987.

61.   Verjee LS, Verhoekx JS, Chan JK, Krausgruber T, Nicolaidou V, Izadi D, et al. Unraveling

the signaling pathways promoting fibrosis in Dupuytren's disease reveals TNF as a therapeutic

target. Proc Natl Acad Sci U S A. 2013;110(10):E928–37. Epub 2013/02/23. doi:

10.1073/pnas.1301100110. PubMed PMID: 23431165; PubMed Central PMCID:

PMCPMC3593900.

62.   Eaton C. Dupuytren Disease. In: Wolfe S, Pederson W, Kozin S, Cohen M, editors.

Green's Operative Hand Surgery. 1. Eighth ed. Philadelphia: Elsevier; 2022. p. 148–81.

63.   Akbar M, Garcia-Melchor E, Chilaka S, Little KJ, Sood S, Reilly JH, et al. Attenuation of

Dupuytren's fibrosis via targeting of the STAT1 modulated IL-13Ralpha1 response. Sci Adv.

2020;6(28):eaaz8272. Epub 2020/07/23. doi: 10.1126/sciadv.aaz8272. PubMed PMID:

32695877; PubMed Central PMCID: PMCPMC7351483.

64.   Kida H, Jiang JJ, Matsui Y, Takahashi I, Hasebe R, Kawamura D, et al. Dupuytren's

contracture-associated SNPs increase SFRP4 expression in nonimmune cells including

fibroblasts to enhance inflammation development. Int Immunol. 2023. Epub 20230131. doi:

10.1093/intimm/dxad004. PubMed PMID: 36719100.

65.   Riesmeijer SA, Kamali Z, Ng M, Drichel D, Piersma B, Becker K, et al. A genome-wide

association meta-analysis implicates Hedgehog and Notch signaling in Dupuytren's disease.

Nat Commun. 2024;15(1):199. Epub 20240103. doi: 10.1038/s41467-023-44451-0. PubMed

PMID: 38172110; PubMed Central PMCID: PMCPMC10764787.

66.   Schultz HS, Nitze LM, Zeuthen LH, Keller P, Gruhler A, Pass J, et al. Collagen induces

maturation of human monocyte-derived dendritic cells by signaling through osteoclast-

associated receptor. J Immunol. 2015;194(7):3169–79. Epub 20150227. doi:

10.4049/jimmunol.1402800. PubMed PMID: 25725106; PubMed Central PMCID:

PMCPMC4367160.

67.   Bakowska JC, Jupille H, Fatheddin P, Puertollano R, Blackstone C. Troyer syndrome

protein spartin is mono-ubiquitinated and functions in EGF receptor trafficking. Mol Biol Cell.

2007;18(5):1683–92. Epub 20070301. doi: 10.1091/mbc.e06-09-0833. PubMed PMID:

17332501; PubMed Central PMCID: PMCPMC1855030.

68.   Augoff K, Tabola R, Kula J, Gosk J, Rutowski R. Epidermal growth factor receptor (EGF-

R) in Dupuytren's disease. J Hand Surg Br. 2005;30(6):570–3. Epub 2005/08/02. doi:

10.1016/j.jhsb.2005.06.008. PubMed PMID: 16055243.

69.   Wang ML, Rajpar I, Ruggiero NA, Fertala J, Steplewski A, Beredjiklian PK, et al.

Circulating inflammatory cytokines alter transcriptional activity within fibrotic tissue of

Dupuytren's disease patients. J Orthop Res. 2021. Epub 2021/04/30. doi: 10.1002/jor.25059.

PubMed PMID: 33913534.

70.   Wilbrand S. Dupuytren´s Contracture : Features and Consequences. Uppsala: Acta

Universitatis Upsaliensis; 2002.

71.   Kuo RYL, Ng M, Prieto-Alhambra D, Furniss D. Dupuytren's Disease Predicts Increased

All-Cause and Cancer-Specific Mortality: Analysis of a Large Cohort from the U.K. Clinical

Practice Research Datalink. Plast Reconstr Surg. 2020;145(3):574e–82e. Epub 2020/02/26. doi:

10.1097/PRS.0000000000006551. PubMed PMID: 32097318; PubMed Central PMCID:

PMCPMC7043723.

72.   van Rijssen AL, ter Linden H, Werker PM. Five-year results of a randomized clinical trial

on treatment in Dupuytren's disease: percutaneous needle fasciotomy versus limited

fasciectomy. Plast Reconstr Surg. 2012;129(2):469–77. Epub 2011/10/12. doi:

10.1097/PRS.0b013e31823aea95. PubMed PMID: 21987045.

73.   Wilbrand S, Ekbom A, Gerdin B. The sex ratio and rate of reoperation for Dupuytren's

contracture in men and women. J Hand Surg Br. 1999;24(4):456–9. Epub 1999/09/03. doi:

10.1054/jhsb.1999.0154. PubMed PMID: 10473157.

74.   Kozanoglu E, Mete FS, Akalin BE, Aksoyler D, Berkoz O, Emekli U, et al. The

epidemiological, etiological, and clinical comparisons of primary and recurrent Dupuytren's

contractures. Ulus Travma Acil Cerrahi Derg. 2025;31(2):133–9. doi: 10.14744/tjtes.2024.95049.

PubMed PMID: 39963920; PubMed Central PMCID: PMCPMC11843423.

75.   Hauksson IT, Havdal MB, Benth JS, Hoelsbrekken SE, Randsborg PH. Collagenase

Clostridium histolyticum Versus Needle Fasciotomy for Primary Metacarpophalangeal

Dupuytren Contracture: Five-Year Results from a Randomized Controlled Trial. JB JS Open

Access. 2024;9(4). Epub 20241120. doi: 10.2106/JBJS.OA.24.00038. PubMed PMID:

39569009; PubMed Central PMCID: PMCPMC11573333.

76.   Karjalainen VL, Soikkeli J, Raisanen MP, Leppanen OV, Reito A, Stjernberg-Salmela S,

et al. Progression of Dupuytren's contracture: a randomized controlled trial comparing surgery,

needle fasciotomy, and collagenase injection. Plast Reconstr Surg. 2025. Epub 20251204. doi:

10.1097/PRS.0000000000012666. PubMed PMID: 41334972.

77.   Raykha C, Crawford J, Gan BS, Fu P, Bach LA, O'Gorman DB. IGF-II and IGFBP-6

regulate cellular contractility and proliferation in Dupuytren's disease. Biochim Biophys Acta.

2013;1832(10):1511–9. Epub 2013/04/30. doi: 10.1016/j.bbadis.2013.04.018. PubMed PMID:

23623986.

78.   O'Gorman DB. The Extracellular Matrix in Dupuytren Disease. Dupuytren Disease and

Related Diseases - The Cutting Edge 2017. p. 43–54.

79.   Pannecoeck R, Serruys D, Benmeridja L, Delanghe JR, van Geel N, Speeckaert R, et al.

Vascular adhesion protein-1: Role in human pathology and application as a biomarker. Crit Rev

Clin Lab Sci. 2015;52(6):284–300. Epub 20150818. doi: 10.3109/10408363.2015.1050714.

PubMed PMID: 26287391.

80.   Yepuri G, Velagapudi S, Xiong Y, Rajapakse AG, Montani JP, Ming XF, et al. Positive

crosstalk between arginase-II and S6K1 in vascular endothelial inflammation and aging. Aging

Cell. 2012;11(6):1005–16. Epub 20120918. doi: 10.1111/acel.12001. PubMed PMID: 22928666.

81.   Volz J, Kusch C, Beck S, Popp M, Vogtle T, Meub M, et al. BIN2 orchestrates platelet

calcium signaling in thrombosis and thrombo-inflammation. J Clin Invest. 2020;130(11):6064–

79. doi: 10.1172/JCI136457. PubMed PMID: 32750041; PubMed Central PMCID:

PMCPMC7598067.

82.   Eaton C. The Next Stage of Clinical Dupuytren Research: Biomarkers and Chronic

Disease Research Tools. In: al. PMNWe, editor. Dupuytren Disease and Related Diseases - The

Cutting Edge. Switzerland: Springer; 2017. p. 391–407.

83.   Yanase Y, Matsuo Y, Takahagi S, Kawaguchi T, Uchida K, Ishii K, et al. Coagulation

factors induce human skin mast cell and basophil degranulation via activation of complement 5

and the C5a receptor. J Allergy Clin Immunol. 2021;147(3):1101–4 e7. Epub 20200902. doi:

10.1016/j.jaci.2020.08.018. PubMed PMID: 32888945.

84.   Romejko K, Markowska M, Niemczyk S. The Review of Current Knowledge on

Neutrophil Gelatinase-Associated Lipocalin (NGAL). Int J Mol Sci. 2023;24(13). Epub

20230621. doi: 10.3390/ijms241310470. PubMed PMID: 37445650; PubMed Central PMCID:

PMCPMC10341718.

85.   Verhoekx JS, Beckett KS, Bisson MA, McGrouther DA, Grobbelaar AO, Mudera V. The

mechanical environment in Dupuytren's contracture determines cell contractility and associated

MMP-mediated matrix remodeling. J Orthop Res. 2013;31(2):328–34. Epub 2012/09/19. doi:

10.1002/jor.22220. PubMed PMID: 22987740.

86.   Hammerman M, Blomgran P, Dansac A, Eliasson P, Aspenberg P. Different gene

response to mechanical loading during early and late phases of rat Achilles tendon healing. J

Appl Physiol (1985). 2017;123(4):800–15. Epub 20170713. doi:

10.1152/japplphysiol.00323.2017. PubMed PMID: 28705996.

87.   Stocks M, Walter AS, Akova E, Gauglitz G, Aszodi A, Boecker W, et al. RNA-seq

unravels distinct expression profiles of keloids and Dupuytren's disease. Heliyon.

2024;10(1):e23681. Epub 20231213. doi: 10.1016/j.heliyon.2023.e23681. PubMed PMID:

38187218; PubMed Central PMCID: PMCPMC10770622.

88.   Wilutzky B, Berndt A, Katenkamp D, Koshmehl H. Programmed cell death in nodular

palmar fibromatosis (Morbus Dupuytren). Histol Histopathol. 1998;13(1):67–72. Epub

1998/02/26. doi: 10.14670/HH-13.67. PubMed PMID: 9476635.

89.   Tanaka T, Biancotto A, Moaddel R, Moore AZ, Gonzalez-Freire M, Aon MA, et al. Plasma

proteomic signature of age in healthy humans. Aging Cell. 2018;17(5):e12799. Epub

2018/07/12. doi: 10.1111/acel.12799. PubMed PMID: 29992704; PubMed Central PMCID:

PMCPMC6156492.

90.   Park TJ, Curran T. Essential roles of Crk and CrkL in fibroblast structure and motility.

Oncogene. 2014;33(43):5121–32. Epub 20131028. doi: 10.1038/onc.2013.453. PubMed PMID:

24166500.

91.   Kanazawa T, Michida H, Uchino Y, Ishihara A, Zhang S, Tabata S, et al. Cell shape-

based chemical screening reveals an epigenetic network mediated by focal adhesions. FEBS J.

2021;288(19):5613–28. Epub 20210422. doi: 10.1111/febs.15840. PubMed PMID: 33768715.

92.   van Beuge MM, Ten Dam EJ, Werker PM, Bank RA. Wnt pathway in Dupuytren disease:

connecting profibrotic signals. Transl Res. 2015;166(6):762–71 e3. Epub 2015/10/17. doi:

10.1016/j.trsl.2015.09.006. PubMed PMID: 26470681.

93.   Satish L, LaFramboise WA, Johnson S, Vi L, Njarlangattil A, Raykha C, et al. Fibroblasts

from phenotypically normal palmar fascia exhibit molecular profiles highly similar to fibroblasts

from active disease in Dupuytren's Contracture. BMC Med Genomics. 2012;5:15. Epub

2012/05/09. doi: 10.1186/1755-8794-5-15. PubMed PMID: 22559715; PubMed Central PMCID:

PMCPMC3375203.

94.   Verhoekx JSN, Verjee LS, Izadi D, Chan JKK, Nicolaidou V, Davidson D, et al. Isometric

contraction of Dupuytren's myofibroblasts is inhibited by blocking intercellular junctions. J Invest

Dermatol. 2013;133(12):2664–71. Epub 2013/05/09. doi: 10.1038/jid.2013.219. PubMed PMID:

23652794.

95.   Moch M, Schieren J, Leube RE. Cortical tension regulates desmosomal morphogenesis.

Front Cell Dev Biol. 2022;10:946190. Epub 20221004. doi: 10.3389/fcell.2022.946190. PubMed

PMID: 36268507; PubMed Central PMCID: PMCPMC9577410.

96.   Ryu YH, Lee YJ, Kim KJ, Lee SJ, Han YN, Rhie JW. Epidermal Growth Factor (EGF)-

Like Repeats and Discoidin I-Like Domains 3 (EDIL3): A Potential New Therapeutic Tool for the

Treatment of Keloid Scars. Tissue Eng Regen Med. 2017;14(3):267–77. Epub 20170407. doi:

10.1007/s13770-017-0034-5. PubMed PMID: 30603483; PubMed Central PMCID:

PMCPMC6171597.

97.   Ferrero GB, Howald C, Micale L, Biamino E, Augello B, Fusco C, et al. An atypical

7q11.23 deletion in a normal IQ Williams-Beuren syndrome patient. Eur J Hum Genet.

2010;18(1):33–8. doi: 10.1038/ejhg.2009.108. PubMed PMID: 19568270; PubMed Central

PMCID: PMCPMC2987159.

98.   Millesi H, Reihsner R, Hamilton G, Mallinger R, Menzel EJ. Biomechanical properties of

normal tendons, normal palmar aponeuroses and palmar aponeuroses from patients with

Dupuytren's disease subjected to elastase and chondroitinase treatment. Connect Tissue Res.

1995;31(2):109–15. doi: 10.3109/03008209509028398. PubMed PMID: 15612326.

99.   Deng L, Feng J, Broaddus RR. The novel estrogen-induced gene EIG121 regulates

autophagy and promotes cell survival under stress. Cell Death Dis. 2010;1(4):e32. doi:

10.1038/cddis.2010.9. PubMed PMID: 21072319; PubMed Central PMCID: PMCPMC2976047.

100.   Jemec B, Grobbelaar AO, Wilson GD, Smith PJ, Sanders R, McGrouther DA. Is

Dupuytren's disease caused by an imbalance between proliferation and cell death? J Hand

Surg Br. 1999;24(5):511–4. Epub 1999/12/22. doi: 10.1054/jhsb.1999.0251. PubMed PMID:

10597921.

101.   Kaku M, Yamauchi M. Mechano-regulation of collagen biosynthesis in periodontal

ligament. J Prosthodont Res. 2014;58(4):193–207. Epub 2014/10/15. doi:

10.1016/j.jpor.2014.08.003. PubMed PMID: 25311991; PubMed Central PMCID:

PMCPMC4253671.

102.   Li T, Qiu J, Jia T, Liang Y, Zhang K, Yan W, et al. G3BP2 regulates oscillatory shear

stress-induced endothelial dysfunction. Genes Dis. 2022;9(6):1701–15. Epub 20211119. doi:

10.1016/j.gendis.2021.11.003. PubMed PMID: 36157502; PubMed Central PMCID:

PMCPMC9485288.

103.   Wei SC, Fattet L, Tsai JH, Guo Y, Pai VH, Majeski HE, et al. Matrix stiffness drives

epithelial-mesenchymal transition and tumour metastasis through a TWIST1-G3BP2

mechanotransduction pathway. Nat Cell Biol. 2015;17(5):678–88. Epub 20150420. doi:

10.1038/ncb3157. PubMed PMID: 25893917; PubMed Central PMCID: PMCPMC4452027.

104.   Modjeski KL, Ture SK, Field DJ, Cameron SJ, Morrell CN. Glutamate Receptor

Interacting Protein 1 Mediates Platelet Adhesion and Thrombus Formation. PLoS One.

2016;11(9):e0160638. Epub 20160915. doi: 10.1371/journal.pone.0160638. PubMed PMID:

27631377; PubMed Central PMCID: PMCPMC5025166.

105.   Tomatsu S, Montano AM, Dung VC, Grubb JH, Sly WS. Mutations and polymorphisms in

GUSB gene in mucopolysaccharidosis VII (Sly Syndrome). Hum Mutat. 2009;30(4):511–9. doi:

10.1002/humu.20828. PubMed PMID: 19224584; PubMed Central PMCID: PMCPMC3048808.

106.   Chakraborty A, Edkins AL. HSP90 as a regulator of extracellular matrix dynamics.

Biochem Soc Trans. 2021;49(6):2611–25. doi: 10.1042/BST20210374. PubMed PMID:

34913470.

107.   de la Mare JA, Jurgens T, Edkins AL. Extracellular Hsp90 and TGFbeta regulate

adhesion, migration and anchorage independent growth in a paired colon cancer cell line model.

BMC Cancer. 2017;17(1):202. Epub 20170316. doi: 10.1186/s12885-017-3190-z. PubMed

PMID: 28302086; PubMed Central PMCID: PMCPMC5356307.

108.   Terzuoli E, Corti F, Nannelli G, Giachetti A, Donnini S, Ziche M. Bradykinin B2 Receptor

Contributes to Inflammatory Responses in Human Endothelial Cells by the Transactivation of

the Fibroblast Growth Factor Receptor FGFR-1. Int J Mol Sci. 2018;19(9). Epub 20180906. doi:

10.3390/ijms19092638. PubMed PMID: 30200598; PubMed Central PMCID:

PMCPMC6163484.

109.   Vandersleyen V, Grosber M, Wilgenhof S, De Kock J, Neyns B, Gutermuth J.

Vemurafenib-associated Dupuytren- and Ledderhose palmoplantar fibromatosis in metastatic

melanoma patients. J Eur Acad Dermatol Venereol. 2016;30(7):1133–5. Epub 20150824. doi:

10.1111/jdv.13268. PubMed PMID: 26303964.

110.   Higuchi Y, Ogata T, Nakanishi N, Nishi M, Tsuji Y, Tomita S, et al. Cavin-2 promotes

fibroblast-to-myofibroblast trans-differentiation and aggravates cardiac fibrosis. ESC Heart Fail.

2023. Epub 20231024. doi: 10.1002/ehf2.14571. PubMed PMID: 37872863.

111.   Gokel JM, Hübner G. Occurrence of Myofibroblasts in the Different Phases of Morbus

Dupuytren (Dupuytren's Contracture). Beiträge zur Pathologie. 1977;161(2):166–75. doi:

10.1016/s0005-8165(77)80095-4.

112.   Chothani S, Schafer S, Adami E, Viswanathan S, Widjaja AA, Langley SR, et al.

Widespread Translational Control of Fibrosis in the Human Heart by RNA-Binding Proteins.

Circulation. 2019;140(11):937–51. Epub 20190709. doi:

10.1161/CIRCULATIONAHA.119.039596. PubMed PMID: 31284728; PubMed Central PMCID:

PMCPMC6749977.

113.   Drazic A, Aksnes H, Marie M, Boczkowska M, Varland S, Timmerman E, et al. NAA80 is

actin's N-terminal acetyltransferase and regulates cytoskeleton assembly and cell motility. Proc

Natl Acad Sci U S A. 2018;115(17):4399–404. Epub 20180326. doi: 10.1073/pnas.1718336115.

PubMed PMID: 29581253; PubMed Central PMCID: PMCPMC5924898.

114.   Bergenudd H, LindgÄRde F, Nilsson BE. Prevalence of Dupuytren's Contracture and its

Correlation with Degenerative Changes of the Hands and Feet and with Criteria of General

Health. Journal of Hand Surgery. 2016;18(2):254–7. doi: 10.1016/0266-7681(93)90123-w.

115.   Flint MH. The genesis of the palmar lesion. In: McFarlane RM, McGrouther DA, Flint MH,

editors. Dupuytren's Disease Biology and Treatment. Edinburg Churchill Livingstone; 1990.

116.   Worku MG, Seretew WS, Angaw DA, Tesema GA. Prevalence and Associated Factor of

Brown Adipose Tissue: Systematic Review and Meta-Analysis. Biomed Res Int.

2020;2020:9106976. Epub 20200616. doi: 10.1155/2020/9106976. PubMed PMID: 32685543;

PubMed Central PMCID: PMCPMC7317326.

117.   Hamamoto H, Ueba Y, Sudo Y, Sanada H, Yamamuro T, Takeda T. Dupuytren's

contracture: morphological and biochemical changes in palmar aponeurosis. Hand.

1982;14(3):237–47. Epub 1982/10/01. doi: 10.1016/s0072-968x(82)80056-9. PubMed PMID:

6130030.

118.   Shih B, Brown JJ, Armstrong DJ, Lindau T, Bayat A. Differential gene expression

analysis of subcutaneous fat, fascia, and skin overlying a Dupuytren's disease nodule in

comparison to control tissue. Hand (N Y). 2009;4(3):294–301. Epub 2009/02/03. doi:

10.1007/s11552-009-9164-0. PubMed PMID: 19184239; PubMed Central PMCID:

PMCPMC2724615.

119.   Tripkovic I, Ogorevc M, Vukovic D, Saraga-Babic M, Mardesic S. Fibrosis-Associated

Signaling Molecules Are Differentially Expressed in Palmar Connective Tissues of Patients with

Carpal Tunnel Syndrome and Dupuytren's Disease. Biomedicines. 2022;10(12). Epub

20221211. doi: 10.3390/biomedicines10123214. PubMed PMID: 36551969; PubMed Central

PMCID: PMCPMC9775445.

120.   Docheva D, Vaerst B, Deiler S, Giunta RE, Schieker M, Volkmer E. [Alterations of cell

phenotype in Dupuytren's disease--an in vitro analysis]. Handchir Mikrochir Plast Chir.

2012;44(2):59–66. Epub 2012/04/13. doi: 10.1055/s-0032-1309025. PubMed PMID: 22495955.

121.   Devine MJ, Birsa N, Kittler JT. Miro sculpts mitochondrial dynamics in neuronal health

and disease. Neurobiol Dis. 2016;90:27–34. Epub 20151219. doi: 10.1016/j.nbd.2015.12.008.

PubMed PMID: 26707701.

122.   Garcia-Martinez I, Garcia-Mesa Y, Garcia-Piqueras J, Martinez-Pubil A, Cobo JL, Feito J,

et al. Sensory innervation of the human palmar aponeurosis in healthy individuals and patients

with palmar fibromatosis. J Anat. 2021. Epub 20211208. doi: 10.1111/joa.13609. PubMed PMID:

34881452.

123.   Nanchahal J, Chan JK. Treatments for early-stage Dupuytren's disease: an evidence-

based approach. J Hand Surg Eur Vol. 2023:17531934221131373. Epub 20230113. doi:

10.1177/17531934221131373. PubMed PMID: 36638105.

124.   Joshi H, Morley SC. Cells under stress: The mechanical environment shapes

inflammasome responses to danger signals. J Leukoc Biol. 2019;106(1):119–25. Epub

20190115. doi: 10.1002/JLB.3MIR1118-417R. PubMed PMID: 30645000; PubMed Central

PMCID: PMCPMC6784827.

125.   Ali MK, Kim RY, Brown AC, Donovan C, Vanka KS, Mayall JR, et al. Critical role for iron

accumulation in the pathogenesis of fibrotic lung disease. J Pathol. 2020;251(1):49–62. Epub

20200330. doi: 10.1002/path.5401. PubMed PMID: 32083318.

126.   Ushijima M, Tsuneyoshi M, Enjoji M. Dupuytren type fibromatoses. A clinicopathologic

study of 62 cases. Acta Pathol Jpn. 1984;34(5):991–1001. Epub 1984/09/01. doi:

10.1111/j.1440-1827.1984.tb07630.x. PubMed PMID: 6507097.

127.   Verjee LS, Midwood K, Davidson D, Essex D, Sandison A, Nanchahal J. Myofibroblast

distribution in Dupuytren's cords: correlation with digital contracture. J Hand Surg Am.

2009;34(10):1785–94. Epub 2009/11/17. doi: 10.1016/j.jhsa.2009.08.005. PubMed PMID:

19910144.

128.   Kischer CW, Speer DP. Microvascular changes in Dupuytren's contracture. J Hand Surg

Am. 1984;9A(1):58–62. doi: 10.1016/s0363-5023(84)80185-9. PubMed PMID: 6693745.

129.   Aarabi S, Bhatt KA, Shi Y, Paterno J, Chang EI, Loh SA, et al. Mechanical load initiates

hypertrophic scar formation through decreased cellular apoptosis. FASEB J. 2007;21(12):3250–

61. Epub 2007/05/17. doi: 10.1096/fj.07-8218com. PubMed PMID: 17504973.

130.   Sood RF, Hocking AM, Muffley LA, Ga M, Honari S, Reiner AP, et al. Genome-wide

Association Study of Postburn Scarring Identifies a Novel Protective Variant. Ann Surg.

2015;262(4):563–9. doi: 10.1097/SLA.0000000000001439. PubMed PMID: 26366535; PubMed

Central PMCID: PMCPMC9632222.

131.   Viji Babu PK, Rianna C, Belge G, Mirastschijski U, Radmacher M. Mechanical and

migratory properties of normal, scar, and Dupuytren's fibroblasts. J Mol Recognit.

2018;31(9):e2719. Epub 2018/04/28. doi: 10.1002/jmr.2719. PubMed PMID: 29701269.

132.   Teckchandani A, Toida N, Goodchild J, Henderson C, Watts J, Wollscheid B, et al.

Quantitative proteomics identifies a Dab2/integrin module regulating cell migration. J Cell Biol.

2009;186(1):99–111. Epub 20090706. doi: 10.1083/jcb.200812160. PubMed PMID: 19581412;

PubMed Central PMCID: PMCPMC2712992.

133.   Johnston P, Chojnowski AJ, Davidson RK, Riley GP, Donell ST, Clark IM. A complete

expression profile of matrix-degrading metalloproteinases in Dupuytren's disease. J Hand Surg

Am. 2007;32(3):343–51. Epub 2007/03/06. doi: 10.1016/j.jhsa.2006.12.010. PubMed PMID:

17336841.

134.   D'Amico F, Candido S, Libra M. Interaction between matrix metalloproteinase-9 (MMP-9)

and neutrophil gelatinase-associated lipocalin (NGAL): A recent evolutionary event in primates.

Dev Comp Immunol. 2021;116:103933. Epub 20201124. doi: 10.1016/j.dci.2020.103933.

PubMed PMID: 33245981.

135.   Luo L, Deng L, Chen Y, Ding R, Li X. Identification of Lipocalin 2 as a Ferroptosis-

Related Key Gene Associated with Hypoxic-Ischemic Brain Damage via STAT3/NF-kappaB

Signaling Pathway. Antioxidants (Basel). 2023;12(1). Epub 20230112. doi:

10.3390/antiox12010186. PubMed PMID: 36671050; PubMed Central PMCID:

PMCPMC9854551.

136.   Wipff PJ, Hinz B. Integrins and the activation of latent transforming growth factor beta1 -

an intimate relationship. Eur J Cell Biol. 2008;87(8-9):601–15. Epub 2008/03/18. doi:

10.1016/j.ejcb.2008.01.012. PubMed PMID: 18342983.

137.   Dong G, Huang X, Jiang S, Ni L, Chen S. Simvastatin Mitigates Apoptosis and

Transforming Growth Factor-Beta Upregulation in Stretch-Induced Endothelial Cells. Oxid Med

Cell Longev. 2019;2019:6026051. Epub 20191217. doi: 10.1155/2019/6026051. PubMed PMID:

31934265; PubMed Central PMCID: PMCPMC6942893.

138.   Lyons RM, Gentry LE, Purchio AF, Moses HL. Mechanism of activation of latent

recombinant transforming growth factor beta 1 by plasmin. J Cell Biol. 1990;110(4):1361–7. doi:

10.1083/jcb.110.4.1361. PubMed PMID: 2139036; PubMed Central PMCID: PMCPMC2116088.

139.   Flevaris P, Vaughan D. The Role of Plasminogen Activator Inhibitor Type-1 in Fibrosis.

Semin Thromb Hemost. 2017;43(2):169–77. Epub 20160824. doi: 10.1055/s-0036-1586228.

PubMed PMID: 27556351.

140.   Howard JC, Varallo VM, Ross DC, Faber KJ, Roth JH, Seney S, et al. Wound healing-

associated proteins Hsp47 and fibronectin are elevated in Dupuytren's contracture. J Surg Res.

2004;117(2):232–8. Epub 2004/03/30. doi: 10.1016/j.jss.2004.01.013. PubMed PMID:

15047128.

141.   Hosokawa N, Hohenadl C, Satoh M, Kuhn K, Nagata K. HSP47, a collagen-specific

molecular chaperone, delays the secretion of type III procollagen transfected in human

embryonic kidney cell line 293: a possible role for HSP47 in collagen modification. J Biochem.

1998;124(3):654–62. doi: 10.1093/oxfordjournals.jbchem.a022162. PubMed PMID: 9722680.

142.   Li Z, Ivanov AA, Su R, Gonzalez-Pecchi V, Qi Q, Liu S, et al. The OncoPPi network of

cancer-focused protein-protein interactions to inform biological insights and therapeutic

strategies. Nat Commun. 2017;8:14356. Epub 20170216. doi: 10.1038/ncomms14356. PubMed

PMID: 28205554; PubMed Central PMCID: PMCPMC5316855.

143.   van den Broek LJ, van der Veer WM, de Jong EH, Gibbs S, Niessen FB. Suppressed

inflammatory gene expression during human hypertrophic scar compared to normotrophic scar

formation. Exp Dermatol. 2015;24(8):623–9. Epub 20150526. doi: 10.1111/exd.12739. PubMed

PMID: 25939875.

1285

1286

	C&C	ECM	ENZ	MBR	PPL	SGN
ACAN		X				
AKR1A1		X				
BIN2		X				
C5	X	X				X
C6	X	X				X
CALCB						X
CASP3			X			
CGA		X				X
CGB3		X				
CGB7		X				
CPB1			X			
CRKL				X	X	X
CSNK1G2			X			X
DSG2		X				X
EDIL3		X				
FKBP5					X	
GUSB			X			
HGS				X		X
HSP90AA1						X
KNG1	X	X	X			
MAP3K11						X
NRG4		X				
PCSK9				X		
PRKAR2B			X			X
RHOT1				X		
SERPINC1	X	X				
SOCS3		X			X	
SYK				X	X	X
TF				X	X	
USP8			X			
WFIKKN2			X			
YWHAB		X				

	CAN	C&C	ECM	INL	INF	MBR	MFB	PPL	SGN	WNT
ACAN			X							
CASP3	X		X	X	X		X	X	X	X
COL1A1	X	X	X		X	X	X	X	X	X
CSNK1G2								X		X
DAB2							X	X		X
DDR2			X				X	X		
HPX			X				X	X		
KNG1	X	X	X				X	X		X
LCN2				X						
MAP2K1	X		X	X	X	X	X	X	X	X
PLAT	X	X	X							
POSTN			X							
PRKCA	X	X	X		X	X	X	X	X	X
SERPINH1			X							
SFRP4							X	X		X
SMAD1							X	X		
STAT1	X		X		X		X	X	X	
STAT3	X		X		X		X	X	X	
TF	X	X					X	X		
USP8							X	X		
YWHAZ	X	X			X		X	X	X	

A

G3BP2 EIF4H NRG4 C6 SERPINC1 TF

C5 KNG1

POSTN FKBP5 HSP90AA1 CPB1 PRSS1

SYK CRKL GP1BB

SOCS3 MBNL2 MBNL1

ACAN MAP3K11 OSCAR GUSB

CASP3 AKR1A1

PCSK9 DSG2 YWHAB USP8 HGS CALCB CGA CGB3

B

C