Precipitation tolerance determines elevational migration strategies of resident birds of the
Western Ghats of India.

Authors: V.A. Akshay[1, 2, 6*], Caitlin J. Campbell[3], Bette Loiselle[4, 5], and Robert Guralnick[6]

1. School of Natural Resources and Environment, University of Florida, Gainesville, FL.

2. Biodiversity Institute, University of Florida, Gainesville, FL.

3. Bat Conservation International, Austin, TX.

4. Center for Latin American Studies, University of Florida, Gainesville, FL.

5. Department of Wildlife Ecology and Conservation, University of Florida, Gainesville,
FL.

6. Florida Museum of Natural History, University of Florida, Gainesville, FL.

*Corresponding author:
V.A. Akshay, Florida Museum of Natural History,
University of Florida, Gainesville, FL 32611
akshayvinodanand@ufl.edu
16

Acknowledgements

We would like to thank the citizen scientists and eBirders of India, without whom this work
would not be possible. We would also like to thank Dr. Michael Belitz and Dr. Scott Robinson
for methodological and theoretical discussion on this work. This work was partially funded by
the NSF: RCN: Cross-Scale Processes Impacting Biodiversity (DEB-1745562) and the
Biodiversity Institute at the University of Florida, for which we would like to thank Dr. Ana
Carnaval and Dr. Pamela Soltis respectively.

Data Accessibility Statement

Raw data was downloaded from eBird and can be accessed by visiting:
https://science.ebird.org/en/use-ebird-data/download-ebird-data-products. The source code and
data sets utilized in this study are available on https://zenodo.org/records/14291657.

Conflict of Interest Statement: The authors declare no conflict of interest.

Abstract:

Aim: Avian elevational migration is a common, but less studied form of short-distance migration. Numerous hypotheses have been proposed to explain this behavior, yet few studies have tested those comprehensively. We examine four non mutually exclusive hypotheses—climatic constraint, limited foraging opportunities, body size, and dispersal ability—using a comparative framework to evaluate their support in explaining avian elevational migration in the monsoon-dominant Western Ghats.

Location: Western Ghats.

Time period: 2013 - 2023.

Major taxa studied: Birds.

Methods: We used citizen science data from eBird to quantify elevational migration patterns of resident birds of the Western Ghats. Using robust data curation and new bias correction methods, we modeled migration patterns of 164 species across summer, monsoon and winter seasons. We built seasonal phylogenetic logistic regression models to test the effect of a species' environmental tolerances, diet, body-size and dispersal ability on elevational migration likelihood.

Results: 43% of resident birds exhibited elevational migration in at least one season. Most species migrated between winter and summer (57), followed by migrations between summer and monsoon (39) and then monsoon and winter (38). Species predominantly moved upslope in summer (82%) and downslope in monsoon (87%), with no discernible pattern in winter. Species with broader temperature tolerances were more likely to migrate across all seasons. Additionally, species associated with wetter environments were more likely to migrate in summer and winter. Lastly, species with narrower precipitation tolerances were more likely to migrate during the monsoon, influenced by diet.

Main conclusions: We find support for the climatic constraint and limited foraging opportunities hypotheses in explaining both the direction and likelihood of elevational migration. Precipitation is a particularly strong climatic driver for elevational migrants, especially under monsoon conditions. This suggests a complex relationship between precipitation tracking and the associated changes in resource availability determining elevational migration strategies in the region.

Key words: citizen science, climatic constraint hypothesis, eBird, elevational migrations, limited foraging opportunities hypothesis, precipitation, species traits.

Introduction

Animal migration is one of nature's most spectacular phenomena, yet we are just beginning to understand the patterns of migration and underlying eco-evolutionary dynamics that determine those patterns. Migration is thought to be favored when the fitness benefits of exploiting spatiotemporally varying resources outweigh the associated costs (Zurell et al., 2018). Migrating animals can take advantage of multiple habitats where suitability (e.g., survival, food availability, predation risk, and reproductive success) differs across seasons and regions (Hsiung et al., 2018; Milner-Gulland et al., 2013). Migration also involves significant costs, such as increased energy expenditure, heightened predation risk during transit, and potentially shorter breeding seasons due to time spent migrating (Alves et al., 2013; Boyle, 2017; Milner-Gulland et al., 2013). These trade-offs influence the evolution of migratory behaviors, where organisms optimize their fitness by balancing the rewards of resource acquisition with the risks and energy costs associated with movement (Bowlin et al., 2010; Shaw, 2016).

While some migrants travel very long distances, many make shorter journeys. Short distance migrants commonly perform seasonal back-and-forth movements between regions with altering favorable and unfavorable conditions, including breeding areas. Short distance migrants are often less time-constrained, thus able to focus on minimizing energetic costs during their seasonal transitions (Dingle & Drake, 2007; Rappole, 2013). Elevational migration is a form of short-distance migration where animals move between breeding and non-breeding grounds that differ in elevation in a cyclical annual pattern (Rappole, 2013). In birds, elevational migration is relatively common, with ~10% of species showing seasonal elevational shifts (Barçante et al., 2017). Although a majority (44%) of avian elevational migrants occur in the neotropics (Barçante et al., 2017), elevational migration occurs worldwide, with, ~ 20% of North American bird species (Boyle, 2017), 55 – 65 % of Himalayan species (Menon et al., 2023), and ~58% of Taiwanese species (Tsai et al., 2021) known to be elevational migrants.

The underlying intrinsic and extrinsic mechanisms explaining elevational migration in birds has been a focus of ornithological research, especially recently (Williamson & Witt, 2021). This has resulted in numerous non-exclusive hypotheses to explain these seasonal elevational shifts. We note four key ones that have garnered empirical support. The first is the 'climatic constraint hypothesis'; here seasonal climatic regimes can expose individuals to unsuitable temperatures, precipitation and wind speeds at their breeding elevations during the non-breeding season. Thus, imposing strong constraints on a species' ability to maintain optimal physiological conditions (Hsiung et al., 2018). To minimize these direct weather-driven physiological constraints, species can move to elevations with more suitable conditions. The central expectation of this hypothesis is that species with narrower tolerances to harsh weather conditions will move downslope during the winter season and upslope during the breeding season to track suitable climate conditions. Recent work by Tsai et al. (2021) and Menon et al. (2023) support this hypothesis and have demonstrated that Taiwanese and Himalayan birds that

are more cold intolerant make larger elevational migrations between their breeding and non-
breeding grounds.

Another hypothesis stems from formative work on elevational migrants in the neotropics,
where food availability has long been thought to be the driver of seasonal movements of birds
(Levey, 1988; Loiselle & Blake, 1991; Stiles, 1988). The limited foraging opportunities (LFO)
hypothesis suggests that during heavy wet-season storms, weather-driven reductions in foraging
opportunities, not intrinsic food availability, drive taxa with higher foraging needs to migrate to
avoid starvation (Boyle, 2008b). This hypothesis, unlike the climatic constraint hypothesis, does
not predict that species with narrower climatic tolerances are more likely to migrate. Initial tests
of the LFO hypothesis, on the White-ruffed Manakin (*Corapipo altera*) in Costa Rica showed
increased captures of elevational migrants at lower elevations following heavy rainfall (Boyle et
al., 2010). Community-level studies later confirmed that storms at higher elevations led to a
higher abundance of migrants in the lowlands, with body size and diet influencing responses
(Boyle, 2011). These studies suggest that species will shift locations to maximize foraging
success during periods of poor conditions, indicating an adaptive flexibility in response to
changing conditions. In sum, the LFO hypothesis posits an indirect effect of harsh weather on
elevational migrants, where species with the greatest foraging need are predicted to make
downslope movements during wet seasons to exploit improved foraging opportunities and
resource availability. Support exists for both the above hypotheses (i.e. "climate constraint" and
"LFO"), the explanatory power of which may depend on system-, season- and species-specific
factors. Using a comparative approach across multiple species can provide a means to determine
if one or the other (or both) hypotheses are most supported.

Two other hypotheses focus more on intrinsic morphological traits: The first is the 'body-
size hypothesis', which proposes that larger individuals can better endure severe weather and
scarce food resources, and thus should make shorter migrations from high-altitude breeding
grounds (Ketterson & Nolan, 1976). The second hypothesis focuses on dispersal ability
(Claramunt et al., 2012; Sheard et al., 2020), and posits that better dispersers are likely to show
stronger elevational migration. Still, the body of research conducted on avian elevational
migration remains limited to shifts between breeding and non-breeding grounds, with little
attention to movements over complete annual cycles (but see (Rueda-Uribe et al., 2024).
Transitional seasons, between breeding and non-breeding seasons, may pose particular
challenges for birds in highly seasonal, tropical systems. Improving our understanding of
elevational movements of birds during these transitional seasons could lend further support to
traditional hypotheses explaining this complex and understudied behavior.

A significant challenge in studying avian elevational migration over annual cycles is the
need for year-round monitoring over broad geographic extents. Citizen science initiatives, such
as eBird, can help overcome some of these challenges via assembling semi-structured checklists
in a common format submitted by an extensive network of citizen scientists (Kelling et al., 2019;
Sullivan et al., 2009). Still, these data resources, however extensively verified or complete,

contain sources of bias including uneven sampling over space and time (Sullivan et al., 2014). In the context of avian elevational migration, biases in citizen science data can skew results towards elevations where sampling has been more frequent. Researchers have employed various statistical methods to correct these biases, specifically resampling bird occurrences within elevational bands to obtain more uniform sampling across elevations (Menon et al., 2023; Tsai et al., 2021). Yet, there is scope for further improvement in correcting biases from semi-structured checklists, in particular to shift from case specific methods to a more generalizable framework.

In this study we used bird occurrence data obtained from the eBird portal to quantify elevational migration of resident birds in the Western Ghats of India and test key hypotheses explaining this migratory behavior. Our overall goals are to (1) quantify elevational migration distance and identify patterns of movement of resident birds of the Western Ghats over a complete annual seasonal cycle, and (2) test the effect of body size, environmental tolerances, dispersal ability and diet on a species' likelihood of being a seasonal elevational migrant, effectively evaluating the expectations of the hypotheses outlined above. These drivers are not mutually exclusive, but rather may interact to affect species' seasonal elevational movements. Thus, we expect species to demonstrate upslope movements in the summer and downslope movements in the winter season, consistent with the expectations of the "climatic constraint hypothesis." Further we expect that an interaction of the LFO and "climatic constraint" hypotheses will provide an explanatory framework for monsoon season elevational movements. Harsh monsoon conditions, with large amounts of rainfall and higher wind speeds, are most likely to limit foraging opportunities in the higher elevations of the Western Ghats. For this reason, we expect species with narrower tolerances to precipitation and wind speed to have a higher likelihood of being an elevational migrant in this season and will likely move towards elevations where resources are more available, i.e., to the lower reaches of the elevational gradient. Further we expect the temperature fluctuations associated with the summer and winter seasons to match the expectations of the "climatic constraint hypothesis." Thus, we expect species with narrower tolerances to temperature to have a higher likelihood of being a seasonal elevational migrant in these seasons. Lastly, we expect species with small body sizes and high dispersal ability to have a higher likelihood of being a seasonal elevational migrant, as per the "body-size" and "dispersal ability" hypotheses. In sum, we expect the four hypotheses outlined above to have varied, yet season- and clade-specific support in this study. Finally, we present a generalizable method for correcting spatial and temporal sampling biases in citizen science data and provide evidence of its effectiveness compared to previous methods.

Materials and Methods

Study site

The Western Ghats are a 2200 km long, ancient mountain range extending along the western coast of the Indian subcontinent, covering tropical latitudes of 8 N - 21 N (Das et al., 2006); and Fig 1a) and form a significant portion of the Western Ghats and Sri Lanka

biodiversity hotspot (Myers et al., 2000). These mountains are topographically, climatically and
biologically diverse (Gadgil & Meher - Homji, 1990, Pascal, 1988) and include an elevational
gradient of 40 - 2625 m (Fig 1a), with significant temperature variation between the plains and
higher elevations (Fig 1b) and lower temperature seasonality than the eastern parts of the
subcontinent (Fig 1c). This mountain range also experiences extreme monsoon conditions during
the months of June to October as a result of the annual south-west monsoon. The topography of
the Western Ghats interacts with the south-west monsoon to create a longitudinal rainfall
gradient (Fig 1d), with higher precipitation seasonality in the northern, lower, portions of these
hills as compared to the southern, higher parts (Fig 1e).

eBird occurrence records and data filtering.

We downloaded eBird checklist data from India during the time period from 2013 - 2023.
We first applied a spatial filter to these data to include checklists that fell within the bounds of
the Western Ghats. Next, we filtered these data to include records between March 2013 to March
2023, resulting in 10 full years of occurrence data. The ten years of data were aggregated for all
further analysis. We retained occurrence records associated with complete checklists, where
observer(s) recorded all identifiable species. To standardize sampling protocol, we included
checklists that followed the "stationary" or "traveling" protocols based on the distance moved by
the observer. To reduce the variation in sampling effort, checklists with a sampling duration of
>5 hours, sampling distance of >1 km and with >10 observers were excluded, following "Best
practices for Using eBird Data" (Strimas-Mackey et al., 2020) and Tsai et al. (2021).
Furthermore, the dataset was filtered to include only resident species belonging to the orders:
Passeriformes, *Bucerotiformes*, *Trogoniformes*, *Piciformes*, *Coraciiformes*, *Cuculiformes*,
Psittaciformes, and *Columbiformes*. We excluded raptors (*Accipitriformes* and *Falconidae*),
swifts (*Apodiformes*) and swallows (*Hirundinidae*) since these groups are generally observed in-
flight and associating elevation to these records is difficult (Menon et al., 2023; Ramesh et al.,
2022). Checklists were assigned a value for season based on the observation month (summer:
March – May; monsoon: June – October; Winter: November – February) and elevation of each
checklist was obtained from the ASTER Global Digital Elevation Model, at 30 m resolution
(Abrams et al., 2020). To account for geographic location uncertainty, elevations were averaged
using a 100 m buffer around each checklist location (Tsai et al., 2021) and rounded to two
decimal places. Elevation values and season categories were joined to each species' occurrence
record resulting in a long-format dataset of species occurrences with associated elevation and
season. All data filtering was performed in the programming environment R v. 4.3.3 (R Core
Team, 2024), using the packages 'auk' (Strimas-Mackey et al., 2023) and 'sf' (Pebesma, 2018).

Bias correction and method validation

A key challenge of eBird data is spatial sampling bias (Sullivan et al., 2014). eBird
checklists are often located closer to roads (Ramesh et al., 2022), lower elevations are sampled
more often than higher elevations (Menon et al., 2023; Tsai et al., 2021) and sampling is

seasonally biased as well. We examined the distribution of occurrence records and checklists as a
function of elevation and season to understand sampling effort across our dataset. Sampling was
uneven across elevations and seasons; the majority of occurrence records were reported from
elevations below 1200 m (Figure S1.1a) and sampling was significantly higher in the winter
when conditions are most pleasant (Figure S1.1b). This elevational and seasonal bias in reporting
has the potential to skew migration quantifications towards the lower elevations with more
robust estimates for the winter. To account for this bias, we developed a resampling approach
that helps correct for these issues. We first calculated the cumulative minutes of sampling for
each season/elevation combination in our filtered dataset as a measure of sampling effort. We
then fit Generalized Additive Models (GAMs; Hastie & Tibshirani, 1986) for each season to
predict sampling effort across the elevational range of occurrence records in our dataset. GAMs
were built using the 'mgcv' (Wood, 2017) package in the R programming environment. Next, we
resampled each species – season combination 20,000 times, with replacement, weighted by the
inverse of the predicted value of sampling effort. We used the inverse of sampling effort to up-
sample elevations with lower effort and down-sample elevations with higher effort. This
resampling approach provides continuous elevational distributions of each species per season
under more equal sampling effort. Using this bias adjusted distribution we extracted each
species' median (50th percentile) elevation per season. Species with fewer than 10 occurrence
records for any one season were excluded.

To test the effectiveness of our approach, we validated our method against the effort
correction protocol used in Tsai et al. (2021), who used a different resampling methodology.
Briefly, those authors binned sampling effort across elevational band and season and resampled
within those bins. In contrast, our approach smooths noise in sampling via fitted GAMs across a
continuous range of elevations. Further, we took care to reduce the effect of outliers in our
resampling approach by including species with at least 10 occurrence records in each season. We
replicated the analyses of Tsai et al. (2021) and then applied our method for bias correction on
their data to obtain estimates of median elevation for each species in their study. In comparison,
we found that our method produced consistent and reasonably conservative estimates of median
elevation (Figure S1.2 & S1.3) and has distinct advantages with respect to execution time (Figure
S1.4).

Quantifying elevational migration

We used our method for bias correction on the filtered bird occurrence data and extracted
median elevation values for all species in each season. We classified a species as an elevational
migrant if the 95% confidence intervals around the median elevation estimate were non-
overlapping between two consecutive seasons. This resulted in a binary variable for each season
denoting whether a species was an elevational migrant. We then calculated elevational migration
distance for a species as the difference between median elevation values between the focal and
the previous season. For example, the monsoon season elevational migration distance was
calculated as the difference between median elevation in the monsoon and median elevation in

the preceding season, summer (median$_{monsoon}$ – median$_{summer}$). This was repeated for all species
and consecutive seasons to arrive at three elevational migration distances per species (Table
S1.1).

Species trait data

We calculated each species' tolerance to temperature, precipitation and wind speed in our
filtered Western Ghats dataset. To do so, we used the mean monthly temperature, precipitation
and wind speed layers, downloaded from CHELSA (Karger et al., 2017), and masked them to the
bounds of the Western Ghats. We focused on calculating environmental tolerances within the
Western Ghats rather than across a species' global distribution because we were interested in a
species' regional adaptations to climate and their associated effect on elevational movements.
Next, for each occurrence point of a species in our dataset, we extracted temperature,
precipitation and wind speed values from the masked environmental layers. Due to a mismatch
between the time period of occurrence records in our dataset (2013 - 2023) and CHELSA data
availability (up to 2018), we used CHELSA layers from 2008–2018. We believe this mismatch
did not impact our calculations of species' environmental tolerances since we used
environmental data across ten years that were matched to our occurrence records to the best of
our ability. We calculated each species' 2.5th, 50th (median), and 97.5th percentile values for
temperature, precipitation and wind speed using the extracted occurrence specific environmental
data. The difference between the 97.5th and 2.5th percentile was considered as the species'
environmental tolerance range. Both the tolerance range and median were used as measures of a
species' environmental tolerance characteristics in downstream analyses.

Additionally, we extracted body mass (as a proxy for body size), diet preference and
Hand-Wing Index (HWI; as a proxy for dispersal ability) from the AVONET global dataset
(Tobias et al., 2022). HWI has been shown to be strongly associated with dispersal ability
(Sheard et al., 2020), with the expectation that species with long, pointed wings and high HWI
have a higher dispersal potential than species with short, rounded wings and low HWI
(Claramunt et al., 2012; Freeman et al., 2023).

Statistical analyses

To identify the patterns of movement of species in our study, we used a chi-square test to
evaluate the intra-seasonal variation in movement direction (upslope vs downslope). For each
season, we calculated the observed frequencies of direction for species demonstrating significant
elevational movement and compared these with expected frequencies of equal distribution
between direction classes. Chi-square tests were performed using the 'chisq.test' function in R.

We then fit seasonal phylogenetic logistic regression models (Ives & Garland, 2010) to
understand the effect of environmental tolerances, body mass, dispersal ability, and diet on a
species' elevational migration status. We modeled elevational migration status (migrant or non-
migrant) for a given season using a binomial distribution (see the "Quantifying elevational

migration" section above). We performed analyses separately for each season as our expectations were season specific. We employed a backward stepwise model selection approach to select the most parsimonious combination of predictor variables that explain the elevational migration status of a species in a season. We used a consensus Western Ghats bird tree as the basis for determining phylogenetic autocorrelation in our logistic regression models (built using Jetz et al., 2012). The phylogenetic logistic regression model was built using the 'phyloglm' function from the 'phylolm' package (Ho & Ane, 2014) in R and stepwise model selection was performed using the 'phyloglmstep' function from the same package. We also built seasonal Phylogenetic Generalized Least Squares (PGLS) regression models (Revell, 2010) to test the effect of our candidate predictor variables on absolute elevational migration distance (see Table S1.2 for more details).

Results

Patterns of elevational migration

We analyzed 946,282 occurrence records from 105,725 eBird checklists to quantify elevational migration of 164 resident bird species of the Western Ghats over three seasons. Our results revealed that 71 (43%) species showed significant elevational migration in at least one season. Summers had the highest number of elevational migrants with 57 (35%) species, of which 47 displayed upslope movement (Figure 2). This was followed by the monsoon with 39 (24%) species, of which 34 exhibited downslope movements (Figure 2). Winters had a marginally lower number of elevational migrants with 38 (23%) species and did not reveal a clear pattern of movement (Figure 2). Chi-square tests for the summer and monsoon seasons indicated a significant difference in migration direction, with more species demonstrating upslope movements in the summer and downslope movements in the monsoon than expected by chance (Table 1). However, chi-square test results for the winter did not show a statistically significant difference in movement direction (Table 1). We identified only 12 (7%) species that exhibited significant movement in all three seasons, however 34 species were common between the summer and monsoon followed by 26 species between the winter and summer, and 15 species between the monsoon and winter. In sum, our results show a consistent pattern of movement of elevational migrants between the three seasons, with the majority of species moving upslope in the summer and downslope in the monsoon, and no clear or dominant pattern of movement up or downslope in the winter (Figure 2; Figure S1.5).

Association between species traits and elevational migration

Our backward model selection approach for the seasonal phylogenetic logistic regression models shows that temperature and precipitation tolerance characteristics were the strongest predictors of elevational migration status in the summer and winter seasons (Figure 3; Table S1.3). Specifically, we found that summer and winter migration likelihood increased with broader temperature tolerances and an association with wetter conditions (Figure 3; Table S1.3). In contrast, monsoon season models revealed a different set of associations: elevational

migration status was positively associated with temperature tolerance, but negatively associated with precipitation tolerance (Figure 3; Table S1.3). We also found a significant effect of diet in the monsoon season. In particular, frugivorous species were less likely to be elevational migrants, indicating that migration strategies may be influenced by specific dietary needs under monsoon conditions. Lastly, we found no significant association between elevational migration status and wind speed tolerance characteristics, body-size or dispersal ability in any season (Table S1.3).

Elevational migration distance models fit under a PGLS framework, partially aligned with the above results. We found that species associated with wetter conditions performed larger elevational movements in all three seasons (Table S1.2). Additionally, species that moved to a larger elevational extent were associated with higher wind speeds in the monsoon and winter, and had a broader tolerance for temperature in the summer season (Table S1.2). However, these models generally exhibited only modest fit to our data based on adjusted R^2 scores (Table S1.4), and we chose to exclude these from further interpretation.

Discussion

This study quantified the elevational migration strategies of 164 resident bird species in the Western Ghats of India. Utilizing a comprehensive citizen science dataset and robust analytical methods, we identified patterns of elevational movement and tested hypotheses regarding this behavior along one of the world's most biodiverse elevational gradients. Our findings indicate that approximately 41% of resident birds exhibit seasonal elevational movements, a figure comparable to the 44% observed in other tropical regions (Barçante et al., 2017). We found significant directionality in these movements, with species predominantly migrating upslope during the breeding summer and downslope during the monsoon. These strongly consistent movements by nearly all species who make seasonal migrations fit expectations as we discuss below. We more directly tested hypotheses related to climatic constraints and, more indirectly, foraging, via phylogenetically informed logistic regression models. However, the predictions of some of the key hypotheses did not fully align with our results. Surprisingly, we found that species with broader tolerances to temperature were more likely to be elevational migrants, regardless of the season. Furthermore, these species are associated with generally wetter conditions during the summer and winter. Lastly, we found the monsoon season posed significant and unique challenges to elevational migrants. During this season, species with narrower tolerances to precipitation were more likely to be elevational migrants, and these species were less likely to be frugivores. These results suggest that depending on the specific question posed, the season in question, and dietary guild, either the "climatic constraint" or LFO hypotheses may more effectively explain the elevational migration strategies of resident birds in the region.

The first question we addressed is whether there are seasonal variations in the movement direction of elevational migrant species. We identified a clear pattern of movement during the

summer and monsoon seasons. In the breeding summer, the majority of elevational migrants
moved upslope from their wintering elevations. Breeding upslope migrations are well
documented in species worldwide (Boyle, 2017; Levey, 1988; Menon et al., 2023; Tsai et al.,
2021) and are expected under the "climatic constraint hypothesis." Furthermore, evidence
suggests that nest predation rates may be lower at higher elevations, as observed in the
neotropics (Boyle, 2008a; Skutch, 1985). Therefore, elevational migrants in the Western Ghats
may be moving upslope during the breeding summer season to elevations where breeding
success is likely to be higher. In the monsoon season, we observed a marked downslope
movement of resident birds from their breeding elevations. Heavy and prolonged rainfall - a
defining characteristic of the monsoon in the Western Ghats - is known to reduce foraging
abilities in small-bodied consumers like birds and bats (Boyle, 2010; Boyle et al., 2020; Foster,
1974; Voigt et al., 2011). Consequently, intense rainfall during the Western Ghats monsoon likely
limits foraging opportunities at higher elevations, prompting species to seek refuge at lower
elevations where resources are more readily available, consistent with studies examining the
elevational response of birds to heavy wet-season storms (Boyle, 2011, 2017; Boyle et al., 2010).
This analysis failed to identify a pattern of elevational movement in the winter. Winters in the
Western Ghats are the most temperate season, with favorable temperatures across the elevational
gradient and limited precipitation. During this season, birds are likely responding to other
confounding drivers, like food resource phenology, that were not included in this study. We
further tested the drivers of these elevational movements in association with key predictors, as
we discuss in more detail below.

Although summer upslope movements may align with breeding site selection and
monsoon season downslope movements with the LFO hypothesis, the reasons behind some
species moving in unexpected directions remain unclear. For example, the Large Gray Babbler
(*Argya malcolmi*), a widely distributed species across the Indian subcontinent (Collar & Robson,
2021), exhibited downslope movement in summer and upslope movement during the monsoon
season (Figure 2). Another species, the Palani Laughingthrush (*Montecincla fairbanki*), a range-
restricted endemic of the Western Ghats (Collar et al., 2020; Robin et al., 2017), demonstrated a
similar pattern of movement (Figure 2). We found eight other species moved downslope in
summer and three more species moved upslope during the monsoon season. These species
exhibit varied life histories, morphological traits, and range distributions, making it difficult to
identify a common driver for these 'against-the-norm' movements through this analysis. We
suggest a more species-specific examination of their migration patterns, which could improve
understanding of why current hypotheses fail to explain a small subset of species' seasonal
elevational movements.

The next question addressed in this study was which factors determine whether a species
makes seasonal elevational migration, regardless of direction. Our results support both the
"climatic constraint" and LFO hypotheses, depending on the season and variable examined.
Species with broader temperature tolerances were more likely to be migratory across all seasons,

which contrasts with the "climatic constraint hypothesis." Recent studies on avian elevational
migration in the Himalayas and Taiwan suggest that species with narrower thermal regimes or
limited cold tolerance make longer elevational movements (Menon et al., 2023; Tsai et al., 2021),
as these regions experience substantial seasonal temperature shifts across steeper gradients. By
contrast, the Western Ghats experience low temperature seasonality (Figure 1c), which likely
reduces temperature constraints on migratory behavior, allowing species with broader tolerances
to remain within a broad thermal niche year-round. This flexibility may allow them to respond to
other seasonal drivers, including precipitation patterns, by tracking habitats and resources that
fluctuate seasonally. Indeed, we found that precipitation, a climate variable with far greater
seasonality than temperature (Figure 1e), had a stronger influence on the likelihood of being
migratory. Species favoring wetter conditions were more likely to migrate in both summer and
winter, suggesting a tracking of the precipitation dimension of their eco-climatic niche. This
result aligns with the "climatic constraint hypothesis," reflecting the need to track wetter regions
along the elevational gradient during key breeding (summer) and potentially more resource-
limited (winter) seasons. During the monsoon, migrants tended to be species with overall
narrower precipitation tolerances, whether they preferred more wet or dry conditions, further
supporting the "climatic constraint hypothesis." Collectively, these findings suggest that while
temperature constraints are moderate, other seasonal conditions, particularly those varying with
precipitation, play a significant role in structuring migratory strategies in the Western Ghats.

Precipitation is known to influence plant and insect phenology (Chown et al., 2011;
Mendoza et al., 2017; Newell et al., 2023), which in turn affects food availability for higher
trophic levels, providing a potential indirect driver of migratory responses. In the monsoon-
dominant Western Ghats, tracking this dimension of a species' eco-climatic niche is crucial for
meeting seasonal energy requirements. During the drier seasons (summer and winter), species
tend to migrate toward wetter environments where resources may be more abundant. However,
during prolonged heavy rainfall events in the monsoon, resources may be abundant but less
accessible due to foraging limitations. Consequently, species are likely to avoid areas with
excessive precipitation and limited foraging opportunities, favoring lower elevations with more
accessible resources. Furthermore, we observed a dietary guild response among monsoon season
elevational migrants, with frugivores showing lower odds of migration. On average, frugivorous
species in this study were of higher body mass than most other foraging guilds (Figure S1.6),
which may make them better equipped to endure resource limitations during the monsoon season
(Boyle, 2008b).

In sum, we find strong support for the LFO hypothesis, especially in the monsoon season.
Elevational migrants in the Western Ghats appear to track regions of higher precipitation and
associated changes in resource abundance during summer and winter. In contrast, larger-bodied
frugivores are less likely to migrate during the monsoon, while species that do migrate tend to
avoid areas of extreme precipitation where foraging opportunities may be compromised. Our
findings reveal the complexity of elevational migration strategies in the region, showing that

species respond to a diverse set of environmental factors, each shifting across seasons in ways
that are not always synchronized with breeding periods. These multifaceted and complex
relationships may not align with traditional hypotheses, showing how species have adapted to the
hypervariable conditions of the Western Ghats.

Caveats and Conclusions:

This work relied solely on citizen science data collected from the Western Ghats by
thousands of citizen scientists across India. Despite rigorous filtering, inherent biases persisted
(Figure S1.1). We developed an approach to correct these biases by using fitted GAMs to predict
sampling effort and resample citizen science occurrence records based on these predictions. The
strength of this method lies in resampling across the gradient of occurrence records, weighted by
a user-controlled scale of sampling effort. Furthermore, resampling results in a continuous
distribution of occurrence records across the gradient of interest. In our case, we applied this
method across a gradient of elevation; however, it can be used for various ecological and
environmental gradients as long as there is some a priori estimate of sampling effort. We
acknowledge that despite bias correction methods, biases in citizen science data may still arise
due to its opportunistic nature. For this reason, we did not attempt to estimate the elevational
boundaries of a species in this study due to the extremely limited number of high-elevation
occurrence records. Instead, we estimated median elevations of a species, which are relatively
less prone to error. Further empirical testing of the method is required to ascertain its efficiency
in predicting robust estimates of boundary quantiles. By using measures of central tendency, we
believe our estimates of species' seasonal median elevation were not influenced by latent
sampling biases and were reliable enough for drawing inferences. We also acknowledge that the
variables used in our analyses were proxies for traits hypothesized to impact elevational
migration, and thus we do not fully or directly test the previously outlined hypotheses. While we
provide a reliable foundation for ecological inferences on the drivers of elevational migration, we
encourage further research that directly tests these hypotheses, such as quantifying foraging
opportunities throughout the year and their impact on elevational migrants or obtaining more
direct measures of the physiological tolerances of elevational migrants and their correlation with
migratory behavior.

In conclusion, our study highlights the complex drivers behind seasonal elevational
migration in the Western Ghats. We found that broader temperature tolerances, rather than
narrower ones, were associated with migratory behavior, suggesting that temperature constraints
are less impactful in this region. Precipitation, with its greater seasonality and intensity
especially during the monsoon season, played a stronger role in shaping migration strategies.
Species tended to track wetter regions or avoid extreme rainfall, indicating that resource needs,
influenced indirectly by precipitation patterns, drive elevational migration. We also highlight
how the monsoon, as a single challenging season, uniquely influences elevational migration
strategies throughout the year, driven by factors distinct from those shaping migrations in other

seasons. This emphasizes the importance of including these transition seasons, between the breeding and non-breeding seasons, in future avian elevational migration studies. Summing across seasons, our findings align with both the "climatic constraint" and Limited Foraging Opportunities hypotheses and provide a foundation for further empirical testing of these ideas. The consistent movement of species along the elevational gradient across all three seasons underscores the necessity of preserving these key breeding habitats and elevational refuges to ensure species survival in the region. Given the accelerating land-use changes and biological invasions affecting the Western Ghats (Arasumani et al., 2019; Lele et al., 2020), conserving these vital habitats is essential for maintaining the resilience of bird populations and ensuring the persistence of biodiversity in this ecologically rich area.

References:

Abrams, M., Crippen, R., & Fujisada, H. (2020). ASTER Global Digital Elevation Model (GDEM) and ASTER Global Water Body Dataset (ASTWBD). *Remote Sensing*, *12*(7), 1156. https://doi.org/10.3390/rs12071156

Alves, J. A., Gunnarsson, T. G., Hayhow, D. B., Appleton, G. F., Potts, P. M., Sutherland, W. J., & Gill, J. A. (2013). Costs, benefits, and fitness consequences of different migratory strategies. *Ecology*, *94*(1), 11–17. https://doi.org/10.1890/12-0737.1

Arasumani, M., Khan, D., Vishnudas, C. K., Muthukumar, M., Bunyan, M., & Robin, V. V. (2019). Invasion compounds an ecosystem-wide loss to afforestation in the tropical grasslands of the Shola Sky Islands. *Biological Conservation*, *230*, 141–150. https://doi.org/10.1016/j.biocon.2018.12.019

Barçante, L., M. Vale, M., & S. Alves, M. A. (2017). Altitudinal migration by birds: A review of the literature and a comprehensive list of species. *Journal of Field Ornithology*, *88*(4), 321–335. https://doi.org/10.1111/jofo.12234

Bowlin, M. S., Bisson, I.-A., Shamoun-Baranes, J., Reichard, J. D., Sapir, N., Marra, P. P., Kunz, T. H., Wilcove, D. S., Hedenstrom, A., Guglielmo, C. G., Akesson, S., Ramenofsky, M., & Wikelski, M. (2010). Grand Challenges in Migration Biology. *Integrative and Comparative Biology*, *50*(3), 261–279. https://doi.org/10.1093/icb/icq013

Boyle, W. A. (2008a). Can variation in risk of nest predation explain altitudinal migration in tropical birds? *Oecologia*, *155*(2), 397–403. https://doi.org/10.1007/s00442-007-0897-6

Boyle, W. A. (2008b). Partial migration in birds: Tests of three hypotheses in a tropical lekking frugivore. *Journal of Animal Ecology*, *77*(6), 1122–1128. https://doi.org/10.1111/j.1365-2656.2008.01451.x

Boyle, W. A. (2010). Does food abundance explain altitudinal migration in a tropical frugivorous bird? *Canadian Journal of Zoology*, *88*(2), 204–213. https://doi.org/10.1139/Z09-133

Boyle, W. A. (2011). Short-distance partial migration of Neotropical birds: A community-level test of the foraging limitation hypothesis. *Oikos*, *120*(12), 1803–1816. https://doi.org/10.1111/j.1600-0706.2011.19432.x

Boyle, W. A. (2017). Altitudinal bird migration in North America. *The Auk*, *134*(2), 443–465. https://doi.org/10.1642/AUK-16-228.1

Boyle, W. A., Norris, D. R., & Guglielmo, C. G. (2010). Storms drive altitudinal migration in a tropical bird. *Proceedings of the Royal Society B: Biological Sciences*, *277*(1693), 2511–2519. https://doi.org/10.1098/rspb.2010.0344

Boyle, W. A., Shogren, E. H., & Brawn, J. D. (2020). Hygric Niches for Tropical Endotherms. *Trends in Ecology & Evolution*, *35*(10), 938–952. https://doi.org/10.1016/j.tree.2020.06.011

Chown, S. L., Sørensen, J. G., & Terblanche, J. S. (2011). Water loss in insects: An environmental change perspective. *Journal of Insect Physiology*, *57*(8), 1070–1084. https://doi.org/10.1016/j.jinsphys.2011.05.004

Claramunt, S., Derryberry, E. P., Remsen, J. V., & Brumfield, R. T. (2012). High dispersal ability inhibits speciation in a continental radiation of passerine birds. *Proceedings of the Royal Society B: Biological Sciences*, *279*(1733), 1567–1574. https://doi.org/10.1098/rspb.2011.1922

Collar, N., & Robson, C. (2021). Large Gray Babbler (*Argya malcolmi*). In S. M. Billerman, B. K. Keeney, P. G. Rodewald, & T. S. Schulenberg (Eds.), *Birds of the World*. Cornell Lab of Ornithology. https://doi.org/10.2173/bow.lagbab2.01.1

Collar, N., Robson, C., & Christie, D. (2020). Palani Laughingthrush (*Montecincla fairbanki*). In S. M. Billerman, B. K. Keeney, P. G. Rodewald, & T. S. Schulenberg (Eds.), *Birds of the World*. Cornell Lab of Ornithology. https://doi.org/10.2173/bow.kerlau2.01

Das, A., Krishnaswamy, J., Bawa, K. S., Kiran, M. C., Srinivas, V., Kumar, N. S., & Karanth, K. U. (2006). Prioritisation of conservation areas in the Western Ghats, India. *Biological Conservation*, *133*(1), 16–31. https://doi.org/10.1016/j.biocon.2006.05.023

Dingle, H., & Drake, V. A. (2007). What Is Migration? *BioScience*, *57*(2), 113–121. https://doi.org/10.1641/B570206

Farr, T. G., Rosen, P. A., Caro, E., Crippen, R., Duren, R., Hensley, S., Kobrick, M., Paller, M., Rodriguez, E., Roth, L., Seal, D., Shaffer, S., Shimada, J., Umland, J., Werner, M., Oskin, M.,

Burbank, D., & Alsdorf, D. (2007). The Shuttle Radar Topography Mission. *Reviews of Geophysics*, *45*(2). https://doi.org/10.1029/2005RG000183

Foster, M. S. (1974). Rain, Feeding Behavior, and Clutch Size in Tropical Birds. *The Auk*, *91*(4), 722–726. https://doi.org/10.2307/4084724

Freeman, B. G., Montgomery, G. A., Heavyside, J., Moncrieff, A. E., Johnson, O., & Winger, B. M. (2023). On the predictability of phenotypic divergence in geographic isolation. *Evolution*, *77*(1), 26–35. https://doi.org/10.1093/evolut/qpac040

Gadgil, M., & Meher-Homji, V. (1986). Localities of great significance to conservation of India's biological diversity. *Proceedings of the Indian Academy of Science*: 165–180.

Hastie, T., & Tibshirani, R. (1986). Generalized Additive Models. *Statistical Science*, *1*(3). https://doi.org/10.1214/ss/1177013604

Ho, L. S. T. and Ane, C. (2014) A linear-time algorithm for Gaussian and non-Gaussian trait evolution models. *Systematic Biology*, 63(3):397-408.

Hsiung, A. C., Boyle, W. A., Cooper, R. J., & Chandler, R. B. (2018). Altitudinal migration: Ecological drivers, knowledge gaps, and conservation implications: Animal altitudinal migration review. *Biological Reviews*, *93*(4), 2049–2070. https://doi.org/10.1111/brv.12435

Ives, A. R., & Garland, T. (2010). Phylogenetic Logistic Regression for Binary Dependent Variables. *Systematic Biology*, *59*(1), 9–26. https://doi.org/10.1093/sysbio/syp074

Jetz, W., Thomas, G. H., Joy, J. B., Hartmann, K., & Mooers, A. O. (2012). The global diversity of birds in space and time. *Nature*, *491*(7424), 444–448. https://doi.org/10.1038/nature11631

Karger, D. N., Conrad, O., Böhner, J., Kawohl, T., Kreft, H., Soria-Auza, R. W., Zimmermann, N. E., Linder, H. P., & Kessler, M. (2017). Climatologies at high resolution for the earth's land surface areas. *Scientific Data*, *4*(1), 170122. https://doi.org/10.1038/sdata.2017.122

Kelling, S., Johnston, A., Bonn, A., Fink, D., Ruiz-Gutierrez, V., Bonney, R., Fernandez, M., Hochachka, W. M., Julliard, R., Kraemer, R., & Guralnick, R. (2019). Using Semistructured Surveys to Improve Citizen Science Data for Monitoring Biodiversity. *BioScience*, *69*(3), 170–179. https://doi.org/10.1093/biosci/biz010

Ketterson, E. D., & Nolan, V. (1976). Geographic Variation and Its Climatic Correlates in the Sex Ratio of Eastern-Wintering Dark-Eyed Juncos (*Junco Hyemalis Hyemalis*). *Ecology*, *57*(4), 679–693. https://doi.org/10.2307/1936182

Lele, A., Arasumani, M., Vishnudas, C. K., Joshi, V., Jathanna, D., & Robin, V. V. (2020).
Elevation and landscape change drive the distribution of a montane, endemic grassland bird.
Ecology and Evolution, *10*(14), 7755–7767. https://doi.org/10.1002/ece3.6500

Levey, D. J. (1988). Spatial and Temporal Variation in Costa Rican Fruit and Fruit-Eating Bird
Abundance. *Ecological Monographs*, *58*(4), 251–269. https://doi.org/10.2307/1942539

Loiselle, B. A., & Blake, J. G. (1991). Temporal Variation in Birds and Fruits Along an
Elevational Gradient in Costa Rica. *Ecology*, *72*(1), 180–193. https://doi.org/10.2307/1938913

Mendoza, I., Peres, C. A., & Morellato, L. P. C. (2017). Continental-scale patterns and climatic
drivers of fruiting phenology: A quantitative Neotropical review. *Global and Planetary Change*,
148, 227–241. https://doi.org/10.1016/j.gloplacha.2016.12.001

Menon, T., Ramesh, V., & Barve, S. (2023). Himalayan birds that show the greatest elevational
shifts remain within the narrowest thermal regimes. *Global Ecology and Biogeography*, *32*(12),
2111–2121. https://doi.org/10.1111/geb.13761

Milner-Gulland, E.J., John M. Fryxell, and Anthony R.E. Sinclair (eds), *Animal Migration: A
Synthesis* (Oxford, 2011; online edn, Oxford Academic, 17 Dec.
2013), https://doi.org/10.1093/acprof:oso/9780199568994.001.0001

Myers, N., Mittermeier, R. A., Mittermeier, C. G., Da Fonseca, G. A. B., & Kent, J. (2000).
Biodiversity hotspots for conservation priorities. *Nature*, *403*(6772), 853–858.
https://doi.org/10.1038/35002501

Newell, F. L., Ausprey, I. J., & Robinson, S. K. (2023). Wet and dry extremes reduce arthropod
biomass independently of leaf phenology in the wet tropics. *Global Change Biology*, *29*(2), 308–
323. https://doi.org/10.1111/gcb.16379

Pascal, J. P. (1988). Wet evergreen forests of the Western Ghats of India: ecology, structure,
floristic composition and succession. *Institut français de Pondichéry.*

Pebesma, E., 2018. Simple Features for R: Standardized Support for Spatial Vector Data. The R
Journal 10 (1), 439-446, https://doi.org/10.32614/RJ-2018-009

Ramesh, V., Gupte, P. R., Tingley, M. W., Robin, V. V., & DeFries, R. (2022). Using citizen
science to parse climatic and land cover influences on bird occupancy in a tropical biodiversity
hotspot. *Ecography*, *2022*(9). https://doi.org/10.1111/ecog.06075

Rappole, J. (2013). *The Avian Migrant: The Biology of Bird Migration*. New York Chichester,
West Sussex: Columbia University Press. https://doi.org/10.7312/rapp14678

R Core Team (2024). *R: A Language and Environment for Statistical Computing*. R Foundation
for Statistical Computing, Vienna, Austria. https://www.R-project.org/.

Revell, L. J. (2010). Phylogenetic signal and linear regression on species data: Phylogenetic
regression. *Methods in Ecology and Evolution*, *1*(4), 319–329. https://doi.org/10.1111/j.2041-
210X.2010.00044.x

Robin, V. V., Vishnudas, C. K., Gupta, P., Rheindt, F. E., Hooper, D. M., Ramakrishnan, U., &
Reddy, S. (2017). Two new genera of songbirds represent endemic radiations from the Shola Sky
Islands of the Western Ghats, India. *BMC Evolutionary Biology*, *17*(1), 31.
https://doi.org/10.1186/s12862-017-0882-6

Rueda-Uribe, C., Herrera-Alsina, L., Lancaster, L. T., Capellini, I., Layton, K. K. S., & Travis, J.
M. J. (2024). Citizen science data reveal altitudinal movement and seasonal ecosystem use by
hummingbirds in the Andes Mountains. *Ecography*, *2024*(3), e06735.
https://doi.org/10.1111/ecog.06735

Shaw, A. K. (2016). Drivers of animal migration and implications in changing environments.
Evolutionary Ecology, *30*(6), 991–1007. https://doi.org/10.1007/s10682-016-9860-5

Sheard, C., Neate-Clegg, M. H. C., Alioravainen, N., Jones, S. E. I., Vincent, C., MacGregor, H.
E. A., Bregman, T. P., Claramunt, S., & Tobias, J. A. (2020). Ecological drivers of global
gradients in avian dispersal inferred from wing morphology. *Nature Communications*, *11*(1),
2463. https://doi.org/10.1038/s41467-020-16313-6

Skutch, A. F. (1985). Clutch Size, Nesting Success, and Predation on Nests of Neotropical Birds,
Reviewed. *Ornithological Monographs*, *36*, 575–594. https://doi.org/10.2307/40168306

Stiles, F. G. (1988). Altitudinal movements of birds on the Caribbean slope of Costa Rica:
implications for conservation. *Tropical rainforests: diversity and conservation*, *12*, 243.

Strimas-Mackey, M., Hochachka, W. M., Ruiz-Gutierrez, V., Robinson, O. J., Miller, E. T., Auer,
T., Kelling, S., Fink, D., & Johnston, A. (2020). *Best Practices for Using eBird Data v1.0*
(Version v1.0). Zenodo. https://doi.org/10.5281/ZENODO.3620739

Strimas-Mackey, M., Miller, E., & Hochachka, W. (2023). auk: eBird Data Extraction and
Processing with AWK. R package version 0.7.0. https://cornelllabofornithology.github.io/auk/.

Sullivan, B. L., Aycrigg, J. L., Barry, J. H., Bonney, R. E., Bruns, N., Cooper, C. B., Damoulas,
T., Dhondt, A. A., Dietterich, T., Farnsworth, A., Fink, D., Fitzpatrick, J. W., Fredericks, T.,
Gerbracht, J., Gomes, C., Hochachka, W. M., Iliff, M. J., Lagoze, C., La Sorte, F. A., … Kelling,
S. (2014). The eBird enterprise: An integrated approach to development and application of

citizen science. *Biological Conservation*, *169*, 31–40.
https://doi.org/10.1016/j.biocon.2013.11.003

Sullivan, B. L., Wood, C. L., Iliff, M. J., Bonney, R. E., Fink, D., & Kelling, S. (2009). eBird: A
citizen-based bird observation network in the biological sciences. *Biological Conservation*,
142(10), 2282–2292. https://doi.org/10.1016/j.biocon.2009.05.006

Tobias, J. A., Sheard, C., Pigot, A. L., Devenish, A. J. M., Yang, J., Sayol, F., Neate-Clegg, M. H.
C., Alioravainen, N., Weeks, T. L., Barber, R. A., Walkden, P. A., MacGregor, H. E. A., Jones, S.
E. I., Vincent, C., Phillips, A. G., Marples, N. M., Montaño-Centellas, F. A., Leandro-Silva, V.,
Claramunt, S., … Schleuning, M. (2022). AVONET: Morphological, ecological and geographical
data for all birds. *Ecology Letters*, *25*(3), 581–597. https://doi.org/10.1111/ele.13898

Tsai, P., Ko, C., Chia, S. Y., Lu, Y., & Tuanmu, M. (2021). New insights into the patterns and
drivers of avian altitudinal migration from a growing crowdsourcing data source. *Ecography*,
44(1), 75–86. https://doi.org/10.1111/ecog.05196

Voigt, C. C., Schneeberger, K., Voigt-Heucke, S. L., & Lewanzik, D. (2011). Rain increases the
energy cost of bat flight. *Biology Letters*, *7*(5), 793–795. https://doi.org/10.1098/rsbl.2011.0313

Williamson, J. L., & Witt, C. C. (2021). Elevational niche-shift migration: Why the degree of
elevational change matters for the ecology, evolution, and physiology of migratory birds.
Ornithology, *138*(2). https://doi.org/10.1093/ornithology/ukaa087

Wood, S.N. (2017) Generalized Additive Models: An Introduction with R (2nd edition).
Chapman and Hall/CRC.

Zurell, D., Gallien, L., Graham, C. H., & Zimmermann, N. E. (2018). Do long-distance
migratory birds track their niche through seasons? *Journal of Biogeography*, *45*(7), 1459–1468.
https://doi.org/10.1111/jbi.13351

674   **Tables**

675   Table 1: Chi-square test results examining the intra-seasonal variation in elevational migration
676   direction of resident birds of the Western Ghats. Seasonal percentages of species demonstrating
677   downslope (Downslope %) or upslope (Upslope %) movement are reported, along with chi-
678   square statistics (χ^2) and P values for each seasonal test.

Season	Downslope %	Upslope %	χ^2	P value
Summer	18	83	24.018	<0.001
Monsoon	87	13	21.546	<0.001
Winter	53	47	0.105	0.746

Figures

Figure 1: The Western Ghats of India (boundary shown as black line) are topographically and
climatically unique. (a) Latitudinal extent and elevational range of the Western Ghats. (b) Mean
annual temperature in the region ranges from 12°C at higher elevations to 25°C in the plains. (c)
Temperature seasonality is lower along the length of these hills and increases moving eastward
across the Indian subcontinent. (d) The Western Ghats creates a strong rain shadow effect,
leading to a longitudinal precipitation gradient. (e) The topography of the Western Ghats, results
in a decrease in precipitation seasonality moving from north to south. Elevation is built from 30
seconds resolution SRTM data (Farr et al., 2007) and climatic variables are visualized from
CHELSA data at 1 km resolution (Karger et al., 2017).

Figure 2: Patterns of elevational migration in this study for species that exhibit significant
elevational movement between the seasons of interest. In the summer species moved
predominantly upslope followed by downslope movement in the monsoon. The winter saw an
almost even distribution of upslope and downslope elevational movements. The distance species
move in meters (m) is represented as the length of the arrow.

Figure 3: Parameter estimates for our seasonal phylogenetic logistic regression models
determining the effect of our candidate predictors on elevational migration status (migrant vs
non-migrant). Dots represent model estimates with their 95% confidence intervals as error bars.
Asterisk denotes significance ($p < 0.05$). For diet categories, the reference level was set as
species with an omnivorous diet.

718