Headwaters to Valley: Water quality in rivers transitioning from forest to agricultural

bottomland

Doug Graber Neufeld, Isaac Alderfer, Zachary Bauman, Micah Buckwalter

Department of Biology, Eastern Mennonite University, Harrisonburg, Virginia, United States of America

Abstract

Many waterways flow out of forestlands, which tend to maintain higher water quality, into agricultural lands,

which tend to degrade water quality. The roles of these two types of land cover in impacting key water quality

parameters (phosphorus as orthophosphate, nitrogen as nitrate, total suspended sediment (TSS), and indicator

bacteria) were investigated for the watershed of the North and South Fork of the Shenandoah River, Virginia. This

area has a particularly sharp boundary between heavily forested and heavily agricultural regions. Two datasets

were analyzed: 1) a large number of datapoints spanning a 20-year range in the Water Quality Portal (WQP)

database, and 2) transects along three representative rivers systems over the span of a 4-year period. Overall

nutrient levels were significantly better in forested regions than agricultural regions; while TSS and indicator

bacteria showed some trends, but less consistently, of higher water quality in forested regions. All parameters

had significant variation in both forested and agricultural areas. Transects showed that waterways flowing out of

forestland maintained higher water quality for approximately the first 8 km in agricultural land. Both transect and

WQP data indicated higher water quality when the percent of forested land cover in a drainage basin was about

70-80%. Thus, forestland does mitigate the impacts of agriculture on water quality to some degree, but this effect

rapidly diminishes as forest cover of the watershed lessens. Furthermore, forests themselves have degraded

water quality at certain times and places; for instance, nutrients level were in the medium to high stress level for

aquatic life in approximately 15% of samples. This study illustrates general trends of land cover effects on water

quality, while also highlighting both site-specific variability, and the dynamics of water quality as water flows out

of forested areas into agricultural areas.

Introduction

Aquatic ecosystems from headwaters to larger waterways downstream have been a major focus of efforts at

maintaining and improving environmental quality, due to their combined importance in providing ecosystem

services, and their integral role in the lives of people [1]. Indeed, maintenance of water quality is directly related

to important planetary boundaries such as the geochemical cycling of nitrogen and phosphorus [2], which have

clearly exceeded safe operating zones. Numerous studies have explored key water quality parameters elucidating

both patterns that are generalizable across waterways, and patterns that are specific to local spatial and temporal

conditions.

Land cover, and especially continued pressure from human activities that alter land cover, are fundamental in

driving water quality of watersheds. Most notable is how natural vegetation such as forests can improve water

quality, while agricultural activities (both pastureland and cropland) can degrade water quality. Protection of

forested lands is seen as crucial to maintaining clean water [3,4], notably illustrated by a recent study in the

southeast United States found that forests had the highest water quality of all landscape types [5]. In parallel to

land cover effects, headwater regions (which often are more forested) have unique characteristics compared to

downstream waters [6-8]. There are clearly multiple factors beyond just land cover type that control water

quality [5], and more information is needed on how water quality changes as waterways traverse different

landscapes. In particular, it is not clear whether there is a sharp transition in water quality as it traverses forested

and agricultural land, and how the impacts of the two landscapes combine to influence downstream water

quality.

Watersheds draining into the Chesapeake Bay are the focus of a long running effort to understand how stressors

such as nutrients and sediments contribute to estuarine health, and how these inputs can be reduced [9-11].

Efforts in the Chesapeake watershed have been mixed, illustrating both the ongoing challenge of reducing

agricultural inputs, and the need for more information on what exactly drives nutrient trends [9,11,12]. The

Shenandoah River watershed of Virginia and West Virginia is a major tributary system of the Chesapeake Bay, and

stands out as having both some of the region's largest protected forestlands, and also the highest density of

livestock and poultry operations in Virginia, along with significant cropland. Furthermore, there is a generally

distinct boundary between the relatively flat valley bottomland which is dominated by agriculture and

predominantly of limestone origin, compared to the surrounding mountains which are heavily forested, federally

protected as national forest and national park land, and predominantly of sandstone origin. This watershed

therefore represents a unique opportunity to study a sharp transition of waterways between heavy forestland of

headwaters, and intensive agriculture downstream.

The present study combines data from a large public database (Water Quality Portal; WQP) with targeted

sampling on representative river systems to compare water quality in adjacent clear heavily forested and

intensively agricultural lands, and to characterize the pattern of change as rivers flow between these two types of

land cover. We focus especially on nutrients (nitrate and orthophosphate), sediments, and bacteria as parameters

determining both local aquatic health, and downstream impacts [13,14].

Methods

Ethics Statement

Site access outside of national forest was either at publicly accessible right-of-way along state or county highways

(usually bridges), or by landowner permission. Within the national forest, permission for site access for the

purpose of water sampling was granted by the US Forest Service.

North and South Fork Shenandoah River Watershed

Water quality data for the combined watersheds of the North and South Forks of the Shenandoah River

watershed were accessed from the Water Quality Portal (WQP), an open access compilation of water quality

results from multiple government, academic and nonprofit organizations. Since land downstream of the

confluence of the North and South Forks (north of Front Royal, Virginia) does not have protected National Forest

or National Park land (as dominates forestland upstream from this point), we did not analyze the watershed of the

Shenandoah River downstream of the confluence. Samples from within the delineated watershed were selected

for the 20 year period from Jan 1, 2004 to Dec 31, 2023. The number of sample values varied with parameter and

ranged between 9,742 and 29,897. Since most sites had multiple measurements of a parameter for that period,

we used the mean parameter value for each site. Outliers were identified and removed for samples having Cook's

distance of 4/n (0.005 for this dataset), and $p<0.05$. The number of outliers varied with parameter, ranging from

1.4% to 3.4% of samples, and the resultant number of sites for each parameter ranged from 639 to 738.

Spatial analysis was performed using ArcGIS Pro 3.3 and QGIS 3.38.0. Since in other aquatic systems, water quality

is shown to be driven by overall land use across the entire drainage basin for a collection site [15], we delineated

watersheds from USGS 1/3 arc-second Digital Elevation Maps (3DEP product in The National Map), and land

use/land cover (LULC) from the USGS National Land Cover Database (2021 data). Geology was mapped as

Generalized Lithography in the State Geologic Map Compilation (SGMC) Geodatabase of the Conterminous United

States (ver. 1.1, August 2017). To calculate a general forest boundary, we calculated the average percent forest

within a 200m radius of each pixel (pixel resolution 25m) of the LULC raster using the focal statistics tool of ArcGIS

Pro. Pixels with at least 75% forest in this radius were counted as forest. This produced a smoothed general forest

boundary that depicts the visual forest transition as seen on satellite images. River kilometers were then

calculated upstream and downstream from the point at which the river crosses this boundary.

River Transect Sites

Three river systems were chosen for water sampling as transects – the Dry River, Briery Branch, and the North

River. All three rivers originate in the mountains on the western edge of the Shenandoah watershed, converge as

the lower North River, and then flow together into the South Fork of the Shenandoah River which joins the

Potomac River before entering the Chesapeake Bay. The headwaters are at approximately 3000 feet, and flow

through the oak-hickory forest of George Washington National Forest before abruptly entering the Shenandoah

valley agricultural region.

Water sample sites were selected at approximately 5km intervals, although specific sites were dependent on

availability of river access. In addition, river sections in the middle interior of the valley dried completely during

periods of low rainfall, especially in midsummer. In other cases of high rainfall or winter, sites in the national

forest were inaccessible for sampling. The final number of samples per site therefore varied. Samples were

collected between March 2020 and July 2024 and were taken at all times of the year. As a general indication of

river level in the region, flow discharge was recorded from a USGS station (#01622000) near Burketown on the

North River, and varied from 114 to 1430 ft^3 per second, representing a range of discharge from 18[th] to 97[th]

percentiles for the period samples were taken. Samples thus represent a wide range of seasons and river flow

conditions.

At each site, parameters were either measured onsite (temperature and conductivity), or water samples were

collected in dedicated water sample bottles (nitrate, orthophosphate, bacteria, sediments). All measurements or

bottle samples were collected in the main column below the water surface, upstream from any disturbances

(wading or previous sampling). Field blanks were collected at one site per transect for nitrate, orthophosphate,

and bacteria. A field spike bottle was also collected at one site per transect for nitrate and orthophosphate. Field

duplicates were collected for all nitrate, orthophosphate and bacteria samples, and for one sediment sample per

transect. All samples were kept on ice (for transport) or at 4^{o}C storage until analysis.

River Transect Parameters

Water samples were analyzed for sediment, nitrate, orthophosphate, temperature, conductivity, and general and

fecal coliforms. A Quality Assurance Project Plan, previously approved for earlier NFWF and Virginia DEQ grant

sampling, guided analysis, and was based on USGS, Virginia DEQ, or EPA approved procedures (e.g. U.S. EPA 2007.

SW-846).

Sediments were measured gravimetrically as per EPA method 160.2. In brief, 1 L samples were filtered through

pre-rinsed, dried and weighted 47 mm diameter GF/F filters (0.7 μM pore-size), and then re-dried and re-weighed.

Conductivity was measured using a Tracer Pocketester (LaMotte) calibrated to a 84 μS/cm standard. Probe

calibration was verified after each transect to ensure calibration did not drift during the measurement period.

Temperature was measured using a Vernier temperature probe, or the internal thermometer on the conductivity

probe. General and fecal coliforms were measured using Coliform Easygel (Micrology, Inc; Goshen, Indiana, USA).

1 ml samples were plated on agar petri dishes, incubated at 37^{o}C until colony color development (generally 36-

48h), and then colony identity determined by color.

Nitrogen and phosphorus were assessed as nitrate and orthophosphate, respectively. While other forms would be

present, these are the most common and relevant forms in these waters. For instance, nitrate is the majority of

total nitrogen in the region [14]. Likewise, the study of Farthing et al [16] on the effect of forestland on

agriculturally impacted rivers found most nitrogen as nitrate, and most phosphorus as phosphate. Both nitrate

and orthophosphate were measured with a colorimetric assay (Hach; for NitraVer5 and PhosVer3, respectively)

according to manufacturer instructions. Duplicate sample bottles for each site were measured and the mean used

as the final value. Absorbance values at appropriate wavelengths were compared to a calibration series, with

blank and standard tested every 10th sample.

Results

Water Quality Patterns in the North and South Forks of Shenandoah River Watershed

Water quality data from the WQP for the entire watershed of the North and South Forks of the Shenandoah River

were compared with documented land cover. Parameters of concern generally correlated with the degree of

forest cover in the watersheds represented by the sample sites. Nitrate, orthophosphate, and TSS all had higher

concentrations outside of areas that were primarily forested (Figs 1a,b,c). This pattern was less obvious for *E.coli*,

with many sites in forestland showing higher indicator bacteria concentrations (Fig 1d). Conductivity was higher in

the valley (Fig 1e), due to the underlying karst geology of the valley, whereas the mountain areas are mostly

silicoclastic (or basaltic, in a few regions of the blue ridge) (Fig 2). The forest boundary corresponded closely to

the geological boundary, most likely because forest was historically cleared in the more agriculturally productive

limestone/dolostone areas. (Fig 2). Because conductivity seemed so clearly correlated with the dramatic

difference in geology that likely overwhelmed any anthropogenic increases in conductivity, we did not focus on

conductivity in the analyses. The percent forestland in the drainage basin represented by sample sites reflects

both site location (forested vs nonforested areas), and the area upstream draining into the point (Fig 1f).

A. Nitrate

Nitrate (ppm)
- 3.40 - 5.00
- 2.29 - 3.39
- 1.54 - 2.28
- 1.02 - 1.53
- 0.67 - 1.01
- 0.43 - 0.66
- 0.26 - 0.42
- 0.15 - 0.25
- 0.07 - 0.14
- 0.01 - 0.06
- Forested areas

0 5 10   20 Miles

B. Orthophosphate

Phosphate (ppm)
- 0.220 - 0.466
- 0.107 - 0.219
- 0.055 - 0.106
- 0.032 - 0.054
- 0.021 - 0.031
- 0.016 - 0.020
- 0.013 - 0.015
- 0.012 - 0.012
- 0.010 - 0.011
- 0.004 - 0.009
- Forest area

0 5 10   20 Miles

C. Total Suspended Solids

Nitrate (ppm)
TSS
- 18.53 - 32.70
- 11.20 - 18.52
- 7.41 - 11.19
- 5.45 - 7.40
- 4.44 - 5.44
- 3.91 - 4.43
- 3.64 - 3.90
- 3.12 - 3.63
- 2.10 - 3.11
- 0.13 - 2.09
- ShenRivers_Clip
- StudyAreaShenValley
- Forested areas

0 5 10   20 Miles

D. *E. coli*

E. coli (cfu/100ml)
Ecoli
- 34545 - 80000
- 14912 - 34544
- 6431 - 14911
- 2768 - 6430
- 1186 - 2767
- 503 - 1185
- 207 - 502
- 80 - 206
- 25 - 79
- 0 - 24
- Forested areas

0 5 10   20 Miles

Fig. 1. Water quality parameters in watershed North and South Forks of Shenandoah River of Virginia. Data from Water

Quality Portal for 2004-2023.

Fig. 2. Underlying lithology of the Shenandoah River watershed compared with areas with substantial forestland (areas with

75% forest cover within 200m radius; see methods).

We quantified how water quality correlates with forest cover first by comparing each water quality parameter to

the percent forest cover of the calculated watershed draining into the sample sites. Linear regression of site

watershed forest cover compared with parameters of concern indicated a significant relationship for nitrate and

TSS, but not for orthophosphate or *E. coli* (Table 1). Given that parameters may relate nonlinearly to the

proportion of forest cover, we further analyzed the relationship by grouping watersheds into 5 bins representing

20% increments of forest cover. Statistical analysis (ANOVA, with Tukey's HSD post-hoc) indicated significant

differences with forest cover for all parameters (Fig. 3). For nitrate and orthophosphate, nutrient levels were

significantly lower in the highest category of forest cover (80-100%), whereas the other categories of forest cover

(60-80%, and lower) had higher and similar levels. Nitrate and orthophosphate data therefore suggested that the

greatest decrease in water quality occurred when rivers moved from heavily forested land to moderate or low

forest cover. This pattern was different for TSS, which showed a more gradual increase in concentration with

decreased forest cover, and for *E. coli*, which showed a less consistent pattern with forest cover.

Table 1

Parameter	Relationship of parameter (y) to forest cover (x)	r^2	significance (p value)	sample size (N)
Nitrate	y=1.26x + 1.80	0.17	**p<0.001**	759
Orthophosphate	y=-0.036 + 0.064	0.03	p=0.373	708
TSS	y=-7.87 + 11.94	0.13	**p<0.001**	674
E. coli	y=-0.005 + 2.077	<0.01	p>0.500	648

Fig. 3. Water quality in forest percent categories. Significant differences between categories indicated by letters on error bars

(upper case p<0.05; lower case 0.05<p<0.1).

Cluster and outlier analysis (Anselin Local Moran's I) confirmed that clusters of low nitrate, orthophosphate and

TSS tended to be located in heavily forested land, but in some cases were just outside the forestland, or along

major waterways that would take significant volumes of water into the valley (Fig. 4; light blue points). Clusters of

high values were located in the valley, usually at some distance from the forested areas (Fig. 4; light red points).

For all parameters, there were outliers (high values in low cluster areas, dark red points; or low values in high

cluster areas, dark blue points), pointing to significant localized variation.

Low clusters for *E. coli* tended to be on the edge of forested land, either from waterways flowing out of the forest,

or main the branches of the rivers that flow along the edge of forest land in the northern section of the

watershed. *E. coli* in forest land was almost all classified as not clustered and not outliers – in other words, an

even distribution of moderate values.

Fig. 4. Cluster and outlier analysis of water quality parameters for North and South Fork Shenandoah River, Virginia. "High-

High" and "Low-Low" are points in high and low cluster areas, respectively. "High-Low" and "Low-High" are high values in low

cluster areas, and low values in high cluster areas, respectively.

Overall, these results suggested that any effects of forestland on water quality extend into nonforested land, and

that there is significant localized variation in all parameters.

Water Quality in Transects of Three Rivers Flowing from Forest to Agricultural Bottomland

We measured water quality along transects for three specific rivers – the Dry River, Briery Branch, and North River

– located on the western side of the Shenandoah watershed. All three rivers flow from significant distances

through national forest land, and then enter areas primarily used for agriculture (as both pasture and cropland),

allowing us to explore more how water quality changes with the transition from forest to agriculture (Fig. 5).

There was an increase in orthophosphate, nitrate, and general and fecal coliforms (Figs. 6a,b,d,e) at 8-10 km

downstream of the forest boundary. This pattern was less obvious for TSS (Fig. 6c), which varied more at sample

sites, and had some higher concentrations in the forestland. When we examined temperature differentials (how

much temperature differed from the average temperature across the transect for that sampling run), we found a

different pattern – temperature appeared to more gradually increase rather showing the abrupt shift at 8 river

kms (see supplemental data). There was significant variation for all variables, indicating (as did the previous

analysis of sites across the Shenandoah watershed; Figs 1,4) that there are significant localized effects that impact

water at any particular point and time.

Conductivity also showed a clear change after rivers moved from forest to agricultural lands (Fig. 6f), with the

increase happening about 8-10 river kilometers downstream from the forest edge. Since the boundary between

silicoclastic and karst geology corresponded closer to the forest edge, it appeared that enough water was flowing

out of forestland that ions from the karst geology were not impacting water quality until the 8-10 km mark. This

increase corresponds to when sample site watersheds began to represent a more significant area of agricultural

land (although the total area in watersheds was still mostly forestland) (Fig. 6f).

Fig. 5. River transect sites relative to land cover designations.

A. Nitrate

B. Orthophosphate

C. Total Suspended Solids

D. Fecal coliform

E. General coliforms

Fig. 6. Water quality parameters along river transects, from forest headwaters to agricultural bottomland. Blue, orange and

green points represent parameter mean ± standard deviation values for the North River, Dry River, and Briary Branch,

respectively. Points are Open points represent percent forest fraction for each corresponding parameter point.

Since there was an abrupt shift in some parameters at the 8-10 km mark, we pooled samples from the transects

into three categories: forestland, edge of valley (up to 8 river kms), and interior of the valley (beyond 8 river kms).

Given the known impacts of flow rate on water quality parameters, we furthered binned samples into three flow

rate categories (low, medium and high). Two-way ANOVA did not show any effects of flow rate ($p>0.05$ for all

parameters). We therefore dropped the flow rate bins, and used one-way ANOVA with post-hoc to investigate

the effect of location. For nitrate, orthophosphate, and general and fecal coliforms, forestland and edge of valley

generally did not statistically differ from each other, whereas a significant shift occurred once water was in the

interior of the valley (>8 km downstream) (Fig. 7). This pattern did not hold for sediments.

We note that the 8 km mark, where percent of overall watershed generally drops to less than 80%, corresponds

with the previous analysis of all Shenandoah valley sites indicating the biggest change in parameters likewise

occurring with this change from heavily forested to moderately forested watersheds (Fig. 3). The water quality

benefits of forested land continue for some distance into agricultural valley land, but this likely depended on

there being a large area of forest in the drainage basin (thus the overall watershed was still mostly forest). If there

isn't a large area forest (such as some of the watersheds in the valley represented by sampling points in the

overall Shenandoah watershed analysis), there are sites of higher water quality, but they are apparently

determined by other factors besides forest cover.

Fig. 7. Binned water quality parameters from transects, characterized by sites in forestland, sites within 8km river miles

downstream from the forest/valley transition (valley edge), and sites more than 8km downstream of the transition (valley

interior). Values are mean ± standard deviation; significant differences between categories indicated by letters on error bars

($p<0.05$).

Finally, although sediment, nitrate and orthophosphate were lower in mountain regions, there were still levels of

these parameters (and coliforms) that would be considered impaired, indicating forested regions are still a source

of these pollutants at times. Orthophosphate and nitrate were in the medium to high stress level (0.05 ppm for

phosphate, and 1 ppm for total nitrogen) in 17.6% and 13.5% of samples, respectively, at forest sites. Comparable

percentages in the interior of valley samples were 45.2% of orthophosphate and 61.0% of nitrate.

Discussion

The current results confirm what is generally understood about water quality in naturally vegetated (in this case

forested) compared to agricultural areas – water quality is consistently better in watersheds which are well

covered with natural vegetation such as forests [5], and agricultural lands can have water quality impacts [16]. It is

evident, however, that within this general trend is important variability that occurs at localized areas. This

variability is important not only for local stream health, but also collectively drives downstream impacts. Further,

many waterways transect both forested and agricultural areas, and the interaction between these two is less well

characterized. By looking at a large public dataset, and focusing in on several representative watersheds, this

study identifies some distinct patterns of water quality in a large watershed with distinct forest and agricultural

segments.

1. Key Water Quality Parameters in Forested vs Agricultural Land

When analyzed across the entire Shenandoah watershed, the clearest pattern of forested vs agricultural land

occurred for nutrients (orthophosphate and nitrate) and conductivity. As key drivers of the downstream health of

aquatic ecosystems, and in this case specifically as central to the major efforts to restore Chesapeake Bay health,

phosphorus and nitrogen are a focus of management efforts in upstream watersheds such as the Shenandoah. In

fact, the Shenandoah/Potomac watershed system is one of the most impacted in Virginia for these nutrients [17].

In our study, both transect sampling and analysis of nutrients in the water quality portal database indicated a

general pattern of higher nutrient levels in agricultural regions that clearly agrees with the well-studied potential

of agricultural practices to impact water quality through anthropogenic activities. We did not explore specific

drivers of nutrient dynamics in the agricultural area, but note that the relationship between agricultural practices

and nutrient levels is complex. For instance, the large scale analysis of Metson et al. [18] found that there was no

single explanatory variable that determines riverine phosphorus levels, and other studies have found variable

impact of best management practices [19].

We note here three observations on nutrient trends from our data. First, our results are roughly consistent with

those found in other studies that focused on this region, both in average concentrations [17, 20, 21], and in the

presence of significant variation on the local level [8]. Second, many of these values were above estimated

threshold criteria for algal and benthic macroinvertebrate impairment [22], and are outside of the EPA suggested

criteria for ecological health [23]. Mean nutrient values for mountain and valley regions were in the low and

medium categories for probability of stress to aquatic organisms, respectively, according to Virginia stressor

analysis [17]. Third, values in areas of low impact (forested regions in the mountains) also reached levels of

concern at times, consistent with Smith et al [8] who found large variation in natural background of phosphorus

and nitrogen, with many stream and river segments in this ecoregion exceeding EPA-proposed criteria.

Conductivity was also clearly different between mountains and valley, however this reflects the strong correlation

with underlying geology rather than land cover differences. Although there are human activities of concern for

conductivity (such as road salting), studies in nearby Appalachian mountain areas point towards agricultural

having relatively low effect on conductivity [24], and conductivity is the parameter of least concern as a stressor in

the region [14,17]. In our study, conductivity then effectively acts as a tracer of water source, with low

conductivity indicating water originating from silicoclastic (mountain) areas, and high conductivity indicating an

origin or longer residence time in limestone or dolostone (valley) areas. In all cases, the mean conductivity was

below 300 μS/cm, generally recognized as the range of no or low probability for stress to aquatic life [17,23,25].

However, some individual samples fell in the high stressor categories (>500 μS/cm), and could be a driver of which

organisms make up aquatic communities in these areas. These high conductivity values occurred during low-flow

periods in valley areas only, usually during mid-summer.

Two other measured parameters – sediment (as total suspended solids) and bacteria – are of major concern

nationwide as drivers of water quality, and have clear associations with agricultural activity. For instance,

sediment is the largest cause of invertebrate-impaired waters nationwide [25], and bacteria is the most common

reason for streams receiving an impairment designation in Virginia [17]. Although these parameters also showed

some patterns of higher levels in agricultural lands, the pattern was less consistent than for nutrients and

conductivity. Fecal and general coliforms were higher in agricultural segments of the watershed in our transect

measurements, but *E. coli* across the watershed (from the water quality portal database) did not correlate as

clearly with forest vs agricultural areas. Conversely, sediments (TSS) were highly variable in the transect data and

did not correlate with land cover, but TSS across the watershed was higher in drainage basins with less forest

cover. Overall, this points to some higher sediment and bacteria values in forested land (see discussion below),

and/or some waterways in primarily agricultural regions that are less impacted by sediments and bacteria. Thus,

although forest cover and agricultural activities are critical for water quality, other factors are driving significant

local variation in these parameters.

2. Water Quality Impacts in Forested Land

Although water quality was better in forested land, our data indicated that nitrogen, orthophosphate, TSS and

indicator bacteria all exceeded levels of concern at particular times and locations. These results are consistent

with Bricker et al [7] who likewise found similar nitrate levels in a nearby mountain area, and with Price and Leigh

[26] who found TSS levels in forested areas of the southern Appalachians as high or higher than the levels

throughout our study area. Smith et al's [8] nationwide study also found some sites and times with levels of

concern in relatively unimpacted areas of our ecoregion.

We did not specifically investigate the source of these pollutant levels in forestlands, although our observations

pointed towards three general categories of possible human-related contributors: 1) existing stable structures, 2)

various types of disturbance, and 3) legacy effects. The contribution of existing stable structures is indicated by

consistently high levels of measured parameters (except conductivity) in retention structures such as a small pond

located in a clearing at the headwater of the North River. This pond was in a small sedimented bowl on the

ridgeline, and is characteristic of numerous small retention basins purposefully maintained in the forest (usually

for supporting game animals). Larger reservoirs are also present in all three watersheds for the purpose of flood

control and municipal water supplies. Recent evidence suggests that ponds can serve as nutrient sinks in

agricultural areas [27]; to our knowledge, there are no similar studies on the water quality role of ponds or lakes

in forested areas.

Disturbance is recognized as critical for mobilization of nutrients and sediments. For instance, forests in this area

are considered to show low nitrogen export unless disturbed [28]. While heavily forested, there are a variety of

processes which impact forests in a way that could mobilize nutrients and sediments. These regions have a

network of roads which are used for recreation, forest service activities (*e.g.* prescribed burns or firefighting), and

resource extraction (*e.g*. timbering). These roads often parallel waterways, are usually unpaved, and in some

areas have low-water crossings rather than bridges. Roads are a small fraction of the total area, but potentially a

significant contributor to water quality, and deserve future attention to understand their role forested regions.

Timber harvesting and prescribed burns are regular localized activities that can have significant impacts on

streams. In addition, our observation is that some streams in the forest areas have eroded banks, either part of

natural processes or as a legacy of alterations previously made, which would be a likely source of nutrients and

sediment. Bare soil from both management activities or streambank erosion are known drivers of water quality

and quantity in forested areas [3,29], and thus are likely contributors to the occasionally high levels of nutrients

and sediments in our study.

Candidate drivers of some water quality parameters include not only present day activities, but also legacy

activities. In addition to management activities occurring over the course of many decades, this area was heavily

logged in the late 1800's, and then used for livestock grazing before forests were restored in the early 1900's. The

role of legacy nutrients as drivers of watershed nutrients has received increased attention as a key driver of water

qaulity (*e.g.* [30]) as studies demonstrate their importance in a variety of watersheds [31-33]. Legacy nutrients

probably have variable residence times depending on local conditions, but are likely on the order of many

decades [33,34]. Mobilization from soil is particularly important during storm events following freeze-thaw cycles

[32], such as the spring rains often experienced in this central Appalachian region. Although these observations all

point toward a potential role of legacy nutrients in forested land, as of the present there is no direct data to guide

an understanding of how much legacy nutrients currently contribute to elevated nutrients in this area.

Finally, other "natural" processes can impact water quality in less disturbed watersheds. Atmospheric deposition

is shown to be important for both nitrogen [8] and phosphorus [35] in relatively undisturbed landscapes. Insect

outbreaks can dramatically increase nitrate from vegetation inputs, as shown in studies from Shenandoah

National Park, which borders the eastern side of this watershed [28]. Wildlife have been shown to be significant

sources of fecal coliforms in watershed analyses [36], and are likely a significant source of this parameter in our

forested areas.

In summary, although a national survey of landscapes found forests with the highest water quality among all land

cover types [5], a narrative that waterways in forested landscapes as invariably "clean" masks some interesting

localized events, and should be distinguished from the general observation that waterways are cleaner on

average. In fact, the recent study identifying forests as having the best water quality also emphasized

considerable variation in sediment and nutrient levels [5]. Forestlands in this region are a complex amalgam of

natural and human processes – as would be the case for most forests in the United States (and indeed globally).

Further studies should extend the general forest vs nonforest paradigm to focus on what categories of activities

impact water quality more or less.

3. Patterns of Water Quality Transition as Rivers flow from Forested to Agricultural Lands

Although this area is notable for a sharp forest/agriculture transition, transect samples indicated water which

flowed out of forested land did not show an immediate change in water quality parameters. Rather, there was a

clear lag period as the water quality parameters remained relatively unchanged until about 8 river km

downstream of the forest boundary (Figs 6,7). Although residence time (in this case, time in an agricultural area)

is a known driver of water quality [16], residence alone would more likely produce a gradual degradation of water

quality. Rather, the decrease in water quality was fairly abrupt, most notably for conductivity, which would track

less with agricultural activities directly and more with underlying geology. When percent forest area of

watersheds for each sampling point were calculated, forested area for the corresponding drainage basins also

showed a sharp decrease at the 8 km mark to about 70-80%. WQP database results were similar, with

parameters generally showing the largest drop in quality when forested areas dropped below 80% (Fig 3).

This is the point at which tributaries draining mostly agricultural areas become a larger part of the total drainage

basin. In turn, this suggests that cumulative inputs from agriculturally derived tributaries are not substantial until

the river is 8km into the agricultural area, and that the pattern may be driven by inputs from the broader tributary

watershed rather than gradual inputs from agricultural activities immediately adjacent to the river. Thus, it seems

that the first 8-10km remains unimpacted because the drainage area from non-forested regions is low. Once

tributaries join, the water quality begins to drop. Furthermore, while the benefits of the forested region stretched

well into the agricultural region, it took only 20-30% of a watershed to be nonforested in order for water quality

to be impacted. Price and Leigh [26] likewise found that even moderate changes in the degree of forestation can

have impacts on water quality variables. In aggregate, these data imply that 1) the value of forest in preserving

water quality is substantial, and extends downstream out of the forested area, but 2) that the ecological service of

maintaining water quality drops off once 20-30% of a total watershed was agricultural.

Although we did not focus on temperature, it is a parameter of critical concern in the area, as long-term

measurements in Shenandoah National Park show an increase that is expected in the presence of climate change

[38]. Although we measured water temperature, interpretation of single temperature readings taken at different

times of the day and season limit conclusions from the transects. However, a rough estimate of transect trends

can be estimated by standardizing individual readings as the deviation from the average of a sampling run along a

transect. Temperature showed a more gradual increase across the length of the transect. Temperature steadily

rose as water moves from headwaters to the end of the sampling transect (supplemental data), contrasting with

the abrupt transitions seen in other parameters. The slow increase would be expected for temperature which is

driven by complex energy exchanges that rely on factors (such as time with exposure to sun) beyond land cover

[39].

Conclusions

The current study confirms the crucial role of substantial forested areas regions to the water quality of

watersheds. While the importance of headwater regions to downstream water quality is recognized [10,40], this

study combined multiple datasets to explore how water quality changes with the transition of waterways from

forested areas, which generally protect water quality, to agricultural areas, which generally degrade water quality.

Key insights which add detail to the general understanding of these landscapes are: 1) substantial variability in

water quality in both agricultural and forest areas points towards important local factors governing pollutants, 2)

while forests clearly had higher water quality than agricultural areas, a narrative of pristine water in forested

areas does not reflect the reality of parameters reaching levels of concern at certain locations and times, and 3)

forested regions provide benefits substantial distances downstream of transitions to agricultural regions, but

water quality rapidly decreased when cumulative forest cover of a corresponding watershed dropped by 20 to

30%. These results contribute to a general understanding of what drives water quality in mosaiced landscapes

such as the Chesapeake Bay watershed, which has shown complex and variable responses to management efforts

[41].

While the current study helps draw a more complete picture of water quality across a diverse landscape, we did

not investigate several other factors that are critical to understanding when and how degradation of water quality

occurs – notably the contribution of storm events and seasonal cycles. Flow rates, for instance, are a known and

crucial determinant of loading [10, 42, 43], and nutrients can follow seasonal trends [7]. Preisendanz et al [44]

argue that "hot moments" are as important as "hot spots" – and that insufficient attention has been directed

towards these temporal events as opposed to the spatial pattern. Whether the patterns seen here represent

those found during peak storm flow, and across different seasons, is a further avenue of fruitful study, particularly

in light of future climate changes [45]. Future increases in precipitation are highly relevant given that downstream

water quality in the Chesapeake Bay is tied closely to flow rates [10], and recent estimates are that climate change

threatens to increase nutrient and sediment loads in the Chesapeake Bay watershed even more than changes in

agricultural intensity, population or development [46].

References

1. Palmer MA, Hondula KL, Koch BJ. (2014) Ecological restoration of streams and rivers: shifting strategies and

shifting goals. Annual Review of Ecology, Evolution, and Systematics. 23;45(1):247-69.

2. Richardson K, Steffen W, Lucht W, Bendtsen J, Cornell SE, Donges JF, Drüke M, Fetzer I, Bala G, Von Bloh W,

Feulner G. (2023) Earth beyond six of nine planetary boundaries. Science advances. 13;9(37):eadh2458.

3. Neary DG, Ice GG, Jackson CR. (2009) Linkages between forest soils and water quality and quantity. Forest

ecology and management. 30;258(10):2269-81.

4. Clinton BD, Vose JM. (2006) Variation in stream water quality in an urban headwater stream in the southern

Appalachians. Water, air, and soil pollution. 169:331-53.

5. Caldwell PV, Martin KL, Vose JM, Baker JS, Warziniack TW, Costanza JK, Frey GE, Nehra A, Mihiar CM. (2023)

Forested watersheds provide the highest water quality among all land cover types, but the benefit of this

ecosystem service depends on landscape context. Science of The Total Environment. 15;882:163550.

6. Richardson, J. S. (2019). Biological diversity in headwater streams. Water, 11(2), 366.

7. Bricker, Suzanne & Rice, Karen & III, Owen. (2014). From Headwaters to Coast: Influence of Human Activities on

Water Quality of the Potomac River Estuary. Aquatic Geochemistry. 20. 10.1007/s10498-014-9226-y.

8. Smith, R. A., Alexander, R. B., & Schwarz, G. E. (2003). Natural background concentrations of nutrients in

streams and rivers of the conterminous United States.

9. Ator SW, Blomquist JD, Webber JS, Chanat JG. (2020) Factors driving nutrient trends in streams of the

Chesapeake Bay watershed. Journal of Environmental Quality.49(4):812-34.

10. Murphy RR, Keisman J, Harcum J, Karrh RR, Lane M, Perry ES, Zhang Q. (2021) Nutrient improvements in

Chesapeake Bay: Direct effect of load reductions and implications for coastal management. Environmental Science

& Technology. 21;56(1):260-70.

11. Zhang Q, Blomquist JD, Fanelli RM, Keisman JL, Moyer DL, Langland MJ. (2023) Progress in reducing nutrient

and sediment loads to Chesapeake Bay: Three decades of monitoring data and implications for restoring complex

ecosystems. Wiley Interdisciplinary Reviews: Water. 10(5):e1671.

12. Clune, J.W., and Capel, P.D., eds., (2021), Nitrogen in the Chesapeake Bay watershed—A century of change,

1950–2050: U.S. Geological Survey Circular 1486, 168 p., https://doi.org/10.3133/cir1486.

13. Caraco, N. F., & Cole, J. J. (1999). Human impact on nitrate export: an analysis using major world rivers. Ambio,

28(2), 167-170.

14. Zheng L, Gerritsen J, Beckman J, Ludwig J, Wilkes S. (2008) Land use, geology, enrichment, and stream Biota in

the Eastern Ridge and Valley Ecoregion: implications for nutrient criteria development 1. JAWRA Journal of the

American Water Resources Association. 44(6):1521-36.

15. Mwaijengo GN, Msigwa A, Njau KN, Brendonck L, Vanschoenwinkel B. (2020) Where does land use matter

most? Contrasting land use effects on river quality at different spatial scales. Science of the Total Environment.

1;715:134825.

16. Farthing, T., Rintsch, E., Larson, O., Grudzinski, B.P., Fisher, T.J., & McCarty, J.L. (2024). Effects of a forested

state park on stream nutrient concentrations in an agriculturally dominated watershed in the U.S. Midwest.

JAWRA Journal of the American Water Resources Association.

17. [VDEQ] Virginia Department of Environmental Quality. 2017. Stressor analysis in Virginia: Data collection and

stressor thresholds. VDEQ Technical Bulletin WQA/2017-001. Richmond (VA): VA Department of Environmental

Quality Water Quality Monitoring, Biological Monitoring, and Water Quality Assessment Programs. 478 p.

https://www.deq.virginia.gov/home/showpublisheddocument/4313/637461491358800000

18. Metson, G. S., Lin, J., Harrison, J. A., & Compton, J. E. (2017). Linking terrestrial phosphorus inputs to riverine

export across the United States. Water research, 124, 177-191.

19. Ator SW, Blomquist JD, Webber JS, Chanat JG. (2020). Factors driving nutrient trends in streams of the

Chesapeake Bay watershed. Journal of Environmental Quality. 49(4):812-34.

20. Omernik, J., Paulsen, S., Griffith, G., & Weber, M. (2016). Regional patterns of total nitrogen concentrations in

the National Rivers and Streams Assessment. Journal of Soil and Water Conservation, 71(3), 167-181.

21. Clune JW, Crawford JK, Boyer EW. (2020) Nitrogen and phosphorus concentration thresholds toward

establishing water quality criteria for Pennsylvania, USA. Water. 17;12(12):3550.

22. Evans-White, M. A., Haggard, B. E., & Scott, J. T. (2013). A review of stream nutrient criteria development in

the United States. Journal of environmental quality, 42(4), 1002-1014.

23. USEPA. (2000). Ambient Water Quality Criteria Recommendations Information Supporting the Development

of State and Tribal Nutrient Criteria Rivers and Streams in Nutrient Ecoregion XI, USEPA-822-B-00–020. USEPA,

Washington, DC

24. Cormier, S. M., Wilkes, S. P., & Zheng, L. (2013). Relationship of land use and elevated ionic strength in

Appalachian watersheds. Environmental Toxicology and Chemistry, 32(2), 296-303.

25. Govenor H, Krometis LA, Willis L, Angermeier PL, Hession WC. (2019) Macroinvertebrate sensitivity thresholds

for sediment in Virginia streams. Integrated Environmental Assessment and Management. 15(1):77-92.

26. Price, K., & Leigh, D. S. (2006). Comparative water quality of lightly-and moderately-impacted streams in the

southern Blue Ridge Mountains, USA. Environmental monitoring and assessment, 120, 269-300.

27. Liu W, Li S, Shen W, Zhuang Y, Li X, Ling F, Zhang L. (2024) Small water bodies influence river water quality in

agricultural watersheds. Agricultural Water Management. 1;304:109070.

28. Eshleman, K. N., Fiscus, D. A., Castro, N. M., Webb, J. R., & Deviney Jr, J. (2001). Computation and visualization

of regional-scale forest disturbance and associated dissolved nitrogen export from Shenandoah National Park,

Virginia. The Scientific World Journal, 1, 539-547.

29. Gellis AC, Gorman Sanisaca L. (2018) Sediment fingerprinting to delineate sources of sediment in the

agricultural and forested Smith Creek watershed, Virginia, USA. JAWRA Journal of the American Water Resources

Association. 54(6):1197-221.

30. Golden HE, Evenson GR, Christensen JR, Lane CR. (2023) Advancing watershed legacy nitrogen modeling to

improve global water quality. Environmental science & technology. 17;57(7):2691-7.

31. Cravotta III CA, Tasker TL, Smyntek PM, Blomquist JD, Clune JW, Zhang Q, Schmadel NM, Schmer NK. (2024)

Legacy sediment as a potential source of orthophosphate: Preliminary conceptual and geochemical models for the

Susquehanna River, Chesapeake Bay watershed, USA. Science of the Total Environment. 20;912:169361.

32. Inamdar S, Johnson E, Rowland R, Warner D, Walter R, Merritts D. (2018) Freeze–thaw processes and intense

rainfall: the one-two punch for high sediment and nutrient loads from mid-Atlantic watersheds. Biogeochemistry.

141:333-49.

33. Van Meter, K. J., Van Cappellen, P., & Basu, N. B. (2018). Legacy nitrogen may prevent achievement of water

quality goals in the Gulf of Mexico. Science, 360(6387), 427-430.

34. McDowell R, Dodd R, Pletnyakov P and Noble A (2020) The Ability to Reduce Soil Legacy Phosphorus at a

Country Scale. Front. Environ. Sci. 8:6. doi: 10.3389/fenvs.2020.00006

35. Stoddard JL, Van Sickle J, Herlihy AT, Brahney J, Paulsen S, Peck DV, Mitchell R, Pollard AI. (2016) Continental-

scale increase in lake and stream phosphorus: are oligotrophic systems disappearing in the United States?.

Environmental science & technology. 5;50(7):3409-15.

36. Nguyen KH, Senay C, Young S, Nayak B, Lobos A, Conrad J, Harwood VJ. (2018) Determination of wild animal

sources of fecal indicator bacteria by microbial source tracking (MST) influences regulatory decisions. Water

research. 1;144:424-34.

37. Varela WL, Mundahl ND, Bergen S, Staples DF, Cochran-Biederman J, Weaver CR. (2023) Physical and

biological stream health in an agricultural watershed after 30+ years of targeted conservation practices. Water.

2;15(19):3475.

38. Jastram, J.D., Snyder, C.D., Hitt, N.P., and Rice, K.C., (2013) Synthesis and interpretation of surface-water

quality and aquatic biota data collected in Shenandoah National Park, Virginia, 1979–2009: U.S. Geological Survey

Scientific Investigations Report 2013–5157, 77 p., http://pubs.usgs.gov/sir/2013/5157/.

39. Leach JA, Kelleher C, Kurylyk BL, Moore RD, Neilson BT. (2023) A primer on stream temperature processes. Wiley Interdisciplinary Reviews: Water. 10(4):e1643.

40. Alexander RB, Boyer EW, Smith RA, Schwarz GE, Moore RB. (2007) The role of headwater streams in downstream water quality 1. JAWRA Journal of the American Water Resources Association. 43(1):41-59.

41. Webber, J.S., Chanat, J.G., Clune, J.W., Devereux, O., Hall, N., Sabo, R.D., & Zhang, Q. (2024). Evaluating water-quality trends in agricultural watersheds prioritized for management-practice implementation. JAWRA Journal of the American Water Resources Association. https://doi.org/10.1111/1752-1688.13197.

42. Lewis, W. M., Melack, J. M., McDowell, W. H., McClain, M., & Richey, J. E. (1999). Nitrogen yields from undisturbed watersheds in the Americas. New Perspectives on Nitrogen Cycling in the Temperate and Tropical Americas: Report of the International SCOPE Nitrogen Project, 149-162.

43. Ezzati G, Kyllmar K, Barron J. (2023) Long-term water quality monitoring in agricultural catchments in Sweden: Impact of climatic drivers on diffuse nutrient loads. Science of the Total Environment. 15;864:160978.

44. Preisendanz HE, Veith TL, Zhang Q, Shortle J. (2020) Temporal inequality of nutrient and sediment transport: a decision-making framework for temporal targeting of load reduction goals. Environmental Research Letters. 18;16(1):014005.

45. Yang Y, Tilman D, Jin Z, Smith P, Barrett CB, Zhu YG, Burney J, D'Odorico P, Fantke P, Fargione J, Finlay JC. (2024) Climate change exacerbates the environmental impacts of agriculture. Science. 6;385(6713):eadn3747.

46. Bhatt G, Linker L, Shenk G, Bertani I, Tian R, Rigelman J, Hinson K, Claggett P. (2023) Water quality impacts of

climate change, land use, and population growth in the Chesapeake Bay watershed. JAWRA Journal of the

American Water Resources Association. 59(6):1313-41.

Supplemental Data, Figure S1

Fig. S1. Temperature differential gradient parameters along river transects, from forest headwaters to agricultural

bottomland. Temperature differential for each sample calculated as site temperature minus average temperature across its

run (samples for the one river and one date). Blue, orange and green points represent parameter mean ± standard deviation

values for the North River, Dry River, and Briary Branch, respectively. Asterisks with higher temperature differentials are

pond or reservoirs.