Headwaters to valley: Water quality in rivers transitioning from forest

to agricultural bottomland

Short title: Water quality in forest and agricultural lands.

Doug Graber Neufeld[1], Isaac Alderfer[1], Zachary Bauman[1], Micah Buckwalter[1]*

[1]Department of Biology, Eastern Mennonite University, Harrisonburg, Virginia, United States of America

*Corresponding author

E-mail: neufeldd@emu.edu

Abstract

Many waterways flow out of forestlands, which tend to maintain higher water quality, into agricultural lands,

which tend to degrade water quality. The roles of land cover in impacting key water quality parameters

(phosphorus , nitrogen, total suspended solids, bacteria, and conductivity) were investigated for the watershed of

the North and South Fork of the Shenandoah River, Virginia. This area has a particularly sharp boundary between

heavily forested and heavily agricultural regions. Two datasets were analyzed: 1) a large number of datapoints

spanning a 20-year range in the Water Quality Portal (WQP) database, and 2) transects along three representative

rivers systems over the span of a 4-year period. All parameters trended better in forested regions than

agricultural regions. This was particularly true for nitrogen and conductivity; phosphorus, TSS and bacteria

showed more local variability, especially in the agricultural region. Periods of high flow increased phosphorus,

sediment and bacteria concentrations, and decreased conductivity, but not when drainage basin forest cover was

less than 80%. Transects showed that waterways flowing out of forestland maintained higher water quality for

approximately the first 8 km in agricultural land. Both transect and WQP data indicated higher water quality when

the percent of forested land cover in a drainage basin was about 70-80%. Thus, forestland does mitigate the

impacts of agriculture on water quality to some degree, but this effect rapidly diminishes as forest cover of the

watershed lessens. Furthermore, forests themselves have degraded water quality at certain times and places; for

instance, nutrients level were in the medium to high stress level for aquatic life in approximately 15% of samples.

This study illustrates general trends of land cover effects on water quality, while also highlighting both site-

specific variability, and the dynamics of water quality as water flows out of forested areas into agricultural areas.

Introduction

Aquatic ecosystems from headwaters to larger waterways downstream have been a major focus of efforts at

maintaining and improving environmental quality, due to their combined importance in providing ecosystem

services, and their integral role in the lives of people [1]. Indeed, maintenance of water quality is directly related

to important planetary boundaries such as the geochemical cycling of nitrogen and phosphorus [2], which have

clearly exceeded safe operating zones. Numerous studies have explored key water quality parameters elucidating

both patterns that are generalizable across waterways, and patterns that are specific to local spatial and temporal

conditions.

Nutrients (nitrogen and phosphorus), sediments, and bacteria are recognized as key parameters determining both

local aquatic health, and downstream impacts [3,4]. While water quality studies have elucidated key insights on

the dynamics of these pollutants in watersheds, these studies also point towards a continued need to understand

these dynamics better. This need is highlighted by observations that concentrations of these pollutants generally

remain high, despite long-running efforts to reduce the loading of these water contaminants through

management actions. For instance, a recent study looked at thousands of water sample measurements found that

the high nutrients were due to the presence of persistent sources, including the potential role of legacy sources or

inadequate current management practices [5]. That study highlighted the role of local ecological conditions as

potentially key to controlling watershed level pollution, and confirmed the important role of headwaters in

determining downstream concentrations. Studies using large datasets have noted the sustained increase at the

continental scale, such as the study of Stoddard et al. [6] finding the percentage of stream length with <10ug/L

total phosphorus (TP) decreased from 24.5 to 1.6% in the 2004 to 2014 period. Likewise, more focused use of

local datasets note a similar lack of progress at the level of local watersheds, such as the study on three

Chesapeake Bay watersheds finding a lack of improvement in nutrient and sediment levels despite the

implementation of multiple management practices [7]. The slow progress in improving water quality highlights

the need to for continued understanding of sources, transport, fate and impacts of these pollutants. In particular,

understanding the key drivers of these processes, and the accompanying management practices to mitigate water

quality impacts, is an ongoing need.

Land cover, and especially continued pressure from human activities that alter land cover, are fundamental in

driving water quality of watersheds. Most notable is how natural vegetation such as forests can improve water

quality, while agricultural activities (both pastureland and cropland) can degrade water quality through a variety

of processes. In the Chesapeake Bay watershed, agriculture is the primary suspected cause of degraded water

quality [8], whereas forest cover can improve water quality [9]. Protection of forested lands is generally seen as

crucial to maintaining clean water [10,11], notably illustrated by a recent study in the southeast United States

finding that forests had the highest water quality of all landscape types [12]. In parallel to land cover effects,

headwater regions (which often are more forested) have unique characteristics compared to downstream waters

[13-15]. There are clearly multiple factors beyond just land cover type that control water quality [12], and more

information is needed on how water quality changes as waterways traverse different landscapes. In particular, it

is not clear whether there is a sharp transition in water quality as it traverses forested and agricultural land, and

how the impacts of the two landscapes combine to influence downstream water quality.

Watersheds draining into the Chesapeake Bay are the focus of a long running effort to understand how stressors

such as nutrients and sediments contribute to estuarine health, and how these inputs can be reduced [16-18].

Efforts in the Chesapeake watershed have been mixed, illustrating both the ongoing challenge of reducing

agricultural inputs, and the need for more information on what exactly drives nutrient trends [16, 18, 19]. Land

cover modification has been highlighted as one of the key drivers in the Chesapeake Bay watershed [9]. The

Shenandoah River watershed of Virginia and West Virginia is a major tributary system of the Chesapeake Bay, and

stands out as having both some of the region's largest protected forestlands, and also the highest density of

livestock and poultry operations in Virginia, along with significant cropland. Furthermore, there is a generally

distinct boundary between the relatively flat valley bottomland which is dominated by agriculture and

predominantly of limestone origin, compared to the surrounding mountains which are heavily forested, federally

protected as national forest and national park land, and predominantly of sandstone origin. This watershed

therefore represents a unique opportunity to study a sharp transition of waterways between heavy forestland of

headwaters, and intensive agriculture downstream.

The present study is unique in combining data from a large public database (Water Quality Portal; WQP) with

targeted sampling on representative river systems to understand the dynamics of key water contaminants in a

headwaters watershed of the Chesapeake Bay with distinct regions of land cover. We used this system to focus on

several interrelated questions that add novel insights to ongoing efforts at understanding nutrient, sediment and

bacteria dynamics: 1) What characterizes water quality in heavily forested headwater regions? 2) How does this

water quality change when transitioning from forested to agriculturally intensive lands? and 3) What are

similarities and differences between different water quality parameters in this pattern of landscape?

Methods

Ethics Statement

Site access outside of national forest was either at publicly accessible right-of-way along state or county highways (usually bridges), or by landowner permission. Within the national forest, permission for site access for the purpose of water sampling was granted by the US Forest Service.

North and South Fork Shenandoah River Watershed

Water quality data for the combined watersheds of the North and South Forks of the Shenandoah River watershed were accessed from the Water Quality Portal (WQP; https://www.waterqualitydata.us/), an open access compilation of water quality results from multiple government, academic and nonprofit organizations. Additional publicly available data was added from the Friends of the Shenandoah River (https://fosr.org/state-of-the-river/compiled-water-quality-data/), which runs a Virginia state approved laboratory and has collected significant water quality data for the region. Since land downstream of the confluence of the North and South Forks (north of Front Royal, Virginia) does not have protected National Forest or National Park land (as dominates forestland upstream from this point), we did not analyze the watershed of the Shenandoah River downstream of the confluence. Samples from within the delineated watershed were selected for the 20 year period from Jan 1, 2004 to Dec 31, 2023. Since most sites had multiple measurements of a parameter for that period, we used the mean parameter value for each site.

In order to separate the effects of flow (*e.g.* base flow vs storm), we analyzed data from USGS water stations (USGS National Water Dashboard; https://dashboard.waterdata.usgs.gov/app/nwd/en/) at the base of the North Fork (#01634000) and South Fork (#01635500). Although flow at water sample points on subwatersheds depends on the local conditions, these gauges roughly correlated with flow at gauges upstream in both watersheds

(upstream North Fork #01632000 correlated with an r^2 of 0.55, and upstream South Fork #01620500 correlated

with an r^2 of 0.42), suggesting they are representative of the watershed. Thus, in the absence of flow data at all of

the many water sample points, the downstream flow rates provided an estimate of when there were high flow

periods throughout the watershed. Water sample measurements were selected as "high flow" when the

corresponding large watershed (North or South Fork) had a discharge in the 75th percentile or above for the

corresponding USGS water gauge (percentiles reported by USGS for a 99 year period). Most of these times (80%

and 77% in North and South Forks, respectively) occurred during the December to May period. The remaining

samples (below the 75th percentile) were designated "low flow".

Outliers of parameters were identified and removed for samples having Cook's distance of 4/n (0.005 for this

dataset), and p<0.05. The cause of outlying data was not known, it could have been inaccurate measurements or

genuinely unusual high or low levels that only occurred occasionally. Given the varied sources of data in the

Water Quality Portal and large number of samples, removal of these outliers ensured that occasional quality

control issues (inaccurate measurements or mis-recorded data), or unusually large fluctuations of parameters did

not greatly skew measurements representing the majority of water flow periods for a waterway.

Spatial analysis was performed using ArcGIS Pro 3.3 and QGIS 3.38.0. Since in other aquatic systems, water quality

is shown to be driven by overall land use across the entire drainage basin for a collection site [20], we delineated

watersheds from USGS 1/3 arc-second Digital Elevation Maps (3DEP product in The National Map). The digital

elevation maps were processed in QGIS to fill and remove sinks, followed by flow direction and accumulation

calculations. The unnest watersheds function (Whitebox) in QGIS was used to delineate watersheds representing

the entire drainage basin for each water sample point. Land use/land cover (LULC) from the USGS National Land

Cover Database was compared between 2003 and 2021, and showed a low degree of change in forested vs

unforested land (in either direction) during this time (0.96%). We therefore used the 2021 land cover data set (Fig

1a) for calculating percent forest in all watersheds, rather than attempting to match year of water collection with

the respective year of LULC data. Geology was mapped as Generalized Lithography in the State Geologic Map

Compilation (SGMC) Geodatabase of the Conterminous United States (ver. 1.1, August 2017), and was categorized

as karst or non-karst geology (Fig 1b). Soil Erodibility was mapped as the K factor from the USA soil service

geographic database (SSURGO) of the Natural Resources Conservation Service (accessed as ArcGIS Online layer)

(Fig 1c). The percent forestland and karst, and the average erodibility, in a drainage basin represented by sample

sites reflects both site location (for instance, whether it is in a forested vs nonforested area), and the area

upstream draining into the point. Water quality values were compared to watershed forest coverage, karst

coverage, and average soil erodibility (K factor) using multiple linear regression. Further comparisons of

parameters in discrete forest coverage categories (in 20% increments) were performed with a Kruskal Wallis and

Dunn's post-hoc test (non-parameter statistics were used due to unequal variances in categories).

A. Land Cover

B. Erodibility

C. Underlying lithology

Fig 1. Land cover (A), erodibility (B) and underlying lithology (C) in study area. These three parameters

represent candidate drivers of water quality, and highlight the difference between the valley area (central,

running southwest - northeast), and the mountain areas (on the western and eastern edges, and in the central

Massanutten ridge region). Study area represents watersheds of the North and South Forks of the Shenandoah

River.

To calculate a general forest boundary in the study area, we calculated the average percent forest within a 200m

radius of each pixel (pixel resolution 25m) of the land cover raster using the focal statistics tool of ArcGIS Pro.

Pixels with at least 75% forest in this radius were counted as forest. This value was chosen as it produced a

smoothed general forest boundary that depicts the visual forest transition as seen on satellite images. River

kilometers for the transect study (see below) were then calculated upstream and downstream from the point at

which the river crosses this boundary.

Cluster and outlier identification was performed in ArcGIS using Anselin Local Moran's I using 999 permutations.

Anselin Local Moran's I is a spatial statistical method used to identify areas of interest in other studies of water

quality [21,22]. Statistically significant clusters and outliers at the 95 percent confidence level are shown as

clusters of high values (HH), clusters of low values (LL), outliers where a high value is mostly surrounded by low

values (HL), and outliers where low value is mostly surrounded by high values (LH).

River Transect Sites

Three river systems were chosen for water sampling as transects – the Dry River, Briery Branch, and the North

River (Fig 2). All three rivers originate in the mountains on the western edge of the Shenandoah watershed,

converge as the lower North River, and then flow together into the South Fork of the Shenandoah River which

joins the Potomac River before entering the Chesapeake Bay. The headwaters are at approximately 3000 feet, and

flow through the oak-hickory forest of George Washington National Forest before abruptly entering the

Shenandoah valley agricultural region. The land cover characteristics are generally similar between the three river

systems, with the overall watersheds dominated by forest and agriculture, with only a small fraction of each

watershed as developed land or wetlands (Fig 2).

Fig 2. Location and characteristics of river transect. Map of river transect sites, located on the western side of the

overall watershed study area, along with land cover categories for the region. Percentages of land cover

categories in each river watershed are shown in the legend, indicating the overall similarity of the three

watersheds used in the transect study.

Water sample sites were selected at approximately 5km intervals, although specific sites were dependent on

availability of river access (see Fig 2). In addition, river sections in the middle interior of the valley dried

completely during periods of low rainfall, especially in midsummer. In other cases of high rainfall or winter, sites

in the national forest were inaccessible for sampling. The final number of samples per site therefore varied.

Samples were collected between March 2020 and July 2024 and were taken at all times of the year (see Table S1

for specific dates). As a general indication of river level in the region, flow discharge was recorded from a USGS

station (#01622000) near Burketown on the North River, and varied from 114 to 1430 ft^3 per second, representing

a range of discharge from 18th to 97th percentiles for the period samples were taken. Samples thus represent a

wide range of seasons and river flow conditions.

At each site, parameters were either measured onsite (temperature and conductivity), or water samples were

collected in dedicated water sample bottles (nitrate, orthophosphate, bacteria, sediments). All measurements or

bottle samples were collected in the main column below the water surface, upstream from any disturbances

(wading or previous sampling). Field blanks were collected at one site per transect for nitrate, orthophosphate,

and bacteria. A field spike bottle was also collected at one site per transect for nitrate and orthophosphate. Field

duplicates were collected for all nitrate, orthophosphate and bacteria samples, and for one sediment sample per

transect. All samples were kept on ice (for transport) or at 4^{o}C storage until analysis.

River Transect Parameters

Water samples were analyzed for sediment, nitrate, orthophosphate, temperature, conductivity, and general and

fecal coliforms. A Quality Assurance Project Plan, previously approved for earlier NFWF and Virginia DEQ grant

sampling, guided analysis, and was based on USGS, Virginia DEQ, or EPA approved procedures (e.g. U.S. EPA 2007.

SW-846).

Sediments were measured gravimetrically as per EPA method 160.2. In brief, 1 L samples were filtered through

pre-rinsed, dried and weighted 47 mm diameter GF/F filters (0.7 μM pore-size), and then re-dried and re-weighed.

Conductivity was measured using a Tracer Pocketester (LaMotte) calibrated to a 84 μS/cm standard. Probe

calibration was verified after each transect to ensure calibration did not drift during the measurement period.

Temperature was measured using a Vernier temperature probe, or the internal thermometer on the conductivity

probe. General and fecal coliforms were measured using Coliform Easygel (Micrology, Inc; Goshen, Indiana, USA).

1 ml samples were plated on agar petri dishes, incubated at $37^{o}C$ until colony color development (generally 36-

48h), and then colony identity determined by color.

Nitrogen and phosphorus were assessed as nitrate and orthophosphate, respectively. While other forms would be

present, these are the most common and relevant forms in these waters. For instance, nitrate is the majority of

total nitrogen (TN) in the region [4]. Likewise, the study of Farthing et al [23] on the effect of forestland on

agriculturally impacted rivers found most nitrogen as nitrate, and most phosphorus as phosphate. Both nitrate

and orthophosphate were measured with a colorimetric assay (Hach; for NitraVer5 and PhosVer3, respectively)

according to manufacturer instructions. Duplicate sample bottles for each site were measured and the mean used

as the final value. Absorbance values at appropriate wavelengths were compared to a calibration series, with

blank and standard tested every 10th sample.

Results

Water Quality Patterns in the North and South Forks of Shenandoah River Watershed

Water quality data for the entire watershed of the North and South Forks of the Shenandoah River were

compared with several candidate drivers of water quality – documented land cover changes (USGS National Land

Cover Database) due to human activities (represented by percent forest coverage), underlying geology

(represented by karst coverage, as a rock type in the area having a large impact on hydrology), and basic soil

composition (represented by erodibility, as a measure of how likely soil constituents are to enter waterways) (Fig

1). Multiple linear regression identified forest coverage as more important for driving water quality, compared to

karst or soil erodibility (Table 1) for all parameters except conductivity, which highly correlated with both forest

coverage and soil erodibility, and moderately correlated with karst. The relatively low adjusted r^2 values point

towards a complex system with a large for additional factors contributing to water quality. None-the-less, the

relationship of parameters to forest coverage was highly statistically significant, whereas it was not for karst or

soil erodibility (aside from conductivity). The predominance of land cover effects compared with measured

geology and soil differences confirms other studies noting the importance of land cover, and further analysis

focused on forest coverage as a major driver of water quality.

Table 1. Multiple Linear Regression of water quality parameters compared with land cover (forest coverage),

geology (karst coverage), and soil (average erodibility) in corresponding watersheds.

Water Quality Parameter	sample size (N)	adjusted r^2	Forest coverage		Karst		Erodibility	
			significance (p value)	coefficient	significance (p value)	coefficient	significance (p value)	coefficient
Nitrate	297	0.59	**p<0.001**	-2.8	p=0.12	-0.41	p=0.65	-0.003
TN	230	0.54	**p<0.001**	-3.34	p=0.06	-0.67	p=0.11	-0.026

Orthophosphate	211	0.07	**p<0.01**	-0.06	p=0.32	-0.02	p=0.91	-0.0001
TP	266	0.07	**p<0.01**	-0.07	p=0.18	-0.07	p=0.30	-0.001
TSS	209	0.22	**p<0.001**	-16.8	p=0.21	-4.83	p=0.40	-0.15
E. coli	265	0.18	**p<0.01**	-769	p=0.87	42.8	p=0.07	-24.6
Conductivity	636	0.75	**p<0.001**	-563	p=0.08	46.1	**p<0.001**	2.71

Mapped concentration averages at the sample sites highlights several general patterns (Fig 3). First,

concentrations of all parameters were consistently lower in forested regions, although there was variability with

some sites having moderate concentrations. Second, the highest concentrations of all parameters were clearly

centered in the valley, although the pattern differed between parameters. For some (nitrate, TN, and

conductivity), the concentrations in the valley were consistently high, whereas for the other parameters they

were more variable in the valley. Third, there was a gradient with the highest values most often located more

centrally in the valley, rather than on the forest edges. In addition, some low concentrations along the valley edge

(the boundary with forests) were associated with waterways entering the valley from the forest. Conductivity

showed the most consistent gradient, showing a gradual increase with distance from the forest into the valley.

Finally, different parameters often showed similar trends at sites (for instance, a site with a high concentration of

one parameter often had high concentrations of other parameters) suggesting generalized water quality issues at

specific sites or areas of the valley.

A. Nitrate

N=403 sites, 9992 cumulative sample measurements

B. Total Nitrogen

N=310 sites, 20388 cumulative sample measurements

C. Orthophosphate

N=288 sites, 9788 cumulative sample measurements

D. Total Phosphorus

N=335 sites, 13565 cumulative sample measurements

F. Total Suspended Solids

N=269 sites, 14673 cumulative sample measurements

F. E. coli

N=391 sites, 13947 cumulative sample measurements

G. Conductivity

N=810 sites, 30876 cumulative sample measurements

Fig 3. Water quality parameters in watershed North and South Forks of Shenandoah River of Virginia. Data are averages for

sample sites for 2004-2023. Sample sizes are given as number of sites, and the total number of samples represented

cumulatively at those sites.

Cluster and outlier analysis (Anselin Local Moran's I) was performed to statistically investigate these spatial

distributions. This confirmed that clusters of low concentrations tended to be located in heavily forested land, but

in some cases were just outside the forestland, or along major waterways that would take significant volumes of

water into the valley (Fig 4; light blue points). Clusters of high values were located in the valley, usually at some

distance from the forested areas (Fig 4; light red points). Cluster analysis also confirmed the more consistent

pattern of nitrogen (nitrate, and TN) and conductivity relative to land cover, whereas other parameters showed

more variability (for instance, with clusters of low concentrations located in the valley). For all parameters,

Anselin Local Moran's I identified some outliers (high values in low cluster areas, dark red points; or low values in

high cluster areas, dark blue points), pointing to significant localized variation. In particular, there were low outlier

sites (Fig 4; dark blue points) scattered in the valley for different parameters, indicating some local areas of higher

water quality among a region with generally low water quality. Outlier high values (Fig 4; dark red points) were

less common, suggesting there were not as many point sources that only impacted a small area – problem areas

tended to be across a stretch of waterway or region (such as the central area of the valley).

A. Nitrate

B. Total Nitrogen

C. Orthophosphate

D. Total Phosphorus

Fig 4. Cluster and outlier analysis of water quality parameters for North and South Fork Shenandoah River, Virginia. "High-High" and "Low-Low" are points in high and low cluster areas, respectively. "High-Low" and "Low-High" are high values in low cluster areas, and low values in high cluster areas, respectively.

Given that parameters may relate nonlinearly to the proportion of forest cover, we further analyzed the

relationship by grouping watersheds into 5 bins representing 20% increments of forest cover. Furthermore, since

storm events are typically major drivers for loads of the water contaminants, we analyzed data separately for high

flow (discharge volumes at or above the 75th percentile), and for low flow (discharge volumes below the 75th

percentile). Statistical analysis (Kruskal Wallis test, followed by post-hoc Dunn's test) indicated significant

differences with forest cover for all parameters (Fig 5) at both high and low flows. The greatest statistical

difference was between the highest category of forest cover (80-100%) and the other forest cover categories. This

was especially evident, for instance, with Nitrate, TP and TSS, which generally had statistically similar

concentrations between the forest cover categories, except for the highest category (80-100%) (Fig 5A,D,E). For

other parameters, such as TN, orthophosphate and *E. coli* (Figs 5B,C,F), there was a somewhat more gradual

decrease with increased forest cover, although there was still notably improved water quality at the highest forest

cover categories. Conductivity showed the most gradual pattern of decreased concentrations with forest cover

(Fig 5G). Overall, these results suggested that any effects of forestland on water quality extend into nonforested

land, that the greatest decrease in water quality for these parameters occurred when rivers moved from heavily

forested land to moderate or low forest cover, and that there is significant localized variation in all parameters.

A. Nitrate

B. Total Nitrogen

C. Orthophosphate

D. Total Phosphorus

Fig 5. Water quality in forest percent categories. Light grey bars are for samples that were below the 75[th] percentile discharge volume, dark grey bars are for samples that were 75[th] percentile or above for discharge volume. Significant differences between categories indicated by letters on error bars. (lower case, $p < 0.05$ for lower 75[th] percentile of discharge volume; upper case, $p < 0.05$ for 75[th] percentile or above of discharge volume).

Comparison of low and high flow conditions indicated that high flow mobilized TP, TSS, and *E. coli*. This was expected, as TP and *E. coli* are known to be associated with sediments, and high flow rates would mobilize sediments through processes such as bank erosion. Conversely, but also unsurprisingly, conductivity decreased in high flow conditions. The effects of high flow were absent or diminished at the higher forest cover categories, consistent with the protective role for forests in buffering flow rates and preventing erosion.

Water Quality in Transects of Three Rivers Flowing from Forest to Agricultural Bottomland

We measured water quality along transects for three specific rivers – the Dry River, Briery Branch, and North River

– located on the western side of the Shenandoah watershed. All three rivers flow from significant distances

through national forest land, and then enter areas primarily used for agriculture (as both pasture and cropland),

allowing us to explore more how water quality changes with the transition from forest to agriculture (Fig 2).

There was an increase in orthophosphate, nitrate, and general and fecal coliforms (Fig 6A,B,D,E) at 8-10 km

downstream of the forest boundary. This pattern was less obvious for TSS (Fig 6C), which varied more at sample

sites, and had some higher concentrations in the forestland. When we examined temperature differentials (how

much temperature differed from the average temperature across the transect for that sampling run), we found a

different pattern – temperature appeared to more gradually increase rather showing the abrupt shift at 8 river

kms (Fig S1). There was significant variation for all variables, indicating (as did the previous analysis of sites across

the Shenandoah watershed; Figs 3,4) that there are significant localized effects that impact water at any particular

point and time.

A. Nitrate

B. Orthophosphate

D. Fecal Coliform

North River • Dry River • Briery Branch • Fraction forested for drainage basin of sample site

E. General Coliform

North River • Dry River • Briery Branch • Fraction forested for drainage basin of sample site

● North River ● Dry River ● Briery Branch ○ Fraction forested for drainage basin of sample site

Fig 6. Water quality parameters along river transects, from forest headwaters to agricultural bottomland. Blue,

orange and green points represent parameter mean ± standard deviation values for the North River, Dry River,

and Briary Branch, respectively. Open points represent percent forest fraction for each corresponding parameter

point.

Conductivity also showed a clear change after rivers moved from forest to agricultural lands (Fig 6F), with the

increase also happening about 8-10 river kilometers downstream from the forest edge. The increase was

characterized by larger variability in the valley compared to the forested region. This increase corresponds to

when sample site watersheds began to represent a more significant area of agricultural land (although the total

area in watersheds was still mostly forestland) (open circles, Fig 6F).

Since there was an abrupt shift in some parameters at the 8-10 km mark, we pooled samples from the transects

into three categories: forestland, edge of valley (up to 8 river kms), and interior of the valley (beyond 8 river kms).

Given the known impacts of flow rate on water quality parameters, we furthered binned samples into three flow

rate categories (low, medium and high). Two-way ANOVA did not show any effects of flow rate ($p > 0.05$ for all

parameters). We therefore dropped the flow rate bins, and used one-way ANOVA with post-hoc to investigate

the effect of location. For nitrate, orthophosphate, and general and fecal coliforms, forestland and edge of valley

generally did not statistically differ from each other, whereas a significant shift occurred once water was in the

interior of the valley (>8 km downstream) (Fig 7). This pattern did not hold for sediments.

Fig 7. Binned water quality parameters from transects. Data is delineated by sites in forestland, sites within 8km

river miles downstream from the forest/valley transition (valley edge), and sites more than 8km downstream of

the transition (valley interior). Values are mean ± standard deviation; significant differences between categories

indicated by letters on error bars (p<0.05).

We note that the 8 km mark, where percent of overall watershed generally drops to less than 80%, corresponds

with the previous analysis of all Shenandoah valley sites indicating the biggest change in parameters likewise

occurring with this change from heavily forested to moderately forested watersheds (Fig 5). The water quality

benefits of forested land continue for some distance into agricultural valley land, but this likely depended on

there being a large area of forest in the drainage basin (thus the overall watershed was still mostly forest). If there

isn't a large area forest (such as some of the watersheds in the valley represented by sampling points in the

overall Shenandoah watershed analysis), there are sites of higher water quality, but they are apparently

determined by other factors besides forest cover.

Finally, although sediment, nitrate and orthophosphate were lower in mountain regions, there were still levels of

these parameters (and coliforms) that would be considered impaired, indicating forested regions are still a source

of these pollutants at times. Orthophosphate and nitrate were in the medium to high stress level (0.05 ppm for

phosphate, and 1 ppm for TN) in 17.6% and 13.5% of samples, respectively, at forest sites. Comparable

percentages in the interior of valley samples were 45.2% of orthophosphate and 61.0% of nitrate.

Discussion

The current results confirm what is generally understood about water quality in naturally vegetated (in this case

forested) compared to agricultural areas – water quality is consistently better in watersheds which are well

covered with natural vegetation such as forests [12], and agricultural lands can have water quality impacts [23]. It

is evident, however, that within this general trend is important variability that occurs at localized areas. This

variability is important not only for local stream health, but also collectively drives downstream impacts. Further,

many waterways transect both forested and agricultural areas, and the interaction between these two is less well

characterized. By looking at a large public dataset, and focusing in on several representative watersheds, this

study identifies some distinct patterns of water quality in a large watershed with distinct forest and agricultural

segments.

1. Key Water Quality Parameters in Forested vs Agricultural Land

When analyzed across the entire Shenandoah watershed, there was a clear pattern of forested vs agricultural land

occurred for all parameters. Nutrients are key drivers of the downstream health of aquatic ecosystems, and in

this case are central to the major efforts to restore Chesapeake Bay health. Phosphorus and nitrogen are both a

focus of management efforts in upstream watersheds such as the Shenandoah. In fact, the Shenandoah/Potomac

watershed system is one of the most impacted in Virginia for these nutrients [24]. In our study, both transect

sampling and analysis of nutrients in the water quality portal database indicated a general pattern of higher

nutrient levels in agricultural regions that clearly agrees with the well-studied potential of agricultural practices to

impact water quality through anthropogenic activities. We did not explore specific drivers of nutrient dynamics in

the agricultural area, but note that the relationship between agricultural practices and nutrient levels is complex.

For instance, the large scale analysis of Metson et al. [25] found that there was no single explanatory variable that

determines riverine phosphorus levels, and other studies have found variable impact of best management

practices [26].

We note here four observations on nutrient trends from our data. First, our results are roughly consistent with

those found in other studies that focused on this region, both in average concentrations [24,27,28], and in the

presence of significant variation on the local level [15]. Second, many of these values were above estimated

threshold criteria for algal and benthic macroinvertebrate impairment [29], and are outside of the EPA suggested

criteria for ecological health [30]. Mean nutrient values for mountain and valley regions were in the low and

medium categories for probability of stress to aquatic organisms, respectively, according to Virginia stressor

analysis [24]. Third, values in areas of low impact (forested regions in the mountains) also reached levels of

concern at times, consistent with Smith et al [15] who found large variation in natural background of phosphorus

and nitrogen, with many stream and river segments in this ecoregion exceeding EPA-proposed criteria. Fourth,

phosphorus (orthophosphate and TP) were more variable in the valley than nitrogen (nitrate and TN) – nitrogen

was consistently high in the valley, whereas phosphorus was high at some sites, and low at others. Although our

data do not indicate what governs this difference (e.g. source, transport, adsorption, etc), it is consistent with the

observation in another study that TP has a smaller patch size than TN [5].

Two other measured parameters – sediment (as total suspended solids) and bacteria – are of major concern

nationwide as drivers of water quality, and have clear associations with agricultural activity. For instance,

sediment is the largest cause of invertebrate-impaired waters nationwide [31] and the primary stressor in the

Chesapeake Bay as a whole [8], and bacteria is the most common reason for streams receiving an impairment

designation in Virginia [24]. These parameters also showed some patterns of higher levels in agricultural lands,

although the pattern was more variable than for nitrogen and conductivity. For instance, sediments (TSS) were

highly variable in the transect data and did not correlate with land cover, but TSS across the watershed was higher

in drainage basins with less forest cover. Bacteria did correlate with land cover, but generally showed more

variability in both the transect and total watershed data. Overall, this points to some higher sediment and

bacteria values in forested land (see discussion below), and/or some waterways in primarily agricultural regions

that are less impacted by sediments and bacteria. Thus, although forest cover and agricultural activities are

critical for water quality, other factors are driving significant local variation in these parameters.

Conductivity was also clearly different between mountains and valley, and showed a more linear gradient with

change in forest cover. Conductivity was the single parameter that showed a correlation with multiple potential

drivers (Table 1) – at high significance for both forest cover and erodibility, and moderate significance (p<0.1) for

karst. It is likely, therefore, that some of this pattern is driven by the change from silicoclastic to karst geology as

rivers move from forest to valley. Although there are human activities of concern for conductivity (such as road

salting, or in areas with industry or mining [Fanelli]), studies in nearby Appalachian mountain areas point towards

agricultural having relatively low effect on conductivity [32], and conductivity is the parameter of least concern as

a stressor in the region [4,24]. In all cases, the mean conductivity was below 300 μS/cm, generally recognized as

the range of no or low probability for stress to aquatic life [24,30,31]. However, some individual samples fell in

the high stressor categories (>500 μS/cm), and could be a driver of which organisms make up aquatic

communities in these areas. These high conductivity values occurred during low-flow periods in valley areas only,

usually during mid-summer.

2. Water Quality Impacts in Forested Land

Although water quality was better in forested land, our data indicated that nitrogen, phosphorus, TSS and

indicator bacteria all exceeded levels of concern at particular times and locations. These results are consistent

with Bricker et al [14] who likewise found similar nitrate levels in a nearby mountain area, and with Price and

Leigh [33] who found TSS levels in forested areas of the southern Appalachians as high or higher than the levels

throughout our study area. Smith et al's [15] nationwide study also found some sites and times with levels of

concern in relatively unimpacted areas of our ecoregion.

We did not specifically investigate the source of these pollutant levels in forestlands, although our observations

pointed towards three general categories of possible human-related contributors: 1) existing stable structures, 2)

various types of disturbance, and 3) legacy effects. The contribution of existing stable structures is indicated by

consistently high levels of measured parameters (except conductivity) in retention structures such as a small pond

located in a clearing at the headwater of the North River. This pond was in a small sedimented bowl on the

ridgeline, and is characteristic of numerous small retention basins purposefully maintained in the forest (usually

for supporting game animals). Larger reservoirs are also present in all three watersheds for the purpose of flood

control and municipal water supplies. Recent evidence suggests that ponds can serve as nutrient sinks in

agricultural areas [34]; to our knowledge, there are no similar studies on the water quality role of ponds or lakes

in forested areas.

Disturbance is recognized as critical for mobilization of nutrients and sediments. For instance, forests in this area

are considered to show low nitrogen export unless disturbed [35]. While heavily forested, there are a variety of

processes which impact forests in a way that could mobilize nutrients and sediments. These regions have a

network of roads which are used for recreation, forest service activities (*e.g.* prescribed burns or firefighting), and

resource extraction (*e.g.* timbering). These roads often parallel waterways, are usually unpaved, and in some

areas have low-water crossings rather than bridges. Roads are a small fraction of the total area, but potentially a

significant contributor to water quality, and deserve future attention to understand their role forested regions.

Timber harvesting and prescribed burns are regular localized activities that can have significant impacts on

streams. In addition, our observation is that some streams in the forest areas have eroded banks, either part of

natural processes or as a legacy of alterations previously made, which would be a likely source of nutrients and

sediment. Bare soil from both management activities or streambank erosion are known drivers of water quality

and quantity in forested areas [10,36], and thus are likely contributors to the occasionally high levels of nutrients

and sediments in our study.

Candidate drivers of some water quality parameters include not only present day activities, but also legacy

activities. In addition to management activities occurring over the course of many decades, this area was heavily

logged in the late 1800's, and then used for livestock grazing before forests were restored in the early 1900's. The

role of legacy nutrients as drivers of watershed nutrients has received increased attention as a key driver of water

quality (*e.g.* [37]) as studies demonstrate their importance in a variety of watersheds [38-40]. Legacy nutrients

probably have variable residence times depending on local conditions, but are likely on the order of many

decades [40,41]. Mobilization from soil is particularly important during storm events following freeze-thaw cycles

[39], such as the spring rains often experienced in this central Appalachian region. Although these observations all

point toward a potential role of legacy nutrients in forested land, at present there is no direct data to guide an

understanding of how much legacy nutrients currently contribute to elevated nutrients in this area.

Finally, other "natural" processes can impact water quality in less disturbed watersheds. Atmospheric deposition

is shown to be important for both nitrogen [15] and phosphorus [6] in relatively undisturbed landscapes. Insect

outbreaks can dramatically increase nitrate from vegetation inputs, as shown in studies from Shenandoah

National Park, which borders the eastern side of this watershed [35]. Wildlife have been shown to be significant

sources of fecal coliforms in watershed analyses [42], and are likely a significant source of this parameter in our

forested areas.

In summary, although a national survey of landscapes found forests with the highest water quality among all land

cover types [12], a narrative that waterways in forested landscapes as invariably "clean" masks some interesting

localized events, and should be distinguished from the general observation that waterways are cleaner on

average. In fact, the recent study identifying forests as having the best water quality also emphasized

considerable variation in sediment and nutrient levels [12]. Forestlands in this region are a complex amalgam of

natural and human processes – as would be the case for most forests in the United States (and indeed globally).

Further studies should extend the general forest vs nonforest paradigm to focus on what categories of activities in

forests impact water quality more or less.

3. Patterns of Water Quality Transition as Rivers flow from Forested to

Agricultural Lands

Although this area is notable for a sharp forest/agriculture transition, transect samples indicated water which

flowed out of forested land did not show an immediate change in water quality parameters. Rather, there was a

clear lag period as the water quality parameters remained relatively unchanged until about 8 river km

downstream of the forest boundary (Figs 6,7). Although residence time (in this case, time in an agricultural area)

is a known driver of water quality [23], residence alone would more likely produce a gradual degradation of water

quality. Rather, the decrease in water quality was fairly abrupt, most notably for conductivity, which tracks with

multiple factors including underlying geology which shifts at the valley edge. When percent forest area of

watersheds for each sampling point were calculated, forested area for the corresponding drainage basins also

showed a sharp decrease at the 8 km mark to about 70-80%. WQP database results were similar, with many

parameters showing the most significant drop in quality when forested areas dropped below 80% (Fig 5).

The 8 km mark is the point at which tributaries draining mostly agricultural areas become a larger part of the total

drainage basin. In turn, this suggests that cumulative inputs from agriculturally derived tributaries are not

substantial until the river is 8km into the agricultural area, and that the pattern may be driven by inputs from the

broader tributary watershed rather than gradual inputs from agricultural activities immediately adjacent to the

river. This is consistent with the observation that sediment loads are correlated with bank erosion, rather than

primarily with runoff from adjacent land. Thus, it seems that the first 8-10km remains unimpacted because the

drainage area from non-forested regions is low. Once tributaries join, or substantial subsurface water from the

valley joins, the water quality begins to drop. Furthermore, while the benefits of the forested region stretched

well into the agricultural region, it took only 20-30% of a watershed to be nonforested in order for water quality

to be impacted. Price and Leigh [33] likewise found that even moderate changes in the degree of forestation can

have impacts on water quality variables. In aggregate, these data imply that 1) the value of forest in preserving

water quality is substantial, and extends downstream out of the forested area, but 2) that the ecological service of

maintaining water quality drops off once 20-30% of a total watershed was agricultural.

Although we did not focus on temperature, it is a parameter of critical concern in the area, as long-term

measurements in Shenandoah National Park show an increase that is expected in the presence of climate change

[43]. We did measure water temperature, but the interpretation of single temperature readings taken at different

times of the day and season limits conclusions from the transects. However, a rough estimate of transect trends

can be estimated by standardizing individual readings as the deviation from the average of a sampling run along a

transect. Temperature showed a more gradual increase across the length of the transect. Temperature steadily

rose as water moves from headwaters to the end of the sampling transect (Fig S1), contrasting with the abrupt

transitions seen in other parameters. The slow increase would be expected for temperature which is driven by

complex energy exchanges that rely on atmospheric factors (such as time with exposure to sun) rather than land

cover or other surface/subsurface factors [44].

Two other factors are key at governing the dynamics of water quality parameters, and are relevant to this

watershed – underground flow and storm events. Although we did not study the dynamics of underground flow,

which would require additional methodology such as flow tracing, others have pointed out the importance of

subsurface flow in this region [45,46]. The valley area is predominantly (but not entirely) karst, where

underground flow would be particularly important for impacting water quality. For instance, two of our transect

rivers are known to be interconnected - the Dry River flows underground into a tributary (Beaver Creek) of Briery

Branch [45]. This highlights how water quality at any point in fact likely represents a cumulative impact beyond

the drainage basin itself. Futhermore, karst may govern contaminants in other ways, such as through the

adsorption of TP [47]. Underground processes could be crucial to management efforts in systems such as Smith

Creek, an extensively studied watershed in our study area where the contribution of underground discharge is

greater than rainfall runoff [46]. For instance, mitigation interventions such as vegetated buffers along streams

may be less effective at intercepting contaminant inputs if water bypasses the buffer through underground flow.

High flow rates produced by storm events can drive the largest portion of loads [17,48,49], as demonstrated for

specific watersheds in the Chesapeake Bay system [46]. That study found that 90% of the sediment load in Smith

Creek in a three year period was due to eight large storms. In addition to increased streambank erosion in

headwaters, storm events initiate flow in the large number of as ephemeral streams, which mobilizes previously

stable pollutants [50]. We performed an initial investigation on the impact of storms by categorizing WQP data

generally as high and low flow, and we highlight several specific insights from our data that contribute to an

understanding the role of storms in water quality: 1) The effects of high flow varied with parameter. For instance,

phosphorus was mobilized more than nitrogen, which is consistent with observations in other studies [46], and

has implications for the impacts of high nutrient levels on algae growth [51]. Bacteria and sediment were also

mobilized, whereas conductivity decreased. Although nitrogen concentrations did not increase, the increased flow

rates would result in an increased nitrogen load, but to a lower degree than for phosphorus, bacteria and

sediment (where both concentration and flow rate increased). 2) The effects of high flow varied with forest

coverage. Notably, periods of high flow where forest coverage was high (>80%) did not cause the increased

phosphorus, bacteria and sediment concentrations noted above. Thus, heavily forest areas moderated the

increases in pollutant loads that occur during storm events, whereas drainage basins with more moderate or low

forest levels had higher loads through increased contaminant concentrations. Seasonal impacts would further

complicate these dynamics, as changes in management practices (such as manure or fertilizer application), and/or

"natural" sources and sinks such as leaf drop and algae growth, change the pollutant source and sinks. These

results highlight the continued need to understand the role of storm events on water quality.

Conclusions

The current study confirms the crucial role of substantial forested areas regions to the water quality of

watersheds. While the importance of headwater regions to downstream water quality is recognized [17,52], this

study combined multiple datasets to explore how water quality changes with the transition of waterways from

forested areas, which generally protect water quality, to agricultural areas, which generally degrade water quality.

Key insights which add detail to the general understanding of these landscapes are: 1) substantial variability in

water quality in both agricultural and forest areas points towards important local factors governing pollutants, 2)

while forests clearly had higher water quality than agricultural areas, a narrative of pristine water in forested

areas does not reflect the reality of parameters reaching levels of concern at certain locations and times, and 3)

forested regions provide benefits substantial distances downstream of transitions to agricultural regions, but

water quality rapidly decreased when cumulative forest cover of a corresponding watershed dropped by 20 to

30%. These results contribute to a general understanding of what drives water quality in mosaiced landscapes

such as the Chesapeake Bay watershed, which has shown complex and variable responses to management efforts

[7].

While the current study helps draw a more complete picture of water quality across a diverse landscape, we did

not investigate all possible drivers of water quality. Several other factors that are critical to understanding when

and how degradation of water quality occurs – notably the contribution of underground flow, storm events and

seasonal cycles. Flow rates, for instance, are a known and crucial determinant of loading [17,48,49], and nutrients

can follow seasonal trends [14]. Preisendanz et al [53] argue that "hot moments" are as important as "hot spots"

– and that insufficient attention has been directed towards these temporal events as opposed to the spatial

pattern. Whether the patterns seen here represent those found during peak storm flow, and across different

seasons, is a further avenue of fruitful study, particularly in light of future climate changes [54]. Future increases

in precipitation are highly relevant given that downstream water quality in the Chesapeake Bay is tied closely to

flow rates [17], and recent estimates are that climate change threatens to increase nutrient and sediment loads in

the Chesapeake Bay watershed even more than changes in agricultural intensity, population or development [55],

as nonpoint sources become more important [56].

References

1. Palmer MA, Hondula KL, Koch BJ. Ecological restoration of streams and rivers: shifting strategies and shifting

goals. Annual Review of Ecology, Evolution, and Systematics. 2014 Nov 23;45(1):247-69.

2. Richardson K, Steffen W, Lucht W, Bendtsen J, Cornell SE, Donges JF, Drüke M, Fetzer I, Bala G, Von Bloh W,

Feulner G. Earth beyond six of nine planetary boundaries. Science advances. 2023 Sep 13;9(37):eadh2458.

3. Caraco, N. F., & Cole, J. J. (1999). Caraco NF, Cole JJ. Human impact on nitrate export: an analysis using major

world rivers. Ambio. 1999 Mar 1;28(2):167-70.

4. Zheng L, Gerritsen J, Beckman J, Ludwig J, Wilkes S. Land use, geology, enrichment, and stream Biota in the

Eastern Ridge and Valley Ecoregion: implications for nutrient criteria development 1. JAWRA Journal of the

American Water Resources Association. 2008 Dec;44(6):1521-36.

5. Frei RJ, Lawson GM, Norris AJ, Cano G, Vargas MC, Kujanpää E, Hopkins A, Brown B, Sabo R, Brahney J, Abbott

BW. Limited progress in nutrient pollution in the US caused by spatially persistent nutrient sources. PLoS One.

2021 Nov 29;16(11):e0258952.

6. Stoddard JL, Van Sickle J, Herlihy AT, Brahney J, Paulsen S, Peck DV, Mitchell R, Pollard AI. (2016) Continental-

scale increase in lake and stream phosphorus: are oligotrophic systems disappearing in the United States?

Environmental science & technology. 5;50(7):3409-15.

7. Webber, J.S., Chanat, J.G., Clune, J.W., Devereux, O., Hall, N., Sabo, R.D., & Zhang, Q. (2024). Evaluating

water-quality trends in agricultural watersheds prioritized for management-practice implementation. JAWRA

Journal of the American Water Resources Association. https://doi.org/10.1111/1752-1688.13197.

8. Fanelli RM, Cashman MJ, Porter AJ. Identifying key stressors driving biological impairment in freshwater

streams in the Chesapeake Bay Watershed, USA. Environmental management. 2022 Dec;70(6):926-49.

9. Delia, K.A.; Haney, C.R.; Dyer, J.L.; Paul, V.G. Spatial Analysis of a Chesapeake Bay Sub-Watershed: How Land

Use and Precipitation Patterns Impact Water Quality in the James River. Water 2021, 13, 1592.

10. Neary DG, Ice GG, Jackson CR. Linkages between forest soils and water quality and quantity. Forest ecology

and management. 2009 Oct 30;258(10):2269-81.

11. Clinton BD, Vose JM. Variation in stream water quality in an urban headwater stream in the southern

Appalachians. Water, air, and soil pollution. 2006 Jan;169:331-53.

12. Caldwell PV, Martin KL, Vose JM, Baker JS, Warziniack TW, Costanza JK, Frey GE, Nehra A, Mihiar CM. Forested

watersheds provide the highest water quality among all land cover types, but the benefit of this ecosystem service

depends on landscape context. Science of the Total Environment. 2023 Jul 15;882:163550.

13. Richardson JS. Biological diversity in headwater streams. Water. 2019 Feb 21;11(2):366.

14. Bricker SB, Rice KC, Bricker OP. From headwaters to coast: influence of human activities on water quality of

the Potomac River Estuary. Aquatic Geochemistry. 2014 May;20:291-323.

15. Smith RA, Alexander RB, and Schwarz GE. Environmental Science & Technology 2003. 37 (14), 3039-3047 DOI:

10.1021/es020663b

16. Ator SW, Blomquist JD, Webber JS, Chanat JG. Factors driving nutrient trends in streams of the Chesapeake

Bay watershed. Journal of Environmental Quality. 2020 Jul;49(4):812-34.

17. Murphy RR, Keisman J, Harcum J, Karrh RR, Lane M, Perry ES, Zhang Q. Nutrient improvements in Chesapeake

Bay: Direct effect of load reductions and implications for coastal management. Environmental Science &

Technology. 2021 Dec 21;56(1):260-70.

18. Zhang Q, Blomquist JD, Fanelli RM, Keisman JL, Moyer DL, Langland MJ. Progress in reducing nutrient and

sediment loads to Chesapeake Bay: Three decades of monitoring data and implications for restoring complex

ecosystems. Wiley Interdisciplinary Reviews: Water. 2023 Sep;10(5):e1671.

19. Clune JW, Capel PD, Miller MP, Burns DA, Sekellick AJ, Claggett PR, Coupe RH, Fanelli RM, Garcia AM,

Raffensperger JP, Terziotti S. Nitrogen in the Chesapeake Bay watershed—A century of change, 1950–2050. US

Geological Survey; 2021. 168 p., https://doi.org/10.3133/cir1486.

20. Mwaijengo GN, Msigwa A, Njau KN, Brendonck L, Vanschoenwinkel B. Where does land use matter most?

Contrasting land use effects on river quality at different spatial scales. Science of the Total Environment. 2020

May 1;715:134825.

21. Segrè Cohen A, Slavik CE, Kurani S, Árvai J. Mapping risks of water injustice and perceptions of privatized

drinking water in the United States: A mixed methods approach. Risk Analysis. 2025.

https://doi.org/10.1111/risa.70012

22. Ijumulana J, Ligate F, Bhattacharya P, Mtalo F, Zhang C. Spatial analysis and GIS mapping of regional hotspots

and potential health risk of fluoride concentrations in groundwater of northern Tanzania. Science of the Total

Environment. 2020 Sep 15;735:139584.

. Farthing T, Rintsch E, Larson O, Grudzinski BP, Fisher TJ, McCarty JL. Effects of a forested state park on stream

nutrient concentrations in an agriculturally dominated watershed in the US Midwest. JAWRA Journal of the

American Water Resources Association. 2024 Aug;60(4):851-64.

24. [VDEQ] Virginia Department of Environmental Quality. Stressor analysis in Virginia: Data collection and

stressor thresholds. VDEQ Technical Bulletin WQA/2017-001. Richmond (VA): VA Department of Environmental

Quality Water Quality Monitoring, Biological Monitoring, and Water Quality Assessment Programs. 2017. 478 p.

https://www.deq.virginia.gov/home/showpublisheddocument/4313/637461491358800000

25. Metson GS, Lin J, Harrison JA, Compton JE. Linking terrestrial phosphorus inputs to riverine export across the

United States. Water Research. 2017 Nov 1;124:177-91.

26. Ator SW, Blomquist JD, Webber JS, Chanat JG. Factors driving nutrient trends in streams of the Chesapeake

Bay watershed. Journal of Environmental Quality. 2020 Jul;49(4):812-34.

27. Omernik J, Paulsen S, Griffith G, Weber M. Regional patterns of total nitrogen concentrations in the National

Rivers and Streams Assessment. Journal of Soil and Water Conservation. 2016 May 1;71(3):167-81.

28. Clune JW, Crawford JK, Boyer EW. Nitrogen and phosphorus concentration thresholds toward establishing

water quality criteria for Pennsylvania, USA. Water. 2020 Dec 17;12(12):3550.

29. Evans-White MA, Haggard BE, Scott JT. A review of stream nutrient criteria development in the United States.

Journal of environmental quality. 2013 Jul;42(4):1002-14.

30. USEPA. Ambient Water Quality Criteria Recommendations Information Supporting the Development of State

and Tribal Nutrient Criteria Rivers and Streams in Nutrient Ecoregion XI, 2000 USEPA-822-B-00–020. USEPA,

Washington, DC

31. Govenor H, Krometis LA, Willis L, Angermeier PL, Hession WC. Macroinvertebrate sensitivity thresholds for

sediment in Virginia streams. Integrated Environmental Assessment and Management. 2019 Jan 1;15(1):77-92.

32. Cormier SM, Wilkes SP, Zheng L. Relationship of land use and elevated ionic strength in Appalachian

watersheds. Environmental Toxicology and Chemistry. 2013 Feb 1;32(2):296-303.

33. Price K, Leigh DS. Comparative water quality of lightly-and moderately-impacted streams in the southern Blue

Ridge Mountains, USA. Environmental Monitoring and Assessment. 2006 Sep;120:269-300.

34. Liu W, Li S, Shen W, Zhuang Y, Li X, Ling F, Zhang L. Small water bodies influence river water quality in

agricultural watersheds. Agricultural Water Management. 2024 Nov 1;304:109070.

35. Eshleman KN, Fiscus DA, Castro NM, Webb JR, Deviney JJ. Computation and Visualization of Regional-Scale

Forest Disturbance and Associated Dissolved Nitrogen Export from Shenandoah National Park, Virginia. The

Scientific World Journal. 2001;1(2):539-47.

36. Gellis AC, Gorman Sanisaca L. Sediment fingerprinting to delineate sources of sediment in the agricultural and

forested Smith Creek watershed, Virginia, USA. JAWRA Journal of the American Water Resources Association.

2018 Dec;54(6):1197-221.

37. Golden HE, Evenson GR, Christensen JR, Lane CR. Advancing watershed legacy nitrogen modeling to improve

global water quality. Environmental science & technology. 2023 Feb 17;57(7):2691-7.

38. Cravotta III CA, Tasker TL, Smyntek PM, Blomquist JD, Clune JW, Zhang Q, Schmadel NM, Schmer NK. Legacy

sediment as a potential source of orthophosphate: Preliminary conceptual and geochemical models for the

Susquehanna River, Chesapeake Bay watershed, USA. Science of the Total Environment. 2024 Feb 20;912:169361.

39. Inamdar S, Johnson E, Rowland R, Warner D, Walter R, Merritts D. Freeze–thaw processes and intense rainfall:

the one-two punch for high sediment and nutrient loads from mid-Atlantic watersheds. Biogeochemistry. 2018

Dec;141:333-49.

40. Van Meter KJ, Van Cappellen P, Basu NB. Legacy nitrogen may prevent achievement of water quality goals in

the Gulf of Mexico. Science. 2018 Apr 27;360(6387):427-30.

41. McDowell R, Dodd R, Pletnyakov P, Noble A. The ability to reduce soil legacy phosphorus at a country scale.

Frontiers in Environmental Science. 2020 Jan 30;8:6.

42. Nguyen KH, Senay C, Young S, Nayak B, Lobos A, Conrad J, Harwood VJ. Determination of wild animal sources

of fecal indicator bacteria by microbial source tracking (MST) influences regulatory decisions. Water research.

2018 Nov 1;144:424-34.

43. Jastram JD, Snyder CD, Hitt NP, Rice KC. Synthesis and interpretation of surface-water quality and aquatic

biota data collected in Shenandoah National Park, Virginia, 1979-2009. US Geological Survey; 2013–5157, 77 p.,

http://pubs.usgs.gov/sir/2013/5157/.

44. Leach JA, Kelleher C, Kurylyk BL, Moore RD, Neilson BT. A primer on stream temperature processes. Wiley

Interdisciplinary Reviews: Water. 2023 Jul;10(4):e1643.

45. Doctor DH, Orndorff W, Maynard J, Heller MJ, Casile GJ. Ch.6 Karst geomorphology and hydrology of the

Shenandoah Valley near Harrisonburg, Virginia. In: Bailey CM, Coiner LV, editors. Elevating Geoscience in the

Southeastern United States: New Ideas about Old Terranes. Geological Society of America; 2014. pp.161-213.

46. Hyer KE, Denver JM, Langland MJ, Webber JS, Böhlke JK, Hively WD, Clune JW. Spatial and temporal variation

of stream chemistry associated with contrasting geology and land-use patterns in the Chesapeake Bay

watershed—Summary of results from Smith Creek, Virginia; Upper Chester River, Maryland; Conewago Creek,

Pennsylvania; and Difficult Run, Virginia, 2010–2013. US Geological Survey; 2016.

47. Li, L.;Wang,W.; Jiang, Z.; Luo, A. Phosphate in Aqueous Solution Adsorbs on Limestone Surfaces and Promotes

Dissolution. Water 2023, 15, 3230. https://doi.org/10.3390/w15183230

48. Lewis WM, Melack JM, McDowell WH, McClain M, Richey JE. Nitrogen yields from undisturbed watersheds in

the Americas. New Perspectives on Nitrogen Cycling in the Temperate and Tropical Americas: Report of the

International SCOPE Nitrogen Project. 1999:149-62.

49. Ezzati G, Kyllmar K, Barron J. Long-term water quality monitoring in agricultural catchments in Sweden: Impact

of climatic drivers on diffuse nutrient loads. Science of the Total Environment. 2023 Mar 15;864:160978.

50. Brinkerhoff CB, Gleason CJ, Kotchen MJ, Kysar DA, Raymond PA. Ephemeral stream water contributions to

United States drainage networks. Science. 2024 Jun 28;384(6703):1476-82.

51. Kelly PT, Renwick WH, Knoll L, Vanni MJ. Stream nitrogen and phosphorus loads are differentially affected by

storm events and the difference may be exacerbated by conservation tillage. Environmental Science &

Technology. 2019 Mar 12;53(10):5613-21.

52. Alexander RB, Boyer EW, Smith RA, Schwarz GE, Moore RB. The role of headwater streams in downstream

water quality 1. JAWRA Journal of the American Water Resources Association. 2007 Feb;43(1):41-59.

53. Preisendanz HE, Veith TL, Zhang Q, Shortle J. (2020) Preisendanz HE, Veith TL, Zhang Q, Shortle J. Temporal

inequality of nutrient and sediment transport: a decision-making framework for temporal targeting of load

reduction goals. Environmental Research Letters. 2020 Dec 18;16(1):014005.

54. Yang Y, Tilman D, Jin Z, Smith P, Barrett CB, Zhu YG, Burney J, D'Odorico P, Fantke P, Fargione J, Finlay JC.

Climate change exacerbates the environmental impacts of agriculture. Science. 2024 Sep 6;385(6713):eadn3747.

55. Bhatt G, Linker L, Shenk G, Bertani I, Tian R, Rigelman J, Hinson K, Claggett P. Water quality impacts of climate

change, land use, and population growth in the Chesapeake Bay watershed. JAWRA Journal of the American

Water Resources Association. 2023 Dec;59(6):1313-41.

56. Easton ZM, Hanson J, Bock E, Asfaw BW. A Review of Chesapeake Bay Climate Change: Potential Impacts on

Watershed Hydrology and Nutrient and Sediment Cycling and Export. JAWRA Journal of the American Water

Resources Association. 2025 Jun;61(3):e70030.

Supporting Information

817 **S1 Table. Dates and flow categories of transect sampling events.**

River	Date	Flow category
Dry River	2/9/2020	high
Briery Branch	3/8/2020	low
North River	10/24/2020	low
Briery Branch	11/8/2020	medium
Dry River	11/21/2020	medium
Briery Branch	12/31/2020	high
Dry River	1/23/2021	medium
Briery Branch	3/13/2021	high
North River	3/20/2022	medium
Dry River	4/3/2022	medium
North River	6/7/2022	medium
Dry River	7/20/2022	low
Briery Branch	7/21/2022	low
North River	8/3/2022	low
Dry River	8/19/2022	low
North River	9/4/2022	low
Briery Branch	6/30/2023	medium
North River	7/3/2023	medium
Dry River	7/5/2023	medium
North River	5/22/2024	high
Dry River	5/23/2024	high
Briery Branch	5/23/2024	high
North River	6/24/2024	low
Dry River	6/24/2024	low
Briery Branch	6/24/2024	low

S1 Figure. Temperature differential gradient parameters along river transects, from forest headwaters to

agricultural bottomland. Temperature differential for each sample calculated as site temperature minus average

temperature across its run (samples for the one river and one date). Blue, orange and green points represent

parameter mean ± standard deviation values for the North River, Dry River, and Briary Branch, respectively.

Asterisks with higher temperature differentials are pond or reservoirs.