Structurally distinct manganese-sensing riboswitch aptamers

regulate diverse expression platform architectures

Authors:

Christine N. Stephen[1], Danea E. Palmer[1], and Tatiana V. Mishanina[1*]

[1]Department of Chemistry and Biochemistry, University of California San Diego, 9500

Gilman Dr, La Jolla, CA 92093

*Corresponding author information:

Phone: (858) 246-5883

Email: tmishanina@ucsd.edu

Abstract

Manganese (Mn)-sensing riboswitches protect bacteria from Mn toxicity by upregulating expression of Mn exporters. The Mn aptamers share key features but diverge in other important elements, including within the metal-binding core. Although X-ray crystal structures of isolated aptamers exist, these structural snapshots lack crucial details about how the aptamer communicates the presence or absence of ligand to the expression platform. In this work, we investigated the Mn-sensing translational riboswitches in *E. coli* (*mntP* and *alx*), which differ in aptamer secondary structure, nucleotide sequence, and pH-dependence of Mn response. We performed co-transcriptional RNA chemical probing, allowing us to visualize RNA folding intermediates that form and resolve *en route* to the final folded riboswitch. For the first time, we report that sampling of metal ions by the RNA begins before the aptamer synthesis and folding are complete. At a single-nucleotide resolution, we pinpoint the transcription window where "riboswitching" occurs in response to Mn binding and uncover key differences in how the *alx* and *mntP* riboswitches fold. Finally, we describe riboswitch-specific effects of pH, providing insights into how two members of the same riboswitch family differentially sense two distinct environmental cues: concentration of Mn and pH.

Introduction

Transition metal ions play key cellular roles (e.g., serving as enzyme cofactors) but become harmful in excess due to misincorporation into biomolecules and metal-catalyzed formation of reactive oxygen species[1,2]. To match the need for metal ion transporters, cells use the metal's intracellular concentration to regulate expression of the metal's transporters at the transcription and/or translation level[3]. Examples of this regulation include metal-binding transcription regulators[4,5] or direct interaction of the metal with a riboswitch encoded in the mRNA[6–8]. Riboswitches are 5'-untranslated regions of mRNA that promote or inhibit expression of the downstream gene by changing their folding upon binding a ligand[9]. A riboswitch is comprised of a ligand-binding aptamer overlapping with an expression platform, whose folding determines the transcriptional and/or translational fate of the mRNA[9]. Many classes of metal-sensing riboswitches have been identified, such as magnesium[7], nickel[8], cobalt[8], and manganese (Mn)[6,10].

The *yybP-ykoY* Mn-sensing riboswitch family is the largest metal-sensing family, with >1,000 unique representatives in different bacterial phyla[11]. Crystal structures, single-molecule FRET (smFRET), and molecular dynamics (MD) simulations of isolated Mn-sensing aptamers provided valuable insights into the overall RNA fold, metal binding site, and RNA dynamics in the presence or absence of Mn[12,13]. However, these studies lack the expression platform of the riboswitch – the "business end" that determines the gene expression outcome. Consequently, it remains unknown how Mn binding to the aptamer actuates the differential folding of the downstream RNA to affect its transcription or

translation. Further, studies to date were performed on pre-synthesized RNA, missing the

aspect of active transcription that is critical to riboswitch folding[14–17].

We addressed these gaps by investigating the two Mn-sensing translational

riboswitches in *E. coli* which regulate translation of the *alx* and *mntP* genes, with the *alx*

riboswitch upregulating gene expression in response to alkaline pH and increased Mn

concentration ([Mn])[18,19] (Fig. 1A). Binding of Mn to either the *mntP* or *alx* aptamers

promotes a translationally active conformation in the expression platform, increasing

expression of the Alx and MntP Mn exporters[10,19,20] (Fig. 1A-B). Multiple structures of Mn-

sensing riboswitch aptamers established their core structural features, consisting of two

coaxially stacked "legs" connected by a four-way junction (4WJ)[6,12,21] (Fig. 1C). The right

leg (L1, P1.1, P1.2, and P2; Fig. 1C) extrudes a conserved adenosine in L1 to form a

cross-helix A-minor interaction with P3.1 in the left leg (L3, P3.1, P3.2, and P4; Fig. 1C).

The L3 loop in the left leg stacks below the conserved adenosine in the A-minor

interaction, forming a phosphate-rich metal-binding hot spot (Fig. 1C). X-ray crystal

structures show two metal-binding sites in the aptamer, with one occupied by Mg^{2+} (M_A)

and the other by Mn^{2+} (M_B)[6,12,21] (Fig. 1C). The P1.1 "switch helix" is a key element of the

yybP-ykoY aptamer as its 3'-half alternatively pairs within the aptamer or the expression

platform depending on whether Mn is bound, with consequences to folding of the rest of

the expression platform.[12] (see Fig. 1B for *alx* and *mntP* alternative P1.1 pairing).

Despite sharing conserved features for Mn-sensing, the *alx* and *mntP* riboswitches

differ in their global architectures (Fig. 2A). The *alx*-like riboswitches are exclusively

present in the *Enterobacteriales* order, members of which are often found in environments

that can alkalinize, such as water, soil, and in association with other organisms[22,23]. Upon

alkaline pH challenge, the *E. coli* cytosol alkalinizes and then recovers to pH 7.8[24]; in turn,

cytosol alkalinization increases cytoplasmic [Mn][19], potentially explaining *alx* translational

activation at alkaline pH. Initial studies of the *alx* and *mntP* riboswitches put forward

alternative secondary structure models based on chemical probing with pre-synthesized

RNA[10,18], prompting us to ask (i) how these alternative structures fold and respond to Mn

during transcription and (ii) why alkaline pH uniquely affects *alx* structure. To answer

these fundamental questions, we applied modern RNA chemical probing methods to *alx*

and *mntP*, where ribonucleotides are modified by either a selective 2'-hydroxyl (SHAPE)

reagent or dimethyl sulfate (DMS) and detected by high-throughput mutational profiling

(MaP)[25].

In this work, we performed DMS-MaP in a co-transcriptional format using established

methods[26,27], getting to the heart of how Mn binding to the aptamer is "communicated" to

the expression platform. Unexpectedly, we uncovered evidence that both *alx* and *mntP*

aptamer folding intermediates begin sampling for divalents before the full aptamer domain

has emerged from RNAP, challenging the notion that a fully folded aptamer is required

for ligand sensing. Further, a critical RNA loop that participates in Mn sensing diverges in

sequence between *alx* and *mntP* and may hold the key to how pH differentially affects

riboregulation by these riboswitches. Using gene reporter assays, we show that alkaline

pH alone cannot rearrange the *alx* riboswitch into a translationally active conformation;

instead, the *alx* riboswitch senses the increased [Mn] brought about by alkaline pH. Lastly,

our co-transcriptional chemical probing data uncovered distinct structural features in the

alx and *mntP* expression platforms, which are potentially coupled to the cellular need for

the Alx and MntP exporter proteins under increased cytosolic [Mn].

Results

The Mn-sensing aptamers of the *alx* and *mntP* riboswitches differ structurally

In Fig. 2A, we present secondary structure models of the *alx* and *mntP* riboswitch aptamer

domains based on prior experimental data for the full-length riboswitches[10,18], which we

updated to reflect the overall aptamer architecture and conserved base-pairs identified by

X-ray crystal structures[6,12,21]. The model of the *mntP* aptamer in Fig. 2A is supported by

an X-ray crystal structure of the aptamer in the apo state (PDB ID: 4Y1M)[6]. Since no

detailed structural information exists for the *alx* aptamer, we mapped its secondary

structure with DMS/SHAPE-MaP using the traditional approach of *in vitro* synthesis by T7

RNA polymerase (RNAP) followed by denaturation and refolding.

pH and Mn effect on the alx aptamer structure. The aptamer RNA was refolded at either

neutral (7.2) or alkaline pH (8.5) data and *in vitro* transcription experiments[18,28] and the

alkaline pH value aligns with published gene reporter and *in vitro* transcription

experiments[19,28]. The aptamer RNAs were probed with either 1-methyl-7-nitroisatoic

anhydride (1M7) or DMS. The resulting secondary structure models of the *alx* aptamer

agree with the published 2D predictions[18] and 3D structure of an *L. lactis* chimera with

the *alx* L3 sequence[21] (Fig. 2B and SI Fig. 1A-C). Comparison of the nucleotide (nt)

reactivities as a function of refolding pH revealed clear flexibility differences in L3 and L1,

with multiple nt (e.g., A37 and A11) showing decreased reactivity towards 1M7 or DMS

at alkaline pH (Fig. 2C).

We next refolded the *alx* aptamer RNA at either neutral (7.2) or alkaline pH (8.5) in the

presence of 1 mM $MnCl_2$ and probed with either DMS or 1M7. While the total physiological

[Mn] (free and bound) in *E. coli* is estimated to be 15-21 μM[29–31], we chose a saturating

concentration of ligand to maximize the ligand-induced shift in the RNA conformational

ensemble, consistent with prior chemical probing experiments on other riboswitches[27,32].

$MgCl_2$ was present at 5 mM in the RNA refolding buffer, thus Mg remained in excess of

Mn, as is the case *in vivo*[33]. We observed L3 generally becomes less reactive towards

1M7 in the presence of Mn, consistent with reduced flexibility upon Mn binding (SI Fig.

1D). Interestingly, in the presence of Mn, pH-dependent reactivity changes for multiple

L1/L3 nt are abolished (e.g., A37 and A11, Fig. 2C), supporting that these nt are buried

in the metal-sensing core with Mn and consistent with published Mn-bound aptamer

structures[6,12,21] (see Supplementary Note 1 and SI Fig. 1E for additional analysis).

Structural comparison of the alx and mntP aptamers. Despite belonging to the same

family, the *alx* and *mntP* riboswitch aptamers harbor intriguing structural differences. First,

the *alx* aptamer contains an intrinsic termination site in the P4 helix[28], which occurs before

the metal-sensing pocket forms (Fig. 2A). The *alx* aptamer lacks the P2 helix, resulting in

a three-way junction (3WJ) rather than a 4WJ in the *mntP* aptamer and all other aptamers

studied by X-ray crystallography. Lastly, the L3 sequences in *alx* and *mntP* metal-sensing

cores differ (Fig. 1C), raising the possibility that L3 tunes Mn affinity of the aptamer.

Transcription by native RNA polymerase is required for *alx* and *mntP* riboswitch-mediated translational regulation *in vivo*

We next wondered whether *alx* and *mntP* riboregulation is kinetically controlled. To test this, we constructed *alx* and *mntP* translational *lacZ* reporters cloned with either the T7 promoter or native *alx/mntP* promoter into single copy plasmids (Fig. 2D). The effect of extracellular alkaline pH or elevated [Mn] on gene expression was measured by β-galactosidase assays in either an *E. coli* strain (MC4100) lacking *alx* (Δ*alx*), to preclude negative autoregulation[19], or the Δ*alx* strain with T7 RNAP expressed from the *lac* promoter (chromosomally encoded, Δ*alx*T7). The plasmids containing the native promoter translational reporters were transformed into the Δ*alx* strain and the plasmids containing the T7 promoter translational reporters were transformed into the Δ*alx*T7 strain. The strains were cultivated in i) neutral pH (LBK pH 6.8) or alkaline pH (LBK pH 8.4) media, and ii) neutral pH (LBK pH 6.8) media with supplemented $MnCl_2$.

We observed the expected changes in *alx* and *mntP* translation in response to pH and supplemented $MnCl_2$ when reporter fusions were transcribed from their native promoters[19] (Fig. 2E-F). While only *alx* translation was induced by alkaline pH, translation of both genes increased with supplemented $MnCl_2$ (Fig. 2E-F). Strikingly, transcription by T7 RNAP abolished the response to alkaline pH and/or Mn (Fig. 2E-F), demonstrating that faster transcription of the *alx* and *mntP* riboswitches (~220 nt/s; T7 RNAP[34] vs. ~20-90 nt/s; *E. coli* RNAP[35]) prevents riboswitching in response to alkalinization and/or elevated [Mn] in the cytosol. In sum, these data establish that transcription by *E. coli*

RNAP is required for ligand-coupled structural rearrangements in the *alx* and *mntP*

expression platforms.

Analysis of the *mntP* aptamer folding during transcription using co-transcriptional

chemical probing

To investigate both pH and Mn-mediated structural changes in the *alx* and *mntP*

riboswitch folding intermediates, we performed co-transcriptional chemical probing

(variable length Transcription Elongation Complex RNA structure probing TECprobe-VL

method, see SI Fig. 2 for construct designs)[27]. Co-transcriptional chemical probing

experiments were performed at both neutral (7.2) and alkaline pH (8.5) ± 1 mM $MnCl_2$

(referred to as "+Mn" or "–Mn" condition, respectively) with 10 mM $MgCl_2$ present to

prevent compromised transcription fidelity from Mn-coordination by RNAP, as observed

for DNA polymerases[36] and reverse transcriptases[37]. Since published work established

that pH≥ 8 renders all 4 nucleobases accessible to DMS modification[38], we first present

data for *alx* and *mntP* DMS-probed folding intermediates at pH 8.5.

The two legs of the Mn aptamer fold differently. We initially mapped *mntP* aptamer folding

transitions and determined the specific transcript lengths where helices in the aptamer

become stably folded (Fig. 3A-B and SI Fig. 3). P3.1, P3.2, and P4 in the left leg of the

aptamer fold first, as detected with coordinated drops in reactivity of nucleotides that are

base-paired in these helices (Fig. 3C). In stark contrast, the right leg of the aptamer folds

into transient, low-melting temperature (T_m) stem-loops that form and resolve until the

synthesis of the aptamer is completed (~nt 124, assuming a 14-nt RNAP footprint, Fig.

3B). While the 5' RNA stretch for these aptamer elements (L1, P1.1, and P1.2) is

transcribed early on, their 3' pairing partner does not emerge from RNAP until late in

transcription. Thus, the weak stem-loop intermediates may prevent RNA from being

trapped in stable but not biologically relevant folds, consistent with other riboswitches[16,27].

P2 in the right leg folds towards the end of aptamer transcription, even though the

sequence encoding P2 emerges at transcript length ~50 nt, indicating that folding of the

entire right leg occurs at the end of aptamer synthesis (Fig. 3C).

RNA folding intermediates sample for Mn. Surprisingly, we observed Mn-dependent

differences in reactivity of several nucleotides *before* complete *mntP* aptamer has been

synthesized (Fig. 3D). For example, A48, the first nucleotide in the L3 loop, demonstrates

markedly high reactivity ±Mn, with its reactivity trending higher +Mn starting at transcript

length ~115 nt – before the aptamer synthesis finished ~124 nt (Fig. 3D). We hypothesize

that this is indicative of an RNA folding intermediate sampling divalent metal ions in

solution in search for Mn, with its L3 and partially formed L1/P1.1. Such partial binding of

Mn was observed in a crystal structure and molecular dynamics (MD) simulations of the

full-length *X. oryzae* aptamer, referred to as "conformer 2", which lacks the ordered

stacking pattern in L3 and contains a partially dehydrated Mn in the M_B site[12] (Fig. 3D).

The partially dehydrated Mn lacks half of the contacts seen in "conformer 1" (coordinated

to fully dehydrated Mn); however, the inner sphere contact between the phosphate of the

first nucleotide in L3 and the metal is observed (A48 in *E. coli mntP* aptamer vs. A46 in

X. oryzae aptamer)[12]. Therefore, A48 can contact a partially dehydrated Mn before the

full aptamer is synthesized, explaining its Mn-dependent reactivity differences in *mntP*

aptamer folding intermediates. U54 in L3 and A103 in L1 also show Mn-dependent

reactivity differences in aptamer intermediates, with their reactivities trending higher –Mn

(Fig. 3D). These reactivity differences can be explained by greater conformational

flexibility in the metal-binding site (see Fig. 3D for U54 and A103 positioning in *X. oryzae*

conformers 1 and 2). Additionally, we observed multiple RNAP pauses preceding *mntP*

aptamer completion that are enhanced by the presence of Mn and may mediate partial

Mn binding/ sampling of divalents in the environment by the RNA (see Supplementary

Note 2 and SI Figs. 4-5 for further analysis).

To identify the intermediates whose folding is truly controlled by transcription kinetics

vs. thermodynamic stability of the folds, we applied a modified TECprobe-VL workflow

wherein intermediate RNA transcripts are extracted from the stalled transcription

elongation complexes and equilibrium refolded before DMS probing. These experiments

identified clear structural differences between co-transcriptionally folded and equilibrium

refolded *mntP* intermediates and provided additional evidence of divalent sampling by

aptamer folding intermediates (see Supplementary Note 3 and SI Fig. 6 for further

analysis).

Analysis of the *alx* aptamer folding during transcription using co-transcriptional

chemical probing

We next investigated the *alx* aptamer folding transitions at pH 8.5 (Fig. 4A-B and SI Fig.

7). Here, P3.1, P3.2, and P4 in the left leg fold first as the aptamer is extended from ~80

nt to ~125 nt (Fig. 4B-4C). Like in the *mntP* aptamer, the 5' end of the right leg folds into

transient, low-T_m stem-loops that resolve once the full-length aptamer RNA has emerged

from RNAP (Fig. 4B), highlighting a potentially conserved mechanism of the *yybP-ykoY*

aptamer folding.

L3 flexibility differs between alx and mntP aptamers. Like with *mntP*, we captured Mn-

dependent reactivity differences for L3 in *alx* aptamer folding intermediates (Fig. 4D). For

example, the reactivity of U35, the first nt in the *alx* L3, trends higher +Mn starting at

transcript lengths as short as ~50 nt (Fig. 4D). This trend mimics that seen with the first

nt in *mntP* L3, A48, with a few key differences. First, the absolute reactivity values ±Mn

are greater for A48 in *mntP* L3 (Fig. 3D). Second, the increased reactivity +Mn occurs

much earlier for *alx* aptamer intermediate transcript lengths (~50 nt for *alx* vs. ~115 nt for

mntP). The absolutely conserved adenosine in *alx* L3 that directly coordinates Mn (A37)

is more reactive –Mn for aptamer folding intermediates starting at transcript length ~70 nt

(Fig. 4D). The higher A37 reactivity –Mn is consistent with greater L3 flexibility in this

condition and further supports that the L3 loop can begin sampling for divalents before

the aptamer domain is fully folded. Curiously, the absolutely conserved adenosine in

mntP L3 (A50) does not demonstrate this Mn-dependent reactivity difference (Fig. 3C),

pointing to a difference in L3 dynamics between *alx* and *mntP* aptamer folding

intermediates. Similar to *mntP,* we observed clear structural differences between co-

transcriptionally folded and equilibrium refolded *alx* intermediates (see Supplementary

Note 4 and SI Fig. 8 for further analysis).

Mn binding is a prerequisite for *alx* riboswitch pH response

Early work characterized the *alx* riboswitch as a pH sensor[18]; however, recent studies

have called this into question[19]. Since the intracellular [Mn] increases in *E. coli* at alkaline

pH[19], we wondered whether the *alx* riboswitch is actually sensing the increase in [Mn] as

a consequence of cytosol alkalinization. To test this, we performed translational *lacZ*

reporter assays on the A37U *alx* mutant, where the absolutely conserved adenosine that

directly coordinates Mn in L3 is replaced with a uridine. This point mutation was previously

shown to ablate Mn-sensing in the *X. oryzae* aptamer[12] making it the best candidate

mutation to decouple the effect of Mn and alkaline pH on *alx* riboregulation *in vivo.*

Strikingly, the 68-fold increase in *alx* reporter translation observed upon a shift from pH

6.8 to 8.4 for the wild-type riboswitch was abolished by the A37U mutation (Fig. 5A).

These data provide the first evidence that the *alx* riboswitch is not a true pH sensor;

instead, the riboswitch senses increased cytosolic [Mn], which is an indirect effect of

cytosol alkalinization. Published *alx* translational reporter data showed an additive effect

of alkaline pH and Mn[19]. We thus hypothesize that alkaline pH enhances the affinity of

the *alx* aptamer for Mn, rather than alters the *alx* riboswitch folding, unifying observations

from early studies on this system[18,39] with a recently published genetic study[19]. A

combination of alkaline pH and high [Mn] produced higher *mntP* reporter translation

compared to the neutral pH condition with added Mn, suggesting that alkaline pH may

also enhance Mn affinity for the *mntP* aptamer, but to a lesser degree than for the *alx*

aptamer[19]. To explore this intriguing point further, we DMS-probed the *alx* and *mntP*

folding intermediates at pH 7.2 ±Mn with the TECprobe-VL method, as described in the

following section.

pH effects on the *alx* and *mntP* aptamer folding: L3

We largely observed the same co-transcriptional folding transitions for the two legs in the

alx and *mntP* aptamers at pH 7.2 as described for pH 8.5 (SI Fig. 9A-B). Although the *alx*

and *mntP* aptamers exhibit the same global folding transitions at pH 7.2, key pH-

dependent reactivity differences in their metal-sensing cores emerged. Here, we focused

on A and C nucleotide reactivities since G and U nucleotides are protonated at the

respective N1 and N3 positions at neutral pH and are thus unreactive towards DMS (pKa

~ 9.2[40]). First, the absolutely conserved adenosine in the *alx* L3 (A37) exhibits a general

Mn-dependent reactivity decrease during aptamer folding at both pH (Fig. 5B). However,

the magnitude of this reactivity decrease is clearly pH-dependent, with A37 consistently

less reactive at pH 8.5 +Mn compared to other conditions tested (Fig. 5B), suggesting a

potentially more stable A37-Mn coordination at alkaline pH.   In stark contrast, the

absolutely conserved adenosine in *mntP* L3 (A50) shows no drop in reactivity at pH 8.5

+Mn; instead, A50 lacks clear pH-dependent reactivity trends during aptamer folding (Fig.

5C). The first L3 nucleotide in the *mntP* aptamer (A48) is more reactive +Mn vs. –Mn

regardless of pH (Fig. 5C), supporting that A48 stacked with A11 in an Mn-bound, ordered

metal-sensing core increases its DMS reactivity. Interestingly, the absolute reactivity

values for A48 are markedly lower at pH 7.2 ±Mn (Fig. 5C), suggesting that A48 has an

upwardly shifted pK$_a$ close to physiological pH (~pH 7.4-7.8 for *E. coli*[23]). We thus

hypothesize that deprotonation of A48 at alkaline pH promotes its stacking with A11 in

the cross-helix A-minor interaction, favoring docking of the helical legs in the aptamer

(see Supplementary Note 5 and SI Figs. 9C-D,10, and 11 for further analysis of pH-

dependent structural changes in *alx/mntP* L1 and L3 in co-transcriptionally folded vs.

equilibrium refolded aptamer intermediates).

We wondered whether L3 sequences in *Enterobacteriales yybP-ykoY* family members

show a unique trend that could explain the pH-dependent reactivity differences in the *alx*

and *mntP* aptamers. We aligned sequences of *yybP-ykoY* riboswitch family members

within the *Enterobacteriales* order, revealing two distinctive groups: an *alx*-like group

(consensus: UCAUUAC, Fig. 5D) and an *mntP*-like group (consensus: ACAUACU, Fig.

5D). We then compared the above sequences to the *yybP-ykoY* riboswitch family L3

consensus sequence generated from 1,063 aligned sequences (consensus: ACAUACU,

Fig. 5D). Generally, the first nucleotide in L3 is an A – an important position since this

nucleotide stacks with the conserved A11 in L1 that forms the cross-helix A-minor

interaction. Therefore, this position being strictly a U in *alx*-like members suggests a

functional role. Since the N3 of U has a close-to-physiological pKa of ~9.2[40], it is possible

that a shift from neutral to alkaline pH alters its stacking with the conserved A11 and

consequently Mn-coupled structural rearrangements in the expression platform.

Analysis of *mntP* expression platform folding during transcription

Cross-helix A-minor disruption at the riboswitching point. The expression platform for the *mntP* riboswitch contains a single helix that begins folding at transcript length ~175 nt and was detected by coordinated drops in reactivity for its constituent nucleotides (Fig. 6A and Fig. S12A). In the absence of Mn, the aptamer remains stably folded for most of expression platform synthesis until the transcript is ~205 nt long (Fig. 6B). At this critical decision point – the riboswitching point – the L1 and P1.1 elements of the aptamer are disrupted if no Mn is available, allowing for alternative pairing of P1.1 switch helix in the expression platform (Fig. 6C). The disruption of P1.1 in the absence of Mn is consistent with previously published single-molecule fluorescence experiments with an isolated aptamer that show that this helix is less stable without Mn[12]. Analysis of single-nucleotide trajectories revealed that A11 reactivity transiently spikes at ~205 nt, whether or not Mn is included in the transcription reaction, but the amplitude of the spike is Mn-dependent (Fig. 6D). The spike in A11 reactivity coincides with P1.1 strand invasion and its reduced amplitude +Mn indicates A11 remains trapped in the long-range A-minor interaction in Mn-bound aptamer. The L1/P1.1 reactivity difference heatmap ±Mn highlights A11 as the nucleotide most sensitive to the presence of the ligand, serving as a marker for the riboswitching point (Fig. 6D). We also assessed reactivity changes for A48 in L3 during expression platform folding, since A48 stacks with A11 and helps anchor the cross-helix A-minor interaction. A48 reactivity drops ±Mn starting at transcript length ~185 nt, with a larger decrease –Mn (Fig. 6E). A48 reactivity is consistently lower –Mn for the remainder of expression platform folding (Fig. 6E), likely due to broken A48-A11 stacking.

We observed similar reactivity trends in *mntP* expression platform folding at pH 7.2 as

pH 8.5 (e.g., A11 reactivity spike, SI Fig. 12B-C) and uncovered additional evidence of

P1.1/L1 disruption in the vicinity of the riboswitching point (see Supplementary Note 6

and SI Fig. 12D). Taken together, our observations highlight that a global structural

change occurs in the *mntP* expression platform at transcript lengths ~200-205 nt, which

is likely prompted by the start of strand invasion at P1.1 in the aptamer (see

Supplementary Note 6 and SI Fig. 12D for additional analysis of pH- and Mn-dependent

reactivity changes in *mntP* L1, L3, and P1.1 during expression platform folding).

Co-transcriptional binding of Mn is required for riboswitching. We next evaluated the

differences between co-transcriptionally folded and equilibrium refolded *mntP*

intermediate transcripts that contain both the aptamer domain and expression platform,

to assess whether co-transcriptional folding is required to observe Mn-coupled structural

rearrangements in the expression platform. While we observed similar P5 folding

transitions (SI Fig. 13A-B), the Mn-dependent spike in A11 reactivity is abolished when

the RNA is equilibrium refolded ±Mn at either pH 7.2 or 8.5 (SI Fig. 13C). These data

support that co-transcriptional Mn-binding is critical for stabilization of the A-minor motif

that docks the two legs of the aptamer together and for prevention of the P1.1/L1 invader

strand from disrupting the aptamer (see Supplementary Note 7 and SI Fig. 13D for

reactivity analysis of the A11 stacking partner, A48, in equilibrium refolded intermediates).

Further supporting the above observations, we found that T7-synthesized, equilibrium

refolded *mntP* riboswitch adopts a translationally inactive conformation ±Mn (DMS/1M7-

probed, described further in Supplementary Note 8 and SI Fig. 14).

Analysis of *alx* expression platform folding during transcription

Distinct features of the alx expression platform folding. The expression platform for the

alx riboswitch contains multiple helices that begin folding at transcript length ~155 nt, as

detected by coordinated drops in reactivity for the constituent nucleotides in each helix

(Fig. 7A and SI Fig. 15A). The *alx* expression platform contains two helices (termed here

P5 and P6) in its translationally inactive conformation that are predicted to fold –Mn,

resulting in the ribosome binding sequence (RBS) base-paring with a disrupted P1.1/L1

(Fig. 7B-D). On the other hand, when P1.1/L1 remain base-paired in the Mn-bound *alx*

aptamer, an alternative helix in the expression platform (termed here P5/P6) is predicted

to fold, and the downstream RBS-containing strand is single-stranded (Fig. 7E-G).

Unlike the *mntP* riboswitch, the *alx* P1.1/L1 rising stem is highly dynamic during the

entirety of *alx* expression platform folding both ±Mn, which was detected as consistent

reactivity towards DMS for multiple nt (Fig. 7A). The strand predicted to invade P1.1/L1

in the *alx* aptamer emerges at transcript length 155 nt, resulting in formation of the P5

helix (Fig. 7A). Once the expression platform is extended to transcript length 190 nt, the

P6 intrinsic terminator helix folds (Fig. 7A). Interestingly, the DMS reactivities for

nucleotides in the *alx* expression platform were largely similar ±Mn, resulting in similar

secondary structure predictions for expression platform folding intermediates (SI Fig.

15B-D).

Despite the global structural similarities in the *alx* expression platform secondary

structural models ±Mn, we detected multiple Mn-dependent reactivity differences for

specific nucleotides that indicate a modest ligand-dependent shift in the conformational

ensemble. P5 stem-loop nucleotides are more reactive –Mn (Fig. 7C), consistent with a

population bias towards the inactive conformer. A137 (bulged in active conformer vs.

base-paired in inactive conformer) demonstrates markedly high reactivity in general and

exhibits higher reactivity +Mn (Fig. 7F), indicating a shift in the conformational ensemble

towards the *alx* active conformer.

pH effects during folding of the alx expression platform. We observed similar DMS

reactivity patterns in the *alx* expression platform at neutral vs. alkaline pH, such as folding

of P5 and P6 (SI Fig. 16A-B), indicating that a shift to alkaline pH impacts metal-sensing

rather than directly affecting folding of the expression platform and thus riboswitching.

However, we observed interesting pH-dependent reactivity trends for A137, which was

more reactive +Mn at both pH and most reactive overall at pH 8.5 +Mn (SI Fig. 16B),

supporting that the active conformer is sampled more frequently under this condition.

We next assessed pH- and Mn-dependent reactivity trends in the *alx* aptamer during

expression platform folding (Fig. 8A-C). Contrasting *mntP*, A11 remains accessible for

DMS methylation across all conditions tested with no significant pH and Mn-dependent

differences (Fig 8B and SI Fig. 16C). Since A11 is conserved in *alx* and *mntP* L1, it is

curious that A11 exhibits distinctive reactivity trends between the two riboswitches. Unlike

in *mntP*, the absolutely conserved A37 in *alx* aptamer remains least reactive at pH 8.5

+Mn for the duration of expression platform folding (supportive of more stable Mn binding

at alkaline pH), with multiple transient spikes in reactivity indicating a change in L3

dynamics at pH 8.5 +Mn during expression platform folding (Fig. 8C and SI Fig. 16C).

Additionally, we observed interesting pH- and Mn-dependent reactivity trends for *alx*

expression platform nt in equilibrium refolded intermediates (see Supplementary Note 9

and SI Fig. 17).

The *mntP* and *alx* RBS regions are dynamic in co-transcriptionally displayed RNA

The RNA surrounding the *mntP* RBS did not demonstrate a clear Mn-dependent reactivity

trend, with DMS-constrained structure models predicting low-T_m hairpins interspersed

with single-stranded regions (Fig. 18A). We thus conclude that the RNA around the RBS

is dynamic, and any Mn-dependent structural differences are likely averaged out. Prior

smFRET study observed such RBS dynamics for the preQ$_1$ translational riboswitch with

ligand-dependent accessibility bursts in the RBS region[41]. Therefore, characterization of

the *mntP* RBS dynamics requires complementary, e.g., single-molecule methods.

Likewise, heatmap visualization of the *mntP* RBS nucleotides at pH 7.2 or 8.5 ±Mn or in

equilibrium refolded intermediates at both pH showed no clear reactivity difference that

would indicate increased RBS accessibility +Mn (SI Fig. 18B-C).

For *alx*, the presence of the P6 intrinsic terminator stem resulted in a stark decrease

in representation for longer transcripts containing the RBS (SI Fig. 19A), precluding

accurate modeling of the RNA structure in this region. To boost representation of full-

length *alx*, we performed a modified co-transcriptional chemical probing experiment with

a terminally biotinylated DNA template and the fast-acting SHAPE reagent benzoyl

cyanide (BzCN, reaction time ~1s) to better capture the short-lived co-transcriptional, Mn-

dependent structural changes in the RBS. Consistent with *mntP*, the single-length BzCN

probing experiment revealed no clear Mn-dependent differences in the reactivities of RBS

nucleotides (Fig. 19B), supporting that the *alx* RBS is dynamic. These dynamics may

serve a functional role; for instance, the RBS may be rapidly docking/undocking with

another tertiary structural element in *alx*, with the docking rate fine-tuning RBS

accessibility. Underscoring the importance of co-transcriptional chemical probing, we

found that T7-synthesized, equilibrium refolded *alx* riboswitch adopts a translationally

inactive conformation ±Mn (DMS/1M7-probed, described further in Supplementary Note

10 and SI Fig. 20). Taken together, while *alx* and *mntP* differ structurally, they may share

common dynamic features that gate RBS accessibility.

Discussion

In this work, we performed a suite of chemical probing experiments on the Mn-sensing

translational riboswitches in *E. coli* that precede the *mntP* and *alx* genes encoding Mn

exporters. While static structural snapshots are available for Mn-sensing aptamers, these

structures do not answer the question of how binding of Mn to the aptamer is

"communicated" to the expression platform – the question at the heart of riboswitch

function. Because the *mntP* and *alx* aptamers display unique architectures, they are an

interesting test case of how structurally distinct aptamers in the same riboswitch family

"talk" to the expression platform. We demonstrated that co-transcriptional folding with the

native *E. coli* RNAP is essential for Mn-coupled structural rearrangements in the *mntP*

and *alx* riboswitches *in vivo*, which prompted us to perform co-transcriptional chemical

probing experiments.

The co-transcriptional chemical probing revealed that 5'-proximal halves in both *alx*

and *mntP* aptamers traverse low-T_m RNA hairpin intermediates during folding,

presumably as a safeguard against premature folding of RNA into stable but non-

functional folds. Surprisingly, we found that both riboswitches show Mn-dependent

reactivity changes *before* the full aptamer domain has folded. The Mn-dependent

structural changes in aptamer folding intermediates map to nucleotides in the *alx* and

mntP metal-sensing cores, indicating that a partially formed metal-sensing core can

interrogate the environment for Mn. Curiously, these Mn-dependent differences emerge

at much earlier transcript lengths for *alx* aptamer folding, which may permit a rapid

response to the subtle increase in [Mn] prompted by cytosol alkalinization[19].

Notably, other riboswitches that have been investigated with co-transcriptional

SHAPE-seq or SHAPE-MaP (ZTP, ppGpp, and fluoride) did not show ligand-dependent

structural changes for folding intermediates[27]. Ligands such as ZTP and ppGpp are larger

than an Mn ion and require a network of interactions to bind their cognate aptamers[42,43].

The resulting structural complexity of these aptamer-ligand interactions likely prevents

their stable interaction with an aptamer folding intermediate. In the case of fluoride

riboswitch, binding of fluoride anion to the aptamer is mediated by a network of Mg^{2+} ions

to overcome steric repulsion by the phosphate backbone[44], making fluoride less likely to

bind an aptamer folding intermediate. Additionally, it will be interesting to see whether

other divalent-sensing riboswitches (e.g., Mg^{2+}- and Ni^{2+}/Co^{2+}-sensing riboswitches) can

begin sampling for ligand before the complete aptamer has folded.

To get at the molecular basis of the divergent pH response of the *alx* and *mntP*

riboswitches, we performed co-transcriptional DMS probing of RNA at both neutral and

alkaline pH ±Mn. We found that pH uniquely affects the structure of the *alx* and *mntP*

metal-sensing cores, providing the first clues to how pH differentially affects Mn sensing.

The *alx* and *mntP* aptamers differ in their global architecture, forming a 3WJ and a 4WJ,

respectively. Future study should investigate how nucleobase protonation states in the

alx and *mntP* metal-sensing cores are coupled to junction dynamics, which controls the

docking of the aptamer's helical legs. Prior smFRET study on the *X. oryzae* aptamer, a

4WJ aptamer like *mntP*, illustrated how Mn binding promotes docking of the two helical

legs that in turn stabilizes the P1.1 switch helix[12]. The dynamics of *alx*-like aptamers has

not yet been studied but may play a role in the differential pH effect on *alx* vs. *mntP* Mn

sensing and consequent riboswitching.

The stark differences in the flexibility, architecture, and pH/Mn response of the

mntP vs *alx* riboswitches may be coupled to the cellular function of their respective Mn

exporters. We showed that alkaline pH alone does not rearrange the *alx* riboswitch into a

translationally active conformer; rather, the *alx* riboswitch senses the increased cytosolic

[Mn] brought about by cytosol alkalinization. We hypothesize that alkaline pH lowers the

K_D of the *alx* aptamer for Mn, explaining why an alkaline pH shift and consequent increase

of [Mn] from 28 to 42 μM (free and bound) activates expression of *alx* but not *mntP*[19] (Fig.

9). The physiological functions of such pH-enhanced expression of Mn exporters could

be multi-fold. For example, Mn(II) is oxidized at alkaline pH[45,46], deeming it unusable by

Mn-dependent enzymes. Further, oxidized Mn(IV) is an extremely reactive agent able to

oxidize carbon-carbon double bonds in nucleobases, the capability exploited in

permanganate footprinting of protein-nucleic acid complexes[47,48]. Removal of Mn by

exporters to minimize the pool of oxidized Mn could thus be a protective measure for the

cell experiencing alkaline stress.

Limitations of this study. There are multiple limitations to the experimental approaches

used here. First, co-transcriptional chemical probing experiments are an endpoint

measurement and do not capture the true structural intermediates traversed by an RNA

during transcription. Further, the DMS probing reaction is 5 min., meaning that the co-

transcriptionally displayed RNA can equilibrate to some degree. As a control, we

performed DMS probing experiments on equilibrium refolded intermediates, which

showed structural differences. However, we cannot rule out the possibility that the RNA

folds observed with co-transcriptional DMS probing are not always representative of a

true folding intermediate. Lastly, since the DMS probing reactions were performed *in vitro*,

the absence of ribosomal proteins may have altered the observed structural ensemble,

which should be addressed in future work.

Methods

The source and catalog number (if applicable) for all key reagents are provided as a Supplementary Table 1.

All oligonucleotides were purchased from either Integrated DNA Technologies or Eton Bioscience. A summary of all oligonucleotides used in this work including sequences, modifications, and purifications is presented in Supplementary Table 2.

Procedures for protein preparations, construction of plasmids and bacterial strains, and preparation of DNA templates are provided in the Supplemental Methods.

Single-round *in vitro* transcription for TECprobe-VL experiments

To prepare the holoRNAP, *E. coli* RNAP (6 μM) and σ^{70} (12 μM) were combined in 1x TECprobe transcription buffer (100 mM Tris-Cl, pH 7.2 or 8.5, 50 mM KCl, and 1 mM DTT) and incubated for 30 min at 37 °C. The single-round *in vitro* transcription reactions for TECprobe-VL experiments were assembled by first combining 1x transcription buffer, 40 nM DNA template (*alx* riboswitch, Template 3, Supplementary Table 3) or 50 nM DNA template (*mntP* riboswitch, Template 6, Supplementary Table 3) with randomly distributed internal biotin-11 dNMPs (AAT BioQuest/Biotium), 0.5 μM holoRNAP, and 500 μM ATP, UTP, CTP, and GTP. The transcription buffer omitted magnesium, which was a part of the start solution added later to initiate transcription elongation. Tris-Cl was added at a final concentration of 100 mM to minimize pH changes during the chemical probing reaction, as described previously[27]. The reaction was incubated for 10 min at 37 °C to form open promoter complexes. Next, SAvPhire Monomeric Streptavidin (Sigma-Aldrich) was added to the reaction at a final concentration of 5 μM, followed by incubation for 10 min at 37 °C to allow streptavidin binding to biotinylation sites on the DNA template. Transcription was initiated by adding 10X start solution (100 mM MgCl$_2$, 100 μg/ml rifampicin) to the reaction to a final concentration of 10 mM MgCl$_2$ and 10 μg/ml rifampicin to inhibit reinitiation. After addition of start solution, the final reaction volume was 60 μl. To test the effect of the Mn ligand on riboswitch structural rearrangements, MnCl$_2$ was added to a final concentration of 1 mM in the transcription reactions and was introduced alongside the start solution. The transcription reaction was incubated for 30 s at 37 °C before proceeding to the chemical probing, which is described in the next section.

Single-round *in vitro* transcription for the single-length TECprobe-SL experiments followed the same protocol as outlined above for variable length TECprobe-VL experiments. The DNA templates used for TECprobe-SL experiments contained a terminal biotin modification that was introduced by a 5' biotinylated reverse primer in the PCR (*alx* riboswitch, Template 7, Supplementary table 3).

RNA chemical probing during TECprobe-VL experiments

Benzoyl cyanide (BzCN, Sigma-Aldrich) probing was performed by splitting the 60-μl transcription reaction into 25 μl aliquots and mixing with either 2.78 μl of 400 mM BzCN dissolved in anhydrous DMSO (Invitrogen) to a final concentration of 17.7 mM [(+) sample] or with anhydrous DMSO [(-) sample]. Since the BzCN probing reaction is complete within ~1 s, no additional incubation was required before quenching the reactions with 250 μl of TRIzol reagent (Invitrogen).

DMS probing was performed by splitting the 60-μl transcription reaction into 25 μl aliquots and mixing with either 2.78 μl of 6.5% v/v DMS (Sigma-Aldrich) in 200 proof ethanol [(+) channel] or with 200 proof ethanol [(-) channel]. The samples were incubated for 5 min at 37 °C. The DMS probing reaction was quenched by adding 6.67 μl of 14.5 μM 2-mercaptoethanol and incubating the sample at 37 °C for 1 min. *In vitro* transcription reactions were then quenched by addition of 250 μl of TRIzol reagent.

RNA purification post-TECprobe-VL probing

The RNA was purified from the quenched *in vitro* transcription reactions using the following protocol. First, 50 μl of chloroform was added to the samples and the samples were mixed by vortexing followed by centrifugation at 18,000g and 4 °C for 5 min. Next, the aqueous layer was transferred to a new tube and nucleic acids were precipitated by addition of 105 μl cold isopropanol and 1 μl GlycoBlue Coprecipitant (Invitrogen). The samples were incubated at room temperature for 15 min and centrifuged at 18,000g and 4 °C for 15 min. The supernatant was discarded and the pellet containing the precipitated nucleic acids was washed by addition of 200 μl 70% ethanol. Following the 70% ethanol wash, the samples were centrifuged at 18,000g and 4 °C for 2 min and the supernatant was discarded. The pelleted nucleic acids were resuspended in 25 μl of 1x DNase buffer (10 mM Tris-Cl, pH 7.5, 2.5 mM MgCl$_2$) and mixed with 0.75 μl of Turbo DNase (Invitrogen) to remove residual DNA template. The samples were incubated at 37 °C for 15 min and DNase digestion was quenched by addition of 250 μl TRIzol reagent (Invitrogen). A second TRIzol extraction was performed as described above, except the pellet containing the DMS-modified RNA transcripts was resuspended in 5 μl of 10% (v/v) DMSO.

RNA 3' adapter ligation onto the TECprobe-VL purified transcripts

The 9N_VRA3 adapter oligonucleotide (0139, Supplementary Table 2) was pre-adenylated with the 5' DNA Adenylation Kit (New England Biolabs) using the following protocol. A 100 μl master mix was prepared that consisted of the 1X DNA Adenylation Buffer (New England Biolabs), 100 μM ATP, 5 μM 9N_VRA3 oligo, and 5 μM Mth RNA Ligase (New England Biolabs). The 100 μl master mix was split into 4 aliquots (25 μl each) in PCR tubes and incubated in a thermal cycler set to 65 °C for 1 h. To clean up the adenylation reactions, reactions were pooled to 50 μl followed by addition of 500 μl TRIzol (Invitrogen) and 100 μl of chloroform to each 50 μl pool. The samples were

vortexed to mix and centrifuged at 18,000g and 4 °C for 5 min. The aqueous phase was
transferred to a new tube and the adapter was ethanol precipitated. The pellet was
resuspended in Tris-Cl, pH 8.0, and its concentration was quantified with the Qubit ssDNA
Assay Kit (Invitrogen) and a Qubit 3 Fluorometer. The pre-adenylated adapter was diluted
to 1 μM, divided into small aliquots, and stored at -20 °C.

The RNA 3' adapter ligation reactions were performed using the following protocol.
Purified RNA redissolved in 5 μl of 10% (v/v) DMSO was combined with 1x T4 RNA Ligase
Buffer (New England Biolabs), 0.5 U/μl SuperaseIN (Invitrogen), 0.05 μM pre-adenylated
9N_VRA3 adapter, 15 % PEG 8000, and 5 U/μl T4 RNA Ligase 2, truncated KQ (New
England Biolabs). The final volume for the ligation reactions was 20 μl and the reactions
were incubated at 25 °C for 2 h.

SPRI bead purification of adapter-ligated RNA

A solid-phase reversible immobilization (SPRI) bead purification protocol was used to
deplete excess 9N_VRA3 adapter oligonucleotide. The SPRI beads were prepared with
SpeedBead Magnetic Carboxylate Beads (Cytiva) using the protocol from Jolivet and
Foley[50]. The ligation reactions were combined with 17.5 μl of water, 40 μl anhydrous
isopropanol, and 22.5 μl of the SPRI beads. The samples were vortexed and incubated
at room temperature for 5 min. The samples were then placed on a magnetic separation
rack and the beads collected on the tube wall before disposing of the supernatant.
Keeping the tubes on the magnetic stand, the beads were washed twice with 200 μl of
70% ethanol. After discarding the second ethanol wash, the tubes were placed in a dry
bath set to 37 °C to evaporate the residual ethanol. The beads were resuspended in 25
μl of elution buffer (20 mM Tris-Cl pH 8.0, 50 mM KCl, 1 mM DTT, 0.1 mM EDTA, 0.1%
Triton X-100, 10 mM MgCl$_2$) and the samples were incubated for 3 minutes at room
temperature to elute the RNA. The samples were placed on a magnetic rack and beads
collected on the tube wall before aliquoting the supernatant containing the RNA into a
fresh tube.

TECprobe-VL cDNA synthesis and cleanup

The reverse transcription reactions were performed via capture of RNA transcripts on
streptavidin beads using a 5' biotinylated reverse transcription primer (0159,
Supplementary table 2), which anneals to the 9N_VRA3 sequence that was appended to
the intermediate riboswitch transcripts. Dynabeads MyOne Streptavidin C1 beads
(Invitrogen) were equilibrated as described previously[27]. 5 μl of 500 nM RT primer was
combined with the purified RNA samples and annealed in a thermal cycler using the
following program: 70 °C for 5 min, ramp to 50 °C at 0.1 °C /s, 50 °C for 5 min, ramp to
40 °C at 0.1 °C /s, 40 °C for 5 min, cool to 25 °C. The equilibrated streptavidin beads were
placed on a magnetic stand to allow the beads to collect on the tube wall and the
supernatant was discarded. The streptavidin beads were then resuspended with the
samples containing RNA annealed to the 5' biotinylated RT primer and incubated at room

temperature for 15 min on an end-over-end rotator. The samples were placed on a
magnetic stand and the beads were allowed to collect on the tube wall before discarding
the supernatant. The beads were resuspended in 19.5 μl of reverse transcription master
mix and incubated in a thermal cycler set to 42 °C for 2 min before addition of the reverse
transcriptase. Upon addition of 0.5 μl of SuperScript II Reverse Transcriptase (Invitrogen)
to each sample, the 20 μl RT reactions contained 1x first strand buffer (50 mM Tris-Cl,
pH 8.0, 75 mM KCl), 500 μM dNTPs, 10 mM DTT, 2 % formamide, 10 ng/μl Extreme
Thermostable Single-Stranded DNA Binding Protein (ET SSB) (New England Biolabs),
0.1 % Triton-X-100, and 3 mM $MnCl_2$. The reverse transcription reactions were incubated
at 42 °C for 180 min followed by a 70 °C incubation for 50 min to inactivate the SuperScript
II and then cooled to 12 °C. Next, the RNAs were degraded by adding 1.25 U RNase H
(New England Biolabs) and 12.5 U RNase I_f (New England Biolabs) to each sample and
the samples were incubated at 37 °C for 20 min followed by a 70 °C incubation for 20 min
to inactivate the RNases. The samples were placed on a magnetic stand and the beads
were allowed to collect on the tube wall. The supernatant was discarded and the beads
were resuspended in 75 μl of Tris Wash Buffer (10 mM Tris-Cl, pH 8.0, 0.05% Triton X-
100) to wash the beads. The samples were placed on a magnetic stand and the wash
was discarded. The bead-bound cDNA was resuspended in 25 μl of the Tris Wash Buffer
and stored at -20 °C.
677
678 **Chemical probing of equilibrium-refolded intermediate transcripts**
679
Single-round *in vitro* transcription using randomly biotinylated DNA templates was
executed as described for TECprobe-VL experiments above. To quench transcription,
600 μl of TRIzol was added to the reactions. Next, 120 μl of chloroform was added to the
samples and the samples were vortexed to mix followed by centrifugation at 18,000g and
4 °C for 5 min. The aqueous layer was transferred into a new tube and nucleic acids were
isopropanol precipitated as described for TECprobe-VL experiments except a larger
volume of isopropanol was used to account for the larger sample volume. The pellets
containing the intermediate RNA transcripts were resuspended in 30 μl of 1x TECprobe
transcription buffer (100 mM Tris-Cl, pH 7.2 or 8.5, 50 mM KCl, 1 mM DTT), placed on a
heat block set to 95 °C for 2 min, and snap-cooled on ice for 1 min. To each sample, 30
μl of 2x equilibration buffer (200 mM Tris-Cl, pH 7.2 or 8.5, 100 mM KCl, 2 mM DTT, 20
mM $MgCl_2$) was added, which contained either no Mn or 2 mM $MnCl_2$ (1 mM final in 60 μl
reaction), and the samples were incubated at 37 °C for 20 min. The RNAs were then
probed with DMS as described for TECprobe-VL experiments, and the DMS-probed RNA
was purified as described for TECprobe-VL experiments.
695
696 **Preparation of TECprobe-VL dsDNA libraries for sequencing**
697
cDNA libraries were PCR-amplified for high throughput sequencing using the following
protocol. Separate PCR reactions were performed for the (+) and (-) samples, which
contained 1x Q5 Reaction Buffer, 1x Q5 GC Enhancer, 200 μM dNTPs, 250 nM RPI
Indexing Primer (0163-0178,0235-0248, Supplementary table 2), 250 nM

dRP1_NoMod.R reverse primer (0162, Supplementary Table 2), 10 nM SC1Brdg_MINUS
or SC1Brdg_PLUS channel barcode oligo (0160,0161, Supplementary Table 2), 12 μl of
bead-bound cDNA library, and 0.02 U/μl Q5 High-Fidelity DNA Polymerase (New England
Biolabs). Amplification was performed with the following thermal cycler program: 98 °C
for 30 s, [98 °C for 10s, 62 °C for 20 s, 72 °C for 20s] x 22 cycles, 72 °C for 2 min, hold
at 12 °C.
708
709 **High-throughput DNA sequencing of TECprobe-VL libraries**
710
711 Sequencing of the TECprobe-VL and equilibrium refolded intermediate transcript libraries
712 was performed by Novogene Co. on a Novaseq X Plus system with 2 x 150 Paired End
713 (PE) reads and a 20% PhiX spike in. The TECprobe-VL and equilibrium refolded
714 intermediate transcript libraries were sequenced at a depth of ~30-60 million PE reads.
715
716 Sequencing of TECprobe-SL libraries was performed by Novogene Co. on a Novaseq X
717 Plus system with 2 x 150 PE reads. No PhiX spike in was added since these libraries
718 were sequenced in a partial lane. The TECprobe-SL libraries were sequenced at a depth
719 of ~10 million PE reads.
720
721 **TECprobe-VL data processing and visualization**
722
Sequence read pre-processing, target generation using cotrans_preprocessor, and
shapemapper2 run script generation were performed exactly as described in the
published protocol using the publicly available TECprobe data processing tools
(https://github.com/e-strobel-lab/TECtools). TECprobe visualization tools was used to
assemble reactivity matrices and analyze the correlation between replicate data sets
(https://github.com/e-strobel-lab/TECprobe_visualization). The compile_sm2_output
script was used to assemble the reactivity profiles for intermediate transcripts into matrix
format and to normalize the dataset. The normalized reactivity matrices were visualized
in heatmap format using GraphPad Prism.
732
For the *alx* riboswitch, replicate data sets were required to obtain adequate sequencing
depth for longer transcripts, as there was a strong bias in representation towards shorter
transcripts for the amplified cDNA libraries. Plots with fraction of aligned reads per
transcript length and alignment rates for *alx* co-transcriptional TECprobe-VL experiments
are presented in SI Fig. 21. Additionally, the presence of two intrinsic termination sites in
the *alx* riboswitch sequence (110 and 190 nt) results in a clear drop in sequencing depth
for intermediate transcripts past these sites (SI Fig. 19A). For the co-transcriptional DMS-
MaP libraries, 5-7 independent replicates were pooled per chemical probing condition to
increase data quality for longer transcript lengths (see SI Figs 22-25 for individual and
concatenated heatmaps). Plots with fraction of aligned reads per transcript length and
alignment rates for *alx* equilibrium refolded TECprobe-VL experiments are presented in
SI Fig. 26. For the equilibrium refolded *alx* TECprobe-VL experiments, 2 independent
replicates were pooled per chemical probing condition to increase data quality for longer

transcript lengths (see SI Figs 27-30 for individual and concatenated heatmaps). To
further increase sequencing depth when necessary, multiple independent PCR
amplifications were performed with the same cDNA library and sequenced; reads were
subsequently pooled together for data processing.
750
For the *mntP* riboswitch, two replicate data sets were concatenated and analyzed
together after assessing the correlation between replicates using the
plot_cotrans_correlation script available in TECprobe visualization tools. Plots with
fraction of aligned reads per transcript length and alignment rates for co-transcriptional
mntP TECprobe-VL experiments are presented in SI Fig. 31 Plots with fraction of aligned
reads per transcript length and alignment rates for equilibrium refolded *mntP* TECprobe-
VL experiments are presented in SI Fig. 32. To further increase sequencing depth when
necessary, multiple independent PCR amplifications were performed with the same cDNA
library and sequenced; reads were subsequently pooled together for data processing.
760
761
762 **In vitro synthesis of full-length riboswitch RNAs with T7 RNAP and cleanup for**
763 **SHAPE-MaP and DMS-MaP**
764
RNA for *in vitro* SHAPE/DMS-MaP was synthesized using the HiScribe T7 High Yield
RNA Kit (New England Biolabs). 100 μL reactions were assembled according to
manufacturer instructions and incubated at 37 °C overnight. RNA was purified and
concentrated using the RNA Clean and Concentrator kit (Zymo Research). For the *alx*
riboswitch and *alx* aptamer, RNAs were gel-purified to isolate full-length transcripts away
from premature transcription termination products using a denaturing PAGE (8%
acrylamide, 8 M urea, 0.5x TBE (Fisher Scientific)). For gel-purification, RNAs were
diluted 1:5 in buffer and mixed with an equal volume 2x formamide loading dye (95%
formamide, 0.5 mM EDTA, 0.025% bromophenol blue, 0.025% xylene cyanol), heated at
95 °C for 2 mins, and loaded on a pre-heated gel. Gel was run at 50 W, stained with 1x
SYBR Green II RNA Gel Stain (Invitrogen) in 1X TBE for 10 mins, and visualized using a
Edvotek TruBlu 2 Blue Transilluminator. Bands corresponding to full-length RNA were
excised and eluted in 10 mL TE buffer (10 mM Tris pH 8.0, 1 mM EDTA pH 8.0) at 4 °C
overnight. The eluted solution of RNA was spun to pellet acrylamide, and RNA was
concentrated using an Amicon Ultra Centrifugal filter (3 kDa molecular weight cutoff).
RNAs were ethanol precipitated, stored at -20 °C, and resuspended in RNase-free H_2O.
For the *mntP* riboswitch, there are no terminated transcripts, eliminating the need for gel
purification. After transcription and cleanup, *mntP* RNAs were ethanol precipitated, stored
at -20 °C, and resuspended in RNase free H_2O.
784
SHAPE-MaP and DMS-MaP on the full-length equilibrium refolded *alx* and *mntP*
riboswitches
5 pmol of RNA in added RNase-free H_2O to a volume of 6 μL was heat denatured for 2
mins at 95 °C, cooled at 4 °C for 2 mins, and then 3 μL buffer was added for a final volume
of 9 μL. The final buffer concentration was 100 mM HEPES, 100 mM NaCl, and 5 mM

MgCl$_2$ at either pH 7.2 or pH 8.4 and supplemented with 1 mM MnCl$_2$ where indicated.
RNA was incubated at 37 °C for 20 mins to allow refolding. For modification with 1M7, 9
μL of RNA was mixed with 1 μL of 100 mM 1M7 ("+" modified sample) or 1 μL DMSO ("-
" negative control), incubated for 75 s at 37 °C, and placed on ice. For the denatured
control, 5 pmol RNA (3 μL) was mixed with 5 μL formamide and 1 μL 10x denaturing
buffer (500 mM HEPES, 40 mM EDTA) at pH 7.2 or 8.4. RNA was denatured for 1 min at
95 °C, mixed with 1 μL of 100 mM 1M7, incubated at 95 °C for 1 min, and placed on ice.
All RNAs were purified and eluted in 50 μL of RNase-free H$_2$O using MicroSpin G-25
columns (Cytiva) and stored at -20 °C. For modification with DMS, 9 μL of RNA was mixed
with 1 μL of 6.5% v/v DMS ("+" modified sample) or 1 μL ethanol ("-" negative control),
incubated for 5 minutes at 37 °C, quenched with 2.4 μL of 30% v/v 2-mercaptoethanol
solution at 37 °C for 1 min before placing on ice. For the denatured control, 5 pmol RNA
(3 μL) was mixed with 5 μL formamide and 1 μL 10x denaturing buffer (500 mM HEPES,
40 mM EDTA) at pH 7.2 or 8.4. Then RNA was denatured for 1 min at 95 °C, mixed with
1 μL of 6.5% v/v DMS, incubated at 95 °C for 1 min, quenched with 2.4 μL of 30% 2-
mercaptoethanol solution at 37 °C for 1 min before placing on ice. All RNAs were purified
and eluted in 50 μL of RNase-free H$_2$O using MicroSpin G-25 columns (Cytiva) and stored
at -20 °C.
808
Reverse transcription (RT) of 10 μL of RNA was performed by annealing 1 μL RT primer
(0277) for 5 min at 65 °C. Then, 8 μL RT buffer was added (50 mM Tris-Cl pH 8, 75 mM
KCl, 10 mM DTT, 0.5 mM dNTPs, and MnCl$_2$ added to 15 mM before use). Samples were
incubated for 2 min at 42 °C before introducing Superscript II reverse transcriptase
(Thermo Fisher Scientific). RT reactions were incubated for 3 h at 42 °C. The reverse
transcriptase was inactivated for 15 min at 70 °C before storing the reactions at 4 °C.
cDNA was purified and eluted in 50 μL of RNase-free H$_2$O using MicroSpin G-25 columns
(Cytiva) and stored at -20 °C. The first PCR (step 1) amplified cDNA through 20
amplification cycles using primers 0278 and 0279. Excess primer was removed using the
Monarch PCR & DNA Cleanup Kit (New England Biolabs). The second PCR (step 2)
further amplified the first PCR amplicon through an additional 10 cycles. The step 2 PCR
used primers 0149 and an RPI indexing primer (0163-0178 and 0235-0248). Step 2 PCR
was cleaned up using Agencourt AMPure XP beads (Beckman Coulter), and final DNA
concentration was measured using the Invitrogen Quant-IT dsDNA High-Sensitivity HS
Assay kit quantified with a BioTek Synergy H1 Multimode Microplate Reader before
pooling libraries for sequencing. Sequencing of these pooled libraries was performed by
Novogene Co. on a NovaSeq 6000 system with 100M 150 PE reads.
826
827 ***In vivo* DMS-MaP of the full-length *alx* riboswitch**
828
The plasmid pMIS518 was transformed into MC4100Δ*alx* cells, and a 5-mL culture was
grown at 37 °C overnight with shaking at 200 rpm. A 1:100 dilution of overnight culture
into 20 mL of fresh potassium-modified LB medium (LBK) media (LBK at pH 7.2 or 8.4

and with supplemented MnCl$_2$ where indicated) was grown at 37 °C with shaking at 200
rpm to an optical density of 0.3-0.5 at 600 nm (OD$_{600}$). The LBK was prepared by replacing
NaCl with equimolar KCl, and was buffered with 100 mM N-[Tris(hydroxymethyl)methyl]-
3-amino propane sulfonic acid (TAPS) and pH adjusted with KOH to 8.4 for the alkaline
medium. For the DMS-probed sample, 10 mL of culture was exposed to 10 μL of ≥99.5%
DMS (Sigma-Aldrich), shaken at 200 rpm at 30 °C for 5 min, and quenched with 1.6 mL
of 0.6 M 2-mercaptoethanol. For the untreated sample, 10 mL of culture was shaken at
200 rpm at 30 °C for 5 min, and quenched with 1.6 mL of 0.6 M 2-mercaptoethanol. The
cultures were spun at 4 °C and the pellets were resuspended in 0.6 mL of hot total RNA
lysis buffer pre-heated to 60 °C (1 mM EDTA, 20 mM Na acetate pH 5.5, and 0.5% SDS).
The sample was phenol-chloroform extracted twice, followed by two chloroform
extractions and ethanol precipitation. The pellet was then resuspended in 100 μL RNase-
free H$_2$O, treated with 10 U Turbo DNase for 1 h at 37 °C, and again phenol-chloroform
extracted, followed by chloroform extraction, ethanol precipitation, and pellets were
resuspended in 100 μL RNase-free H$_2$O. 0.5 μL purified total RNA was used for RT, and
all subsequent library preparation steps were identical to the above *in vitro* DMS-MaP
protocol.
849
850 **RNA structure prediction and visualization**
851
RNA structure prediction for co-transcriptionally probed riboswitch folding intermediates
was performed using the RNAstructure V6.4[51] Fold command with default settings,
adding the DMS outputs as a constraint, and forcing the 3'-most 11 to 14 nt to be single-
stranded to account for the *E. coli* RNAP footprint. The set of minimum free-energy
structures were visualized in VARNA[52] to assess agreement with the DMS probing data,
and structure diagrams were colored in accordance with DMS reactivity. Additionally,
existing X-ray crystal structures for Mn-sensing aptamer domains were visualized with
UCSF Chimera[53] and used to guide modeling of conserved secondary and tertiary
structure motifs within the *alx* and *mntP* aptamers. RNA structure prediction for
equilibrium refolded riboswitch folding intermediates was performed as described above;
however, the 3' most nt were not kept single-stranded since the intermediate RNA
transcripts were extracted from stalled transcription elongation complexes before DMS
probing.
865
866 The denatured and equilibrium-refolded full-length *mntP* riboswitch*, alx* riboswitch*, and
867 *alx* aptamer RNA structures were predicted using the RNAStructure V6.4[51] Fold
868 command with default settings, adding the SHAPE or DMS outputs as a constraint. The
869 minimum free energy structures were visualized in VARNA[52] and colored in accordance
870 with nucleotide reactivity.
871
872
873
874

β-galactosidase assays

Overnight culture of each strain was diluted 1:100 into LBK with appropriate antibiotic at either pH 6.8 or pH 8.4, with or without 1 mM supplemented $MnCl_2$. The cultures were grown to mid-log phase and supplemented with 1 mM IPTG where necessary for expression of T7 RNAP. β-galactosidase assays were performed using the Miller method[54]. Briefly, bacterial growth was measured using OD_{600} before cultures were placed on ice. 100 μL cells were lysed via vortexing in 900 μL sodium phosphate buffer (100 mM sodium phosphate pH 7, 10 mM KCl, 2 mM $MgSO_4$, 38 mM 2-mercaptoethanol), 20 μL 0.1% SDS, and 20 μL chloroform. Substrate O-nitrophenyl-β-D-galactopyranoside (ONPG) was added and the reaction was stopped with 1 M Na_2CO_3 when a yellow color appeared, or after 35 min. After centrifuging 5 min at 16,000 g, absorbance was measured at 420 nm (A_{420}) to calculate activity using Miller units. Each culture was tested in duplicate and A_{420} values were averaged. Each final reported value is an average of three replicates with standard deviation.

Sequence analysis for *yybP-ykoY* riboswitches

The sequences for *yybP-ykoY* riboswitches were downloaded from the Rfam database (Rfam reference number RF00080) and aligned using the Kalign multi sequence alignment program (https://www.ebi.ac.uk/jdispatcher/msa). The sequence alignment and consensus L3 sequence was visualized using Jalview[55]. The *yybP-ykoY* riboswitch sequences found within the Enterobacteriales order were accessed from both the Rfam database and from *yybP-ykoY* regulog collection (https://regprecise.lbl.gov/collection_rfam.jsp?riboswitch_id=13). These sequences were aligned with the Muscle multi sequence alignment program (https://www.ebi.ac.uk/jdispatcher/msa). The sequence alignment and consensus L3 sequence was visualized using Jalview[55].

Data Availability

The raw sequencing data generated in this study will be deposited into the Sequencing Read Archive (https://www.ncbi.nlm.nih.gov/sra) with the BioProject accession code PRJNA1197522. Individual BioSample accession codes will be provided in a Supplementary Table. All processed reactivity data will be deposited in the RNA Mapping Database (https://rmdb.stanford.edu/) and the individual accession codes will be provided in a Supplementary Table.

References

1.   Imlay, J. A. The Mismetallation of Enzymes during Oxidative Stress. *J. Biol. Chem.* **289**,

28121–28128 (2014).

2.   Imlay, J. A. The molecular mechanisms and physiological consequences of oxidative

stress: lessons from a model bacterium. *Nat. Rev. Microbiol.* **11**, 443–454 (2013).

3.   Chandrangsu, P., Rensing, C. & Helmann, J. D. Metal homeostasis and resistance in

bacteria. *Nat. Rev. Microbiol.* **15**, 338–350 (2017).

4.   Lee, J.-W. & Helmann, J. D. Functional specialization within the Fur family of

metalloregulators. *BioMetals* **20**, 485–499 (2007).

5.   Glasfeld, A., Guedon, E., Helmann, J. D. & Brennan, R. G. Structure of the manganese-

bound manganese transport regulator of *Bacillus subtilis*. *Nat. Struct. Mol. Biol.* **10**,

652–657 (2003).

6.   Price, I. R., Gaballa, A., Ding, F., Helmann, J. D. & Ke, A. Mn^{2+}-Sensing Mechanisms of

yybP-ykoY Orphan Riboswitches. *Mol. Cell* **57**, 1110–1123 (2015).

7.   Dann, C. E. *et al*. Structure and Mechanism of a Metal-Sensing Regulatory RNA. *Cell*

130, 878–892 (2007).

8.   Furukawa, K. *et al*. Bacterial Riboswitches Cooperatively Bind Ni^{2+} or Co^{2+} Ions and

Control Expression of Heavy Metal Transporters. *Mol. Cell* **57**, 1088–1098 (2015).

9.   Serganov, A. & Nudler, E. A Decade of Riboswitches. *Cell* **152**, 17–24 (2013).

10.   Dambach, M. *et al*. The Ubiquitous *yybP-ykoY* Riboswitch Is a Manganese-Responsive

Regulatory Element. *Mol. Cell* **57**, 1099–1109 (2015).

11. Kavita, K. & Breaker, R. R. Discovering riboswitches: the past and the future. *Trends*

Biochem. Sci. **48**, 119–141 (2023).

12. Suddala, K. C. *et al.* Local-to-global signal transduction at the core of a Mn^{2+} sensing

riboswitch. *Nat. Commun.* **10**, 4304 (2019).

13. Sung, H.-L. & Nesbitt, D. J. Single-Molecule FRET Kinetics of the Mn^{2+} Riboswitch:

Evidence for Allosteric Mg^{2+} Control of "Induced-Fit" vs "Conformational Selection"

Folding Pathways. *J. Phys. Chem. B* **123**, 2005–2015 (2019).

14. Wickiser, J. K., Winkler, W. C., Breaker, R. R. & Crothers, D. M. The Speed of RNA

Transcription and Metabolite Binding Kinetics Operate an FMN Riboswitch. *Mol. Cell*

18, 49–60 (2005).

15. Chauvier, A. *et al.* Transcriptional pausing at the translation start site operates as a

critical checkpoint for riboswitch regulation. *Nat. Commun.* **8**, 13892 (2017).

16. Strobel, E. J., Cheng, L., Berman, K. E., Carlson, P. D. & Lucks, J. B. A ligand-gated strand

displacement mechanism for ZTP riboswitch transcription control. *Nat. Chem. Biol.* **15**,

1067–1076 (2019).

17. Chauvier, A., Dandpat, S. S., Romero, R. & Walter, N. G. A nascent riboswitch helix

orchestrates robust transcriptional regulation through signal integration. *Nat.*

Commun. **15**, 3955 (2024).

18. Nechooshtan, G., Elgrably-Weiss, M., Sheaffer, A., Westhof, E. & Altuvia, S. A pH-

responsive riboregulator. *Genes Dev.* **23**, 2650–2662 (2009).

19. Sharma, R. & Mishanina, T. V. A riboswitch-controlled TerC family transporter Alx tunes

intracellular manganese concentration in *Escherichia coli* at alkaline pH. *J. Bacteriol.*

206, e00168-24 (2024).

20. Zeinert, R. *et al.* Structure–function analysis of manganese exporter proteins across

bacteria. *J. Biol. Chem.* **293**, 5715–5730 (2018).

21. Bachas, S. T. & Ferré-D'Amaré, A. R. Convergent Use of Heptacoordination for Cation

Selectivity by RNA and Protein Metalloregulators. *Cell Chem. Biol.* **25**, 962-973.e5

(2018).

22. Adeolu, M., Alnajar, S., Naushad, S. & S. Gupta, R. Genome-based phylogeny and

taxonomy of the '*Enterobacteriales*': proposal for *Enterobacterales* ord. nov. divided

into the families *Enterobacteriaceae*, *Erwiniaceae* fam. nov., *Pectobacteriaceae* fam.

nov., *Yersiniaceae* fam. nov., *Hafniaceae* fam. nov., *Morganellaceae* fam. nov., and

Budviciaceae fam. nov. *Int. J. Syst. Evol. Microbiol.* **66**, 5575–5599 (2016).

23. Padan, E., Bibi, E., Ito, M. & Krulwich, T. A. Alkaline pH homeostasis in bacteria: New

insights. *Biochim. Biophys. Acta BBA - Biomembr.* **1717**, 67–88 (2005).

24. Hartmann, F. S. F. *et al.* Visualizing the pH in *Escherichia coli* Colonies via the Sensor

Protein mCherryEA Allows High-Throughput Screening of Mutant Libraries. *mSystems*

7, e00219-22 (2022).

25. Smola, M. J., Rice, G. M., Busan, S., Siegfried, N. A. & Weeks, K. M. Selective 2'-hydroxyl

acylation analyzed by primer extension and mutational profiling (SHAPE-MaP) for

direct, versatile and accurate RNA structure analysis. *Nat. Protoc.* **10**, 1643–1669

(2015).

26. Szyjka, C. E. & Strobel, E. J. Cotranscriptional RNA Chemical Probing. in *Riboregulator*

Design and Analysis (eds. Chappell, J. & Takahashi, M. K.) vol. 2518 291–330 (Springer

US, New York, NY, 2022).

27. Szyjka, C. E. & Strobel, E. J. Observation of coordinated RNA folding events by

systematic cotranscriptional RNA structure probing. *Nat. Commun.* **14**, 7839 (2023).

28. Stephen, C. & Mishanina, T. V. Alkaline pH has an unexpected effect on transcriptional

pausing during synthesis of the *Escherichia coli* pH-responsive riboswitch. *J. Biol.*

Chem. **298**, 102302 (2022).

29. Kaur, G. *et al.* Novel MntR-Independent Mechanism of Manganese Homeostasis in

Escherichia coli by the Ribosome-Associated Protein HflX. *J. Bacteriol.* **196**, 2587–2597

(2014).

30. Kaur, G. *et al.* Affected energy metabolism under manganese stress governs cellular

toxicity. *Sci. Rep.* **7**, 11645 (2017).

31. Martin, J. E., Waters, L. S., Storz, G. & Imlay, J. A. The *Escherichia coli* Small Protein

MntS and Exporter MntP Optimize the Intracellular Concentration of Manganese. *PLOS*

Genet. **11**, e1004977 (2015).

32. Siegfried, N. A., Busan, S., Rice, G. M., Nelson, J. A. E. & Weeks, K. M. RNA motif

discovery by SHAPE and mutational profiling (SHAPE-MaP). *Nat. Methods* **11**, 959–965

(2014).

33. Papp-Wallace, K. M. & Maguire, M. E. Magnesium Transport and Magnesium

Homeostasis. *EcoSal Plus* **3**, 10.1128/ecosalplus.5.4.4.2 (2008).

34. Anand, V. S. & Patel, S. S. Transient State Kinetics of Transcription Elongation by T7 RNA

Polymerase. *J. Biol. Chem.* **281**, 35677–35685 (2006).

35. Wang, M. D. *et al.* Force and Velocity Measured for Single Molecules of RNA

Polymerase. *Science* **282**, 902–907 (1998).

36. Balint, E. & Unk, I. For the Better or for the Worse? The Effect of Manganese on the

Activity of Eukaryotic DNA Polymerases. *Int. J. Mol. Sci.* **25**, 363 (2023).

37. Zubradt, M. *et al.* DMS-MaPseq for genome-wide or targeted RNA structure probing in

vivo. *Nat. Methods* **14**, 75–82 (2017).

38. Mustoe, A. M., Lama, N. N., Irving, P. S., Olson, S. W. & Weeks, K. M. RNA base-pairing

complexity in living cells visualized by correlated chemical probing. *Proc. Natl. Acad.*

Sci. **116**, 24574–24582 (2019).

39. Nechooshtan, G., Elgrably-Weiss, M. & Altuvia, S. Changes in transcriptional pausing

modify the folding dynamics of the pH-responsive RNA element. *Nucleic Acids Res.* **42**,

622–630 (2014).

40. Saenger, W. *Principles of Nucleic Acid Structure*. (Springer New York, New York, NY,

1984). doi:10.1007/978-1-4612-5190-3.

41. Liberman, J. A. *et al.* Structural analysis of a class III preQ$_1$ riboswitch reveals an

aptamer distant from a ribosome-binding site regulated by fast dynamics. *Proc. Natl.*

Acad. Sci. **112**, (2015).

42. Jones, C. P. & Ferré-D'Amaré, A. R. Recognition of the bacterial alarmone ZMP through

long-distance association of two RNA subdomains. *Nat. Struct. Mol. Biol.* **22**, 679–685

(2015).

43. Peselis, A. & Serganov, A. *ykkC* riboswitches employ an add-on helix to adjust

specificity for polyanionic ligands. *Nat. Chem. Biol.* **14,** 887–894 (2018).

44. Ren, A., Rajashankar, K. R. & Patel, D. J. Fluoride ion encapsulation by Mg^{2+} ions and

phosphates in a fluoride riboswitch. *Nature* **486,** 85–89 (2012).

45. Morgan, J. J. Kinetics of reaction between O_2 and Mn(II) species in aqueous solutions.

Geochim. Cosmochim. Acta **69**, 35–48 (2005).

46. Hu, S. *et al.* Hematite-mediated Mn(II) abiotic oxidation under oxic conditions: pH

effect and mineralization. *J. Colloid Interface Sci.* **636**, 267–278 (2023).

47. Spicuglia, S., Kumar, S., Chasson, L., Payet-Bornet, D. & Ferrier, P. Potassium

permanganate as a probe to map DNA–protein interactions *in vivo*. *J. Biochem.*

Biophys. Methods **59**, 189–194 (2004).

48. Davis, C. A., Bingman, C. A., Landick, R., Record, M. T. & Saecker, R. M. Real-time

footprinting of DNA in the first kinetically significant intermediate in open complex

formation by *Escherichia coli* RNA polymerase. *Proc. Natl. Acad. Sci.* **104**, 7833–7838

(2007).

49. Saba, J. *et al.* The elemental mechanism of transcriptional pausing. *eLife* **8**, e40981

(2019).

50. Jolivet, P. & Foley,J. SPRI bead mix. *protocols.io* (2020) doi:https://doi.org/10.

17504/protocols.io.bnz4mf8w.

51. Reuter, J. S. & Mathews, D. H. RNAstructure: software for RNA secondary structure

prediction and analysis. *BMC Bioinformatics* **11**, 129 (2010).

52. Darty, K., Denise, A. & Ponty, Y. VARNA: Interactive drawing and editing of the RNA

secondary structure. *Bioinformatics* **25**, 1974–1975 (2009).

53. Pettersen, E. F. *et al.* UCSF Chimera—A visualization system for exploratory research

and analysis. *J. Comput. Chem.* **25**, 1605–1612 (2004).

54. Miller, JH. *A Short Course in Bacterial Genetics: A Laboratory Manual and Handbook for*

Escherichia Coli and Related Bacteria. (Cold Spring Harbor Laboratory, Cold Spring

Harbor, NY, 1992).

55. Waterhouse, A. M., Procter, J. B., Martin, D. M. A., Clamp, M. & Barton, G. J. Jalview

Version 2—a multiple sequence alignment editor and analysis workbench.

Bioinformatics **25**, 1189–1191 (2009).

1048

1049

1050

1051

1052

1053

1054

1055

1056

1057

1058

1059

Figures

1061
1062
1063
Figure 1. Introduction to the *E. coli alx* and *mntP* Mn-sensing translational
riboswitches. A. Cartoon model of *alx* and *mntP* riboregulation of their respective genes
in *E. coli*. (Top panel) At alkaline pH, the cytosolic [Mn] increases in *E. coli*, resulting in
the *alx* riboswitch turning "on" expression of Alx, an Mn exporter. (Bottom panel) An
increase in environmental [Mn] results in the *mntP* riboswitch turning "on" expression of
MntP, the major Mn exporter in *E. coli*. **B.** Cartoon models of the *alx* and *mntP*
riboswitches in their translationally active vs. inactive conformations. L3 (purple), L1
(orange), P1.1 (gray), and RBS/start codon (red) are highlighted. **C.** *L. lactis* Mn-sensing

aptamer crystal structure with the *E. coli mntP* L3 sequence (PDB 1D: 6CC3). "Left leg"
and "right leg" refer to coaxially stacked helices in the aptamer. L3 (purple), L1 (orange),
P1.1 (gray), and G-C receptor base pair in cross-helix A-minor (magenta) are highlighted.
Mg^{2+} in the M_A site and Cd^{2+} in the M_B site (captured instead of Mn^{2+}) are green and
yellow spheres, respectively. Sequence alignment of *alx* and *mntP* L1 and L3 are orange
and yellow, respectively.
1078

Figure 2. Aptamer architectures and riboswitch-controlled translational response
in vivo. **A.** *mntP* and *alx* aptamer structure models showing the *mntP* 4WJ and *alx* 3WJ
elements. The key Mn-sensing loops (L1 is orange and L3 is purple) and the P1.1 switch
helix (gray) are highlighted. **B.** Secondary structure model of equilibrium-refolded (pH 7.2
–Mn) *alx* aptamer modeled based on reactivity with 1M7. **C.** Individual nucleotide
reactivity differences when modified with 1M7 at neutral (7.2) vs alkaline pH (8.5),
measured in the presence or absence of Mn. **D.** Translational reporter construct design
for *alx* and *mntP* translational reporters with the native or T7 promoter. **E.** Translational
response of *alx* to supplemented Mn (left) or alkaline pH (right) as a function of promoter.
Statistical analysis done using a parametric unpaired t-test to assess significance. **F.**
Translational response of *mntP* to supplemented Mn (left) or alkaline pH (right) as a
function of promoter. Statistical analysis done using a parametric unpaired t-test to assess
significance.

Figure 3. Co-transcriptional DMS probing of the *mntP* riboswitch aptamer at alkaline pH. A. TECprobe-VL DMS reactivity matrix for the *mntP* aptamer at pH 8.5. Reactivities shown were normalized with a single, whole-dataset calculated normalization factor. Data are from two independent replicates that were concatenated and analyzed together. **B.** Secondary structures of *mntP* aptamer folding intermediates colored by DMS reactivity of the indicated transcript at pH 8.5. **C.** Transcript length-dependent reactivity changes showing folding transitions of the *mntP* aptamer at pH 8.5 ± 1 mM Mn. DMS data for 0 mM Mn (orange) and 1 mM Mn (purple) conditions from Fig. 3A and SI Fig. 3, respectively. *E. coli mntP* apo aptamer (PDB ID: 4Y1M) shown with labeled loops and helical domains. Nt highlighted in orange correspond to the nt positions with reactivities plotted in the adjacent graph. **D.** Plots show transcript length- and Mn-dependent reactivity changes in *mntP* aptamer folding intermediates at pH 8.5 ± 1 mM Mn. DMS data for 0 mM Mn (orange) and 1 mM Mn (purple) conditions from Fig. 3A and SI Fig. 3, respectively. Vertical dotted line marks completion of aptamer folding. *X. oryzae* aptamer metal-sensing core (PDB ID: 6N2V) with Mg^{2+} (green sphere) and Mn^{2+} (purple sphere) highlighted. Conformer 1 shows Mn^{2+} fully bound and Conformer 2 shows Mn^{2+} partially bound. Nt highlighted in purple correspond to nt reactivity plots above.

Figure 4. Co-transcriptional DMS probing of the *alx* riboswitch aptamer at alkaline
pH. A. TECprobe-VL DMS reactivity matrix for the *alx* aptamer at pH 8.5. Reactivities
shown were normalized with a single, whole-dataset calculated normalization factor. Data
are from seven independent replicates that were concatenated and analyzed together. **B.**
Secondary structures of *alx* aptamer folding intermediates, which are colored by the DMS
reactivity of the indicated transcript at pH 8.5. **C.** Transcript length-dependent reactivity
changes showing folding transitions of the *alx* aptamer at pH 8.5 ± 1 mM Mn. DMS data
for 0 mM Mn (orange) and 1 mM Mn (purple) conditions from Fig. 4A and SI Fig. 7,
respectively. Secondary structure model of *alx* aptamer with labeled structural domains.
L1 (orange), L3 (purple), P1.1 (gray), and intrinsic termination site (pink) are highlighted.
Nt boxed in red correspond to adjacent nt reactivity plots. **D.** *L. lactis* Mn-sensing aptamer
with *E. coli alx* L3 sequence (PDB ID: 6CC1). L3 nt (purple), cross-helix A-minor nt: A11
(orange) and G63-C33 (magenta), Mg^{2+} (green sphere), and Cd^{2+} (yellow sphere) are
highlighted. Plots show transcript length- and Mn-dependent reactivity changes in *alx*
aptamer folding intermediates at pH 8.5 ± 1 mM Mn. DMS data for 0 mM (orange) and 1
mM (purple) conditions from Fig. 4A and SI Fig. 7, respectively.
1183
1184

Figure 5. Decoupling the effects of alkaline pH and Mn on *alx* riboswitching *in vivo* and pH-dependent reactivity changes in the divergent *alx* and *mntP* L3 sequences. A. (Top panel) *E. coli* cell with increased cytosolic [Mn] resulting from an alkaline pH challenge (left) and construct design for *alx* translational reporter (right). (Bottom panel) *L. lactis* Mn-sensing aptamer with *E. coli alx* L3 sequence (PDB ID: 6CC1, left). Conserved A37 (purple), Mg^{2+} (green sphere), and Cd^{2+} (yellow sphere) are highlighted. (Bar graph) Loss of *alx* translational response to either Mn or pH upon mutation of the Mn-coordinating adenosine. **B-C**. Reactivity difference heatmaps and reactivity changes in aptamer intermediates for *alx* L3 nt (**B**) and *mntP* L3 nt (**C**) during aptamer folding at pH 7.2 vs. 8.5 ± 1 mM Mn. *alx* L3 and *mntP* L3 (purple), L1 (orange), L3 (purple), P1.1 (gray), and cross-helix A-minor (pink) are highlighted. DMS data for 0 mM Mn (orange) and 1 mM Mn (purple) conditions from Fig. 4A, SI Fig. 7, and SI Fig. 9B (**B**, *alx*) and Fig. 3A, SI Fig. 3, and SI Fig. 9A (**C**, *mntP*). Vertical dotted line marks completion of aptamer folding. **D.** L3 consensus sequences from *yybP-ykoY* family members in the Rfam database (top) and *mntP*-like (middle)/ *alx*-like (bottom) members in the *Enterobacteriales* order.

Key *mntP* expression platform folding transitions

Figure 6. Co-transcriptional DMS probing of the *mntP* riboswitch during expression platform folding. A-B. (Left panel) TECprobe-VL DMS reactivity matrices of *mntP* riboswitch at pH 8.5. Reactivities shown were normalized with a single, whole-dataset calculated normalization factor. Data are from two independent replicates that were concatenated and analyzed together. (Right panel) Secondary structures of *mntP* expression platform folding intermediates colored by DMS reactivity of the indicated transcript at pH 8.5. **C.** (Top panel) Truncated secondary structure models of *mntP* riboswitch with a stabilized L1/P1.1 (Mn-bound) or disrupted L1/P1.1 (Mn-unbound). L1(orange), L3 (purple), P1.1 (gray), and cross-helix A-minor interaction (pink) are highlighted. (Bottom panel) *X. oryzae* aptamer metal-sensing core (PDB ID: 6N2V) with Mg^{2+} (green sphere) and Mn^{2+} (purple sphere) highlighted. Conformer 1 shows Mn^{2+} fully bound and Conformer 2 shows Mn^{2+} partially bound. A11 (orange) and A48 (purple) are highlighted. **D-E.** (Top panels) Transcript length- and Mn-dependent reactivity changes in A11 (**D**) and A48 (**E**) showing cross-helix A-minor disruption. (Bottom panels) Reactivity difference heatmaps for P1.1/L1 (**D**) and L3 (**E**) at pH 8.5 ± 1 mM Mn. DMS data for 0 mM and 1 mM Mn conditions from Fig. 6A-B and SI Fig. 12A, respectively. Vertical or horizontal dotted lines mark indicated RNA folding transitions.

Key *alx* expression platform folding transitions

E. coli alx riboswitch: folding of the expression platform cotranscriptionally probed, DMS
pH 8.5 0 mM Mn

Mn-dependent differences in *alx* expression platform structure

Figure 7. Co-transcriptional DMS probing of the *alx* riboswitch during expression platform folding. A. (Left panel) TECprobe-VL DMS reactivity matrix of the *alx* expression platform (top) and L1/P1.1 rising stem (bottom) at pH 8.5. Reactivities shown were normalized with a single, whole-dataset calculated normalization factor. Data are from seven independent replicates that were concatenated and analyzed together. (Right panel) Secondary structures of *alx* folding intermediates colored by DMS reactivity of the indicated transcript at pH 8.5. **B.** Truncated secondary structure model of the *alx* aptamer with a disrupted L1/P1.1 (Mn-unbound). L1 (orange), L3 (purple), and P1.1 (gray) are highlighted. **C.** Secondary structure model of *alx* expression platform at transcript length 190 nt –Mn. L1 falling stem (orange) and P1.1 falling stem (gray) paired in P5 are highlighted. Adjacent to nt boxed in pink are plots with their respective transcript-length and Mn-dependent reactivity changes. DMS data for 0 mM (orange) and 1 mM Mn (purple) conditions are from Fig. 7A and SI Fig. 15A, respectively. **D.** Secondary structure model of full-length, translationally inactive *alx* expression platform with same aptamer color scheme as in panels A-B. RBS/start codon (red) and P6 intrinsic terminator stem (pink) are highlighted. **E.** Truncated secondary structure model of the *alx* aptamer with a stabilized L1/P1.1 (Mn-bound) with same aptamer color scheme as in panel A. Cross-helix A-minor motif (pink) is highlighted. **F.** Secondary structure model of *alx* expression platform at transcript length 190 nt +Mn. Adjacent to nt boxed in pink are plots with their respective transcript-length and Mn-dependent reactivity changes. DMS data for 0 mM (orange) and 1 mM Mn (purple) conditions are from Fig. 7A and SI Fig. 15A, respectively. **G.** Secondary structure model of full-length, translationally active *alx* expression platform with same aptamer color scheme as in panel D. RBS/start codon (red) and intrinsic termination site in inactive conformer (pink) are highlighted.

Figure 8. Changes in the *alx* aptamer during expression platform folding. A. (Left panel) TECprobe-VL DMS reactivity matrix of the *alx* riboswitch aptamer during expression platform folding at pH 8.5. Reactivities shown were normalized with a single, whole-dataset calculated normalization factor. Data are from seven independent replicates that were concatenated and analyzed together. (Right panel) Secondary structure model of *alx* aptamer with labeled structural domains. L1 (orange), L3 (purple), P1.1 (gray), and intrinsic termination site (pink) are highlighted. **B-C.** Reactivity difference heatmaps for L1/P1.1 (**B**) and L3 (**C**) at pH 8.5 ± 1 mM Mn. Adjacent to heatmaps are transcript length- and Mn-dependent reactivity changes in L1/P1.1 nt (**B**) and L3 nt (**C**). DMS data for 0 mM and 1 mM Mn conditions from Fig. 8A and SI Fig. 15A, respectively. Vertical or horizontal dotted lines/arrows mark indicated RNA folding transitions. *L. lactis* Mn-sensing aptamer with *E. coli alx* L3 sequence (PDB ID: 6CC1) shown in panel B with A37 (purple), cross-helix A-minor nt: A11 (orange) and G63-C33 (magenta), Mg^{2+} (green sphere), and Cd^{2+} (yellow sphere) highlighted.

1288
Figure 9. Model for *alx* and *mntP* riboregulation in *E. coli*. Cartoon model of *alx* and
mntP riboregulation of their respective genes in *E. coli*. At alkaline pH, the cytosolic [Mn]
increases, resulting in the *alx* riboswitch turning "on" expression of Alx to prevent toxic
accumulation of Mn in the cytosol. The modest increase in cytosolic [Mn] brought about
by alkaline pH is insufficient to shift the *mntP* riboswitch towards its translationally active
conformation, leaving *mntP* expression "off".

1295
1296
1297

1298

1299

1300

1301

1302

1303

1304

1305

1306

Acknowledgements

We thank Dr. Eric Strobel and Dr. Courtney Szyjka for protocols and helpful discussions regarding TECprobe-VL experiments. We also thank members of our research group for their valuable feedback during the course of this study. This work was supported by the National Institutes of Health (NIGMS ESI grant R35GM142785), Burroughs Wellcome Fund CASI (grant no.: 1016945) to T.V.M., UCSD institutional support, and Yinan Wang Memorial Chancellor's Endowed Junior Faculty Fellowship to T.V.M. This material is based upon work supported by the National Science Foundation Graduate Research Fellowship Program under Grant No. DGE-2038238 to D.P. Any opinions, findings, and conclusions or recommendations expressed in this material are those of the author(s) and do not necessarily reflect the views of the National Science Foundation.

Author contributions

C.S. and T.V.M, conceptualization; C.S. and D.P., methodology; C.S. and D.P., investigation; C.S. and D.P., formal analysis; C.S. and D.P., visualization; C.S. and D.P., writing-initial draft; C.S., D.P., and T.V.M, writing-review and editing; T.V.M, supervision; T.V.M., funding acquisition.

Competing Interests

The authors declare no competing interests.