# Charged peptides enriched in aromatic residues decelerate condensate ageing driven by cross-β-sheet formation

Ignacio Sanchez-Burgos[1,2], Andres R. Tejedor[2,3], Alejandro Castro[3],
Alejandro Feito[3], Rosana Collepardo-Guevara[1,2,4*] and Jorge R. Espinosa[1,3*]

[1] Maxwell Centre, Cavendish Laboratory, Department of Physics,
University of Cambridge, J J Thomson Avenue, Cambridge CB3 0HE, United Kingdom.
[2] Yusuf Hamied Department of Chemistry, University of Cambridge,
Lensfield Road, Cambridge CB2 1EW, United Kingdom
[3] Department of Physical-Chemistry, Universidad Complutense de Madrid, Av. Complutense s/n, 28040, Madrid, Spain.
[4] Department of Genetics, University of Cambridge, Cambridge CB2 3EH, United Kingdom.
* = To whom correspondence should be sent.  email: rc597@cam.ac.uk, jorgerene@ucm.es
(Dated: March 27, 2025)

Biomolecular condensates, formed through liquid-liquid phase separation, play wide-ranging roles in cellular compartmentalization and biological processes. However, their transition from a functional liquid-like phase into a solid-like state—usually termed as condensate ageing—represents a hallmark associated with the onset of multiple neurodegenerative diseases. In this study, we design a computational pipeline to explore potential candidates, in the form of small peptides, to regulate ageing kinetics in biomolecular condensates. By combining equilibrium and non-equilibrium simulations of a sequence-dependent residue-resolution force field, we investigate the impact of peptide insertion—with different composition, patterning, and net charge—in the condensate phase diagram and ageing kinetics of archetypal proteins driving condensate ageing: TDP-43 and FUS. We reveal that small peptides composed by a specific balance of aromatic and charged residues can substantially decelerate ageing up to two orders of magnitude. The mechanism is controlled through condensate density reduction induced by peptide self-repulsive electrostatic interactions that specifically target protein regions prone to form cross-β-sheet fibrils. Our work proposes an efficient computational framework to rapidly scan the impact of small molecule insertion in condensate ageing and develop novel pathways for controlling phase transitions relevant to disease prevention.

## I. INTRODUCTION

Cell compartmentalization is a cornerstone of cellular organization, enabling spatiotemporal regulation of biochemical reactions with remarkable precision [1]. While membrane-bound organelles have been recognized as the primary means of achieving compartmentalization, transformative experiments have revealed liquid-liquid phase separation (LLPS) as a highly dynamic and versatile mechanism for organizing cellular content [2–4]. Through LLPS, biomolecular condensates are formed via multivalent and dynamic interactions among proteins, nucleic acids, and other macromolecules, creating membraneless phase-separated compartments with unique biochemical properties [1, 5–9]. For instance, the formation of the nucleoli—where ribosomal RNA is synthesized and processed—is driven by LLPS of key constituents including fibrillarin and nucleophosmin that facilitate the assembly of ribosomal subunits [10–15]. Moreover, the emergence of stress granules, which appear in response to cellular stress, concentrating mRNAs and proteins [16, 17], or the nucleation of P granules, which play essential roles in germ cell development as found in *Caenorhabditis elegans* [18, 19], represent fundamental examples of intracellular organization via phase-separation. Importantly, condensates can rapidly assemble and disassemble, which provides a unique flexible mechanism to regulate essential cellular processes such as gene expression, stress responses, and signal transduction among others [20–22].

Nevertheless, dysregulation of intracellular LLPS has been implicated in the onset of multiple neurodegenerative disorders [23–25] including amyotrophic lateral sclerosis (ALS), Alzheimer's disease, Parkinson's disease, and certain types of dementia—where condensate phase separation eventually leads to the formation of aberrant solid-like aggregates [26–29]. In ALS, mutations in proteins such as the Transactive response DNA binding protein of 43 kDa (TDP-43) and Fused in Sarcoma (FUS) disrupt their functional phase behaviour, resulting in the formation of toxic solid-like aggregates that compromise cellular function [30–36]. Similarly, in frontotemporal dementia, the misfolding and aggregation of Tau protein and other disease-associated proteins are linked to altered LLPS [37–39], contributing to neuronal dysfunction and cell death. Beyond neurodegenerative diseases, LLPS has also been associated with cancer, where aberrant condensate formation influences oncogenic signalling pathways and promotes tumorigenesis [40–45]. The progressive transition of functional liquid-like biomolecular condensates into pathological solid-like states (e.g., ageing) has been suggested to be originated by the gradual accumulation of inter-protein β-sheets within the condensates [39, 46–48]. These inter-protein structures can be formed over time when biomolecules are concentrated to high levels, as illustrated in Fig. 1(a), and are capable of transforming transient liquid-like interactions into long-lived highly energetic cross-links characteristic of kinetically arrested states such as glasses or gels. The progressive accumulation

of cross-β-sheet structures over time has explained the gradual decrease in molecular mobility that numerous biomolecular condensates experience over time [7, 49–54]. Therefore, understanding the molecular underpinnings of LLPS in these disorders not only elucidates a fundamental mechanism of pathogenesis, but also underscores the urgent need to develop strategies to regulate LLPS due to its critical role in health and disease.

To address this challenge, a potential therapeutic strategy has involved the development of small molecules or peptides that specifically modulate LLPS [55–57]. Multiple studies have focused on using different molecules to dissolve already formed condensates [57–61]. Examples include dissolution of DDX3 [62], PARP1 [63], and PI3K [64] stress granules with the use of synthetic molecules, as well as the dissolution of Tau [65] and p53 [66] condensates with the addition of small peptides. Another considerable body of literature focuses on the inhibition or delay in the condensates ageing, rather than dissolving them. Particularly, delaying or inhibiting aggregation in Tau [67], Aβ42 [68–70], TDP-43 [56, 71] and α-synuclein [72–74] condensates has been attempted in presence of LLPS modulating small organic molecules. Nevertheless, while short peptides offer a promising strategy for therapeutic intervention due to their specificity and potential for cell-penetration [75], the precise protein-peptide interactions required to control the aggregation and ageing pathways remains largely unknown [18, 60, 76].

Computational approaches represent a powerful tool for expediting the screening and optimization of suitable peptide sequences for this goal, and contribute to the development of effective therapeutics against condensate-related disorders such as neurodegenerative diseases [77]. Here, we perform non-equilibrium Molecular Dynamics (MD) simulations at the coarse-grained residue-resolution level [78] to investigate the inhibition of condensate ageing through the insertion of small peptides at relatively low peptide concentrations. As a proof-of-concept, we focus on the low-complexity domains of FUS (referred to as FUS LCD) and TDP-43 (referred to as TDP-43 LCD) due to their well-established relevance in condensate misregulation and ageing driven by inter-protein β-sheet accumulation [30–36]. In Fig. 1(b), we highlight the charged and aromatic residues within both protein sequences, as well as the low-complexity aromatic-rich kinked segments (LARKS) that have been found capable of forming inter-protein β-sheet structures [46, 79]. By means of MD simulations, we perform a high-resolution analysis of peptide-protein interactions over time, capturing detailed dynamics and structural changes within condensates that experimental techniques are unable to detect [5, 80–92]. We employ the residue-level resolution CALVADOS2 model [78] to explicitly represent each amino acid, thereby capturing essential sequence-dependent effects in condensate phase behaviour while enabling us to perform simulations over longer timescales and larger systems than would be possible with fully atomistic models [93–95]. The CALVADOS2 model [78] accurately predicts the conformational properties and propensities to undergo LLPS for diverse sequences and solution conditions [78, 90] while providing fair predictions for condensate material properties despite being far beyond its parametrization scope [90].

We first investigate how the stability of FUS LCD and TDP-43 LCD condensates varies as a function of the concentration of the inserted peptides, and later we examine how the ageing rate is modulated by their recruitment into the condensate. Our goal extends from designing promising peptides that decelerate ageing to uncover the underlying mechanisms and interactions behind their inhibitory effect, which can further guide the design of functional sequences for inhibiting condensate ageing. To achieve this, we analyse inter-protein contact frequency maps, interrogate how condensate density and stability regulate ageing kinetics, and explore potential peptide-protein specific interactions that influence the hardening process. Overall, our work proposes a computational scheme to rapidly scan potential inhibitors of liquid-to-solid condensate transitions and provides insights into the underlying molecular interactions controlling the phase behaviour and time-dependent material properties of protein condensates.

## II. RESULTS AND DISCUSSION

### A. A coarse-grained modelling approach for designing peptide sequences controlling condensate stability and ageing

Our method involves a multistep approach in which we first analyze how the phase diagram of protein condensates is influenced by the amino acid composition of the designed peptides to preliminarily identify candidates that are capable of slowing down condensate ageing. It has been shown both experimentally [51, 52] and computationally [50, 96, 97] that an increase in the density of the condensates enhances local high-density fluctuations of protein LARKS, which in turn, promote higher rates of cross-β-sheet transitions. In contrast, recruitment of partner molecules (e.g., IDRs, RNAs, or other biomolecules) which decrease condensate concentration reduces the probability of cross-β-sheet fibril formation [49]. To predict the variation in condensate density as a function of the concentration of the inserted peptides, we employ the CALVADOS2 residue-resolution model [78]. In this model, each amino acid across the sequence is represented by a single bead with its own chemical identity. The low-complexity domains of both FUS and TDP-43 are considered as fully flexible polymers in which subsequent amino acids are connected by harmonic bonds. The different residue-residue interactions in the model consist of a combination of electrostatic and hydrophobic interactions [78, 98] implemented through a Coulomb/Debye-Hückel potential and an Ashbaugh/Hatch potential, respectively (further technical details of the

Figure 1: (a) Schematic representation of the accumulation of inter-protein β-sheets within a bulk condensate simulation. The zoomed images depict the nature of the initially intrinsically disordered protein regions (left) and the cross-β-sheet structures formed over time (right). (b) TDP-43 LCD and FUS LCD sequence representation, highlighting the charged and aromatic residues, as well as the low-complexity aromatic-rich kinked segments (LARKS) as indicated in the legend. (c) Storage (G') and shear loss (G") moduli as a function of frequency (ω) for non-aged (top) and aged (bottom) FUS LCD condensates. (d) Time-evolution of cross-β-sheet concentration for FUS LCD and TDP-43 LCD condensates. The continuous lines depict simulation data while the dashed lines represent fits to the data according to a secondary nucleation dominated mechanism (see Section II C for details on these fits). (e) Simulation of a FUS LCD condensate in bulk conditions containing small peptides that hinder the nucleation of cross-β-sheet structures.

model can be found in the Supplementary Material (SM) Section SI).

Once determined the condensate density (or concentration) variation upon peptide insertion through short NpT Molecular Dynamics simulations (further details on these simulations are provided in Section II B), we perform non-equilibrium simulations—using our ageing dynamic algorithm [50, 99]—both in the presence and absence of peptides, in which condensates can gradually transform their material properties due to the emergence of inter-protein β-sheet structures. Within these simulations, the intermolecular interactions between the amino acids that are part of the LARKS (as those highlighted in black in Fig. 1(b)) can be transformed over time, mimicking the disorder-to-order transitions taking place in these protein regions [46]. Further details on this algorithm, which has been parametrized based on all-atom calculations of binding free energy differences between disordered and structured LARKS in both FUS [99] and TDP-43 [96] are detailed in Sections II C and SII of the SM. In close

agreement with experimental findings, our results show that the formation of inter-protein β-sheets progressively modifies the viscoelastic properties of the condensates [7, 49–53]. We characterize this change on the condensate material properties in Fig. 1(c) by measuring the storage (G') and loss (G") moduli as a function of frequency (ω) for FUS LCD condensates before and after the formation of cross-β-sheet structures. In the non-aged condensate (Top Panel), the system displays liquid-like behaviour, characterized by the fact that the magnitude of G" is higher than G' at low frequencies [97]. However, once the condensate accumulates cross-β-sheet structures, and establishes long-lived inter-protein interactions (e.g., of 20-45 $k_B T$ as found for FUS and TDP-43 LARKS in Refs. [50, 96]), elasticity dominates over flow, and therefore, the system exhibits a solid-like behaviour (G' > G" at lower frequencies). Such behaviour, experimentally reported for both FUS [52, 54, 100] and TDP-43 [79, 101] among many other proteins [102], emerges from a percolated network of cross-β-sheets assemblies which restrains the protein mobility and impedes its diffusion across the condensate

[50].

In our scheme, we analyse the kinetics of this process to determine how peptides modify the nucleation time and growth rate at which inter-protein β-sheet structures are formed. In Fig. 1(d), we show the time evolution of the relative concentration of LARKS forming cross-β-sheets in pure condensates of both FUS LCD and TDP-43 LCD. These curves allow direct determination of the different parameters that serve as a metric for assessing the ageing rate. By performing multiple independent simulations, we determine the characteristic timescales of cross-β-sheet formation in each protein condensate. These parameters (detailed in Section IIC) determine whether there is an effective deceleration of the ageing kinetics as a function of the composition, patterning, and concentration of the inserted peptides with respect to the pure protein condensate. In what follows we probe this method for protein condensates in presence of small peptides, with the aim that their recruitment within the condensate frustrates the protein high-density fluctuations leading to cross-β-sheet transitions. We provide a schematic explanation in Fig. 1(e) of how small peptides (coloured in red) are intertwined within the protein liquid network (coloured in navy blue), and potentially inhibit the formation of inter-protein β-sheets, either by lowering the overall protein density, or by establishing selective interactions with aggregation-prone protein regions.

### B. Phase boundaries of FUS LCD and TDP-43 LCD condensates in presence of small peptide modulators

To prevent the formation of inter-protein β-sheet structures without significantly altering the stability and material properties of the condensates, peptides must only hinder local LARKS high-density fluctuations (driving cross-β-sheet formation) while preserving the central interactions between protein domains that dictate the saturation concentration and critical solution temperature of the condensate. For that purpose, the concentration of peptides within the condensates shall be kept moderately low. We begin by estimating the phase diagram of pure FUS LCD condensates (Fig. 2(a)) using Direct Coexistence (DC) simulations [5]. Through the combination of DC simulations and the laws of rectilinear diameters and critical exponents [103], we determine the coexistence densities and critical solution temperature for FUS LCD pure condensates (Fig. 2(a); black filled and empty symbol, respectively). We define the critical solution temperature for LLPS as our metric of biomolecular condensate stability [90] when comparing between different protein/peptide systems. Since imposing a given concentration of peptides within a protein condensate is hardly attainable through DC simulations [83], we perform NpT (constant number of particles (N), pressure (p) and Temperature (T)) simulations for protein/peptide mixtures, where the critical solution temperature ($T_c$) is estimated to be between the highest temperature at which the condensate is stable at

p=0 atm and the lowest one at which the condensed phase evolves into the diluted phase (Fig. 2(a), bottom; further details on this method are provided in Section SIII of the SM). This approach—validated in Ref. [83]—additionally allows to evaluate the condensate density as a function of temperature for different peptide sequences and/or protein/peptide stoichiometries (Fig. 2(a); red and green symbols). Our simulations are performed at physiological NaCl concentration (150 mM), which is imposed through the Debye length of the screened Coulombic potential [78, 104] (see SM section SI for further details). Strikingly, we find that increasing concentrations of short polyArginine peptides of 24 residues ($R_{24}$) enhance the stability of FUS LCD condensates despite their highly self-repulsive character. On the other hand, addition of $(SYSYKRKK)_3$ peptides, which are also positively charged, reduces condensate density and $T_c$ upon its recruitment within the condensate (Fig. 2(a)). We attribute such opposite behaviour to the strong cation-π interactions that $R_{24}$ peptides are capable of establishing with the aromatic residues of FUS LCD sequence at low concentrations; in contrast to the less energetic contacts that lysines (K) and serines (S) can form with FUS LCD.

To further investigate the impact of peptide regulation on FUS LCD phase separation propensity, we perform a systematic study of different representative sequences that present specific variations of patterning and composition (Fig. 2(b)). We note that there are millions of possible peptide combinations that may be used to modulate biomolecular condensate' phase behaviour; however, we design 15 different sequences of 24 residues per chain enriched in specific amino acids that frequently appear in RNA-binding proteins (RBPs) such as FUS or TDP-43. This strategy enables us to maintain as much as possible the protein condensate composition of the pure protein system. The chosen residues are serine (S), tyrosine (Y), and glycine (G) which ensure substantial binding affinity to the aromatic-rich regions in FUS LCD and TDP-43 LCD sequences due to π-π contacts mediated by tyrosine. Moreover, we also introduce charged residues across the peptide sequences: lysine (K), arginine (R), and glutamic acid (E) to generate peptide-peptide self-repulsion that partially destabilize LARKS-LARKS high-density fluctuations leading to cross-β-sheet clustering. The choice of including both R and K when designing positively charged peptides is their different roles in phase separation. Arginine is a context-dependent sticker [105] due to its capacity to form strong cation-π interactions, whereas K mostly acts as an spacer only favouring electrostatically oppositely charged contacts [106, 107]. In contrast, both E and aspartic acid (D) have been shown to play comparable roles when establishing LLPS intermolecular interactions [107], and thus for simplicity, we only consider E when designing negatively charged peptides.

In Fig. 2(b), we show how different peptide sequences and concentrations decisively regulate $T_c$ in FUS

Figure 2: (a) Normalized temperature (where $T_{c,FUS\ LCD}$=322 K) *vs.* density phase diagram for FUS LCD in presence and absence of two peptides modulating its phase behaviour. Increasing concentrations of $R_{24}$ augment the condensate density and critical solution temperature, while $(SYSYKRKK)_3$ induces the opposite behaviour. Filled points represent direct measurements under coexistence conditions, while the empty point depicts the critical temperature for LLPS. The continuous lines are shown as a guide for the eye. At the bottom we represent typical snapshots of the NpT simulations employed to compute the phase diagram in presence of small peptides, where at temperatures below the critical one, the condensed phase remains stable at p=0 bar, while above $T_c$, the system forms a low density phase. (b) Normalized critical temperature of FUS LCD mixtures with different peptide sequences (as indicated in the legend) as a function of the peptide concentration. (c) Normalized critical temperature (where $T_{c,TDP-43\ LCD}$=308 K) of TDP-43 LCD mixtures with different peptides as a function of the peptide concentration. The green and red shaded areas in panels (b) and (c) represent an increase and decrease in $T_c$, respectively.

LCD/peptide mixtures as compared to pure condensates ($T_{c,\ FUS\ LCD}$) at constant physiological salt concentration. The first two peptides we explore are the tryptophan (W)-tyrosine (Y) 12-repeat sequence $((WY)_{12})$ and $R_{24}$. $(WY)_{12}$ is a purely aromatic sequence formed of 'sticker' residues [108–110] which are capable of forming strong π-π interactions with FUS LCD. $R_{24}$ only contains arginine, and displays high binding affinity to FUS LCD aromatic residues such as Y and phenylalanine (F) through cation-π interactions, as well as moderate repulsive interactions with other positively charged residues [107]. Interestingly, despite being $(WY)_{12}$ a purely self-attractive peptide and $R_{24}$ a self-repulsive peptide, both of them monotonically increase the critical solution temperature for phase separation across the concentration range studied here (Fig. 2(b); brown and red symbols). Such increase in $T_c$ implicitly produces an increase in the density of the condensate (Fig. 1(a)) which implies higher local density fluctuations in FUS LCD LARKS with respect to the pure condensate, thus

potentially promoting more frequent cross-β-sheet transitions and faster ageing. The LLPS enhancement observed in presence of $R_{24}$ is consistent with experimental findings [111] reporting an increased propensity of RNA-binding proteins such as TDP-43 for condensation and insolubility in presence of arginine-rich peptides (i.e., proline-arginine dipeptide repeats such as $(PR)_{25}$). Therefore, short peptides aiming to partially reduce LARKS high density fluctuations must contain less 'sticker' amino acids so the overall condensate density is moderately reduced while still enabling LLPS (Fig. 2(a); red symbols). Similarly to these two peptides, we test $(SYGSYYGS)_3$ in which charged residues are excluded (cyan curve). Although this sequence contains fewer aromatic residues than $(WY)_{12}$, still increases the critical solution temperature for phase separation due to the absence of self-repulsive interactions that contribute to reducing condensate density, hence it will not be further considered as a potential ageing inhibitor.

We next investigate the effect of a combination of 'sticker' and 'spacer' amino acids. For this purpose, $(PHSY)_6$ peptides are added to FUS LCD condensates, where proline (P), serine (S) and histidine (H), act as spacers. Remarkably, we find how this sequence is able to lower the critical temperature in a marginal extent up to relatively high concentrations ($>0.075$ mg peptide/mg FUS LCD). Nevertheless, since a high concentration of $(PHSY)_6$ is required to slightly reduce the density of the condensate, this peptide is not ideal to preserve: (1) the material properties of FUS LCD condensates; and (2) the overall condensate composition, which in turn impacts other physicochemical parameters such as the surface tension (Fig. 2(b)). We also test the influence of partially charged peptides on the stability of the condensates. We first insert strong negatively charged peptides to destabilize FUS LCD homotypic intermolecular interactions near the LARKS regions via electrostatic-driven binding affinity between K and E residues from FUS LCD and the peptides. $E_{24}$ (only glutamic acid, not shown in Fig. 2(b)) peptides destabilize the condensate by at least 0.1 $T_c/T_{c,FUS\ LCD}$ at the lowest concentration measured, making it not suitable for this purpose since $T_c/T_{c,FUS\ LCD} = 0.1$ implies a reduction of $\sim 30K$ in the critical solution temperature.

We then design four additional peptide sequences with lower content of glutamic acid ($(SYSYEEEE)_3$, $(SYGSYEGS)_3$, $(SYGESYGSYGEG)_2$, and $(SYGSYGSYEGEG)_2$), each of them with a different patterning and composition of negatively charged residues (E), stickers (Y), and spacers (G, S). We find that $(SYSYEEEE)_3$ still destabilizes FUS LCD condensates to a great extent by lowering the critical solution temperature at significantly low concentrations ($>0.05$ $T_c/T_{c,FUS\ LCD}$ with respect to pure FUS LCD condensates at mass ratios $< 0.05$ mg peptide/mg FUS LCD). While certainly a decrease in the critical temperature and density of the condensates is intended, the concentration at which that effect is achieved is too low, and therefore, we do not consider $(SYSYEEEE)_3$ as a potential cross-β-sheet inhibitor. Its strong capacity to disrupt the FUS LCD liquid network connectivity is not optimal if its relative concentration within the condensate is not at least 0.05 mg peptide/mg FUS LCD. This is because the concentration of the inserted peptides must be high enough to interact with a significant number of protein LARKS to effectively disrupt high density fluctuations over time leading to cross-β-sheet transitions. By testing the three other sequences with a lower concentration of E ($(SYGSYEGS)_3$, $(SYGESYGSYGEG)_2$ and $(SYGSYGSYEGEG)_2$), we find a moderate destabilizing effect of 5-10 K in $T_c$ within a concentration range of 0.05 to 0.1 mg peptide/mg FUS LCD. Such destabilizing driving force meets a reasonable compromise between a moderate reduction in condensate density and stability while ensuring a sufficient concentration of peptides to target FUS LCD-prone aggregation domains. We note that a similar behaviour is expected to be achieved if aspartic acid instead of E would have been sequenced across the

peptides due to their physicochemical resemblance at modulating protein LLPS as reported in Refs. [78, 107, 112, 113].

Lastly, we evaluate how the presence of positively charged peptides impacts the critical solution temperature of FUS LCD. We design two sequences of peptides containing a common segment SYSY—with high associative binding for LARKS and adjacent regions [96]—and different positively charged domains (KRKS and KRKK, respectively) to display a certain degree of electrostatic self-repulsion. These peptides are: $(SYSYKRKS)_3$ and $(SYSYKRKK)_3$. Examining FUS LCD phase behaviour as a function of the peptide concentration shows that only $(SYSYKRKK)_3$ meets the criterion of inducing a moderate destabilization effect in the critical solution temperature ($\sim 0.04$ $T_c/T_{c,FUS\ LCD}$) at low mass ratios (i.e., $\sim 0.075$). Such difference in the behaviour between both peptides stems from the increased abundance of lysine in $(SYSYKRKK)_3$, which enhances electrostatic self-repulsion. Consequently, this reduces condensate density and stability, akin to the impact experimentally [114, 115] and computationally [83, 116] reported when high concentrations of poly-Uridine (negatively charged in this case) are recruited into RNA-binding protein condensates.

Furthermore, we have conducted a similar analysis involving TDP-43 LCD condensates (Fig. 2(c)). We probe the stability of the condensates in the presence of the peptide sequences found to be successful in reducing the density and stability of FUS LCD condensates: $((SYSYKRKS)_3$, $(SYSYKRKK)_3$, $(SYGSYEGS)_3$, $(SYGESYGSYGEG)_2$ and $(SYGSYGSYEGEG)_2$). As reported in Fig. 2(c), the five sequences achieve a reasonable compromise between a moderate reduction in condensate density and stability at a sufficient concentration of peptides capable of targeting multiple TDP-43 proteins at once. In addition, we test one more peptide, $(SYGKSYGSYGKG)_2$, in which the content of lysine has been reduced. Since TDP-43 LCD is slightly positively charged (e.g., $+2e$ across its sequence), this peptide destabilizes the condensate within the desired range of concentration (Fig. 2(c); red symbols) through a combination of homotypic self-repulsion and a subtle balance of heterotypic interactions: electrostatic repulsion (mediated by positively charged residues) and π-π attraction (induced by aromatic amino acids). The same destabilizing effect mediated via electrostatic repulsion is thought to be responsible for the RNA-driven re-entrant behaviour of RBP condensate stability [114–117]. Finally, as in FUS LCD condensates, we also evaluate the impact of recruiting non-charged peptides enriched in sticker residues: $((SYGSYYGS)_3$ and $(SYGYSYGSYGYG)_2$). Consistent with our findings for FUS LCD, both peptide sequences monotonically enhance condensate stability across the studied range of concentrations, therefore they are unable to prevent LARKS high-density fluctuations driving condensate hardening.

Overall, from our simulations of both FUS LCD and

TDP-43 LCD mixtures we conclude that only self-repulsive charged peptides with a certain degree of aromatic residues and neutral spacers enable mild variations on condensate stability while remaining at low stoichiometries (e.g., 1 peptide per every 10-20 proteins forming the condensate). Recent computational work [118] has shown how electrostatic and hydrophobic residues play a crucial role in the regulation of condensate stability through non-specific interactions, in agreement with our findings for FUS LCD and TDP-43 LCD. Moreover, computational studies have also predicted that the inclusion of arginine-rich peptides enhances the stability of protein condensates [119, 120] (at least for a certain concentration range), as found in our FUS LCD/$R_{24}$ mixtures (Fig. 2(b)). This behaviour is influenced by the cation-π interactions which are established between peptides and proteins, as reported through atomistic simulations in Refs. [119, 121]. Furthermore, we note that another recent computational study has also predicted the suppression of hnRNPA1 phase separation after the addition of glutamic acid enriched peptides [122], which is in close agreement with our observations (Fig. 2(b)-(c)) and previous simulations studies of hnRNPA1 in the presence of polyUridine [50, 107, 116]. However, while the factors which modulate the stability of protein condensates in presence of small molecules and peptides is beginning to be understood, their impact on the condensate time-dependent material properties remains unclear.

### C. Charged peptides enriched in aromatic residues decelerate condensate ageing driven by inter-protein structural transitions

To elucidate the impact of peptide inclusion in condensate material properties and ageing, we now perform non-equilibrium MD simulations incorporating our ageing algorithm, which accounts for the description of structural and energetic transformation of disordered LARKS into inter-protein β-sheets [50, 96]. As input for the free-energy binding difference between the structured vs. disordered LARKS, we use atomistic potential-of-mean-force (PMF) calculations from Refs. [99] and [96] for FUS LCD and TDP-43 LCD, respectively. In these studies, PMF calculations for the dissociation of the inter-protein β-sheet clusters were evaluated based on the structures from Refs. [46, 79], resolved via Cryo-Electron Microscopy (Cryo-EM). PMF calculations were performed using an atom-level force field [124], considering both structured conformations of Refs. [46, 79] and the intrinsically disordered ensemble predicted by the all-atom model [124].

Our algorithm continuously evaluates LARKS high-density fluctuations across the simulation using a distance criterion based on a local order parameter developed by us [99]. When at least four LARKS from different protein replicas meet specific proximity conditions (i.e., within a cut-off distance consistent with the free energy minima of the PMF calculations), the interactions between these segments are adjusted according to the PMF structured binding

free energy of the given sequence domain. These new interactions become significantly stronger (i.e., from 3-6 $k_BT$ per LARKS when being disordered to 30-50 $k_BT$ upon the structural transition [96]) mimicking the emergence of an inter-protein β-sheet structure (Fig. 1(a)). While LARKS cross-β-sheets are not usually thermoresistant, as amyloid fibrils [52, 125], these structures are still highly stable as experimentally [39, 46–49] and computationally [50, 99, 126] found. Moreover, besides the energetic implications of the structural change, our algorithm also imposes an angular potential between consecutive amino acids to reflect the increased rigidity of the newly formed structures, thus replicating the conformational change across the involved segment. For further details on the local order parameter, cut-off distances, magnitude of the high-density fluctuations, and binding energies associated to the structural transitions, please see Section SII of SM.

We perform non-equilibrium simulations in bulk conditions (at the corresponding pre-ageing condensate density) for both FUS LCD and TDP-43 LCD condensates in the presence of the previously selected peptides at a normalized temperature of $T/T_c=0.95$ (293 and 305 K for TDP-43 LCD and FUS LCD, respectively). The inserted peptides are at the maximum concentration which lowers the critical temperature for phase separation in ∼0.03 $T/T_c$ (the absolute numerical value of each concentration is specific for each system and is reported in the SM section SV). The inter-protein β-sheet concentration as a function of time for different FUS LCD trajectories in absence vs. presence of different peptide sequences is shown in Fig. 3(a). Since the nucleation of cross-β-sheet stacks is a stochastic process [127], we carry out 6 independent trajectories of each system with different initial velocity seeds. The inter-protein β-sheet concentration is normalized by the maximum number of cross-β-sheets formed (at the plateau) once the protein mobility within the condensate becomes fully arrested (which is approximately at 90% of the total possible β-sheets formed). While pure FUS LCD condensates (light blue curve) rapidly develop inter-protein structural transitions over time, condensates including peptides such as (SYGSYEGS)$_3$ (brown curve) display considerably longer lag times (∼100 times slower) before the formation of β-sheets assemblies. The ageing curves for all the studied systems containing the different independent trajectories per system are reported in Fig. S1. According to our simulation setup, and based on the obtained trends, the process follows a nucleation-elongation mechanism [127]. The nucleation stage is the rate-limiting step of the process as compared to the cross-β-sheet growth. Notably, until the first structured assembly is formed, proteins display liquid-like behaviour characterized by transient and weak interactions. However, once a stable nucleus emerges, the elongation step rapidly evolves, giving rise to a sharp increase in the cross-β-sheet concentration (Fig. 3(a)). Remarkably, we find that fitting our results to a secondary nucleation-dominated kinetic model [128] (dashed curves fitted with Amylofit software [123]) matches the

Figure 3: (a) Relative cross-β-sheet concentration—determined from the amount of inter-protein β-sheets formed—as a function of time for FUS LCD condensates in absence vs. presence of different peptide sequences as indicated in the legend. Filled points represent measured β-sheet concentration from non-equilibrium simulations, while dashed lines depict fits to a secondary nucleation dominated kinetic model performed with Amylofit [123]. (b) Half-time ($t_{1/2}$) for the different FUS LCD condensate mixtures studied. $t_{1/2}$ is defined as the point where the relative β-sheet concentration value is halfway between the initial value and final plateau value. In the box plot representations, the average value is represented with a cross symbol, while the box boundaries are obtained from the 25 and 75 percentile values from our data range. (c) Relative cross-β-sheet concentration as a function of time for TDP-43 LCD condensates in absence vs. presence of different peptide sequences as indicated in the legend. Filled points and dashed curves represent the same quantities as in panel (a). (d) Half-time for the different TDP-43 LCD mixtures studied.

simulation data. Hence, our simulations suggest that both cross-β-sheet growth and secondary nucleation represent two fundamental mechanisms by which kinetically arrested states are reached.

To further characterize the ageing kinetics, we evaluate the time required to reach half of the cross-β-sheets formed in the condensate (hereafter referred to as the half-time, or $t_{1/2}$). $t_{1/2}$ is a standard metric for quantifying the kinetics of aggregation from condensates and/or supersaturated protein solutions [69, 72, 129–131]. In Fig. 3(b), we report the box plots for the average half-times (from independent trajectories) obtained for the different peptide/FUS LCD mixtures. The average value is indicated with a cross symbol, the boxes represent the 25-75 quartile range, and the error bars indicate the highest and lowest values. We conclude that ageing kinetics are substantially decelerated in the presence of most of the chosen peptides selected

based on the density-stability criterion discussed in Section II B. From all the tested sequences, only (SYSYKRKK)$_3$ is not capable of significantly deferring the nucleation time. On the other hand, (SYGSYEGS)$_3$, (SYGESYGSYGEG)$_2$, (SYGSYGSYEGEG)$_2$, and (SYSYKRKS)$_3$ (in a milder extent) can substantially disrupt LARKS high-density fluctuations triggering FUS LCD progressive solidification. Importantly, we find that negatively charged peptides perform significantly better than positively charged ones in decreasing $t_{1/2}$. The FUS LCD sequence only contains 2 negatively charged residues out of 163 amino acids, nevertheless, such additional degree of destabilization through electrostatic repulsive forces that negatively charged peptides induce in FUS LCD condensates contributes to further hindering condensate ageing. We also find that including peptides with the same amino acid composition but different sequence patterns, such as (SYGESYGSYGEG)$_2$ and (SYGSYGSYEGEG)$_2$, appears to play a minor role in

modulating the onset of cross-β-sheet formation. Similarly, the change in E abundance between (SYGSYEGS)$_3$ and both (SYGESYGSYGEG)$_2$ and (SYGSYGSYEGEG)$_2$ does not show a major impact in varying the ageing propensity as long as the balance between peptide self-repulsion and protein binding affinity for being recruited into the condensate is maintained. In Section II D, we will perform a molecular contact analysis of the condensate mixtures to uncover which major intermolecular interactions are responsible for the deceleration of cross-β-sheet formation.

From the box plot analysis, we also reveal that ageing kinetics is governed by the nucleation lag time, which initially prevents the rigidification cascade from occurring. Since the nucleation step is a stochastic event that takes place with a given probability (depending on the condensate composition, density, and thermodynamic conditions), it is expected that the median (horizontal line) is lower than the average (black cross), and the distribution is wider for a lower probability of nucleation—as it happens in presence of negatively charged peptides. The t$_{1/2}$ data conforming the box plots shown in Fig. 3(b) follow a Poisson distribution [132], which also reaffirms the role of the nucleation stage as the crucial limiting step in condensate ageing driven by cross-β-sheet ordering. We note that apart from the ageing half-time, the nucleation lag time (defined as the time at which the first inter-protein β-sheet is formed), and log($k_n$) (where $k_n$ is the primary nucleation rate constant [123]) can also be used to quantify the extent to which the ageing rate is altered by the presence of peptides. These additional analyses showing how the nucleation time, log($k_n$) and t$_{1/2}$ provide equivalent conclusions for the ageing kinetics of these systems are detailed in Section SVI of the SM (Figs. S3(a) and S4(a)).

Next, we perform non-equilibrium simulations of TDP-43 LCD condensates in the absence and presence of the peptides that partially destabilized LLPS in Fig. 2(c). The time-evolution of the cross-β-sheet concentration for multiple TDP-43 LCD mixtures is shown in Fig. 3(c). We consistently observe an increase in the nucleation time when small peptides are added compared to the pure protein condensate. However, the growth of cross-β-sheet clusters occurs on a similar rate across all systems. From these curves, along with the collective analysis of the different independent trajectories for each system (Fig. S2), we conclude that the ageing kinetics is limited by the nucleation and dominated by the growth, as found for FUS LCD systems in Figs. 3(a-b). To quantify the effectiveness of the selected peptides in slowing down ageing, we calculate the ageing half-time. In Fig. 3(d), t$_{1/2}$ is displayed in box plots for all systems. Impressively, all chosen peptides significantly decelerate ageing; with multiple trajectories showing half-times up to two orders of magnitude larger than that of pure TDP-43 LCD condensates, particularly when incorporating the (SYGSYEGS)$_3$ and (SYGESYGSYGEG)$_2$ sequences. In contrast to FUS LCD condensates, both positively and negatively charged peptides delay the nucle-

ation stage similarly: the box plots for t$_{1/2}$ in the presence of (SYGESYGSYGEG)$_2$ and (SYGKSYGSYGKG)$_2$— where E and K residues are swapped—overlap within the uncertainty. Nevertheless, when arginine residues instead of lysines are included in the peptide, as in (SYSYKRKK)$_3$, the ability to hinder inter-protein structural transitions is significantly reduced. We also evaluate the role of residue patterning by inserting (SYGESYGSYGEG)$_2$ and (SYGSYGSYEGEG)$_2$ peptides. In contrast to FUS LCD condensates, the (SYGESYGSYGEG)$_2$ sequence enables longer lag times before cross-β-sheet fibril formation than (SYGSYGSYEGEG)$_2$. Hence, pattern variations are likely to be significant depending on the specific sequence features of the protein undergoing condensate ageing. Furthermore, the degree of clustering of aromatic vs. charged residues in the peptides can remarkably influence their impact on ageing kinetics, as shown in Fig. 3(d). We also evaluate the primary nucleation rate constant ($k_n$) and the average nucleation lag time in Figs. S3(b) and S4(b). As for FUS LCD mixtures, both analysis yield the same conclusions as through the analysis of t$_{1/2}$: the addition of partially— positively or negatively—charged peptides enriched in aromatic residues effectively delay the emergence of cross-β-sheet structures (Fig. 3(d)).

### D. Molecular factors governing ageing deceleration via peptide insertion

To decipher the mechanisms by which peptide insertion delays condensate ageing, we compute the intermolecular contact frequency maps for the different systems studied in Section II C before inter-protein β-sheets are formed. We consider a residue-residue contact to effectively contribute to LLPS when two amino acids are found at a distance below 1.2σ$_{ij}$, being σ$_{ij}$ the average molecular diameter of the two residues $i$ and $j$, and 1.2 a slightly larger distance to the minimum in the interaction potential (∼1.12σ$_{ij}$) between the $i$th and $j$th amino acids (further details on these calculations are described in Section SVII of the SM). In Fig. 4(a)-(b), we show the contact frequency of all the different residues for FUS LCD condensates in the absence (a) vs. presence of (SYGSYEGS)$_3$ peptides (b). From the comparison of both maps, we find that the overall intermolecular contact frequency between FUS LCD proteins is marginally altered upon peptide insertion. However, some of the residues that establish more prominent interactions, such as the subdomains that range between the 36th to the 40th residue, from the 53rd to the 58th, and from the 77th to the 82nd, which correspond to LARKS and LARKS adjacent segments, reduce their contact frequency when (SYGSYEGS)$_3$ peptides are recruited. In Fig. S5 we report the contact frequency maps for the rest of inserted peptides, which are relatively similar to that shown in Fig. 4(b), i.e., the overall inter-protein contact valency remains similar to that in pure FUS LCD condensates, but contacts between LARKS and LARKS adjacent regions are reduced.

Figure 4: Intermolecular frequency contact maps for FUS LCD homotypic contacts in pure FUS LCD condensates (a) and FUS LCD/(SYGSYEGS)₃ mixtures (b). The pairwise residue-residue contact frequency is expressed in percentage in all panels. (c) Intermolecular frequency contact maps of the different inserted peptides with the FUS LCD sequence in the condensate. The contacts are calculated between all residues of the inserted peptide and the FUS LCD sequence. Intermolecular frequency contact maps for TDP-43 LCD homotypic contacts in pure TDP-43 LCD condensates (d) and TDP-43 LCD/(SYGSYEGS)₃ mixtures (e). (f) Intermolecular frequency contact maps of the different inserted peptides with TDP-43 LCD. In panels (c) and (f), the LARKS of each protein sequence are highlighted with dashed squares.

Furthermore, we evaluate the contact frequencies between the inserted peptides and FUS LCD proteins (Fig. 4(c)). The three LARKS capable of forming inter-protein β-sheet structures are highlighted by black dashed squares. We find that positively charged peptides ((SYSYKRKS)₃ and (SYSYKRKK)₃) show a very strong binding affinity for the two first LARKS of the sequence and the negatively charged residues located at its N-terminal domain. However, they do not display specific binding affinity for other subregions of the sequence. Conversely, negatively charged peptides ((SYGSYEGS)₃, (SYGESYGSYGEG)₂, and (SYGSYGSYEGEG)₂) exhibit more homogeneous cross-interactions throughout the protein sequence, except for the negatively charged residues in the N-terminal domain. This suggests that, even for a sequence with only two negatively charged residues and no positively charged amino acids, as FUS LCD, a potential strategy to preserve liquid-like behaviour is to insert charged molecules of the same sign as the protein that establish frequent interactions throughout the sequence (e.g., as π-π contacts) while enhancing electrostatic repulsion partially disrupting inter-protein connectivity. Based on our results from Fig. 3(b) and 4(a)-(c), we reveal that a balance of moderate electrostatic self-repulsion and non-specific attractive interactions—i.e., mediated by π-π (or cation-π) interactions—contributes to decreasing LARKS

high-density fluctuations triggering cross-β-sheet stacking. Importantly, the addition of the aforementioned peptides reduces protein condensate concentration by approximately 5% while largely maintaining their overall stability (Fig. 2(b)).

It is also remarkable that negatively charged peptides are capable of interacting in higher extent with the ₇₇STGGYG₈₂ LARKS, while positively charged sequences ((SYSYKRKS)₃ and (SYSYKRKK)₃) display much stronger binding affinity for the first two LARKS of the sequence (₃₇SYSGYS₄₂ and ₅₄SYSSYGQS₆₁). According to our residue-level simulations, this is a cooperative effect due to the presence of the N-terminal negatively charged residues near the location of these two LARKS, which establish attractive electrostatic interactions that further strengthen π-π contacts among the peptides and the adjacent LARKS. Nevertheless, we note that while positively charged peptides engage more locally with the first two LARKS of the sequence compared to negatively charged peptides, their impact on ageing kinetics is moderately lower (Fig. 3(b)).

Furthermore, we quantify whether there is any major conformational change in FUS LCD proteins when peptides are recruited into the condensate. To that goal, we evaluate the FUS LCD radius of gyration ($R_g$) in condensate bulk

conditions. Interestingly, we do not find substantial changes in $R_g$ when different peptide sequences are added with respect to the pure protein condensate (Fig. S7). Hence, a mechanism in which peptide recruitment decelerates ageing by inducing a conformational switch in FUS LCD favouring intra-molecular contacts vs. inter-molecular interactions is discarded for these systems. Moreover, we have computed the diffusion coefficient ($D$) of FUS LCD within the condensates upon peptide addition (Fig. S9). We find no correlation of slower $D$ with greater $t_{1/2}$. Therefore, condensate density reduction upon peptide inclusion appears to be the dominating mechanism by which ageing is decelerated in the studied mixtures.

We also examine the impact on ageing kinetics when peptides increase condensate density rather than reduce it. To that aim, we select $(WY)_{12}$, which is not self-repulsive and it significantly enhances condensate stability and protein concentration inside the condensate upon its recruitment (Fig. 2(b)). For this mixture, we observe a modest deceleration in the nucleation time (Fig. S10), which is comparable to the case of FUS LCD + $(SYSYKRKS)_3$. Strikingly, in FUS LCD/$(WY)_{12}$ condensates, the $R_g$ of FUS decreases with respect to the previous systems (Fig. S10), suggesting that a conformational change resulting into a more compact configurational ensemble can also modulate the ageing rate. As a consequence of such closer conformations, the ratio of intra-molecular contacts vs. inter-molecular ones becomes lower in FUS LCD. This result shows how an alternative mechanism might also regulate ageing kinetics in protein condensates. Nevertheless, the highest deceleration rate in cross-β-sheet formation is observed when peptides partially reduce condensate protein density instead of promoting more compact LCD conformations.

Next, we evaluate the contact frequency maps for TDP-43 LCD mixtures before ageing occurs. The inter-protein TDP-43 LCD contact maps in the absence (d) vs. presence of $(SYGSYEGS)_3$ peptides (e) are shown in Fig. 4. We find in this case that the overall inter-protein connectivity is reduced upon peptide addition as a consequence of the partial density reduction driven by self-repulsive electrostatic interactions among peptides. Crucially, the most frequent contacts, which occur between residues 61-65, that corresponds to the largest LARKS in TDP-43 (Fig. 1(b)), are significantly reduced by approximately 40%. Conversely to FUS LCD mixtures, where only minor subdomains reduce their intermolecular connectivity in advantage to other subregions which increase their connectivity (Fig. 4(a)-(b)), in TDP-43 LCD mixtures we observe a broad diminishment of inter-protein contacts across the entire sequence. In Fig. S6, we show the contact frequency maps for TDP-43 LCD—TDP-43 LCD in presence of the other inserted peptides, which notably resemble to that shown for $(SYGSYEGS)_3$.

We plot the intermolecular contacts of TDP-43 LCD with all peptides that decelerate ageing kinetics (Fig. 3(c)) as different panels in Fig. 4(f). TDP-43 LCD possesses an overall positive charge of $+2e$ with two negatively charged amino acids (1D and 1E) and four positively charged residues (3R and 1K). In TDP-43 LCD condensates, $(SYSYKRKK)_3$ insertion produces the lowest ageing deceleration of the whole set since the cross-interactions among proteins and peptides are only partially localized across the LARKS domain positioned from the 60th to the 70th residue, while the rest of the sequence mildly establishes substantial interactions with the peptides (Fig. 4(f); first panel). On the other hand, $(SYGKSYGSYGKG)_2$, which contains less than half of the positive charges compared to $(SYSYKRKK)_3$, is able to foster higher binding connectivity to TDP-43 LCD, and as a consequence, this sequence is one of the most effective in retarding ageing. Our results evidence how peptides with stronger affinity for the target protein possess much greater effectiveness in frustrating cross-β-sheet formation. Ageing deceleration relies on the affinity in protein-peptide interactions, and the partial condensate density reduction induced by peptide self-repulsion. Importantly, the inserted negatively charged peptides further support these observations ($(SYGSYEGS)_3$, $(SYGESYGSYGEG)_2$, and $(SYGSYGSYEGEG)_2$) since their strong impact on the cross-β-sheet nucleation time is triggered by the prominent heterotypic protein-peptide interactions established across the sequence, with most long-lived contacts taking place in the LARKS and near-LARKS regions: $_{27}GNNQGS_{32}$, $_{39}NFGAFS_{44}$, $_{55}AALQSS_{60}$, and $_{60}SWGAAGALASQ_{70}$. In that sense, the fact that more widespread peptide-protein contacts accumulate within (or around) the LARKS in TDP-43 LCD mixtures compared to FUS LCD condensates may explain why $t_{1/2}$ increases in greater extent in TDP-43 LCD with respect to FUS LCD (Fig. 3). Nevertheless, the nucleation lag time in FUS LCD mixtures can still be delayed up to a factor of 50 even when peptides display broad binding affinity for the entire sequence (Fig. 3(c)). We have also quantified if major conformational changes occur in TDP-43 LCD when peptides are added into the condensate. As found in FUS LCD mixtures, no significant changes in $R_g$ are observed upon the addition of peptide sequences which partially reduce condensate stability and density (Fig. S8). Likewise, we compute the diffusion coefficient of TDP-43 LCD for the different mixtures, and no correlation has been detected between $D$ and $t_{1/2}$ (Fig. S9).

Overall, based on our multiple computational analyses, we conclude that a combination of factors are involved in preventing the nucleation stage of cross-β-sheet structures: 1) A decrease of condensate density upon peptide insertion, which is associated to a lower critical solution temperature for phase separation (Fig. 2); and 2) Substantial binding affinity between peptides and aromatic-rich protein domains (Fig. 4(c)-(f)). Our simulations of FUS LCD and TDP-43 LCD condensate mixtures demonstrate how electrostatic repulsion between peptides and proteins is needed for decreasing condensate density (Fig. 2(b)). However,

when peptides exhibit excessively strong repulsion toward one another, they cannot effectively prevent LARKS high-density fluctuations since only at low concentration maintain LLPS, and at those conditions multiple condensate regions lack effective protein-peptide interactions. If peptides specifically bind to protein LARKS whilst promoting electrostatically-driven low-density regions, structural transitions become highly frustrated (Fig. 4(f)).

## III.   CONCLUSIONS

In this study, we provide a detailed exploration of the key molecular parameters that enable small peptides to decelerate the ageing kinetics of protein condensates. By combining equilibrium and non-equilibrium simulations of a residue-resolution protein model [78], we examine the potential that multiple peptide sequences possess in modulating the phase diagram, ageing kinetics, diffusion coefficient, conformational ensemble, and intermolecular interactions of two protein low-complexity domains (TDP-43 LCD and FUS LCD) whose liquid-to-solid transition has been associated to several neurodegenerative disorders [25, 36, 101]. Our simulations reveal that an effective strategy for decelerating ageing consists of partially reducing protein condensate density while minimally altering the stability of the condensate. That can be achieved through the inclusion of small peptides—24 amino acids long in this study—which combine aromatic residues acting as 'stickers', polar and uncharged residues behaving as 'spacers', and charged residues of the same sign enforcing peptide electrostatic self-repulsion. The balance between stickers, spacers, and charged residues is critical to ensure binding affinity to the protein aromatic regions susceptible to undergo cross-β-sheet transitions, and peptide self-repulsion to lower the condensate local density near the protein LARKS. We discover that peptides containing 10-15% of charged residues—either negatively charged, e.g., D or E, or positively charged, K—exhibit electrostatic self-repulsion and reduce condensate density without heavily compromising its stability (Fig. 2(b)). We also reveal that arginine, despite being positively charged, acts as a sticker since the target binding domains for the peptides are rich in aromatic residues, and it establishes cation-π interactions instead of contributing to condensate destabilization via electrostatic self-repulsion. By lowering the density of the condensates in approximately 5-10%, these peptides strongly reduce the probability of disordered LARKS transitioning into cross-β-sheets, a hallmark of condensate ageing and pathological aggregation.

We characterize the peptide-driven modulation of condensate ageing by measuring the half-time of cross-β-sheet formation (which reflects the duration of liquid-like behaviour within the condensates before turning into kinetically arrested assemblies; Fig. 3), and the nucleation lag-time (which corresponds to the onset before the emergence of the first cross-β-sheet nucleus; Fig. S3).

We find that most of the selected peptides, based on the condensate stability criterion (Fig. 2), are capable of substantially delaying the nucleation lag-time. Such deceleration in the nucleation stage is essential, as it is the rate-limiting step in solidification kinetics given that the cross-β-sheet growth barely varies upon peptide inclusion (Fig. 2(a)-(c)). Our findings for TDP-43 LCD condensates (Fig. 3(f)) reveal that peptides with strong binding affinity for LARKS and adjacent regions are particularly effective in delaying the nucleation lag-time, drastically reducing the likelihood of LARKS high-density fluctuations. Remarkably, we find that these peptides also preserve the diffusion coefficient of proteins in non-aged pure condensates, and their average conformational ensemble, quantified through their radius of gyration under condensed phase conditions. The resemblance of the protein-peptide mixtures to pure protein systems in terms of critical solution temperature, protein diffusion, and protein conformational ensemble suggests that peptides can be ideal molecular regulators of pathological liquid-to-solid transitions in biomolecular condensates. Moreover, our simulations show how small sequence variations in terms of composition, amino acid patterning, and net peptide charge radically determine the impact of peptide inclusion on condensate ageing kinetics.

Taken together, our work highlights three primary factors that contribute to a peptide's inhibitory potential of condensate hardening: (1) a reduction in condensate density without altering condensate stability and protein mobility; (2) moderate but consistent peptide-protein binding affinity (i.e., driven by π-π or cation-π interactions) that partially disrupts inter-protein molecular connectivity; and (3) selective interactions between peptides and aromatic-rich regions (e.g., LARKS), that further destabilize LARKS-rich environments through self-repulsive interactions. These, a priori, intuitive conclusions have been reached thanks to the possibility of exploring through realistic and efficient computational models how a large set of peptide sequences govern the stability and ageing of protein condensates. Experimentally testing such amount of sequence variations can be extremely challenging [102]. However, our work proposes an efficient computational pipeline by which the impact of sequence variations can be easily tested in condensate stability, viscoelastic properties, and ageing kinetics. Extending the sampling of the studied sequences to a considerable larger set can further contribute to decelerate, or even fully inhibit, the emergence of cross-β-sheet transitions in protein condensates. In that sense machine-learning algorithms can be extremely helpful to optimize sequence iterations which maximize the nucleation lag-time of cross-β-sheet aggregation. Therefore, our study provides a promising proof-of-concept for further exploration of potential sequences as modulators of condensate-related pathologies, offering insights that may guide future therapeutic designs. Refining peptide composition to enhance the specificity of interactions with LARKS and other key regions, can potentially offer even more robust strategies to control biocondensate phase behaviour and their strong

link to health and disease.

## IV. ACKNOWLEDGEMENTS

I. S.-B. acknowledges funding from Derek Brewer scholarship of Emmanuel College and EPSRC Doctoral Training Programme studentship, number EP/T517847/1, Ramon y Cajal fellowship (awarded to J.R.E.), as well as the UKRI EPSRC under the UK Government's guarantee scheme (EP/Z002028/1), following successful evaluation by the ERC (Consolidator Grant awarded to R.C.G.) under the European Union's Horizon Europe research and innovation programme. A. R. T. acknowledges funding from the European Union Horizon 2020 research and innovation programme (grant agreement 803326 to R.C.-G.). J. R. E. acknowledges funding from Emmanuel College, the University of Cambridge, the Ramon y Cajal fellowship (RYC2021-030937-I), and the Spanish scientific plan and committee for research reference PID2022-136919NA-C33. This work has been performed using resources provided by the Cambridge Tier-2 system operated by the University of Cambridge Research Computing Service (http://www.hpc.cam.ac.uk) funded by EPSRC Tier-2 capital grant EP/P020259/1-CS170. This work has also been performed using resources provided by Archer2 (https://www.archer2.ac.uk/) funded by EPSRC Tier-2 capital grant EP/P020259/e829. The authors also thankfully acknowledge RES computational resources provided by Mare Nostrum 5 through the activity 2024-3-0001.

[1] Y. Shin and C. P. Brangwynne, "Liquid phase condensation in cell physiology and disease," *Science*, vol. 357, no. 6357, 2017.

[2] C. P. Brangwynne, C. R. Eckmann, D. S. Courson, A. Rybarska, C. Hoege, J. Gharakhani, F. Jülicher, and A. A. Hyman, "Germline p granules are liquid droplets that localize by controlled dissolution/condensation," *Science*, vol. 324, no. 5935, pp. 1729–1732, 2009.

[3] S. Sanulli, M. Trnka, V. Dharmarajan, R. Tibble, B. Pascal, A. Burlingame, P. Griffin, J. Gross, and G. Narlikar, "Hp1 reshapes nucleosome core to promote phase separation of heterochromatin," *Nature*, vol. 575, no. 7782, pp. 390–394, 2019.

[4] G. Narlikar, S. Sanulli, J. D. Gross, P. Griffin, and M. Trnka, "The role of phase-separation in heterochromatin," *Biophysical Journal*, vol. 116, no. 3, p. 161a, 2019.

[5] J. R. Espinosa, J. A. Joseph, I. Sanchez-Burgos, A. Garaizar, D. Frenkel, and R. Collepardo-Guevara, "Liquid network connectivity regulates the stability and composition of biomolecular condensates with many components," *Proceedings of the National Academy of Sciences*, vol. 117, no. 24, pp. 13238–13247, 2020.

[6] S. F. Banani, H. O. Lee, A. A. Hyman, and M. K. Rosen, "Biomolecular condensates: organizers of cellular biochemistry," *Nature reviews Molecular cell biology*, vol. 18, no. 5, pp. 285–298, 2017.

[7] A. Molliex, J. Temirov, J. Lee, M. Coughlin, A. P. Kanagaraj, H. J. Kim, T. Mittag, and J. P. Taylor, "Phase separation by low complexity domains promotes stress granule assembly and drives pathological fibrillization," *Cell*, vol. 163, no. 1, pp. 123–133, 2015.

[8] A. C. Murthy, G. L. Dignon, Y. Kan, G. H. Zerze, S. H. Parekh, J. Mittal, and N. L. Fawzi, "Molecular interactions underlying liquid- liquid phase separation of the fus low-complexity domain," *Nature structural & molecular biology*, vol. 26, no. 7, pp. 637–648, 2019.

[9] H.-R. Li, W.-C. Chiang, P.-C. Chou, W.-J. Wang, and J.-r. Huang, "Tar dna-binding protein 43 (tdp-43) liquid–liquid phase separation is mediated by just a few aromatic residues," *Journal of Biological Chemistry*, vol. 293, no. 16, pp. 6090–6098, 2018.

[10] L. Latonen, "Phase-to-phase with nucleoli–stress responses, protein aggregation and novel roles of rna," *Frontiers in cellular neuroscience*, vol. 13, p. 151, 2019.

[11] M. Feric, N. Vaidya, T. S. Harmon, D. M. Mitrea, L. Zhu, T. M. Richardson, R. W. Kriwacki, R. V. Pappu, and C. P. Brangwynne, "Coexisting liquid phases underlie nucleolar subcompartments," *Cell*, vol. 165, no. 7, pp. 1686–1697, 2016.

[12] D. Mitrea, C. JA, G. CS, D. Ban, P. Banerjee, S. CB, N. A, D. AA, and R. Kriwacki, "Nucleophosmin integrates within the nucleolus via multi-modal interactions with proteins displaying r-rich linear motifs and rrna," *eLife Sciences*, 02 2016.

[13] D. M. Mitrea, J. A. Cika, C. B. Stanley, A. Nourse, P. L. Onuchic, P. R. Banerjee, A. H. Phillips, C.-G. Park, A. A. Deniz, and R. W. Kriwacki, "Self-interaction of npm1 modulates multiple mechanisms of liquid–liquid phase separation," *Nature communications*, vol. 9, no. 1, pp. 1–13, 2018.

[14] S. C. Weber and C. P. Brangwynne, "Inverse size scaling of the nucleolus by a concentration-dependent phase transition," *Current Biology*, vol. 25, no. 5, pp. 641–646, 2015.

[15] F. Frottin, F. Schueder, S. Tiwary, R. Gupta, R. Körner, T. Schlichthaerle, J. Cox, R. Jungmann, F. Hartl, and M. Hipp, "The nucleolus functions as a phase-separated protein quality control compartment," *Science*, vol. 365, no. 6451, pp. 342–347, 2019.

[16] P. Yang, C. Mathieu, R.-M. Kolaitis, P. Zhang, J. Messing, U. Yurtsever, Z. Yang, J. Wu, Y. Li, Q. Pan, *et al.*, "G3bp1 is a tunable switch that triggers phase separation to assemble stress granules," *Cell*, vol. 181, no. 2, pp. 325–345, 2020.

[17] N. Bley, M. Lederer, B. Pfalz, C. Reinke, T. Fuchs, M. Glaß, B. Möller, and S. Hüttelmaier, "Stress granules are dispensable for mrna stabilization during cellular stress," *Nucleic acids research*, vol. 43, no. 4, pp. e26–e26, 2015.

[18] R. Oliva, S. K. Mukherjee, Z. Fetahaj, S. Möbitz, and R. Winter, "Perturbation of liquid droplets of p-granule protein laf-1 by the antimicrobial peptide ll-iii," *Chemical Communications*, vol. 56, no. 78, pp. 11577–11580, 2020.

[19] D. S. Protter, B. S. Rao, B. Van Treeck, Y. Lin, L. Mizoue, M. K. Rosen, and R. Parker, "Intrinsically disordered regions can contribute promiscuous interactions to rnp granule assembly," *Cell reports*, vol. 22, no. 6, pp. 1401–1412, 2018.

[20] F. Niss, L. Piñero-Paez, W. Zaidi, E. Hallberg, and A.-L. Ström, "Key modulators of the stress granule response tia1, tdp-43, and g3bp1 are altered by polyglutamine-expanded atxn7," *Molecular Neurobiology*, vol. 59, no. 8, pp. 5236–5251, 2022.

[21] J. Gal, J. Chen, D.-Y. Na, L. Tichacek, K. R. Barnett, and H. Zhu, "The acetylation of lysine-376 of g3bp1 regulates rna binding and stress granule dynamics," *Molecular and cellular biology*, 2019.

[22] W.-C. Tsai, S. Gayatri, L. C. Reineke, G. Sbardella, M. T. Bedford, and R. E. Lloyd, "Arginine demethylation of g3bp1 promotes stress granule assembly," *Journal of Biological Chemistry*, vol. 291, no. 43, pp. 22671–22685, 2016.

[23] A. Pichet Binette, A. Mammana, L. Wisse, M. Rossi, O. Strandberg, R. Smith, N. Mattsson-Carlgren, S. Janelidze, S. Palmqvist, ADNI, *et al.*, "Associations between misfolded alpha-synuclein aggregates and alzheimer's disease pathology in vivo," *Alzheimer's & Dementia*, 2024.

[24] V. N. Uversky, "Neuropathology, biochemistry, and biophysics of $\alpha$-synuclein aggregation," *Journal of neurochemistry*, vol. 103, no. 1, pp. 17–37, 2007.

[25] D. Ghosh, S. Mehra, S. Sahay, P. K. Singh, and S. K. Maji, "$\alpha$-synuclein aggregation and its modulation," *International journal of biological macromolecules*, vol. 100, pp. 37–54, 2017.

[26] W. M. Babinchak and W. K. Surewicz, "Liquid–liquid phase separation and its mechanistic role in pathological protein aggregation," *Journal of molecular biology*, vol. 432, no. 7, pp. 1910–1925, 2020.

[27] J. Li, M. Zhang, W. Ma, B. Yang, H. Lu, F. Zhou, and L. Zhang, "Post-translational modifications in liquid-liquid phase separation: a comprehensive review," *Molecular biomedicine*, vol. 3, no. 1, p. 13, 2022.

[28] M. Li, Y. Fan, Q. Li, X. Wang, L. Zhao, and M. Zhu, "Liquid-liquid phase separation promotes protein aggregation and its implications in ferroptosis in parkinson's disease dementia," *Oxidative Medicine and Cellular Longevity*, vol. 2022, no. 1, p. 7165387, 2022.

[29] C. J. Webber, S. E. Lei, and B. Wolozin, "The pathophysiology of neurodegenerative disease: Disturbing the balance between phase separation and irreversible aggregation," *Progress in molecular biology and translational science*, vol. 174, pp. 187–223, 2020.

[30] I. P. Blair, K. L. Williams, S. T. Warraich, J. C. Durnall, A. D. Thoeng, J. Manavis, P. C. Blumbergs, S. Vucic, M. C. Kiernan, and G. A. Nicholson, "Fus mutations in amyotrophic lateral sclerosis: clinical, pathological, neurophysiological and genetic analysis," *Journal of Neurology, Neurosurgery & Psychiatry*, vol. 81, no. 6, pp. 639–645, 2010.

[31] Y. Shang and E. J. Huang, "Mechanisms of fus mutations in familial amyotrophic lateral sclerosis," *Brain research*, vol. 1647, pp. 65–78, 2016.

[32] H. Deng, K. Gao, and J. Jankovic, "The role of fus gene variants in neurodegenerative diseases," *Nature Reviews Neurology*, vol. 10, no. 6, pp. 337–348, 2014.

[33] M. A. Gitcho, R. H. Baloh, S. Chakraverty, K. Mayo, J. B. Norton, D. Levitch, K. J. Hatanpaa, C. L. White III, E. H. Bigio, R. Caselli, *et al.*, "Tdp-43 a315t mutation in familial motor neuron disease," *Annals of Neurology: Official Journal of the American Neurological Association and the Child Neurology Society*, vol. 63, no. 4, pp. 535–538, 2008.

[34] E. Kabashi, P. N. Valdmanis, P. Dion, D. Spiegelman, B. J. McConkey, C. V. Velde, J.-P. Bouchard, L. Lacomblez, K. Pochigaeva, F. Salachas, *et al.*, "Tardbp mutations in individuals with sporadic and familial amyotrophic lateral sclerosis," *Nature genetics*, vol. 40, no. 5, pp. 572–574, 2008.

[35] S. Lattante, G. A. Rouleau, and E. Kabashi, "Tardbp and fus mutations associated with amyotrophic lateral sclerosis: summary and update," *Human mutation*, vol. 34, no. 6, pp. 812–826, 2013.

[36] J. L. Carey and L. Guo, "Liquid-liquid phase separation of tdp-43 and fus in physiology and pathology of neurodegenerative diseases," *Frontiers in molecular biosciences*, vol. 9, p. 826719, 2022.

[37] S. Wegmann, B. Eftekharzadeh, K. Tepper, K. M. Zoltowska, R. E. Bennett, S. Dujardin, P. R. Laskowski, D. MacKenzie, T. Kamath, C. Commins, *et al.*, "Tau protein liquid–liquid phase separation can initiate tau aggregation," *The EMBO journal*, vol. 37, no. 7, p. e98049, 2018.

[38] R. E. Tato, A. Frank, and A. Hernanz, "Tau protein concentrations in cerebrospinal fluid of patients with dementia of the alzheimer type.," *Journal of Neurology, Neurosurgery & Psychiatry*, vol. 59, no. 3, pp. 280–283, 1995.

[39] M. Von Bergen, S. Barghorn, L. Li, A. Marx, J. Biernat, E.-M. Mandelkow, and E. Mandelkow, "Mutations of tau protein in frontotemporal dementia promote aggregation of paired helical filaments by enhancing local $\beta$-structure," *Journal of Biological Chemistry*, vol. 276, no. 51, pp. 48165–48174, 2001.

[40] D. C. Costa, G. A. de Oliveira, E. A. Cino, I. N. Soares, L. P. Rangel, and J. L. Silva, "Aggregation and prion-like properties of misfolded tumor suppressors: is cancer a prion disease?," *Cold Spring Harbor perspectives in biology*, vol. 8, no. 10, p. a023614, 2016.

[41] S. Alberti and D. Dormann, "Liquid–liquid phase separation in disease," *Annual review of genetics*, vol. 53, pp. 171–194, 2019.

[42] S. Spannl, M. Tereshchenko, G. J. Mastromarco, S. J. Ihn, and H. O. Lee, "Biomolecular condensates in neurodegeneration and cancer," *Traffic*, vol. 20, no. 12, pp. 890–911, 2019.

[43] A. Navalkar, S. Ghosh, S. Pandey, A. Paul, D. Datta, and S. K. Maji, "Prion-like p53 amyloids in cancer," *Biochemistry*, vol. 59, no. 2, pp. 146–155, 2019.

[44] G. A. de Oliveira, Y. Cordeiro, J. L. Silva, and T. C. Vieira, "Liquid-liquid phase transitions and amyloid aggregation in proteins related to cancer and neurodegenerative diseases," *Advances in protein chemistry and structural biology*, vol. 118, pp. 289–331, 2019.

[45] K. Taniue and N. Akimitsu, "Aberrant phase separation and cancer," *The FEBS journal*, vol. 289, no. 1, pp. 17–39, 2022.

[46] M. P. Hughes, M. R. Sawaya, D. R. Boyer, L. Goldschmidt, J. A. Rodriguez, D. Cascio, L. Chong, T. Gonen, and D. S. Eisenberg, "Atomic structures of low-complexity protein segments reveal kinked $\beta$ sheets that assemble networks," *Science*, vol. 359, no. 6376, pp. 698–701, 2018.

[47] L. Emmanouilidis, E. Bartalucci, Y. Kan, M. Ijavi, M. E. Pérez, P. Afanasyev, D. Boehringer, J. Zehnder, S. H. Parekh, M. Bonn, et al., "A solid beta-sheet structure is formed at the surface of fus droplets during aging," *Nature Chemical Biology*, pp. 1–9, 2024.

[48] M. G. Iadanza, M. P. Jackson, E. W. Hewitt, N. A. Ranson, and S. E. Radford, "A new era for understanding amyloid structures and disease," *Nature reviews Molecular cell biology*, vol. 19, no. 12, pp. 755–773, 2018.

[49] M. Linsenmeier, L. Faltova, C. Morelli, U. Capasso Palmiero, C. Seiffert, A. M. Küffner, D. Pinotsi, J. Zhou, R. Mezzenga, and P. Arosio, "The interface of condensates of the hnrnpa1 low-complexity domain promotes formation of amyloid fibrils," *Nature chemistry*, vol. 15, no. 10, pp. 1340–1349, 2023.

[50] A. R. Tejedor, I. Sanchez-Burgos, M. Estevez-Espinosa, A. Garaizar, R. Collepardo-Guevara, J. Ramirez, and J. R. Espinosa, "Protein structural transitions critically transform the network connectivity and viscoelasticity of rna-binding protein condensates but rna can prevent it," *Nature communications*, vol. 13, no. 1, p. 5717, 2022.

[51] Y. Shen, A. Chen, W. Wang, Y. Shen, F. S. Ruggeri, S. Aime, Z. Wang, S. Qamar, J. R. Espinosa, A. Garaizar, et al., "The liquid-to-solid transition of fus is promoted by the condensate surface," *Proceedings of the National Academy of Sciences*, vol. 120, no. 33, p. e2301366120, 2023.

[52] S. Chatterjee, Y. Kan, M. Brzezinski, K. Koynov, R. M. Regy, A. C. Murthy, K. A. Burke, J. J. Michels, J. Mittal, N. L. Fawzi, et al., "Reversible kinetic trapping of fus biomolecular condensates," *Advanced Science*, vol. 9, no. 4, p. 2104247, 2022.

[53] A. Patel, H. O. Lee, L. Jawerth, S. Maharana, M. Jahnel, M. Y. Hein, S. Stoynov, J. Mahamid, S. Saha, T. M. Franzmann, et al., "A liquid-to-solid phase transition of the als protein fus accelerated by disease mutation," *Cell*, vol. 162, no. 5, pp. 1066–1077, 2015.

[54] L. Jawerth, E. Fischer-Friedrich, S. Saha, J. Wang, T. Franzmann, X. Zhang, J. Sachweh, M. Ruer, M. Ijavi, S. Saha, et al., "Protein condensates as aging maxwell fluids," *Science*, vol. 370, no. 6522, pp. 1317–1323, 2020.

[55] R. J. Wheeler, H. O. Lee, I. Poser, A. Pal, T. Doeleman, S. Kishigami, S. Kour, E. N. Anderson, L. Marrone, A. C. Murthy, et al., "Small molecules for modulating protein driven liquid-liquid phase separation in treating neurodegenerative disease," *BioRxiv*, p. 721001, 2019.

[56] M. Y. Fang, S. Markmiller, A. Q. Vu, A. Javaherian, W. E. Dowdle, P. Jolivet, P. J. Bushway, N. A. Castello, A. Baral, M. Y. Chan, et al., "Small-molecule modulation of tdp-43 recruitment to stress granules prevents persistent tdp-43 accumulation in als/ftd," *Neuron*, vol. 103, no. 5, pp. 802–819, 2019.

[57] I. Maruri-Lopez and M. Chodasiewicz, "Involvement of small molecules and metabolites in regulation of biomolecular condensate properties," *Current Opinion in Plant Biology*, vol. 74, p. 102385, 2023.

[58] B. Dai, T. Zhong, Z.-X. Chen, W. Chen, N. Zhang, X.-L. Liu, L.-Q. Wang, J. Chen, and Y. Liang, "Myricetin slows liquid–liquid phase separation of tau and activates atg5-dependent autophagy to suppress tau toxicity," *Journal of Biological Chemistry*, vol. 297, no. 4, 2021.

[59] V. A. Korobeynikov, A. K. Lyashchenko, B. Blanco-Redondo, P. Jafar-Nejad, and N. A. Shneider, "Antisense oligonucleotide silencing of fus expression as a therapeutic approach in amyotrophic lateral sclerosis," *Nature medicine*, vol. 28, no. 1, pp. 104–116, 2022.

[60] L. Palanikumar, L. Karpauskaite, M. Al-Sayegh, I. Chehade, M. Alam, S. Hassan, D. Maity, L. Ali, M. Kalmouni, Y. Hunashal, et al., "Protein mimetic amyloid inhibitor potently abrogates cancer-associated mutant p53 aggregation and restores tumor suppressor function," *Nature communications*, vol. 12, no. 1, p. 3962, 2021.

[61] M. Derbis, E. Kul, D. Niewiadomska, M. Sekrecki, A. Piasecka, K. Taylor, R. K. Hukema, O. Stork, and K. Sobczak, "Short antisense oligonucleotides alleviate the pleiotropic toxicity of rna harboring expanded cgg repeats," *Nature communications*, vol. 12, no. 1, p. 1265, 2021.

[62] B. C. Cui, V. Sikirzhytski, M. Aksenova, M. D. Lucius, G. H. Levon, Z. T. Mack, C. Pollack, D. Odhiambo, E. Broude, S. B. Lizarraga, et al., "Pharmacological inhibition of dead-box rna helicase 3 attenuates stress granule assembly," *Biochemical pharmacology*, vol. 182, p. 114280, 2020.

[63] G. Catara, G. Grimaldi, L. Schembri, D. Spano, G. Turacchio, M. Lo Monte, A. R. Beccari, C. Valente, and D. Corda, "Parp1-produced poly-adp-ribose causes the parp12 translocation to stress granules and impairment of golgi complex functions," *Scientific reports*, vol. 7, no. 1, p. 14035, 2017.

[64] A. M. Heberle, P. R. Navas, M. Langelaar-Makkinje, K. Kasack, A. Sadik, E. Faessler, U. Hahn, P. Marx-Stoelting, C. A. Opitz, C. Sers, et al., "The pi3k and mapk/p38 pathways control stress granule assembly in a hierarchical manner," *Life science alliance*, vol. 2, no. 2, 2019.

[65] K. Hou, P. Ge, M. R. Sawaya, J. L. Dolinsky, Y. Yang, Y. X. Jiang, L. Lutter, D. R. Boyer, X. Cheng, J. Pi, et al., "How short peptides can disassemble ultra-stable tau fibrils extracted from alzheimers disease brain by a strain-relief mechanism," *bioRxiv*, pp. 2024–03, 2024.

[66] H. Zhang, S. Zhang, H. He, W. Zhao, J. Chen, and R.-g. Shao, "Gap 161 targets and downregulates g 3 bp to suppress cell growth and potentiate cisplaitin-mediated cytotoxicity to colon carcinoma hct 116 cells," *Cancer science*, vol. 103, no. 10, pp. 1848–1856, 2012.

[67] S. Jonchhe, W. Pan, P. Pokhrel, and H. Mao, "Small molecules modulate liquid-to-solid transitions in phase-separated tau condensates," *Angewandte Chemie International Edition*, vol. 61, no. 23, p. e202113156, 2022.

[68] J. Habchi, P. Arosio, M. Perni, A. R. Costa, M. Yagi-Utsumi, P. Joshi, S. Chia, S. I. Cohen, M. B. Müller, S. Linse, et al., "An anticancer drug suppresses the primary nucleation reaction that initiates the production of the toxic a$\beta$42 aggregates linked with alzheimer's disease," *Science advances*, vol. 2, no. 2, p. e1501244, 2016.

[69] J. Habchi, S. Chia, R. Limbocker, B. Mannini, M. Ahn, M. Perni, O. Hansson, P. Arosio, J. R. Kumita, P. K. Challa, et al., "Systematic development of small molecules to inhibit specific microscopic steps of a$\beta$42 aggregation in alzheimer's disease," *Proceedings of the National Academy of Sciences*, vol. 114, no. 2, pp. E200–E208, 2017.

[70] G. T. Heller, F. A. Aprile, T. C. Michaels, R. Limbocker, M. Perni, F. S. Ruggeri, B. Mannini, T. Löhr, M. Bonomi, C. Camilloni, et al., "Small-molecule sequestration of amyloid-$\beta$ as a drug discovery strategy for alzheimer's disease," *Science Advances*, vol. 6, no. 45, p. eabb5924, 2020.

[71] A. Girdhar, V. Bharathi, V. R. Tiwari, S. Abhishek, W. Deeksha, U. S. Mahawar, G. Raju, S. K. Singh, G. Prabusankar, E. Rajakumara, *et al.*, "Computational insights into mechanism of aim4-mediated inhibition of aggregation of tdp-43 protein implicated in als and evidence for in vitro inhibition of liquid-liquid phase separation (llps) of tdp-432c-a315t by aim4," *International journal of biological macromolecules*, vol. 147, pp. 117–130, 2020.

[72] S. T. Dada, Z. Toprakcioglu, M. P. Cali, A. Röntgen, M. C. Hardenberg, O. M. Morris, L. K. Mrugalla, T. P. Knowles, and M. Vendruscolo, "Pharmacological inhibition of $\alpha$-synuclein aggregation within liquid condensates," *Nature Communications*, vol. 15, no. 1, p. 3835, 2024.

[73] R. Staats, Z. F. Brotzakis, S. Chia, R. I. Horne, and M. Vendruscolo, "Optimization of a small molecule inhibitor of secondary nucleation in $\alpha$-synuclein aggregation," *Frontiers in Molecular Biosciences*, vol. 10, p. 1155753, 2023.

[74] S. Chia, Z. Faidon Brotzakis, R. I. Horne, A. Possenti, B. Mannini, R. Cataldi, M. Nowinska, R. Staats, S. Linse, T. P. Knowles, *et al.*, "Structure-based discovery of small-molecule inhibitors of the autocatalytic proliferation of $\alpha$-synuclein aggregates," *Molecular Pharmaceutics*, vol. 20, no. 1, pp. 183–193, 2022.

[75] I. W. Hamley, "Small bioactive peptides for biomaterials design and therapeutics," *Chemical reviews*, vol. 117, no. 24, pp. 14015–14041, 2017.

[76] R. Oliva, S. K. Mukherjee, L. Ostermeier, L. A. Pazurek, S. Kriegler, V. Bader, D. Prumbaum, S. Raunser, K. F. Winklhofer, J. Tatzelt, *et al.*, "Remodeling of the fibrillation pathway of $\alpha$-synuclein by interaction with antimicrobial peptide ll-iii," *Chemistry–A European Journal*, vol. 27, no. 46, pp. 11845–11851, 2021.

[77] B. Szała-Mendyk, T. M. Phan, P. Mohanty, and J. Mittal, "Challenges in studying the liquid-to-solid phase transitions of proteins using computer simulations," *Current opinion in chemical biology*, vol. 75, p. 102333, 2023.

[78] G. Tesei and K. Lindorff-Larsen, "Improved predictions of phase behaviour of intrinsically disordered proteins by tuning the interaction range," *bioRxiv*, pp. 2022–07, 2022.

[79] E. L. Guenther, Q. Cao, H. Trinh, J. Lu, M. R. Sawaya, D. Cascio, D. R. Boyer, J. A. Rodriguez, M. P. Hughes, and D. S. Eisenberg, "Atomic structures of tdp-43 lcd segments and insights into reversible or pathogenic aggregation," *Nature structural & molecular biology*, vol. 25, no. 6, pp. 463–471, 2018.

[80] J. Wessén, S. Das, T. Pal, and H. S. Chan, "Analytical formulation and field-theoretic simulation of sequence-specific phase separation of protein-like heteropolymers with short-and long-spatial-range interactions," *The Journal of Physical Chemistry B*, vol. 126, no. 45, pp. 9222–9245, 2022.

[81] I. Sanchez-Burgos, J. A. Joseph, R. Collepardo-Guevara, and J. R. Espinosa, "Size conservation emerges spontaneously in biomolecular condensates formed by scaffolds and surfactant clients," *Scientific Reports*, vol. 11, no. 1, p. 15241, 2021.

[82] I. Sanchez-Burgos, J. R. Espinosa, J. A. Joseph, and R. Collepardo-Guevara, "Rna length has a non-trivial effect in the stability of biomolecular condensates formed by rna-binding proteins," *PLoS computational biology*, vol. 18, no. 2, p. e1009810, 2022.

[83] I. Sanchez-Burgos, L. Herriott, R. Collepardo-Guevara, and J. R. Espinosa, "Surfactants or scaffolds? rnas of varying lengths control the thermodynamic stability of condensates differently," *Biophysical Journal*, vol. 122, no. 14, pp. 2973–2987, 2023.

[84] J. A. Joseph, J. R. Espinosa, I. Sanchez-Burgos, A. Garaizar, D. Frenkel, and R. Collepardo-Guevara, "Thermodynamics and kinetics of phase separation of protein-rna mixtures by a minimal model," *Biophysical Journal*, vol. 120, no. 7, pp. 1219–1230, 2021.

[85] I. Sanchez-Burgos, J. R. Espinosa, J. A. Joseph, and R. Collepardo-Guevara, "Valency and binding affinity variations can regulate the multilayered organization of protein condensates with many components," *Biomolecules*, vol. 11, no. 2, p. 278, 2021.

[86] L. M. Pietrek, L. S. Stelzl, and G. Hummer, "Hierarchical ensembles of intrinsically disordered proteins at atomic resolution in molecular dynamics simulations," *Journal of chemical theory and computation*, vol. 16, no. 1, pp. 725–737, 2019.

[87] D. Fusco, J. J. Headd, A. De Simone, J. Wang, and P. Charbonneau, "Characterizing protein crystal contacts and their role in crystallization: rubredoxin as a case study," *Soft matter*, vol. 10, no. 2, pp. 290–302, 2014.

[88] P. C. Souza, L. P. Borges Araujo, C. Brasnett, R. A. Moreira, F. Grunewald, P. Park, L. Wang, H. Razmazma, A. C. Borges-Araujo, L. F. Cofas-Vargas, *et al.*, "Gōmartini 3: From large conformational changes in proteins to environmental bias corrections," *BioRxiv*, pp. 2024–04, 2024.

[89] M. Tsanai, P. W. Frederix, C. F. Schroer, P. C. Souza, and S. J. Marrink, "Coacervate formation studied by explicit solvent coarse-grain molecular dynamics with the martini model," *Chemical Science*, vol. 12, no. 24, pp. 8521–8530, 2021.

[90] A. Feito, I. Sanchez-Burgos, A. Rey, R. Collepardo-Guevara, J. R. Espinosa, and A. R. Tejedor, "Capturing single-molecule properties does not ensure accurate prediction of biomolecular phase diagrams," *Molecular Physics*, p. e2425757, 2024.

[91] F. Dar and R. Pappu, "Phase separation: Restricting the sizes of condensates," *Elife*, vol. 9, p. e59663, 2020.

[92] W. Guo, T. Lu, R. Crisci, S. Nagao, T. Wei, and Z. Chen, "Determination of protein conformation and orientation at buried solid/liquid interfaces," *Chemical Science*, vol. 14, no. 11, pp. 2999–3009, 2023.

[93] M. Paloni, R. Bailly, L. Ciandrini, and A. Barducci, "Unraveling molecular interactions in liquid–liquid phase separation of disordered proteins by atomistic simulations," *The Journal of Physical Chemistry B*, vol. 124, no. 41, pp. 9009–9016, 2020.

[94] J.-E. Shea, R. B. Best, and J. Mittal, "Physics-based computational and theoretical approaches to intrinsically disordered proteins," *Current opinion in structural biology*, vol. 67, pp. 219–225, 2021.

[95] K. Sarthak, D. Winogradoff, Y. Ge, S. Myong, and A. Aksimentiev, "Benchmarking molecular dynamics force fields for all-atom simulations of biological condensates," *Journal of Chemical Theory and Computation*, vol. 19, no. 12, pp. 3721–3740, 2023.

[96] S. Blazquez, I. Sanchez-Burgos, J. Ramirez, T. Higginbotham, M. M. Conde, R. Collepardo-Guevara, A. R. Tejedor, and J. R. Espinosa, "Location and concentration of aromatic-rich segments dictates the percolating inter-molecular network and viscoelastic properties of ageing condensates," *Advanced Science*, vol. 10, no. 25, p. 2207742, 2023.

[97] A. R. Tejedor, R. Collepardo-Guevara, J. Ramírez, and J. R. Espinosa, "Time-dependent material properties of aging biomolecular condensates from different viscoelasticity measurements in molecular dynamics simulations," *The Journal of Physical Chemistry B*, vol. 127, no. 20, pp. 4441–4459, 2023.

[98] F. Cao, S. von Bülow, G. Tesei, and K. Lindorff-Larsen, "A coarse-grained model for disordered and multi-domain proteins," *Protein Science*, vol. 33, no. 11, p. e5172, 2024.

[99] A. Garaizar, J. R. Espinosa, J. A. Joseph, G. Krainer, Y. Shen, T. P. Knowles, and R. Collepardo-Guevara, "Aging can transform single-component protein condensates into multiphase architectures," *Proceedings of the National Academy of Sciences*, vol. 119, no. 26, p. e2119800119, 2022.

[100] N. A. Erkamp, I. A. Sanchez-Burgos, A. Zhou, T. J. Krug, S. S. Qamar, T. Sneideris, E. Zhang, K. Nakajima, A. Chen, R. Collepardo-Guevara, *et al.*, "Dynamically arrested condensate fusion creates complex structures with varying material properties," *bioRxiv*, pp. 2024–11, 2024.

[101] B. Portz, B. L. Lee, and J. Shorter, "Fus and tdp-43 phases in health and disease," *Trends in biochemical sciences*, vol. 46, no. 7, pp. 550–563, 2021.

[102] S. Alberti and A. A. Hyman, "Biomolecular condensates at the nexus of cellular stress, protein aggregation disease and ageing," *Nature Reviews Molecular Cell Biology*, vol. 22, no. 3, pp. 196–213, 2021.

[103] J. S. Rowlinson and B. Widom, *Molecular theory of capillarity*. Courier Corporation, 2013.

[104] G. Tesei, T. K. Schulze, R. Crehuet, and K. Lindorff-Larsen, "Accurate model of liquid–liquid phase behavior of intrinsically disordered proteins from optimization of single-chain properties," *Proceedings of the National Academy of Sciences*, vol. 118, no. 44, p. e2111696118, 2021.

[105] A. Bremer, M. Farag, W. M. Borcherds, I. Peran, E. W. Martin, R. V. Pappu, and T. Mittag, "Deciphering how naturally occurring sequence features impact the phase behaviors of disordered prion-like domains," *bioRxiv*, 2021.

[106] G. Krainer, T. J. Welsh, J. A. Joseph, J. R. Espinosa, S. Wittmann, E. de Csilléry, A. Sridhar, Z. Toprakcioglu, G. Gudiškytė, M. A. Czekalska, *et al.*, "Reentrant liquid condensate phase of proteins is stabilized by hydrophobic and non-ionic interactions," *Nature Communications*, vol. 12, no. 1, pp. 1–14, 2021.

[107] A. R. Tejedor, A. Aguirre Gonzalez, M. J. Maristany, P. Y. Chew, K. Rusell, J. Ramirez, J. R. Espinosa, and R. Collepardo-Guevara, "Chemically-informed coarse-graining of electrostatic forces in charge-rich biomolecular condensates," *bioRxiv*, pp. 2024–07, 2024.

[108] J. Wang, J.-M. Choi, A. S. Holehouse, H. O. Lee, X. Zhang, M. Jahnel, S. Maharana, R. Lemaitre, A. Pozniakovsky, D. Drechsel, *et al.*, "A molecular grammar governing the driving forces for phase separation of prion-like rna binding proteins," *Cell*, vol. 174, no. 3, pp. 688–699, 2018.

[109] Y. Yang, H. B. Jones, T. P. Dao, and C. A. Castaneda, "Single amino acid substitutions in stickers, but not spacers, substantially alter ubqln2 phase transitions and dense phase material properties," *The Journal of Physical Chemistry B*, vol. 123, no. 17, pp. 3618–3629, 2019.

[110] D. De Sancho and X. Lopez, "Crossover in aromatic amino acid interaction strength: Tyrosine vs. phenylalanine in biomolecular condensates," *bioRxiv*, pp. 2024–10, 2024.

[111] S. Hutten, S. Usluer, B. Bourgeois, F. Simonetti, H. M. Odeh, C. M. Fare, M. Czuppa, M. Hruska-Plochan, M. Hofweber, M. Polymenidou, *et al.*, "Nuclear import receptors directly bind to arginine-rich dipeptide repeat proteins and suppress their pathological interactions," *Cell reports*, vol. 33, no. 12, 2020.

[112] G. L. Dignon, W. Zheng, Y. C. Kim, R. B. Best, and J. Mittal, "Sequence determinants of protein phase behavior from a coarse-grained model," *PLoS computational biology*, vol. 14, no. 1, p. e1005941, 2018.

[113] J. A. Joseph, A. Reinhardt, A. Aguirre, P. Y. Chew, K. O. Russell, J. R. Espinosa, A. Garaizar, and R. Collepardo-Guevara, "Physics-driven coarse-grained model for biomolecular phase separation with near-quantitative accuracy," *Nature Computational Science*, vol. 1, no. 11, pp. 732–743, 2021.

[114] S. Maharana, J. Wang, D. K. Papadopoulos, D. Richter, A. Pozniakovsky, I. Poser, M. Bickle, S. Rizk, J. Guillén-Boixet, T. M. Franzmann, M. Jahnel, L. Marrone, Y.-T. Chang, J. Sterneckert, P. Tomancak, A. A. Hyman, and S. Alberti, "Rna buffers the phase separation behavior of prion-like rna binding proteins," *Science*, vol. 360, no. 6391, pp. 918–921, 2018.

[115] P. R. Banerjee, A. N. Milin, M. M. Moosa, P. L. Onuchic, and A. A. Deniz, "Reentrant phase transition drives dynamic substructure formation in ribonucleoprotein droplets," *Angewandte Chemie*, vol. 129, no. 38, pp. 11512–11517, 2017.

[116] A. R. Tejedor, A. Garaizar, J. Ramírez, and J. R. Espinosa, "'rna modulation of transport properties and stability in phase-separated condensates," *Biophysical Journal*, vol. 120, no. 23, pp. 5169–5186, 2021.

[117] I. Alshareedah, T. Kaur, J. Ngo, H. Seppala, L.-A. D. Kounatse, W. Wang, M. M. Moosa, and P. R. Banerjee, "Interplay between short-range attraction and long-range repulsion controls reentrant liquid condensation of ribonucleoprotein–rna complexes," *Journal of the American Chemical Society*, vol. 141, no. 37, pp. 14593–14602, 2019.

[118] C. Wang, H. R. Kilgore, A. P. Latham, and B. Zhang, "Nonspecific yet selective interactions contribute to small molecule condensate binding," *Journal of Chemical Theory and Computation*, 2024.

[119] K. Kamagata, M. Ariefai, H. Takahashi, A. Hando, D. R. G. Subekti, K. Ikeda, A. Hirano, and T. Kameda, "Rational peptide design for regulating liquid–liquid phase separation on the basis of residue–residue contact energy," *Scientific Reports*, vol. 12, no. 1, p. 13718, 2022.

[120] A. Netzer, A. Baruch Leshem, S. Veretnik, I. Edelstein, and A. Lampel, "Regulation of peptide liquid–liquid phase separation by aromatic amino acid composition," *Small*, p. 2401665, 2024.

[121] S. Torrino, W. M. Oldham, A. R. Tejedor, I. S. Burgos, L. Nasr, N. Rachedi, K. Fraissard, C. Chauvet, C. Sbai, B. P. O'Hara, *et al.*, "Mechano-dependent sorbitol accumulation supports biomolecular condensate," *Cell*, 2023.

[122] J. Zhong, X. Xu, and C. Wang, "Computer-aided design of peptides for suppressing liquid-liquid phase separation," *Journal of Electrical Systems*, vol. 20, no. 3, pp. 2298–2303, 2024.

[123] G. Meisl, J. B. Kirkegaard, P. Arosio, T. C. Michaels, M. Vendruscolo, C. M. Dobson, S. Linse, and T. P. Knowles, "Molecular mechanisms of protein aggregation from global fitting of kinetic models," *Nature protocols*,

vol. 11, no. 2, pp. 252–272, 2016.

[124] P. Robustelli, S. Piana, and D. E. Shaw, "Developing a molecular dynamics force field for both folded and disordered protein states," *Proceedings of the National Academy of Sciences*, vol. 115, no. 21, pp. E4758–E4766, 2018.

[125] A. Šarić, Y. C. Chebaro, T. P. Knowles, and D. Frenkel, "Crucial role of nonspecific interactions in amyloid nucleation," *Proceedings of the National Academy of Sciences*, vol. 111, no. 50, pp. 17869–17874, 2014.

[126] A. Garaizar, J. R. Espinosa, J. A. Joseph, and R. Collepardo-Guevara, "Kinetic interplay between droplet maturation and coalescence modulates shape of aged protein condensates," *Scientific reports*, vol. 12, no. 1, p. 4390, 2022.

[127] P. Arosio, M. Vendruscolo, C. M. Dobson, and T. P. Knowles, "Chemical kinetics for drug discovery to combat protein aggregation diseases," *Trends in pharmacological sciences*, vol. 35, no. 3, pp. 127–135, 2014.

[128] S. I. Cohen, M. Vendruscolo, C. M. Dobson, and T. P. Knowles, "Nucleated polymerization with secondary path-ways. ii. determination of self-consistent solutions to growth processes described by non-linear master equations," *The Journal of chemical physics*, vol. 135, no. 6, 2011.

[129] S. I. Cohen, M. Vendruscolo, C. M. Dobson, and T. P. Knowles, "From macroscopic measurements to microscopic mechanisms of protein aggregation," *Journal of molecular biology*, vol. 421, no. 2-3, pp. 160–171, 2012.

[130] C. Dammers, M. Schwarten, A. Buell, and D. Willbold, "Pyroglutamate-modified a$\beta$ (3-42) affects aggregation kinetics of a$\beta$ (1-42) by accelerating primary and secondary pathways," *Chemical science*, vol. 8, no. 7, pp. 4996–5004, 2017.

[131] K. Zhaliazka and D. Kurouski, "Nanoscale structural characterization of amyloid $\beta$ 1–42 oligomers and fibrils grown in the presence of fatty acids," *ACS Chemical Neuroscience*, vol. 15, no. 18, pp. 3344–3353, 2024.

[132] R. D. Yates and D. J. Goodman, *Probability and stochastic processes: a friendly introduction for electrical and computer engineers*. John Wiley & Sons, 2014.