Intensive transmission in wild, migratory birds drove rapid geographic dissemination and repeated spillovers of H5N1 into agriculture in North America

Lambodhar Damodaran[1], Anna Jaeger[1], Louise H. Moncla[1]

[1]Department of Pathobiology, School of Veterinary Medicine, University of Pennsylvania

Corresponding Author: Louise H. Moncla
Email: lhmoncla@upenn.edu

Abstract

Since late 2021, a panzootic of highly pathogenic H5N1 avian influenza virus has driven significant morbidity and mortality in wild birds, domestic poultry, and mammals. In North America, infections in novel avian and mammalian species suggest the potential for changing ecology and establishment of new animal reservoirs. Outbreaks among domestic birds have persisted despite aggressive culling, necessitating a re-examination of how these outbreaks were sparked and maintained. To recover how these viruses were introduced and disseminated in North America, we analyzed 1,818 Hemagglutinin (HA) gene sequences sampled from North American wild birds, domestic birds and mammals from November 2021-September 2023 using Bayesian phylodynamic approaches. Using HA, we infer that the North American panzootic was driven by ~8 independent introductions into North America via the Atlantic and Pacific Flyways, followed by rapid dissemination westward via wild, migratory birds. Transmission was primarily driven by Anseriformes, shorebirds, and Galliformes, while species such as songbirds, raptors, and owls mostly acted as dead-end hosts. Unlike the epizootic of 2015, outbreaks in domestic birds were driven by ~46-113 independent introductions from wild birds, with some onward transmission. Backyard birds were infected ~10 days earlier on average than birds in commercial poultry production settings, suggesting that they could act as "early warning signals" for transmission upticks in a given area. Our findings support wild birds as an emerging reservoir for HPAI transmission in North America and suggest continuous surveillance of wild Anseriformes and shorebirds as crucial for outbreak inference. Future prevention of agricultural outbreaks may require investment in strategies that reduce transmission at the wild bird/agriculture interface, and investigation of backyard birds as putative early warning signs.

Introduction

Highly pathogenic avian influenza (HPAI) viruses pose persistent challenges for human and animal health. Since emerging in 1996, highly pathogenic H5N1 viruses of the A/goose/Guangdong lineage have spread globally via endemic transmission among domestic birds in Asia and Africa coupled with long-distance dispersal by wild migrating birds (1,2). In 2005, introduction of poultry-derived H5N1 viruses into wild birds in China led to viral dispersal across Northern Africa and Asia, establishing new lineages of endemic circulation in poultry (3,4). In 2014, wild, migratory birds carried highly pathogenic H5N8 viruses from Europe to North America, sparking an outbreak that resulted in the culling of over 50.5 million commercial birds (5). While this outbreak substantially impacted the agriculture industry, aggressive culling quelled the outbreak, and North America remained free of HPAI for years.

Since December 2021, clade 2.3.4.4b HPAI H5N1 viruses have spread across the Americas, causing a panzootic of significant morbidity and mortality in wild and domestic animals. These viruses were likely first introduced into North America in late 2021 by migratory birds flying across the Arctic Circle from Europe (6,7), after which reassortment with endemic, low-pathogenicity avian influenza (LPAI) North American H5Nx (where Nx refers to various Neuraminidase (NA) subtypes) viruses produced a virus with altered tissue tropism in mammals (8). In contrast to past epizootics, morbidity and mortality has been widespread across a broad range of wild avian species not usually impacted by HPAI (9) such as raptors, owls, passerines (1,10), and Sandwich Terns (11–13). Infections have also occurred in mammal species not typically associated with HPAI, such as foxes, skunks, raccoons, harbor seals, dolphins, bears (1,14), and recently, domestic goats and dairy cattle (15). Putative transmission among marine mammals and domestic dairy cattle pose new challenges for animal health and biosecurity, and highlight the need to understand the ecological factors that lead to spillover (15,16).

Historically, H5N1 transmission has been linked to poultry production, with occasional cross-continental movement by wild birds of the Anseriformes (waterfowl such as ducks and geese) and Charadriiformes (Shorebirds) orders (17–19). Unlike the North American epizootic in 2014-2015, widespread culling of domestic birds has not halted detections in North America, suggesting that patterns of transmission since 2022 may be distinct from past epizootics. Prior work has posited that clade 2.3.4.4b viruses may be better able to infect and transmit among wild bird species, leading to persistent, seasonal circulation in European wild birds (11,13). Early genomic analysis of the United States epizootic linked outbreaks in poultry to wild birds, though the robustness of these results to differences in sampling between wild and domestic birds was not directly examined (10). Designing effective surveillance and intervention strategies hinges on delineating which species are driving transmission in North America, a

90   question that remains understudied. The broad range of affected wild species in this
91   panzootic raises the possibility that new reservoir hosts could be established,
92   necessitating an evaluation of which species should be actively surveilled. Finally, while
93   it is currently thought that cases in mammals likely stem from infections in wild birds,
94   work to formally link infections across species has been sparse.
95
Viral phylodynamic approaches are emerging as critical tools for outbreak
reconstruction (20,21). Viral genomes contain molecular records of transmission
histories, allowing them to be used to trace how outbreaks begin and spread. Here we
use Bayesian phylogeographic approaches paired with rigorous controls for sampling
bias (22,23) to trace how highly pathogenic H5N1 viruses were introduced and
disseminated across North America. We capitalize on a dataset of 1,818 Hemagglutinin
gene sequences sampled from North American birds and mammals in 2021-2023, and
curate additional metadata on geography, migratory flyways, domestic/wild status, host
taxonomic order, and migratory behavior to reconstruct transmission between these
groups. Using this dataset of HA sequences, we show that the epizootic in North
America was driven by ~ 8 independent introductions that descend from outbreaks in
Europe and Asia, though only a single introduction spread successfully across the
continent. The initial wave of H5N1 transmission spread from east to west by wild,
migratory birds between adjacent migratory flyways. Transmission from non-canonical
avian species like songbirds and owls was limited and resulted in dead-end
transmission chains, suggesting that these species are unlikely to establish as
reservoirs. Instead, transmission was primarily sustained by Anseriformes, shorebirds,
and Galliformes. In contrast to the outbreak in 2014/2015, outbreaks in agriculture were
seeded by ~46-113 independent introductions from wild birds, with some onward
transmission. Backyard birds were infected slightly more frequently, and earlier on
average, than commercial birds, suggesting that increased backyard bird surveillance
could serve as early warning signs for increased incidence. Together, these data
pinpoint surveillance in wild aquatic birds as critical for contextualizing outbreaks in
mammals and agriculture. Given the increasing role of wild bird transmission in North
America, investment in interventions that reduce interactions between domestic and
wild animals may now be crucial for limiting future outbreaks in agriculture.

Results

Viral sequence data capture seasonal variation of HPAI detections

Figure 1. Detections of HPAI in North America show distinct epidemic waves following introduction events in late 2021. A) Detections of HPAI in wild birds, domestic birds, and non-human mammals. B) The Log-scaled Effective population size (N_e) estimates estimated in BEAST using the Bayesian SkyGrid coalescent for sequences collected between Sep 2021 and Aug 2023. C) Correlation plot of log(N_e) vs HPAI detections by week, spearman correlation displayed.

In the United States, the United State Department of Agriculture Animal and Plant Health Inspection Service (APHIS) manages HPAI surveillance and testing in wild birds via investigation of reported morbidity and mortality events, hunter-harvested game birds/waterfowl, sentinel species/live bird collection, and environmental sampling of water bodies and surfaces (24,25). As of September 30th 2024, most HPAI detections have been reported in wild birds, which were sampled via testing of sick and dead (5,611), hunter harvested (3,340), and live wild birds (1,127) (Figure S1a). APHIS also surveilles domestic birds using several reporting methods: mandatory testing through the National Poultry Improvement Plan, coordination with state agricultural agencies,

routine testing in high-risk areas, and backyard flock surveillance (26). Data on
domestic bird detections are reported with information on poultry type (e.g., duck,
chicken) and by whether the farm is classified as a commercial operation or backyard
flock. Backyard flocks are categorized by the USDA as operations with fewer than 1,000
birds (27,28) and by the World Organization for Animal Health (WOAH) as any birds
kept in captivity for reasons other than for commercial production (29). Among domestic
birds, detections (1,177 total) came predominantly from commercial chickens (9.3%),
commercial turkeys (28.5%), commercial breeding operations (species unspecified)
(15.3%), and birds designated WOAH Non-Poultry which refers to backyard birds
(42.3%) (Figure S1b). Other domestic bird detections occurred in game bird raising
operations (2.5%) and commercial ducks (2.0%). This panzootic has notably impacted a
broad range of mammalian hosts, with detections (399) reported in red foxes (24.3%),
mice (24.1%), skunks (12.2%), and domestic cats (13.2%). Other mammalian hosts
(26.2%) represent a wide range of species including harbor seals, bobcats, fishers, and
bears (Figure S1c).
The first detection of HPAI H5N1 viruses in the United States occurred in a wild
American wigeon in South Carolina on December 30th, 2021. From January – May
2022, a wave of 2,510 total detections were reported across 43 states and 91 different
species (Figure 1). Following a lull in the summer, cases rose again in August 2022,
leading to a larger epizootic wave that lasted until March 2023, totaling 8,001 detections
across all 48 contiguous U.S. states and Alaska. Case detections peaked in the fall and
spring, coinciding roughly with seasonal migration timing for birds migrating between
North and South America (30,31). Seasonal case variation could arise due to seasonal
bird migration, fluctuating virus prevalence in wild birds, or fledging times of susceptible
chicks (32), though continued monitoring is necessary to determine whether these
patterns persist in future years.
Sequence data sampled in North America is heavily skewed toward the first 6
months of the outbreak, with 74% of all available sequences sampled from January-July
2022 (Figure S2). To evaluate whether sequence data reflect case detections, we
inferred the viral effective population size (Ne), a measure that approximates viral
incidence (21). Using 6 datasets of sequences subsampled by host taxonomic order
(see Methods for details), we infer that Ne is modestly correlated with detections
(highest Spearman rank correlation: 0.65, p=4.4e-11) (Figure 1c)(Figure S3-4), and that
peaks in Ne precede peaks in detections by ~1 week (Figure S5), likely reflecting the
lag between viral transmission and case detection. These data suggest that despite
uneven sequence acquisition across time, the diversity of sampled sequences reflect
the amplitude of H5N1 cases. We therefore opted to use sequence data for the entire
sampling period for broad inferences on introductions and geographic spread across
North America, but supplement these analyses with a series of controls for sampling
differences between groups. For more intensive reconstructions of transmission
patterns between wild birds, commercial poultry, and backyard birds we focus on the
initial 6-month period with the most densely sampled data, coupled with experiments to
assess the impacts of sampling on results.

200
201 *Wild, migratory birds drove expansion of H5N1 across the continent*
202
To reconstruct the number and timings of H5N1 introductions into North America, we
constructed a dataset that including HA sequences from North America from domestic
and wild birds (n= 1,327 unique isolates, identical sequences were removed), along with
contextual sequences from other continents with ongoing outbreaks in 2021 and 2022
(Asia = 294, Europe = 300). We then inferred the number and timings of introductions
into North America using a discrete trait phylogeographic model (see Methods for
details) (33).
210

211
Figure 2. Introductions from Europe facilitated the majority of ongoing transmission
while Asian introductions did not result in continent-wide onward transmission. A)
Bayesian phylogenetic reconstruction of n=1,921 globally sampled sequences of HPAI clade
2.3.4.4b colored by continent of isolation. Opacity of branches corresponds to posterior support
for the discrete trait inferred for a given branch, the thickness corresponds to the number of
descendent tips the given branch produces. B) A close-up view of the starred section of the tree
in A, focusing on introductions from Asia. C) We inferred the number of transitions from Asia to
North America across the posterior set of 9,000 trees. The x-axis represents the number of
introductions, and the y-axis represents the proportion of trees across the posterior set with that
number of inferred transitions.
222

Most sequenced infections in North America (98.5% of tips) descend from a single
introduction from Europe in late 2021 (95% Highest Posterior Density (HPD),
September 9th – October 7th 2021) (Figure 2a), consistent with reports of infected
migratory gulls in Newfoundland and Labrador Canada in November 2021, and
subsequent mortality in farmed birds in December of 2021 (6,7,10). Previous
surveillance of migratory birds traveling from Europe to North America in October of
2021 recorded detections of HPAI of European origin from wigeons, geese, skua, and
gulls (6). We also infer 7 (median = 7, 95% HPD: (6,8)) additional introductions between
February and September 2022 that nest within the diversity of viruses circulating in Asia
(Figure 2B-C). These introductions represent infections sampled in Alaska, Oregon,
California, Wyoming and British Columbia, Canada that failed to disseminate widely and
persisted for short periods of time (0.024 – 6.9 months). The western location of these
tips suggest potential introduction via the Pacific flyway, consistent with previous reports
documenting incursions into North America from Japan via Alaska and the upper Pacific
(34)(Figure S6).
238
Recent analyses of global H5N1 circulation patterns suggest wild birds as increasingly
important reservoirs for clade 2.3.4.4b virus evolution and transmission (35). In the
Americas, avian migratory routes are classified by the U.S. Fish and Wildlife Service
(USFWS) into 4 major flyways: the Atlantic, Mississippi, Central, and Pacific (36). If the
epizootic were spread predominantly by wild, migratory birds, we reasoned that viruses
sampled from the same, or neighboring flyways, should cluster together more closely
than viruses sampled from non-adjacent flyways. To test this, we assigned avian
sequences to the migratory flyway matching the US state of sampling, assembled a
dataset of 250 sequences randomly subsampled for each USFWS flyway (total
n=1,000), and implemented a discrete trait diffusion model to estimate transition rates
between flyways, a proxy for transmission.
250

251
Figure 3. Migratory birds rapidly disseminated H5N1 via migratory flyways. A)
Phylogenetic reconstruction of n=1,000 sequences colored by continental flyway. Inset is the
results of the PACT analysis for persistence in each flyway (how long a tip takes to leave its
sampled location going backwards on the tree), excluding the Pacific clade to show persistence
following Atlantic introduction. B) U.S. Fish and Wildlife Service waterfowl flyways map, with
arrows annotated to represent rates with Bayes factor support of at least 100. Here the size of
the arrow corresponds to the magnitude of the mean transition rate. C) Mean transition rates
from the Bayesian Stochastic Search Variable Selection (BSSVS) of USFWS flyways where
color of the bar corresponds to the source population and the opacity corresponds to the Bayes
Factor (BF) support (where white corresponds to BF < 3 and full color corresponds to BF >
100). D) Mean transition rates from the BSSVS of migratory behavior of birds.

Tips that descend from viruses circulating in Asia (those in Figure 2B) cluster together
as a basal clade inferred in the Pacific flyway (orange cluster at top of tree, posterior
probability = 0.98), consistent with introduction into the West Coast. The primary
introduction from Europe occurred via the Atlantic Flyway, and then spread rapidly
across the US (Figure 3A,B). From the inferred time of introduction in the Atlantic flyway
between September 9th – October 7th 2021, viruses descending from this introduction
had disseminated and been sampled in every major flyway within ~4.8 months.

Sequences clustered strongly by flyway, grouping most closely with others sampled
within the same or geographically adjacent flyway (Figure 3A). Transmission was most
efficient between adjacent flyways, and primarily proceeded from east to west (Figure
3B-C, Table S1). We calculate the Bayes Factor (BF) support for each possible
transition rate, which is calculated by dividing the posterior odds a transition rate is non-
zero by the equivalent prior odds. Generally, BF > 30 are considered strong support,
indicating that a given rate is 30x more likely to be included in the diffusion network, in
this analysis we primarily focus on BF >= 100, or 100x more likely to be included (see
Methods for details). We infer the highest supported rates (BF > 100) from the
Mississippi to Central flyway (4.69 transitions/year, 95% HPD: (1.67,7.87), Atlantic to
Mississippi flyway (2.3 transitions/year, 95% HPD: (0.74,4.08)), and Central to Pacific
flyway (1.93 transitions/year, 95% HPD: (0.64,3.48))(Figure 3C, Table S1). Though the
Pacific flyway experienced the highest number of introductions during the epizootic,
transitions from the Pacific flyway elsewhere were inferred with low magnitude and
weak support, and very little transmission occurred from west to east. Indeed, only a
single statically supported rate was inferred from the Pacific flyway to the adjacent
Central flyway (BF = 3, 0.69 transitions/year, 95% HPD: (0.004, 2.35)). Quantification of
the length of times that lineages persisted in each flyway showed slightly longer
persistence within the Atlantic and Pacific flyways, potentially due to the habitat and
species richness in each flyway allowing for greater interaction of hosts (37). Previous
work has shown that within-flyway transmission occurs far more rapidly than
transmission between flyways, which may occur over longer time spans (> 5 years) (38–
40). We speculate that the strong signal of east to west diffusion could be explained by
rapid, exponential spread among naïve wild birds in North America during early
panzootic expansion. Limited transmission from the Pacific flyway could also be
explained by differential fitness of the lineages introduced into the Pacific vs. Atlantic
flyways, ecological isolation of the Pacific flyway, or by differences in host distribution at
the time of incursion. Future work will be necessary to differentiate among these
hypotheses.
300
The strong clustering of sequences by flyways is consistent with long-range
transmission by wild, migratory birds, but is not a direct measurement of it. To directly
test this, we classified wild bird sequences by whether they were sampled from a wild
bird considered migratory, partially migratory, or sedentary using the AVONET database
(41). We then modeled discrete trait diffusion across 5 categories: wild migratory birds
(includes most ducks and geese), wild partially migratory birds (some ducks, raptors,
and vultures), wild sedentary birds (owls crows), domestic birds, and non-human
mammals. Consistent with the flyways analysis, wild, migratory birds are inferred across
the entire tree backbone with high statistical support, indicating that these birds played
an important role in sustained transmission and geographic dissemination (Figure S7).
Transitions from wild migratory birds were inferred with the highest number and most
strongly supported rates (BF > 3000), indicating that migrating wild birds were critical to
seeding infections in other species (Figure 3D, Table S2). Taken together, our results
show that HPAI H5N1 viruses were repeatedly introduced into North America
throughout the epizootic, with incursions into both the Atlantic and Pacific flyways.
Following introduction into the Atlantic flyway, H5N1 viruses were rapidly disseminated

from east to west by migrating wild birds. These results highlight the capacity of
migratory birds to rapidly transmit these viruses across vast geographic areas in North
America.
320
321 *Epizootic transmission is sustained by canonical host species*
322
Previous outbreaks of highly pathogenic H5N1 viruses have been facilitated by wild
Anseriformes (waterfowl), wild Charadriiformes (shorebirds), and domestic Galliformes
(42–45), though the role of these hosts varies across outbreaks. In the current
panzootic, die-offs have occurred across non-canonical host orders like Accipitriformes
(raptors, condors, vultures), Strigiformes (owls), and Passeriformes (sparrows, crows,
robins, etc.) (9), raising the possibility that these new species could establish as
reservoirs that merit surveillance. To determine whether particular host groups played
outsized roles in driving transmission in the epizootic, we classified sequences by
taxonomic orders that were most well-sampled and modeled transmission between
them using a discrete trait model. We consolidated sequences of two orders of raptors,
Accipitriformes and Falconiformes, hereafter referred to as "raptors". We also
consolidated two orders of pelagic birds, Charadriiformes and Pelecaniformes, hereafter
referred to as "shorebirds". Following classification, we defined 7 host order groups:
Anseriformes, Shorebirds, Strigiformes, Passeriformes, Raptors, Galliformes, and non-
human mammals.
338
Discrete trait approaches assume that the number of sequences in a dataset are
representative of the underlying distribution of cases in an outbreak, resulting in faulty
inference when this assumption is violated (23,33,46) and bias when groups are
unevenly sampled (22,23). To account for differential sampling among these host
groups, we therefore considered two, distinct subsampling approaches. The first is a
proportional sampling regime in which sequences are sampled proportional to the
detections in each host group each month. This common sampling regime assumes that
case detections in each group are the closest proxy for the case distribution in the
outbreak, and attempts to align sampling with underlying model assumptions. However,
this approach may not be appropriate if case detection is heavily biased between
groups. For HPAI H5N1 in North America, detections in wild birds are primarily identified
when humans report sick or dead birds to wildlife health authorities or wildlife rescues
(Figure S1A), which may skew detections towards birds with dedicated rescue services
or birds that reside in closer proximity to humans. For example, Anseriformes and
raptors comprised 50.2% and 20.3% of all sequences, respectively, which could arise
from high case intensity or a higher rate of case acquisition. A second, complementary
subsampling approach is to sample sequences equally, meaning that sequences are
sampled from each group in perfectly equal numbers. By forcing the number of
sequences from each group to be equal, the transmission inference must be driven by
the underlying sequence diversity in each group rather than by sampling differences.
Given the high variation among detections within each host group, we opted to pursue
an equal sampling regime. We performed 3 independent subsamples, each comprised
of a dataset of 100 randomly sampled sequences per host group. To account for
variation across subsampled datasets, we combined the results for the 3 independent

 subsamples to summarize statistical support (Figure S9, Table S3-4). Due to similar
tree topologies across replicates, we visualize the phylogeny of the subsample with the
highest posterior support (equal order subsample 1) below and make the results of all
subsamples available in supplement (see Figure S8, Table S5-S10).

Figure 4. Anseriformes drove outbreak transmission, while new host species represent dead-end infections. A) Bayesian phylogenetic reconstruction of n=655 sequences subsampled by host order with equal proportions of each host. The color of tips and branches represents taxonomic order, and opacity represents the posterior support for the inferred host group. Thickness of branches correspond to the number of tips descending from a given branch. B) Transition rates from the host group on the left (labelled "From") to the host on the right (labelled "To") as inferred from the combined results of three equal orders subsamples. The x location of the dot represents the inferred mean transition rate, and the black lines (whiskers) represent the 95% HPD. The color of each bar represents the "From" host. The opacity represents the Bayes factor support for the inclusion of the rate in the diffusion network. White (opacity of 0) represents any Bayes factors inferred to be less than 3, while a full color (opacity of 1) represents any Bayes factors inferred to be greater than or equal to 100. C) Results of the PACT analysis for persistence in each host order for phylogeny shown in panel A.

The first introduction into North America is comprised of infections sampled in great
black-backed gulls (inferred as a large clade of shorebirds, posterior probability = 0.69),
consistent with previous evidence of migratory gulls facilitating transmission from
Europe (Figure 4A). This inferred-shorebirds cluster contains 6 sequences from harbor
seals sampled from outbreaks in New England, resulting in a highly supported transition
rate (BF = 537, posterior probability= 0.99) from shorebirds to non-human mammals
(2.09 transitions/year, 95% HPD: (0.99, 4.63), aligning with suggestions that these
outbreaks are linked to scavenging or environmental contamination by infected shore-
birds (1,16). After this initial cluster of infections, the remainder of the phylogeny
backbone is inferred in Anseriformes with high posterior support (0.99), indicating that
Anseriformes played an important role in driving sustained transmission and dispersal
across North America. We infer Anseriformes as the predominant hosts seeding
infections into other species (Figure S10) (Figure 4B) (Table S5-10), with the highest
rates to Galliformes (4.49 transitions/year (95% HPD: (1.84, 7.21), BF = 1691, posterior
probability = 0.99) and Strigiformes (3.41 transitions/year (95% HPD: (1.24, 6.14), BF =
232, posterior probability = 0.99). Aligning with speculation following mortality events in
bald eagles (47), we also infer a highly supported transition rate (BF = 127, posterior
probability = 0.95) from Anseriformes to Raptors, consistent with putative links between
raptors and the waterfowl they predate. These patterns were preserved in each
independent subsample, indicating high robustness to sampling (Figure S8-9). We also
infer support for transmission from Galliformes to Anseriformes (2.47 transitions/year
(95% HPD: (0.45, 4.88), BF = 147, posterior probability = 0.96), Strigiformes, and
nonhuman mammals. In this dataset, Galliformes primarily represent domesticated
poultry (98% of sequences), suggesting that transmission from domestic birds back to
wild birds and mammals may also have occurred, a hypothesis we investigate in more
depth below. However, lineages in Galliformes tended to be short-lived, persisting for
0.26 years on average (95% HPD: 0.07, 0.33 years). In contrast, viral lineages persisted
for the longest in Anseriformes, with a mean persistence time of 0.71 years (95% HPD:
0.42, 0.88 years) (Figure 4C), and Shorebirds, with a mean persistence time of 0.654
years (95% HPD: 0.18, 1.04 years). Therefore, while Anseriformes, Shorebirds, and
Galliformes all contributed to transmission events to other species, longer-term
persistence was primarily driven by transmission in Anseriformes and Shorebirds.
416
One surprising result was that we inferred raptors as a strongly supported source
population to Anseriformes (1.87 transitions/year (95%HPD: (0.18, 3.94), BF = 39,
posterior probability = 0.87). Previous characterizations of HPAI in Raptors during the
2014/2015 outbreak in North America showed mortality events and neurological
symptoms in wild raptors (48). Serological evidence of infections in bald eagles have
indicated exposure to influenza A viruses in 5% of birds tested between 2006 and 2010
(49). In the ongoing panzootic, raptors represent the third most prevalent group in wild
bird detections in Europe (12% of detections) and second most detected group in North
America (20.3%) (13,50). Future work to establish whether the high number of cases
among raptors, and potential link to Anseriformes, is driven by efficient case detection
vs. changing patterns of viral transmission will be necessary for formulating wildlife
management strategies.
429

We found that Strigiformes (owls) exhibited primarily sink-like behavior with only two
supported source transition rates to Passeriformes (BF= 82, posterior probability =
0.93), and Anseriformes (BF=61, posterior probability = 0.91). Surveillance efforts show
evidence of transmission of HPAI in owls during previous outbreaks of HPAI globally
and have typically been sampled alongside other bird species across several different
host orders, primarily from the order Anseriformes (51–53). Overall, we found limited
support for the non-canonical host groups songbirds and nonhuman mammals seeding
infections in other species. Inference of transitions from these three host groups tended
to be less supported and lower in magnitude, suggesting that these species did not play
major roles in driving transmission across the continent or to other species (Figure 4B).
Instead, these species tended to act as sinks, forming short, terminal transmission
chains that did not lead to long-term persistence (Figure 4C and Figure S11). Similarly,
mammals served as sinks for viral diversity, supporting very short persistence times of
0.22 years (95%HPD:(0.088, 0.328)), and resulting in only one strongly supported
transition rate, to Anseriformes (BF = 53, posterior probability = 0.89). Finally, mammal
sequences cluster across the entire diversity of the phylogeny (Figure 4A), and are not
associated with one particular cluster of viruses, indicating that mammal infections were
not confined to a particular viral lineage. Instead, these findings are most compatible
with a model in which wild mammals are infected by direct interaction with wild birds,
likely related to scavenging and predation behaviors (14).
450
451 *Agricultural outbreaks were seeded by repeated introductions from wild birds*
452
Since 2022, the US has culled over 104.4 million domestic birds with agricultural losses
estimated between $2.5 to $3 billion USD (54). Understanding the degree to which
outbreaks in agriculture have been driven by repeated introductions from wild birds vs.
sustained transmission in agriculture is critical for improving surveillance and biosecurity
practices. However, differences in sampling between wild and domestic birds challenge
this goal. While domestic birds comprise 11% of all detections, they comprise 23.2% of
sequences, making them overrepresented in available sequence data. In contrast,
though a higher number of detections and sequences have been deposited for wild
birds, wild birds are likely to be heavily under sampled due to the challenge of sampling
wildlife (9,24). Finally, while each detection in wild birds represents a single infected
animal, domestic bird detections usually represent a single infected farm, where the true
number of infected animals is unknown. Given these challenges, we designed a
"titration" analysis to measure the impact of varying degrees of sampling on
transmission inference between wild and domestic birds. We first generated a dataset
composed of equal numbers of domestic and wild bird sequences (using all 270
available domestic bird sequences and 270 randomly sampled wild bird sequences)
sampled between November 2021 and August 2023. By setting the number of
sequences from each group equal, we force the inference to be driven by the sequence
data itself, rather than the sampling regime. Next, we added in progressively more wild
bird sequences until we reached a final ratio of domestic to wild sequences equal to 1:3,
which approximates the ratio of detections in domestic and wild birds during the
epizootic (1026 domestic detections vs. 3078 wild detections for study time period). In
total, we generated 5 datasets with the following ratios of domestic to wild bird

sequences: 1:1, 1:1.5,1:2,1:2.5, and 1:3 (see Methods for more details). For each
dataset, we applied a discrete trait diffusion model to infer transmission between wild
and domestic birds.
479

480
Figure 5. **Outbreaks in domestic birds were seeded by repeated introductions**
from wild birds, with some onward transmission A) Phylogenetic reconstruction
where taxa and branches colored by wild or domestic host status containing a 1:3 ratio
of domestic to wild bird sequences (n=1080). B) Number of transitions from a given trait
to another trait inferred through ancestral state reconstruction for each titration. C)
Results of the PACT analysis for persistence in domestic and wild birds for each
titration.
488
When domestic/wild sequences were included in equal proportions, the backbone of the
phylogeny and majority of internal nodes are inferred as wild birds, suggesting that wild
birds are inferred as the primary source in the outbreak regardless of sampling (Figure
S12A). Under equal sampling, this result is likely driven by higher genetic diversity
among viruses sampled from wild birds, consistent with a large, source population.
Within the background of wild bird sequences are multiple nested clusters of domestic
bird sequences, consistent with some transmission between domestic birds.
Transmission is inferred bi-directionally, with similar magnitudes of transmission inferred
from domestic to wild birds, and from wild to domestic birds (Figure 5B, Figure S13,

Table S11). If these patterns represent the true transmission history in the epizootic, we
reasoned that these patterns should remain intact even when additional wild bird
sequences are added to the tree. If not, then we hypothesized that some of domestic
bird clusters may be disrupted as more wild bird sequences are added into the tree.
502
As wild sequences were progressively added into the tree, most domestic-only clusters
became smaller, broken up by wild sequences that interspersed within these clades
(Figure S12A-E). The "breaking up" of these domestic clusters results in inference of
more transitions from wild to domestic birds, and less transmission between domestic
birds (Figure 5B-C, Figure S13). The phylogeny of the final dataset (using a 1:3 ratio of
domestic to wild sequences) shows 106 clusters of domestic sequences (Figure 5A,
Figure S14-15, Table S11) each inferred as a unique introduction of H5N1 from wild
birds into domestic birds. Among these domestic-only clusters, we calculate that
lineages persisted for ~4.5 months on average (95% HPD: 2.7, 5.63). In contrast, we
infer limited transmission from domestic birds back to wild birds (Figure 5B, Table S11),
with only 4 inferred introductions from domestic birds to wild birds (Table S11). Viral
lineages in wild birds also persisted for over twice as long as those in domestic birds,
persisting for ~10 months (95% HPD: 5.7, 14.07) (Figure 5C).
516
Commercial turkey operations have been heavily impacted during the epizootic,
comprising 53.7% of all detections on commercial farms (55). However, the presence of
wild turkeys throughout North America makes categorizing turkey sequences as
domestic or wild status ambiguous. 98% of all turkey sequences are not associated with
metadata on domestic/wild status, and thus were excluded from the previous analysis.
To determine whether this decision could have biased our results, we performed an
additional analysis in which any turkey sequence not labeled as "wild turkey" was
categorized as "domestic" and combined with the other domestic bird sequences. In this
dataset, ~20% of domestic bird sequences were turkeys. We then created a second
titration analysis using these domestic sequences to generate datasets of 1:1,1:1.5, and
1:2 ratios of domestic:wild sequences. We then evaluated the impact of the inclusion of
turkey sequences on inferred transmission patterns between wild and domestic birds.
529
Inclusion of turkey sequences did not substantially change inferred transition rates
between wild and domestic birds but did increase the inferred persistence times of
domestic bird clusters (Figure S13, Table S11). As wild bird sequences were added to
the tree, we observed the same "breakup" of domestic clades as in the above analysis,
with more wild sequences leading to more inferred introductions into domestic birds,
and very few into wild birds. In both titration experiments, the final number of inferred
transmission events from domestic to wild birds was 4 (Table S11), indicating minimal
transmission back to wild species, regardless of whether turkeys were included (Figure
S16, Table S11). Inclusion of turkey sequences resulted in slightly longer inferred
persistence times in domestic birds, increasing inferred persistence by 1.29 and 1.54
months in the 1:1.5 and 1:2 titrations (Figure S18). While turkey sequences tended to
cluster closely with other domestic sequences, they did form several large turkey-only
clusters on the tree (Figure S17), indicating some degree of separation between
commercial turkey and non-turkey operations. To more directly estimate the role turkeys

played in transmission we built a final 1:2 (domestic:wild) dataset where the number of turkey and domestic poultry where equal which conformed to both a uniform and case proportional dataset. Surprisingly, we found that while most introductions into turkey populations stemmed from wild birds (42 transitions), we infer a high number of transmission events between turkeys and other domestic birds. We estimate ~38 introductions from turkeys to other domestic birds, and 18 in the opposite direction (Figure S19, Table S12), indicating some degree of transmission between distinct poultry operation types. The high degree of transitions from wild birds to turkeys, and from turkeys to other domestic birds suggest a putative role for turkeys in mediating transmission between wild birds and other types of domestic poultry operations. In line with the high number of turkey detections during the epizootic, we infer comparatively larger clusters in turkeys than in other domestic birds, suggesting that turkeys may have played a key role in transmission among domestic populations.

Together, these data suggest a few important conclusions. First, wild birds are inferred strongly as the major drivers of transmission. Wild birds were inferred as the major source of viral dispersal regardless of sampling regime and independent of whether or not turkeys were included in the analysis, indicating strong support for their role in broad dissemination of H5N1. Second, regardless of sampling regime and presence/absence of turkey sequences, we infer that outbreaks in agricultural birds were driven by repeated, independent introductions from wild birds, with some onward transmission between domestic operations. While the exact number of inferred introductions vary across analyses (Table S11, S12), we infer no fewer than 46, and as many as 113 independent introductions into domestic birds. When allowing sampling frequencies to approximate detections (the 1:3 ratio analysis), we resolve a higher number of introductions into domestic birds with shorter transmission chains, though lineages still persisted for an estimated 4-6 months. Analysis of turkey and non-turkey data indicate some degree of transmission between agricultural operations, and a potential role for turkeys in mediating transmission between wild birds and non-turkey domestic birds. Together, these results indicate that while both the 2014/2015 epizootic and the 2022 epizootic involved transmission between domestic premises, that transmission since 2022 has been fundamentally distinct. While the epizootic of 2014/2015 was started by a small number of introductions that rapidly propagated between commercial operations, the epizootic since 2022 has been driven by intensive and persistent transmission among wild birds, resulting in continuous incursions into domestic bird populations that have continuously sparked new outbreaks in agriculture. These results suggest that wild birds may now play an increasing role in H5N1 transmission in North America, potentially necessitating updates to biosecurity, surveillance, and outbreak control.

Spillovers to backyard birds occur earlier and slightly more often than those to commercial birds

The 2014/2015 H5Nx epizootic in the United States was driven by extensive transmission in commercial poultry, prompting a series of biosecurity updates for commercial poultry farms (5,56). However, not all domestic birds are raised in

commercial settings. Rearing domesticated poultry in the home setting has become
increasingly popular in the United States, with an estimated 12 million Americans
owning "backyard birds" in 2022 (57). These birds have been heavily impacted during
the ongoing epizootic, with some evidence for distinct transmission chains circulating in
backyard birds vs. commercial poultry (10). Because backyard birds generally
experience less biosecurity than commercial birds and are more likely to be reared
outdoors (27), we hypothesized that spillovers into backyard birds may be more
frequent than spillovers directly into commercial poultry.
598
To test this hypothesis, we used a subset of sequences sampled between January and
May of 2022 that contained additional metadata specifying whether they were collected
from commercial poultry or from backyard birds (n= 275 from commercial poultry, n=85
from backyard birds). We then built a tree that included an approximately equal number
of sequences sampled from domestic and wild birds, but in which the domestic
sequences were evenly split between commercial poultry and backyard birds
(commercial birds = 85, backyard bird = 85, wild birds=193). As with the previous
analysis, we infer wild birds as the primary source population and infer repeated
introductions into commercial and backyard birds (Figure S20A). Unexpectedly though,
backyard bird sequences appeared to cluster more basally than commercial poultry
sequences, and sometimes fell directly ancestral to clusters of commercial poultry
sequences (Figure S20A). While every introduction into backyard birds was inferred to
descend from wild birds, 10 out of 26 introductions into commercial poultry were inferred
to descend from backyard birds (Figure S21A). This pattern was reproducible across
multiple replicate subsamples, indicating that it was independent of the exact subset of
wild bird sequences included in the tree. We developed two hypotheses that could
explain this pattern. The first is that backyard birds "mediated" transmission between
wild birds and commercial birds, possibly due to their greater likelihood of outdoor
rearing. Under this model, spillovers into backyard birds could be spread to commercial
populations via shared personnel, clothing, or equipment, resulting in sequences from
backyard birds preferentially nesting between wild and commercial bird sequences on
the tree. Alternatively, backyard birds could have simply been infected earlier on
average than commercial birds. If transmission in wild birds is persistent and high, and
backyard birds have a higher risk of exposure due to lessened biosecurity and
increased interactions with wildlife, then it could take less time for a successful spillover
event to occur and be detected in these birds, resulting in clustering that is more basal
in the tree.
626
To differentiate between these hypotheses, we performed a second titration analysis.
We started with the phylogeny including equal numbers of sequences from commercial
and backyard birds, allowing us to directly compare introduction patterns in these two
groups. Then, we added in progressively more wild bird sequences into the tree until all
available wild bird sequences were added into the tree. We added sequences in
increments of 25% (where % refers to percentage of total available wild bird sequences
in the time period), resulting in 3 additional analyses that included 50%, 75%, and 100%
of all available wild bird sequences. For example, the final dataset included 942
sequences, comprising 85 commercial bird, 85 backyard bird, and 772 wild bird

sequences. For each dataset, we inferred the number and timings of transmission
events between wild birds, commercial birds, and backyard birds across the posterior
set of trees. If backyard birds acted as mediators to outbreaks in commercial birds
(hypothesis 1), then the relationship between backyard birds and commercial birds
should remain unchanged as more wild bird sequences are added into the tree.
Alternatively, if backyard birds and commercial birds were infected independently
(hypothesis 2), then additional wild bird sequences should disrupt these clusters, and
intersperse between commercial and backyard bird sequences, resulting in more
independent introductions that occur earlier in backyard birds.

646
Figure 6. Backyard birds are infected by wild birds earlier than commercial birds.
A) Phylogenetic reconstruction of sequences collected between Jan 2022 and May
2023 with all available wild bird sequences and equal proportions of commercial and
backyard birds (n= 942) where taxa and branches are colored by host domesticity
status. B) Exploded tree view of the phylogeny showing the branches of transmission in
each domestic bird type following transmission from wild birds where subtrees represent

the traversal of a tree from the root to the tip where the state is unchanged from the
initial state (given by the large dot on left) to the tips represented by the smaller dots
representing continuous chains of transmission within a given state. C) Proportion of
trees from the posterior tree set with a given number of transitions from wild birds to
backyard birds and commercial birds (100% available wild sequences). D) Markov
rewards trunk proportion for domesticity status showing the waiting time for a given
status across branches of the phylogeny over time. E) Cumulative Markov jumps from a
given bird type to another over time where each line represents a single phylogeny from
the posterior sample of trees.
662
Throughout the experiment, wild bird sequences attached throughout the phylogeny,
disrupting nearly every backyard bird-commercial bird cluster, and dissolving the signal
of backyard bird to commercial bird transmission originally observed (Figure S20). The
final tree that included all available wild bird sequences resulted in inference of ~82
independent introductions from wild birds to domestic birds, with most clusters
containing only commercial (39 clusters) or backyard bird (43 clusters) sequences
(Figure 6A-B, Figure S20-21), suggesting that outbreaks in these groups were likely
seeded independently. Indeed, of the initial 10 transmission events inferred from
backyard birds to commercial birds, only 2 remained undisturbed in the final tree (Figure
6B, Figure S21). These two events represent outbreaks that occurred in the same state
within 6 days of each other, so it is plausible that these outbreaks are directly linked.
However, all other clusters were disrupted. As wild bird sequences were added into the
tree, the number of inferred introductions into backyard birds and commercial birds
diverged across the posterior trees for each titration (Figure S22), with backyard birds
experiencing slightly more introductions (mean = 43 introductions, 95% HPD: (35, 49))
than commercial poultry (mean = 39 introductions, 95% HPD: (32, 44)) (Figure 6C). This
is consistent with the rate of transmission inferred from wild birds into each domestic
population with a slightly higher rate to backyard birds (1.83 transitions/year, 95% HPD:
(0.282, 3.79), BF = 18,000, posterior probability = 0.99), then to commercial birds (1.6
transitions/year, 95% HPD: (0.23, 3.36), BF = 18,000, posterior probability = 0.99)
(Table S13-S15). We infer a low transition rate from backyard birds to commercial birds
(0.64 transitions/year (95% HPD: 3.1E-06, 2.05, BF = 4.58, posterior probability = 0.7),
and transitions from domestic birds back to wild birds were not statistically well-
supported.
687
To directly determine whether spillovers into backyard birds occurred earlier than those
into commercial poultry, we estimated the number of transitions between hosts across
the phylogeny ("Markov jumps") and the amount of time that is spent in each host
between transitions ("Markov rewards") (58,59). We infer the highest mean duration in
wild birds, representing 87.7% of Markov rewards. Backyard bird and commercial bird
sequences showed lower, and similar reward time percentages of 5.3% and 7.0%
respectively. Calculation of the Markov reward trunk proportion, a proxy for when
transmission occurred in each group, showed that early in the epizootic, transmission in
backyard birds preceded transmission in commercial poultry (Figure 6D, Figure S21).
Enumeration of the cumulative number of transitions between hosts over time across
the posterior set of trees (Markov jumps (59)), again revealed that backyard birds

experienced slightly more jumps than commercial poultry (backyard birds = 43
introductions, 95% HPD: 36, 50; commercial birds = 39 introductions, 95% HPD: 32,
44), and that these introductions occurred earlier on average (Figure 6E). The lag time
between the cumulative transitions for backyard birds and commercial birds was ~9.6
days, indicating that backyard birds may have been infected ~9 days earlier than
commercial birds. Comparison of detections and sequence availability in commercial
birds vs backyard birds show no apparent skewing in availability of samples for each
group (Figure S23). Taken together, these data confirm that in the first 6 months of the
epizootic, outbreaks in backyard bird and commercial bird populations were generally
seeded independently, with limited evidence for transmission between them. Spillovers
into backyard birds occurred slightly more frequently and about 9 days earlier on
average than spillovers into commercial poultry, suggesting that backyard bird
populations could potentially act as early warning signals for upticks in transmission.
Given the increasing role of wild birds in H5N1 transmission, backyard birds could
potentially act as useful sentinel species for gauging increasing transmission in wildlife
and increased risk of spillovers into agriculture. Future work will be necessary to
investigate the utility of this hypothesis.
716
717
718 **Discussion**
719
The 2022 panzootic of HPAI H5N1 has impacted wildlife health, agriculture, and human
pandemic risk across the Americas. This panzootic has been distinguished by the high
number of infections in wildlife not usually impacted by HPAI, its rapid dissemination
across the Americas, and for its persistence despite aggressive culling of domestic
birds. In this study, we used a dataset of 1,818 HA sequences paired with curated
metadata to reconstruct how H5N1 viruses were introduced and spread throughout
North America. We show that H5N1 viruses were introduced ~8 independent times into
North America, with repeated incursions into the Pacific flyway that failed to
disseminate. A single introduction into the Atlantic flyway spread across the continent
within ~4.8 months, transmitting from east to west by wild, migratory birds. Long-range
dispersal and persistence were primarily driven by transmission in Anseriforme species,
with infections in wild birds like songbirds and owls primarily serving as dead-end hosts.
Finally, we show through repeated subsampling experiments that unlike the epizootic of
2014/2015, outbreaks in agriculture were driven by repeated, independent introductions
by wild birds, with some onward transmission. Backyard birds experienced slightly more
introductions than commercial poultry, and these introductions occurred ~9 days earlier
on average. Taken together, our results highlight continuous genomic surveillance in
wildlife as critical for accurate outbreak reconstruction, and suggest wild, migratory birds
as emerging reservoirs for highly pathogenic H5N1 viruses in North America. Our
findings implicate surveillance in these wild, aquatic species as critical for ongoing
tracking and response, and suggest that preventing future outbreaks in agriculture may
now require layered interventions beyond culling.
742
743 We infer that most epizootic transmission descends from a single introduction into the
744 Atlantic flyway in the fall of 2021 (September-October), in line with work describing the

first confirmed detections on an exhibition farm in Newfoundland and Labrador Canada (6) and prior work that identified introductions into North America from Europe and Asia (7,60). Though H5N1 was first confirmed in captive birds in early December (December 9, 2021), retrospective testing revealed infection in a great black-backed gull from a nearby pond on November 26, 2021, confirming circulation in wild birds prior to detection on the farm (6). Our analyses suggest that highly pathogenic H5N1 viruses may have been circulating in wild birds in North America as early as September – October of 2021, one to two months prior to the first reported detection. Detailed analysis of species-specific migration patterns have been used to suggest that H5N1 may have been introduced in the autumn migration (6), a hypothesis supported by our data. Clade 2.3.4.4b viruses cause varying degrees of symptoms across avian species, so circulation in birds that experience milder symptoms could have obscured early detection and allowed for some degree of cryptic transmission (61). Species such as herring gulls show obvious neurological deficits such as paralysis, while other species such as Eurasian teals show no detectable clinical signs (11,62). Following introduction into the Atlantic flyway, we estimate that H5N1 viruses were transmitted from east to west, spreading from coast to coast in ~4.8 months. Compared to other avian-transmitted viruses, this is quite rapid. For example, West Nile Virus spread from the Northeast across the continent over the course of 4 years (63). We speculate that the relative rapidity of geographic spread could be explained by one or a combination of the following factors: high inherent transmissibility of clade 2.3.4.4b viruses in wild birds; rapid, long-range migration of wild bird species driving transmission; and rapid expansion across an immunologically naïve population in North America (61). Though limited work has been published on seroprevalence in wild birds, exposure to the current clade 2.3.4.4b viruses in North American wildlife was likely extremely limited prior to 2022 (64), potentially allowing for rapid, exponential growth following a successful incursion. Future work to examine the impacts of prior exposure to endemic, low-pathogenicity H5 viruses to susceptibility to clade 2.3.4.4b viruses may provide insights into future patterns of spread. Continuous surveillance among wild birds will also inform whether the rapid degree of transmission observed early in the panzootic will lessen over time as immunity builds in avian species.

Though a single introduction into the Atlantic flyway accounted for most transmission in the epizootic, the Pacific flyway experienced more independent incursions (~7), and it remains unclear why these failed to disseminate onward. Compared to other studies, we infer more incursions (5 more than previous estimates) into the Pacific flyway, likely due to the higher number of sequences from the Pacific region included in our analysis (65). Early after introduction, the European lineage introduced into the Atlantic flyway reassorted with endemic, low-pathogenicity H5 viruses in North America, resulting in a virus that exhibited altered tissue tropism in mammals (66). Whether this early reassortment event, or subsequent reassortments, also resulted in viruses with enhanced fitness in wild birds is currently unknown, but could explain the success of the Atlantic flyway introduction. Alternatively, the failure of Pacific incursions to spread onward could be explained by ecological isolation of the Pacific flyway, potentially due to land features like the Rocky Mountains. Prior studies of low-pathogenicity avian influenza have shown limited transmission between flyways and strong influence of

geography on the restriction of virus dispersal across the North American continent (67–69). In line with this hypothesis, we infer very minimal transmission occurring out of the Pacific flyway, even when inference was performed with an equal number of sequences per flyway. Finally, Pacific flyway introductions could have failed to disseminate due to a lack of suitable host species at the times and locations of the incursions, or simply by chance. Future work that examines the relative fitness of reassortants in wild birds, and that incorporates migratory movement data with the distribution of suitable species across time and space, will likely be necessary to distinguish between these hypotheses.

Our work demonstrates the impact of migratory bird movement on the rapid spread across the North American continent. In North America, transmission was most efficient between neighboring flyways, suggesting strong dissemination through geographic space that correlates to flyway areas. Prior work mapping migratory movements of Anseriformes (specifically, Mallards, Northern pintails, American Green-winged Teals, and Canada geese) showed that these birds exhibited North-South migratory pathways with some overlap between flyways (38,70). Future work to characterize the impact of migration in the western hemisphere is critically important as the range of many migratory birds encompasses several regions which are poorly sampled such as the Caribbean, Central America, and the Amazon basin. Follow-up studies to link finer-scale movement of wild birds to avian influenza transmission pathways may improve resolution of geographic spread and risk modeling, particularly for agricultural areas. Portions of the Atlantic and Mississippi flyways encompass high-density production areas for broiler chickens and turkeys, collectively producing nearly 7.32 billion broiler chickens and 182.2 million turkeys in the US annually (71), areas that have been heavily impacted by the panzootic. Our analysis of viral persistence times showed that viral lineages persisted for the shortest periods of time in the Mississippi and Central flyways and in Galliformes. Combined with our analyses showing that most agricultural outbreaks were driven by repeated introductions from wild birds, these findings suggest that infections in these domestic populations were generally transient. Our data suggest future work to track the movement of highly pathogenic H5 viruses across annual migratory cycles in wild birds as critical for pinpointing time periods that pose the greatest risk for spillover into agricultural settings. Successfully integrating migration timings of major species of Anseriformes, such as mallards, into risk models could allow biosecurity updates to be proactively deployed at times of year predicted to be critical based on time and geography (72,73).

Despite widespread infections in non-canonical avian species, we show that long-range, persistent transmission in this epizootic was driven by classical hosts for avian influenza: Anseriformes. Recent modeling of HPAI risk in Europe identified *Anatinae* and *Anserinae* (within the order Anseriformes) species prevalence as the most consistent predictors of HPAI detection across seasons (74), and future work will be necessary to determine whether those patterns hold in the Americas. Anseriformes and Shorebirds are primarily thought to inhabit wetland and shore habitats respectively but have been observed co-mingling in human-made habitats, like urban environments and waste disposal sites (75,76). We observe supported source transmission patterns from

raptors, birds which have previously shown susceptibility to clade 2.3.4.4 viruses and exposure to avian influenza viruses more generally (48,49). The severity and exposure risk in bird species in North America may also impacted by LPAI circulation which has been consistent and widespread for decades (75,77,78). The co-circulation of HPAI and LPAI in canonical host species can lead to novel reassortants which increase the pathogenicity of the virus. While this study focused on hemagglutinin sequences, future analysis should also include other gene segments and in particular focus on the reassortment of the virus between different NA subtypes as well as with LPAI viruses. The role of reassortment in the greater North American outbreak is still understudied and could provide important insights into transmission dynamics across different species.

In this study, we find that outbreaks in agriculture were seeded by repeated introductions from wild birds. This pattern held true regardless of sampling regime, and aligns with anecdotal observations that clade 2.3.4.4b viruses are increasingly being maintained by transmission among wild bird species, including now in North America (35,79). These findings contrast with genomic and epidemiologic investigation of the epizootic in 2014/2015, which implicated transmission between commercial poultry operations as the major source of dissemination (5,80). During that epizootic, transmission between farms was putatively linked to virus movement via personnel, equipment, and clothing, prompting updates to recommended biosecurity protocols for large, commercial operations (5,56). That epizootic was also well-controlled by culling domestic flocks, and after culling 50.5 million domestic birds, the epizootic died out. In contrast, at the time of writing, detections in wild and domestic birds in North America have continued despite culling nearly 104.4 million domestic birds. We show that despite some persistence among domestic bird populations, that introductions into domestic populations ultimately led to transmission chains that died out, and that they generally did not result in re-introduction into wild birds. Though detailed epidemiologic analyses are necessary to pinpoint the precise series of events that led to these outbreaks, our data imply that transmission in agricultural settings was fundamentally distinct from past outbreaks. Multiple, independent analyses suggest that transmission was efficient and persistent in wild birds, leading to repeated spillovers into agriculture. Taken together, our findings implicate wild birds as an emerging reservoir for highly pathogenic H5N1 viruses in North America. Continuous transmission in wild birds suggests a plausible explanation for rapid cross-continental spread, and for the lack of control in agriculture despite aggressive culling. In combination with other studies, our data suggest that future prevention of agricultural outbreaks may now require layered interventions that seek to reduce interactions between wild and domestic birds, paired with aggressive biosecurity between farms (18,81). Continuous reseeding from wild birds may also reduce the effectiveness of culling as the primary strategy for control. As highly pathogenic H5N1 viruses continue to circulate in North American wild birds, investment in control methods that reduce successful transmission between wild birds and agricultural animals, including potentially vaccination, should be explored.

We find that spillovers into backyard birds occurred slightly earlier and more frequently than those into commercial farms. Though commercial poultry operations generally

operate with higher degrees of biosecurity than backyard flocks, there are some
documented locations that could allow for interaction between domestic birds and
wildlife (27). Retention ponds on commercial poultry farms are frequently visited by wild
waterfowl (82), while natural features such as water bodies and vegetation near
residential coops and commercial production sites could also act as potential points of
wild to domestic transmission. Though recent data on backyard bird rearing in the US is
limited, a large survey of backyard bird populations from 2004 showed that backyard
bird flocks often contain multiple species, usually have outdoor access, and that 60-75%
of backyard flocks experience regular contact with wild birds (27). 38% of surveyed
backyard flocks were on a property that contained a pond that attracted wild waterfowl,
providing a clear, plausible link to water bird interaction. The same survey showed that
biosecurity precautions tend to be much more limited in backyard bird populations, with
88% of backyard flocks using no precautions (shoe covers, footbaths, clothing changes)
at all (27). Analysis of seropositivity to avian influenza in backyard poultry in Maryland
showed that exposure to waterfowl resulted in a ~3x higher likelihood of harboring anti-
influenza antibodies compared to birds not exposed to waterfowl (83). Early infections
during the 2014/2015 epizootic were recorded in backyard birds (84), and backyard
birds have been heavily impacted during this epizootic. Because backyard birds are
typically reared for egg and meat consumption, infections in backyard birds could also
pose a risk for human exposure. Putative links between infected locally reared birds and
human infections have been made in Mali and Egypt where backyard flocks are
common, serving as an indicator for the potential for exposure to occur in North America
(85,86). Given backyard birds' enhanced likelihood of interaction with wild birds, and our
inference of earlier spillovers in those groups, backyard bird populations could be
investigated as early warning sentinels for increasing transmission in local wild bird
populations. 93% of backyard flocks contain <100 birds, and most backyard bird owners
report raising birds for enjoyment and eggs (27). A subset of backyard bird owners may
therefore present an opportunity for community engagement that could potentially help
identify early detections. If engagement were successful, prompt reporting for illness
and infections could be used to alert other nearby operations that highly pathogenic
avian influenza is circulating locally, allowing for more advanced warning for heightened
biosecurity.
915
Sampling bias is pervasive across viral outbreak datasets, and no modeling approach
can completely overcome inherent biases in data acquisition. In this study, we opted to
explore multiple sampling regimes and pair them with experiments to directly assess the
impacts of sampling on our results. By combining case proportional subsampling and
equal sampling regimes with titration experiments, we attempt to evaluate the biases
between case detection and sequence availability that exist in these data, and to report
conclusions that were robust to these biases. Even so, an important caveat of our work
is that our inferences are limited by the availability of sequence data, and results could
change if future data become available. Our results also highlight the necessity of
consistent surveillance data from wildlife species. The inferences we make in this study
rely on intensive sampling of wild birds from the same time period as agricultural
outbreaks. Accurately distinguishing between hypotheses of epizootic spread (e.g.,
whether agricultural outbreaks are driven by introductions from wild birds or by from

farm-to-farm spread) depends on adequate sequence data from wild birds, without
which transmission inference is impossible. In our examination of avian host groups,
fine-scale analysis of individual species was limited by data availability, necessitating
grouping birds into higher level taxonomic orders. As H5N1 viruses continue to evolve
and spread globally, investment in surveillance strategies that capture circulating
diversity among wild birds will likely be critical for accurately tracking viral evolution,
prioritizing vaccine strains, and contextualizing new emergence events, like the recent
outbreaks in dairy cattle.
937
The H5N1 panzootic of 2022 has severely impacted wildlife ecology, agriculture, and
human pandemic risk across the Americas. Developing and deploying successful
interventions requires disentangling how these viruses spread across the Americas and
spilled into new species. Our findings point to wild, migratory birds as emerging
reservoirs for highly pathogenic H5N1 viruses in North America, a fundamental shift in
ecology with implications for biosecurity and pandemic risk mitigation. Our results
highlight the utility of wild bird surveillance for accurately distinguishing hypotheses of
epizootic spread, and suggest continuous surveillance as critical for preventing and
dissecting future outbreaks. Our data underscore that continued establishment of H5N1
in North American wildlife may now necessitate a shift in risk management and
mitigation, with interventions focused on reducing risk within the context of endemic
circulation in wild birds. At the time of writing, outbreaks in dairy cattle highlight the
critical importance of modeling the ecological interactions within and between wild birds
and domestic production. Future work to effectively model viral evolution and spread
hinges critically on effective surveillance across wild and domestic species to capture
key transmission pathways across large geographic scales. Ultimately, these data are
essential for informing biosecurity, outbreak response, and vaccine strain selection.
955
956
957 **Materials and Methods**
958
959 <u>Dataset collection and processing</u>
960
961 *Genomic data processing and initial phylogenetics*
962
We downloaded all available nucleotide sequence data and associated meta-data for
the Hemagglutinin protein of all HPAI clade 2.3.4.4b H5Nx viruses from the GISAID
database on 2023-11-25 (87). For each subset of the data described for further
phylodynamic modeling the following process was followed. We first aligned sequences
using MAFFT v7.5.20, sequence alignments were visually inspected using Geneious
and sequences causing significant gaps were removed and nucleotides before the start
codon and after the stop codon were removed (88,89). We de-duplicated identical
sequences collected on the same day (retaining identical sequences that occurred on
different days). We identified and removed temporal outliers for all genomic datasets by
performing initial phylogenetic reconstruction in a maximum likelihood framework using
IQtree v.1.6.12 and used the program TimeTree v 0.11.2 was used to remove temporal
outliers and to assess the clockliness of the dataset prior to Bayesian phylogenetic

reconstruction (90,91). This resulted in a dataset of 1818 sequences that were used in
further analyses (Figure S24).
977
978 *AVONET database*
979
We downloaded the AVONET database for avian ecology data and merged it to
available host metadata from GISAID for each sequence (41). We used the species if
provided to match the species indicated in the AVONET database. If host metadata in
GISAID was defined using common name for a bird, we determined the taxonomic
species name and used that for further merging with the AVONET data (e.g. "Mallard"
was replaced with *Anas platyrhynchos*) for the given region to match the species to its
respective ecological data. We additionally determined the domesticity status of a given
using the metadata provided.
988
989 *Detections by USDA*
990
991 Data for detections of HPAI in North America were collected from USDA APHIS.
992 Reports for mammals, wild birds, and domestic poultry were all downloaded (download
993 date: 2023-11-25) (50).
994
995 Phylodynamic analysis
996
997 The following Bayesian phylogenetic reconstructions and analyses were performed
998 using BEAST v.1.10.4 (92).
999
1000 *Empirical tree set estimation and coalescent analysis.*
1001
We performed Bayesian phylogenetic reconstruction for each dataset prior to discrete
trait diffusion modeling to estimate a posterior set of empirical trees. The following priors
and settings were used for each subset of the sequence data. We used the HKY
nucleotide substitution model with gamma-distributed rate variation among sites and
lognormal relaxed molecular clock model (93,94). The Bayesian SkyGrid coalescent
was used with the number of grid points corresponding to the number of weeks between
the earliest and latest collected sample (e.g for a dataset collected between 2021-11-04
and 2023-08-11 we would set 92 grid points) (95). We initially ran four independent
MCMC chains with a chain-length of 100 million states logging every 10000 states. We
diagnosed the combined results of the independent runs diagnosed Tracer v1.7.2. to
ensure adequate ESS (ESS > 200) and reasonable estimates for parameters (92). If
ESS was inadequate additional independent MCMC runs were run increasing chain
length to 150 million states, sampling every 15000 states were performed. We
combined the tree files from each independent MCMC run removing 10-30% burn-in
and resampling to get a tree file with between 9000 and 10000 posterior trees using
Logcombiner v1.10.4. A posterior sample of 500 trees was extracted and used as
empirical tree sets in discrete trait diffusion modeling.
1019
1020 *Discrete trait diffusion analysis*

1021

1022 *Dataset subsampling and definition of discrete traits*

1023

1024 We characterized the geographic introduction of HPAI into North America by randomly
1025 sampling 100 sequences from Europe and Asia for each year between 2021-2023 (total
1026 300 non-North American) and all available North American sequences across the study
1027 period. Following removal of temporal outliers this resulted in a dataset of n= 1921
1028 sequences annotated by continent of origin.

1029

To characterize geographic transmission within North America, following introduction,
we constructed a dataset of sequences subsampled based on migratory flyway. We
used place of isolation data to match the US state or Canadian province the sequence
was collected from with the respective U.S. Fish and Wildlife Service Migratory Bird
Program Administrative Flyway (36). We subsampled 250 sequences for each flyway
(Atlantic, Mississippi, Central, and Pacific) to create a dataset of 1000 sequences
collected between November 2021 and August 2023.

1037

We classified sequences by host taxonomic order, inferring the host species using
designations in the strain name and/or metadata to match species records in AVONET
(41). To ensure that each discrete trait had an adequate number of samples for the
discrete trait analysis of host orders we combined orders in two instances based on
taxonomic and behavioral similarity. The order Falconiformes (n=14), which represents
falcons, was added to Accipitriformes (n=363), which includes other raptors such as
eagles, hawks, and vultures. Pelecaniformes (n=34) which includes pelicans were
grouped with Charadriiformes (n = 74, shorebirds and waders) due to their similar
aquatic lifestyles and behaviors. Mammals were kept as a broad non-human
classification as most samples were of the order carnivora (foxes, skunks, bobcats etc.),
apart from samples of dolphins (Artiodactyla) and Virginia opossum (Didelphimorphia).
The following orders were omitted due to low number of sequences: Rheaforimes
(n=2), Casuariiformes (n=1), Apodiformes (n=2), Suliformes (n=7), Gaviiformes (n=1),
Gruiformes (n=1), Podicipediformes (n=1).

1052

We randomly subsampled 100 sequences for each host order between 2021-11-04 and
2023-08-11 resulting in a dataset of n=655 sequences where all isolates for host orders
with less than 100 samples, Passeriformes (n = 57) and Strigiformes (n=99) (removing
one temporal outlier), were used (Figure S25). We repeated this random subsampling
three times resulting in three separate datasets. We additionally performed three sub-
samples of sequences based on the proportion of detections in each host order group
which were collected between 2021-11-04 and 2023-08-11. Three random proportional
samples were taken each with the following number of sequences for each group:
Accipitriformes =133, Anseriformes = 342, Passeriformes = 12, Nonhuman-mammal =
16, Galliformes = 83, Charadriiformes = 40, Strigiformes = 29 (total n=655 sequences).

1063

We defined discrete traits for use in discrete trait diffusion modeling based on the
available sequence metadata and merged AVONET data. In addition to taxonomic
order, we defined migratory behavior. Birds were classified as sedentary (staying in

each location and not showing any major migration behavior), Partially migratory (e.g.
small proportion of population migrates long distances, or population undergoes short-
distance migration, nomadic movements, distinct altitudinal migration, etc.), or Migratory
(majority of population undertakes long-distance migration).
1071
1072 *Discrete trait modeling framework*
1073
For each discrete trait dataset, we used an asymmetric continuous time Markov chain
discrete trait diffusion model and implemented the Bayesian stochastic search variable
selection (BSSVS) to determine the most parsimonious diffusion network (33). We
inferred the history of changes from a given trait to another across branches of the
phylogeny, providing a rate of transitions from A to B/year for each pair of trait states.
When reporting these results, we refer to state A as the source population/state and B
as the sink population/state. We implemented the Bayesian Stochastic Search Variable
Selection which allows us to determine which rates have the highest posterior support
by using a stochastic binary operator which turns on and off rates to determine their
contribution to the diffusion network. For each pairwise transition rate, we calculate the
level of Bayes Factor (BF) support that the given rate has. The BF represents the
support of a given rate. The BF is calculated as the ratio of the posterior odds of the
given rate being non-zero divided by the equivalent prior odds which is set as a Poisson
prior with a 50% prior probability on the minimal number of rates possible (33). We use
the support definitions by Kass and Rafferty to interpret the BF support where a BF > 3
indicates little support, a BF between 3 and 10 indicates substantial support, a BF
between 10 and 100 indicates strong support, and a BF greater than 100 indicates very
strong support (96). Empirical sets were used with the discrete traits defined for each
sequence to perform discrete trait diffusion modeling. Each discrete trait model was
implemented using three independent MCMC chains with a chain length of 10 million
states, logging every 1000 states. Runs were combined using LogCombiner v.1.10.4.
subsampling a posterior sample of 10,000 trees/states. The Bayes Factor support for
transition rates were calculated using the program SPREAD3 (97). Maximum clade
credibility trees were constructed using TreeAnnotator v1.10.4.
1098
1099 *Domestic/Wild titration analysis*
1100
To study the impact of sampling of wild birds on the estimation of rates between
domestic and wild birds we created five separate datasets with varying numbers of wild
birds for sequences collected between 2021 and 2023. We randomly sampled 270
domestic sequences and 270 wild sequences as the initial 1:1 ratio dataset. We then
made four more datasets increasing the number of wild sequences by a factor of 0.5
(adding 135 wild sequences) resulting in a final "titration" of 1:3 domestic to wild
sequences (n=1080). We applied a two-state asymmetric CTMC discrete trait diffusion
model where sequences were labeled as domestic or wild. All priors and model
parameters selected are the same as those described in the empirical tree set
description above. To study the impact of the inclusion of turkeys in the transmission
between domestic and wild populations we annotated all unannotated sequences
collected from turkeys as domestic. We then created three datasets starting with 525

domestic and 525 wild bird sequences, adding 263 sequences to successive titrations
resulting in 1:1,1:1.5, and 1:2 (domestic:wild) sequence datasets with a final titration
size of 1,575 sequences. We again applied a two-state asymmetric CTMC discrete trait
diffusion model where sequences were labeled as domestic or wild with all priors and
model parameters selected are the same as those described in the empirical tree set
description above. To determine whether the proportion of turkeys to other domestic
birds would impact the results of the previously described titration analysis we built a
dataset with where the domestic bird group had equal numbers of turkey and domestic
(non-turkey) sequences. This dataset included 173 turkey, 173 domestic bird, and 692
wild bird sequences totaling 1038 sequences. We applied an asymmetric CTMC
discrete trait diffusion model using a BSSVS for a three-trait model with the following
states: wild birds, domestic birds (not turkey), and turkey. We performed three
independent runs of this analysis using the models and parameters described in the
empirical tree analysis section above. All titration replicates were performed using an
MCMC chain length of 100 million states sampling every 10,000 states.
1128
1129 *Commercial, backyard, wild bird titration analysis*
1130
Metadata and annotated sequences were made available describing sequences as
being from backyard birds for sequences collected in early 2022 which distinguished
them from commercial poultry (previously all sequences being determined domestic)
(10). We used this metadata to create a dataset with equally sampled backyard birds
and commercial birds (n= 85 for each bird type) and then added all available wild birds
(n=722) in 25% increments creating four separate datasets for sequences collected
between Jan 2022 and June 2022. This resulted in a final dataset of n= 942 sequences.
We performed discrete trait diffusion modeling using an asymmetric CTMC diffusion
model for sequences labeled as backyard bird, commercial bird, and wild bird. We
employed the Markov Jump analysis to observe the number of jumps between discrete
states across the posterior set of trees and estimated the Markov Rewards to determine
the waiting time for a given discrete trait state in the phylogeny (58,59).
1143
1144 *Extraction of phylogenetic metrics*
1145
1146 We calculated the transitions between states across branches of phylogenies estimated
1147 from ancestral state reconstructions using the Baltic python package (22). To calculate
1148 the persistence of a given discrete trait we used the program PACT v0.9.5. which
1149 calculates the persistence of a trait by traversing the phylogenetic tree backwards and
1150 measuring the amount of time a tip takes to leave its sampled state (98).
1151
1152 **Data and code availability:**
1153 All analytical scripts, metadata annotations, and BEAST XMLs used in this analysis can
1154 be found at the following GitHub repository: https://github.com/moncla-lab/North-
1155 American-HPAI
1156

All data that was used in this analysis were sourced from public databases. Acknowledgement table for GISAID isolates used in this analysis can be found in Table S13.

Several of the analyses presented have also been publicly made available using a maximum likelihood framework through the Nextstrain pipeline and a narrative of this work can be found in the following link: https://nextstrain.org/community/narratives/moncla-lab/nextstrain-narrative-hpai-north-america@main/HPAI-in-North-America

Acknowledgements

We would like to thank Mia Kim Torchetti for her helpful feedback and discussion about our results. This work was supported by NIH R00-AI147029-05 and by funding from the Centers of Excellence for Influenza Research and Response (CEIRR), funded by NIH 75N93021C00015. LHM is a Pew Biomedical Scholar and is supported by NIH R00-AI147029-05. AJ is supported by NIH R00-AI147029-05, and LD is supported by NIH 75N93021C00015.

Citations:

1. Puryear WB, Runstadler JA. High-pathogenicity avian influenza in wildlife: a changing disease dynamic that is expanding in wild birds and having an increasing impact on a growing number of mammals. J Am Vet Med Assoc. 2024 Apr 10;1–9.

2. Lycett SJ, Pohlmann A, Staubach C, Caliendo V, Woolhouse M, Beer M, et al. Genesis and spread of multiple reassortants during the 2016/2017 H5 avian influenza epidemic in Eurasia. Proc Natl Acad Sci. 2020 Aug 25;117(34):20814–25.

3. Smith GJD, Vijaykrishna D, Ellis TM, Dyrting KC, Leung YHC, Bahl J, et al. Characterization of Avian Influenza Viruses A (H5N1) from Wild Birds, Hong Kong, 2004–2008. Emerg Infect Dis. 2009 Mar;15(3):402–7.

4. Alexander DJ. Summary of Avian Influenza Activity in Europe, Asia, Africa, and Australasia, 2002–2006. Avian Dis. 2007 Mar;51(s1):161–6.

5. Final Report for the 2014–2015 Outbreak of Highly Pathogenic Avian Influenza (HPAI) in the United States [Internet]. Veterinary Services Surveillance, Preparedness, and Response Services Animal and Plant Health Inspection Service: United States Department of Agriculture; 2016 Aug [cited 2024 Apr 1]. Available from: https://www.aphis.usda.gov/media/document/2086/file

6. Caliendo V, Lewis NS, Pohlmann A, Baillie SR, Banyard AC, Beer M, et al. Transatlantic spread of highly pathogenic avian influenza H5N1 by wild birds from Europe to North America in 2021. Sci Rep. 2022 Jul 11;12(1):11729.

7. Gass JD, Hill NJ, Damodaran L, Naumova EN, Nutter FB, Runstadler JA. Ecogeographic
Drivers of the Spatial Spread of Highly Pathogenic Avian Influenza Outbreaks in Europe and
the United States, 2016–Early 2022. Int J Environ Res Public Health. 2023 Jun 1;20(11):6030.

8. Erdelyan CNG, Kandeil A, Signore AV, Jones MEB, Vogel P, Andreev K, et al. Multiple
transatlantic incursions of highly pathogenic avian influenza clade 2.3.4.4b A(H5N5) virus
into North America and spillover to mammals. Cell Rep. 2024 Jul;43(7):114479.

9. Klaassen M, Wille M. The plight and role of wild birds in the current bird flu panzootic. Nat
Ecol Evol. 2023 Aug 16;7(10):1541–2.

10. Youk S, Torchetti MK, Lantz K, Lenoch JB, Killian ML, Leyson C, et al. H5N1 highly pathogenic
avian influenza clade 2.3.4.4b in wild and domestic birds: Introductions into the United
States and reassortments, December 2021–April 2022. Virology. 2023 Oct;587:109860.

11. Spackman E, Pantin-Jackwood MJ, Lee SA, Prosser D. The pathogenesis of a 2022 North
American highly pathogenic clade 2.3.4.4b H5N1 avian influenza virus in mallards (*Anas*
platyrhynchos). Avian Pathol. 2023 May 4;52(3):219–28.

12. Knief U, Bregnballe T, Alfarwi I, Ballmann MZ, Brenninkmeijer A, Bzoma S, et al. Highly
pathogenic avian influenza causes mass mortality in Sandwich Tern *Thalasseus sandvicensis*
breeding colonies across north-western Europe. Bird Conserv Int. 2024;34:e6.

13. European Food Safety Authority, European Centre for Disease Prevention and Control,
European Union Reference Laboratory for Avian Influenza, Adlhoch C, Fusaro A, Gonzales
JL, et al. Avian influenza overview December 2022 – March 2023. EFSA J [Internet]. 2023
Mar [cited 2024 Apr 25];21(3). Available from:
https://data.europa.eu/doi/10.2903/j.efsa.2023.7917

14. Elsmo E, Wünschmann A, Beckmen K, Broughton-Neiswanger L, Buckles E, Ellis J, et al.
Pathology of natural infection with highly pathogenic avian influenza virus (H5N1) clade
2.3.4.4b in wild terrestrial mammals in the United States in 2022 [Internet]. 2023 [cited
2024 Apr 26]. Available from: http://biorxiv.org/lookup/doi/10.1101/2023.03.10.532068

15. Nguyen TQ, Hutter C, Markin A, Thomas M, Lantz K, Killian ML, et al. Emergence and
interstate spread of highly pathogenic avian influenza A(H5N1) in dairy cattle [Internet].
2024 [cited 2024 May 26]. Available from:
http://biorxiv.org/lookup/doi/10.1101/2024.05.01.591751

16. Puryear W, Sawatzki K, Hill N, Foss A, Stone JJ, Doughty L, et al. Outbreak of Highly
Pathogenic Avian Influenza H5N1 in New England Seals [Internet]. 2022 [cited 2024 Jun 5].
Available from: http://biorxiv.org/lookup/doi/10.1101/2022.07.29.501155

17. Kaplan BS, Webby RJ. The avian and mammalian host range of highly pathogenic avian
H5N1 influenza. Virus Res. 2013 Dec;178(1):3–11.

18. Hill NJ, Bishop MA, Trovão NS, Ineson KM, Schaefer AL, Puryear WB, et al. Ecological
divergence of wild birds drives avian influenza spillover and global spread. Murcia PR,
editor. PLOS Pathog. 2022 May 19;18(5):e1010062.

19. Sonnberg S, Webby RJ, Webster RG. Natural history of highly pathogenic avian influenza
H5N1. Virus Res. 2013 Dec;178(1):63–77.

20. Volz EM, Koelle K, Bedford T. Viral Phylodynamics. Wodak S, editor. PLoS Comput Biol. 2013
Mar;9(3):e1002947.

21. Frost SDW, Volz EM. Viral phylodynamics and the search for an 'effective number of
infections.' Philos Trans R Soc B Biol Sci. 2010 Jun 27;365(1548):1879–90.

22. Dudas G, Carvalho LM, Rambaut A, Bedford T. MERS-CoV spillover at the camel-human
interface. eLife. 2018 Jan 16;7:e31257.

23. De Maio N, Wu CH, O'Reilly KM, Wilson D. New Routes to Phylogeography: A Bayesian
Structured Coalescent Approximation. Pritchard JK, editor. PLOS Genet. 2015 Aug
12;11(8):e1005421.

24. Bevins SN, Pedersen K, Lutman MW, Baroch JA, Schmit BS, Kohler D, et al. Large-Scale Avian
Influenza Surveillance in Wild Birds throughout the United States. Yoon KJ, editor. PLoS
ONE. 2014 Aug 12;9(8):e104360.

25. Shriner SA, Root JJ, Lutman MW, Kloft JM, VanDalen KK, Sullivan HJ, et al. Surveillance for
highly pathogenic H5 avian influenza virus in synanthropic wildlife associated with poultry
farms during an acute outbreak. Sci Rep. 2016 Nov 4;6(1):36237.

26. Bokma BH, Hall C, Siegfried LM, Todd Weaver J. Surveillance for Avian Influenza in the
United States. Ann N Y Acad Sci. 2006 Oct;1081(1):163–8.

27. Poultry 2004 Part I: Reference of Health and Management of Backyard / Small Production
Flocks in the United States, 2004 [Internet]. U.S. Department of Agriculture. Animal and
Plant Health Inspection Service. Veterinary Services. National Animal Health Monitoring
System (NAHMS); Available from:
https://www.aphis.usda.gov/sites/default/files/poultry04_dr_parti.pdf

28. Garber L, Hill G, Rodriguez J, Gregory G, Voelker L. Non-commercial poultry industries:
Surveys of backyard and gamefowl breeder flocks in the United States. Prev Vet Med. 2007
Jul 16;80(2):120–8.

29. 2022 OIE - Terrestrial Animal Health Code [Internet]. WOAH; 2022 Mar. Available from:
https://www.woah.org/fileadmin/Home/eng/Health_standards/tahc/current/glossaire.pdf

30. McKellar AE. Phenological Synchrony and Bird Migration: Changing Climate and Seasonal
Resources in North America **Phenological Synchrony and Bird Migration: Changing Climate**

and Seasonal Resources in North America edited by Eric M. Wood and Jherime L.
Kellermann. 2015. CRC Press, Boca Raton, Florida, USA. xiv + 228 pages, 8 color and 53
black-and-white illustrations. $116.96 (hardcover). ISBN 978-1-4822-4030-6. The Auk. 2016
Jan;133(1):113–4.

31. Marra PP, Francis CM, Mulvihill RS, Moore FR. The influence of climate on the timing and
rate of spring bird migration. Oecologia. 2005 Jan;142(2):307–15.

32. Kent CM, Ramey AM, Ackerman JT, Bahl J, Bevins SN, Bowman AS, et al. Spatiotemporal
changes in influenza A virus prevalence among wild waterfowl inhabiting the continental
United States throughout the annual cycle. Sci Rep. 2022 Jul 29;12(1):13083.

33. Lemey P, Rambaut A, Drummond AJ, Suchard MA. Bayesian Phylogeography Finds Its Roots.
Fraser C, editor. PLoS Comput Biol. 2009 Sep;5(9):e1000520.

34. Ramey AM, Scott LC, Ahlstrom CA, Buck EJ, Williams AR, Kim Torchetti M, et al. Molecular
detection and characterization of highly pathogenic H5N1 clade 2.3.4.4b avian influenza
viruses among hunter-harvested wild birds provides evidence for three independent
introductions into Alaska. Virology. 2024 Jan;589:109938.

35. Xie R, Edwards KM, Wille M, Wei X, Wong SS, Zanin M, et al. The episodic resurgence of
highly pathogenic avian influenza H5 virus. Nature. 2023 Oct 26;622(7984):810–7.

36. Migratory Bird Program Administrative Flyways | U.S. Fish & Wildlife Service [Internet].
2023 [cited 2024 Apr 28]. Available from: https://www.fws.gov/partner/migratory-bird-
program-administrative-flyways

37. Gergely, K.J., Boykin, K.G., McKerrow, A.J., Rubino, M.J., Tarr, N.M., and Williams, S.G. Gap
Analysis Project (GAP) Terrestrial Vertebrate Species Richness Maps for the Conterminous
U.S. [Internet]. 2019. (U.S. Geological Survey Scientific Investigations Report 2019–5034).
Available from: https://doi.org/10.3133/sir20195034.

38. Bahl J, Krauss S, Kühnert D, Fourment M, Raven G, Pryor SP, et al. Influenza A Virus
Migration and Persistence in North American Wild Birds. Andino R, editor. PLoS Pathog.
2013 Aug 29;9(8):e1003570.

39. Fourment M, Darling AE, Holmes EC. The impact of migratory flyways on the spread of avian
influenza virus in North America. BMC Evol Biol. 2017 Dec;17(1):118.

40. Prosser DJ, Chen J, Ahlstrom CA, Reeves AB, Poulson RL, Sullivan JD, et al. Maintenance and
dissemination of avian-origin influenza A virus within the northern Atlantic Flyway of North
America. Fouchier RAM, editor. PLOS Pathog. 2022 Jun 6;18(6):e1010605.

41. Tobias JA, Sheard C, Pigot AL, Devenish AJM, Yang J, Sayol F, et al. AVONET: morphological,
ecological and geographical data for all birds. Coulson T, editor. Ecol Lett. 2022
Mar;25(3):581–97.

42. Arnal A, Vittecoq M, Pearce-Duvet J, Gauthier-Clerc M, Boulinier T, Jourdain E. Laridae: A
neglected reservoir that could play a major role in avian influenza virus epidemiological
dynamics. Crit Rev Microbiol. 2015 Oct 2;41(4):508–19.

43. Lee DH, Torchetti MK, Hicks J, Killian ML, Bahl J, Pantin-Jackwood M, et al. Transmission
Dynamics of Highly Pathogenic Avian Influenza Virus A(H5Nx) Clade 2.3.4.4, North America,
2014–2015. Emerg Infect Dis. 2018 Oct;24(10):1840–8.

44. Wille M. Ecology and Evolution of Avian Influenza A Viruses in Wild Birds. In: Genetics and
Evolution of Infectious Diseases [Internet]. Elsevier; 2024 [cited 2024 Jul 29]. p. 863–98.
Available from: https://linkinghub.elsevier.com/retrieve/pii/B9780443288180000057

45. Munster VJ, Baas C, Lexmond P, Waldenström J, Wallensten A, Fransson T, et al. Spatial,
Temporal, and Species Variation in Prevalence of Influenza A Viruses in Wild Migratory
Birds. Kawaoka Y, editor. PLoS Pathog. 2007 May 11;3(5):e61.

46. Hong SL, Lemey P, Suchard MA, Baele G. Bayesian Phylogeographic Analysis Incorporating
Predictors and Individual Travel Histories in BEAST. Curr Protoc. 2021 Apr;1(4):e98.

47. Nemeth NM, Ruder MG, Poulson RL, Sargent R, Breeding S, Evans MN, et al. Bald eagle
mortality and nest failure due to clade 2.3.4.4 highly pathogenic H5N1 influenza a virus. Sci
Rep. 2023 Jan 5;13(1):191.

48. Shearn-Bochsler VI, Knowles S, Ip H. Lethal Infection of Wild Raptors with Highly Pathogenic
Avian Influenza H5N8 and H5N2 Viruses in the USA, 2014–15. J Wildl Dis. 2019 Jan
1;55(1):164.

49. Redig PT, Goyal SM. Serologic Evidence of Exposure of Raptors to Influenza A Virus. Avian
Dis. 2012 Jun;56(2):411–3.

50. 2022–2024 Detections of Highly Pathogenic Avian Influenza [Internet]. [cited 2024 Apr 28].
Available from: https://www.aphis.usda.gov/livestock-poultry-disease/avian/avian-
influenza/hpai-detections

51. Kim HK, Kim HJ, Noh JY, Van Phan L, Kim JH, Song D, et al. Serological evidence of H5-
subtype influenza A virus infection in indigenous avian and mammalian species in Korea.
Arch Virol. 2018 Mar;163(3):649–57.

52. Komar N, Olsen B. Avian Influenza Virus (H5N1) Mortality Surveillance. Emerg Infect Dis.
2008 Jul;14(7):1176–8.

53. Jennelle CS, Carstensen M, Hildebrand EC, Cornicelli L, Wolf P, Grear DA, et al. Surveillance
for Highly Pathogenic Avian Influenza Virus in Wild Birds during Outbreaks in Domestic
Poultry, Minnesota, 2015. Emerg Infect Dis. 2016 Jul;22(7):1278–82.

54. Farahat RA, Khan SH, Rabaan AA, Al-Tawfiq JA. The resurgence of Avian influenza and
human infection: A brief outlook. New Microbes New Infect. 2023 Jun;53:101122.

55. Patyk KA, Fields VL, Beam AL, Branan MA, McGuigan RE, Green A, et al. Investigation of risk
factors for introduction of highly pathogenic avian influenza H5N1 infection among
commercial turkey operations in the United States, 2022: a case-control study. Front Vet
Sci. 2023 Aug 30;10:1229071.

56. APHIS FOREIGN ANIMAL DISEASE FRAMEWORK RESPONSE STRATEGIES [Internet]. United
States Department of Agriculture; Available from:
https://www.aphis.usda.gov/sites/default/files/fadprep_manual_2.pdf

57. APPA NATIONAL PET OWNERS SURVEY 2021-2022. American Pet Products Association;

58. Minin VN, Suchard MA. Counting labeled transitions in continuous-time Markov models of
evolution. J Math Biol. 2007 Nov 30;56(3):391–412.

59. Minin VN, Suchard MA. Fast, accurate and simulation-free stochastic mapping. Philos Trans
R Soc B Biol Sci. 2008 Dec 27;363(1512):3985–95.

60. Alkie TN, Lopes S, Hisanaga T, Xu W, Suderman M, Koziuk J, et al. A threat from both sides:
Multiple introductions of genetically distinct H5 HPAI viruses into Canada via both East Asia-
Australasia/Pacific and Atlantic flyways. Virus Evol. 2022 Sep 10;8(2):veac077.

61. Graziosi G, Lupini C, Catelli E, Carnaccini S. Highly Pathogenic Avian Influenza (HPAI) H5
Clade 2.3.4.4b Virus Infection in Birds and Mammals. Animals. 2024 May 2;14(9):1372.

62. Tarasiuk K, Kycko A, Knitter M, Świętoń E, Wyrostek K, Domańska-Blicharz K, et al.
Pathogenicity of highly pathogenic avian influenza H5N8 subtype for herring gulls (Larus
argentatus): impact of homo- and heterosubtypic immunity on the outcome of infection.
Vet Res. 2022 Dec 14;53(1):108.

63. Kramer LD, Ciota AT, Kilpatrick AM. Introduction, Spread, and Establishment of West Nile
Virus in the Americas. Reisen W, editor. J Med Entomol. 2019 Oct 28;56(6):1448–55.

64. Stallknecht DE, Carter DL, Sullivan-Brügger L, Link P, Ferraro E, McCarty C, et al. Highly
Pathogenic H5N1 Influenza A Virus (IAV) in Blue-Winged Teal in the Mississippi Flyway Is
Following the Historic Seasonal Pattern of Low-Pathogenicity IAV in Ducks. Pathogens. 2024
Nov 19;13(11):1017.

65. Ahlstrom CA, Torchetti MK, Lenoch J, Beckmen K, Boldenow M, Buck EJ, et al. Genomic
characterization of highly pathogenic H5 avian influenza viruses from Alaska during 2022
provides evidence for genotype-specific trends of spatiotemporal and interspecies
dissemination. Emerg Microbes Infect. 2024 Dec 31;13(1):2406291.

66. Kandeil A, Patton C, Jones JC, Jeevan T, Harrington WN, Trifkovic S, et al. Rapid evolution of
A(H5N1) influenza viruses after intercontinental spread to North America. Nat Commun.
2023 May 29;14(1):3082.

67. Lam TT, Ip HS, Ghedin E, Wentworth DE, Halpin RA, Stockwell TB, et al. Migratory flyway
and geographical distance are barriers to the gene flow of influenza virus among North
American birds. Ecol Lett. 2012 Jan;15(1):24–33.

68. Krauss S, Obert CA, Franks J, Walker D, Jones K, Seiler P, et al. Influenza in Migratory Birds
and Evidence of Limited Intercontinental Virus Exchange. Kawaoka Y, editor. PLoS Pathog.
2007 Nov 9;3(11):e167.

69. Girard YA, Runstadler JA, Aldehoff F, Boyce W. Genetic structure of Pacific Flyway avian
influenza viruses is shaped by geographic location, host species, and sampling period. Virus
Genes. 2012 Jun;44(3):415–28.

70. Buhnerkempe MG, Webb CT, Merton AA, Buhnerkempe JE, Givens GH, Miller RS, et al.
Identification of migratory bird flyways in North America using community detection on
biological networks. Ecol Appl. 2016 Apr;26(3):740–51.

71. National Agricultural Statistics Service - [Internet]. USDA; Available from:
https://www.nass.usda.gov/Charts_and_Maps/Poultry/

72. Pantin-Jackwood MJ, Costa-Hurtado M, Shepherd E, DeJesus E, Smith D, Spackman E, et al.
Pathogenicity and Transmission of H5 and H7 Highly Pathogenic Avian Influenza Viruses in
Mallards. Schultz-Cherry S, editor. J Virol. 2016 Nov;90(21):9967–82.

73. Trovão NS, Suchard MA, Baele G, Gilbert M, Lemey P. Bayesian Inference Reveals Host-
Specific Contributions to the Epidemic Expansion of Influenza A H5N1. Mol Biol Evol. 2015
Sep 3;msv185.

74. Hayes S, Hilton J, Mould-Quevedo J, Donnelly C, Baylis M, Brierley L. Ecology and
environment predict spatially stratified risk of highly pathogenic avian influenza in wild
birds across Europe [Internet]. 2024 [cited 2024 Jul 30]. Available from:
http://biorxiv.org/lookup/doi/10.1101/2024.07.17.603912

75. Sims LD, Weaver J, Swayne DE. Epidemiology of avian influenza in agricultural and other
man-made systems. In: Swayne DE, editor. Animal Influenza [Internet]. 1st ed. Wiley; 2016
[cited 2024 Apr 28]. p. 302–36. Available from:
https://onlinelibrary.wiley.com/doi/10.1002/9781118924341.ch12

76. Rasmussen EA, Czaja A, Cuthbert FJ, Tan GS, Lemey P, Nelson MI, et al. Influenza A viruses
in gulls in landfills and freshwater habitats in Minnesota, United States. Front Genet. 2023
May 9;14:1172048.

77. Fouchier R, Munster V. Epidemiology of low pathogenic avian influenza viruses in wild birds:
-EN- Epidemiology of low pathogenic avian influenza viruses in wild birds -FR- Epidemiologie
des virus de l'influenza aviaire faiblement pathogène dans l'avifaune -ES- Epidemiología de
la influenza aviar de baja patogenicidad en aves silvestres. Rev Sci Tech OIE. 2009 Apr
1;28(1):49–58.

78. Hicks JT, Edwards K, Qiu X, Kim DK, Hixson JE, Krauss S, et al. Host diversity and behavior
determine patterns of interspecies transmission and geographic diffusion of avian influenza
A subtypes among North American wild reservoir species. Stern A, editor. PLOS Pathog.
2022 Apr 13;18(4):e1009973.

79. Yang Q, Wang B, Lemey P, Dong L, Mu T, Wiebe RA, et al. Synchrony of Bird Migration with
Global Dispersal of Avian Influenza Reveals Exposed Bird Orders. Nat Commun. 2024 Feb
6;15(1):1126.

80. Hicks JT, Lee DH, Duvvuri VR, Kim Torchetti M, Swayne DE, Bahl J. Agricultural and
geographic factors shaped the North American 2015 highly pathogenic avian influenza
H5N2 outbreak. Nelson MI, editor. PLOS Pathog. 2020 Jan 21;16(1):e1007857.

81. Verhagen JH, Fouchier RAM, Lewis N. Highly Pathogenic Avian Influenza Viruses at the
Wild–Domestic Bird Interface in Europe: Future Directions for Research and Surveillance.
Viruses. 2021 Jan 30;13(2):212.

82. Sullivan JD, McDonough AM, Lescure LM, Prosser DJ. Identifying an Understudied Interface:
Preliminary Evaluation of the Use of Retention Ponds on Commercial Poultry Farms by Wild
Waterfowl. Korennoy F, editor. Transbound Emerg Dis. 2024 Apr 3;2024:1–9.

83. Madsen JM, Zimmermann NG, Timmons J, Tablante NL. Avian Influenza Seroprevalence and
Biosecurity Risk Factors in Maryland Backyard Poultry: A Cross-Sectional Study. Leung FCC,
editor. PLoS ONE. 2013 Feb 20;8(2):e56851.

84. Bertran K, Lee DH, Pantin-Jackwood MJ, Spackman E, Balzli C, Suarez DL, et al. Pathobiology
of Clade 2.3.4.4 H5Nx High-Pathogenicity Avian Influenza Virus Infections in Minor
Gallinaceous Poultry Supports Early Backyard Flock Introductions in the Western United
States in 2014-2015. Schultz-Cherry S, editor. J Virol. 2017 Nov;91(21):e00960-17.

85. Molia S, Traoré I, Kamissoko B, Diakité A, Sidibé MS, Sissoko KD, et al. Characteristics of
commercial and traditional village poultry farming in Mali with a focus on practices
influencing the risk of transmission of avian influenza and Newcastle disease. Acta Trop.
2015 Oct;150:14–22.

86. Sheta BM, Fuller TL, Larison B, Njabo KY, Ahmed AS, Harrigan R, et al. Putative human and
avian risk factors for avian influenza virus infections in backyard poultry in Egypt. Vet
Microbiol. 2014 Jan;168(1):208–13.

87. Shu Y, McCauley J. GISAID: Global initiative on sharing all influenza data – from vision to
reality. Eurosurveillance [Internet]. 2017 Mar 30;22(13). Available from:
https://www.eurosurveillance.org/content/10.2807/1560-7917.ES.2017.22.13.30494

88. Katoh K, Standley DM. MAFFT Multiple Sequence Alignment Software Version 7:
Improvements in Performance and Usability. Mol Biol Evol. 2013 Apr;30(4):772–80.

89. Kearse M, Moir R, Wilson A, Stones-Havas S, Cheung M, Sturrock S, et al. Geneious Basic: An
integrated and extendable desktop software platform for the organization and analysis of
sequence data. Bioinformatics. 2012 Jun;28(12):1647–9.

90. Minh BQ, Schmidt HA, Chernomor O, Schrempf D, Woodhams MD, von Haeseler A, et al. IQ-
TREE 2: New Models and Efficient Methods for Phylogenetic Inference in the Genomic Era.
Teeling E, editor. Mol Biol Evol. 2020 May;37(5):1530–4.

91. Sagulenko P, Puller V, Neher RA. TreeTime: Maximum-likelihood phylodynamic analysis.
Virus Evol [Internet]. 2018 Jan [cited 2021 Jul 25];4(1). Available from:
http://academic.oup.com/ve/article/doi/10.1093/vex042/4794731

92. Suchard MA, Lemey P, Baele G, Ayres DL, Drummond AJ, Rambaut A. Bayesian phylogenetic
and phylodynamic data integration using BEAST 1.10. Virus Evol [Internet]. 2018 Jan [cited
2021 Sep 12];4(1). Available from:
https://academic.oup.com/ve/article/doi/10.1093/ve/vey016/5035211

93. Shapiro B, Rambaut A, Drummond AJ. Choosing Appropriate Substitution Models for the
Phylogenetic Analysis of Protein-Coding Sequences. Mol Biol Evol. 2006 Jan 1;23(1):7–9.

94. Drummond AJ, Ho SYW, Phillips MJ, Rambaut A. Relaxed Phylogenetics and Dating with
Confidence. Penny D, editor. PLoS Biol. 2006 Mar;4(5):e88.

95. Gill MS, Lemey P, Faria NR, Rambaut A, Shapiro B, Suchard MA. Improving Bayesian
Population Dynamics Inference: A Coalescent-Based Model for Multiple Loci. Mol Biol Evol.
2013 Mar 1;30(3):713–24.

96. Kass RE, Raftery AE. Bayes Factors. J Am Stat Assoc. 1995 Jun;90(430):773–95.

97. Bielejec F, Baele G, Vrancken B, Suchard MA, Rambaut A, Lemey P. SpreaD3: Interactive
Visualization of Spatiotemporal History and Trait Evolutionary Processes. Mol Biol Evol.
2016 Aug;33(8):2167–9.

98. Bedford T, Riley S, Barr IG, Broor S, Chadha M, Cox NJ, et al. Global circulation patterns of
seasonal influenza viruses vary with antigenic drift. Nature. 2015 Jul;523(7559):217–20.

1468