1 **Intensive transmission in wild, migratory birds drove rapid geographic**
2 **dissemination and repeated spillovers of H5N1 into agriculture in North America**
3

4 Lambodhar Damodaran[1], Anna Jaeger[1], Louise H. Moncla[1]
5
6 [1]Department of Pathobiology, School of Veterinary Medicine, University of Pennsylvania
7
8 Corresponding Author: Louise H. Moncla
9 Email: lhmoncla@upenn.edu

10
11 **Abstract**
12

Since late 2021, a panzootic of highly pathogenic H5N1 has devastated wild birds,
agriculture, and mammals. Analysis of 1,818 Hemagglutinin sequences from wild birds,
domestic birds and mammals revealed that the North American panzootic was driven by
~9 introductions into the Atlantic and Pacific Flyways, followed by rapid dissemination
via wild, migratory birds. Transmission was primarily driven by Anseriformes, while non-
canonical species acted as dead-end hosts. Unlike the epizootic of 2015, outbreaks in
domestic birds were driven by ~46-113 independent introductions from wild birds that
persisted for up to 6 months. Backyard birds were infected ~9 days earlier on average
than commercial poultry, suggesting potential as "early warning signals" for
transmission upticks. We find that wild birds are an emerging source of H5N1 in North
America, necessitating enhanced, continuous surveillance. Prevention of agricultural
outbreaks may now require novel strategies that reduce transmission at the wild
bird/agriculture interface.

26
27
28
29 **Introduction**
30
Highly pathogenic avian influenza (HPAI) viruses pose persistent challenges for human
and animal health. Since emerging in 1996, highly pathogenic H5N1 viruses of the
A/goose/Guangdong lineage have spread globally via endemic transmission among
domestic birds in Asia and Africa coupled with long-distance dispersal by wild migrating
birds[1,2]. In 2005, introduction of poultry-derived H5N1 viruses into wild birds in China led
to viral dispersal across Northern Africa and Asia, establishing new lineages of endemic
circulation in poultry[3,4]. In 2014, wild, migratory birds carried highly pathogenic H5N8
viruses from Europe to North America, sparking an outbreak that resulted in the culling
of over 50.5 million commercial birds in the United States [5]. While this outbreak
substantially impacted the agriculture industry, these viruses did not establish persistent
transmission in wild birds. Following aggressive culling of domestic birds, the epizootic
was extinguished and North America remained free of HPAI for years.
43
In December 2021, clade 2.3.4.4b HPAI H5N1 viruses were introduced and spread
across the Americas, causing a panzootic of significant morbidity and mortality in wild
and domestic animals. These viruses were likely first introduced into North America in
late 2021 by migratory birds flying across the Arctic Circle from Europe[6,7], after which

reassortment with low-pathogenicity avian influenza (LPAI) viruses endemic to North
America produced a reassortant with altered neurotropism in experimentally infected
mammals[8]. In contrast to past North American epizootics, morbidity and mortality has
been widespread across a broad range of wild avian species not usually impacted by
HPAI[9] such as raptors, owls, passerines [1,10], and Sandwich Terns[11–13]. Infections have
also occurred in mammal species not typically associated with HPAI, such as foxes,
skunks, raccoons, harbor seals, dolphins, bears[1,14], domestic goats, and dairy cattle[15].
The broad range of affected wild avian species raises the possibility that new reservoir
hosts (hosts that would sustain continuous transmission chains) could be established,
necessitating an evaluation of which species should be actively surveilled. Putative
transmission among marine mammals and domestic dairy cattle pose new challenges
for animal health and biosecurity, and highlight the need to understand the ecological
factors that lead to spillover in mammals and agriculture[15,16].
61
Historically, H5N1 transmission has been linked to enzootic transmission in domestic
poultry[17,18], paired with occasional cross-continental movement by wild birds of the
Anseriformes (waterfowl such as ducks and geese) and Charadriiformes (Shorebirds)
orders[19–21]. Unlike the North American epizootic in 2014-2015, widespread culling of
domestic birds has not halted detections in North America, suggesting that patterns of
transmission since 2022 may be distinct from past North American epizootics. Prior
work has posited that clade 2.3.4.4b viruses may be better able to infect and transmit
among wild bird species, leading to persistent, seasonal circulation in European wild
birds and widespread reassortment[2,11,13]. In Europe, clade 2.3.4.4b viruses caused
repeated incursions into European wild and domestic birds from 2016-2020, with a
notable shift towards seasonal outbreaks and a broader range in infected wild species
since 2020[22]. Thus, rather than acting as transient hosts that facilitate long-range
movement, wild birds are now thought to play a greater role in maintaining and
disseminating these viruses across Europe and Asia. However, the role of wild vs.
domestic birds in driving transmission in North America has not been robustly or
comprehensively studied, limiting informed enactment of surveillance and outbreak
intervention strategies.
79
Previous genomic analysis of the United States epizootic linked outbreaks in poultry to
wild birds, though the robustness of these results to differences in sampling between
wild and domestic birds was not directly examined[10]. Other studies have reported
surveillance and localized outbreak data to identify new incursions in wild bird
populations[6,7,23], but do not address the role of different species in contributing to cross-
continental epizootic spread, transmission between species, or outbreaks in agriculture.
As of 2025, highly pathogenic avian influenza viruses are classified as foreign animal
diseases by the United States and Canada, with outbreak control plans primarily
focused on biosecurity to reduce spread between farms, and rapidly culling domestic
birds[24,25]. If the epizootic in North America reflects the changing ecology of clade
2.3.4.4b viruses towards wild bird driven transmission observed elsewhere, then
surveillance, policy, and outbreak mitigation strategies may need to be fundamentally

reformulated. Surveillance activities should focus on the host groups most important for
viral dispersal, while control plans should be formulated to prevent incursions and
spread in agriculture. Disentangling which species drove the North American epizootic,
and whether transmission was driven by wild birds vs. domestic poultry, are therefore
key for formulating effective surveillance and intervention strategies, but currently
understudied. Finally, while it is currently thought that cases in mammals likely stem
from infections in wild birds, work to formally link infections across species has been
sparse.
100
Viral phylodynamic approaches are emerging as critical tools for outbreak
reconstruction[26,27]. Viral genomes contain molecular records of transmission histories,
allowing them to be used to trace how outbreaks begin and spread. Here we use
Bayesian phylogeographic approaches paired with rigorous controls for sampling
bias[28,29] to trace how highly pathogenic H5N1 viruses were introduced and
disseminated during the first 18 months of the epizootic across North America. We
capitalize on a dataset of 1,824 Hemagglutinin gene sequences sampled from North
American birds and mammals in 2021-2023, and curate additional metadata on
geography, migratory flyways, domestic/wild status, host taxonomic order, and
migratory behavior to reconstruct transmission between these groups. Using this
dataset of HA sequences, we show that the epizootic in North America was driven by ~9
independent introductions that descend from outbreaks in Europe and Asia, though only
a single introduction spread successfully across the continent. The initial wave of H5N1
transmission spread from East to West by wild, migratory birds between adjacent
migratory flyways. Transmission from non-canonical avian species like songbirds and
owls was limited and resulted in dead-end transmission chains, suggesting that these
species are unlikely to establish as reservoirs. Instead, transmission was primarily
sustained by Anseriformes, suggesting these species as useful targets for surveillance
activities. In contrast to the outbreak in 2014/2015, outbreaks in agriculture were
seeded by ~46-113 independent introductions from wild birds, with some onward
transmission. Backyard birds were infected slightly more frequently, and earlier on
average, than commercial birds, suggesting that increased backyard bird surveillance
could be investigated as early warning signs for increased incidence. Together, these
data pinpoint wild birds as emerging sources for H5N1 virus transmission in North
America, capable of rapid, and long-distance viral dispersal and repeated reseeding of
outbreaks in agriculture. Our results implicate continuous surveillance in wild aquatic
migratory birds (particularly Anseriformes) as critical for contextualizing outbreaks in
mammals and agriculture, and suggest that viral evolution in North America may now be
increasingly governed by wild bird movement, ecology, and reassortment. Investment in
interventions beyond culling that reduce interactions between domestic and wild
animals may now be crucial for limiting future outbreaks in agriculture.
132
133
134 **Results**
135
136 *Viral sequence data capture seasonal variation of HPAI detections*

Figure 1. Detections of HPAI in North America show distinct epidemic waves following
introduction events in late 2021. A) Detections of HPAI in wild birds, domestic birds, and non-
human mammals. B) The Log-scaled Effective population size (N_e) estimates estimated in
BEAST using the Bayesian SkyGrid coalescent for sequences collected between Sep 2021 and
Aug 2023. C) Correlation plot of log(N_e) vs HPAI detections by week, spearman correlation
displayed.

Most sequence data from North America represent detections in the United States and
Canada (United States: 1590, Canada: 224, Central America: 8), with a "detection"
classified as a positive PCR test from a collected sample. In Canada, surveillance in
wild and domestic populations are coordinated by a collaborative effort from the
Canadian Food Inspection Agency, Environment Canada, the Public Health Agency of
Canada, and the Canadian Wildlife Health Centre, which perform year-round
surveillance in live and dead wild birds, and case investigation for suspected poultry
outbreaks[30]. In the United States, the United State Department of Agriculture Animal
and Plant Health Inspection Service (APHIS) manages HPAI surveillance and testing in
wild birds via investigation of reported morbidity and mortality events, hunter-harvested
game birds/waterfowl, sentinel species/live bird collection, and environmental sampling
of water bodies and surfaces[31,32]. USDA APHIS also surveilles domestic birds using

several reporting methods: mandatory testing through the National Poultry Improvement
Plan, coordination with state agricultural agencies, routine testing in high-risk areas, and
backyard flock surveillance[33].
Throughout the epizootic and during the time period analyzed in this study
(November 2021-September 2023), most HPAI detections in the US have been
reported in wild birds sampled via testing of sick and dead (5,611), hunter harvested
(3,340), and live wild birds (1,127) (Figure S1a). Data on domestic bird detections are
reported with information on poultry type (e.g., duck, chicken) and by whether the farm
is classified as a commercial operation or backyard flock. Backyard flocks are
categorized by the USDA as operations with fewer than 1,000 birds[34,35] and by the
World Organization for Animal Health (WOAH) as any birds kept in captivity for reasons
other than for commercial production[36]. Among domestic birds, detections (1,177 total)
came predominantly from commercial chickens (9.3%), commercial turkeys (28.5%),
commercial breeding operations (species unspecified) (15.3%), and birds designated
WOAH Non-Poultry which refers to backyard birds (42.3%) (Figure S1b). Other
domestic bird detections occurred in game bird raising operations (2.5%) and
commercial ducks (2.0%). The North American epizootic has impacted a broad range of
mammalian hosts, with detections (399) reported in red foxes (24.3%), mice (24.1%),
skunks (12.2%), and domestic cats (13.2%). Other mammalian hosts (26.2%) represent
a wide range of species including harbor seals, bobcats, fishers, and bears (Figure
S1c).
The first detection of HPAI H5N1 in North America was reported in migratory
gulls in Newfoundland and Labrador Canada in November 2021, followed by the first
detection in the United States in a wild American wigeon in South Carolina on
December 30th, 2021[6]. From January – May 2022, a wave of 2,510 total detections
were reported across 43 states and 91 different species (Figure 1). Following a lull in
the summer of 2022, cases rose again in August 2022, leading to a larger epizootic
wave that lasted until March 2023, totaling 8,001 detections across all 48 contiguous
U.S. states and Alaska. Case detections peaked in the fall and spring, coinciding
roughly with seasonal migration timing for birds migrating between North and South
America[37,38]. Seasonal case variation could arise due to seasonal bird migration,
fluctuating virus prevalence in wild birds, or fledging times of susceptible chicks[39],
though continued monitoring is necessary to determine whether these patterns persist
in future years.
Sequence data sampled in North America is heavily skewed toward sequences
from the United States (United States: 1590, Canada: 224, Central America: 8), and
from the first 6 months of the outbreak, with 74% of all available sequences sampled
from January-July 2022 (Figure S2). To evaluate whether sequence data reflect case
detections, we inferred the viral effective population size (Ne), a measure of viral
genetic diversity shown to be mathematically related to disease prevalence and the
disease transmission rate[26]. We inferred Ne using a nonparametric population model
(Skygrid), which captures relative changes in genetic diversity and the variability of
growth rate in the virus population over time, providing a proxy for epidemic dynamics
as previously described[26,40]. Using 6 datasets of sequences subsampled by host

taxonomic order (see Methods for details), we infer that *Ne* is modestly correlated with
detections (highest Spearman rank correlation: 0.65, p=4.4e-11) (Figure 1c)(Figure S3-
4), and that peaks in *Ne* precede peaks in detections by ~1 week (Figure S5), likely
reflecting the lag between viral transmission and case detection. Thus, despite uneven
sequence acquisition across time, the diversity of sampled sequences roughly reflect
the amplitude of H5N1 cases. We therefore opted to use sequence data for the entire
sampling period for broad inferences on introductions and geographic spread, but
supplement these analyses with controls for sampling differences between groups. For
more intensive reconstructions of transmission patterns between wild birds, commercial
poultry, and backyard birds, we focus on the initial 6-month period with the most
densely sampled data, coupled with experiments to assess the impacts of sampling on
results. Finally, though we retained data from Canada and Central America for all
subsequent analyses, our results are likely most informative about transmission within
the United States due to the heavy skewing of data towards the United States.
216
217 *Highly pathogenic H5N1 was introduced multiple times from Europe and Asia*
218
Though early case reports have described early introductions into the Atlantic and
Pacific, a comprehensive and geographically representative analysis of the number and
timings of introductions into North America across the continent has not been published
[6,8,23]. To reconstruct the number and timings of H5N1 introductions into North America
during the first 18 months of the epizootic, we constructed a dataset that includes HA
sequences from North America from domestic and wild birds (n= 1,327 unique isolates,
identical sequences were removed), along with contextual sequences from other
continents with ongoing outbreaks in 2021 and 2022 (Asia = 294, Europe = 300). We
then inferred the number and timings of introductions into North America using a
discrete trait phylogeographic model (see Methods for details)[41].
229

Figure 2. Introductions from Europe facilitated the majority of ongoing transmission
while Asian introductions did not result in continent-wide onward transmission. A)
Bayesian phylogenetic reconstruction of n=1,927 globally sampled sequences of HPAI clade
2.3.4.4b colored by continent of isolation. Opacity of branches corresponds to posterior support
for the discrete trait inferred for a given branch, the thickness corresponds to the number of
descendent tips the given branch produces. B) A close-up view of the starred section of the tree
in A, focusing on introductions from Asia. C) We inferred the number of transitions from Asia to
North America across the posterior set of 9,000 trees. The x-axis represents the number of
introductions, and the y-axis represents the proportion of trees across the posterior set with that
number of inferred transitions.

Most sequenced infections in North America (98.5% of tips) descend from a single
introduction from Europe in late 2021 (95% Highest Posterior Density (HPD),
September 9th – October 7th 2021) (Figure 2a), consistent with reports of infected
migratory gulls in Newfoundland and Labrador Canada in November 2021, and
subsequent mortality in farmed birds in December of 2021[6,7,10]. We also recapitulate a
second, short-lived introduction from Europe in 2022 as previously described by
Erdelyan et al[8]. However, we also infer 7 (median = 7, 95% HPD: (6,8)) additional
introductions between February and September 2022 that nest within the diversity of
viruses circulating in Asia (Figure 2B-C), 5 of which have not been previously described.
These introductions represent infections sampled in Alaska, Oregon, California,
Wyoming and British Columbia, Canada that showed limited dissemination and persisted
for short periods of time (0.024 – 6.9 months). The western location of these tips
suggest potential introduction via the Pacific flyway, consistent with previous reports
documenting incursions into North America from Japan via Alaska and the upper
Pacific[42] (Figure S6). Though none of these Pacific introductions had sampled

descendants in the time period analyzed (up to September 2023), additional data
available at time of writing indicate that one of these introductions re-emerged in late
2024 as the D1.1 lineage (annotated in Figure 2a)[43]. Though it remains unclear why this
HA lineage was not detected from mid-2023 to 2024, the novel introductions
documented here, and the eventual outgrowth of one of these lineages, highlights the
importance of surveillance in the Pacific region for capturing viral importations. These
data suggest that H5N1 viruses were introduced into North America at least 9 times,
and that viral flow into the Pacific coast may be far more common than previously
documented.
266
267 *Wild, migratory birds drove rapid expansion of H5N1 across the continent*
268
Recent analyses of global H5N1 circulation patterns suggest wild birds as increasingly
important sources for clade 2.3.4.4b virus evolution and transmission in Europe and
Asia[17]. In the Americas, avian migratory routes are classified by the U.S. Fish and
Wildlife Service (USFWS) into 4 major flyways: the Atlantic, Mississippi, Central, and
Pacific[44]. If the epizootic were spread predominantly by wild, migratory birds, we
reasoned that viruses sampled from the same, or neighboring flyways, should cluster
together more closely than viruses sampled from non-adjacent flyways. To test this, we
assigned avian sequences to the migratory flyway matching the US state of sampling,
assembled a dataset of 250 sequences randomly subsampled for each USFWS flyway
(total n=1,000), and implemented a discrete trait diffusion model to estimate transition
rates between flyways as a proxy for viral movement between regions. To determine
whether sequences clustered more strongly by flyway than expected by chance, we
calculated the Association Index (AI), a measure of how strongly a trait is associated
with a phylogenetic tree[45]. Finally, to determine whether movement between flyways
was better supported than movement across other adjacent geographic areas, we
quantified transitions between 4 North American regions stratified by latitude (see
Methods for details).

286
Figure 3. Migratory birds rapidly disseminated H5N1 via migratory flyways. A)
Phylogenetic reconstruction of n=1,000 sequences colored by continental flyway. Inset is the
results of the PACT analysis for persistence in each flyway (how long a tip takes to leave its
sampled location going backwards on the tree), excluding the Pacific clade to show persistence
following Atlantic introduction. B) Mean and 95% HPD for the number of markov jumps between
USFWS flyways where the color of bar on the right of each jump pair corresponds to the source
population and the height of the bar corresponds to the Bayes Factor (BF) support (where white
corresponds to BF < 3 and full color corresponds to BF > 100). C) U.S. Fish and Wildlife Service
waterfowl flyways map, with arrows annotated to represent rates with Bayes factor support of at
least 100. Here the size of the arrow corresponds to the magnitude of the mean transition rate.
D) Posterior distribution of the number of Markov jumps between flyways in the eastward or
westward direction and between adjacent and distant flyways. E) Chord diagram of discrete trait
diffusion based on migratory status going from the source population on the left to the sink
population on the right where the thickness of the chord represents the mean transition rate and
the color represents the Bayes Factor support.
302
Tips that descend from viruses circulating in Asia (those in Figure 2B) cluster together
as a basal clade inferred in the Pacific flyway (orange cluster at top of tree, posterior
probability = 0.98), consistent with introduction into the West Coast. The primary
introduction from Europe occurred via the Atlantic Flyway, and subsequently spread
rapidly across North America (Figure 3A,C). From the inferred time of introduction in the
Atlantic coast (September 9th – October 7th 2021), viruses descending from this
introduction had been sampled in every other flyway within ~4.8 months, indicating
markedly faster spread than observed for other avian-transmitted viruses[46]. Sequences
clustered strongly by flyway (AI =10.563, p=0.00199), grouping most closely with those
sampled within the same or geographically adjacent flyway (Figure 3A, Table S18).

Transitions (inferred as "Markov jumps") between adjacent flyways were ~10 times
more frequent (mean = 239, 95% HPD: 216, 262) than those between distant flyways
(mean = 24, 95% HPD: 12, 33)(Figure 3D, Figure S7), indicating a strong signal of
dissemination via geographic proximity. Transitions between geographic groups
stratified by latitude showed a similar trend towards transitions between adjacent (mean
= 111, 95% HPD: 97, 125) vs distant (mean = 39, 95% HPD: 31, 46) regions, though
the effect was less pronounced (Figure S9, Table S2), suggesting transmission that
proceed most strongly among flyway regions. Transitions were predominantly inferred
from East to West (Figure 3C, D, Table S1), with East to West jumps inferred ~4.4 times
more frequently (mean = 214, 95% HPD: (196, 232)) than West to East jumps (mean =
49, 95% HPD: (38, 57)) (Figure 3D), and 2.3-3.8 times more frequently than jumps
along the North-South axis (Figure S9, Table S2).
325
We next calculated the Bayes Factor (BF) support for each between-flyway transition
rate, and describe those very high statistical support (BF > 100, see Methods for
details). We infer the highest and most well supported rates from the Mississippi to
Central flyway (56.301 markov jumps/year, 95% HPD: 47.85, 64.33), Atlantic to
Mississippi flyway (37.34 markov jumps/year, 95% HPD: 30.84, 43.065), and Central to
Pacific flyway (13.127 markov jumps/year, 95% HPD: 7.975, 18.077)(Figure 3B, Figure
S8, Table S1). Though the Pacific flyway experienced the highest number of
introductions during the time period analyzed, transitions from the Pacific flyway
elsewhere were inferred with low magnitude and weak support, consistent with the
limited transmission we infer from West to East. Indeed, only a single statistically
supported rate was inferred from the Pacific flyway to the adjacent Central flyway (BF =
3, 11.236 markov jumps/year, 95% HPD: 7.975, 13.292). Quantification of the length of
time that each lineage persisted in each flyway showed slightly longer persistence
within the Atlantic and Pacific flyways, though the estimates were variable (Figure 3A).
We speculate that this pattern could reflect higher habitat and species richness within
coastal flyways[47], or that coastal flyways each only border 1 other flyway for viruses to
transition to, potentially resulting in longer within-flyway persistence.
343
The strong clustering by flyways is consistent with long-range transmission by wild,
migratory birds. To measure this, we next classified wild bird sequences by whether
they were sampled from a wild bird considered migratory, partially migratory, or
sedentary using the AVONET database[48]. We then modeled discrete trait diffusion
across a subsample of 1000 sequences with equal proportions of the following 5
categories: wild migratory birds (includes most ducks and geese), wild partially
migratory birds (some ducks, raptors, and vultures), wild sedentary birds (owls crows),
domestic birds, and non-human mammals. Wild migratory and partially migratory birds
are inferred at the root of the tree with probabilities substantially higher than expected
based on their sampling frequencies (Figure S10 and Table S19), indicating a role for
these species in sustained transmission across the epizootic. Transitions from wild
migratory birds were inferred with the highest number and most strongly supported
transition rates (BF > 3000), indicating that migrating wild birds were critical sources for
infections in other species (Figure 3E, Table S3). In contrast, transitions from non-
migratory wild birds were inferred with low magnitudes and weak support (Figure 3E,
Table S3). Taken together, H5N1 viruses were then rapidly disseminated from East to
West following a single introduction on the Atlantic coast, spreading between adjacent

migratory flyways by wild, migrating wild birds. Viral dispersal was strongly associated
with geographic proximity, with the strongest evidence for viral movement that
proceeded westward. These results highlight the capacity of migratory birds to rapidly
disseminate novel viral incursions across continental North America.
365
366   *Epizootic transmission was sustained by canonical host species*
367
Previous outbreaks of highly pathogenic H5N1 viruses have been facilitated by wild
Anseriformes (waterfowl) and Charadriiformes (shorebirds), and domestic species
(Galliformes and Anseriformes)[18,49–52], though the role of these hosts varies across
outbreaks. While domestic ducks have played critical roles in bridging interactions
between wild and domestic populations in past outbreaks in Asia, domestic ducks
account for only 2% of all detections in the US, and the vast majority of cases in the
North American panzootic have been in wild birds and Gallinaceous poultry (turkeys
and chickens), potentially reflecting differences in poultry production between regions[53].
In the current panzootic, die-offs have occurred across a range of wild, non-canonical
host orders, including Accipitriformes (raptors, condors, vultures), Strigiformes (owls),
and Passeriformes (sparrows, crows, robins, etc.)[9], raising the possibility that these new
species could establish as reservoirs that merit surveillance. Pinpointing which species
contributed to epizootic spread is thus critical for designing effective surveillance
strategies. To determine whether particular host groups played outsized roles in driving
transmission in the epizootic, we classified sequences by taxonomic orders that were
most well-sampled and modeled transmission between them using a discrete trait
model. We consolidated sequences of two orders of raptors, Accipitriformes and
Falconiformes, hereafter referred to as "raptors". We also consolidated two orders of
pelagic birds, Charadriiformes and Pelecaniformes, hereafter referred to as
"shorebirds". Following classification, we defined 7 host order groups: Anseriformes,
Shorebirds, Strigiformes, Passeriformes, Raptors, Galliformes, and non-human
mammals.
390
Discrete trait approaches assume that the number of sequences in a dataset are
representative of the underlying distribution of cases in an outbreak, resulting in faulty
inference when this assumption is violated[29,41,54] and bias when groups are unevenly
sampled[28,29]. To account for differential sampling among these host groups, we
therefore considered two, distinct subsampling approaches. The first is a proportional
sampling regime in which sequences are sampled proportional to the detections in each
host group each month. This common sampling regime assumes that case detections in
each group are the closest proxy for the case distribution in the outbreak, and attempts
to align sampling with underlying model assumptions. However, this approach may not
be appropriate if case detection is heavily biased between groups. For HPAI H5N1 in
North America, detections in wild birds are primarily identified when humans report sick
or dead birds to wildlife health authorities or wildlife rescues (Figure S1A), which may
skew detections towards birds with dedicated rescue services or birds that reside in
closer proximity to humans. For example, Anseriformes and raptors comprised 50.2%
and 20.3% of all sequences, respectively, which could arise from high case intensity or
a higher rate of case acquisition. A second, complementary subsampling approach is to
sample sequences equally, meaning that sequences are sampled from each group in
perfectly equal numbers. By forcing the number of sequences from each group to be

equal, the transmission inference must be driven by the underlying sequence diversity
in each group rather than by sampling differences. Given the high variation among
detections within each host group, we opted to pursue both sampling regimes and focus
on results that were concordant in both. We first performed an Association Index test to
confirm that clustering was sufficient for discrete trait inference (Table S18). Next, for
each regime, we performed 3 independent subsamples, where the dataset was
sampled either proportional to cases, or equally (see Methods for details). For the equal
sampling regime, each dataset included 100 randomly sampled sequences per host
group, except for Passeriformes for which only 57 sequences were available. To
account for variation across subsampled datasets, we combined the results for the 3
independent subsamples to summarize statistical support (Figure S12, Table S4-5).
Due to similar tree topologies across replicates, we visualized the phylogeny of the
dataset with the highest posterior support (equal order subsample 1) below and make
the results of all analyses available in supplement (see Figure S11, Table S6-S11).
Finally, to measure the effects of potential sampling bias on the inferred transition rates,
we performed a modified "tip shuffle" analysis. We generated 100 datasets in which the
host tip assignments were randomly shuffled, re-inferred the host group at internal
nodes, and infer a mean root state probability for each host across the 100 shuffled
datasets. We then compared the root state probability in the empirical data to that
inferred in the shuffled data as a measure of the impact of sampling on the results as
previously described (see Methods for details)[55].

**Figure 4. Anseriformes drove outbreak transmission, while new host species
represent dead-end infections.** A) Bayesian phylogenetic reconstruction of n=655
sequences subsampled by host order with equal proportions of each host. The color of
tips and branches represents taxonomic order, and opacity represents the posterior
support for the inferred host group. Thickness of branches correspond to the number of
tips descending from a given branch. B) Mean number of Markov jumps and 95% HPD
from the host group on the left (labelled "From") to the host on the right (labelled "To")
as inferred from the combined results of three equal orders subsamples. The dot
represents the mean number of Markov jumps and the black lines (whiskers) represent
the 95% HPD. The corresponding bar plot shows the bayes factor support for each
jump pair and color of each bar represents the "From" host. The opacity of the bar
represents the Bayes factor support for the inclusion of the rate in the diffusion network.
White (opacity of 0) represents any Bayes factors inferred to be less than 3, while a full
color (opacity of 1) represents any Bayes factors inferred to be greater than or equal to
100. C) Results of the PACT analysis for persistence in each host order for phylogeny
shown in panel A. D) For each host, we computed the proportion of Markov jumps
involving that host order in which that host was inferred as a source (jump coming from
that order) or as a sink (jump going to that order). The bars represent the variability
across the 3 replicates of equal orders subsamples.

The first introduction into North America is comprised of infections sampled in great
black-backed gulls (large shorebirds cluster, posterior probability = 0.69), consistent
with evidence of migratory gulls facilitating transmission from Europe (Figure 4A). This
shorebirds cluster contains 6 sequences from harbor seals sampled from outbreaks in
New England, resulting in a highly supported transition rate (BF = 537, posterior
probability= 0.99) from shorebirds to non-human mammals (6.07 markov jumps/year,
95% HPD: (3.21, 8.91), aligning with suggestions that these outbreaks are linked to
scavenging or environmental contamination by infected shore-birds[1,16]. After this initial
cluster of infections, multiple deep, internal nodes on the phylogeny are inferred in
Anseriformes with high posterior support (0.99), indicating that Anseriformes played an
important role in driving sustained transmission and dispersal across North America.
Across all replicates in both sampling regimes (6 total analyses), Anseriformes are
inferred at the root 2-3 times more frequently than in null, shuffled datasets (Table S19),
providing strong support for Anseriformes as critical drivers of epizootic transmission
beyond what is expected from their frequency in the data. We infer Anseriformes as the
predominant hosts seeding infections into other species (Figures 4B and 4D, Figure
S13, Table S6-11), with the highest rates to Galliformes (17.81 markov jumps/year
(95% HPD: (9.27, 26.02), BF = 1691, posterior probability = 0.99) and Strigiformes
(13.51 markov jumps/year, 95% HPD: 5.35, 22.87, BF = 232, posterior probability =
0.99). Aligning with speculation following mortality events in bald eagles[56], we also infer
a highly supported transition rate (BF = 127, posterior probability = 0.95) from
Anseriformes to Raptors, consistent with putative links between raptors and the
waterfowl they predate. Each of these patterns were preserved in each independent
subsample in both sampling regimes, indicating high robustness to sampling (Figure
S11-12).

We also infer support for transmission from Galliformes to Anseriformes (7.86 markov
jumps/year (95% HPD: (0.71, 14.97), BF = 147, posterior probability = 0.96),
Strigiformes, and nonhuman mammals. In this dataset, Galliformes primarily represent
domesticated poultry (98% of sequences), suggesting that transmission from domestic
birds back to wild birds and mammals may also have occurred, a hypothesis we
investigate in more depth below. However, lineages in Galliformes tended to be short-
lived, persisting for 0.26 years on average (95% HPD: 0.07, 0.33 years). Additionally,
Galliformes were inferred at the root less frequently than expected for their sampling
frequency (Table S19), and tended to cluster together more strongly than expected by
chance (p=0.0099, Table S18), consistent with a more limited role in transmission that
may have been confined to localized agricultural outbreaks. In contrast, viral lineages
persisted for the longest in Anseriformes (mean persistence time = 0.71 years, 95%
HPD: 0.42, 0.88 years) (Figure 4C), and Shorebirds (mean persistence time = 0.654
years, 95% HPD: 0.18, 1.04 years). Shorebird sequences were very highly clustered
with each other (AI=8.008, null=2.324, p=0.00999), suggesting some degree of
separation between viruses circulating in Shorebird populations, consistent with
ecological partitioning of low pathogenicity avian influenza viruses in these hosts [57]. Tip
shuffle analyses show mixed results for the Shorebirds, indicating a lack of consistent
evidence for their role in transmission relative to their sampling frequency in the dataset.
These data suggest that while Anseriformes, shorebirds, and Galliformes may all have
contributed to transmission events to other species, that Anseriformes were the

predominant drivers of longer-term persistence and spread to other hosts in the time
period analyzed.
500
One surprising result was inference of raptors as a low-frequency, but statistically well
supported source population to Anseriformes (5.18 markov jumps/year (95%HPD:
(0.36, 9.27), BF = 39, posterior probability = 0.87). Previous characterizations of HPAI in
Raptors during the 2014/2015 outbreak in North America showed mortality events and
neurological symptoms in wild raptors[58], while serological evidence of infections in bald
eagles have indicated exposure to influenza A viruses in 5% of birds tested between
2006 and 2010[59]. In the ongoing panzootic, raptors represent the third most prevalent
group in wild bird detections in Europe (12% of detections) and second most detected
group in North America (20.3%)[13,60]. Tip shuffle results for both sampling regimes
indicate that raptors are generally less probable at the root than expected based on
their sampling frequency, indicating that the clustering of genetic sequences supports a
limited role for epizootic transmission. Thus, while raptors may have been both heavily
impacted and sampled at a high frequency, these data suggest that raptors were
unlikely key drivers of epizootic spread. Future work to better establish the reasons for
such high case numbers among raptors and to investigate their potential links to
Anseriformes will be necessary for formulating wildlife management strategies.
517
We found limited support for non-canonical host groups (songbirds, owls, and
nonhuman mammals) in seeding infections in other species. Passeriformes (songbirds),
Strigiformes (owls), and mammals each primarily served as sinks for viral diversity
(Figure 4B,D), with transitions inferred with low magnitude and weak support (Figure
4B). Summing the number of jumps originating from wild canonical (Anseriformes,
shorebirds), wild noncanonical (Passeriformes, Strigiformes, raptors, mammals), and
Galliforme (domestic) hosts confirm that noncanonical hosts primarily acted sinks that
were far likelier to receive virus than propagate it onward (Figure S14), supporting short,
terminal transmission chains that did not lead to long-term persistence (Figure 4C and
Figure S15). Mammal sequences cluster across the entire diversity of the phylogeny
(Figure 4A) and are not associated with one particular cluster of viruses, indicating that
mammal infections were not confined to a particular viral lineage, supporting very short
persistence times of 0.22 years (95%HPD:(0.088, 0.328)), and only one strongly
supported transition rate to Anseriformes (BF = 53, posterior probability = 0.89).
Instead, these findings are most compatible with a model in which wild mammals and
other non-canonical species are infected by direct interaction with wild birds, likely
related to scavenging and predation behavior[14].
535
Taken together, these data suggest that despite high case numbers in several unusual
wild hosts, non-canonical species generally played minor roles in transmission across
the continent or to other species. Instead, epizootic transmission was most strongly
supported in Anseriformes. We infer Anseriformes as predominant drivers of longer-
term persistence and spread to other hosts across multiple independent analyses using
distinct sampling regimes, supporting surveillance in these species for capturing trends
in viral diversity and spread. Our data also suggest some role for shorebirds in
supporting persistent viral transmission. Future work to disentangle the relationship
between viral spread in these two key host groups may further resolve their utilities for
surveillance and monitoring.

Agricultural outbreaks were seeded by repeated introductions from wild birds

From 2022 to mid-2025, the US culled over 160 million domestic birds, with agricultural losses estimated between $2.5 to $3 billion USD [61]. Understanding the degree to which outbreaks in agriculture have been driven by repeated introductions from wild birds vs. sustained transmission in agriculture is critical for improving surveillance and biosecurity practices. However, differences in sampling between wild and domestic birds challenge this goal. While domestic birds comprise 11% of all detections, they comprise 23.2% of sequences, making them overrepresented in available sequence data. In contrast, though a higher number of detections and sequences have been deposited for wild birds, wild birds are likely to be heavily under sampled due to the challenge of sampling wildlife[9,31]. Finally, while each detection in wild birds represents a single infected animal, domestic bird detections usually represent a single infected farm, where the true number of infected animals is unknown. Given these challenges, we designed a "titration" analysis to measure the impact of varying degrees of sampling on transmission inference between wild and domestic birds. We first generated a dataset composed of equal numbers of domestic and wild bird sequences (using all 270 available domestic bird sequences and 270 randomly sampled wild bird sequences) sampled between November 2021 and August 2023. By setting the number of sequences from each group equal, we force the inference to be driven by the sequence data itself, rather than the sampling regime. Next, we added in progressively more wild bird sequences until we reached a final ratio of domestic to wild sequences equal to 1:3, which approximates the ratio of detections in domestic and wild birds during the epizootic (1026 domestic detections vs. 3078 wild detections for study time period). In total, we generated 5 datasets with the following ratios of domestic to wild bird sequences: 1:1, 1:1.5,1:2,1:2.5, and 1:3 (see Methods for more details). For each dataset, we applied a discrete trait diffusion model to infer transmission between wild and domestic birds. The goal of this analysis was two-fold. First, we aimed to determine whether either domestic or wild birds would be inferred as the primary source population, and whether that inference would be robust to our choice in sampling regime. Second, by titrating in sequences at varying degrees, we hoped to assess whether the inferred number of transitions between hosts stabilized at a certain ratio. If inferred transitions increase linearly with sequences and never stabilize, this would indicate that more surveillance data is necessary. If not, this provides evidence that adding additional data does not result in altered results, suggesting that the currently available data may be sufficient for estimating dynamics within this time period.

583
Figure 5. **Outbreaks in domestic birds were seeded by repeated introductions**
from wild birds, with some onward transmission A) Phylogenetic reconstruction
where taxa and branches colored by wild or domestic host status containing a 1:3 ratio
of domestic to wild bird sequences (n=1080). B) Number of transitions from a given trait
to another trait inferred through ancestral state reconstruction for each titration. C)
Results of the PACT analysis for persistence in domestic and wild birds for each
titration.
591
When domestic/wild sequences were included in equal proportions, the root of the
phylogeny and majority of internal nodes are inferred as wild birds, suggesting that wild
birds are inferred as the primary source in the outbreak even when sampled evenly
(Figure S16A). Wild birds are inferred at the root of the tree at a far higher probability
than expected from their sampling (posterior probability = 0.895 in empirical data vs.
0.482 in tip shuffled data), while domestic birds are underrepresented (Table S19).
Under equal sampling, this result is likely driven by higher genetic diversity among
viruses sampled from wild birds, consistent with a large, source population. Within the
background of wild bird sequences, domestic bird sequences form highly clustered
groups (AI=23.096, p=0.0019, Table S18), consistent with some transmission between
domestic birds. Transmission is inferred bi-directionally, with similar magnitudes of
transmission inferred from domestic to wild birds, and from wild to domestic birds
(Figure 5B, Figure S17, Table S12). If these patterns represent the true transmission

history in the epizootic, we reasoned that these patterns should remain intact even
when additional wild bird sequences are added to the tree. If not, then we hypothesized
that some of domestic bird clusters may be disrupted as more wild bird sequences are
added into the tree.
609
As wild sequences were progressively added into the tree, most domestic-only clusters
became smaller, broken up by wild sequences that interspersed within these clades
(Figure S16A-E). The "breaking up" of these domestic clusters results in inference of
more transitions from wild to domestic birds, and fewer transitions from domestic to wild
birds (Figure 5B-C, Figure S17). The largest changes in the inferred transitions occurred
between the 1:1 and 1:2.5 titrations, with minimal to no changes observed between
transitions inferred in the 1:2.5 and 1:3 datasets, suggesting stability in the inferred
transitions towards the end of the experiment (Table S12). The phylogeny of the final
dataset (using a 1:3 ratio of domestic to wild sequences) shows 106 clusters of
domestic sequences (Figure 5A, Figure S18-19, Table S12) each inferred as a unique
introduction of H5N1 from wild birds into domestic birds. Among these domestic-only
clusters, we calculate that lineages persisted for ~4.5 months on average (95% HPD:
2.7, 5.63). In contrast, we infer limited transmission from domestic birds back to wild
birds (Figure 5B, Table S12), with only 4 inferred introductions from domestic birds to
wild birds (Table S12). Viral lineages in wild birds also persisted for over twice as long
as those in domestic birds, persisting for ~10 months (95% HPD: 5.7, 14.07) (Figure
5C).
627
Commercial turkey operations have been heavily impacted during the epizootic,
comprising 53.7% of all detections on commercial farms[62]. However, the presence of
wild turkeys throughout North America makes categorizing turkey sequences as
domestic or wild status ambiguous. 98% of all turkey sequences are not associated with
metadata on domestic/wild status, and thus were excluded from the previous analysis.
However, epidemiologic data provides some hint that most deposited turkey sequences
likely stem from domestic outbreaks. Among case detections during the study period,
only 139 were reported in wild turkeys, representing 1.5% of all wild bird detections. In
contrast, commercial turkey outbreaks comprised 28.5% of all domestic detections in
the study period, suggesting that unlabelled turkey sequences are most likely to have
come from domestic birds. While this data is not conclusive, we performed an additional
analysis to determine whether our exclusion of turkey sequences (that are likely
domestic) may have biased our results. In this analysis, any turkey sequence not
labeled as "wild turkey" was categorized as "domestic" and combined with the other
domestic bird sequences, resulting in a dataset in which ~20% of domestic bird
sequences were turkeys. We then created a second titration analysis using these
domestic sequences to generate datasets of 1:1,1:1.5, and 1:2 ratios of domestic:wild
sequences. We then evaluated the impact of the inclusion of turkey sequences on
inferred transmission patterns between wild and domestic birds.
647
Inclusion of turkey sequences did not substantially change inferred transition rates
between wild and domestic birds but did increase the inferred persistence times of
domestic bird clusters (Figure S17, Table S12). As wild bird sequences were added to
the tree, we observed the same "breakup" of domestic clades as in the above analysis,
with more wild sequences leading to more inferred introductions into domestic birds,

and very few into wild birds. In both titration experiments, the final number of inferred
transmission events from domestic to wild birds was 4 (Table S12), indicating minimal
transmission back to wild species, regardless of whether turkeys were included (Figure
S20, Table S12). Inclusion of turkey sequences resulted in slightly longer inferred
persistence times in domestic birds, increasing inferred persistence by 1.29 and 1.54
months in the 1:1.5 and 1:2 titrations (Figure S22). While turkey sequences tended to
cluster closely with confirmed domestic sequences, they did form several large turkey-
only clusters on the tree (Figure S21), indicating some degree of separation between
turkey and non-turkey operations. To more directly estimate the role turkeys played in
transmission we built a final 1:2 (domestic:wild) dataset where the number of turkey and
domestic poultry were equal, which conformed to both a uniform and case proportional
dataset. Surprisingly, we found that while most introductions into turkey populations
stemmed from wild birds (42 transitions), we infer a high number of transmission events
between turkeys and other domestic birds. We estimate ~38 introductions from turkeys
to other domestic birds, and 18 in the opposite direction (Figure S23, Table S13),
indicating some degree of transmission between distinct poultry operation types. The
high degree of transitions from wild birds to turkeys, and from turkeys to other domestic
birds suggest a putative role for turkeys in mediating transmission between wild birds
and other types of domestic poultry operations. In line with the high number of turkey
detections during the epizootic, we infer comparatively larger clusters in turkeys than in
other domestic birds, suggesting that turkeys may have played a key role in
transmission among domestic populations.
675
Together, these data suggest a few important conclusions. First, wild birds are inferred
strongly as the major drivers of transmission. Wild birds were inferred as the major
source of viral dispersal even when heavily downsampled to be equal to domestic birds,
and independent of whether or not turkeys were included in the analysis, indicating
strong support for their role in H5N1 transmission. Second, regardless of sampling
regime and presence/absence of turkey sequences, we infer that outbreaks in
agricultural birds were driven by repeated, independent introductions from wild birds,
with some onward transmission between domestic operations. While the exact number
of inferred introductions vary across analyses (Table S12-13), we infer no fewer than
46, and as many as 113 independent introductions into domestic birds. When allowing
sampling frequencies to approximate detections (the 1:3 ratio analysis), we resolve a
higher number of introductions into domestic birds with shorter transmission chains,
though lineages still persisted for an estimated 4-6 months. Analysis of turkey and non-
turkey data indicate some degree of transmission between agricultural operations, and
a potential role for turkeys in mediating transmission between wild birds and non-turkey
domestic birds. Together, these results indicate that while both the 2014/2015 epizootic
and the 2022 epizootic involved transmission between domestic premises, that
transmission since 2022 has been fundamentally distinct. While the epizootic of
2014/2015 was started by a small number of introductions that rapidly propagated
between commercial operations, the epizootic since 2022 has been driven by intensive
and persistent transmission among wild birds, resulting in continuous incursions into
domestic bird populations that have continuously sparked new outbreaks in agriculture.
Wild birds thus played a critical role in agricultural outbreaks in North America from
2021-2023, marking a significant departure from past epizootics that may necessitate
updates to biosecurity, surveillance, and outbreak control.

701
702 *Spillovers to backyard birds occur earlier and slightly more often than those to*
703 *commercial birds*
704
The 2014/2015 H5Nx epizootic in the United States was driven by extensive
transmission in commercial poultry, prompting a series of biosecurity updates for
commercial poultry farms[5,63]. However, not all domestic birds are raised in commercial
settings. Rearing domesticated poultry in the home setting has become increasingly
popular in the United States, with an estimated 12 million Americans owning "backyard
birds" in 2022[64]. These birds have been heavily impacted during the ongoing epizootic,
579,005 backyard birds killed/euthanized, with some evidence for distinct transmission
chains circulating in backyard birds vs. commercial poultry[10]. Because backyard birds
generally experience less biosecurity than commercial birds and are more likely to be
reared outdoors[34], we hypothesized that spillovers into backyard birds may be more
frequent than spillovers directly into commercial poultry.
716
To test this hypothesis, we used a subset of sequences sampled between January and
May of 2022 that contained additional metadata specifying whether they were collected
from commercial poultry or from backyard birds (n= 275 from commercial poultry, n=85
from backyard birds). We then built a tree that included an approximately equal number
of sequences sampled from domestic and wild birds, but in which the domestic
sequences were evenly split between commercial poultry and backyard birds
(commercial birds = 85, backyard bird = 85, wild birds=193). As with the previous
analysis, we infer wild birds as the primary source population and infer repeated
introductions into commercial and backyard birds (Figure S24A). Unexpectedly though,
backyard bird sequences appeared to cluster more basally than commercial poultry
sequences, and sometimes fell directly ancestral to clusters of commercial poultry
sequences (Figure S24A). While every introduction into backyard birds was inferred to
descend from wild birds, 10 out of 26 introductions into commercial poultry were inferred
to descend from backyard birds (Figure S25A). This pattern was reproducible across
multiple replicate subsamples, indicating that it was independent of the exact subset of
wild bird sequences included in the tree. Given the debated role of backyard birds in
outbreaks in commercial poultry[65], we further explored two hypotheses that could
explain this pattern. The first is that backyard birds "mediated" transmission between
wild birds and commercial birds, possibly due to their greater likelihood of outdoor
rearing. Under this model, spillovers into backyard birds could be spread to commercial
populations via shared personnel, clothing, or equipment, resulting in sequences from
backyard birds preferentially nesting between wild and commercial bird sequences on
the tree. Alternatively, backyard birds could have simply been infected earlier on
average than commercial birds. If transmission in wild birds is persistent and high, and
backyard birds have a higher risk of exposure due to lessened biosecurity and
increased interactions with wildlife, then it could take less time for a successful spillover
event to occur and be detected in these birds, resulting in clustering that is more basal
in the tree.
745
To differentiate between these hypotheses, we performed a second titration analysis.
We started with the phylogeny including equal numbers of sequences from commercial
and backyard birds, allowing us to directly compare introduction patterns in these two

groups. Then, we added in progressively more wild bird sequences into the tree until all
available wild bird sequences were added into the tree. We added sequences in
increments of 25% (where % refers to percentage of total available wild bird sequences
in the time period), resulting in 3 additional analyses that included 50%, 75%, and 100%
of all available wild bird sequences. For example, the final dataset included 942
sequences, comprising 85 commercial bird, 85 backyard bird, and 772 wild bird
sequences. For each dataset, we inferred the number and timings of transmission
events between wild birds, commercial birds, and backyard birds across the posterior
set of trees. If backyard birds acted as mediators to outbreaks in commercial birds
(hypothesis 1), then the relationship between backyard birds and commercial birds
should remain unchanged as more wild bird sequences are added into the tree.
Alternatively, if backyard birds and commercial birds were infected independently
(hypothesis 2), then additional wild bird sequences should disrupt these clusters, and
intersperse between commercial and backyard bird sequences, resulting in more
independent introductions that occur earlier in backyard birds.

765
Figure 6. Backyard birds are infected by wild birds earlier than commercial birds.
A) Phylogenetic reconstruction of sequences collected between Jan 2022 and May
2023 with all available wild bird sequences and equal proportions of commercial and
backyard birds (n= 942) where taxa and branches are colored by host domesticity
status. B) Exploded tree view of the phylogeny showing the branches of transmission in
each domestic bird type following transmission from wild birds where subtrees represent

the traversal of a tree from the root to the tip where the state is unchanged from the
initial state (given by the large dot on left) to the tips represented by the smaller dots
representing continuous chains of transmission within a given state. C) Proportion of
trees from the posterior tree set with a given number of transitions from wild birds to
backyard birds and commercial birds (100% available wild sequences). D) Markov
rewards trunk proportion for domesticity status showing the waiting time for a given
status across branches of the phylogeny over time. E) Cumulative Markov jumps from a
given bird type to another over time where each line represents a single phylogeny from
the posterior sample of trees.
781
Throughout the experiment, wild bird sequences attached throughout the phylogeny,
disrupting nearly every backyard bird-commercial bird cluster, and dissolving the signal
of backyard bird to commercial bird transmission originally observed (Figure S24). The
final tree that included all available wild bird sequences resulted in inference of ~82
independent introductions from wild birds to domestic birds, with most clusters
containing only commercial (39 clusters) or backyard bird (43 clusters) sequences
(Figure 6A-B, Figure S24-25), suggesting that outbreaks in these groups were likely
seeded independently. Indeed, of the initial 10 transmission events inferred from
backyard birds to commercial birds, only 2 remained undisturbed in the final tree (Figure
6B, Figure S25). These two events represent outbreaks that occurred in the same state
within 6 days of each other, so it is plausible that these outbreaks are directly linked.
However, all other clusters were disrupted. As wild bird sequences were added into the
tree, the number of inferred introductions into backyard birds and commercial birds
diverged across the posterior trees for each titration (Figure S26), with backyard birds
experiencing slightly more introductions (mean = 42 introductions, 95% HPD: (35, 49))
than commercial poultry (mean = 39 introductions, 95% HPD: (32, 44)) (Figure 6C). This
is consistent with the rate of transmission inferred from wild birds into each domestic
population with a slightly higher rate to backyard birds (48.963 markov jumps/year, 95%
HPD: 40.163, 56.228, BF = 18,000, posterior probability = 0.99), then to commercial
birds (46.48 markov jumps/year, 95% HPD: 40.163, 52.785, BF = 18,000, posterior
probability = 0.99) (Table S14-S17). We infer a low transition rate from backyard birds to
commercial birds (2.678 markov jumps/year, 95% HPD:1.148, 4.59, BF = 4.58, posterior
probability = 0.7), and transitions from domestic birds back to wild birds were not
statistically well-supported.
806
To determine whether spillovers into backyard birds occurred earlier than those into
commercial poultry, we estimated the number of transitions between hosts across the
phylogeny ("Markov jumps") and the amount of time that is spent in each host between
transitions ("Markov rewards") [66,67]. We infer the highest mean duration in wild birds,
representing 87.7% of Markov rewards. Backyard bird and commercial bird sequences
showed lower, and similar reward time percentages of 5.3% and 7.0% respectively. The
Markov Reward proportion is calculated as the proportion of the phylogeny at a given
time being a given discrete state. By looking at the proportion a given state is over time
across the phylogeny we can provide a proxy for how long transmission has occurred in
each group between transition events. We find that early in the epizootic, transmission
in backyard birds slightly preceded transmission in commercial poultry (Figure 6D,
Figure S25). Enumeration of the cumulative number of transitions between hosts over
time across the posterior set of trees (Markov jumps[66]), again revealed that backyard

birds experienced slightly more jumps than commercial poultry (backyard birds = 43
introductions, 95% HPD: 36, 50; commercial birds = 39 introductions, 95% HPD: 32,
44), and that these introductions occurred earlier on average (Figure 6E). The lag time
between the cumulative transitions for backyard birds and commercial birds was ~9.6
days, indicating that transitions to backyard birds were inferred ~9 days earlier than
those to commercial birds. Comparison of detections and sequence availability in
commercial birds vs backyard birds show no apparent skewing in availability of samples
for each group in that time period, suggesting that this pattern is not simply due to an
abundance of earlier cases in backyard birds at that time (Figure S27). While this
pattern could also arise if cases in backyard birds are systematically reported earlier
than commercial poultry, data on testing turnarounds and enrollment in the US
indemnity payment register suggest that this is unlikely to be the case. Commercial and
backyard bird farms have almost identical lag times between case reporting and
confirmation (2.15 days for commercial birds and 2.4 days for backyard birds)[68], and
commercial poultry operations generally depopulate completely within 24 hours[69],
indicating testing and response in commercial operations that is efficient and slightly
earlier than in backyard birds. Comparison of the proportion of farms that received
indemnity payments (as a proxy for the percentage that submit to regular testing) shows
that while 511 of 168,048 commercial operations (0.3%) reported cases, that only 656
out of ~12 million backyard bird owners (0.0055%) were enrolled in the program[64,68,70].
While complete data on the distribution of backyard bird operations within North
America are sparse, these data suggest that backyard birds are likely undersampled
relative to commercial poultry, with somewhat lagged testing, making the early pattern
we observe compelling. Future studies utilizing expanded datasets across future
epizootic waves are necessary to confirm this pattern more broadly.
845
These data confirm that in the first 6 months of the epizootic, outbreaks in backyard bird
and commercial bird populations were generally seeded independently, with limited
evidence for transmission between them. We show phylogenetic evidence that
spillovers into backyard birds may have occurred slightly more frequently and earlier on
average than spillovers into commercial poultry, despite epidemiologic data that testing
is slower and more limited in backyard birds. Should future studies support these
findings more broadly, then backyard bird populations could potentially be investigated
as early warning signals for upticks in transmission. Future work will be necessary to
investigate the utility of this hypothesis.
855
856
Discussion
The panzootic of HPAI H5N1 that began in 2021 has impacted wildlife health,
agriculture, and human pandemic risk across the Americas. The North American
epizootic has been distinguished by the high number of infections in wildlife not usually
impacted by HPAI, its rapid dissemination across the Americas, and for its persistence
despite aggressive culling of domestic birds. In this study, we used a dataset of 1,824
HA sequences paired with curated metadata to reconstruct how H5N1 viruses were
introduced and spread. We show that H5N1 viruses were introduced ~9 independent
times into North America, with repeated incursions into the Pacific flyway that mostly
persisted transiently. A single introduction into the Atlantic flyway spread across the
continent within ~4.8 months, transmitting from East to West by wild, migratory birds.

Long-range dispersal and persistence were primarily driven by transmission in Anseriforme species, with infections in wild bird species like songbirds and owls primarily serving as dead-end hosts. Finally, we show through repeated subsampling experiments that unlike the epizootic of 2014/2015, the epizootic since 2021 was fundamentally distinct because transmission was driven by wild, migrating birds. Wild birds repeatedly re-seeded outbreaks in agriculture, sparking new introductions that persisted despite improvements in biosecurity and rapid culling responses. Outbreaks in commercial poultry and backyard birds were generally seeded independently, with introductions into backyard birds occurring ~9 days earlier on average, though future work is necessary to confirm these patterns more broadly. Together, our results suggest a critical shift in the ecology of highly pathogenic avian influenza viruses in North America. As the primary source of transmission shifts from poultry to wild migratory birds, the ecology of clade 2.3.4.4b viruses in North America may now follow patterns unfolding globally, where evolution is increasingly governed by wild bird movement, ecology, and reassortment. Our results implicate surveillance in wild Anseriforme species as potentially fruitful targets for ongoing tracking and response, and highlight the necessity of continued surveillance in wild birds for accurate outbreak reconstruction. Finally, our results provide explanations for the rapid expansion across the continent and for why culling domestic birds may no longer be sufficient for preventing outbreaks in agriculture. Instead, layered interventions, like improved biosecurity, separation of wild and domestic birds, and domestic animal vaccination may now be necessary for reducing future spillovers to agriculture, and by extension, humans.

Our study develops multiple lines of evidence that collectively support wild birds as a critical emerging source of highly pathogenic avian influenza transmission in North America. By directly modeling transitions between host groups based on domestic/wild classification, taxonomic order, and migratory behavior, paired with geographic analyses showing strong dispersal across flyways, we show that wild birds were key drivers of the epizootic. These results imply that continuous surveillance in wild birds may now be key for viral tracking and outbreak reconstruction, with our data pointing to Anseriformes as good potential targets. Recent modeling of HPAI risk in Europe identified *Anatinae* and *Anserinae* (within the order Anseriformes) species prevalence as the most consistent predictors of HPAI detection across seasons[71], in line with our findings. Our data suggest that non-canonical hosts acted as sinks for viral diversity, played limited roles in epizootic spread, and are unlikely to establish as long-term reservoirs, aligning with a model in which these species were opportunistically infected from more classical species. Test positivity data from fall 2022 showed test positivity in wild ducks of 81.8% in Newfoundland and Labrador, indicating incredibly widespread transmission in these species early on[72]. Thus, given limited resources for wildlife surveillance, our data suggest wild, migrating Anseriformes as a useful target for surveillance for risk assessment, vaccine strain selection, and tracking of viral evolution and spread.

Utilizing a large and geographically representative dataset, we estimate the number and timings of introductions in North America, and highlight the capacity of migratory birds to rapidly disseminate these viruses across the continent. Our analyses suggest that highly pathogenic H5N1 viruses may have been circulating in wild birds in North

America as early as September – October of 2021, one to two months prior to the first
reported detection. Following successful introduction into the Atlantic flyway, initial
epizootic spread proceeded rapidly between adjacent migratory flyways, spreading from
East to West within ~4.8 months. This rapid speed of dissemination contrasts with other
avian-transmitted viruses like West Nile Virus, which spread from the Northeast across
the continent over the course of 4 years[46]. We speculate that the rapid geographic
spread and strong signal of East to West diffusion could be explained by one or a
combination of the following factors: high inherent transmissibility of clade 2.3.4.4b
viruses in wild birds; rapid, long-range migration of wild bird species driving
transmission; and exponential spread among an immunologically naïve population of
wild birds in North America during early panzootic expansion[73,74]. Previous work has
shown that within-flyway viral population structure is more observable within short time
periods (< 5 years)[75], while transmission assessed over longer time scales is more likely
to span multiple flyways[76,77]. Continuous surveillance of wild species will be necessary
to determine whether the rapid degree of transmission, and strong clustering by flyways
observed early in the panzootic will lessen over time as immunity builds in avian hosts.
932
Though a single introduction into the Atlantic flyway accounted for most transmission in
the epizootic, the Pacific flyway experienced more independent incursions (~7).
Compared to other studies, we infer more 5 previously undescribed incursions[78] into the
Pacific, suggesting frequent viral flow between Asia and the Pacific coast of North
America that could be an important target for surveillance efforts. Despite this high
degree of flow, these introductions mostly did not persist long-term. Limited
transmission from the Pacific flyway could be explained by differential fitness of the
lineages introduced into the Pacific vs. Atlantic flyways, ecological isolation of the
Pacific flyway, or by differences in host distributions at the locations and times of these
incursions. Early after introduction, the European lineage introduced into the Atlantic
flyway reassorted with endemic, low-pathogenicity H5 viruses in North America,
resulting in a virus that exhibited altered tissue tropism in mammals[79]. Whether this
early reassortment event, or subsequent reassortments, produced viruses with
enhanced fitness in wild birds is currently unknown, but could explain the success of the
Atlantic flyway introduction. Alternatively, the failure of Pacific incursions to spread
onward could be explained by ecological isolation of the Pacific flyway, potentially due
to land features like the Rocky Mountains. Prior studies of low-pathogenicity avian
influenza show a strong influence of geography on restricting virus dispersal across the
North American continent, providing plausible support for this hypothesis[80–82]. Finally,
Pacific flyway introductions could have persisted for less time due to a lack of suitable
host species at the times and locations of the incursions, or simply by chance. Future
work will be necessary to differentiate among these hypotheses.
In this study, we find that outbreaks in agriculture were seeded by repeated
introductions from wild birds. This pattern held true regardless of sampling regime, and
aligns with anecdotal observations that clade 2.3.4.4b viruses are increasingly being
maintained by transmission among wild bird species more globally, including now in
North America[17,83]. These findings contrast with genomic and epidemiologic
investigation of the epizootic in 2014/2015, which implicated transmission between
commercial poultry operations via personnel, equipment, and clothing as the major
source of dissemination[5,84], prompting updates to recommended biosecurity
protocols[5,63]. Because the viruses circulating in 2014/2015 did not establish in local wild

bird populations, that epizootic was well-controlled by culling domestic flocks, and after culling 50.5 million domestic birds, the epizootic died out. In contrast, at the time of writing, detections in wild and domestic birds in North America have continued despite culling over 160 million domestic birds. We show that despite some persistence among domestic bird populations, that introductions into domestic populations ultimately led to transmission chains that died out, and that they generally did not result in re-introduction into wild birds. These relatively short transmission chains likely reflect improved biosecurity plans developed since 2015[5], and far more rapid and aggressive culling of domestic birds. Compared to 2014/2015, the median time from case detection to depopulation in this epizootic ranged from 4-51 hours[69], vs. 6 days previously [5], indicating a far more rapid culling response. Despite these improvements in biosecurity, the number of poultry outbreaks, and corresponding human spillovers, has been staggering since 2021. Our results provide an explanation for these phenomena. Compared to past outbreaks, transmission in agricultural settings in North America since 2021 have been fundamentally distinct because these viruses circulated efficiently in wild birds, allowing for rapid dispersal and continuous outbreak re-seeding. While improvement in biosecurity may have reduced farm-to-farm spread, the establishment of continuously circulating lineages in wild birds likely made this epizootic far more challenging to prevent and control. While current US and Canadian policy classifies H5N1 as a foreign animal disease that prioritizes culling, these new dynamics challenge the effectiveness of culling as the primary strategy for control. Our data suggest that future prevention of agricultural outbreaks may now require layered interventions that seek to reduce interactions between wild and domestic birds, paired with aggressive biosecurity between farms[20,22]. As highly pathogenic H5N1 viruses continue to circulate in North American wild birds, investment in control methods that reduce successful transmission between wild birds and agricultural animals, including potentially vaccination, should be explored.

Using a small dataset from the first 6 months of the epizootic, we find phylogenetic evidence that spillovers into backyard birds may have occurred slightly earlier and more frequently than those into commercial farms. We also show through extensive subsampling experiments that outbreaks on commercial and backyard premises were generally seeded independently, with limited viral flow between them. Though commercial poultry operations generally operate with higher degrees of biosecurity than backyard flocks, there are some documented locations that could allow for interaction between commercial poultry and wildlife[34]. Retention ponds on commercial poultry farms are frequently visited by wild waterfowl[85], while natural features such as water bodies and vegetation near residential coops and commercial production sites could also act as potential points of wild to domestic transmission. Though the role of backyard birds on avian influenza outbreaks on commercial farms is debated[65], our data suggest that backyard birds could potentially be investigated as sentinel species for transmission in wild birds. In the United States, commercial poultry operations are more likely to be enrolled in voluntary testing programs and indemnity programs that guarantee access to funds following depopulation. As a result, commercial poultry are generally more well surveilled and have more rapid response times for testing and depopulation. Though recent data on backyard bird rearing in the US is limited, a large survey of backyard bird populations from 2004 showed that backyard bird flocks often contain multiple species, usually have outdoor access, and that 60-75% of backyard

flocks experience regular contact with wild birds[34]. 38% of surveyed backyard flocks
were on a property that contained a pond that attracted wild waterfowl, and 40% have a
wild bird feeder on the property, providing clear, plausible links to water bird interaction.
Multiple surveys have shown that biosecurity precautions tend to be much more limited
in backyard bird populations, with 88% of backyard flocks using no precautions (shoe
covers, footbaths, clothing changes) at all[34]. Analysis of seropositivity to avian influenza
in backyard poultry in Maryland showed that exposure to waterfowl resulted in a ~3x
higher likelihood of harboring anti-influenza antibodies compared to birds not exposed
to waterfowl[86]. Early infections during the 2014/2015 epizootic were recorded in
backyard birds[87], though the overall rate of backyard bird detections have been
substantially higher in the current epizootic[88], presumably due to the difference in the
predominant species driving transmission (e.g., commercial poultry vs wild birds). Given
backyard birds' enhanced likelihood of interaction with wild birds, and our inference of
earlier spillovers in those groups, expanded studies may be useful for investigating
whether the patterns of earlier spillovers in these populations hold true more broadly. If
so, then backyard bird populations could be investigated as early warning sentinels for
increasing transmission in local wild bird populations. 93% of backyard flocks contain
<100 birds, and most backyard bird owners report raising birds for enjoyment and
eggs[34]. A subset of backyard bird owners may therefore present an opportunity for
community engagement that could potentially help identify early detections. If
engagement were successful, prompt reporting for illness and infections could be used
to alert other nearby operations that highly pathogenic avian influenza is circulating
locally, allowing for more advanced warning for heightened biosecurity.
1035
Sampling bias is pervasive across viral outbreak datasets, and no modeling approach
can completely overcome inherent biases in data acquisition. In this study, we
employed multiple, overlapping analyses to control for sampling biases, providing a
framework for performing phylodynamic analyses in the presence of uncertain sampling.
Highly pathogenic avian influenza data is rife with sampling issues. In the US, avian
influenza sampling of wild birds employs 3 distinct methods (hunter harvest, sick and
dead, and live birds), each with unmeasured biases. Only wild Anseriformes are
sampled live or hunter harvested, while all other host groups are sampled sick or dead,
likely skewing detections towards birds with dedicated rescue services and those
located near humans. In domestic birds, detection depends on producer identification of
illness and reporting, and subsequent testing, which likely varies across production
types, locations, and premises. While we accumulated as much data as possible to
contextualize how cases are sampled and sequenced in North America, it is impossible
to know the true case burden in either domestic or wild birds. Because of this, we opted
to investigate multiple distinct subsampling approaches (equal and proportional), report
results that are consistent between them, and to employ statistical tests for trait bias to
most accurately estimate transmission dynamics. Equal sampling regimes rely
inherently on the diversity in sampled sequences to make inferences about sources of
viral diversity, while proportional regimes bring the data in line with model assumptions
that sequence numbers are proportional to cases. For analyses of host orders and
wild/domestic bird transitions (via titration tests), we employ both. Beginning with an
equal sampling regime, we show that the genetic diversity of viruses circulating in
Anseriformes and wild birds is sufficient to infer them as source populations, patterns
which remained under proportional sampling regimes. In the titration tests, we show that

while the central conclusions remained true, that the precise number of transitions
between wild and domestic birds depends heavily on sampling numbers, providing a
clear argument for continuous surveillance in wildlife, and a warning for over-confidence
in estimating the particular numbers of transitions between groups. Finally, we
quantified the level of imbalance between sampled traits across each discrete trait
analysis, and find limited evidence that sampling biases drove our inferences (Table
S17-18). Still, these methods have important caveats. An equal subsampling regime
might over-represent a truly under-impacted group, while a proportional subsampling
might over-represent a more well sampled group. Though use of replicates allowed us
greater confidence in our results, all phylodynamic inferences are limited by sequence
data availability, and results could change if future data become available. Additional
sequence data could result in inference of more introductions into North America, or
altered numbers of transitions between domestic and wild birds. Our analyses only
employ HA sequences, a caveat that could result in slightly different numbers of inferred
transitions if full genomes were used. Accurately distinguishing between hypotheses of
epizootic spread (e.g., whether agricultural outbreaks are driven by introductions from
wild birds or by from farm-to-farm spread) depends on adequate sequence data from
wild birds, without which transmission inference is impossible. As H5N1 viruses
continue to evolve and spread globally, investment in surveillance strategies that
capture circulating diversity among wild birds will likely be critical for accurately tracking
viral evolution, prioritizing vaccine strains, and contextualizing new emergence events,
like the recent outbreaks in dairy cattle.
1082
Taken together, our findings implicate a critical shift in highly pathogenic avian influenza
ecology in North America, with wild birds playing the central role in transmission and
dispersal in the 2021-2023 epizootic. Persistent and intensive transmission in wild birds
provides an explanation for the rapid cross-continental spread, and continued
agricultural outbreaks despite aggressive culling. Our results highlight the utility of wild
bird surveillance for accurately distinguishing hypotheses of epizootic spread, and
suggest continuous surveillance as critical for preventing and dissecting future
outbreaks. Our data underscore that continued establishment of H5N1 in North
American wildlife may necessitate a shift in risk management and mitigation, with
interventions focused on reducing risk within the context of enzootic circulation in wild
birds. At the time of writing, outbreaks in dairy cattle highlight the critical importance of
modeling the ecological interactions within and between wild birds and domestic
production. Future work to effectively model viral evolution and spread hinges critically
on effective surveillance across wild and domestic species to capture key transmission
pathways across large geographic scales. Ultimately, these data are essential for
informing biosecurity, outbreak response, and vaccine strain selection.
1099
1100
1101 **Materials and Methods**
1102
1103 Dataset collection and processing
1104
1105 *Genomic data processing and initial phylogenetics*
1106

We downloaded all available nucleotide sequence data and associated meta-data for
the Hemagglutinin protein of all HPAI clade 2.3.4.4b H5Nx viruses from the GISAID
database on 2023-11-25[89]. For each subset of the data described for further
phylodynamic modeling the following process was followed. We first aligned sequences
using MAFFT v7.5.20, sequence alignments were visually inspected using Geneious
and sequences causing significant gaps were removed and nucleotides before the start
codon and after the stop codon were removed[90,91]. We de-duplicated identical
sequences collected on the same day (retaining identical sequences that occurred on
different days). We identified and removed temporal outliers for all genomic datasets by
performing initial phylogenetic reconstruction in a maximum likelihood framework using
IQtree v.1.6.12 and used the program TimeTree v 0.11.2 was used to remove temporal
outliers and to assess the clockliness of the dataset prior to Bayesian phylogenetic
reconstruction[92,93]. This resulted in a dataset of 1824 sequences that were used in
further analyses (Figure S24).

1121
1122 *AVONET database*
1123
We downloaded the AVONET database for avian ecology data and merged it to
available host metadata from GISAID for each sequence[48]. We used the species if
provided to match the species indicated in the AVONET database. If host metadata in
GISAID was defined using common name for a bird, we determined the taxonomic
species name and used that for further merging with the AVONET data (e.g. "Mallard"
was replaced with *Anas platyrhynchos*) for the given region to match the species to its
respective ecological data. Domesticity status (whether a sequence was isolated from a
wild host or a domestic host) was determined using available metadata downloaded
from GISAID using the 'Note' and 'Domestic_Status' fields in sequence associated
metadata. Additionally, if a given sequence strain name (in the field 'Isolate_Name')
indicated domestic status (e.g. A/domestic_duck/2022) these sequences were labeled
as belonging to domestic hosts.

1136
1137 *Detections by USDA*
1138
1139 Data for detections of HPAI in North America were collected from USDA APHIS.
1140 Reports for mammals, wild birds, and domestic poultry were all downloaded (download
1141 date: 2023-11-25)[60].

1142
1143 <u>Phylodynamic analysis</u>
1144
1145 The following Bayesian phylogenetic reconstructions and analyses were performed
1146 using BEAST v.1.10.4[94].

1147
1148 *Empirical tree set estimation and coalescent analysis.*
1149
We performed Bayesian phylogenetic reconstruction for each dataset prior to discrete
trait diffusion modeling to estimate a posterior set of empirical trees. The following priors
and settings were used for each subset of the sequence data. We used the HKY
nucleotide substitution model with gamma-distributed rate variation among sites and
lognormal relaxed molecular clock model[95,96]. The Bayesian SkyGrid coalescent was

used with the number of grid points corresponding to the number of weeks between the
earliest and latest collected sample (e.g for a dataset collected between 2021-11-04
and 2023-08-11 we would set 92 grid points)[40]. We initially ran four independent MCMC
chains with a chain-length of 100 million states logging every 10000 states. We
diagnosed the combined results of the independent runs diagnosed Tracer v1.7.2. to
ensure adequate ESS (ESS > 200) and reasonable estimates for parameters[94]. If ESS
was inadequate additional independent MCMC runs were run increasing chain length to
150 million states, sampling every 15000 states were performed. We combined the tree
files from each independent MCMC run removing 10-30% burn-in and resampling to get
a tree file with between 9000 and 10000 posterior trees using Logcombiner v1.10.4. A
posterior sample of 500 trees was extracted and used as empirical tree sets in discrete
trait diffusion modeling.
1167
1168 *Discrete trait diffusion analysis*
1169
1170 *Dataset subsampling and definition of discrete traits*
1171
We characterized the geographic introduction of HPAI into North America by randomly
sampling 100 sequences from Europe and Asia for each year between 2021-2023 (total
300 non-North American) and all available North American sequences across the study
period. Following removal of temporal outliers this resulted in a dataset of n= 1927
sequences annotated by continent of origin. The sequence data available from North
America broken down by country is as follows: United States (1590), Canada (224),
Honduras (2), Costa Rica (5), and Panama (1).
1179
To characterize geographic transmission within North America, following introduction,
we constructed a dataset of sequences subsampled based on migratory flyway. We
used place of isolation data to match the US state or Canadian province the sequence
was collected from with the respective U.S. Fish and Wildlife Service Migratory Bird
Program Administrative Flyway[44]. We subsampled 250 sequences for each flyway
(Atlantic, Mississippi, Central, and Pacific) to create a dataset of 1000 sequences
collected between November 2021 and August 2023. In addition to USFWS flyways we
defined 4 geographic regions going North to South based on latitude lines with the
following delineations for each group. We divided North America into 4 regions
segregated by latitude, with the northernmost group above the 49N parallel and the
southernmost group below the 36N parallel. We then sampled 916 sequences evenly
across these categories and inferred transitions between these regions.
1192
We classified sequences by host taxonomic order, inferring the host species using
designations in the strain name and/or metadata to match species records in
AVONET[48]. To ensure that each discrete trait had an adequate number of samples for
the discrete trait analysis of host orders we combined orders in two instances based on
taxonomic and behavioral similarity. The order Falconiformes (n=14), which represents
falcons, was added to Accipitriformes (n=363), which includes other raptors such as
eagles, hawks, and vultures. Pelecaniformes (n=34) which includes pelicans were
grouped with Charadriiformes (n = 74, shorebirds and waders) due to their similar
aquatic lifestyles and behaviors. Mammals were kept as a broad non-human
classification as most samples were of the order carnivora (foxes, skunks, bobcats etc.),

apart from samples of dolphins (Artiodactyla) and Virginia opossum (Didelphimorphia).
The following orders were omitted due to low number of sequences: Rheaforimes
(n=2), Casuariiformes (n=1), Apodiformes (n=2), Suliformes (n=7), Gaviiformes (n=1),
Gruiformes (n=1), Podicipediformes (n=1).
1207
We randomly subsampled 100 sequences for each host order between 2021-11-04 and
2023-08-11 resulting in a dataset of n=655 sequences where all isolates for host orders
with less than 100 samples, Passeriformes (n = 57) and Strigiformes (n=99) (removing
one temporal outlier), were used (Figure S25). We repeated this random subsampling
three times resulting in three separate datasets. We additionally performed three sub-
samples of sequences based on the proportion of detections in each host order group
which were collected between 2021-11-04 and 2023-08-11. Three random proportional
samples were taken each with the following number of sequences for each group:
Accipitriformes =133, Anseriformes = 342, Passeriformes = 12, Nonhuman-mammal =
16, Galliformes = 83, Charadriiformes = 40, Strigiformes = 29 (total n=655 sequences).
1218
We defined discrete traits for use in discrete trait diffusion modeling based on the
available sequence metadata and merged AVONET data. In addition to taxonomic
order, we defined migratory behavior. Birds were classified as sedentary (staying in
each location and not showing any major migration behavior), Partially migratory (e.g.
small proportion of population migrates long distances, or population undergoes short-
distance migration, nomadic movements, distinct altitudinal migration, etc.), or Migratory
(majority of population undertakes long-distance migration). We subsampled sequences
based on migratory behavior including nonhuman-mammals and domestic birds to
create a subsample of 500 sequences with equal sampling across behavior groups.
1228
1229 *Discrete trait modeling framework*
1230
For each discrete trait dataset, we used an asymmetric continuous time Markov chain
discrete trait diffusion model and implemented the Bayesian stochastic search variable
selection (BSSVS) to determine the most parsimonious diffusion network[41]. We inferred
the history of changes from a given trait to another across branches of the phylogeny,
providing a rate of transitions from A to B/year for each pair of trait states. When
reporting these results, we refer to state A as the source population/state and B as the
sink population/state. We implemented the Bayesian Stochastic Search Variable
Selection which allows us to determine which rates have the highest posterior support
by using a stochastic binary operator which turns on and off rates to determine their
contribution to the diffusion network. In addition to the discrete trait diffusion rate
employed a Markov Jump analysis to observe the number of jumps between discrete
states across the posterior set of trees and estimated the Markov Rewards to determine
the waiting time for a given discrete trait state in the phylogeny[66,67] . We calculate the
transition rate as a realization of the CTMC process by dividing the number of markov
jumps by the tree height (branch length from the earliest tip to the root of the tree) , and
separately, by tree length (sum of all branch lengths). For each pairwise transition rate,
we calculate the level of Bayes Factor (BF) support that the given rate has. The BF
represents the support of a given rate. The BF is calculated as the ratio of the posterior
odds of the given rate being non-zero divided by the equivalent prior odds which is set
as a Poisson prior with a 50% prior probability on the minimal number of rates

possible[41]. We use the support definitions by Kass and Rafferty to interpret the BF
support where a BF > 3 indicates little support, a BF between 3 and 10 indicates
substantial support, a BF between 10 and 100 indicates strong support, and a BF
greater than 100 indicates very strong support[97].
Empirical sets were used with the discrete traits defined for each sequence to perform
discrete trait diffusion modeling. Each discrete trait model was implemented using three
independent MCMC chains with a chain length of 10 million states, logging every 1000
states. Runs were combined using LogCombiner v.1.10.4. subsampling a posterior
sample of 10,000 trees/states. The Bayes Factor support for transition rates were
calculated using the program SPREAD3 [98]. Maximum clade credibility trees were
constructed using TreeAnnotator v1.10.4.
1262
1263 *Domestic/Wild titration analysis*
1264
To study the impact of sampling of wild birds on the estimation of rates between
domestic and wild birds we created five separate datasets with varying numbers of wild
birds for sequences collected between 2021 and 2023. We randomly sampled 270
domestic sequences and 270 wild sequences as the initial 1:1 ratio dataset. We then
made four more datasets increasing the number of wild sequences by a factor of 0.5
(adding 135 wild sequences) resulting in a final "titration" of 1:3 domestic to wild
sequences (n=1080). We applied a two-state asymmetric CTMC discrete trait diffusion
model where sequences were labeled as domestic or wild. All priors and model
parameters selected are the same as those described in the empirical tree set
description above. To study the impact of the inclusion of turkeys in the transmission
between domestic and wild populations we annotated all unannotated sequences
collected from turkeys as domestic. We then created three datasets starting with 525
domestic and 525 wild bird sequences, adding 263 sequences to successive titrations
resulting in 1:1,1:1.5, and 1:2 (domestic:wild) sequence datasets with a final titration
size of 1,575 sequences. We again applied a two-state asymmetric CTMC discrete trait
diffusion model where sequences were labeled as domestic or wild with all priors and
model parameters selected are the same as those described in the empirical tree set
description above. To determine whether the proportion of turkeys to other domestic
birds would impact the results of the previously described titration analysis we built a
dataset with where the domestic bird group had equal numbers of turkey and domestic
(non-turkey) sequences. This dataset included 173 turkey, 173 domestic bird, and 692
wild bird sequences totaling 1038 sequences. We applied an asymmetric CTMC
discrete trait diffusion model using a BSSVS for a three-trait model with the following
states: wild birds, domestic birds (not turkey), and turkey. We performed three
independent runs of this analysis using the models and parameters described in the
empirical tree analysis section above. All titration replicates were performed using an
MCMC chain length of 100 million states sampling every 10,000 states.
1292
1293 *Commercial, backyard, wild bird titration analysis*
1294
Metadata and annotated sequences were made available describing sequences as
being from backyard birds for sequences collected in early 2022 which distinguished
them from commercial poultry (previously all sequences being determined domestic)[10].
We used this metadata to create a dataset with equally sampled backyard birds and

commercial birds (n= 85 for each bird type) and then added all available wild birds
(n=722) in 25% increments creating four separate datasets for sequences collected
between Jan 2022 and June 2022. This resulted in a final dataset of n= 942 sequences.
We performed discrete trait diffusion modeling using an asymmetric CTMC diffusion
model described in the previous section for sequences labeled as backyard bird,
commercial bird, and wild bird.
1305
1306   *Extraction of phylogenetic metrics*
1307
1308   We calculated the transitions between states across branches of phylogenies estimated
1309   from ancestral state reconstructions using the Baltic python package[28]. To calculate the
1310   persistence of a given discrete trait we used the program PACT v0.9.5. which calculates
1311   the persistence of a trait by traversing the phylogenetic tree backwards and measuring
1312   the amount of time a tip takes to leave its sampled state[99].
1313
1314
1315   **Assessment of Sampling Bias**
1316
BaTs analysis
To determine if the discrete traits analyzed correlated with shared ancestry in the
phylogeny, we employed tip trait association tests implemented in the Bayesian Tip-
association Significance (BaTs) program v1.0[45]. This program assesses the
phylogenetic structure of discrete traits across viral lineages using three metrics: the
association index (AI), parsimony score (PS), and (maximum monophyletic clade size
(MC). The AI measures the imbalance of internal nodes of a phylogeny for a given set
of traits. The PS calculates the number of state changes in the phylogeny. The MC
measures the maximum number of tips belonging to a monophyletic clade for each
discrete trait of interest. These metrics are calculated for the phylogeny as tips are
randomly swapped to create a null distribution to compare against. Taken together
these metrics quantify the degree of clustering within the phylogeny with Lower AI and
PS values indicating stronger phylogenetic structure, suggesting that closely related
taxa tend to share the same trait, whereas higher values indicate weaker structure and
more frequent transitions between trait states. Statistical significance was assessed by
comparing observed values against a null distribution generated through randomization,
with p-values reported for each test. All discrete trait groupings showed evidence for
clustering by trait, supporting the use of trait modeling across the tree.
1335
1336   *Tip shuffle analysis*
1337
To assess the sensitivity of each of our discrete trait reconstructions to differences in
sampling between groups, we implemented a modified version of a tip swap analysis[55].
As originally developed, a tip swap analysis attempts to assess the impact of trait
sampling on discrete trait measurements. An operator is implemented within the MCMC
chain that randomly picks pairs of tips and swaps their trait values, thus generating a
posterior set of trees among which pairs of trait assignments have been randomly
swapped. The probability of each state at the root is then computed, and compared to
the inferred root state probabilities in the empirical data. Because the root state
probabilities in randomized datasets should primarily reflect the frequency of each trait

in the analysis, empirical results that differ substantially from this null distribution are
interpreted as evidence that the sequence data is informing the analysis beyond what is
expected based on trait frequencies alone. Thus, traits for which the root state
probability differs considerably from the root state probability in the null data are
frequently interpreted as being informed by the data, rather than sampling bias. While
this approach has been shown to perform well on small phylogenies[50,77], the strategy of
swapping single pairs of tips poses challenges for larger trees. In our flyways dataset
which includes ~1000 tips, we found that even with extremely high operator values
(4000), the traditional tip-swap analysis resulted in a posterior set of trees in which the
majority of tips (93.3%) remained assigned to their true state at least 50% of the time,
resulting in a null dataset that was only partially randomized. We believe this is due to
the high number of tips in our analysis, resulting in only an extremely small fraction of
tips randomized at any given step in the MCMC chain. To overcome this limitation, we
instead performed a randomized tip shuffle analysis. Using the empirical set of trees
inferred for each discrete trait analysis, we generated 100 null datasets in which we
shuffled the trait assignments randomly across the tips. In this approach, we preserve
the phylogenetic tree topology and the ratio of samples from each group, but shuffle
their assignments at the tips. For each discrete trait analysis, we generated 100 distinct
shuffled versions of the empirical trees, reran the analysis, and summarized the
resulting posterior distribution by inferring a maximum clade credibility tree. We then
computed the root state probabilities for each trait for each mcc tree, and computed the
mean root state probability across all 100 replicates. This computed mean is reported in
Table S19, in the column labelled "Mean root state probability across 100 datasets with
randomly shuffled tip states." Finally, we compared these null values to the root state
probabilities calculated for each group in the empirical data (reported as "Root stat
probability in Empirical data" in Table S19). As expected, the root state probabilities
inferred in the shuffled datasets are proportional to the number of sequences included
for each group. For the analyses using an equal sampling regime (Migration, Flyway,
Host Orders Equal, and initial titration tests), this leads to approximately equal expected
root state probabilities across groups. In contrast, the root state probabilities in the
empirical data generally differ significantly from expectation, suggesting that the
phylogenetic results are informed by the genetic data rather than from sampling alone.
1379
1380 The results of BaTS analyses and tip shuffling analyses for each discrete trait in this
1381 study can be found in the supplemental material (Figure S30-S39), Table S18-19).
1382
1383
1384 **Data and code availability:**
1385 All analytical scripts, metadata annotations, and BEAST XMLs used in this analysis can
1386 be found at the following GitHub repository: https://github.com/moncla-lab/North-
1387 American-HPAI
1388
1389 All data that was used in this analysis were sourced from public databases.
1390 Acknowledgement table for GISAID isolates used in this analysis can be found in Table
1391 S20.
1392
1393 Several of the analyses presented have also been publicly made available using a
1394 maximum likelihood framework through the Nextstrain pipeline and a narrative of this

work can be found in the following link:

Acknowledgements

We would like to thank Mia Kim Torchetti for her helpful feedback and discussion about our results. This work was supported by NIH R00-AI147029-05 and by funding from the Centers of Excellence for Influenza Research and Response (CEIRR), funded by NIH 75N93021C00015. LHM is a Pew Biomedical Scholar and is supported by NIH R00-AI147029-05. LD is supported by NIH 75N93021C00015.

Citations:

[1]

1. Puryear, W. B. & Runstadler, J. A. High-pathogenicity avian influenza in wildlife: a changing disease dynamic that is expanding in wild birds and having an increasing impact on a growing number of mammals. *J. Am. Vet. Med. Assoc.* 1–9 (2024) doi:10.2460/javma.24.01.0053.

2. Lycett, S. J. *et al.* Genesis and spread of multiple reassortants during the 2016/2017 H5 avian influenza epidemic in Eurasia. *Proc. Natl. Acad. Sci.* **117**, 20814–20825 (2020).

3. Smith, G. J. D. *et al.* Characterization of Avian Influenza Viruses A (H5N1) from Wild Birds, Hong Kong, 2004–2008. *Emerg. Infect. Dis.* **15**, 402–407 (2009).

4. Alexander, D. J. Summary of Avian Influenza Activity in Europe, Asia, Africa, and Australasia, 2002–2006. *Avian Dis.* **51**, 161–166 (2007).

5. *Final Report for the 2014–2015 Outbreak of Highly Pathogenic Avian Influenza (HPAI) in the United States*. https://www.aphis.usda.gov/media/document/2086/file (2016).

6. Caliendo, V. *et al.* Transatlantic spread of highly pathogenic avian influenza H5N1 by wild birds from Europe to North America in 2021. *Sci. Rep.* **12**, 11729 (2022).

7. Gass, J. D. *et al.* Ecogeographic Drivers of the Spatial Spread of Highly Pathogenic

Avian Influenza Outbreaks in Europe and the United States, 2016–Early 2022. *Int. J.*

Environ. Res. Public. Health **20**, 6030 (2023).

8. Erdelyan, C. N. G. *et al.* Multiple transatlantic incursions of highly pathogenic avian

influenza clade 2.3.4.4b A(H5N5) virus into North America and spillover to mammals.

Cell Rep. **43**, 114479 (2024).

9. Klaassen, M. & Wille, M. The plight and role of wild birds in the current bird flu

panzootic. *Nat. Ecol. Evol.* **7**, 1541–1542 (2023).

10.   Youk, S. *et al.* H5N1 highly pathogenic avian influenza clade 2.3.4.4b in wild and

domestic birds: Introductions into the United States and reassortments, December

2021–April 2022. *Virology* **587**, 109860 (2023).

11.   Spackman, E., Pantin-Jackwood, M. J., Lee, S. A. & Prosser, D. The

pathogenesis of a 2022 North American highly pathogenic clade 2.3.4.4b H5N1 avian

influenza virus in mallards (*Anas platyrhynchos*). *Avian Pathol.* **52**, 219–228 (2023).

12.   Knief, U. *et al.* Highly pathogenic avian influenza causes mass mortality in

Sandwich Tern *Thalasseus sandvicensis* breeding colonies across north-western

Europe. *Bird Conserv. Int.* **34**, e6 (2024).

13.   European Food Safety Authority *et al.* Avian influenza overview December 2022

– March 2023. *EFSA J.* **21**, (2023).

14.   Elsmo, E. *et al.* Pathology of natural infection with highly pathogenic avian

influenza virus (H5N1) clade 2.3.4.4b in wild terrestrial mammals in the United States

in 2022. Preprint at https://doi.org/10.1101/2023.03.10.532068 (2023).

15.   Nguyen, T.-Q. *et al.* Emergence and interstate spread of highly pathogenic avian

influenza A(H5N1) in dairy cattle. Preprint at

https://doi.org/10.1101/2024.05.01.591751 (2024).

16.   Puryear, W. *et al.* Outbreak of Highly Pathogenic Avian Influenza H5N1 in New England Seals. Preprint at https://doi.org/10.1101/2022.07.29.501155 (2022).

17.   Xie, R. *et al.* The episodic resurgence of highly pathogenic avian influenza H5 virus. *Nature* **622**, 810–817 (2023).

18.   Wille, M. Ecology and Evolution of Avian Influenza A Viruses in Wild Birds. in *Genetics and Evolution of Infectious Diseases* 863–898 (Elsevier, 2024). doi:10.1016/B978-0-443-28818-0.00005-7.

19.   Kaplan, B. S. & Webby, R. J. The avian and mammalian host range of highly pathogenic avian H5N1 influenza. *Virus Res.* **178**, 3–11 (2013).

20.   Hill, N. J. *et al.* Ecological divergence of wild birds drives avian influenza spillover and global spread. *PLOS Pathog.* **18**, e1010062 (2022).

21.   Sonnberg, S., Webby, R. J. & Webster, R. G. Natural history of highly pathogenic avian influenza H5N1. *Virus Res.* **178**, 63–77 (2013).

22.   Verhagen, J. H., Fouchier, R. A. M. & Lewis, N. Highly Pathogenic Avian Influenza Viruses at the Wild–Domestic Bird Interface in Europe: Future Directions for Research and Surveillance. *Viruses* **13**, 212 (2021).

23.   Alkie, T. N. *et al.* A threat from both sides: Multiple introductions of genetically distinct H5 HPAI viruses into Canada via both East Asia-Australasia/Pacific and Atlantic flyways. *Virus Evol.* **8**, veac077 (2022).

24.   *HIGHLY PATHOGENIC AVIAN INFLUENZA RESPONSE PLAN THE RED BOOK*. https://www.aphis.usda.gov/sites/default/files/hpai_response_plan.pdf.

25.   *Policy on the Importation of Terrestrial Foreign or Emerging Animal Disease Agents into Canada by External Facilities*. https://inspection.canada.ca/en/animal-

health/requirements-handling-pathogens/pathogen-imports/policy-

importation?utm_source=chatgpt.com.

26.　Volz, E. M., Koelle, K. & Bedford, T. Viral Phylodynamics. *PLoS Comput. Biol.* **9**,

e1002947 (2013).

27.　Frost, S. D. W. & Volz, E. M. Viral phylodynamics and the search for an 'effective

number of infections'. *Philos. Trans. R. Soc. B Biol. Sci.* **365**, 1879–1890 (2010).

28.　Dudas, G., Carvalho, L. M., Rambaut, A. & Bedford, T. MERS-CoV spillover at

the camel-human interface. *eLife* **7**, e31257 (2018).

29.　De Maio, N., Wu, C.-H., O'Reilly, K. M. & Wilson, D. New Routes to

Phylogeography: A Bayesian Structured Coalescent Approximation. *PLOS Genet.* **11**,

e1005421 (2015).

30.　*Overview of How Canada Prevents, Prepares and Responds to Bird Flu*

Outbreaks. https://inspection.canada.ca/en/animal-health/terrestrial-

animals/diseases/reportable/avian-influenza/prevention-preparedness-and-response.

31.　Bevins, S. N. *et al.* Large-Scale Avian Influenza Surveillance in Wild Birds

throughout the United States. *PLoS ONE* **9**, e104360 (2014).

32.　Shriner, S. A. *et al.* Surveillance for highly pathogenic H5 avian influenza virus in

synanthropic wildlife associated with poultry farms during an acute outbreak. *Sci.*

Rep. **6**, 36237 (2016).

33.　Bokma, B. H., Hall, C., Siegfried, L. M. & Todd Weaver, J. Surveillance for Avian

Influenza in the United States. *Ann. N. Y. Acad. Sci.* **1081**, 163–168 (2006).

34.　*Poultry 2004 Part I: Reference of Health and Management of Backyard / Small*

Production Flocks in the United States, 2004.

https://www.aphis.usda.gov/sites/default/files/poultry04_dr_parti.pdf.

35.   Garber, L., Hill, G., Rodriguez, J., Gregory, G. & Voelker, L. Non-commercial

poultry industries: Surveys of backyard and gamefowl breeder flocks in the United

States. *Prev. Vet. Med.* **80**, 120–128 (2007).

36.   *2022 OIE - Terrestrial Animal Health Code*.

https://www.woah.org/fileadmin/Home/eng/Health_standards/tahc/current/glossaire.p

df (2022).

37.   McKellar, A. E. Phenological Synchrony and Bird Migration: Changing Climate

and Seasonal Resources in North America **Phenological Synchrony and Bird**

Migration: Changing Climate and Seasonal Resources in North America edited

by Eric M. Wood and Jherime L. Kellermann. 2015. CRC Press, Boca Raton, Florida,

USA. xiv + 228 pages, 8 color and 53 black-and-white illustrations. $116.96

(hardcover). ISBN 978-1-4822-4030-6. *The Auk* **133**, 113–114 (2016).

38.   Marra, P. P., Francis, C. M., Mulvihill, R. S. & Moore, F. R. The influence of

climate on the timing and rate of spring bird migration. *Oecologia* **142**, 307–315

(2005).

39.   Kent, C. M. *et al.* Spatiotemporal changes in influenza A virus prevalence among

wild waterfowl inhabiting the continental United States throughout the annual cycle.

Sci. Rep. **12**, 13083 (2022).

40.   Gill, M. S. *et al.* Improving Bayesian Population Dynamics Inference: A

Coalescent-Based Model for Multiple Loci. *Mol. Biol. Evol.* **30**, 713–724 (2013).

41.   Lemey, P., Rambaut, A., Drummond, A. J. & Suchard, M. A. Bayesian

Phylogeography Finds Its Roots. *PLoS Comput. Biol.* **5**, e1000520 (2009).

42.   Ramey, A. M. *et al.* Molecular detection and characterization of highly pathogenic

H5N1 clade 2.3.4.4b avian influenza viruses among hunter-harvested wild birds

provides evidence for three independent introductions into Alaska. *Virology* **589**,

109938 (2024).

43.   Krammer, F., Hermann, E. & Rasmussen, A. L. Highly pathogenic avian

influenza H5N1: history, current situation, and outlook. *J. Virol.* **99**, e02209-24 (2025).

44.   Migratory Bird Program Administrative Flyways | U.S. Fish & Wildlife Service.

https://www.fws.gov/partner/migratory-bird-program-administrative-flyways (2023).

45.   Parker, J., Rambaut, A. & Pybus, O. G. Correlating viral phenotypes with

phylogeny: Accounting for phylogenetic uncertainty. *Infect. Genet. Evol.* **8**, 239–246

(2008).

46.   Kramer, L. D., Ciota, A. T. & Kilpatrick, A. M. Introduction, Spread, and

Establishment of West Nile Virus in the Americas. *J. Med. Entomol.* **56**, 1448–1455

(2019).

47.   Gergely, K.J., Boykin, K.G., McKerrow, A.J., Rubino, M.J., Tarr, N.M., and

Williams, S.G. *Gap Analysis Project (GAP) Terrestrial Vertebrate Species Richness*

Maps for the Conterminous U.S. https://doi.org/10.3133/sir20195034. (2019).

48.   Tobias, J. A. *et al.* AVONET: morphological, ecological and geographical data for

all birds. *Ecol. Lett.* **25**, 581–597 (2022).

49.   Arnal, A. *et al.* Laridae: A neglected reservoir that could play a major role in avian

influenza virus epidemiological dynamics. *Crit. Rev. Microbiol.* **41**, 508–519 (2015).

50.   Lee, D.-H. *et al.* Transmission Dynamics of Highly Pathogenic Avian Influenza

Virus A(H5Nx) Clade 2.3.4.4, North America, 2014–2015. *Emerg. Infect. Dis.* **24**,

1840–1848 (2018).

51.   Munster, V. J. *et al.* Spatial, Temporal, and Species Variation in Prevalence of

Influenza A Viruses in Wild Migratory Birds. *PLoS Pathog.* **3**, e61 (2007).

52.   Kwon, J. *et al.* Domestic ducks play a major role in the maintenance and spread

of H5N8 highly pathogenic avian influenza viruses in South Korea. *Transbound.*

Emerg. Dis. **67**, 844–851 (2020).

53.   Dhingra, M. S. *et al.* Global mapping of highly pathogenic avian influenza H5N1

and H5Nx clade 2.3.4.4 viruses with spatial cross-validation. *eLife* **5**, e19571 (2016).

54.   Hong, S. L., Lemey, P., Suchard, M. A. & Baele, G. Bayesian Phylogeographic

Analysis Incorporating Predictors and Individual Travel Histories in BEAST. *Curr.*

Protoc. **1**, e98 (2021).

55.   Edwards, C. J. *et al.* Ancient Hybridization and an Irish Origin for the Modern

Polar Bear Matriline. *Curr. Biol.* **21**, 1251–1258 (2011).

56.   Nemeth, N. M. *et al.* Bald eagle mortality and nest failure due to clade 2.3.4.4

highly pathogenic H5N1 influenza a virus. *Sci. Rep.* **13**, 191 (2023).

57.   Bahl, J. *et al.* Ecosystem Interactions Underlie the Spread of Avian Influenza A

Viruses with Pandemic Potential. *PLOS Pathog.* **12**, e1005620 (2016).

58.   Shearn-Bochsler, V. I., Knowles, S. & Ip, H. Lethal Infection of Wild Raptors with

Highly Pathogenic Avian Influenza H5N8 and H5N2 Viruses in the USA, 2014–15. *J.*

Wildl. Dis. **55**, 164 (2019).

59.   Redig, P. T. & Goyal, S. M. Serologic Evidence of Exposure of Raptors to

Influenza A Virus. *Avian Dis.* **56**, 411–413 (2012).

60.   2022–2024 Detections of Highly Pathogenic Avian Influenza.

https://www.aphis.usda.gov/livestock-poultry-disease/avian/avian-influenza/hpai-

detections.

61.   Farahat, R. A., Khan, S. H., Rabaan, A. A. & Al-Tawfiq, J. A. The resurgence of

Avian influenza and human infection: A brief outlook. *New Microbes New Infect.* **53**,

101122 (2023).

62.   Patyk, K. A. *et al.* Investigation of risk factors for introduction of highly pathogenic

        avian influenza H5N1 infection among commercial turkey operations in the United

        States, 2022: a case-control study. *Front. Vet. Sci.* **10**, 1229071 (2023).

63.   *APHIS FOREIGN ANIMAL DISEASE FRAMEWORK RESPONSE*

        STRATEGIES. https://www.aphis.usda.gov/sites/default/files/fadprep_manual_2.pdf.

64.   *APPA NATIONAL PET OWNERS SURVEY 2021-2022*.

65.   Smith, G. & Dunipace, S. How backyard poultry flocks influence the effort

        required to curtail avian influenza epidemics in commercial poultry flocks. *Epidemics*

        3, 71–75 (2011).

66.   Minin, V. N. & Suchard, M. A. Counting labeled transitions in continuous-time

        Markov models of evolution. *J. Math. Biol.* **56**, 391–412 (2007).

67.   Minin, V. N. & Suchard, M. A. Fast, accurate and simulation-free stochastic

        mapping. *Philos. Trans. R. Soc. B Biol. Sci.* **363**, 3985–3995 (2008).

68.   *FOIA Report 24-04961-F_000001*. 24-04961-F_000001

        https://www.aphis.usda.gov/sites/default/files/avian-influenza-incidents-and-

        depopulation-methods-feb-2022-to-june-2024.pdf.

69.   *2022–2023 HIGHLY PATHOGENIC AVIAN INFLUENZA OUTBREAK Summary*

        of Depopulation Methods and the Impact on Lateral Spread.

        https://www.aphis.usda.gov/sites/default/files/hpai-2022-2023-summary-depop-

        analysis.pdf.

70.   *Census of Agriculture*.

        https://www.nass.usda.gov/Publications/AgCensus/2022/#full_report (2022).

71.   Hayes, S. *et al.* Ecology and environment predict spatially stratified risk of highly

pathogenic avian influenza in wild birds across Europe. Preprint at

https://doi.org/10.1101/2024.07.17.603912 (2024).

72.   Wight, J. *et al.* Avian influenza virus circulation and immunity in a wild urban duck

population prior to and during a highly pathogenic H5N1 outbreak. *Vet. Res.* **55**, 154

(2024).

73.   Graziosi, G., Lupini, C., Catelli, E. & Carnaccini, S. Highly Pathogenic Avian

Influenza (HPAI) H5 Clade 2.3.4.4b Virus Infection in Birds and Mammals. *Animals*

14, 1372 (2024).

74.   Stallknecht, D. E. *et al.* Highly Pathogenic H5N1 Influenza A Virus (IAV) in Blue-

Winged Teal in the Mississippi Flyway Is Following the Historic Seasonal Pattern of

Low-Pathogenicity IAV in Ducks. *Pathogens* **13**, 1017 (2024).

75.   Bahl, J. *et al.* Influenza A Virus Migration and Persistence in North American Wild

Birds. *PLoS Pathog.* **9**, e1003570 (2013).

76.   Fourment, M., Darling, A. E. & Holmes, E. C. The impact of migratory flyways on

the spread of avian influenza virus in North America. *BMC Evol. Biol.* **17**, 118 (2017).

77.   Prosser, D. J. *et al.* Maintenance and dissemination of avian-origin influenza A

virus within the northern Atlantic Flyway of North America. *PLOS Pathog.* **18**,

e1010605 (2022).

78.   Ahlstrom, C. A. *et al.* Genomic characterization of highly pathogenic H5 avian

influenza viruses from Alaska during 2022 provides evidence for genotype-specific

trends of spatiotemporal and interspecies dissemination. *Emerg. Microbes Infect.* **13**,

2406291 (2024).

79.   Kandeil, A. *et al.* Rapid evolution of A(H5N1) influenza viruses after

intercontinental spread to North America. *Nat. Commun.* **14**, 3082 (2023).

80. Lam, T. T. *et al.* Migratory flyway and geographical distance are barriers to the

gene flow of influenza virus among North American birds. *Ecol. Lett.* **15**, 24–33

(2012).

81. Krauss, S. *et al.* Influenza in Migratory Birds and Evidence of Limited

Intercontinental Virus Exchange. *PLoS Pathog.* **3**, e167 (2007).

82. Girard, Y. A., Runstadler, J. A., Aldehoff, F. & Boyce, W. Genetic structure of

Pacific Flyway avian influenza viruses is shaped by geographic location, host

species, and sampling period. *Virus Genes* **44**, 415–428 (2012).

83. Yang, Q. *et al.* Synchrony of Bird Migration with Global Dispersal of Avian

Influenza Reveals Exposed Bird Orders. *Nat. Commun.* **15**, 1126 (2024).

84. Hicks, J. T. *et al.* Agricultural and geographic factors shaped the North American

2015 highly pathogenic avian influenza H5N2 outbreak. *PLOS Pathog.* **16**, e1007857

(2020).

85. Sullivan, J. D., McDonough, A. M., Lescure, L. M. & Prosser, D. J. Identifying an

Understudied Interface: Preliminary Evaluation of the Use of Retention Ponds on

Commercial Poultry Farms by Wild Waterfowl. *Transbound. Emerg. Dis.* **2024**, 1–9

(2024).

86. Madsen, J. M., Zimmermann, N. G., Timmons, J. & Tablante, N. L. Avian

Influenza Seroprevalence and Biosecurity Risk Factors in Maryland Backyard Poultry:

A Cross-Sectional Study. *PLoS ONE* **8**, e56851 (2013).

87. Bertran, K. *et al.* Pathobiology of Clade 2.3.4.4 H5Nx High-Pathogenicity Avian

Influenza Virus Infections in Minor Gallinaceous Poultry Supports Early Backyard

Flock Introductions in the Western United States in 2014-2015. *J. Virol.* **91**, e00960-

17 (2017).

88.   *Epidemiologic and Other Analyses of HPAI-Affected Poultry Flocks*.

1644   https://www.aphis.usda.gov/sites/default/files/epi-analyses-avian-flu-poultry-2nd-

1645   interim-rpt.pdf.

1646   89.   Shu, Y. & McCauley, J. GISAID: Global initiative on sharing all influenza data –

1647   from vision to reality. *Eurosurveillance* **22**, (2017).

1648   90.   Katoh, K. & Standley, D. M. MAFFT Multiple Sequence Alignment Software

1649   Version 7: Improvements in Performance and Usability. *Mol. Biol. Evol.* **30**, 772–780

1650   (2013).

1651   91.   Kearse, M. *et al.* Geneious Basic: An integrated and extendable desktop

1652   software platform for the organization and analysis of sequence data. *Bioinformatics*

1653   **28**, 1647–1649 (2012).

1654   92.   Minh, B. Q. *et al.* IQ-TREE 2: New Models and Efficient Methods for

1655   Phylogenetic Inference in the Genomic Era. *Mol. Biol. Evol.* **37**, 1530–1534 (2020).

1656   93.   Sagulenko, P., Puller, V. & Neher, R. A. TreeTime: Maximum-likelihood

1657   phylodynamic analysis. *Virus Evol.* **4**, (2018).

1658   94.   Suchard, M. A. *et al.* Bayesian phylogenetic and phylodynamic data integration

1659   using BEAST 1.10. *Virus Evol.* **4**, (2018).

1660   95.   Shapiro, B., Rambaut, A. & Drummond, A. J. Choosing Appropriate Substitution

1661   Models for the Phylogenetic Analysis of Protein-Coding Sequences. *Mol. Biol. Evol.*

1662   **23**, 7–9 (2006).

1663   96.   Drummond, A. J., Ho, S. Y. W., Phillips, M. J. & Rambaut, A. Relaxed

1664   Phylogenetics and Dating with Confidence. *PLoS Biol.* **4**, e88 (2006).

1665   97.   Kass, R. E. & Raftery, A. E. Bayes Factors. *J. Am. Stat. Assoc.* **90**, 773–795

1666   (1995).

1667   98.   Bielejec, F. *et al.* SpreaD3: Interactive Visualization of Spatiotemporal History

1668   and Trait Evolutionary Processes. *Mol. Biol. Evol.* **33**, 2167–2169 (2016).

1669   99.   Bedford, T. *et al.* Global circulation patterns of seasonal influenza viruses vary

1670   with antigenic drift. *Nature* **523**, 217–220 (2015).