Applying deconstructed sourdough communities and fermentation parameters to low-FODMAP wheat bread production

Anna Elizabeth Wittwer[1], Jiaying Amanda Li [1], Kate Howell [1]

[1]School of Agriculture, Food, and Ecosystem Sciences, Faculty of Science, The University of Melbourne, Victoria, Australia

Corresponding author: Kate Howell (khowell@unimelb.edu.au)

Short running head: Low-FODMAP bread-making with sourdough techniques

Abstract

Aims: Despite being made with the same starting materials, standard and sourdough bread differ in terms of levels of fermentable oligosaccharides, disaccharides, monosaccharides, and polyols (FODMAPs). This study aims to assess the impact of a selected yeast-bacteria co-culture isolated from sourdough starters in conjunction with varying bread-making process parameters on bread FODMAP content and overall quality.

Methods and Results: An investigation of the microbial composition of sourdough starters found that many contain *Maudiozyma humilis* and *Fructilactobacillus sanfranciscensis*. A selected pairing was used for small-scale bread-making, and dough fermentation and bread quality parameters were assessed. The sourdough microbes exerted distinctive effects on dough gassing capacity and FODMAP composition in comparison to baker's yeast. To meet low-FODMAP cutoff values, the presence of a fermenting yeast and an extended dough fermentation time (> 6 h) were necessary. Sensory panellists displayed similar levels of preference for bread fermented by baker's yeast and the sourdough co-culture and tended to prefer and detect more complex aromas in co-culture bread fermented for 24 h than for 6 h.

Conclusions: An extended, low-temperature dough fermentation regimen can be used with a co-culture of common sourdough fermentation microbes to produce low-FODMAP, well-liked bread with standard bread-making materials.

Keywords: FODMAPs, bread, wheat, sourdough, *Maudiozyma humilis*, *Fructilactobacillus sanfranciscensis*

Introduction

Wheat bread is an important and widely consumed staple food of the Western diet. It is a good source of energy, dietary fibre, and micronutrients (Ispiryan et al., 2021), while also being convenient and well-liked. However, certain wheat-based products are unsuitable for consumption by people with functional gastrointestinal disorders such as irritable bowel syndrome (IBS) because they are often rich in fermentable oligosaccharides, disaccharides, monosaccharides and polyols (FODMAPs).

Common examples of FODMAPs include fructans, α-galactooligosaccharides, lactose, fructose in excess of glucose, polyols, and they are found in a broad range of food groups in addition to cereal-based products (Menezes et al., 2018; Varney et al., 2017). These carbohydrates are not properly absorbed in the small intestine, leading to the production of gas and short-chain fatty acids in the large intestine by the gut microbiome. For this reason, many FODMAPs are also considered to be dietary fibres (Yan et al., 2018), but in IBS sufferers these properties lead to adverse gastrointestinal symptoms including abdominal pain, diarrhoea, flatulence, and bloating. Furthermore, certain FODMAPs attract water via osmotic activity, which can lead to luminal distention (Gibson & Shepherd, 2010).

The major FODMAP present in wheat bread is fructans (**Figure 1**), which are β (2→1) and/or β (2→6) fructo-oligosaccharides (FOS) containing a single glucose moiety. Fructan concentrations range from 0.6 to 2.9% in wholewheat flour (Huynh et al., 2008), and 1.4 to 1.7% in white wheat flour (Atzler et al., 2020). The average degree of polymerisation (DP_{av}) is 4, with a maximum of 14-19 (Atzler et al., 2020). Fructans derived from grain products are branched and categorised as graminan-type (Menezes et al., 2018; Pejcz et al., 2020). Other FODMAPs present in wheat bread in smaller quantities can include raffinose, tri- and tetra-fructo-oligosaccharides (e.g., nystose, kestose), excess fructose, and mannitol (Biesiekierski et al., 2011).

Figure 1 Key FODMAPs in standard and sourdough wheat bread fermented for 60 or 80 minutes. Bars represent mean values. Data from Schmidt and Sciurba (2021).

A diet low in FODMAPs is a proven therapeutic measure to reduce or eliminate IBS symptoms in those with functional gastro-intestinal disorders (Halmos et al., 2014), but the exclusion of wheat bread from the diet is not ideal due to its staple food status and the non-FODMAP fibres and micronutrients it provides. Bread can be considered low-FODMAP, however, if in a serving of bread (50 g) the following cutoff values are not exceeded: 0.3 g of oligosaccharides, 0.15 g of excess fructose, 0.4 g of polyols, and a total of 0.5 g of FODMAPs excluding lactose (Varney et al., 2017). Current low-FODMAP breadmaking focuses on either enzymatic pre-treatment of flour to remove FODMAPs, or the use of grains that are naturally low in FODMAPs, such as spelt (Biesiekierski et al., 2011). While both options are viable, they are relatively expensive and are not as broadly accessible as conventional white wheat bread.

Conventional white wheat bread is typically leavened with baker's yeast (industrial strains of *Saccharomyces cerevisiae*) for a short period of time at approx. 30°C (Shewry et al., 2022). Despite the fact that baker's yeast produces invertase, an enzyme that can effect fructan hydrolysis (Laurent et al., 2020), FODMAP levels in conventional wheat bread often exceed low-FODMAP thresholds, mainly because of high fructan content (**Figure 1**).

Sourdough bread, on the other hand, while made with the same ingredients as standard bread, has a distinct and often improved FODMAP composition. Namely, its fructan content is lower, whereas levels of excess fructose and mannitol may be enriched (Schmidt & Sciurba, 2021). Sourdough bread is traditionally leavened with a type I sourdough starter. This is a spontaneously-fermented mixture of

flour and water that via a process of regular re-feeding with fresh ingredients (termed 'backslopping')

develops a community of often non-conventional yeasts and bacteria (Gobbetti, 1998; Rogalski et al.,

2021). A mature sourdough typically contains one or two yeast species and one-several bacterial

species (Van Kerrebroeck et al., 2017), which may interact with one another by various means. One of

the most common sourdough yeasts is *Maudiozyma humilis* (formerly *Kazachstania* or *Candida*

humilis), an ascomycetous yeast that occupies a phylogenetically close position relative to the

Saccharomyces genus (Arora et al., 2021; Liu et al., 2024; Smukowski Heil et al., 2022). This yeast is

commonly isolated from sourdough starters that also contain *Fructilactobacillus sanfranciscensis*, an

obligately heterofermentative lactic acid bacterium (LAB) whose primary ecological niche is

sourdough starters (Altilia et al., 2021; Rogalski et al., 2020). In addition to its varied microbial

profile compared to conventional bread, sourdough bread is often made according to different process

parameters. Depending on the style of sourdough production, dough proofing or fermentation can be

carried out over approx. 24 h and can include several propagation steps (Brandt, 2007).

It is already well-understood that extending dough proofing times leads to increased degradation of

fructans in bread dough (Schmidt & Sciurba, 2021; Ziegler et al., 2016). However, the effects of

proofing temperature on this process and its interaction with proofing time are less well-characterised.

Furthermore, while some yeasts, such as *Saccharomyces cerevisiae*, *Kluyveromyces marxianus*, and

Lachancea fermentati have been investigated for their applicability to low-FODMAP breadmaking

(Ispiryan et al., 2021; Struyf et al., 2018), there remains a lack of information about the potential of

common sourdough microbes in this area, given the low-fructan status of sourdough bread.

Therefore, the aim of this study was to investigate the individual and combined contributions of two

key aspects of sourdough breadmaking to FODMAP content and bread technical quality: functional

metabolic properties of a common sourdough yeast-bacteria pairing, and the dough proofing time-

temperature regime typical of sourdough breadmaking.

Materials and Methods

Growth media preparation

For yeast strain isolation, two media were prepared. Wallerstein Nutrient (WLN) agar was prepared according to the manufacturer's instructions (Oxoid). Yeast-peptone-dextrose (YPD) agar was composed of 10 g/L yeast extract (Oxoid), 20 g/L bacteriological peptone (Oxoid), 20 g/L anhydrous D-glucose (Merck), and 20 g/L w/v bacteriological grade agar (Ajax). When used for strain isolation and viable colony enumeration, yeast growth media were supplemented with 100 μg/mL ampicillin and 30 μg/mL chloramphenicol to prevent bacterial growth.

For bacterial strain isolation, de Man, Rogosa & Sharpe (MRS) agar was prepared according to the manufacturer's specifications (Oxoid). Isolation and enumeration media were supplemented with 10 μg/mL cycloheximide to prevent fungal growth.

Antibiotic solutions were added aseptically to just-autoclaved media at approx. 40-50 °C.

Strain isolation and identification

Nine sourdough starters were donated to this project. Codes, suburbs and isolates from each bakery are given in **Table 1**.

Table 1 *Maudiozyma humilis* and *Fructilactobacillus sanfranciscensis* isolated from sourdough starters. Cells shaded grey indicate starters that contain both species.

Leaven	Origin	Code	Species	Top BLAST identity (%)
Os	Ovens St Bakery, Brunswick VIC	OsY1	*Maudiozyma humilis*	98.9
Osr	Ovens St Bakery, Brunswick VIC	OsrY6	*Maudiozyma humilis*	99.3
Os	Ovens St Bakery, Brunswick VIC	OsY5	*Maudiozyma humilis*	99.4
Osr	Ovens St Bakery, Brunswick VIC	OsrY1	*Maudiozyma humilis*	99.5
Nrk	Natimuk, VIC	NrkY1	*Maudiozyma humilis*	99.7
Nrk	Natimuk, VIC	NrkB1	*Fructilactobacillus sanfranciscensis*	100.0
Os	Ovens St Bakery, Brunswick VIC	OsB1	*Fructilactobacillus sanfranciscensis*	100.0
Os	Ovens St Bakery, Brunswick VIC	OsB2	*Fructilactobacillus sanfranciscensis*	100.0
Osr	Ovens St Bakery, Brunswick VIC	OsrB1	*Fructilactobacillus sanfranciscensis*	100.0
Lbr	Loafer Bread Bakery, Fitzroy North VIC	LbrY1	*Maudiozyma humilis*	98.0
Fb	Falco Bakery, Collingwood VIC	FbY1	*Maudiozyma humilis*	99.7
Lb	Loafer Bread Bakery, Fitzroy North VIC	LbY2	*Maudiozyma humilis*	99.8
Gw	Great Western Granary, Great Western VIC	GwB1	*Fructilactobacillus sanfranciscensis*	100.0
Kb	Ket Baker, Wallington VIC	KbB1	*Fructilactobacillus sanfranciscensis*	100.0
Zb	Zelda Bakery, Ripponlea VIC	ZbY4	*Maudiozyma humilis*	100.0

During sampling, care was taken to avoid contamination such that only microbes present in the sourdough starter itself or the 'house microbiota' (organisms associated with each isolate's respective bakers and bakery environment) of the bakery would be collected (Minervini et al., 2015). Bakers were asked to collect a portion of their sourdough starter(s) in sterile tubes, or to provide a sample in their own container. Samples were stored at room temperature prior to analysis and cultured 24-72 h after the last starter refreshment.

Sourdough starters were first serially diluted by adding approx. 1 g starter to 9 mL 0.1% w/v peptone water. Suspensions were vortexed thoroughly between each dilution step. Serial dilutions were carried out until a factor of 10^{-6} was reached. Aliquots of 100 μL of each 10^{-3}-10^{-6} suspension were spread-plated on antibiotic-supplemented WLN and YPD agar to isolate yeasts, and antibiotic-supplemented MRS agar to isolate LAB.

127 Plates were incubated in aerobic conditions at 28 °C for 2-3 days. Plates with 30-300 countable

128 colonies present were selected for sub-culturing. Well-isolated colonies were selected on the basis of

129 distinct morphological characteristics. Three iterations of streak plating were performed to isolate

130 pure colonies. After the third iteration, each isolate was given an identification code to reflect its

131 origin, and glycerol stocks were stored at -80 °C.

Yeast isolate identification

Genomic DNA was extracted from pure yeast colonies with the MasterPure™ Yeast DNA Purification

Kit (Epicentre®, Madison, WI) according to the manufacturer's instructions. PCR amplifications of

the extracted DNA were carried out using the KAPA HiFi HotStart PCR Kit (Kapa Biosystems,

Wilmington, MA). The reaction used the primers NL-1 and NL-4 to target the D1/D2 region of the

large subunit ribosomal DNA as described in Kurtzman & Robnett (1998). Amplicons were submitted

to the Australian Genome Research Facility (AGRF) for sequencing. Raw sequences were trimmed,

forward and reverse sequences aligned, and the alignments searched against the NCBI nucleotide

database in Geneious Prime 2024.0. In the event of an ambiguous or improbable top result, YeastIP

(Weiss et al., 2013), a curated sequence database, was used to identify isolates.

Bacterial isolate identification

143 Pure bacterial colonies were prepared for identification in Applied Biosystems™ PrepMan® Ultra

144 Sample Preparation Reagent (Thermo Fisher Scientific). Attenuated samples were submitted to the

145 AGRF for taxonomic assignations using the 16s rRNA gene.

Strain selection for dough and bread analyses

147 One sourdough-derived *Maudiozyma humilis* strain was selected for small-scale breadmaking trials

148 based on the following criteria: a) originated from a sourdough starter in which a *Fructilactobacillus*

149 *sanfranciscensis* isolate was also present and b) demonstrated acceptable levels of growth on standard

150 growth carbohydrates and fructan-like carbohydrates.

A defined medium (Yeast Nitrogen Base with amino acids (YNB) Sigma) supplemented individually with 0.5% w/v glucose, fructose, small FOS (DP 2-8), or large FOS (DP > 60) was prepared to test yeast isolate growth on different carbon sources. Small FOS and large FOS (inulin) were purchased from Megazyme (Bray, Ireland)

Yeasts were precultured twice in aliquots of 5 mL of a starvation medium based on the composition of YPD that contained 5% of the standard concentration of glucose (1% w/v yeast extract, 2% w/v peptone, 0.1% w/v glucose). Cell suspension turbidity was standardised to an optical density measurement at 600nm (OD_{600}) of 0.1 in 0.1% peptone. To each well of a U-bottomed 96-well microtitre plate, 198 μL supplemented YNB broth was added. Aliquots of 2 μL of the yeast cell suspensions were added, and 2 μL 0.1% peptone was used for the blanks. Plates were incubated for 24 h at 28 °C, and cell density was measured with a plate reader (Multiskan Go®, ThermoFisher) every hour at 600 nm. Kinetic parameters used to assess yeast growth performance were obtained with the *growthcurver* package (Sprouffske & Wagner, 2016).

Dough and bread analyses

Inoculum preparation

Yeasts and bacteria were maintained on YPD and MRS agar, respectively. A fresh plate was prepared 2-5 days prior to breadmaking to ensure maximum live cells. *M. humilis* OsrY1 and the baker's yeast were precultured in 30mL YPD broth at 28 °C at 220 rpm for 18 h. *F. sanfranciscensis* OsrB1 was precultured in 30mL MRS broth at 29 °C at 18% CO_2 for 72 h. Prior to baking, precultures were centrifuged at 4000 rpm/1789 rcf for 5 min and the growth medium was removed.

To inoculate dough with baker's yeast in the same manner as the sourdough-derived microbes, a commercially available baker's yeast was purchased and cultured in YPD broth. The baker's yeast was maintained on YPD and stored in 80% v/v glycerol.

Dough preparation

Bread dough was prepared in small domestic breadmaking machines (BM2500 Compact Bakehouse, Sunbeam, NSW, Australia). Each batch of dough was composed of 500 g flour, 7.5 g table salt, 5 g canola oil, and 315 g water. Pre-cultured cells were added as shown in **Table 2** with a sterile transfer pipette. White wheat all-purpose baker's flour containing 11.5% protein and 3.0% dietary fibre was used (Wallaby T55 variety, Laucke, South Australia). Doughs were mixed for 5 min on setting 9, then the machines were switched off.

Table 2 Dough fermentation microbe treatments applied to samples.

Treatment, code	Microbe(s)	Inoculum mass (g)
Unstarted dough, None	None	0
Baker's yeast, BY	*Saccharomyces cerevisiae* BY	0.6 ± 0.05
Sourdough yeast, OsrY1	*Maudiozyma humilis* OsrY1	0.6 ± 0.05
Sourdough bacterium, OsrB1	*Fructilactobacillus sanfranciscensis* OsrB1	0.6 ± 0.05
Sourdough co-culture, OsrY1 + OsrB1	*M. humilis* OsrY1 + *F. sanfranciscensis* OsrB1	1.2 ± 0.1

Dough characterisation

Enumeration of yeast and LAB

Quantities of 0.5 ± 0.05 g bread dough were added incubated at 25 or 32°C. At 0, 2, 6, and 24 h proofing, aliquots of 4.5 mL 0.1% w/v peptone water were added. Tubes were vigorously shaken and vortexed to create a 10^{-1} dilution, then the dough slurry was serially diluted to 10^{-5}. Aliquots of 75 μL of the 10^{-3}-10^{-5} dilutions were spread plated on antibiotic-supplemented MRS and YPD agar. The MRS plates were incubated at 27°C under 18% CO_2 and the YPD plates were incubated under aerobic conditions at 28°C for 4 days. Plates containing colonies within the countable range (30-300) were selected for enumeration.

Dough pH and temperature

A solid-state food and dairy pH meter HI199161 (Hanna Instruments) was used to measure dough pH and temperature at each proofing time point. Three separate measures were obtained at each time point by placing the pH probe in three different locations within the dough sample.

Dough gassing capacity

Immediately after mixing, portions of 100 g dough were added to 500 mL Schott bottles and sealed

with gas pressure modules (ANKOM Technology, Macedon, NY). Bottles were kept in incubators set

to 25 or 32°C, and pressure measurements were taken every 15 s over 24 h.

Dough FODMAP quantification

Quantities of approx. 5 g dough were sampled at 0, 6, and 24 h of fermentation. The 2 h timepoint

was omitted from analysis due to unacceptable bread technical qualities (i.e. insufficient gas

production for leavening), but the 0 h was analysed as a baseline. Dough samples were freeze-dried

for 72 h (Biobase freeze dryer BK-FD10P). Dough was then crushed to a powder with a metal spatula,

and FODMAPs were extracted using a hot water method. For fructan analyses, 0.1 g freeze-dried,

powdered dough was added to 5 mL deionised water and incubated in boiling water for 15 min. For

the other FODMAP and dough enzyme analyses, 0.1 g dough was added to 1 mL deionised water and

incubated for 10 minutes at 80°C in a heating block. Fructan, mannitol, and excess fructose were

quantified with UV-based methods using kits from K-FRUC, K-MANOL, and K-FRUGL kits

respectively (Megazyme).

Bread preparation and characterisation

Quantities of 325 g dough were portioned into lightly oiled containers, covered with cling film and

left to prove for 2, 6, or 24 hours in incubators set to 25 or 32°C. One hour before the end of the set

proofing time, doughs were knocked back and shaped into small loaf tins (14.6 x 7.62 cm, Wilton).

For all doughs, the final hour of proving was carried out at 25°C to ensure consistent dough

temperature prior to baking. Loaves were baked in a Convotherm 4 easyDial 6.10 ES oven (Elfging,

Germany) for 30 min at 180°C (fan 4, Crisp & tasty 3, Bake pro 3). After baking, they were cooled,

and a serrated knife was used to make slices of approx. 2 cm thick. The maximum height of the

middle three slices from each loaf was measured.

Two slices from each loaf were photographed with an iPhone 13 using default camera settings in a

light box (Photosimile 50; Ortery, Irvine, CA) for digital image analysis. Photographs were cropped to

show only the largest area of regular crumb patterns. They were then analysed with MATLAB® ver.

R2023b (Mathworks Inc., Matick, MA) to output various parameters, including void fraction, mean

cell area (mm^2), small cell density (per cm^2), and large cell density (per cm^2).

Bread sensory analysis

225 An untrained sensory consumer panel was used to test palatability and acceptability of low-FODMAP

226 wheat breads. Four samples were selected based on a) low-FODMAP status and b) acceptable bread

227 technical quality.

228 **Table 3** Low-FODMAP bread samples chosen for sensory analysis.

Fermentation microbe(s)	Fermentation time (h)	Fermentation temp. (°C)
Baker's yeast (*S. cerevisiae*)	24	25
M. humilis OsrY1	24	25
M. humilis OsrY1 + *F. sanfranciscensis* OsrB1	24	25
M. humilis OsrY1 + *F. sanfranciscensis* OsrB1	6	25

Participants (*n* = 56) aged 18-50 attended four sensory panel sessions held over a 5-day period. One

week prior to testing, bread loaves were baked, cut into 1-cm-thick slices, and frozen. Samples were

defrosted the night before each session. Participants were asked to examine and taste each bread

sample one at a time, labelled with a random 3-digit code. The order in which the four samples were

given was randomised for each participant. They were asked to score each sample based on how it

looked, smelled, and tasted using a 7-point semi-structured scale with described extremes. Approval

for this experiment was given by the University of Melbourne Human Research Ethics Committee

(HREC 2024-30068-56367-3).

Statistical analyses

All experiments were carried out in triplicate, and statistical analyses and data visualisations were

completed in R (R Core Team, 2024). Group means for all data except the bread sensory data were

compared with the Kruskal-Wallis nonparametric test, and *post-hoc* pairwise comparisons were made

with the Dunn's test with Bonferroni's correction applied. Sensory data scores were compared with

one-way ANOVA, and pairwise comparisons conducted with Tukey's HSD test. Statistical

significance was considered at $p < 0.05$.

Results

Yeasts and bacteria from sourdough starters

Based on LSU rDNA gene sequencing results, nine *Maudiozyma humilis* strains were isolated from

the sourdough starters studied (**Table 1**). Pairwise identity for yeasts ranged from 98 to 99.8% and for

bacteria was 100%. Bacteria were identified using the 16s rRNA gene, and the main species identified

was *Fructilactobacillus sanfranciscensis*.

Since *M. humilis* and *F. sanfranciscensis* are a frequently co-occurring pair in sourdough starters

(Rogalski et al., 2021), these species were selected for low-FODMAP bread-making trials. In order to

maximise potential FODMAP degradation, we focused on assessing the functional properties of the

M. humilis strains, as *F. sanfranciscensis* is not among the group of LAB that produce FODMAP-

degrading enzymes such as fructanase (Loponen & Gänzle, 2018). To select the *M. humilis* strain

most capable of bread FODMAP reduction and best suited to bread-making, growth was measured in

defined media individually supplemented with carbohydrates. Strains with growth kinetic parameters

are listed in **Table 4**.

Table 4 Growth kinetic parameters of *Maudiozyma humilis* strains. Growth performance of strains
was assessed in yeast nitrogen base (YNB), a defined medium individually supplemented with fructo-
oligosaccharides (FOS) with a degree of polymerisation (DP) of 2-8, FOS DP > 60, or glucose. Values
shown are mean ± SD.

	Area under the curve (AUC)			Time to mid-exponential phase (h)		
Strain	YNB + FOS DP < 8	YNB + FOS DP > 60	YNB + glucose	YNB + FOS DP < 8	YNB + FOS DP > 60	YNB + glucose
M. humilis NrkY1	11.44 ± 0.69	9.18 ± 0.86	13.98 ± 0.42	11.56 ± 0.52	10.38 ± 0.75	7.48 ± 0.35
M. humilis OsY1	8.65 ± 2.92	6.64 ± 2.59	18.43 ± 0.68	11.05 ± 0.76	10.92 ± 1.17	7.89 ± 0.08
M. humilis OsY5	9.25 ± 4.82	6.35 ± 2.54	15.9 ± 1.51	10.72 ± 0.73	9.32 ± 1.54	8.11 ± 0.32

M. humilis OsrY1	10.77 ± 5.01	9.02 ± 0.32	18.44 ± 0.58	10.67 ± 1.16	8.93 ± 0.37	8.26 ± 0.16
M. humilis OsrY6	13.49 ± 1.16	6.43 ± 2.6	16.46 ± 1.56	11.18 ± 0.3	11.37 ± 0.66	8.13 ± 0.55
S. cerevisiae BY	9.38 ± 3.14	6.12 ± 3.06	17.72 ± 0.9	9.64 ± 2	12.18 ± 1.53	7.91 ± 0.08

262 *M. humilis* OsrY1 was selected for small-scale breadmaking trials because its growth intensity and

263 speed in defined media were either equivalent or superior to those of the other strains, particularly on

264 media supplemented with large FOS (DP > 60, **Table 4**). *F. sanfranciscensis* OsrB1 isolated from the

265 same starter (**Table 1**) was selected for co-culture and bacterial mono-culture tests.

Microbial growth kinetics

Bread dough was prepared with the following five microbe conditions: baker's yeast, *M. humilis*

OsrY1, *F. sanfranciscensis* OsrB1, *M. humilis* OsrY1 + *F. sanfranciscensis* OsrB1, and no inoculum

(unstarted dough). Doughs were fermented in proofing ovens set to either 25 or 32°C.

Internal dough temperature stabilised by 2 h fermentation and did not change thereafter. The mean

internal temperature of doughs fermented at 32°C and 25°C, regardless of microbial treatment, was

29°C and 24°C, respectively (**Table S2**).

Growth and persistence of yeasts and LAB in the dough substrate was investigated through plate

count-based enumeration (**Figure 2**). For all four inoculated treatments, colony-forming unit (CFU)

counts increased significantly over the course of the total 24 h fermentation time. This increase was

only statistically significant for the co-culture dough when fermented at 32°C, however. In the three

mono-culture doughs inoculated with only one strain, no contaminating yeast or bacteria were

detected. By 24 h, the baker's yeast reference dough did not have significantly higher yeast counts

than the *M. humilis* OsrY1 mono- and co-culture doughs.

Figure 2 Enumeration of viable yeast and LAB in dough fermented at 25 and 32°C over time. Dough
samples were serially diluted in 0.1% w/v peptone, and yeast and LAB were enumerated on
antibiotic-supplemented YPD and MRS agar plates respectively. Log_{10}-transformed data presented as
mean ($n = 3$) colony-forming units (CFU) ± SE, * represents $p < 0.05$.

The highest CFU/g dough values were found in the dough inoculated solely with *F. sanfranciscensis*

OsrB1. Proofing temperature was found to have a significant effect on growth. At 6 h, LAB counts

were significantly higher in the co-culture dough fermented at 32°C than those in the 25°C co-culture

dough. No significant differences between LAB counts in the *F. sanfranciscensis* mono-culture dough

proofed at 25 and 32°C were detected. Conversely, for both treatments containing *M. humilis* OsrY1,

yeast counts at 24 h of fermentation were higher at 25°C than at 32°C. In the unstarted dough

treatment, to which no microbes were added during the dough mixing stage, counts of LAB and yeast

increased markedly by 24 h, and this increase was more pronounced at 32°C than at 25°C.

Dough pH was measured with a solid-state meter throughout the fermentation process. Over the 24-

hour measurement period, the pH of all doughs, regardless of inoculum or proofing temperature,

decreased (**Table S1**). pH of the *M. humilis* mono- and co-culture doughs decreased significantly from

2 to 24 h at both temperatures. In all dough fermentation treatments, the 24-hour samples had lower

pH values when fermented at 32°C than when fermented at 25°C.

Fermentation dynamics

An ANKOM gas production system was used to measure each treatment's dough leavening capability in small quantities over 24 h. Area under the curve (AUC) of each treatment-temperature combination was calculated to quantify overall gas production.

Figure 3 Leavening ability of *M. humilis* OsrY1 in mono- and co-culture is comparable to that of baker's yeast. **A** Lines indicate cumulative dough pressure output. **B** Boxplots show each treatment-temperature group AUC value, grey region indicates low gassing capacity. **C** Bars indicate dough cumulative pressure at 2, 6, or 24 h. Mean ($n = 3$) values shown, error bars and ribbons indicate SE. * $p < 0.05$, ** $p < 0.01$.

Dough cumulative pressure output was affected by microbial treatment, proofing temperature, and

individual treatment-temperature combinations. The treatments can be grouped into high- and low-

gassing capacity status according to whether they contain a yeast or not; the latter group's mean AUC

is < 50 (**Figure 3B**). Among these groups, the treatments that demonstrated the highest and lowest

gassing capacity were baker's yeast fermented at 32°C and unstarted dough at 25°C. Overall gas

production at either fermentation temperature did not significantly differ between any of the yeasted

treatments. At 2, 6, and 24 h, dough leavening ability was significantly improved by increased

fermentation temperature (**Figure 3C**).

FODMAP content

317 Changes in concentrations of key dough FODMAPs (fructans, mannitol, excess fructose) over time

318 under different fermentation conditions were measured.

Figure 4 The presence of a fermenting yeast and > 6 h dough fermentation are required to reduce fructan levels to below the low-FODMAP threshold. Shaded light blue region indicates low-FODMAP threshold for oligosaccharides in cereal products as described by Varney et al. (2017). Points indicate mean ($n = 3$) total FODMAP content per 50 g serving ± SD. * $p < 0.05$.

Fructan was the main FODMAP detected (89.6% of total measured FODMAP content), with mannitol

and excess fructose being found in minor proportions (5.3% and 5.1%, respectively). Over 24 h dough

fermentation, *M. humilis* OsrY1 was capable of fructan degradation (and lowering of total FODMAP

content) to the same extent as baker's yeast (**Figure 4**). By 24 h, all three yeasted treatments lowered

fructan levels to well below the 0.3 g per serve threshold required to classify a food as low in

oligosaccharides.

The increase in mannitol concentration in the dough over time was only significant for dough

fermented solely by *F. sanfranciscensis* OsrB1. By the 24 h timepoint, mannitol consisted of approx.

27% total FODMAP content in *F. sanfranciscensis* OsrB1 dough fermented at both 25 and 32°C.

Furthermore, mannitol levels were negligible in all other treatments, including the dough co-cultured

with *M. humilis* OsrY1.

In yeasted doughs fermented at 32°C, excess fructose tended to accumulate around the 6 h timepoint

and deplete as fermentation continued. For *M. humilis* OsrY1, this depletion was statistically

significant ($p = 0.04291$). In general, dough monosaccharides followed an 'accumulate-deplete'

pattern in yeasted doughs (**Figure S2**). Available fructose and glucose levels were not linked to dough

α-amylase content, which did not change in any inoculated doughs (**Figure S1**), so either microbial

enzymes or other forms of endogenous amylases were likely to have been responsible.

At each time point, fermentation temperature demonstrated no significant effect on total FODMAP

content. However, for specific treatment groups, effects related to individual FODMAPs were

observed. At both 6 and 24 h fermentation, total FODMAP levels in *M. humilis* OsrY1 mono-culture

doughs fermented at 32°C were lower than those at 25°C ($p = 0.04953$). *M. humilis* OsrY1 dough

proofed for 6 h at 32°C was the only treatment with fructan levels below the 0.3 g per serve threshold

at this time point. Fructan degradation by baker's yeast was slightly improved at 32°C compared to

25°C at the 24 h time point ($p = 0.04953$), and doughs containing *F. sanfranciscensis* OsrB1 showed

significantly higher mannitol levels at 6 h when fermented at 32°C. In general, temperature-based

comparisons were only marginally significant at the α 0.05 threshold. There were also no

significant differences in excess fructose content according to proofing temperature.

Crumb structure quality parameters

352 To ensure that any low-FODMAP bread produced was sufficiently leavened and had an optimal

353 crumb structure, various loaf quality parameters were measured.

Figure 5 During extended dough proofing, higher temperatures lead to over-proved bread with a more uneven crumb structure. **A** Loaf height, **B** percentage of loaf cross section composed of 'void' or gas pockets (well-leavened bread exceeds 25% void fraction, indicated by the grey shaded region), **C** average gas pocket size, and **D** the ratio of large to small gas pockets. Bars indicate mean ($n = 3$) ± SE. * $p < 0.05$.

An interplay between fermentation microbe, temperature, and extent was observed in the degree of

leavening of the different breads. In the early stages of fermentation (2 h), only loaf heights of yeasted

treatments proofed at 32°C approached acceptable values. The baker's yeast loaves were significantly

taller at 2 and 6 h fermentation at 32 than at 25°C. For doughs baked at later fermentation stages, the

distinction between yeasted and non-yeasted doughs was more evident at 25 than 32°C (**Figure 5A**).

The yeasted doughs fermented at 32°C underwent over-proving (collapse of dough gluten structure

resulting in reduced loaf height). This effect was most pronounced for the baker's yeast bread, whose

height decreased by approx. 10 mm by 24 h compared to 6 h. Bread is considered appropriately

leavened when composed of approximately 25% 'void' or gas pockets (**Figure 5B**). The void fraction

of the *M. humilis* OsrY1 mono- and co-culture loaves was not significantly different from the baker's

yeast loaves under any conditions, nor did the mono- and co-culture loaf void fractions differ from

each other.

The size and variation in size of the gas pockets is also an important crumb structure parameter. The

average gas pocket size increases with fermentation duration (**Figure 5C**). A less uniform crumb

structure was more common in bread made from dough proofed at 25°C, and large gas pockets only

appeared in the 32°C breads once over-proved (**Figure 5D**).

Bread sensory analysis

Finally, to confirm that low-FODMAP bread produced with sourdough co-cultures would be

palatable, we conducted a sensory panel. Un-trained participants were asked to rate various visual,

taste, and aroma aspects of each of the four samples on 7-point scales with defined extremes. **Figure 6**

shows the distribution and mean scores given to each sample in response to the survey.

Figure 6 Extending proofing time of sourdough co-culture breads improves overall liking and distinctive sourdough flavours and aromas. Un-trained study participants ($n = 56$) scored samples on each attribute on a 7-point scale with defined extremes. $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, **** $p < 0.0001$.

Sensory participants gave the highest overall liking scores to baker's yeast bread fermented for 24 h at 25°C, and this was also reflected in their scores for overall flavour, aroma, and texture (**Figure 6**). The

baker's yeast sample only scored significantly higher for overall liking, flavour, and visual than the 6 h

bread co-fermented with *M. humilis* and *F. sanfranciscensis*, not the two 24 h breads fermented with

M. humilis.

Between the two breads fermented with sourdough co-culture, panellists tended to prefer the 24 h

bread, which received higher scores for sour/vinegar and fruity aromas and flavours. The aroma

complexity of the 24 h co-culture bread was also rated significantly higher than the 6 h bread. Both

sour/vinegar attributes were scored higher for the 24 h co-culture bread than the *M. humilis* mono-

culture bread although they did not differ in terms of dough pH (**Table S1**). Fruity aromas and

flavours were also scored as significantly stronger in the co-culture 24 h bread than the 6 h bread.

Of the three 24 h breads, the two containing *M. humilis* were rated as being significantly darker in

colour than the baker's yeast bread, and overall visual scores were lower for these two samples than

the baker's yeast and 6 h co-culture breads.

Discussion

This study isolated and identified yeasts and bacteria in sourdough starters and optimised the use of a

selected sourdough-derived co-culture in the production of low-FODMAP white wheat bread.

Variation in bread dough fermentation process parameters facilitates leavening and FODMAP

degradation to differing extents depending on the fermentation microbe(s) used.

Globally, 80 species of yeast and 82 species of lactobacilli have been identified in multiple instances

in sourdough starters (Arora et al., 2021). In accordance with the findings of our survey of sourdough

starters, the two most common yeast species found in sourdough starters are *Saccharomyces*

cerevisiae and *Maudiozyma humilis* (Bigey et al., 2021; Wittwer et al., 2022). *M. humilis* and *F.*

sanfranciscensis have previously been found to frequently co-occur in sourdough starters and

contribute interdependently to the bread fermentation process (Carbonetto et al., 2018). This pairing

occurred in three of the nine sourdough starters sampled in the present study.

Our quantification of fructans in white wheat dough aligns with previously published values, and

confirms that they are the principal FODMAP found in bread dough (Biesiekierski et al., 2011; Huynh

et al., 2008).

The inulinase activity of *Kluyveromyces marxianus* CBS6014 allows this yeast to effect rapid and

near-complete degradation of fructans in whole wheat flour, whose fructan content is generally higher

than that in white wheat flour (Biesiekierski et al., 2011; Struyf et al., 2018). *M. humilis*, which does

not produce inulinase or any other enzyme specific to dough oligosaccharides relies instead on

invertase. Both our and other published studies of changes in fructan levels in dough fermented by

invertase-producing yeasts like *M. humilis* and *S. cerevisiae* suggest that effectiveness of invertase-

mediated fructan degradation is heavily dependent on fermentation duration (Laurent et al., 2020;

Schmidt & Sciurba, 2021; Ziegler et al., 2016). Mannitol accumulated in the dough inoculated solely

with *Fructilactobacillus sanfranciscensis*. This was expected, as this bacterium is known to use

fructose in bread dough as an external electron acceptor to regenerate NADP, thereby producing

mannitol (Korakli & Vogel, 2003). However, it did not accumulate in the co-culture dough treatment.

It is unlikely that *M. humilis* consumed the mannitol produced, as it is described as negative for

growth or assimilation on mannitol (Lachance et al., 2011). The more likely explanation is that *M.*

humilis consumed the available fructose before *F. sanfranciscensis* could convert it to mannitol.

Dough fermentation co-cultures must perform well from a technical standpoint as well as beneficially

modulating bread FODMAP composition. Differences in fermentative performance according to

dough fermentation temperature suggested that baker's yeast is reliant on higher temperatures for

optimal leavening and FODMAP reduction. This is likely to be a property of *S. cerevisiae* having a

larger growth temperature range (Vaughan-Martini & Martini, 2011). This difference was less

pronounced for *M. humilis*, whose type strain does not grow at temperatures over 30°C (Lachance et

al., 2011). Furthermore, we found that dough over-proving and the subsequent loss of loaf height were

worse at the higher dough fermentation temperature tested. Together, these results suggest that *M.*

humilis is better suited to an extended, cooler dough fermentation regime targeted towards reducing

FODMAP levels while maintaining bread quality. Only one pairing was tested in the present study,

but Carbonetto et al. (2020) found that co-culturing *M. humilis* with LAB strains affected loaf height

in a strain combination-dependent manner. However, as an obligately heterofermentative bacterium,

Fructilactobacillus sanfranciscensis is expected to affect loaf height in either a positive or neutral way

(Carbonetto et al., 2020), so it is possible that other sourdough-derived *F. sanfranciscensis* strains

would interact with *M. humilis* in a similar way due to their physiology.

In addition to their loaf technical properties (i.e., leavening), fermentation microbes from sourdough

contribute desirable aroma and flavour attributes to bread that consumers find appealing. Non-

conventional yeast species are known to generate particular aromas in bread, especially fruity ones

(Aslankoohi et al., 2016). Our panellists only detected these in breads co-fermented with *F.*

sanfranciscensis, suggesting that organic acids were the main volatile metabolites being detected.

More generally, our study found that consumers preferred baker's yeast bread, which echoes previous

findings from similar trials (Winters et al., 2019). However, while panellists' preferences for the

baker's yeast and co-culture breads fermented for 24 h at 25°C were similar, significant differences

between 6 and 24 h sourdough co-culture breads were detected. In terms of overall liking, colour,

sour/vinegar aromas and flavours, fruity aromas and flavours, and aroma complexity, the 24 h

sourdough co-culture breads scored higher. This demonstrates the potential sensory value in extending

dough fermentations. Many sourdough-associated volatile aroma compounds evaporate before or

during the baking process, however (Pétel et al., 2017), which may explain why the sourdough

microbe breads did not receive the highest scores for aroma complexity.

The research presented here builds on the small but growing body of literature focusing on Australian

sourdough starters. Our results show that increasing fermentation temperature slightly improves the

extent of fructan degradation by invertase-producing yeasts. However, this effect is only relevant to

the early stages of dough fermentation (< 6 h). Likewise, higher fermentation temperatures lead to

slightly improved leavening in yeasted doughs, but this leads to over-proving in extended dough

fermentations. This effect was less pronounced for *M. humilis*, whose gas production pattern was

slower, steadier, and less affected by proofing temperature increases. Therefore, we recommend that a

dough fermentation aimed at FODMAP reduction should feature an extended dough fermentation (> 6

h), ambient proofing temperature, and fermenting microbes which do not over-leaven bread in these

conditions.

Acknowledgements

469   We wish to thank bakers of Victoria who donated samples of their sourdough starters to us. For their

470   time and effort, we wish to thank all bread sensory study participants. Julie Hanot assisted in the

471   collection of sourdough starter samples and yeast isolation, and her work is gratefully acknowledged.

Funding

473   This work has been supported by a Research Training Program scholarship provided by the Australian

474   Government and the University of Melbourne to AEW.

References

Altilia, S., Foschino, R., Grassi, S., Antoniani, D., Dal Bello, F., & Vigentini, I. (2021). Investigating the growth kinetics in sourdough microbial associations. *Food Microbiology*, *99*, 103837. https://doi.org/10.1016/j.fm.2021.103837

Arora, K., Ameur, H., Polo, A., Di Cagno, R., Rizzello, C. G., & Gobbetti, M. (2021). Thirty years of knowledge on sourdough fermentation: A systematic review. *Trends in Food Science & Technology*, *108*, 71–83. https://doi.org/10.1016/j.tifs.2020.12.008

Aslankoohi, E., Herrera-Malaver, B., Rezaei, M. N., Steensels, J., Courtin, C. M., & Verstrepen, K. J. (2016). Non-Conventional Yeast Strains Increase the Aroma Complexity of Bread. *PLOS ONE*, *11*(10), e0165126. https://doi.org/10.1371/journal.pone.0165126

Atzler, J. J., Ispiryan, L., Gallagher, E., Sahin, A. W., Zannini, E., & Arendt, E. K. (2020). Enzymatic degradation of FODMAPS via application of β-fructofuranosidases and α-galactosidases- A fundamental study. *Journal of Cereal Science*, *95*, 102993. https://doi.org/10.1016/j.jcs.2020.102993

Biesiekierski, J. R., Rosella, O., Rose, R., Liels, K., Barrett, J. S., Shepherd, S. J., Gibson, P. R., & Muir, J. G. (2011). Quantification of fructans, galacto-oligosacharides and other short-chain carbohydrates in processed grains and cereals: Short-chain carbohydrates in grains and cereals. *Journal of Human Nutrition and Dietetics*, *24*(2), 154–176. https://doi.org/10.1111/j.1365-277X.2010.01139.x

Bigey, F., Segond, D., Friedrich, A., Guezenec, S., Bourgais, A., Huyghe, L., Agier, N., Nidelet, T., & Sicard, D. (2021). Evidence for Two Main Domestication Trajectories in Saccharomyces cerevisiae Linked to Distinct Bread-Making Processes. *Current Biology*, *31*(4), 722-732.e5. https://doi.org/10.1016/j.cub.2020.11.016

Brandt, M. J. (2007). Sourdough products for convenient use in baking. *Food Microbiology*, *24*(2), 161–164. https://doi.org/10.1016/j.fm.2006.07.010

Carbonetto, B., Nidelet, T., Guezenec, S., Perez, M., Segond, D., & Sicard, D. (2020). Interactions

between Kazachstania humilis Yeast Species and Lactic Acid Bacteria in Sourdough.

Microorganisms, *8*(2), 240. https://doi.org/10.3390/microorganisms8020240

Carbonetto, B., Ramsayer, J., Nidelet, T., Legrand, J., & Sicard, D. (2018). Bakery yeasts, a new

model for studies in ecology and evolution. *Yeast*, *35*(11), 591–603.

https://doi.org/10.1002/yea.3350

Gibson, P. R., & Shepherd, S. J. (2010). Evidence-based dietary management of functional

gastrointestinal symptoms: The FODMAP approach. *Journal of Gastroenterology and*

Hepatology, *25*(2), 252–258. https://doi.org/10.1111/j.1440-1746.2009.06149.x

Gobbetti, M. (1998). The sourdough microflora: Interactions of lactic acid bacteria and yeasts. *Trends*

in Food Science & Technology, *9*(7), 267–274. https://doi.org/10.1016/S0924-

2244(98)00053-3

Halmos, E. P., Power, V. A., Shepherd, S. J., Gibson, P. R., & Muir, J. G. (2014). A Diet Low in

FODMAPs Reduces Symptoms of Irritable Bowel Syndrome. *Gastroenterology*, *146*(1), 67-

75.e5. https://doi.org/10.1053/j.gastro.2013.09.046

Huynh, B.-L., Palmer, L., Mather, D. E., Wallwork, H., Graham, R. D., Welch, R. M., & Stangoulis, J.

C. R. (2008). Genotypic variation in wheat grain fructan content revealed by a simplified

HPLC method. *Journal of Cereal Science*, *48*(2), 369–378.

https://doi.org/10.1016/j.jcs.2007.10.004

Ispiryan, L., Borowska, M., Sahin, A. W., Zannini, E., Coffey, A., & Arendt, E. K. (2021). Lachancea

fermentati FST 5.1: An alternative to baker's yeast to produce low FODMAP whole wheat

bread. *Food & Function*, *12*(22), 11262–11277. https://doi.org/10.1039/D1FO01983J

Korakli, M., & Vogel, R. F. (2003). Purification and characterisation of mannitol dehydrogenase from

Lactobacillus sanfranciscensis. *FEMS Microbiology Letters*, *220*(2), 281–286.

https://doi.org/10.1016/S0378-1097(03)00129-0

Kurtzman, C. P., & Robnett, C. J. (1998). Identification and phylogeny of ascomycetous yeasts from

analysis of nuclear large subunit (26S) ribosomal DNA partial sequences. *Antonie van*

Leeuwenhoek, *73*, 331–371.

Lachance, M.-A., Boekhout, T., Scorzetti, G., Fell, J. W., & Kurtzman, C. P. (2011). Candida

Berkhout (1923). In *The Yeasts: A Taxonomic Study* (5th ed.). Elsevier Science & Technology.

https://doi.org/10.1016/B978-0-444-52149-1.00090-2

Laurent, J., Timmermans, E., Struyf, N., Verstrepen, K. J., & Courtin, C. M. (2020). Variability in

yeast invertase activity determines the extent of fructan hydrolysis during wheat dough

fermentation and final FODMAP levels in bread. *International Journal of Food

Microbiology*, *326*, 108648. https://doi.org/10.1016/j.ijfoodmicro.2020.108648

Liu, F., Hu, Z.-D., Yurkov, A., Chen, X.-H., Bao, W.-J., Ma, Q., Zhao, W.-N., Pan, S., Zhao, X.-M.,

Liu, J.-H., Wang, Q.-M., & Boekhout, T. (2024). Saccharomycetaceae: Delineation of fungal

genera based on phylogenomic analyses, genomic relatedness indices and genomics-based

synapomorphies. *Persoonia - Molecular Phylogeny and Evolution of Fungi*.

https://doi.org/10.3767/persoonia.2024.52.01

Loponen, J., & Gänzle, M. (2018). Use of Sourdough in Low FODMAP Baking. *Foods*, *7*(7), 96.

https://doi.org/10.3390/foods7070096

Menezes, L. A. A., Minervini, F., Filannino, P., Sardaro, M. L. S., Gatti, M., & Lindner, J. D. D.

(2018). Effects of Sourdough on FODMAPs in Bread and Potential Outcomes on Irritable

Bowel Syndrome Patients and Healthy Subjects. *Frontiers in Microbiology*, *9*, 1–7.

https://doi.org/10.3389/fmicb.2018.01972

Minervini, F., Lattanzi, A., De Angelis, M., Celano, G., & Gobbetti, M. (2015). House microbiotas as

sources of lactic acid bacteria and yeasts in traditional Italian sourdoughs. *Food

Microbiology*, *52*, 66–76. https://doi.org/10.1016/j.fm.2015.06.009

Pejcz, E., Spychaj, R., & Gil, Z. (2020). Technological methods for reducing the content of fructan in

rye bread. *European Food Research and Technology*, *246*(9), 1839–1846.

https://doi.org/10.1007/s00217-020-03537-5

Pétel, C., Onno, B., & Prost, C. (2017). Sourdough volatile compounds and their contribution to

bread: A review. *Trends in Food Science & Technology*, *59*, 105–123.

https://doi.org/10.1016/j.tifs.2016.10.015

R Core Team. (2024). *R: A Language and Environment for Statistical Computing* (Version 4.4.2)

[Computer software]. https://www.R-project.org/

Rogalski, E., Ehrmann, M. A., & Vogel, R. F. (2020). Role of Kazachstania humilis and

Saccharomyces cerevisiae in the strain-specific assertiveness of Fructilactobacillus

sanfranciscensis strains in rye sourdough. *European Food Research and Technology*, *246*(9),

1817–1827. https://doi.org/10.1007/s00217-020-03535-7

Rogalski, E., Ehrmann, M. A., & Vogel, R. F. (2021). Strain-specific interaction of Fructilactobacillus

sanfranciscensis with yeasts in the sourdough fermentation. *European Food Research and*

Technology, *247*(6), 1437–1447. https://doi.org/10.1007/s00217-021-03722-0

Schmidt, M., & Sciurba, E. (2021). Determination of FODMAP contents of common wheat and rye

breads and the effects of processing on the final contents. *European Food Research and*

Technology, *247*(2), 395–410. https://doi.org/10.1007/s00217-020-03633-6

Shewry, P. R., America, A. H. P., Lovegrove, A., Wood, A. J., Plummer, A., Evans, J., van den Broeck,

H. C., Gilissen, L., Mumm, R., Ward, J. L., Proos, Z., Kuiper, P., Longin, C. F. H., Andersson,

A. A. M., Philip van Straaten, J., Jonkers, D., & Brouns, F. (2022). Comparative compositions

of metabolites and dietary fibre components in doughs and breads produced from bread

wheat, emmer and spelt and using yeast and sourdough processes. *Food Chemistry*, *374*,

131710. https://doi.org/10.1016/j.foodchem.2021.131710

Smukowski Heil, C., Howell, K., & Sicard, D. (2022). Yeasts and Breadmaking. In *Yeasts: From*

Nature to Bioprocesses (Vol. 2, pp. 327–356). Bentham Science Books.

https://doi.org/10.2174/9789815051063122020013

Sprouffske, K., & Wagner, A. (2016). Growthcurver: An R package for obtaining interpretable metrics

from microbial growth curves. *BMC Bioinformatics*, *17*(1), 172.

https://doi.org/10.1186/s12859-016-1016-7

Struyf, N., Vandewiele, H., Herrera-Malaver, B., Verspreet, J., Verstrepen, K. J., & Courtin, C. M.

(2018). Kluyveromyces marxianus yeast enables the production of low FODMAP whole

wheat breads. *Food Microbiology*, *76*, 135–145. https://doi.org/10.1016/j.fm.2018.04.014

Van Kerrebroeck, S., Maes, D., & De Vuyst, L. (2017). Sourdoughs as a function of their species

diversity and process conditions, a meta-analysis. *Trends in Food Science & Technology*, *68*,

152–159. https://doi.org/10.1016/j.tifs.2017.08.016

Varney, J., Barrett, J., Scarlata, K., Catsos, P., Gibson, P. R., & Muir, J. G. (2017). FODMAPs: Food

composition, defining cutoff values and international application: Defining and adapting the

low-FODMAP diet. *Journal of Gastroenterology and Hepatology*, *32*, 53–61.

https://doi.org/10.1111/jgh.13698

Vaughan-Martini, A., & Martini, A. (2011). Saccharomyces Meyen ex Reess (1870). In *The Yeasts: A*

Taxonomic Study (5th ed.). Elsevier Science & Technology. https://doi.org/10.1016/B978-0-

444-52149-1.00061-6

Weiss, S., Samson, F., Navarro, D., & Casaregola, S. (2013). YeastIP: A database for identification

and phylogeny of *Saccharomycotina* yeasts. *FEMS Yeast Research*, *13*(1), 117–125.

https://doi.org/10.1111/1567-1364.12017

Winters, M., Panayotides, D., Bayrak, M., Rémont, G., Viejo, C. G., Liu, D., Le, B., Liu, Y., Luo, J.,

Zhang, P., & Howell, K. (2019). Defined co-cultures of yeast and bacteria modify the aroma,

crumb and sensory properties of bread. *Journal of Applied Microbiology*, *127*(3), 778–793.

https://doi.org/10.1111/jam.14349

Wittwer, A. E., Sicard, D., & Howell, K. S. (2022). Kazachstania humilis. *Trends in Microbiology*.

https://doi.org/10.1016/j.tim.2022.05.007

Yan, Y. L., Hu, Y., & Gänzle, M. G. (2018). Prebiotics, FODMAPs and dietary fiber—Conflicting

concepts in development of functional food products? *Current Opinion in Food Science*, *20*,

30–37. https://doi.org/10.1016/j.cofs.2018.02.009

Ziegler, J. U., Steiner, D., Longin, C. F. H., Würschum, T., Schweiggert, R. M., & Carle, R. (2016).

Wheat and the irritable bowel syndrome – FODMAP levels of modern and ancient species

and their retention during bread making. *Journal of Functional Foods*, *25*, 257–266.

https://doi.org/10.1016/j.jff.2016.05.019