Dissociating Contributions of Theta and Alpha Oscillations from Aperiodic Neural Activity in Human Visual Working Memory

Quirine van Engen[1]✉, Geeling Chau[2], Aaron Smith[1], Kirsten Adam[3], Thomas Donoghue[4],

Bradley Voytek[1, 5-7] ✉

[1]Department of Cognitive Science, University of California San Diego, La Jolla, CA 92093, USA
[2]Computation & Neural Systems, Caltech, Pasadena, CA 91125, USA
[3]Department of Psychological Sciences, Rice University, Houston, TX 77251, USA
[4]Department of Biomedical Engineering, Columbia University, New York, NY 10027, USA
[5]Halıcıoğlu Data Science Institute, University of California San Diego, La Jolla, CA 92093, USA
[6]Neurosciences Graduate Program, University of California San Diego, La Jolla, CA 92093, USA
[7]Kavli Institute for Brain and Mind, University of California San Diego, La Jolla, CA 92093, USA
✉e-mail: quvaneng@ucsd.edu

Number of pages: 19
Number of figures: 4
Number of words for abstract: 235
Number of words for introduction: 649
Number of words for discussion: 1477

Code and Data availability
All data used in this study are publicly available at:
https://osf.io/8xuk3/
Adam, K. C. S., Robison, M. K., & Vogel, E. K. (2018, June 25). Open Data 2018: Contralateral delay activity tracks fluctuations in working memory performance. https://doi.org/10.17605/OSF.IO/8XUK3

All code used for all analyses and plots are publicly available on GitHub at:
https://github.com/voytekresearch/WM_specparam

Acknowledgements
We thank Andrew Bender, Dillan Cellier, Ryan Hammonds, Blanca Martin-Burgos, Michael Preston, Eena Kosik, Sydney Smith and Christian Cazares for their advice and feedback on the manuscript. We also want to thank Kirsten Adam, and the Vogel lab for making their data public. Funding was provided by a NIH National Institute of General Medical Sciences grant R01GM134363-01 (to B.V.).

Author contributions
Q.v.E., K.A., T.D., and B.V. conceived of the experiments and contributed to the intellectual and methodological development of the ideas. Q.v.E., G.C., A.S., and T.D. wrote analysis code and analyzed data. Q.v.E., K.A., T.D., and B.V. wrote and edited the manuscript.

Competing interests
The authors declare no competing interests.

Abstract

While visual working memory (WM) is strongly associated with reductions in occipitoparietal 8-12 Hz alpha power, the role of 4-7 Hz frontal midline theta power is less clear, with both increases and decreases widely reported. Here, we test the hypothesis that this theta paradox can be explained by non-oscillatory, aperiodic neural activity dynamics. Because traditional time-frequency analyses of electroencephalopgraphy (EEG) data conflate oscillations and aperiodic activity, event-related changes in aperiodic activity can manifest as task-related changes in apparent oscillations, even when none are present. Reanalyzing EEG data from two visual WM experiments (n = 74), and leveraging spectral parameterization, we found systematic changes in aperiodic activity with WM load, and we replicated classic alpha, but not theta, oscillatory effects after controlling for aperiodic changes. Aperiodic activity decreased during WM retention, and further flattened over the occipitoparietal cortex with an increase in WM load. After controlling for these dynamics, aperiodic-adjusted alpha power decreased with increasing WM load. In contrast, aperiodic-adjusted theta power increased during WM retention, but because aperiodic activity reduces more, it falsely appears as though theta "oscillatory" power (e.g., bandpower) is reduced. Furthermore, only a minority of participants (31/74) had a detectable degree of theta oscillations. These results offer a potential resolution to the theta paradox where studies show contrasting power changes. We identify novel aperiodic dynamics during human visual WM that mask the potential role that neural oscillations play in cognition and behavior.

Significance statement (120)

Working Memory (WM) is our ability to hold information in mind without it being present in our external environment. Years of research focused on oscillatory brain dynamics to discover the mechanisms of WM. Here, we specifically look at oscillatory and non-oscillatory, aperiodic activity as measured with scalp EEG to test their significance in supporting WM. We challenge earlier findings regarding theta oscillations with our analysis approach, while replicating alpha oscillation findings. Furthermore, aperiodic activity is found to be involved in WM, over frontal regions in a task-general manner, and over anterior regions this activity is reduced with an increase the number of items that are remembered. Thus, we have identified novel aperiodic dynamics during human visual WM.

1. Introduction

We commonly encounter situations where we need to remember the color and identity of objects in our environment, ranging from simple tasks like cooking, to the games we play, to more complex interactions with our environment. Visual working memory (WM) refers to our ability to actively maintain information in mind, for a brief period of time, that is used to guide behavior to execute these tasks (Baddeley, 1992; D'Esposito & Postle, 2015). It is characterized by its limited capacity (Cowan, 2010; Miller, 1956; Oberauer et al., 2016).

Using electroencephalography (EEG), systematic changes in neural activity have been observed while participants maintain items in memory, even in the absence of continuing sensory input. In particular, two oscillatory bands have been identified as playing an important role in WM: frontal midline theta (4-7Hz, hereafter just "theta") (Berger & Sauseng, 2022; Hsieh & Ranganath, 2014; Jensen & Tesche, 2002), and occipitoparietal alpha (8-12Hz, hereafter, "alpha") (D'Esposito & Postle, 2015; Gevins & Smith, 2000; Halgren et al., 2002; Pavlov & Kotchoubey, 2022). Decreases in alpha power have been widely observed with increasing WM load (Fukuda et al., 2015, 2016), whereas hemispheric differences due to load have inconsistent findings (Adam et al., 2018, 2020; Medendorp et al., 2007; Sauseng et al., 2009; van der Werf et al., 2008; van Dijk et al., 2010). Topographical changes in alpha power can be used to decode the location of items in WM (Bae & Luck, 2018; Foster et al., 2016). In contrast, theta's role in

WM is less clear, with studies reporting a mix of WM-related power increases and decreases, even within the same study (Herweg et al., 2020; Pavlov & Kotchoubey, 2022).

This lack of clarity regarding the role of theta might be explained by non-oscillatory, aperiodic neural dynamics. Traditional time-frequency analysis methods – including wavelet analysis, and the filter-Hilbert approach – conflate neural oscillations with non-oscillatory, aperiodic activity (Donoghue, Haller, Peterson, et al., 2020; He, 2014). Historically often treated as task-irrelevant noise, there is a growing consensus that aperiodic activity is dynamic and task-relevant, with recent research highlighting its role in aging (Voytek et al., 2015), sleep (Höhn et al., 2024; Lendner et al., 2020), anesthesia (Colombo et al., 2019; Waschke et al., 2021), and disease (Helson et al., 2023; Karalunas et al., 2022; Smith, Kosik, van Engen, et al., 2023; Smith, Ma, et al., 2023). Preliminary evidence suggests that resting-state aperiodic activity is correlated with WM capacity (McKeon et al., 2024), and task-based aperiodic activity is functionally relevant in WM processes (Frelih et al., 2024; Gyurkovics et al., 2022; Virtue-Griffiths et al., 2022).

Here, we tested the hypothesis that mixed results regarding theta in human WM can potentially be resolved by accounting for event-related aperiodic dynamics. While it is possible the same applies to occipitoparietal alpha activity - it might be driven, in whole or in part, by aperiodic activity - previous work suggests that alpha oscillations, due to typically having higher power, are less likely to be masked by aperiodic dynamics than theta (Donoghue, Dominguez, et al., 2020). We illustrate three scenarios where task-induced changes to the neural power spectrum appear as theta bandpower decreases (**Fig. 1C**). The first scenario illustrates the - potentially incorrectly - standard model, whereby aperiodic activity is stable between task-states while narrowband theta oscillatory power dynamically changes. The second and third scenarios show no difference in relative oscillatory theta power (peak height), even though total theta bandpower decreases relative to baseline. Here, theta band effects are driven by aperiodic dynamics, regardless of whether detectable theta oscillations are present.

Here we reanalyzed two large (n = 74) human EEG datasets of participants performing a visual WM task (**Fig. 1A**) (Adam et al., 2018). Using spectral parameterization (**Fig. 1B**), we replicated traditional alpha effects even when accounting for aperiodic activity; power decreased with increasing WM load. The original paper used the filter-Hilbert method and reported task-related theta power decreases during WM retention. In contrast, we found aperiodic-adjusted theta power increased during retention, but a simultaneous decrease in aperiodic activity overwhelmed this relatively smaller theta oscillatory power increase. To traditional analysis methods it would appear as though theta power reduced when, in fact, the opposite occurred.

By disentangling aperiodic activity from narrowband oscillations, we found aperiodic activity dynamically decreases (spectral flattening) due to task-general demands in the frontal midline and occipitoparietal regions. This effect was task-dependent in occipitoparietal regions, with a higher reduction in aperiodic activity with increased WM load. Consistent with prior research (Bailey et al., 2020; Klimesch, 1999; Mitchell et al., 2008), we found evidence for theta oscillatory activity in less than half (42%) of the participants, whereas almost every participant (91%) showed clear alpha oscillatory activity. Our results demonstrate how implicit assumptions of our analysis methods can yield misleading conclusions. While theta oscillations clearly play a role in rodent hippocampal function relating to behavioral and cognitive tasks (Soltani Zangbar et al., 2020; Vanderwolf, 1969), the evidence for human frontal midline theta is more sparse and mixed (Herweg et al., 2020; Pavlov & Kotchoubey, 2022). Not only did our findings show correlational evidence of aperiodic activities involvement in human visual WM, we also demonstrated new insights into oscillatory dynamics by accounting for event-related aperiodic activity.

2. Methods

2.1. Study design from Adam et al. (2018)

All data used in this manuscript was obtained from OSF (https://osf.io/8xuk3/) (Adam et al., 2018).

2.1.1. Participants
Unique participants were recruited for Experiments 1 and 2. All participants were between the age of 18
and 35, with normal or corrected-to-normal visual acuity and color vision. Data from 31 participants were
collected in Experiment 1, and 48 participants in Experiment 2. For more details on inclusion and
exclusion criteria, see Adam et al. (2018). After EEG artifact rejection, 30 participants were considered for
further analysis in Experiment 1, and 44 in Experiment 2.
149
2.1.2 Task design
In both experiments, participants performed a whole-report visual WM task (**Fig. 1A**), in which they were
asked to remember the colors of differently colored squares across a blank screen retention period and
then recall the colors of all remembered items (Huang et al 2010; Adam et al, 2015). Throughout each
trial, participants were asked to fixate on a dot presented in the middle of the screen, except during the
response. A cue was presented for 1100 ms to indicate whether they had to memorize the items on the
right or the left side of the screen. The memory array was presented for 250 ms followed by a retention
period of 1300 ms, after which a response for each cued item had to be given (**Fig. 1B**). The number of
items were equal on each side to balance the visual stimulation within each visual hemifield. This design
was chosen to ensure hemispheric differences in neural activity were driven by cognitive demands, rather
than perceptual demands (Vogel & Machizawa, 2004).
Each block consisted of 30 trials. For Experiment 1, each block had 10 trials for three different set
sizes, one, three or six, and Experiment 2 contained 30 trials of set size six. The conditions in Experiment
1 were defined as the three set sizes, whereas in Experiment 2 two conditions were defined based on the
performance on each trial: if more than three items were correctly answered, it was deemed a "high
performance" trial, versus having less than three items correct was deemed a "poor performance" trial.
166
2.1.3 EEG measurements
EEG passive electrodes and cap were from ElectroCap International (Eaton, OH). Recordings were done
on an SA Instruments Amplifier running LabView 6.1 (Fife, Scotland). The cap contains 20 electrodes
according to the 10/20 international sites. Two supplementary electrodes were placed over the occipital
lobe: OL was placed between T5 and O1, and OR was placed between T6 and O2. In addition,
electrooculography (EOG) was recorded to measure horizontal eye movements and blinks. For more
detailed information (Adam et al., 2018; Foster et al., 2016).
174
2.1.4 EEG preprocessing
Data were available from the original paper's repository, already pre-processed and epoched, but not
baselined. Pre-processing steps included: automatic removal of trials with eye blinks or eye movements
using a set threshold, automatic removal of trials with excessive noise using a peak-to-peak threshold
rejection, and finally a visual inspection of the data to determine whether the automated steps were
successfully implemented. More details can be found in Adam et al. (2018). Here, we used the non-
baselined epochs.
The same electrode clusters were used as defined by the original paper (Adam et al., 2018) for
replicating their original results, and the novel analysis conducted in this paper. These clusters for
lateralized alpha were left and right occipital and parietal regions (O1, OL, P3, PO3, T5, O2, OR, P4, PO4,
and T6), and for theta a cluster over the frontal-midline (F3, F4, Fz).
186
2.2 Replication of original results
A detailed description of the replication methods and results can be found in the Jupyter notebooks hosted
at GitHub.
To replicate the original results we performed separate bandpass filtering in the theta (4-7Hz) and
alpha (8-12Hz) bands. We then performed a Hilbert transform to compute the analytic amplitude of these
frequency bands over time. This amplitude was baseline corrected to 400 ms before cue onset.
For statistical analysis, theta and alpha power was averaged over the retention period (400 - 1500
ms after memory array onset) per condition per the original approach. A two-way repeated-measures
ANOVA was performed on alpha power with condition (set size for Experiment 1 or performance for
Experiment 2) and lateralization as factors. Note that the factor set size or performance is comparable to a
whole-field alpha approach, and the second factor adds the lateralization effect. A pairwise t-test was used
on theta power to investigate the difference between good and poor performance.

199
2.3 Spectral parameterization: separating oscillatory from aperiodic activity
2.3.1 The specparam model
Spectral parameterization (Donoghue, Haller, Peterson, et al., 2020) models a power spectrum as a
combination of an aperiodic component (fit as a Lorentzian) and putative oscillatory components that rise
above the aperiodic signal (fit as Gaussians). Here, the parameters characterizing the aperiodic
component are the spectral exponent and the offset. Three parameters characterize the Gaussian periodic
components: peak height (relative to the aperiodic signal), center frequency, and bandwidth (**Fig. 1B**).
More details can be found in (Donoghue, Haller, Peterson, et al., 2020).

2.3.2 Identification of participants exhibiting oscillatory activity
Because oscillations are not consistently present (Jones, 2016), spectral parameterization helps to
explicitly identify participants and electrodes where there are putative oscillations (defined here as having
a significant "bump" in the power spectrum within the theta or alpha range above aperiodic activity). This
triggered an additional inclusion criterion, different from the original report, in that oscillatory activity should
only be analyzed from participants who exhibited putative oscillations. Thus, for alpha power in
Experiments 1 and 2, participants were required to have a parameterized spectral peak between 8 and 12
Hz in the occipitoparietal electrodes. For theta power, participants were required to have a peak between
4 and 7 Hz that was distinct from alpha activity on frontal midline electrodes (detailed below). To perform
this analysis, a power spectrum was generated for every trial, for the whole trial length using Welch's
method and a Hanning window of two times the sampling frequency (2*250 samples). The model was
applied to the power spectrum in the frequency range of 2-40 Hz. Peak width limits were set from 2-8 Hz.
To minimize overfitting of relatively flat spectra without peaks, *specparam* requires an amplitude threshold
for determining the height of the Gaussian, relative to the aperiodic signal, to be considered a peak. This
threshold was set at 0.2 μV^2/Hz for the alpha frequency band, and 0.1 μV^2/Hz for theta frequency band.
Different absolute thresholds were chosen for the bands because theta power is typically lower than alpha
power, relative to the aperiodic fit. For both experiments, *specparam* successfully discriminated between
participants who exhibited alpha power and those who did not. This was further confirmed by visual
inspection. However, discriminating participants based on theta power proved to be more difficult because
not all participants exhibited a distinct theta peak. A portion of participants also exhibited alpha power on
the frontal midline electrodes overlapping with the theta frequency range. Therefore, determining whether
a participant exhibits theta power was guided by *specparam* and confirmed with manual visual
classification to avoid lower frequency alpha peaks to be considered theta.

2.3.3 Generating single trial power spectra and applying specparam
A power spectrum was created for each trial, for all electrode groups – contralateral occipitoparietal, the
ipsilateral occipitoparietal, and frontal midline. The same electrodes were used as in the replication
results. A time window from 400 to 1500 ms after memory array onset (1100 ms) was used for the
retention period to produce power spectra, excluding the early perceptual ERP time points (Kałamała et
al., 2024). For the baseline period, we used the cue period from -1100 until memory array onset (1100
ms). This decision is due to the ITI only being 500 ms long. Power spectra were created using Welch's
method and a Hanning window that was equal to the length of the data (1100 ms).
The *specparam* model was applied to these power spectra using the same settings described in section
2.2.3. Four outputs were collected:
     1) The peak height within a frequency range of 8 to 12 Hz for alpha power and 4 to 7 Hz for theta
power, for both the retention period and baseline period. If a peak was absent, the power was considered
"0".
     2) Abundance of oscillations, defined as the percentage of trials within a condition in which a peak
in the alpha or theta range is observed.
     3) The aperiodic exponent.
     4) The aperiodic offset.
     Additionally, bandpower was collected as the average power within the frequency bands of
interest. Resulting in five measurements per trial.
     A baseline correction was performed by subtracting the baseline values of these parameters from
the values of the retention period in a trial-by-trial fashion for all five outputs. Afterwards, each output was
averaged per condition, participant, and across electrodes within each cluster.

255
2.3.4 Inclusion criteria for single-trial model fit
Model fit inspection was done by calculating the mean R^2 per condition and per participant. Trials with a model fit lower or higher than twice the standard deviation of the mean were removed from further analysis. Afterwards, participants were checked again for having enough trials per condition (minimum of 75 in Experiment 1, and minimum of 40 in Experiment 2) to remain included in this study. This resulted in excluding four participants in Experiment 1 and another four in Experiment 2 for the occipitoparietal analysis. No participants were excluded based on this criterion in either experiment for the frontal midline analysis.

2. 4 Behavioral analysis and correlation to EEG metrics
WM capacity was calculated in two ways. The first was based on the main task while EEG was recorded, by calculating the average number of items correctly remembered over all trials. The second method was based on the performance during a baseline WM task which followed the same procedure in both experiments (referred to as the Color Change Detection Task in Adam et al., 2018). This WM estimate was calculated per set size, and then averaged to get a single value representing WM capacity per participant, expressed as an estimate ("K") which is calculated with the following formula:

$K = N \times (H - FA)$
N = set size
H = hit rate (proportion of correct trials)
FA = false alarm rate (proportion of incorrect no-change trials)

2.5 Statistical analysis
The dependent variables in both experiments were the oscillatory power relative to the aperiodic signal, oscillation abundance, aperiodic exponent, aperiodic offset, and bandpower

For the occipitoparietal region, separate two-factor repeated measures ANOVAs were applied on all dependent variables. The factors were set size in Experiment 1 (one, three, six) or performance in Experiment 2 (good, poor) and lateralization (ipsilateral, contralateral). In cases where the ANOVA was significant, pairwise t-tests with Bonferroni correction were applied as post-hoc tests to examine the differences between the levels within each factor. For the frontal midline region, separate repeated measures ANOVA were applied to test for significant differences on all dependent variables between set sizes (Experiment 1), and pairwise t-tests were applied similarly to test statistical significance of performance (Experiment 2). Additionally, paired t-tests were applied on each variable averaged over the conditions to test whether there was a significant difference between the retention and baseline period.

Shapiro-Wilk tests were used to test for normality, Levene's test for equal variance, and Mauchly's test were used to test for sphericity. If sphericity was violated, the Greenhouse-Geisser correction was applied for the factorial repeated measures ANOVAs. If one or more assumptions were violated, a Friedman test was used instead of a repeated measures ANOVA, or the Wilcoxon signed rank test was used instead of a pairwise t-test. Cohen's d was reported as effect size for t-tests, partial eta-squared for ANOVA tests, and the Common-Language Effect Size (CLES) for non-parametric tests.

For our exploratory analysis, we investigated correlations between WM capacity and aperiodic exponent or offset, Pearson correlations were calculated with a Bonferroni threshold correction.

2.6 Software
All data were processed using the Python programming language (version 3.7.1). Signal processing was done with the *neurodsp* module (version 2.1.0) (Cole et al., 2019), and power spectra were fitted with the *specparam* toolbox (version 1.0.0) (Donoghue, Haller, Peterson, et al., 2020). All statistics were performed in Pingouin (version 0.5.2) (Vallat, 2018), and plots were created using Matplotlib (version 3.0.2) (Hunter, 2007) and seaborn (version 0.9.0) (Waskom, 2021).

2.7 Data and code statement
Data used in this project is from Adam et al. (2018), and is available at OSF (https://osf.io/8xuk3/). The code for the analyses carried out in this paper are available at our GitHub page.

3. Results

3.1 Successful replication of the original paper's results

All findings from the original paper (Adam et al., 2018) were first successfully replicated before moving onto the spectral parameterization analysis. These results can be found on our GitHub repository. In short, an increase in WM load was associated with a bigger decrease in lateralized occipitoparietal alpha power, and when separated by trial accuracy, theta power was higher for good WM performance than poor.

3.2 Spectral parameterization: separating oscillatory from aperiodic activity

3.2.1. Participants

For this analysis, only participants exhibiting alpha oscillations over occipitoparietal electrodes or theta power over frontal midline electrodes were included. Furthermore, they needed to pass the minimum number of trial requirements as well. All 30 participants in Experiment 1, and the majority (37 out of 44) in Experiment 2 exhibited clear peaks in the alpha range according to the spectral parameterization criteria. After excluding participants who did not meet the minimum number of trials requirement, we ended with a sample size of 26 in Experiment 1, and 26 in Experiment 2. In contrast, only 12 out of 30 participants in Experiment 1, and 19 out of 44 in Experiment 2 met the criteria for exhibiting putative theta oscillations, identified as peaks in the theta range. The final sample size after excluding participants is 10 in Experiment 1, and 17 in Experiment 2 (**Table 1**).

To increase the sample size and therefore statistical power for the aperiodic analyses on frontal midline electrodes, we decided to include participants that did not exhibit putative theta oscillations. In Experiment 1, 15 from the 18 participants without oscillations satisfied the minimum trial count requirement and were included here, bringing the total to 25 participants. In Experiment 2, 17 from the 27 met these criteria in Experiment 2, bringing that total to 34.

3.2.2 Experiment 1: Frontal midline aperiodic-adjusted theta power and aperiodic activity

All variables violated sphericity. From the 12 participants with putative theta oscillations, 2 were rejected due to low trial count. For the remaining 10 participants included in this analysis concerning theta oscillations, aperiodic-adjusted theta power was not significantly different between set sizes ($W(2) = 0.07$, $p = 0.50$). Theta power over the three set sizes was significantly higher during the retention period compared to baseline ($t(9) = 2.72$, $p = 0.024$, Cohen's $d = 0.54$). Similar to the theta power results, theta abundance was not significantly different between set sizes ($W(2) = 0.01$, $p = 0.90$). In contrast to power, there was a significant decrease in abundance from baseline over all conditions ($W = 7$, $p = 0.037$, CLES $= 0.36$).

Frontal midline aperiodic exponent and offset were calculated for all participants included ($n = 25$), not just the participants that also had theta power. The frontal aperiodic exponent was not significantly different between set sizes ($W(2) = 0.04$, $p = 0.34$), but was significantly decreased from baseline over all conditions ($W = 66$, $p = 5.6*10^{-3}$, CLES $= 0.48$). A similar pattern was observed for the frontal midline offset; no significant differences between set sizes ($W(2) = 0.046$, $p = 0.30$), but aperiodic offset was significantly reduced from baseline over all conditions ($W = 0.046$, $p = 8.3*10^{-5}$, CLES $= 0.44$). Theta bandpower was not significantly different between set sizes ($W = 0.028$, $p = 0.48$), but was also significantly reduced during the retention period over all set sizes ($W = 89$, $p = 2.9*10^{-2}$, CLES $= 0.46$).

These results, leveraging the novel spectral parameterization method, suggest that theta oscillations are less frequently observed over trials in the retention period compared to baseline, but they demonstrate higher power (peak height) when present. Additionally, we found that the aperiodic exponent and offset reduced significantly during the WM retention period, but no differences due to WM load were observed in any of the frontal midline metrics for Experiment 1.

3.2.3 Experiment 2: Frontal midline aperiodic-adjusted theta power and aperiodic activity

Seventeen participants met all inclusion criteria for theta oscillation analyses. Aperiodic-adjusted theta power was not significantly different between poor and good performance ($t(16) = -1.59$, $p = 0.13$, Cohen's $d = -0.48$), nor was theta power different during the retention period from baseline ($t(16) = 2.05$, $p = 0.057$, Cohen's $d = 0.36$). Similarly, theta abundance was not significantly different between poor and good performance ($t(16) = 1.08$, $p = 0.30$, Cohen's $d = 0.41$), nor was it different from baseline to retention ($t(16) = -1.62$, $p = 0.12$, Cohen's $d = 0.23$).

The midline frontal aperiodic exponent and offset were calculated for all included participants ($n=34$), not just the participants that also had theta power. Neither were normally distributed; therefore the

non-parametric Wilcoxon signed-rank test was used to determine statistical significance. The exponent
was not significantly different between performances (W = 280, p = 0.77, CLES = 0.54), but it was
significantly decreased from baseline (t(32) = -2.37, p = 0.024, Cohen's d =0.14). Aperiodic offset was not
significantly different between performances (W = 283, p = 0.81, CLES = 0.52), offset was significantly
reduced from baseline (t(32) = -4.09, p = $2.7*10^{-4}$, Cohen's d = 0.27). Theta bandpower in showed no
difference between performances (t(25) < 1.0), but was significantly reduced during the retention period
from baseline (W = 135, p = $9.6*10^{-3}$, CLES = 0.45).
In contrast to Experiment 1, we found that theta abundance as identified via spectral
parameterization did not differ by task-state (baseline vs. retention) nor by task performance. Similarly, to
Experiment 1, task state (retention vs. baseline) induced a significant reduction in aperiodic exponent and
offset. These results stand in contrast to the original paper that found higher theta activity in trials with
good performance than poor performance, using the traditional filter-Hilbert analysis approach that mixes
aperiodic and oscillatory activity. Thus, it comes as a surprise to not observe differences between
performances for both aperiodic measures, nor for aperiodic-adjusted theta power.
381
3.2.4 Experiment 1: Occipitoparietal aperiodic-adjusted alpha power and aperiodic activity
Using a two-way repeated measures ANOVA, there was a main effect of set size on aperiodic-adjusted
occipitoparietal alpha power (F(2,50) = 6.25, p = $3.8*10^{-3}$, η_p^2 = 0.20), but not for lateralization (F(1,25) =
1.17, p = 0.29, η_p^2 = 0.045). Alpha power in set size one was significantly higher than set size three (t(25)
= 2.10, p = 0.046, Cohen's d = 0.39), and set size six (t(25) = 3.10, p = $4.7*10^{-3}$, Cohen's d = 0.64), but not
different between set size three and six (t(25) = 1.59, p = 0.13, Cohen's d = 0.18). The interaction between
set size and lateralization was not significant (F(2,50) = 2.95, p = 0.061, η_p^2 = 0.11). Alpha abundance was
significantly different between set sizes (F(2,50) 3.59, p = 0.035, η_p^2 = 0.13), with the post-hoc only
showing a significant increase from set size one to six (t(25) = -2.44, p = 0.022, Cohen's d = -0.40). Main
effect of lateralization was not significant, nor in the interaction (F < 1.0).
There was a main effect of set size on occipitoparietal aperiodic exponent such that it decreased
with increasing set size (F(2,50) = 5.99, p = $4.6*10^{-3}$, η_p^2 = 0.19). A post-hoc test shows that the difference
is significant between set sizes one and six (t(25) = 3.08, p = $5.0*10^{-3}$, Cohen's d = 0.52), and between set
sizes three and six (t(25) = 2.35, p = 0.027, Cohen's d = 0.26). No significant effect was found for
lateralization, nor was there an interaction effect between lateralization and set size (F < 1.0).
Furthermore, exponent was significantly decreased from baseline over all conditions (t(25) = -5.43, p =
$1.2*10^{-5}$, Cohen's d =0.22). Occipitoparietal aperiodic offset was also significantly decreased with
increasing set sizes (F(2,50) = 7.68, p = $1.2*10^{-3}$, η_p^2 = 0.23). A post-hoc pairwise t-test shows that this
difference is between set sizes one and six (t(25) = 3.30, p = $2.9*10^{-3}$, Cohen's d = 0.64), and between set
sizes three and six (t(25) = 3.01, p = $5.9*10^{-3}$, Cohen's d = 0.34). Furthermore, over all conditions the
offset was significantly lower during the retention period compared to baseline (t(25) = -6.93, p = $2.9*10^{-7}$,
Cohen's d = 0.32). Bandpower showed a main effect of set size (F(2,50) = 7.18, p = $1.8*10^{-3}$, η_p^2 = 0.22),
but not for lateralization (F(1,25) < 1.0) or their interaction (F(2,50) = 2.98, p = 0.060, η_p^2 = 0.11). Post-hoc
tests showed that all three set sizes were significantly different from each other. Set size one had the
lowest bandpower, lower than set size three (t(25) = 2.15, p = $4.2*10^{-2}$, Cohen's d = 0.36), and set size six
(t(25) = 2.95, p = $6.8*10^{-3}$, Cohen's d = 0.54), and set set size three was lower than set size six (t(25) =
3.74, p = $9.5*10^{-4}$, Cohen's d = 0.15).
In summary, we observed alpha peak height decrease with an increase in WM set size, but no
differences between hemispheres. A lack of lateralization is in contrast with the original paper's findings.
However, lateralization effects in alpha power in the literature are mixed (Adam et al., 2018, 2020;
Medendorp et al., 2007; Sauseng et al., 2009; van der Werf et al., 2008; van Dijk et al., 2010). Aperiodic
exponent and offset showed similar results, a decrease with an increase in WM set size, on top of an
overall decrease during the retention period from baseline. These results show that aperiodic activity is
involved in WM processes over occipitoparietal regions.
416
3.2.5 Experiment 2: Occipitoparietal aperiodic-adjusted alpha power and aperiodic activity
For aperiodic-adjusted alpha oscillations, we observed a significant main effect of performance (F(1,25) =
4.78, p = 0.038, η_p^2 = 0.16), in which power was lower in the good performance condition compared to the
poor performance condition. But lateralization nor the interaction effect were significant (F < 1.0).
Alpha abundance did not differ as a function of performance, lateralization, or their interaction (F < 1.0).

Occipitoparietal aperiodic exponent was not significantly different between performance, nor the
interaction between performance and lateralization (F < 1.0), but it was for lateralization (F(1,25) = 4.34, p
= 0.048, η_p^2 = 0.15) in which the contralateral hemisphere had a lower exponent than the ipsilateral
hemisphere. The exponent was also significantly reduced during the retention period compared to
baseline over all conditions (t(25) = -4.99, p = 3.9*10^{-5}, Cohen's d = 0.24). A similar pattern was observed
for the offset; the contralateral hemisphere had a lower offset than the ipsilateral hemisphere (F(1,25) =
6.08, p = 0.021, η_p^2, = 0.20), and an overall reduction was found compared to baseline (t(25) = -6.41, p =
1.0*10^{-6}, Cohen's d = 0.38). Whereas alpha bandpower showed a main effect of set size in Experiment 1,
no significant results were observed between good or poor performance in Experiment 2 (F < 1.0).
Alpha peak height was lower on good performance trials than poor, which is in line with our results
for set sizes in Experiment 1. However, aperiodic exponent and offset did not show similar patterns as in
Experiment 1, but it was lower over the ipsilateral hemisphere compared to the contralateral hemisphere,
with this being the only time we observe lateralization effects. We did also observe overall reduced
exponent and offset during the retention period from baseline. Hinting at a more WM task-general over
task-specific involvement.
437
3.3 Correlations between working memory capacity and task-states
Given the importance that theta oscillations are said to play in WM, we next sought to examine whether
there was a difference in WM capacity between participants for whom there was, versus was no,
detectable theta oscillations with spectral parameterization. To test this, we looked at WM capacity in two
ways, either as estimated from the performance on the WM task performed during the EEG experiment, or
as measured from a baseline WM task collected prior to EEG recording. Notably, these two metrics of WM
capacity were correlated within participants for both Experiment 1 (r(25) = 0.56, p = 3.37 * 10^{-3}) and
Experiment 2 (r(33) = 0.41, p = 0.018).
We found no significant WM capacity differences between the groups of participants who
exhibited or did not exhibit theta oscillations when measured from the main EEG task or from the baseline
task, in either experiment (p > 0.4 in all instances). This indicates that WM capacity is not reliant on
whether participants exhibited measurable theta oscillations.
In an additional exploratory analysis, we examined whether either measures of WM capacity
correlated to aperiodic activity, exponent or offset from either frontal midline or occipitoparietal regions.
We examined this using average pre-stimulus baseline aperiodic activity, during the retention period, and
in the within-participant change from baseline to retention. This resulted in 24 correlations, with a
Bonferroni corrected threshold of p < 2.08*10^{-3}. None of those correlations were significant under this
criterion.
Finally, we noticed that there appeared to be a difference in the overall aperiodic-adjusted theta
power between Experiments 1 and 2 (Fig. 3 A&G). Notably, in Experiment 1 set size could vary between
one, three or six items, whereas in Experiment 2 the set size was constant at six items. To verify this
visual observation, we performed t-test on relative theta power during the baseline, retention period, and
their difference between Experiment 1 and 2. Relative aperiodic-adjusted theta power was significantly
higher during Experiment 1 for both the baseline (t(15.85) = 2.57, p = 0.020, Cohen's d = 1.09) and
retention periods (t(12.49) = 2.56, p = 0.024, Cohen's d = 1.17). However the difference in theta oscillatory
power between baseline and retention was not significant (t(14.83) = 1.36, p =0.19, Cohen's d = 0.59).
These results indicate that aperiodic-adjusted theta power is overall higher during Experiment 1
with a variable WM set size, compared to Experiment 2 with a constant high WM set size, but the task-
induced changes in relative theta power from baseline are similar. More information and figures can be
found on our GitHub page.
468
4. Discussion
Many theories of human visual WM rely on theta and alpha oscillatory mechanisms for coordinating
information flow between brain regions and as a neural code for memory (Lisman & Jensen, 2013; Palva
& Palva, 2007). However, these theories are built on the flawed assumption that event-related changes in
those frequency bands always capture oscillations. Here, we sought to explicitly examine whether and to
what degree theta and alpha oscillatory activity is distinct from aperiodic activity in WM. Explicitly
controlling for aperiodic activity, we find that fewer than half of participants (42%) had an identifiable theta
peak in their power spectra. While not conclusive evidence for a lack of oscillations, this suggests that if

frontal midline theta oscillations are present in human EEG during WM, they are weak and/or rare.
Therefore, it is likely that previous studies investigating the role of frontal midline theta in WM were more
often than not largely capturing non-oscillatory aperiodic activity in the majority of participants, rather than
true theta oscillations. If that is true, then it means that, while frontal midline theta may play a role in
human WM, it is not a requirement for successful memory performance on any given trial.

Aperiodic-adjusted theta power increased during the retention period compared to baseline, but,
consistent with previous analyses of these data, total theta bandpower actually decreased. This paradox,
whereby theta simultaneously increases and decreases during WM retention, is mirrored by many mixed
results regarding theta in human memory(Herweg et al., 2020; Pavlov & Kotchoubey, 2022). We believe
this paradox can be explained by considering aperiodic activity, since we observed significant aperiodic
decreases during WM retention, but surprisingly not between WM loads. If these aperiodic effects are
ignored, they can mask and muddle the underlying oscillatory theta effects. **Thus, the mixed results**
regarding theta in human WM can partly be resolved by accounting for aperiodic dynamics. The
rarity of theta oscillations is an interesting contrast to the relative ubiquity of alpha oscillations. Even when
accounting for aperiodic activity, we replicated the common finding that alpha power decreases with
increasing WM load. A load-dependent decrease was also found for aperiodic activity. Thus, both ongoing
alpha oscillations and aperiodic activity support visual attention and WM retention.

Importantly, we identified a task-general, and load-dependent decrease of aperiodic activity during
WM retention. Aperiodic activity has potential links to underlying physiological mechanisms, offering a
potential bridge across levels of analysis, relating spiking activity to LFP, and, by extension, scalp EEG.
The aperiodic exponent at least partially reflects the balance of excitation and inhibition (E:I), as has been
shown in a simple computational model involving GABA and AMPA currents (Gao et al., 2017), with
pharmacological interventions (Colombo et al., 2019; Waschke et al., 2021), and using magnetic
resonance spectroscopy imaging (McKeon et al., 2024). However, research using biophysical modeling
challenges this perhaps overly simplistic hypothesis (Brake et al., 2024). Nevertheless, if aperiodic activity
serves as an indicator of E:I, we can make insightful connections with computational models of WM that
highlight the importance of a balanced E:I network to sustain activity for maintaining information (Lu et al.,
2023; Wang, 1999). This sustained spiking activity is mostly asynchronous, and therefore has limited
contribution to oscillating activity (Compte et al., 2000), making aperiodic activity a better candidate to
study. Since these computational models are based on prefrontal cortex anatomy and dynamics, it is
intriguing we only found load-dependent aperiodic decreases in occipitoparietal regions, but not over the
frontal midline.

Thus, the task-general decreases in aperiodic activity observed in both frontal midline and
occipitoparietal regions could indicate a shift in E:I balance that is supported by computational models of
sustained activity during the maintenance period. The task-specific modulations in occipitoparietal regions
support the sensory recruitment theory, in which sensory regions are actively involved in the maintenance
of WM items (D'Esposito & Postle, 2015). In this case, occipitoparietal regions are involved in the
maintenance of visual stimuli through asynchronous activity as well as oscillatory activity, whereas the
frontal midline could be considered a top-down signal (Lara & Wallis, 2015).

While we found aperiodic flattening after stimulus onset on a visual WM task in frontal and anterior
regions, others found steepening or mixed results (Akbarian et al., 2023; Frelih et al., 2024; Kałamała et
al., 2024; Lendner et al., 2023; Virtue-Griffiths et al., 2022). Notably, some of these papers (Akbarian et
al., 2023; Frelih et al., 2024) used the n-back task, which mixes motor preparation responses with WM
maintenance activity (Pavlov & Kotchoubey, 2022). Mixed results were also observed on a visual
continuous recall WM task; increased exponents over the frontal midline, and decreases over
occipitoparietal regions, with an increase in exponent being associated with improved performance on a
battery test (Virtue-Griffiths et al., 2022). Non-WM cognitive tasks suffer a similar fate; some find
decreases in exponent after stimulus onset (Podvalny et al., 2015; Waschke et al., 2021), others find
biphasic responses (Kałamała et al., 2024). These inconsistent results can be partly explained by choice

of task and whether fast motor responses are required, but also potentially driven by a lack of consistency
in how aperiodic activity is measured and reported (Donoghue, 2024).
After correcting for aperiodic activity, we no longer observed a positive relationship between theta
activity and WM load as in the original study. This lack of replication is likely driven by two factors. First,
the large, event-related decrease in aperiodic activity is conflated with theta oscillations dynamics, which
the filter-Hilbert approach cannot disentangle. Second, only 42% of participants had detectable theta
oscillations according to spectral parameterization. Therefore, we could only analyze theta activity related
to performance for a subset of participants. This finding is consistent with other research showing theta
oscillations are surprisingly scarce within, and across, groups of participants (Bailey et al., 2020; Mitchell
et al., 2008; Wilson et al., 2022). Notably, we did not find behavioral differences between participants
exhibiting theta oscillations or not, and thus suggest the ability to detect theta oscillation at the scalp is due
to anatomical (cortical folding, hippocampus orientation) or physiological, rather than behaviorally
functional, differences.
In contrast to theta, almost all participants had detectable alpha oscillations (91%), which might
contribute to the finding that alpha oscillations are less susceptible to masking by aperiodic dynamics than
theta (Donoghue, Dominguez, et al., 2020). Correcting for aperiodic activity, we found an effect of relative
alpha power due to WM load and performance. Furthermore, lateralized effects were only observed during
the experiment using a high load only and separating trials based on WM accuracy, in which relative alpha
power was more suppressed on the contralateral visual cortex of the attended hemifield. This is not a
surprising finding, since the literature on lateralization effects are inconsistent (Adam et al., 2018, 2020;
Fukuda et al., 2016; Sauseng et al., 2009). In summary, alpha oscillations in these datasets were involved
in both visual WM and attention mechanisms (D'Esposito & Postle, 2015; Gevins & Smith, 2000; Halgren
et al., 2002; Pavlov & Kotchoubey, 2022).
Though our results show aperiodic activity is involved in visual WM, there are methodological
limitations to be addressed. First, performing spectral parameterization on task-based data means that we
have shorter time windows than resting-state paradigms (Colombo et al., 2019; Helson et al., 2023;
McKeon et al., 2024; Smith, Kosik, van Engen, et al., 2023; Waschke et al., 2021), resulting in noisy single
trial power spectra. Although still feasible (Frelih et al., 2024; Höhn et al., 2024; Kałamała et al., 2024;
Podvalny et al., 2015; Virtue-Griffiths et al., 2022; Waschke et al., 2021), we minimized this noise by
rejecting trials based on model fits that were too low (indicating a bad fit), or too high (indicating overfitting
noise), while still averaging the model output over a sufficient number of trials. Another issue became
apparent while visually examining power spectra to determine whether participants exhibited oscillatory
activity within the theta or alpha band. These canonical frequency bands are defined adjacent to one
another, and some participants exhibited easily identifiable peaks in both bands, some only in one band,
and others in neither. However, there were edge cases in which a lower center frequency alpha peak bled
into the theta range, making them difficult to distinguish and to determine whether a peak should be
considered alpha, or theta based on *a priori* frequency bands. This is already a concern for bandpass
filters in which neighboring band activity infiltrates the band of interest (Pavlov & Kotchoubey, 2022).
Here, we have provided correlational evidence that aperiodic activity is involved in human visual
WM and shown that it can mask frontal midline theta oscillatory dynamics. One way to causally
disentangle the role of oscillatory versus aperiodic dynamics in human WM would be to leverage non-
invasive neurostimulation. Neurostimulation is commonly used to study the causal role of oscillations, via
transcranial Alternating Current Stimulation (tACS) (Grover et al., 2023). Because aperiodic activity
consists of arrhythmic unstructured fluctuations, transcranial Random Noise Stimulation (tRNS) - where a
white noise spectrum electrical current is introduced into the brain - is a better candidate to target
endogenous aperiodic activity, thought white noise is quite different from endogenous aperiodic activity. It
has been shown that tRNS flattens the spectrum (van Bueren et al., 2023), thereby effectively showing
that aperiodic activity can be targeted with non-invasive electrical stimulation. Thus, it may be possible to

test the causal role of aperiodic activity in human visual WM using aperiodic stimulation that more closely
mimics endogenous aperiodic activity.
While we did not observe load-dependent aperiodic changes over the frontal midline, we did
reveal the possibility that previous frontal midline theta results that do not correct for aperiodic activity
likely do not reflect theta oscillations. It will be interesting to investigate if our results based on extracranial
EEG translate to intracranial EEG and animal models of WM, and if our results are driven by the
imprecision of EEG to detect frontal midline theta oscillations. While our results do not invalidate previous
research into extracranial theta oscillations in relation to WM, they challenge popular and widely held
beliefs regarding the mechanistic role for theta oscillations to group or segregate channels of information
(Lisman & Jensen, 2013). After all, if people can still reliably perform WM tasks in the absence of
detectable theta oscillations, then either our detection ability is severely flawed, or our theories are
incomplete (Van Bree et al., 2024).
586
587

Figure 1: Overview of experiment, analysis, and hypotheses. *A) Visual description of the whole-report*
visual working memory task (adapted from Adam et al., 2018). A cue indicated whether participants were
required to attend to the left or right side of the screen. Both sides contained the same number of colored
squares, but in different locations. In experiment 1, set size was manipulated to consist of one, three or six
items on each side. In experiment 2, the set size was kept constant at six items. After a 1300 ms retention
period, participants were prompted to indicate the color of each item of the attended side. There were nine
discrete colors to choose from. B) Spectral parameterization method. From a power spectrum, the
aperiodic components, the exponent and offset, were quantified separately from the periodic components,
relative power. Total power was captured as bandpower. C) Three scenarios illustrating how task-induced
changes in the power spectrum appear as theta bandpower changes. The theta frequency range (4-7 Hz)
is highlighted in red. Scenario 1 (left) shows the assumed model whereby task-related changes in theta
band activity are driven by changes in narrowband, oscillatory theta activity manifesting above a stable
aperiodic signal. Scenario 2 (middle) and 3 (right) describe situations in which a reduction in total theta
power on the task (green line) from baseline (orange line) is observed, all driven by changes in aperiodic
dynamics whether or not detectable theta oscillations are present.

Figure 2: Grouping participants based on the presence of oscillatory peaks. *Here we show for both experiments (rows), and oscillatory bands (columns) the average (thick lines), and individual participant PSDs (thin lines) averaged over all trials for those who did exhibit oscillatory power (magenta), and those who did not (teal). For theta, we also observe group differences in both aperiodic exponent and offset, in which participants without theta oscillations typically have flatter spectra. These results apply to both experiments. * = p < 0.05*

624
Figure 3: Frontal midline aperiodic activity decreases during the retention period from baseline,
whereas theta oscillations increase in power. *A-F) Experiment 1 results. A) Shows the power spectra*
from baseline (black), and the three different set sizes. B) Theta peak height increases from baseline,
whereas abundance (D) and bandpower (C) decreases. Aperiodic exponent (E) and offset (F) show no
significant differences between set sizes, but are significantly decreased from baseline. G-L) Experiment 2
results. F) Shows the power spectra for the baseline (black), and poor (red) versus good (green) WM
performance. No significant differences were observed in theta peak height (H), nor in abundance (J),
*whereas both bandpower (I), aperiodic exponent (K), and offset (L) decreased from baseline. * = p < 0.05,*
*** = p < 0.005, ***= p < 0.001, n.s. = not significant.*
634
635
636

637
638 *Figure 4: Occipitoparietal aperiodic activity decreases with increase in load, but not performance,*
639 *whereas alpha oscillatory power follows classic findings. A-F) Experiment 1 results. A) Shows the*
640 *power spectra from baseline (black), and three set sizes. B) Occipitoparietal alpha peak height shows a*
641 *significant decrease with an increase in WM load, but no lateralized effect is observed. Similar patterns*
642 *are observed for total alpha bandpower (C), whereas abundance increases with set size (D). Aperiodic*
643 *exponent (E) shows no lateralized effect, but there was a decrease with an increase in WM load. Similar*
644 *patterns were observed for the aperiodic offset (F). G-L) Experiment 2 results. F) Shows the power*
645 *spectra for the baseline (black), and poor (red) versus good (green) WM performance. A decrease in*
646 *alpha peak height was observed for the good condition compared to the poor condition, but again no*
647 *lateralization effect was found (H). Bandpower and (I) abundance (J) did not show any significant*
648 *differences. The aperiodic exponent (K) and offset (L) were both significantly reduced from baseline, and a*
649 *lateralization effect was observed, in which activity in the contralateral hemisphere was more suppressed*
650 *than the ipsilateral hemisphere. * = p < 0.05, ** = p < 0.005, ***= p < 0.001, n.s. = not significant.*
651

652 **5. References**

653 Adam, K. C. S., Robison, M. K., & Vogel, E. K. (2018). Contralateral Delay Activity Tracks Fluctuations in
654   Working Memory Performance. *Journal of Cognitive Neuroscience*, *30*(9), 1229–1240.
655   https://doi.org/10.1162/jocn_a_01233
656 Adam, K. C. S., Vogel, E. K., & Awh, E. (2020). Multivariate analysis reveals a generalizable human
657   electrophysiological signature of working memory load. *Psychophysiology*, *57*(12), e13691.
658   https://doi.org/10.1111/psyp.13691
659 Akbarian, F., Rossi, C., Costers, L., D'hooghe, M. B., D'haeseleer, M., Nagels, G., & Schependom, J. V.
660   (2023). *Stimulus-related modulation in the 1/f spectral slope suggests an impaired inhibition of*
661   *irrelevant information in people with multiple sclerosis* (p. 2023.12.28.573572). bioRxiv.
662   https://doi.org/10.1101/2023.12.28.573572
663 Baddeley, A. (1992). Working Memory. *Science*, *255*(5044), 556.
664 Bae, G.-Y., & Luck, S. J. (2018). Dissociable Decoding of Spatial Attention and Working Memory from
665   EEG Oscillations and Sustained Potentials. *Journal of Neuroscience*, *38*(2), 409–422.
666   https://doi.org/10.1523/JNEUROSCI.2860-17.2017
667 Bailey, N. W., Freedman, G., Raj, K., Spierings, K. N., Piccoli, L. R., Sullivan, C. M., Chung, S. W., Hill, A.
668   T., Rogasch, N. C., & Fitzgerald, P. B. (2020). Mindfulness Meditators Show Enhanced Accuracy

669   and Different Neural Activity During Working Memory. *Mindfulness*, *11*(7), 1762–1781.
670    https://doi.org/10.1007/s12671-020-01393-8
671 Berger, B., & Sauseng, P. (2022). Brain rhythms: How control gets into working memory. *Current Biology*,
672    *32*(10), R479–R481. https://doi.org/10.1016/j.cub.2022.04.036
673 Brake, N., Duc, F., Rokos, A., Arseneau, F., Shahiri, S., Khadra, A., & Plourde, G. (2024). A
674    neurophysiological basis for aperiodic EEG and the background spectral trend. *Nature*
675    *Communications*, *15*(1), 1514. https://doi.org/10.1038/s41467-024-45922-8
676 Cole, S., Donoghue, T., Gao, R., & Voytek, B. (2019). NeuroDSP: A package for neural digital signal
677    processing. *Journal of Open Source Software*, *4*(36), 1272. https://doi.org/10.21105/joss.01272
678 Colombo, M. A., Napolitani, M., Boly, M., Gosseries, O., Casarotto, S., Rosanova, M., Brichant, J.-F.,
679    Boveroux, P., Rex, S., Laureys, S., Massimini, M., Chieregato, A., & Sarasso, S. (2019). The
680    spectral exponent of the resting EEG indexes the presence of consciousness during
681    unresponsiveness induced by propofol, xenon, and ketamine. *NeuroImage*, *189*, 631–644.
682    https://doi.org/10.1016/j.neuroimage.2019.01.024
683 Compte, A., Brunel, N., Goldman-Rakic, P. S., & Wang, X.-J. (2000). Synaptic Mechanisms and Network
684    Dynamics Underlying Spatial Working Memory in a Cortical Network Model. *Cerebral Cortex*,
685    *10*(9), 910–923. https://doi.org/10.1093/cercor/10.9.910
686 Cowan, N. (2010). The Magical Mystery Four: How is Working Memory Capacity Limited, and Why?
687    *Current Directions in Psychological Science*, *19*(1), 51–57.
688    https://doi.org/10.1177/0963721409359277
689 D'Esposito, M., & Postle, B. R. (2015). The Cognitive Neuroscience of Working Memory. *Annual Review*
690    *of Psychology*, *66*(1), 115–142. https://doi.org/10.1146/annurev-psych-010814-015031
691 Donoghue, T. (2024). *A systematic review of aperiodic neural activity in clinical investigations*.
692    https://doi.org/10.1101/2024.10.14.24314925
693 Donoghue, T., Dominguez, J., & Voytek, B. (2020). Electrophysiological Frequency Band Ratio Measures
694    Conflate Periodic and Aperiodic Neural Activity. *eNeuro*, *7*(6).
695    https://doi.org/10.1523/ENEURO.0192-20.2020
696 Donoghue, T., Haller, M., Peterson, E. J., Varma, P., Sebastian, P., Gao, R., Noto, T., Lara, A. H., Wallis,
697    J. D., Knight, R. T., Shestyuk, A., & Voytek, B. (2020). Parameterizing neural power spectra into
698    periodic and aperiodic components. *Nature Neuroscience*, *23*(12), 1655–1665.
699    https://doi.org/10.1038/s41593-020-00744-x
700 Foster, J. J., Sutterer, D. W., Serences, J. T., Vogel, E. K., & Awh, E. (2016). The topography of alpha-
701    band activity tracks the content of spatial working memory. *Journal of Neurophysiology*, *115*(1),
702    168–177. https://doi.org/10.1152/jn.00860.2015
703 Frelih, T., Matković, A., Mlinarič, T., Bon, J., & Repovš, G. (2024). *Modulation of aperiodic EEG activity*
704    *provides sensitive index of cognitive state changes during working memory task* (p.
705    2024.05.13.593835). bioRxiv. https://doi.org/10.1101/2024.05.13.593835
706 Fukuda, K., Kang, M.-S., & Woodman, G. F. (2016). Distinct neural mechanisms for spatially lateralized
707    and spatially global visual working memory representations. *Journal of Neurophysiology*, *116*(4),
708    1715–1727. https://doi.org/10.1152/jn.00991.2015
709 Fukuda, K., Mance, I., & Vogel, E. K. (2015). ⌣ Power Modulation and Event-Related Slow Wave Provide.
710    *The Journal of Neuroscience*, *35*(41), 14009–14016.
711 Gao, R., Peterson, E. J., & Voytek, B. (2017). Inferring synaptic excitation/inhibition balance from field
712    potentials. *NeuroImage*, *158*, 70–78. https://doi.org/10.1016/j.neuroimage.2017.06.078
713 Gevins, A., & Smith, M. E. (2000). Neurophysiological Measures of Working Memory and Individual
714    Differences in Cognitive Ability and Cognitive Style. *Cerebral Cortex*, *10*(9), 829–839.
715    https://doi.org/10.1093/cercor/10.9.829
716 Grover, S., Fayzullina, R., Bullard, B. M., Levina, V., & Reinhart, R. M. G. (2023). A meta-analysis
717    suggests that tACS improves cognition in healthy, aging, and psychiatric populations. *Science*
718    *Translational Medicine*, *15*(697), eabo2044. https://doi.org/10.1126/scitranslmed.abo2044
719 Gyurkovics, M., Clements, G. M., Low, K. A., Fabiani, M., & Gratton, G. (2022). Stimulus-Induced
720    Changes in 1/f-like Background Activity in EEG. *Journal of Neuroscience*, *42*(37), 7144–7151.
721    https://doi.org/10.1523/JNEUROSCI.0414-22.2022
722 Halgren, E., Boujon, C., Clarke, J., Wang, C., & Chauvel, P. (2002). Rapid Distributed Fronto-parieto-
723    occipital Processing Stages During Working Memory in Humans. *Cerebral Cortex*, *12*(7), 710–
724    728. https://doi.org/10.1093/cercor/12.7.710

He, B. J. (2014). Scale-free brain activity: Past, present, and future. *Trends in Cognitive Sciences*, *18*(9), 480–487. https://doi.org/10.1016/j.tics.2014.04.003

Helson, P., Lundqvist, D., Svenningsson, P., Vinding, M. C., & Kumar, A. (2023). Cortex-wide topography of 1/f-exponent in Parkinson's disease. *Npj Parkinson's Disease*, *9*(1), Article 1. https://doi.org/10.1038/s41531-023-00553-6

Herweg, N. A., Solomon, E. A., & Kahana, M. J. (2020). Theta Oscillations in Human Memory. *Trends in Cognitive Sciences*, *24*(3), 208–227. https://doi.org/10.1016/j.tics.2019.12.006

Höhn, C., Hahn, M. A., Lendner, J. D., & Hoedlmoser, K. (2024). Spectral Slope and Lempel–Ziv Complexity as Robust Markers of Brain States during Sleep and Wakefulness. *eNeuro*, *11*(3). https://doi.org/10.1523/ENEURO.0259-23.2024

Hsieh, L.-T., & Ranganath, C. (2014). Frontal midline theta oscillations during working memory maintenance and episodic encoding and retrieval. *NeuroImage*, *85*, 721–729. https://doi.org/10.1016/j.neuroimage.2013.08.003

Hunter, J. D. (2007). Matplotlib: A 2D graphics environment. *Computing in Science \& Engineering*, *9*(3), 90–95.

Jensen, O., & Tesche, C. D. (2002). Frontal theta activity in humans increases with memory load in a working memory task: Frontal theta increases with memory load. *European Journal of Neuroscience*, *15*(8), 1395–1399. https://doi.org/10.1046/j.1460-9568.2002.01975.x

Jones, S. R. (2016). When brain rhythms aren't 'rhythmic': Implication for their mechanisms and meaning. *Current Opinion in Neurobiology*, *40*, 72–80. https://doi.org/10.1016/j.conb.2016.06.010

Kałamała, P., Gyurkovics, M., Bowie, D. C., Clements, G. M., Low, K. A., Dolcos, F., Fabiani, M., & Gratton, G. (2024). Event-induced modulation of aperiodic background EEG: Attention-dependent and age-related shifts in E:I balance, and their consequences for behavior. *Imaging Neuroscience*, *2*, 1–18. https://doi.org/10.1162/imag_a_00054

Karalunas, S. L., Ostlund, B. D., Alperin, B. R., Figuracion, M., Gustafsson, H. C., Deming, E. M., Foti, D., Antovich, D., Dude, J., Nigg, J., & Sullivan, E. (2022). Electroencephalogram aperiodic power spectral slope can be reliably measured and predicts ADHD risk in early development. *Developmental Psychobiology*, *64*(3), e22228. https://doi.org/10.1002/dev.22228

Klimesch, W. (1999). EEG alpha and theta oscillations reflect cognitive and memory performance: A review and analysis. *Brain Research Reviews*, *29*(2–3), 169–195. https://doi.org/10.1016/S0165-0173(98)00056-3

Lara, A. H., & Wallis, J. D. (2015). The Role of Prefrontal Cortex in Working Memory: A Mini Review. *Frontiers in Systems Neuroscience*, *9*, 173. https://doi.org/10.3389/fnsys.2015.00173

Lendner, J. D., Harler, U., Daume, J., Engel, A. K., Zöllner, C., Schneider, T. R., & Fischer, M. (2023). Oscillatory and aperiodic neuronal activity in working memory following anesthesia. *Clinical Neurophysiology*, *150*, 79–88. https://doi.org/10.1016/j.clinph.2023.03.005

Lendner, J. D., Helfrich, R. F., Mander, B. A., Romundstad, L., Lin, J. J., Walker, M. P., Larsson, P. G., & Knight, R. T. (2020). An electrophysiological marker of arousal level in humans. *eLife*, *9*, e55092. https://doi.org/10.7554/eLife.55092

Lisman, J. E., & Jensen, O. (2013). The Theta-Gamma Neural Code. *Neuron*, *77*(6), 1002–1016. https://doi.org/10.1016/j.neuron.2013.03.007

Lu, L., Gao, Z., Wei, Z., & Yi, M. (2023). Working memory depends on the excitatory–inhibitory balance in neuron–astrocyte network. *Chaos: An Interdisciplinary Journal of Nonlinear Science*, *33*(1), 013127. https://doi.org/10.1063/5.0126890

McKeon, S. D., Perica, M. I., Parr, A. C., Calabro, F. J., Foran, W., Hetherington, H., Moon, C.-H., & Luna, B. (2024). Aperiodic EEG and 7T MRSI evidence for maturation of E/I balance supporting the development of working memory through adolescence. *Developmental Cognitive Neuroscience*, *66*, 101373. https://doi.org/10.1016/j.dcn.2024.101373

Medendorp, W. P., Kramer, G. F. I., Jensen, O., Oostenveld, R., Schoffelen, J.-M., & Fries, P. (2007). Oscillatory activity in human parietal and occipital cortex shows hemispheric lateralization and memory effects in a delayed double-step saccade task. *Cerebral Cortex (New York, N.Y.: 1991)*, *17*(10), 2364–2374. https://doi.org/10.1093/cercor/bhl145

Miller, G. A. (1956). The Magical Number Seven, Plus or Minus Two Some Limits on Our Capacity for Processing Information. *Psychological Review*, *63*(2), 81.

Mitchell, D. J., McNaughton, N., Flanagan, D., & Kirk, I. J. (2008). Frontal-midline theta from the perspective of hippocampal "theta." *Progress in Neurobiology*, *86*(3), 156–185. https://doi.org/10.1016/j.pneurobio.2008.09.005

Oberauer, K., Farrell, S., Jarrold, C., & Lewandowsky, S. (2016). What limits working memory capacity? *Psychological Bulletin*, *142*(7), 758–799. https://doi.org/10.1037/bul0000046

Palva, S., & Palva, J. M. (2007). New vistas for α-frequency band oscillations. *Trends in Neurosciences*, *30*(4), 150–158. https://doi.org/10.1016/j.tins.2007.02.001

Pavlov, Y. G., & Kotchoubey, B. (2022). Oscillatory brain activity and maintenance of verbal and visual working memory: A systematic review. *Psychophysiology*, *59*(5), e13735. https://doi.org/10.1111/psyp.13735

Podvalny, E., Noy, N., Harel, M., Bickel, S., Chechik, G., Schroeder, C. E., Mehta, A. D., Tsodyks, M., & Malach, R. (2015). A unifying principle underlying the extracellular field potential spectral responses in the human cortex. *Journal of Neurophysiology*, *114*(1), 505–519. https://doi.org/10.1152/jn.00943.2014

Sauseng, P., Klimesch, W., Heise, K. F., Gruber, W. R., Holz, E., Karim, A. A., Glennon, M., Gerloff, C., Birbaumer, N., & Hummel, F. C. (2009). Brain Oscillatory Substrates of Visual Short-Term Memory Capacity. *Current Biology*, *19*(21), 1846–1852. https://doi.org/10.1016/j.cub.2009.08.062

Smith, S. E., Kosik, E. L., van Engen, Q., Kohn, J., Hill, A. T., Zomorrodi, R., Blumberger, D. M., Daskalakis, Z. J., Hadas, I., & Voytek, B. (2023). Magnetic seizure therapy and electroconvulsive therapy increase aperiodic activity. *Translational Psychiatry*, *13*(1), Article 1. https://doi.org/10.1038/s41398-023-02631-y

Smith, S. E., Ma, V., Gonzalez, C., Chapman, A., Printz, D., Voytek, B., & Soltani, M. (2023). Clinical EEG slowing induced by electroconvulsive therapy is better described by increased frontal aperiodic activity. *Translational Psychiatry*, *13*(1), Article 1. https://doi.org/10.1038/s41398-023-02634-9

Soltani Zangbar, H., Ghadiri, T., Seyedi Vafaee, M., Ebrahimi Kalan, A., Fallahi, S., Ghorbani, M., & Shahabi, P. (2020). Theta Oscillations Through Hippocampal/Prefrontal Pathway: Importance in Cognitive Performances. *Brain Connectivity*, *10*(4), 157–169. https://doi.org/10.1089/brain.2019.0733

Vallat, R. (2018). Pingouin: Statistics in Python. *Journal of Open Source Software*, *3*(31), 1026. https://doi.org/10.21105/joss.01026

Van Bree, S., Levenstein, D., Krause, M., Voytek, B., & Gao, R. (2024). *Decoupling Measurements and Processes: On the Epiphenomenon Debate Surrounding Brain Oscillations in Field Potentials*. https://doi.org/10.31234/osf.io/knjfw

van Bueren, N. E. R. van, Ven, S. H. G. van der, Hochman, S., Sella, F., & Kadosh, R. C. (2023). Human neuronal excitation/inhibition balance explains and predicts neurostimulation induced learning benefits. *PLOS Biology*, *21*(8), e3002193. https://doi.org/10.1371/journal.pbio.3002193

van der Werf, J., Jensen, O., Fries, P., & Medendorp, W. P. (2008). Gamma-Band Activity in Human Posterior Parietal Cortex Encodes the Motor Goal during Delayed Prosaccades and Antisaccades. *Journal of Neuroscience*, *28*(34), 8397–8405. https://doi.org/10.1523/JNEUROSCI.0630-08.2008

van Dijk, H., van der Werf, J., Mazaheri, A., Medendorp, W. P., & Jensen, O. (2010). Modulations in oscillatory activity with amplitude asymmetry can produce cognitively relevant event-related responses. *Proceedings of the National Academy of Sciences*, *107*(2), 900–905. https://doi.org/10.1073/pnas.0908821107

Vanderwolf, C. H. (1969). Hippocampal electrical activity and voluntary movement in the rat. *Electroencephalography and Clinical Neurophysiology*, *26*(4), 407–418. https://doi.org/10.1016/0013-4694(69)90092-3

Virtue-Griffiths, S., Fornito, A., Thompson, S., Biabani, M., Tiego, J., Thapa, T., & Rogasch, N. C. (2022). *Task-related changes in aperiodic activity are related to visual working memory capacity independent of event-related potentials and alpha oscillations* [Preprint]. Neuroscience. https://doi.org/10.1101/2022.01.18.476852

Vogel, E. K., & Machizawa, M. G. (2004). Neural activity predicts individual differences in visual working memory capacity. *Nature*, *428*(6984), 748–751. https://doi.org/10.1038/nature02447

Voytek, B., Kramer, M. A., Case, J., Lepage, K. Q., Tempesta, Z. R., Knight, R. T., & Gazzaley, A. (2015). Age-Related Changes in 1/f Neural Electrophysiological Noise. *Journal of Neuroscience*, *35*(38), 13257–13265. https://doi.org/10.1523/JNEUROSCI.2332-14.2015

835   Wang, X.-J. (1999). Synaptic Basis of Cortical Persistent Activity: The Importance of NMDA Receptors to
836   Working Memory. *Journal of Neuroscience*, *19*(21), 9587–9603.
837   https://doi.org/10.1523/JNEUROSCI.19-21-09587.1999
838   Waschke, L., Donoghue, T., Fiedler, L., Smith, S., Garrett, D. D., Voytek, B., & Obleser, J. (2021).
839   Modality-specific tracking of attention and sensory statistics in the human electrophysiological
840   spectral exponent. *eLife*, *10*, e70068. https://doi.org/10.7554/eLife.70068
841   Waskom, M. L. (2021). *Journal of Open Source Software*, *6*(60), 3021. https://doi.org/10.21105/joss.03021
842   Wilson, L. E., da Silva Castanheira, J., & Baillet, S. (2022). Time-resolved parameterization of aperiodic
843   and periodic brain activity. *eLife*, *11*, e77348. https://doi.org/10.7554/eLife.77348
844