1   **Title**

2   **Reanalysis of metabolomics data reveals that microbiota transplant therapy modulates**

3   **important fecal and plasma metabolite profiles in children with autism spectrum disorders**

Khemlal Nirmalkar[1,# *], Fatir Qureshi[2, 3,!,#], Dae-Wook Kang[4,], Christina K. Flynn[1,5], Juergen Hahn[2,3,6], James B. Adams[1,5], Rosa Krajmalnik-Brown[1,7 *]

[1]Biodesign Center for Health Through Microbiomes, Arizona State University, Tempe, AZ, USA

[2]Department of Biomedical Engineering, Rensselaer Polytechnic Institute, Troy, NY 12180, USA

[3]Center for Biotechnology and Interdisciplinary Studies, Rensselaer Polytechnic Institute, Troy, NY 12180, USA

[4]Department of Civil and Environmental Engineering, The University of Toledo, Toledo, OH 43606, USA

[5]School for Engineering of Matter, Transport, and Energy, Arizona State University, Tempe, AZ 85287, USA

[6]Department of Chemical and Biological Engineering, Rensselaer Polytechnic Institute, Troy, NY 12180, USA

[7]School of Sustainable Engineering and the Built Environment, Arizona State University, Tempe, AZ 85281, USA

# These authors contributed equally

! Current address: Whitehead Institute for Biomedical Research, Cambridge, MA, 02142

*Correspondence: Rosa Krajmalnik-Brown Dr.Rosy@asu.edu and Khemlal Nirmalkar khem@asu.edu

Keywords: Autism Spectrum Disorder (ASD), metabolites, metabolomics, Fecal Microbiota transplant (FMT), Microbiota transplant therapy (MTT)

Running Title: Reanalysis of metabolomics data for microbiota transplant therapy in Autism

Abstract:

While Autism Spectrum Disorder (ASD) is diagnosed through behavioral symptoms and psychometric evaluations, it has also been associated with distinct metabolomic patterns. A previous clinical trial of Microbiome Transplant Therapy (MTT) in children with ASD and gastrointestinal (GI) issues revealed significant differences in plasma metabolomics between children with ASD and their typically developing (TD) counterparts, which diminished after MTT. The objective of this study was to reanalyze the plasma and fecal samples using updated metabolomics libraries at Metabolon, and applying a comprehensive panel of statistical methods. This approach aimed to provide deeper insights into ASD-related metabolic differences and the impact of MTT. The reanalysis identified more statistically significant metabolites and highlighted specific metabolites whose relative peak intensities differed between the ASD and TD groups, as well as metabolites with significant changes in peak intensity post-MTT. Several plasma metabolites, including sarcosine, iminodiacetate, caproate, and caprylate, initially showed significant differences between the ASD and TD groups but shifted to resemble TD levels after MTT treatment. In fecal samples, *p*-cresol sulfate and sphingolipids emerged as metabolites with altered intensities following MTT treatment. Multivariate Fisher's Discriminant Analysis (FDA) with leave-one-out cross-validation revealed that a set of metabolites—including p-cresol sulfate, hydroxyproline, and caprylate—could robustly classify the ASD and TD cohorts pre-treatment. However, after treatment, the same FDA model could not distinguish the two groups, as the FDA scores became similar to those of the TD cohort.

Our findings enhance the understanding of ASD-associated metabolic changes and highlight the potential of MTT to influence these profiles. This underscores the importance of reanalysis using updated databases and robust statistical methods for comparative analysis. Further studies with larger cohorts and placebo-controlled trials are necessary to validate these results and explore the biochemical pathways involved, paving the way for personalized treatment approaches.

Introduction

In the past decade, the prevalence of autism spectrum disorder (ASD) in the United States has significantly increased, with 1 in 31 children now diagnosed (CDC, 2025). Individuals with ASD experience social communication and behavioral impairments (Adams et al., 2011; Lord et al., 2018), and often have persistent gastrointestinal (GI) issues, such as constipation and diarrhea (Adams et al., 2011; Hung et al., 2023). These co-occurring conditions correlate with increased ASD behavioral severity (Adams et al., 2011), prompting research intersecting their relationship with the gut-brain axis and gut microbiome.. Studies reveal significant differences in gut microbiome compositions between ASD individuals and typically developing (TD) children, including higher levels of *Clostridia* species and lower levels of beneficial microbes like *Bifidobacterium* (Kang et al., 2019; Morton et al., 2023).

The complex interaction among microbes, influenced by their metabolites, has been hypothesized to affect GI and behavioral symptoms in ASD (Adams et al., 2011; Krajmalnik-Brown et al., 2014; Kang et al., 2020; Needham et al., 2021; Morton et al., 2023; Hung et al., 2023; Zheng et al., 2023). Differences in metabolite profiles in fecal, urine, and blood samples have been reported in ASD individuals compared to TD peers (Gabriele et al., 2014; Kang et al., 2018; 2020; Needham et al., 2021; Zheng et al., 2021). For example, elevated levels of fecal short-chain fatty acids and blood glutamate, along with increased blood gamma-aminobutyric acid (GABA), have been observed in people with ASD (Shinohe et al., 2006; Kang et al., 2018; El-Ansary et al., 2014; Khalifa et al., 2019). GABA and glutamate, which are critical for cognition and mood regulation, show an altered levels in ASD individuals, suggesting an underlying neuroinflammation (El-Ansary et al., 2014). Additionally, other gut-derived metabolites, such as p-cresol and p-cresol sulfate, are found at higher levels in ASD individuals' urine, feces, or serum (Flynn et al., 2025; Kang et al., 2018; De Angelis et al., 2013; Gabriele et al., 2014).

Given the microbiome-gut-ASD interconnection, Microbiota Transplant Therapy (MTT) has been explored for its potential benefits in ASD (Kang et al., 2017). Microbiota Transplant Therapy, (redefined from microbiota transfer therapy, Takyi et al. 2025) has shown promise in addressing GI issues and reducing behavioral severity in ASD children. An open-label study by

Kang et al. reported an 80% reduction in GI symptoms and a 24% decrease in core ASD

symptoms post-MTT, with sustained improvements over two years (Kang et al., 2017, 2019).

Increases in beneficial microbes such as *Bifidobacterium*, *Prevotella*, and *Desulfovibrio* were

also noted (Kang et al., 2017, 2019).

Building on these findings, untargeted metabolomics of fecal and plasma samples from the same

individuals in the MTT study identified several plasma metabolites significantly lower in the

ASD group at baseline, including nicotinamide riboside, IMP, iminodiacetate, methylsuccinate,

galactonate, valylglycine, sarcosine, and leucylglycine, while caprylate and heptanoate were

higher (Kang et al., 2020). Though no fecal metabolite showed significant baseline differences,

p-cresol sulfate levels were non-significantly higher in ASD children and decreased after MTT

(Kang et al., 2020). Multivariate analysis highlighted significant metabolite changes post-MTT,

with improvements in sarcosine and inosine 5'-monophosphate in plasma (Adams et al., 2019)

and a model of indole, imidazole, propionate, adenosine, and hydroxyproline achieving 96%

accuracy in distinguishing ASD from TD children (Qureshi et al., 2020).

In this study, we reanalyzed the metabolomics data collected for the Kang et al. 2017 and 2019

studies. We extend upon the findings reported by Kang et al. (2020) by expanding the scope of

metabolomics data analyzed through Metabolon's updated metabolomics libraries and robust

statistical methods. Utilizing more up-to-date algorithms and libraries for metabolite detection,

we offer a comprehensive comparison of metabolite profiles, encompassing nucleotides,

neurotransmitters, amino acids, xenobiotics, and lipids derived from human plasma and fecal

samples collected from a carefully matched cohort of individuals with ASD and typically

developing children. Previously, data was analyzed using univariate tests and principal-

component analysis (PCA) and correlation analysis for selective metabolites. In this reanalysis,

we not only replicated the prior analysis techniques, but we also explored other methods such as

contrasting the area under the receiver operating characteristic curve (AUROC) between groups,

random forest, and Fisher discriminant analysis (FDA) to revalidate the findings (Fig. 1). The

results reveal significant differences in metabolite levels between TD and ASD baseline groups

in plasma and feces, which were previously unidentified, and important changes shifting the

ASD metabolon towards TD after MTT.

Methods

Study design, sample collection and data processing

117
118
This study extends the work of Kang et al. (2020) by performing a more thorough reanalysis of
plasma and fecal metabolomics data collected before and after MTT in children with ASD. The
original protocol was approved by the U.S. Food and Drug Administration (FDA)
(investigational new drug number 15886) and the Institutional Review Board of Arizona State
University (ASU IRB protocols #00001053 and #00004890). The trial was registered at
ClinicalTrials.gov (NCT02504554), as described in Kang et al. (2017, 2020). Study participants,
consent, MTT, and sample collections were described in detail in our previous studies (Kang et
al., 2017; 2020). In brief, 20 typically developing (TD) children and eighteen children with ASD
were recruited, in the age range of 7-16 years old. Children with ASD participated in an open-
label clinical trial including vancomycin for 2 weeks, one-day bowel cleanse with MoviPrep,
followed by 7-8 weeks of MTT and Prilosec (a stomach acid suppressant), and a follow-up at 8
weeks post-treatment (i.e., 18 weeks from Day 0) (Kang et al., 2017).

For metabolomic analysis, we collected plasma samples at ASD baseline, week 3 (after
the vancomycin and bowel cleanse and just before the first dose of microbiota), and week 10 (the
end of MTT treatment) (Fig. 1). We had one missing plasma sample in week 3. Fecal samples
were collected at baseline, week 3 (right after an initial dose of microbiota), week 10 and week
18 (8 weeks after the end of MTT). For TD children, we collected plasma and feces only at
baseline as they were not treated. All samples were frozen immediately after collection, shipped
in dry ice to ASU, and stored at -80°C. Demographic, medical data and diet history was
recorded, as described by Kang *et al*., 2017, 2020. Aliquoted frozen samples were analyzed by
Metabolon Inc. (Durham, NC, USA) using ultrahigh-performance liquid chromatography-
tandem mass spectroscopy (UHPLC-MS/MS) instruments for untargeted metabolomics.

This reanalysis leverages upgrades in peak identification algorithms, software, and
libraries available to Metabolon Inc. (https://www.metabolon.com/). In feces, a total of 1,068
metabolites were detected across all samples. Of these, 825 metabolites (77.3%) were chemically
identified and assigned a metabolite name, while the remaining 243 features (22.7%) were
unnamed. Similarly, in the plasma samples, a total of 1,001 metabolites were detected, with 844
(84.3%) chemically identified and 157 (15.7%) remaining unnamed.

Their proprietary software integrates the mass spectrometry data with retention time alignment,

isotope pattern analysis, and database matching to enhance metabolite annotation accuracy. In

total, 5,400 compounds were detectable via untargeted metabolomics, with available annotations.

Raw data underwent imputation and z-score normalization according to the methods described

by Kang et al., 2020. Using these extended approaches new data were generated, resulting in

modified abundances of metabolites (peak area/intensity) and the identification of additional

metabolites within the samples. To focus the analysis on identified compounds for biological

interpretation, metabolites lacking chemical annotations (e.g., unidentified compounds labeled as

'X-###') were excluded from downstream analysis.

It is important to note that, we recorded xenobiotics ~16% in plasma and ~18% in feces which

was 1% higher than our previous study (Kang et al., 2020). The slight increase likely reflects

updates in Metabolon's annotation libraries during reanalysis. Importantly, Metabolon's

xenobiotic category encompasses a broad range of compounds, including not only exogenous

chemicals (e.g., drugs, pollutants) but also microbial co-metabolites, dietary constituents, and

plant-derived compounds. For instance, gut-derived metabolites such as p-cresol and 4-

ethylphenyl sulfate are classified by metabolon as xenobiotics due to their chemical features and

involvement in pathways like benzoate metabolism. Therefore, the relatively high proportion of

xenobiotics observed here may reflect the microbiome-influenced metabolic landscape of

individuals with autism, rather than an overrepresentation of exogenous exposures.

Univariate analysis

Initially, univariate hypothesis testing was employed to compare metabolites between groups.

Depending on the data distribution and type of relationship of the two groups, the most

appropriate statistical test was selected. Mann-Whitney was utilized for comparing non-

parametric data with similar distributions between the TD and ASD samples, while a t-test was

used in alternative cases. When comparing ASD groups at various time points a Wilcoxon sign-

rank test was used for non-parametric distributions and a paired t-test was used for parametric

distributions. False discovery rates (FDR) were determined using the leave-one-out (LOO)

approach. This technique is well-suited for high-dimensional data that has an underlying

correlational structure, as is common in metabolic studies (Arici et al., 2025). This method

applies the same hypothesis testing approach to every combination of a dataset with one sample

excluded. Then, the proportion of significant findings is determined to evaluate the likelihood of

false discovery. Subsequently, findings with $p < 0.05$ and FDR <0.05 were considered

statistically significant.

Area Under the Receiver Operating Characteristic Curve (AUROC)

The area under the receiver operating characteristic curve (AUROC) was used to assess the

degree to which metabolite measurements were discernible between pre-treatment and post-

treatment time points. For a binary classification task, the AUROC is determined by plotting the

true positive rate (TPR) against the false positive rate (FPR) of a model across varying threshold

values. The area under the TPR/FNR plot corresponds to the AUROC and can serve as a metric

to quantify the degree to which different groups are separable. The AUROC value ranges from 0

to 1, where a value of 0.5 indicates no discrimination, a value of 1 represents perfect

discrimination, and a value close to 0 suggests that the model is performing worse than random,

potentially predicting the opposite of the true outcome.

The AUROC was calculated by treating the measurements of each metabolite individually as the

classifier input, which was subject to a threshold value. AUROC was determined for each fecal

and plasma metabolite, with baseline measurements being compared against all other time

points. The measurements collected from the ASD group were also compared against the TD

measurements, both in their entirety and on a time-point basis. By assessing all the AUROC

values generated for each metabolite between baseline, week3, and week10 vs TD across, the

average fecal and plasma AUROC values for each group were determined.

Fisher's Discriminant Analysis (FDA)

FDA is a dimensionality reduction technique that aims to find a transformation of the original

data that accentuates the differences between distinct groups while reducing the variations within

each group. This method projects multidimensional data into a lower dimensional space in such a

way as that dissimilar groups are maximally separated, while members of the same group remain

closely grouped together. By doing so, FDA facilitates the creation of more discriminative

representations (Huberty, 1975).

Metabolites that had been deemed to be statistically significant in both plasma and feces

were used to develop an FDA model for the purpose of classification between ASD and TD

cohorts. To mitigate the risk of overfitting, feature selection was limited to five variables, as

literature suggests this approach optimally suits small sample sizes with similar data structure

properties in ASD prediction. This decision is also supported by the success of prior fecal and

plasma model panels achieving the upper bounds of their predictive accuracy with this number of

features (Howsmon et al., 2017; Vargason et al., 2019; Qureshi et al., 2020). An exhaustive

search of all potential FDA model combinations of five metabolites was determined, using only

the ASD baseline and TD groups as training data. For all possible combinations of five

metabolites, the AUROC was evaluated to determine whether to conduct further testing. Each

metabolite combination model that achieved an AUROC greater than 0.97 was subjected to a

leave-one-out cross-validation (LOO-CV) paradigm. The top-performing model, as quantified by

CV accuracy, was then used to predict the classification of the MTT week3 and week10 groups.

Random Forest Analysis

Random forest (RF) is defined as an ensemble learning method that can be used for

classification. An RF approach to classification operates by aggregating the collective

performance of multiple decision trees and using the plurality of outcomes to predict the group

label of a sample. This algorithm is constructed by first setting aside one-third of the training

data to utilize as testing data for cross-validation purposes. Subsequently, a user-specified

number of decision trees is constructed. The performance of the algorithm is then quantified

based on the classification accuracy on the randomly selected test set. A random forest model

was fit using all significant metabolomic measurements in order to evaluate the mean decrease in

accuracy for the exclusion of each variable. This metric is commonly utilized to provide a

perspective on feature importance, which can be used to more efficiently identify variables better

suited for distinguishing between labeled categories (Altman et al., 2017). Although random

forests are generally more effective with larger datasets, they can still serve as exploratory tools

for feature selection in smaller datasets, provided their limitations are acknowledged and results

are interpreted cautiously (Wouter et al., 2013)

Correlation Coefficient

The Pearson's correlation coefficient is defined by the strength and direction of the linear

relationship between two continuous variables. The Pearson's correlation coefficient was

determined between all metabolite, pathway, and taxa pairs. The significance of the correlation
coefficients was evaluated using hypothesis testing with a significance level set at α = 0.05. The
robustness of significant findings was cross validated using a LOO approach. Employing this
strategy, the significance of the correlation coefficient was determined for all possible
combinations of the input measurement with a single sample removed. If a false discovery rate
(FDR) of less than 0.05 was determined, the finding was deemed to be significant.

All univariate analysis, area under the receiver operating characteristic (AUROC), random forest
analysis and plots were made with Python (v3.8.5) in Jupyter Notebook (v6.1.4) using numpy
(v1.19.2), matplotlib (v3.3.2), seaborn (v0.11.0), pandas (v1.1.3) and scipy (v1.5.2). PCA plot
was made using MetaboAnalyst v6.0 (https://www.metaboanalyst.ca/). BioRender (Biorender,
2022, Licensed, https://biorender.com) and Inkscape (v1.1) were used to create or edit the
figures.

Results
In the previous study (Kang et al., 2020), 621 plasma and 666 fecal metabolites were identified
in the collected samples. However, through reprocessing and reanalysis of the metabolomics data
using new updated reference library, a greater number of biochemical compounds were
identified from the same samples. We identified a total of 844 plasma and 825 fecal metabolites,
and most of the newly identified metabolites in plasma and feces were lipids, amino acids, or
xenobiotics (Fig. 2).

Metabolite profiles changed after MTT in ASD children
Univariate analyses showed that certain metabolites that were initially different between ASD
and TD groups at baseline, and shifted following MTT. These metabolites relative peak intensity
changed post-treatment, and some of these intensities became similar to those in TD in plasma
(Fig. 3A, C) and fecal samples (Fig. 3B, Supplementary S1). In plasma, out of 844 metabolites,
we observed 20 metabolites whose intensity was significantly different between the ASD
baseline and TD and that also changed after MTT (Fig. 3A). We grouped these plasma
metabolites into two clusters; **1)** Cluster P-high (cluster of 14 plasma metabolites), which
includes metabolites that had significantly higher intensity (adjusted $p<0.05$) at ASD baseline vs

TD, but after MTT, the relative intensity of these metabolites decreased and became similar to TD (Fig. 3A). Figure 3A shows 12 plasma metabolites in this category, including: glycerophosphoethanolamine (GPEA), caprylate (8:0), and caproate (6:0). The other two plasma metabolites, taurine and adenosine-5'-monophosphate (AMP), were non-significantly higher ($p>0.05$) at baseline vs TD but significantly decreased after MTT (Fig. 3A). **2)** Cluster P-low (cluster of 6 plasma metabolites) which includes metabolites that had significantly lower intensity (adjusted $p<0.05$) at baseline vs TD, but after MTT, relative intensity of these metabolites significantly increased (adjusted $p<0.05$) and became similar to TD. Cluster P-low includes sarcosine, iminodiacetate, bilirubin, 2-methylserine, and indole propionate.

Similar to plasma, out of 825 fecal metabolites, we observed 22 metabolites that had significantly different intensities from TD or changed after MTT (Fig 3B, Supplementary Fig. S1). Fecal metabolites were also divided into two clusters; **1)** Cluster F-low (cluster of 3 fecal metabolites) where all metabolites had significantly lower intensity (adjusted $p<0.05$) at baseline vs. TD, and **2)** Cluster F-high (cluster of 19 fecal metabolites) where all metabolites had higher intensity (adjusted $p<0.05$) at baseline vs. TD. Cluster F-low (Fig. 3B-) shows 3 fecal metabolites pentadecanoate (15:0), myristate (14:0), and carboxyethyl-GABA whose intensity was significantly lower at baseline vs. TD, and only one metabolite pentadecanoate (15:0) whose intensity significantly increased (adjusted $p<0.05$) after week18 (Fig. 3B). In cluster F-high, 19 metabolites had significantly higher intensity (adjusted $p<0.05$) at baseline vs TD, here the intensity of 5 metabolites: p-cresol sulfate, theobromine, oxalate ethanedioate, dodecanedioate (C12), and Urolithin-A significantly decreased (adjusted $p<0.05$) at week3 vs baseline (Fig. 3B). The intensity of N-palmitoyl-sphingosine (d18:1/16:0), p-cresol sulfate, eicosanoyl-sphingosine (d20:1), hydroxyproline, theobromine, N-stearoyl-sphingosine (d18:1/18:0), N-oleoyl-sphingosine (d18:1/18:1), 3-methyl-2-oxobutyrate, dodecanedioate (C12), and D-urobilin significantly decreased after week18 compared to baseline. We did not observe any significant change for week10 vs. baseline (Fig. 3B).

Analytics and Patterns of Differential Metabolomic Composition Appear Similar between Feces and Plasma

We utilized AUROC, PCA analysis and random forest analysis to categorize the differences in metabolomic composition across groups and timepoints. AUROC was calculated for all 20

plasma and 22 fecal metabolites that were significantly different between ASD baseline and TD.

To examine how metabolites differed post-MTT, AUROC was recalculated at all time points

between the ASD and TD cohorts. The average AUROC for all 20 plasma metabolites was 0.73

comparing ASD baseline vs TD, and ASD baseline vs after week3, and week10 AUROC's were

0.70 and 0.62 respectively (Fig. 4A). When we compared the MTT group with TD, after week3

and week10, we observed that the plasma profile became more similar to TD with AUROC 0.56

for week3 vs TD and 0.59 for week10 vs TD (Fig. 4A). A similar trend was also observed for 22

fecal metabolites, which were initially observed with an average AUROC of 0.74 for ASD

baseline vs TD. After MTT, fecal metabolite profiles became more similar to the TD group, with

AUROC values of 0.58 for week3 vs TD and 0.61 for week10 vs TD (Fig. 4B). Overall, the

observed shift in AUROC values for plasma and fecal metabolites indicates that after MTT, the

metabolic composition in plasma and fecal samples from children with ASD became more

similar to their TD counterparts.

PCA analysis showed that relative intensity of the 20 significant plasma metabolites at

baseline were distinct from TD, but after MTT, became much more similar to TD (Fig. 4C).

Fecal samples exhibited greater variance in the typically developing group compared to their

ASD counterparts in the first two principal components, and this pattern was also observed in the

post treatment ASD group (Supplementary Fig. S2). Random forest was used as a

complimentary method to validate the assessment of each metabolites relative utility in

distinguishing between cohorts. Using the full list of metabolites, it was possible to achieve

accuracies greater than 0.95. The mean decrease in accuracy for the ASD status prediction model

indicated that both fecal and plasma metabolites had a similar influence on model performance

(Supplementary Fig. S3). The top 10 features contributing to mean decrease in accuracy were

split evenly between the plasma and fecal metabolites, potentially offering complementary

insights into discerning ASD and TD samples. (Supplementary Fig. S4).

MTT changed metabolites abundance in ASD towards TD

We selected four plasma metabolites that were highly significantly different at baseline and

showed changes after MTT (Fig. 5).. A lysine derivative sarcosine and iminodiacetate (IDA)

were significantly lower (adjusted $p<0.05$) at baseline vs. TD. However, after week3 (after

vancomycin) and week10 (MTT) the relative intensity was observed to significantly increase

(adjusted $p<0.05$) and became more similar to TD (Fig. 3A, 5A-B). To quantify the changes, we

calculated the AUROC, and sarcosine's intensity showed 0.82 AUROC at baseline vs TD, at

week3 (AUROC=0.73) and at week10 (AUROC=0.85) compared. baseline (Fig. 6A). Notably,

the AUROC at week3 and week10 compared to TD were 0.56 and 0.52, respectively (Fig. 6A),

which indicates that after MTT, sarcosine's relative intensity became similar to TD. Similar

findings were observed for iminodiacetate (Fig. 6B).

In contrast, we observed two medium chain fatty acids in plasma, caproate (6:0) and caprylate

(8:0), that were significantly higher at baseline vs TD, but after week3 and week10 (MTT)

significantly decreased (adjusted $p<0.05$) and their abundances became similar to TD (Fig. 5C-

D). Caproate and caprylate AUROCs were 0.78 for baseline vs TD, which showed further

differences comparing baseline to week3 (AUROC caproate=0.83, caprylate=0.79) and week10

(AUROC caproate=0.88, caprylate=0.90) vs. baseline, but the AUROC at week3 and week10 vs.

TD was between 0.51-0.65 (Fig. 6C-D), which also indicates that after MTT, the relative

intensity of caproate and caprylate became similar to TD.

In feces, phenolic compound p-cresol sulfate, theobromine, and three sphingolipids; N-

palmitoyl-sphingosine (d18:1/16:0), N-stearoyl-sphingosine (d18:1/18:0), and N-oleoyl-

sphingosine (d18:1/18:1) were significantly higher at baseline vs TD but after week3 (right after

an initial dose of microbiota), week10 (MTT) and week18 (after 8 weeks of MTT) significantly

decreased vs. baseline and became similar to TD (Fig 3B, Fig. 7A-D). AUROC calculation

showed that baseline vs. TD had 0.76 for p-cresol sulfate, after baseline vs. week3

AUROC=0.80, vs week10 AUROC=0.66, vs week18 AUROC=0.75, which indicates after MTT,

p-cresol sulfate profile changed in children with ASD (Fig. 8A). Interestingly, p-cresol sulfate

AUROC for TD vs. week3; TD vs. week10; TD vs. week18 were in the range 0.55-0.60 (Fig.

8A), indicating that after MTT, p-cresol sulfate peak relative intensity became similar to TD.

Similarly, theobromine, N-palmitoyl-sphingosine (d18:1/16:0), N-stearoyl-sphingosine

(d18:1/18:0), and N-oleoyl-sphingosine (d18:1/18:1) had higher AUROC after MTT vs. ASD

baseline and lower AUROC values after MTT vs. TD, indicating important changes from

baseline and greater similarity to TD measurements (Fig. 8B-D). Overall, AUROC validate the

findings from univariate analysis and showed the differences between the groups for plasma and

fecal metabolites.

FDA models revealed five key metabolites that notably changed after MTT and became similar
to TD

Univariate hypothesis testing identified 20 plasma metabolites and 22 fecal metabolites whose
intensities significantly differed between TD and ASD cohorts at baseline or changed after
microbiota transplant therapy (MTT) (Fig. 3). Both sets of metabolites were used to separately
construct multivariate statistical models *via* FDA using all possible combinations of 3, 4, and 5
variables to distinguish between ASD and TD cohorts. The top-performing FDA model made up
exclusively of plasma metabolites included, 2-palmitoyl-GPC (16:0), caprylate (8:0), laurate
(12:0), bilirubin, and indolepropionate achieving an AUROC of 0.99 and a leave-one-out cross-
validated (LOO-CV) accuracy of 95%. Similarly, the plasma metabolite data was used to
develop an FDA model, with significant features selected through the same univariate approach.
The optimal plasma model incorporated *p*-cresol sulfate, hydroxyproline, theobromine, N-oleoyl-
sphingosine (d18:1/18:1), and 1-methylhistidine.

Utilizing the complete set of 44 significantly differing metabolites from both plasma and
feces, all possible 5-metabolite FDA models were also constructed. Through LOO-CV, a model
consisting of *p*-cresol sulfate (feces), hydroxyproline (feces), caprylate (plasma),
indolepropionate (plasma), and bilirubin (plasma) was identified as the most robust, with a cross-
validation accuracy of 95% (Fig. 4D). This model exhibited a type I error rate of 3% and a type
II error rate of 0%. Notably, the performance of all multivariate models for classifying between
ASD and TD status significantly declined after MTT, as did the measurements taken at the two-
year follow-up. The metabolite peak intensities in the model showed significant convergence
towards the TD group, with the difference between the median of the five metabolites relative
intensity values decreasing by 88% between week 0 and week18, compared to the TD group.

Fecal p-cresol and p-cresol sulfate are positively correlated

We conducted correlation analyses to examine the relationship between certain fecal and plasma
metabolites. Fecal *p*-cresol and fecal *p*-cresol sulfate were significantly positively correlated
(p<0.001, Pearson-R=0.75) (Supplementary Fig. S5). Fecal *p*-cresol and fecal *p*-cresol sulfate
did not show a significant correlation with plasma *p*-cresol sulfate (Supplementary Fig. S6-7),
which suggests that gut permeability or other factors may affect how much *p*-cresol leaves the
gut. *p*-cresol was not detected in plasma, which is not surprising since it has very low solubility

in aqueous solutions. We performed correlation analyses with previously published taxonomic

and metabolic pathways (KEGG orthologs) (Nirmalkar et al., 2022), but did not observe

statistically significant correlations.

Discussion

A distinct metabolic signature has been consistently observed in plasma and fecal samples

collected from children with ASD (Kang et al., 2020; Needham et al., 2021). While several

underlying biochemical pathways and environmental factors contribute to these differences,

MTT not only improved GI and behavioral symptoms associated with ASD (Kang et al., 2017,

2019) but also leads to a convergence of ASD metabolite profiles towards those observed in TD

samples (Kang et al., 2020). Reanalysis of absorbance peaks by Metabolon Inc. using a more

comprehensive database (see raw data in supplementary file 2), and improvement in our data

analysis pipeline allowed for an increase in the number of metabolites detected in plasma and

feces (Supplementary Fig. S2) compared to our previous measurements (Kang et al., 2020).

Through reanalysis, we identified 844 metabolites in plasma, and 825 metabolites in feces (Fig.

2), which is a substantial increase in identified metabolites from the original study. Furthermore,

several biochemical compounds showed notable shifts in plasma and fecal metabolic profiles

after MTT in ASD children. Following treatment, 20 metabolites in plasma and 22 in feces

whose peak intensities were initially dissimilar to TD became similar to TD (Fig. 3).

For plasma metabolites, we found similar results to Kang et al., 2020 for bilirubin (an

oxidant), iminodiacetate (xenobiotic and a glycine derivative) (Lin et al. 2019), and sarcosine (a

glycine derivative, and beneficial against depression and schizophrenia) (Tsai et al., 2004). The

relative intensities of these compounds were significantly lower at baseline but increased

significantly after MTT in children with ASD (Fig. 2). Also, the intensity of caprylate (8:0, a

medium-chain fatty acid) was higher at baseline and significantly decreased after MTT. The

analysis for these metabolites shows the consistency of the data previously reported by Kang et

al. (2019) and improvement of plasma metabolite profiles after MTT in children with ASD.

However, with this re-analysis we detected some potentially important metabolites, such as

caproate (6:0, a medium-chain fatty acid), phospholipids such as glycerophosphocholine (GPC),

phosphoethanolamine (PEA), glycerophosphoethanolamine (GPEA), and 2-palmitoyl-GPC

(16:0). The relative intensity of these metabolites was higher at baseline and significantly

decreased after MTT in children with ASD (Fig. 3A). Kang et al. postulated that a decrease in

medium-chain fatty acids, such as caprylate to be correlated with an improvement in ASD

symptoms (Kang et al., 2020).

Phospholipids are lipids that are highly abundant in neuronal membranes (Joensuu et al.,

2020), and they are an important source of energy and provide long-chain polyunsaturated fatty

acids (LC-PUFAs). Some research suggests that dysregulated phospholipid metabolism may be a

biological component of neurodevelopmental disorders like ASD (Pastural et al., 2009; Tamji et

al., 2010; El-Ansary et al. 2011; Brown et al., 2011; Ventura et al., 2020; Needham et al., 2021).

Increased sphingolipid-1-PO_4, a type of phospholipid that was observed in ASD serum compared

to TD (Wang et al., 2016). In our fecal samples, relative intensities of sphingosines N-palmitoyl-

sphingosine (d18:1/16:0), N-stearoyl-sphingosine (d18:1/18:0), and N-oleoyl-sphingosine

(d18:1/18:0) (also called ceramides) were significantly higher at ASD baseline compared to TD

but after MTT, they significantly decreased, and their abundance became similar to TD (Fig.

3B). These lipids are important for phospholipid metabolism and mitochondrial function (Tamji

& Crawford, 2011). However, the differential abundance of these mitochondrial markers in fecal

and plasma metabolites in children with ASD suggests potential mitochondrial dysfunction in

children with ASD and may contribute to ASD severity (Needham et al., 2021; Pastural et al,

2009; Rossignol & Frye, 2012; Griffiths & Levy, 2017; Lombard, 1998).

In plasma, GPC (phospholipid) levels were significantly higher in children with ASD

compared to their TD counterparts. While GPC levels decreased following MTT, the change was

not statistically significant (Fig 3A). Similarly, Needham et al reported that GPC was

significantly higher in ASD children's plasma samples and was associated with mitochondrial

dysfunction (Needham et al., 2021). GPC is a phospholipid-containing choline found in human

milk (Dangat & Johsi, 2023) and is an important nutrient for memory, mood, and muscle

regulation. However, administration of GPC has been shown to improve cognition against

dementia and Alzheimer's disease (AD) (Amenta et al., 2014; Sagaro et al., 2023). In animal

studies, administration of GPC was found to release acetylcholine in rat brains, elevate dopamine

and serotonin levels in the frontal cortex, and enhance cognitive function (Traini et al., 2020).

Despite the beneficial roles of phospholipids containing choline, increased GPC levels in our

samples from children with ASD suggest dysregulation in phospholipid metabolism, and it has

been reported that elevated levels of GPC+phosphocholines (PC) can be associated with bipolar

disorder patients (Cao et al., 2017).

Another plasma metabolite, indolepropionate (IPA or Indole-3-propionate), was

significantly lower at baseline compared to TD (Fig. 3A). IPA is a gut microbiota-derived

metabolite of tryptophan, and is increasingly recognized for its neuroprotective and anti-

inflammatory properties. In this study, IPA was significantly lower in children with ASD at

baseline compared to TD controls, suggesting a potential deficiency in beneficial microbial

activity or a disrupted microbial-host interface. Previous studies have reported similarly lower

IPA levels in ASD animal models and in children with intrauterine growth restriction (Wang et

al., 2023). Interestingly, IPA levels did not significantly change following MTT, indicating that

this metabolite may be less responsive to short-term microbial interventions. The persistent

reduction (non-significantly) in IPA highlights the need for further investigation into specific

microbial contributors and whether longer-term or targeted interventions could restore its

production and potential neuroprotective effects in ASD.

In feces, p-cresol and p-cresol sulfate have consistently emerged as metabolites of

interest due to their elevated presence in the urine and feces of children diagnosed with ASD

(Altieri et al., 2011, Chen et al., 2013; Kang et al., 2018; Daneberga et al., 2022; Diémé et al.,

2015; Emond et al., 2013; Gevi et al 2016; Gevi et al., 2020, Li et al., 2018; Mussap et al., 2020;

Osredkar et al., 2023; Perisco & Napolioni 2013, Piras et al., 2022; Tevzadze et al., 2017;

Timperio et al., 2022; Flynn et al., 2025). The primary source of p-cresol in the body is its

production by gut bacteria and act as an antimicrobial compound (Saito et al., 2019). p-cresol is

produced *via* tyrosine metabolism and is considered a uremic toxin (Candleliere et al., 2022). To

eliminate p-cresol from the body, the host liver carries out sulfation andforms$p$-cresol sulfate

which increases its solubility so that it can be more easily excreted from the body (Gryp et al.,

2017; Spanogiannopoulos et al., 2016; Clayton et al., 2009).

We observed that peak intensity of fecal p-cresol sulfate was significantly higher at ASD

baseline compared to TD (Fig. 3B, 7A). After MTT, p-cresol sulfate decreased significantly, and

its intensity became more similar to TD (Fig. 3B, Fig. 7A). However, we did not observe

significant differences or changes between groups for p-cresol in feces, p-cresol sulfate for

plasma, and p-cresol was not detected in plasma. Recent mouse studies have shown that when p-

cresol was administered, mice exhibited social behavior deficits and more repetitive behavior

relative to controls (Bermudez-Martin et al., 2021), and subsequent fecal microbiota transplant

from healthy mice led to normalization of ASD-like symptoms (Bermudez-Martin et al., 2021).

These findings suggest that *p*-cresol and/or *p*-cresol sulfate may contribute to the

pathophysiology of autism. Current results led to a hypothesis that MTT helps to reduce the *p*-

cresol sulfate levels by remodeling the gut microbial ecology in ASD children and allowing the

growth of microbiota from healthy donors. Another sulfated metabolite that has been associated

with ASD in other studies, is 4-ethylphenyl sulfate (4EPS) (Hsiao et al., 2013; Needham et al.,

2021; Stewart Campbell et al., 2022). While we did not detect 4EPS in fecal samples, wealso did

not observe significant differences or changes between groups in plasma samples. Recent clinical

trials in children with ASD using AB-2004, an adsorbent, showed decreased *p*-cresol sulfate and

4EPS levels in urine. These changes occurred alongside improvements in ASD-like behaviors,

suggesting 4EPS and *p*-cresol sulfate could be correlated with behavioral severity in ASD

(Stewart Campbell et al., 2022).

In feces, another interesting microbial metabolite, Urolithin-A was significantly higher at

baseline compared to TD (Fig. 3B). Although Urolithin A is a gut microbiota-derived metabolite

produced from dietary ellagitannins found in fruits such as pomegranates and berries (García-

Villalba et al., 2013; Selma et al., 2014), its elevated levels at baseline in individuals with ASD

may reflect a dysbiotic microbiota enriched in urolithin-producing taxa despite the self-restricted

diets commonly observed in ASD. After MTT, the introduction of a more balanced microbial

community likely reduced these specialized microbes, leading to lower Urolithin-A levels but no

statistical change except at week3 (Fig. 3B). We hypothesize that the high Urolithin-A levels in

ASD may be a marker of dysbiosis rather than of beneficial microbial function, and further

studies are required to confirm the microbial drivers of Urolithin-A production and its biological

relevance in ASD.

It is important to note that certain metabolites such as theobromine and cetirizine (Fig.

3B), while statistically significant between groups, reflect exogenous intake (e.g., chocolate,

allergy medications) rather than underlying biological shifts. Their appearance underscores the

importance of controlling or at least documenting for diet, medication use, and seasonal

variation, which should be explored in future studies with larger, placebo-controlled cohorts and

dietary monitoring.

Our most robust multivariate FDA model was composed of five metabolites from either

plasma or fecal samples; *p*-cresol sulfate (feces), hydroxyproline (feces), caprylate (plasma),

indole-propionate (plasma), and bilirubin (plasma), that were significantly different at baseline

vs TD and became similar to TD after MTT (Fig. 3A-B). Fecal *p*-cresol sulfate emerged as a

recurring component in many of the top-performing multivariate FDA models. Univariate

analysis demonstrated strong differences in the composition of *p*-cresol sulfate between the ASD

baseline and TD groups, achieving an AUROC of 0.76. These differences were utilized in the

linear multivariate models relying on this variable, highlighting *p*-cresol sulfate as a potentially

relevant biomarker for ASD in feces. Higher caprylate (plasma), and lowerindole-propionate

(plasma), and lower bilirubin (plasma) were also observed in the previous study (Kang et al.,

2020) at baseline and after MTT, relative intensity became similar to TD levels. Indole-

propionate, a gut metabolite has been found to be associated with ASD in humans (Yu et al.,

2020) and animal models (Wang et al., 2023; Jiang et al., 2024). Hydroxyproline, another FDA

model-identified metabolite (Fig. 4D), was previously identified at significantly higher relative

intensity in the plasma of children with ASD (Chen et al., 2024).

Combining plasma and fecal metabolite datasets resulted in higher cross-validated accuracy than

using either dataset alone, potentially as a consequence of complimenting biological insight

provided by each source. The FDA models developed for the reanalysis demonstrated an

accuracy of 95%, outperforming previous work with fecal metabolites from the same study,

which achieved an accuracy of 94.7% (Qureshi et al., 2020). Previous assessments of plasma

metabolites have also shown comparable capabilities to distinguish between ASD and TD

cohorts using linear classifier techniques (Howsmon et al., 2017; Adams et al., 2019). This prior

work also highlighted that after MTT treatment, the predictive accuracy and average AUROC of

fecal metabolites decreased substantially (Qureshi et al., 2020). This pattern was observed in the

metabolomic panels composed of both plasma and fecal samples in this study as well, suggesting

that improvements in several metabolites may have compromised their predictive validity.

Although cross-validation demonstrates the robustness of these findings, the influence of

geographic covariates on microbiome composition should be assessed to determine the broader

applicability of these results (Falony et al., 2016). Future studies should incorporate external

validation with different cohorts with a large dataset and longitudinal sampling to confirm the

reproducibility and clinical applicability of these models. No consistent metabolite alterations

were observed across plasma and fecal samples. This may reflect differential host–microbiota

interactions, underscoring the complexity of cross-compartmental signaling. Furthermore,

studies involving larger cohorts with a placebo arm, along with mechanistic investigations, will

be important to better elucidate the metabolic dynamics in ASD and ensure the broader clinical

relevance of MTT's impact.

Although the initial shifts in the metabolic profile observed by week 3 reflect the effects

of vancomycin treatment, our data suggest that these changes alone do not fully explain the

sustained alignment with the TD group seen after MTT. Notably, plasma metabolites remained

similar to TD profiles at week 18, indicating that MTT may help stabilize or maintain these

shifts. Also, we observed only modest changes in fecal metabolites by week 3 (after a high-dose

of microbiota), with clearer transitions toward TD-like profiles occurring after 8 weeks of MTT.

These findings are consistent with Sandler et al. (Sandler et al., 2000), where vancomycin led to

temporary improvements in autism symptoms, but these improvements were lost post-treatment,

potentially due to lack of recolonization by beneficial microbes. In contrast, our results show that

MTT may preserve or reinforce beneficial shifts initiated by antibiotics. This interpretation is

consistent with prior findings in recurrent *Clostridioides difficile* infection (Zhou et al., 2023)

and highlights the need to consider both antibiotic and transplant contributions in interpreting

longitudinal metabolomic responses. In conclusion, the reanalysis of plasma and fecal

metabolites revealed an additional 223 plasma and 159 fecal metabolites compared to the

previous report by Kang et al., 2020. The contrast in metabolite composition before and after

MTT groups provides insight on metabolites with potential biological relevance in children with

autism and/or GI issues. This research underscores the value of re-analysis when databases

improve and the importance of investigating non-behavioral features in ASD diagnosis and

provides valuable insights into the intricate metabolic landscape of this complex disorder.

Using different analysis methods, we consistently observed that sarcosine,

iminodiacetate, caproate, and caprylate in plasma were significantly different in ASD vs TD, as

reported previously. However, we also identified phospholipids such as

glycerophosphoethanolamine (GPE), glycerophosphocholine (GPC), and phosphoethanolamine

in plasma, as well as fecal sphingosines N-palmitoyl-sphingosine (d18:1/16:0), N-stearoyl-

sphingosine (d18:1/18:0), and N-oleoyl-sphingosine (d18:1/18:0), which were significantly

different at baseline compared to typically developing (TD) individuals and/or changed after

MTT. Moreover, we observed that fecal *p*-cresol sulfate was higher at baseline compared to TD

and decreased significantly after MTT.

Notably, *p*-cresol sulfate in feces was consistently found across our different statistical

methods as a potential biomarker before and after MTT in children with ASD. This suggests the

potential of microbially-produced metabolite in affecting ASD symptoms. Larger cohort and

placebo-controlled trials are necessary to validate our findings and unravel the mechanisms

underlying MTT in ameliorating aberrant metabolomic profiles associated with ASD.

Furthermore, insights into the underlying biochemical pathways will facilitate the development

of more tailored treatment strategies.

Additional Information: The Supplementary Material for this article can be found online at:

Data Availability.

The clinical data used in this study are available from the previously published work by Kang et

al. (Kang et al. 2019, 2020). The metabolomics datasets analyzed in this manuscript are provided

in the supplemental material file 2-3.

Code Availability.

No custom code was developed for this study, except for FDA analysis (Fig.4D) which can be

found at https://github.com/FatirQureshi/FDA-Model-Search. All analyses were performed using

standard software packages, as described in the Methods section of the manuscript.

Abbreviations:

1. **4EPS** - 4-Ethylphenyl Sulfate
2. **AMP** - Adenosine-5'-Monophosphate
3. **ASD** - Autism Spectrum Disorder
4. **AUROC** - Area Under the Receiver Operating Characteristic Curve

5. **CV** - Cross validation
6. **FDA** - Fisher's Discriminant Analysis (context-dependent, could also mean Food and Drug Administration)
7. **FDR** - False Discovery Rate
8. **GABA** - Gamma-Aminobutyric Acid
9. **GC-MS** - Gas Chromatography-Mass Spectrometry
10. **GI** - Gastrointestinal
11. **GPC** - Glycerophosphocholine
12. **GPEA** - Glycerophosphoethanolamine
13. **IMP** - Inosine 5'-Monophosphate
14. **KEGG** - Kyoto Encyclopedia of Genes and Genomes
15. **LC-PUFA** - Long-Chain Polyunsaturated Fatty Acid
16. **LOO** - Leave-One-Out
17. **LOO-CV** - Leave-One-Out Cross-Validation
18. **MTT** - Microbiota Transplant Therapy
19. **PC** - Phosphocholine
20. **PCA** - Principal Component Analysis
21. **PEA** - Phosphoethanolamine
22. **TD** - Typically Developing
23. **UHPLC-MS/MS** - Ultrahigh-Performance Liquid Chromatography-Tandem Mass Spectroscopy

Authors Contributions

Conceptualization, K.N., D.W.K., J.B.A., R.K.B.; Data curation, K.N, F.Q., J.B.A., R.K.B.;

Formal analysis, K.N, F.Q., J.B.A., R.K.B.; Funding acquisition, D.W.K., J.B.A., R.K.B.;

Investigation, K.N, F.Q., J.B.A., R.K.B.; Methodology, K.N., J.H., J.B.A., R.K.B.; Project

administration, R.K.B., J.B.A.; Software, K.N.; F.Q.; Supervision, D.W.K., J.H., J.B.A., R.K.B.;

Visualization, K.N., F.Q.; Writing—original draft, K.N., F.Q., J.B.A., R.K.B.; Writing—review

& editing, , K.N., F.Q., D.W.K., J.H., J.B.A., R.K.B.

Disclosure/Conflict of interest

K.N., J.B.A. D. -W.K. and R.K.-B. have pending/approved patents for autism biomarkers and the

use of FMT for various conditions, including autism. J.B.A. and R.K.-B are co-founders of

Autism Diagnostics LLC and Gut-Brain Axis Therapeutics. The other authors declare no

competing interests.

Acknowledgments

We thank all children with ASD and typically developing children for participating in the study as well as the children's families for their support on the study. We thank Biodesign Center for Health Through Microbiomes Start-up at Arizona State University for the financial support and Joel Meraz for assisting with the references. This work was supported by Finch Therapeutics (formerly Crestovo) to obtain the metabolomics data set.

659 **Figure 1**: The study design of microbiota transplant therapy for fecal and plasma metabolomics (A) and

660 the overview of the data analyses for fecal and plasma metabolites (B).

661 Plasma was collected right before high-dose microbiota and feces were collected right after high-dose

662 microbiota.

664
665 **Figure 2**. An overview of plasma metabolomics with the previous A) and re-analyzed B), fecal

666 metabolomics with the previous C) and re-analyzed D) of peaks. The pie charts show the total number of

667 metabolites identified and the type of metabolites in plasma and fecal samples.

Figure 3. Univariate comparison of the (z-score) relative intensity of plasma (A) and fecal (B) metabolites in children with ASD and comparisons after MTT and with TD. For the plasma metabolites (A), a longitudinal paired-wise univariate analysis was performed at baseline vs. week3, week10 and cross-sectional unpaired comparison vs. TD. Similarly, for the fecal metabolites (B), a longitudinal paired-wise univariate analysis was performed (shown in solid blue lines at the bottom of the heatmap) at baseline vs. week3, week10, week18 and cross-sectional unpaired comparison vs. TD (shown in dashed blue line). (C) heatmap (z-score) for each participant with 20 plasma metabolites that were significantly changed or were different in children with ASD at baseline compared with MTT and with TD groups in Figure 3A. Week18 time was not available for the plasma. See Supplementary Fig. S1 for fecal metabolites heatmap for each participant. The median of relative intensities in plasma (A) and feces (B) was used to construct the heatmap for each group. *Single asterisk indicates p <0.05, **double asterisks indicate p <0.01, triple *** asterisks indicate p <0.001. # indicates the statistically significant (p<0.05)

metabolites after log-transformation. All p-values are FDR corrected. ASD: Autism Spectrum Disorders,

TD: Typically Developing.

Figure 4. An average Receiver operating characteristic curves (AUROC) of A) plasma, B) fecal

metabolites, C) principal component analysis (PCA) of plasma metabolites, and D) FDA analysis of fecal

and plasma metabolites (*p*-cresol sulfate, caprylate (C:8) bilirubin, indolepropionate and hydroxyproline)

using probability density curves associated with FDA scores before and after MTT in ASD, and TD

children. For FDA analysis, true positive vs false positive rate was determined across all threshold values

to characterize between ASD Baseline against all possible alternative groups. For AUROC and PCA

plots, 22 fecal and 20 plasma metabolites (presented in Figure 3) were used. Refer to Supplementary

Figure S2 for PCA plot for fecal metabolites. ASD – autism spectrum disorder, MTT - microbiota

transplant therapy, TD - typically developing, FDA - Fisher discriminant analysis.

Figure 5. Univariate comparison of the relative intensity (after log10 transformation) in plasma for A)

Sarcosine, B) Iminodiacetate (IDA), C) Caproate (6:0), and D) Caprylate (8:0) at ASD baseline vs. MTT

(week3, week10) and TD. Each colored dot represents one ASD individual, and grey colored dots

represent TD. Asterisks represent significant differences between ASD baseline, and the other groups

(*Single asterisk indicates $p < 0.05$, **double asterisks indicate $p < 0.01$, ns not significant, all p-values are

FDR corrected). ASD: Autism Spectrum Disorders, TD: Typically Developing.

Figure 6. Receiver operating characteristic curves (AUROC) of A) Sarcosine, B) Iminodiacetate (IDA),

C) Caproate (6:0), and D) Caprylate (8:0) for ASD baseline vs. MTT (week3, week10) and TD. True

positive vs false positive rate was determined across all threshold values to characterize between ASD

Baseline against all possible alternative groups.

Figure 7. Univariate comparison of the relative intensity (after log10 transformation) in feces for A) p-
cresol sulfate, B) N-oleoyl-sphingosine (d18:1/18:1), C) N-palmitoyl-sphingosine (d18:1/16:0), and D) N-
stearoyl-sphingosine (d18:1/18:0) at ASD baseline vs. MTT (week3, week10, week18) and TD. Each
colored dot represents one ASD individual and grey colored dots represent TD. Asterisks represent
significant differences between ASD baseline and the other groups (*Single asterisk indicates p <0.05,
**double asterisks indicate p<0.01, ns not significant, all p-values are FDR corrected). ASD: Autism
Spectrum Disorders, TD: Typically Developing.

Figure 8. Receiver operating characteristic curves (AUROC) of A) p-cresol sulfate, B) (N-oleoyl-sphingosine d18:1/18:1), C) N-palmitoyl-sphingosine (d18:1/16:0), and D) N-stearoyl-sphingosine (d18:1/18:0) at ASD baseline vs. MTT (week3, week10, week18) and TD. True positive vs false positive rate was determined across all threshold values to characterize between ASD baseline against all possible alternative groups. pCS - p-cresol sulfate.

References:

1. Adams JB, Johansen LJ, Powell LD, Quig D, Rubin RA. Gastrointestinal flora and gastrointestinal status in children with autism – comparisons to typical children and correlation with autism severity. BMC Gastroenterol. 2011a. doi:10.1186/1471-230X-11-22.

2. Adams JB, Audhya T, McDonough-Means S, Rubin RA, Quig D, Geis E, Gehn E, Loresto M, Mitchell J, Atwood S, Barnhouse S, Lee W. Nutritional and metabolic status of children with autism vs. neurotypical children, and the association with autism severity. Nutr Metab (Lond). 2011b ;8(1):34. doi: 10.1186/1743-7075-8-34. PMID: 21651783; PMCID: PMC3135510.

3. Adams, J.B.; Vargason, T.; Kang, D.-W.; Krajmalnik-Brown, R.; Hahn, J. Multivariate Analysis of Plasma Metabolites in Children with Autism Spectrum Disorder and Gastrointestinal Symptoms Before and After Microbiota Transfer Therapy. *Processes* **2019**, *7*, 806. https://doi.org/10.3390/pr7110806.

4. Altieri, L., Neri, C., Sacco, R., Curatolo, P., Benvenuto, A., Muratori, F., Santocchi, E., Bravaccio, C., Lenti, C., Saccani, M., Rigardetto, R., Gandione, M., Urbani, A., & Persico, A. M. (2011). Urinary p-cresol is elevated in small children with severe autism spectrum disorder. Biomarkers, 16(3), 252–260. https://doi.org/10.3109/1354750X.2010.548010

5. Altman, N., Krzywinski, M. Ensemble methods: bagging and random forests. Nat Methods 14, 933–934 (2017). https://doi.org/10.1038/nmeth.4438

6. Amenta F, Carotenuto A, Fasanaro AM, Rea R, Traini E. The ASCOMALVA (Association between the Cholinesterase Inhibitor Donepezil and the Cholinergic Precursor Choline Alphoscerate in Alzheimer's Disease) Trial: interim results after two years of treatment. J Alzheimers Dis. 2014;42 Suppl 3:S281-8. doi: 10.3233/JAD-140150

7. Arici H, Qureshi FA, Mulmule J, Hahn J. 2025. Multiple hypothesis testing for pre-screening of factor selection for classification of high-dimensional data. *J Process Control* 152:103469. https://doi.org/10.1016/j.jprocont.2025.103469.

8. Bermudez-Martin, P., Becker, J.A.J., Caramello, N. et al. The microbial metabolite p-Cresol induces autistic-like behaviors in mice by remodeling the gut microbiota. Microbiome 9, 157 (2021). https://doi.org/10.1186/s40168-021-01103-z.

9. Brown CM, Austin DW. Autistic disorder and phospholipids: A review. Prostaglandins Leukot Essent Fatty Acids. 2011 Jan-Feb;84(1-2):25-30. doi: 10.1016/j.plefa.2010.09.007. PMID: 20970971.

10. Cao, B., Stanley, J., Passos, I. *et al.* Elevated Choline-Containing Compound Levels in Rapid Cycling Bipolar Disorder. *Neuropsychopharmacol* **42**, 2252–2258 (2017). https://doi.org/10.1038/npp.2017.39

11. CDC. 2023-ADDM-Factsheet_508; 2023. https://www.cdc.gov/autism/data-research/index.html .

12. Chen, Q., Qiao, Y., Xu, X. J., Tao, Y., & You, X. (2019). Urine organic acids as potential biomarkers for autism-spectrum disorder in chinese children. Frontiers in Cellular Neuroscience, 13. https://doi.org/10.3389/fncel.2019.00150

13. Chen, WX., Chen, YR., Peng, MZ. et al. Plasma Amino Acid Profile in Children with Autism Spectrum Disorder in Southern China: Analysis of 110 Cases. J Autism Dev Disord 54, 1567–1581 (2024). https://doi.org/10.1007/s10803-022-05829-z

14. Clayton TA, Baker D, Lindon JC, Everett JR, Nicholson JK. Pharmacometabonomic identification of a significant host-microbiome metabolic interaction affecting human drug metabolism. Proc Natl Acad Sci U S A. 2009 Aug 25;106(34):14728-33. doi: 10.1073/pnas.0904489106. Epub 2009 Aug 10. PMID: 19667173; PMCID: PMC2731842.

15. Created in BioRender. Krajmalnik-brown, R. (2022) https://BioRender.com/x41x091.

16. Daneberga, Z., Nakazawa-Miklasevica, M., Berga-Svitina, E., Murmane, D., Isarova, D., Cupane, L., Masinska, M., Nartisa, I., Lazdane, A., & Miklasevics, E. (2022). Urinary organic acids spectra in children with altered gut microbiota composition and autistic spectrum disorder. Nordic Journal of Psychiatry, 76(7), 523–529. https://doi.org/10.1080/08039488.2021.2014954

17. Dangat K, Joshi S. *Breast Milk and Cognitive Performance in Children*. Elsevier Inc.; 2023. https://www.sciencedirect.com/science/article/abs/pii/B9780323898348000143

18. De Angelis M., Piccolo M., Vannini L., Siragusa S., De Giacomo A., Serrazzanetti D.I.,
Cristofori F., Guerzoni M.E., Gobbetti M., Francavilla R. Fecal microbiota and
metabolome of children with autism and pervasive developmental disorder not otherwise
specified. PLoS ONE. 2013;8:e76993. doi: 10.1371/journal.pone.0076993.

19. Diémé, B., Mavel, S., Blasco, H., Tripi, G., Bonnet-Brilhault, F., Malvy, J., Bocca, C.,
Andres, C. R., Nadal-Desbarats, L., & Emond, P. (2015). Metabolomics Study of Urine in
Autism Spectrum Disorders Using a Multiplatform Analytical Methodology. Journal of
Proteome Research, 14(12), 5273–5282.

20. El-Ansary AK, Bacha AG, Al-Ayahdi LY. Impaired plasma phospholipids and relative
amounts of essential polyunsaturated fatty acids in autistic patients from Saudi Arabia.
Lipids Health Dis. 2011 Apr 22;10:63. doi: 10.1186/1476-511X-10-63. PMID: 21513514;
PMCID: PMC3107801.

21. El-Ansary, A., Al-Ayadhi, L. GABAergic/glutamatergic imbalance relative to excessive
neuroinflammation in autism spectrum disorders. J Neuroinflammation 11, 189 (2014).
https://doi.org/10.1186/s12974-014-0189-0

22. Emond P, Mavel S, Aïdoud N, Nadal-Desbarats L, Montigny F, Bonnet-Brilhault F,
Barthélémy C, Merten M, Sarda P, Laumonnier F, Vourc'h P, Blasco H, Andres CR. GC-
MS-based urine metabolic profiling of autism spectrum disorders. Anal Bioanal Chem.
2013 Jun;405(15):5291-300. doi: 10.1007/s00216-013-6934-x. Epub 2013 Apr 10.
PMID: 23571465.

23. Falony G, Joossens M, Vieira-Silva S, Wang J, Darzi Y, Faust K, Kurilshikov A, Bonder
MJ, Valles-Colomer M, Vandeputte D, Tito RY, Chaffron S, Rymenans L, Verspecht C,
De Sutter L, Lima-Mendez G, D'hoe K, Jonckheere K, Homola D, Garcia R, Tigchelaar
EF, Eeckhaudt L, Fu J, Henckaerts L, Zhernakova A, Wijmenga C, Raes J. 2016.
Population-level analysis of gut microbiome variation. *Science* 352(6285):560–564.
https://doi.org/10.1126/science.aad3503.

24. Flynn CK, Adams JB, Krajmalnik-Brown R, et al. Review of Elevated Para-Cresol in
Autism and Possible Impact on Symptoms. Int J Mol Sci. 2025;26(4):1513. Published
2025 Feb 11. doi:10.3390/ijms26041513.

25. Gabriele S, Sacco R, Cerullo S, et al. Urinary p -cresol is elevated in young French children with autism spectrum disorder: a replication study. Biomarkers. 2014;19(6):463-470. doi:10.3109/1354750X.2014.936911.

26. García-Villalba R, Beltrán D, Espín JC, Selma MV, Tomás-Barberán FA. Time course production of urolithins from ellagic acid by human gut microbiota. J Agric Food Chem. 2013;61(37):8797-8806. doi:10.1021/jf402498b

27. Gevi, F., Zolla, L., Gabriele, S., & Persico, A. M. (2016). Urinary metabolomics of young Italian autistic children supports abnormal tryptophan and purine metabolism. Molecular Autism, 7, 47. https://doi.org/10.1186/s13229-016-0109-5.

28. Gevi, F., Belardo, A., & Zolla, L. (2020). A metabolomics approach to investigate urine levels of neurotransmitters and related metabolites in autistic children. Biochimica et Biophysica Acta (BBA) - Molecular Basis of Disease, 1866(10), 165859. https://doi.org/10.1016/j.bbadis.2020.165859

29. Golubeva AV, Joyce SA, Moloney G, Burokas A, Sherwin E, Arboleya S, Flynn I, Khochanskiy D, Moya-Pérez A, Peterson V, Rea K, Murphy K, Makarova O, Buravkov S, Hyland NP, Stanton C, Clarke G, Gahan CGM, Dinan TG, Cryan JF. Microbiota-related Changes in Bile Acid & Tryptophan Metabolism are Associated with Gastrointestinal Dysfunction in a Mouse Model of Autism. EBioMedicine. 2017 Oct;24:166-178. doi: 10.1016/j.ebiom.2017.09.020. Epub 2017 Sep 21. PMID: 28965876; PMCID: PMC5652137.

30. Gorrindo P, Williams KC, Lee EB, et al. Gastrointestinal dysfunction in autism: parental report, clinical evaluation, and associated factors. Autism Res 2012; 5(2):101–8.

31. Griffiths KK, Levy RJ, Evidence of Mitochondrial Dysfunction in Autism: Biochemical Links, Genetic-Based Associations, and Non-Energy-Related Mechanisms. Oxid. Med. Cell. Longev 2017 (2017), doi:10.1155/2017/4314025. - DOI - PMC - PubMed

32. Gryp T, Vanholder R, Vaneechoutte M, Glorieux G. p-Cresyl Sulfate. Toxins (Basel). 2017 Jan 29;9(2):52. doi: 10.3390/toxins9020052. PMID: 28146081; PMCID: PMC5331431.

33. Hashimoto, K. Emerging role of the host microbiome in neuropsychiatric disorders: overview and future directions. Mol Psychiatry 28, 3625–3637 (2023). https://doi.org/10.1038/s41380-023-02287-6

34. He, J., Gong, X., Hu, B. et al. Altered Gut Microbiota and Short-chain Fatty Acids in

Chinese Children with Constipated Autism Spectrum Disorder. Sci Rep 13, 19103 (2023).

https://doi.org/10.1038/s41598-023-46566-2

35. Howsmon DP, Kruger U, Melnyk S, James SJ, Hahn J. Classification and adaptive

behavior prediction of children with autism spectrum disorder based upon multivariate

data analysis of markers of oxidative stress and DNA methylation. PLoS Comput Biol.

2017;13(3):e1005385. Published 2017 Mar 16. doi:10.1371/journal.pcbi.1005385.

36. Hsiao EY, McBride SW, Hsien S, Sharon G, Hyde ER, McCue T, Codelli JA, Chow J,

Reisman SE, Petrosino JF, Patterson PH, Mazmanian SK. Microbiota modulate

behavioral and physiological abnormalities associated with neurodevelopmental

disorders. Cell. 2013 Dec 19;155(7):1451-63. doi: 10.1016/j.cell.2013.11.024. Epub 2013

Dec 5. PMID: 24315484; PMCID: PMC3897394.

37. Huberty, C. J. (1975). Discriminant analysis. Review of Educational Research, 45(4),

543-598.

38. Hung, L.Y., Margolis, K.G. Autism spectrum disorders and the gastrointestinal tract:

insights into mechanisms and clinical relevance. Nat Rev Gastroenterol Hepatol (2023).

https://doi.org/10.1038/s41575-023-00857-1

39. Javaneh Tamiji, Dorota A. Crawford. The Neurobiology of Lipid Metabolism in Autism

Spectrum Disorders. *Neurosignals* 2011; 18 (2): 98–112.

https://doi.org/10.1159/000323189

40. Jiang, J., Wang, D., Jiang, Y. et al. The gut metabolite indole-3-propionic acid activates

ERK1 to restore social function and hippocampal inhibitory synaptic transmission in a

16p11.2 microdeletion mouse model. Microbiome 12, 66 (2024).

https://doi.org/10.1186/s40168-024-01755-7

41. Joensuu M, Wallis TP, Saber SH, Meunier FA. Phospholipases in neuronal function: A

role in learning and memory? J Neurochem. 2020;153(3):300-333. doi:

10.1111/jnc.14918. Epub 2020 Feb 7. PMID: 31745996.

42. Kang DW, Adams JB, Gregory AC, et al. Microbiota Transfer Therapy alters gut

ecosystem and improves gastrointestinal and autism symptoms: an open-label study.

Microbiome. 2017;5(1):10. doi:10.1186/s40168-016-0225-7.

43. Kang D.W., Ilhan Z.E., Isern N.G., Hoyt D.W., Howsmon D.P., Shaffer M., Lozupone
C.A., Hahn J., Adams J.B., Krajmalnik-Brown R. Differences in fecal microbial
metabolites and microbiota of children with autism spectrum disorders. Anaerobe.
2018;49:121–131. doi:10.1016/j.anaerobe.2017.12.007.

44. Kang DW, Adams JB, Coleman DM, Pollard EL, Maldonado J, McDonough-Means S,
Caporaso JG, Krajmalnik-Brown R. Long-term benefit of Microbiota Transfer Therapy
on autism symptoms and gut microbiota. Sci Rep. 2019 Apr 9;9(1):5821. doi:
10.1038/s41598-019-42183-0. PMID: 30967657; PMCID: PMC6456593.

45. Kang DW, Adams JB, Vargason T, Santiago M, Hahn J, Krajmalnik-Brown R. Distinct
Fecal and Plasma Metabolites in Children with Autism Spectrum Disorders and Their
Modulation after Microbiota Transfer Therapy. Marco ML, ed. *mSphere*.
2020;5(5):e00314-20. doi:10.1128/mSphere.00314-20.

46. Khalifa D, Shahin O, Salem D, Raafat O. Serum glutamate was elevated in children aged
3-10 years with autism spectrum disorders when they were compared with controls. Acta
Paediatr. 2019 Feb;108(2):295-299. doi: 10.1111/apa.14477. Epub 2018 Jul 16. PMID:
29949195.

47. Krajmalnik-Brown R, Lozupone C, Kang D-W, Adams JB. Gut bacteria in children with
autism spectrum disorders: challenges and promise of studying how a complex
community influences a complex disease. Microb Ecol Health Dis. 2015; 26.
doi:10.3402/mehd.v26.26914.

48. Kumar A, Pramanik J, Goyal N, et al. Gut Microbiota in Anxiety and Depression:
Unveiling the Relationships and Management Options. *Pharmaceuticals (Basel)*.
2023;16(4):565. Published 2023 Apr 9. doi:10.3390/ph16040565

49. Leyrolle Q, Prado-Perez L, Layé S. The gut-derived metabolites as mediators of the
effect of healthy nutrition on the brain. Front Nutr. 2023 Jun 9;10:1155533. doi:
10.3389/fnut.2023.1155533. PMID: 37360297; PMCID: PMC10289296.

50. Li C, Shen K, Chu L, Liu P, Song Y, Kang X. Decreased levels of urinary free amino
acids in children with autism spectrum disorder. J Clin Neurosci. 2018 Aug;54:45-49.
doi: 10.1016/j.jocn.2018.05.001. Epub 2018 May 28. PMID: 29853226.

51. Lin S, Yue X, Wu H, Han T-L, Zhu J, Wang C, et al. Explore potential plasma biomarkers
of acute respiratory distress syndrome (ARDS) using GC-MS metabolomics

52. Lombard J, Autism: a mitochondrial disorder? Med. Hypotheses 50, 497–500 (1998). –
    PubMed

53. Lord C, Elsabbagh M, Baird G, Veenstra-Vanderweele J. Autism spectrum disorder.
    Lancet. 2018 Aug 11;392(10146):508-520. doi: 10.1016/S0140-6736(18)31129-2. Epub
    2018 Aug 2. PMID: 30078460; PMCID: PMC7398158.

54. Martin CR, Osadchiy V, Kalani A, Mayer EA. The Brain-Gut-Microbiome Axis. Cell Mol
    Gastroenterol Hepatol. 2018;6(2):133-148. Published 2018 Apr 12.
    doi:10.1016/j.jcmgh.2018.04.003

55. Morton JT, Jin DM, Mills RH, et al. Multi-level analysis of the gut–brain axis shows
    autism spectrum disorder-associated molecular and microbial profiles. Nat Neurosci.
    2023;26(7):1208-1217. doi:10.1038/s41593-023-01361-0

56. Mottron, L., Bzdok, D. Autism spectrum heterogeneity: fact or artifact?. Mol Psychiatry
    25, 3178–3185 (2020). https://doi.org/10.1038/s41380-020-0748-y.

57. Mussap, M., Siracusano, M., Noto, A., Fattuoni, C., Riccioni, A., Rajula, H. S. R., Fanos,
    V., Curatolo, P., Barberini, L., & Mazzone, L. (2020). The Urine Metabolome of Young
    Autistic Children Correlates with Their Clinical Profile Severity. Metabolites, 10(11),
    476. https://doi.org/10.3390/metabo10110476.

58. Needham BD, Adame MD, Serena G, Rose DR, Preston GM, Conrad MC, Campbell AS,
    Donabedian DH, Fasano A, Ashwood P, Mazmanian SK. Plasma and Fecal Metabolite
    Profiles in Autism Spectrum Disorder. Biol Psychiatry. 2021 Mar 1;89(5):451-462. doi:
    10.1016/j.biopsych.2020.09.025. Epub 2020 Oct 10. PMID: 33342544; PMCID:
    PMC7867605.

59. Osredkar, J., Baškovič, B. Ž., Finderle, P., Bobrowska-Korczak, B., Gątarek, P., Rosiak,
    A., Giebułtowicz, J., Vrhovšek, M. J., & Kałużna-Czaplińska, J. (2023). Relationship
    between Excreted Uremic Toxins and Degree of Disorder of Children with ASD.
    International Journal of Molecular Sciences, 24(8), 7078.
    https://doi.org/10.3390/ijms24087078

60. Pastural E, Ritchie S, Lu Y, Jin W, Kavianpour A, Khine Su-Myat K, Heath D, Wood PL,
    Fisk M, Goodenowe DB. Novel plasma phospholipid biomarkers of autism:
    mitochondrial dysfunction as a putative causative mechanism. Prostaglandins Leukot

Essent Fatty Acids. 2009 Oct;81(4):253-64. doi: 10.1016/j.plefa.2009.06.003. Epub 2009
Jul 15. PMID: 19608392.

61. Peralta-Marzal LN, Prince N, Bajic D, Roussin L, Naudon L, Rabot S, Garssen J,
Kraneveld AD, Perez-Pardo P. The Impact of Gut Microbiota-Derived Metabolites in
Autism Spectrum Disorders. Int J Mol Sci. 2021 Sep 17;22(18):10052. doi:
10.3390/ijms221810052. PMID: 34576216; PMCID: PMC8470471.

62. Peralta-Marzal, L.N., Rojas-Velazquez, D., Rigters, D. et al. A robust microbiome
signature for autism spectrum disorder across different studies using machine
learning. Sci Rep 14, 814 (2024). https://doi.org/10.1038/s41598-023-50601-7.

63. Persico, A. M., & Napolioni, V. (2013). Urinary p-cresol in autism spectrum disorder.
Neurotoxicology and Teratology, 36, 82–90. https://doi.org/10.1016/j.ntt.2012.09.002.

64. Petroff O.A.C. Book Review: GABA and glutamate in the human brain. Neuroscientist.
2002;8:562–573.

65. Petroff OA. GABA and glutamate in the human brain. Neuroscientist. 2002
Dec;8(6):562-73. doi: 10.1177/1073858402238515. PMID: 12467378.

66. Piras, C., Mussap, M., Noto, A., De Giacomo, A., Cristofori, F., Spada, M., Fanos, V.,
Atzori, L., & Francavilla, R. (2022). Alterations of the Intestinal Permeability are
Reflected by Changes in the Urine Metabolome of Young Autistic Children: Preliminary
Results. Metabolites, 12(2), 104. https://doi.org/10.3390/metabo12020104.

67. Qureshi, F.; Adams, J.; Hanagan, K.; Kang, D.-W.; Krajmalnik-Brown, R.; Hahn, J.
Multivariate Analysis of Fecal Metabolites from Children with Autism Spectrum
Disorder and Gastrointestinal Symptoms before and after Microbiota Transfer Therapy. *J.*
Pers. Med. **2020**, *10*, 152. https://doi.org/10.3390/jpm10040152

68. Rossignol DA, Frye RE, Mitochondrial dysfunction in autism spectrum disorders: a
systematic review and meta-analysis. Mol. Psychiatry 17, 290–314 (2012). - PMC –
PubMed

69. Sagaro GG, Traini E, Amenta F. Activity of Choline Alphoscerate on Adult-Onset
Cognitive Dysfunctions: A Systematic Review and Meta-Analysis. J Alzheimers Dis.
2023;92(1):59-70. doi: 10.3233/JAD-221189. PMID: 36683513; PMCID:
PMC10041421.

70. Saito Y, Sato T, Nomoto K, Tsuji H. Identification of phenol- and p-cresol-producing intestinal bacteria by using media supplemented with tyrosine and its metabolites. FEMS Microbiol Ecol. 2018 Sep 1;94(9):fiy125. doi: 10.1093/femsec/fiy125. Erratum in: FEMS Microbiol Ecol. 2019 Apr 1;95(4): Erratum in: FEMS Microbiol Ecol. 2020 Jan 1;96(1): PMID: 29982420; PMCID: PMC6424909.

71. Selma MV, Beltrán D, García-Villalba R, Espín JC, Tomás-Barberán FA. Description of urolithin production capacity from ellagic acid of two human intestinal Gordonibacter species. Food Funct. 2014;5(8):1779-1784. doi:10.1039/c4fo00092g

72. Serger, E., Luengo-Gutierrez, L., Chadwick, J.S. et al. The gut metabolite indole-3 propionate promotes nerve regeneration and repair. Nature 607, 585–592 (2022). https://doi.org/10.1038/s41586-022-04884-x

73. Shinohe A, Hashimoto K, Nakamura K, et al. Increased serum levels of glutamate in adult patients with autism. *Prog Neuropsychopharmacol Biol Psychiatry*. 2006;30(8):1472-1477. doi:10.1016/j.pnpbp.2006.06.013

74. Spanogiannopoulos P, Bess EN, Carmody RN, Turnbaugh PJ. The microbial pharmacists within us: a metagenomic view of xenobiotic metabolism. Nat Rev Microbiol. 2016 Apr;14(5):273-87. doi: 10.1038/nrmicro.2016.17. Epub 2016 Mar 14. PMID: 26972811; PMCID: PMC5243131.

75. Stewart Campbell A, Needham BD, Meyer CR, Tan J, Conrad M, Preston GM, Bolognani F, Rao SG, Heussler H, Griffith R, Guastella AJ, Janes AC, Frederick B, Donabedian DH, Mazmanian SK. Safety and target engagement of an oral small-molecule sequestrant in adolescents with autism spectrum disorder: an open-label phase 1b/2a trial. Nat Med. 2022 Mar;28(3):528-534. doi: 10.1038/s41591-022-01683-9. Epub 2022 Feb 14. PMID: 35165451.

76. Szeligowski T, Yun AL, Lennox BR, Burnet PWJ. The Gut Microbiome and Schizophrenia: The Current State of the Field and Clinical Applications. *Front Psychiatry*. 2020;11:156. Published 2020 Mar 12. doi:10.3389/fpsyt.2020.00156

77. Tamiji J, Crawford DA. The neurobiology of lipid metabolism in autism spectrum disorders. Neurosignals. 2010;18(2):98-112. doi:10.1159/000323189

78. Tamji, R., & Crawford, M. A. (2011). Phospholipid metabolism and mitochondrial function. *International Journal of Developmental Neuroscience*, 29(6), 633-641.

79. Takyi, E., Nirmalkar, K., Adams, J., & Krajmalnik-Brown, R. (2025). Interventions targeting the gut microbiota and their possible effect on gastrointestinal and neurobehavioral symptoms in autism spectrum disorder. Gut Microbes, 17(1). https://doi.org/10.1080/19490976.2025.2499580

80. Tevzadze, G., Shanshiashvilli, L., & Mikeladze, D. (2017). Children with epilepsy and autistic spectrum disorders show similarly high levels of urinary p-cresol. Journal of Biological Physics and Chemistry, 17(2), 77–80. https://doi.org/10.4024/05te17a.jbpc.17.02.

81. Timperio, A. M., Gevi, F., Cucinotta, F., Ricciardello, A., Turriziani, L., Scattoni, M. L., & Persico, A. M. (2022). Urinary Untargeted Metabolic Profile Differentiates Children with Autism from Their Unaffected Siblings. Metabolites, 12(9). https://doi.org/10.3390/metabo12090797.

82. Traini E, Carotenuto A, Fasanaro AM, Amenta F. Volume Analysis of Brain Cognitive Areas in Alzheimer's Disease: Interim 3-Year Results from the ASCOMALVA Trial. J Alzheimers Dis. 2020;76(1):317-329. doi: 10.3233/JAD-190623. PMID: 32508323; PMCID: PMC7369051.

83. Tsai GC, Lane HY, Yang PC, Chong MY, Lange N. 2004. Glycine transporter I inhibitor, N-methylglycine (Sarcosine), added to antipsychotics for the treatment of schizophrenia. Biol Psychiatry 55:452–456.

84. Vargason T, Roth E, Grivas G, Ferina J, Frye RE, Hahn J. Classification of autism spectrum disorder from blood metabolites: Robustness to the presence of co-occurring conditions. Res Autism Spectr Disord. 2020;77:101644. doi:10.1016/j.rasd.2020.101644.

85. Venkatesh M, Mukherjee S, Wang H, Li H, Sun K, Benechet AP, Qiu Z, Maher L, Redinbo MR, Phillips RS, Fleet JC, Kortagere S, Mukherjee P, Fasano A, Le Ven J, Nicholson JK, Dumas ME, Khanna KM, Mani S. Symbiotic bacterial metabolites regulate gastrointestinal barrier function via the xenobiotic sensor PXR and Toll-like receptor 4. Immunity. 2014 Aug 21;41(2):296-310. doi: 10.1016/j.immuni.2014.06.014. Epub 2014 Jul 24. PMID: 25065623; PMCID: PMC4142105.

86. Ventura G, Calvano CD, Porcelli V, et al. Phospholipidomics of peripheral blood mononuclear cells (PBMCs): the tricky case of children with autism spectrum disorder

(ASD) and their healthy siblings. Anal Bioanal Chem. 2020;412(25):6859-6874. doi:10.1007/s00216-020-02817-z

87. Wang, H.; Liang, S.; Wang, M.; Gao, J.; Sun, C.; Wang, J.; Xia, W.; Wu, S.; Sumner, S.J.; Zhang, F.; et al. Potential Serum Biomarkers from a Metabolomics Study of Autism. *J. Psychiatry Neurosci.* **2016**, *41*, 27–37.

88. Wang, T., Chen, B., Luo, M. et al. Microbiota-indole 3-propionic acid-brain axis mediates abnormal synaptic pruning of hippocampal microglia and susceptibility to ASD in IUGR offspring. Microbiome 11, 245 (2023). https://doi.org/10.1186/s40168-023-01656-1

89. Wouter G. Touw, Jumamurat R. Bayjanov, Lex Overmars, Lennart Backus, Jos Boekhorst, Michiel Wels, Sacha A. F. T. van Hijum, Data mining in the Life Sciences with Random Forest: a walk in the park or lost in the jungle?, Briefings in Bioinformatics, Volume 14, Issue 3, May 2013, Pages 315–326, https://doi.org/10.1093/bib/bbs034

90. Yu R, Wu Z, Wang S, Zhang M, Zhou G, Li B. Isolation, identification and characterization of propionic acid bacteria associated with autistic spectrum disorder. Microb Pathog. 2020 Oct;147:104371. doi: 10.1016/j.micpath.2020.104371. Epub 2020 Jul 4. PMID: 32634613. \

91. Zhao, ZH., Xin, FZ., Xue, Y. et al. Indole-3-propionic acid inhibits gut dysbiosis and endotoxin leakage to attenuate steatohepatitis in rats. Exp Mol Med 51, 1–14 (2019). https://doi.org/10.1038/s12276-019-0304-5

92. Zheng Y, Bek MK, Prince NZ, Peralta Marzal LN, Garssen J, Perez Pardo P, Kraneveld AD. The Role of Bacterial-Derived Aromatic Amino Acids Metabolites Relevant in Autism Spectrum Disorders: A Comprehensive Review. Front Neurosci. 2021 Oct 21;15:738220. doi: 10.3389/fnins.2021.738220. PMID: 34744609; PMCID: PMC8568365.

93. Zheng Y, Prince N, van Hattem C, Garssen J, Pardo PP, Kraneveld AD. The interaction between intestinal bacterial metabolites and phosphatase and tensin homolog in autism spectrum disorder. Mol Cell Neurosci. 2023 Mar;124:103805. doi: 10.1016/j.mcn.2022.103805. Epub 2022 Dec 30. PMID: 36592799.

94. Zhou Y, Bi Z, Hamilton MJ, Zhang L, Su R, Sadowsky MJ, Roy S, Khoruts A, Chen C. *p*-Cresol Sulfate Is a Sensitive Urinary Marker of Fecal Microbiota Transplantation

1070 and Antibiotics Treatments in Human Patients and Mouse Models. *International Journal*

1071 *of Molecular Sciences*. 2023; 24(19):14621. https://doi.org/10.3390/ijms241914621