Full Title:

Forage preference in two geographically co-occurring fungus gardening ants: a dietary DNA

approach

Short Title:

Dietary DNA techniques for forage identification in fungus gardening ants

Authors:

Matthew Richards-Perhatch V, Elizabeth Boshers, Matthew Greenwold, Jon N. Seal

Correspondence:

Jon N. Seal, Department of Biology, University of Texas at Tyler, 3900 University Blvd, Tyler,

TX 75799, USA trachymyrmex@gmail.com

ORCID ID:

Matthew Richards-Perhatch V (0009-0004-9403-0717)

Elizabeth Boshers (0000-0002-9166-1407)

Matthew Greenwold (0000-0002-7856-5162)

Jon Seal (0000-0001-8013-1438)

Abstract

Traditional methods of forage identification are impractical with non-leafcutting fungus gardening

ants, making diet-related ecological and life history questions difficult to study. To address this

limitation, we utilized dietary DNA metabarcoding on excavated ant fungus gardens to generate

forage diversity metrics for the two co-occurring species *Trachymyrmex septentrionalis* and

Mycetomoellerius turrifex. Ten fungus garden samples from each species were collected from a

60x70 m plot in East Texas. Each of the colonies we sampled was paired with a colony from the

other species within 3 m of it. Plant forage diversity was assessed with chloroplast *trnL* primers,

and insect frass forage diversity was assessed with mitochondria *COI* primers. DNA

metabarcoding identified a total of 44 plant taxa across all samples, but performed poorly when

characterizing foraged insect frass. Plant beta diversity was significantly different between the

gardens of *T. septentrionalis* and *M. turrifex* colonies, as well as paired colonies. Colony pairs

also had significantly different plant alpha diversity. This indicates that diet preference is likely

driven both by ant species-specific plant preference, and colony location-specific plant resource

availability. Overall, our results show that dietary DNA techniques are a promising tool for the

identification of plant forage in ant fungus gardens, enabling the study of future diet-based

ecological and natural history questions.

Introduction

Fungus Gardening Ants

Fungus-gardening ants are a New World monophyletic group characterized by their cultivation,

and obligate consumption, of fungi. These ants use material foraged by their colony as a

substrate to grow their fungus garden, feeding off the fungus instead of directly consuming the

forage themselves. The more basal fungus-gardening ants forage insect-feces, dry plant debris,

leaf-litter, and seed fragments [1,2]. More derived fungus gardening ants practice "higher

agriculture" and can be subdivided into two main groups: the leafcutting ants, and the non-

leafcutting higher agriculture ants [3,4]. The most specialized of these two groups, the

leafcutting ants, are known to forage extensively on freshly cut vegetative material; while the

non-leafcutting higher agriculture ants often forage considerable amounts of freshly fallen flower

petals, soft leaf tissue, and fruit, in addition to material similar to that collected by more basal

fungus gardening species [2,5].

Traditional Methods of Diet Identification

Visual identification of material carried by foraging workers is a commonly utilized technique in

determining ant diet and forage preferences. However, reliance on this technique presents

several issues. Visual forage identification is time consuming, relies heavily on taxonomic

expertise, and requires foragers to be conspicuous enough for observation [6,7]. In addition,

observing individual foragers only offers a small snapshot of total colony dietary breadth, as a

single colony may send out hundreds or even thousands of workers to forage [8].

Among fungus gardening ants, visual identification of forage material is most practical with the

leafcutting genera *Atta* and *Acromyrmex*, as their foraging strategies predispose them to easy

observation. The characteristic foraging trails created by *Atta* and *Acromyrmex* colonies can

often be cleanly followed from the vegetation they are cutting to their nest/tunnel entrances,

allowing for species level identification of their forage material [9–11]. Unfortunately, visual

identification of forage material falls short with fungus gardening ant species outside of *Atta* and

Acromyrmex. Most non-leafcutting fungus gardening ants do not create true foraging trails,

making it difficult and very time consuming to accurately trace the species of vegetation that

they are foraging from [12]. The foragers of non-leafcutting fungus gardening ants are also

usually smaller and more cryptic than those of leafcutting ants. Additionally, some foragers may

collect items that are unsuitable, if not toxic, and deposit them in their nests but not the fungus

gardens [41]. Thus observations on forager behavior alone are not likely to depict a

comprehensive overview of what ants are collecting and incorporating into their fungus garden.

These factors have contributed to a deficit of literature regarding the diet composition of non-

leafcutting fungus gardening ants, with the little information available primarily consisting of a

handful of forage material observations, often originally published in conjunction with the original

natural history descriptions of the ants [1,2]. Clearly, relying solely on visual identification of

forage material is insufficient for the study of non-leafcutting fungus gardening ants.

Dietary DNA

Dietary DNA (dDNA) techniques present a possible alternative to visual forage identification.

Metabarcoding techniques based on dDNA, DNA extracted from an organism's gut, stomach, or

fecal matter, have emerged as an accurate and resource-effective way to determine animal

diets [7,13]. Previous studies, in multiple animal groups, have successfully utilized dDNA

analysis to determine diet composition when direct observations of feeding behavior and/or

morphological identification of forage material was insufficient [7,13].

Dietary DNA analysis has only been used to examine ant diet composition in a handful of

studies, with even fewer of these studies having attempted to determine broad diet composition

[8,14,15]. One factor complicating dDNA ant research is a yet unidentified compound in the

gaster of adult ants that has been found to inhibit PCR, potentially impacting gut content dDNA

analysis in adult ants [15,16]. As an additional complication, in many species adult ants are

unable to digest solid food, instead feeding it to their brood. The adult ants then consume the

digested liquid nutrients via trophallaxis from their larvae [8,17]. Wulff et al. (2021) leveraged

this adult-larvae digestion dynamic to their advantage when using dDNA analysis to investigate

the diet composition of *Solenopsis invicta* [8]. They did so by sequencing the gut content of *S.*

invicta larvae instead of adults, as the larvae act as a centralized depository of colony prey and

are not known to produce PCR inhibitors [8,15,16].

A similarly innovative approach could be used to analyze the diet composition of fungus

gardening ants, since their fungus gardens also act as a centralized depository for colony

forage. The fungus gardens cultivated by fungus gardening ants function as an external

ancillary digestive system for the colony [2,17–19]. Within this ancillary digestive system, forage

materials used as fungal substrate become highly degraded, due both to physical mastication

by the ants and enzymatic breakdown by the fungus. This degradation necessitates the use of

specialized genetic techniques, like dDNA metabarcoding, that are designed to work with highly

degraded staples [2,5,20].

Evaluation of Diet and Study Species

Using dDNA metabarcoding techniques to identify fungal substrate in ant gardens could be

used to answer a number of ecological and natural history questions. Dietary DNA

metabarcoding has been successfully used to detect niche partitioning, and overlap, in several

animal species [21–25]. These techniques have also been used to understand seasonal and/or

biogeographic based shifts in animal diets [26–32].

Dietary differences have been investigated (without the use of dDNA methods) in several

species of fungus gardening leafcutting ants. In a dry tropical forest in Costa Rica, the co-

occurring leafcutter species *Atta cephalotes* and *Acromyrmex octospinosus* were found to

forage almost entirely different resources, with the only exception being they both foraged

flowers [33]. A similar trend was observed in Panama, with *Atta* species foraging primarily on

tree leaves and co-occurring *Acromyrmex* species foraging primarily on flowers and herbal

leaves [34]. To our knowledge, no previous studies have examined differences in diet between

non-leafcutting fungus gardening ants.

Trachymyrmex septentrionalis and *Mycetomoellerius turrifex* are two higher agriculture non-

leafcutting fungus gardening ant species, with a partial overlap of their geographic distribution

occurring in East Texas [35,36]. The range of *T. septentrionalis* stretches from Texas to Florida,

and as far north as central Illinois and New York; while the range of *M. turrifex* extends from

Texas to northeastern Mexico and western Louisiana [36–38]. Like most non-leafcutting fungus

gardeners, specific information on the preferred forage material of *T. septentrionalis* and *M.*

turrifex is highly limited [2]. What documentation does exist indicates that they both forage

insect frass and plant material, but only *T. septentrionalis* has been observed cutting/foraging

fresh vegetation [2,39–42]. Their geographic range overlap and the similarity of their

documented forage material makes these two species excellent subjects to test the efficacy of

using dDNA analysis to study diet in non-leafcutting fungus gardening ants.

The primary objective of this study was to explore whether dDNA metabarcoding could be used

to identify and describe the diversity of plant material and/or insect frass used as substrate in

the fungus gardens of co-occuring *T. septentrionalis* and *M. turrifex* colonies. The secondary

objective of this study was to evaluate whether diversity metrics based on identified fungal

substrate were sufficient to detect differences in diet between *T. septentrionalis* and *M. turrifex*

colonies.

Materials and Methods

Sample Collection

The fungus gardens of ten *T. septentrionalis* and ten *M. turrifex* colonies were sampled from an

approximately 60x70 m plot on the University of Texas at Tyler campus in Tyler, Texas, USA.

Sampling occurred from May 16-26, 2023. Each of the ten *M. turrifex* colonies we sampled was

paired with a *T. septentrionalis* colony within 3 m of it. These colony pairs, composed of one *M.*

turrifex colony and one nearby *T. septentrionalis* colony, were sampled concurrently. Each

colony was only sampled once.

Fungus gardens were non-destructively sampled from each colony by careful excavation of the

topmost chamber containing an appreciable amount of fungus. Sampling was focused on

chambers closer to the surface since deeper chambers were unlikely to contain fungus until

later in the season when the upper chambers are too hot for fungus garden growth. Sterilized

tools were used when removing fungus gardens from the chamber to minimize DNA

contamination from environmental sources and previously sampled colonies. Excavated fungus

garden samples were placed in sterile vials containing 95% ethanol and stored in a -80°C

freezer.

Sample Metabarcoding

Metabarcoding was conducted by Jonah Ventures (https://jonahventures.com/) using

chloroplast *trnL* (forward primer c: 5' CGAAATCGGTAGACGCTACG 3', reverse primer h: 5'

CCATTGAGTCTCTGCACCTATC 3') and mitochondrial cytochrome c oxidase subunit I (*COI*)

(forward primer ZBJ-ArtF1c: 5' AGATATTGGAACWTTATATTTTATTTTTGG 3', reverse primer

ZBJ-ArtR2c: 5' WACTAATCAATTWCCAAATCCTCC 3') primers [43,44]. Individual fungus

garden samples were shipped overnight in an ice box to Jonah Ventures in Boulder, CO.

At Jonah Ventures, frozen samples were thawed for 1-2 hours before processing. Sample

barcodes were recorded and assigned a well within the 96 well plate or numbered extraction

tube. Under a laminar flow hood, sterile cotton swabs (Fisher, Catalog No. 22-363-173) were

coated with sediment matter, and the swabs were placed in the corresponding extraction plate

or tube. Sterile tweezers and pliers were used to handle cotton swabs and remove the wooden

ends of the cotton swab before extraction. Plates or tubes were immediately processed or

stored in -20°C until the extraction process could be performed.

Genomic DNA from each fungus garden sample was extracted using the DNeasy 96 PowerSoil

Pro Kit (384) (Catalog No. 47017) according to the manufacturer's protocol. Genomic DNA was

eluted into 100 µl and frozen at -20°C.

For *trnL* genetic analysis, a portion of the chloroplast *trnL* intron was PCR amplified from each

genomic DNA sample using the c and h trnL primers. Both forward and reverse primers also

contained a 5' adaptor sequence to allow for subsequent indexing and Illumina sequencing.

Each 25 µL PCR reaction was mixed according to the Promega PCR Master Mix specifications

(Promega, Catalog No. M5133, Madison, WI) which included 0.4 µM of each primer and 1 µl of

gDNA. DNA was PCR amplified using the following conditions: initial denaturation at 94°C for 3

minutes, followed by 40 cycles of 30 seconds at 94°C, 30 seconds at 55°C, and 1 minute at

72°C, and a final elongation at 72°C for 10 minutes.

For *COI* genetic analysis, a portion of the *COI* gene was PCR amplified from each genomic

DNA sample using the ZBJ-ArtF1c and ZBJ-ArtR2c primers. Both forward and reverse primers

also contained a 5' adaptor sequence to allow for subsequent indexing and Illumina sequencing.

Each 25 µL PCR reaction was mixed according to the Promega PCR Master Mix specifications

(Promega, Catalog No. M5133, Madison, WI) which included 12.5 µl Master Mix, 1.0 µM of each

primer, 3.0 µl of gDNA, and 7.5 µl DNase/RNase-free H_2O. DNA was PCR amplified using the

following conditions: initial denaturation at 94°C for 5 minutes, followed by 45 cycles of 30 s at

94°C, 45 s at 45°C, and 45 s at 72°C, and a final elongation at 72°C for 10 minutes.

To determine amplicon size and PCR efficiency, each reaction was visually inspected using a

2% agarose gel with 5 µl of each sample as input. Amplicons were then cleaned by incubating

amplicons with Exo1/SAP for 30 minutes at 37°C followed by inactivation at 95°C for 5 minutes

and stored at -20°C.

A second round of PCR was performed to complete the sequencing library construct, appending

with the final Illumina sequencing adapters and integrating a custom sample-specific 12-

nucleotide index sequence (one index read per amplicon) generated by Jonah Ventures. The

indexing PCR included Promega Master mix, 0.5 µM of each primer and 2 µl of template DNA

(cleaned amplicon from the first PCR reaction) and consisted of an initial denaturation of 95°C

for 3 minutes followed by 8 cycles of 95°C for 30 seconds, 55°C for 30 seconds and 72°C for 30

seconds.

Final indexed amplicons from each sample were cleaned and normalized using SequalPrep

Normalization Plates (Life Technologies, Carlsbad, CA). 25 µl of PCR amplicon was purified and

normalized using the Life Technologies SequalPrep Normalization kit (Catalog No. A10510-01)

according to the manufacturer's protocol. Samples were then pooled together by adding 5µl of

each normalized sample to the pool.

Sample library pools were sent for sequencing on an Illumina MiSeq (San Diego, CA) at the

Texas A&M Agrilife Genomics and Bioinformatics Sequencing Core facility using the v2 500-

cycle kit (Catalog No. MS-102-2003). A PhiX concentration of 15% was used. As a quality

control measure, a bioanalyzer chip was used for concentration adjustment at the sequencing

center prior to sequencing.

Raw sequence data were demultiplexed using pheniqs v2.1.0, enforcing strict matching of

sample barcode indices (i.e, no errors) [45].

Sequence Processing

After receiving the demultiplexed data in the form of paired end FASTQ files from Jonah

Ventures, the files were imported into Qiime2 for further processing (Figure 1) [46].

Fig 1. Flowchart of each major bioinformatic step in the development of our analysis

metrics.

The Qiime2 plugin "cutadapt" was used to remove primers from the sequences. The Qiime2

plugin "dada2" was used to denoise the *trnL* and *COI* sequences, truncate the *COI* sequences,

and merge the forward and reverse *COI* sequence pairs [47]. Only the forward *trnL* sequences

were used because of the short *trnL* amplicon length (143 bp). The *trnL* sequences were not

truncated because their untruncated median quality scores were 30 or greater. The *COI*

sequences were truncated where their median quality scores dropped below 30 (forward: 100

bp, reverse: 87 bp). Rarefaction of the sequences was conducted using Qiime2's 'feature-table

rarefy' command. A sampling depth of 9677 was used for *trnL* sequences (Figure 2). Due to

low *COI* sequencing depth (<1000) in the majority of samples, downstream analysis was not

conducted for *COI* outside of taxonomic classification (Figure 3).

Fig 2. Rarefaction curves for *trnL* sequences. The dashed vertical line shows the chosen

trnL rarefaction depth of 9677.

Fig 3. Rarefaction curves for *COI* sequences. The lower left corner of the plot has been

magnified to show greater detail.

Qiime2's "RESCRIPt" plugin was used to create two reference databases from NCBI; one

specific to *trnL* and a second specific to arthropod *COI* [48,49]. Initial reference pools were

constructed from these downloaded databases via primer-pair search using Qiime2's 'feature-

classifier extract-reads' command. These initial pools were both expanded twice using

RESCRIPt's 'extract-seq-segments' command. During reference pool expansions of the *trnL*

specific database a '--p-perc-identity' value of 0.7 was used. During reference pool expansions

of the *COI* specific database a '--p-perc-identity' value of 0.9 was used. These '--p-perc-identity'

values were chosen because at the end of their respective reference pool expansions they had

extracted sequence segments from ~70% of their original query sequences [49]. Dereplication

during each of these steps was performed using the '--p-mode super' modifier in RESCRIPt's

'dereplicate' command. These expanded reference pools were then used to construct two

classifiers, one for *trnL* and one for *COI*, using RESCRIPt's 'evaluate-fit-classifier' command

(Figure S1).

Taxonomic assignment for *trnL* and *COI* sequences was conducted using the custom

RESCRIPt classifiers described above, in combination with Qiime2's 'feature-classifier classify-

sklearn' command.

Statistical Analysis

Once taxonomic assignment was completed, the Qiime2 feature and metadata files were

imported into R and used to create two phyloseq objects using the R packages "qiime2R" and

"phyloseq" [50–52]. An independent phylogenetic tree was constructed by grafting our classified

trnL sequences onto the GBOTB.extended.WP.tre megatree, reported by Jin and Qian (2022),

using the 'phylo.maker' command from the R package "U.PhyloMaker" [53,54]. Six classified

trnL sequences failed to graft to the tree due to low taxonomic resolution and were discarded.

The resulting phylogenetic tree was converted to a phylo class object using the

'phyloseq::read_tree' command; multichotomies were then resolved using the 'multi2di'

command from the R package "ape" [55]. This tree was merged with our phyloseq object using

the 'phyloseq::merge_phyloseq' command. This updated phyloseq object was used to calculate

alpha diversity for each colony, in the form of Faith's phylogenetic diversity (Faith's PD) scores

and species richness, using the 'calculatePD' command in the R package "biomeUtils" [56].

Beta diversity between each colony was also calculated from the updated phlyoseq object, in

the form of unweighted unique fraction metric (uUniFrac) dissimilarity distances, using the

'phyloseq::distance' command.

These specific diversity metrics were chosen for two primary reasons: 1) their calculations are

based on relative frequency of occurrence (binary presence/absence) of the sequenced taxa,

and 2) their calculations take into account the relative relatedness of the sequenced taxa.

The number of sequencing reads from dDNA often does not accurately represent the biomass

of consumed material [6,13]. This is particularly relevant in studies like ours, where sequenced

material consists of a broad diversity of partially digested multi-cellular samples. It is likely that

digestive processes, the order items are consumed, primer biases, and taxonomic biases in the

reference database influence both the sequencing outcomes and taxonomic classification

[6,57,58]. Due to the known impact of these factors on estimations of abundance, only metrics

based on binary presence/absence of the sequenced taxa (i.e. uUniFrac and Faith's PD) were

used.

Another issue arises from the various states of DNA degradation present in our samples; as

sequences originating from the same species, or even the same individual, may end up

classified at different taxonomic levels. For example, if one group of sequences are classified as

Fagaceae and another group of sequences from the same sample are classified as *Fagaceae*

Quercus, it is not currently possible to determine if this indicates the presence of two separate

genera of *Fagaceae*, or if both sets of sequences come from the same genera. This can result

in the artificial inflation of identified taxa, as sequences from a single taxon could be functionally

double counted. Some previous dDNA studies have attempted to account for this issue by

compressing their post-classified dDNA taxa into diet events: instances where there is

unequivocal evidence from dDNA analysis that an organism consumed at least one individual of

a species [8,59]. While this approach does minimize double counting of taxa, it increases the

risk of undercounting dietary events and can obscure dietary events of taxa unable to be

identified to the species level [59]. To avoid the drawbacks of using a diet event based

approach, we instead utilized metrics that included the phylogenetic relationship of classified

taxa in their calculations (i.e. uUniFrac and Faith's PD). Using these phylogeny-weighted

metrics allows sequences classified at different taxonomic levels to be assessed as separate

taxa, while minimizing the impact of possible double counts.

Our analysis incorporated two main predictor variables: colony ant species (*T. septentrionalis*

and *M. turrifex*) and colony pair ID (Pair-2, Pair-8, Pair-12, Pair-14, Pair-17, Pair-20, Pair-28,

Pair-30, Pair-44, and Pair-50). We tested for differences in fungal substrate plant taxa alpha

diversity (Faith's PD and species richness) and beta diversity (uUniFrac dissimilarity distance).

Faith's PD and species richness distributions were tested for normality via the Shapiro-Wilk

normality tests by using the 'shapiro.test' command in R. To analyze Faith's PD and species

richness against each of our predictor variables, several analysis of variance (ANOVA) analyses

were conducted using the 'aov' command in R.

Permutational analysis of variance (PerMANOVA) analyses were conducted using the 'adonis2'

commands in the R package "vegan" [60]. Our PerMANOVA analyses used 9,999 permutations.

Data visualization was performed with the R packages "tidyverse", "ggtree", and "ggrepel" [61–

63].

Results

Plant Taxa Identified from Fungus Garden Samples

Across all sampled fungus gardens, a total of 44 plant classifications were identified from *trnL*

sequences (Figure 4). Fungal substrate plant taxa composition was derived from 32 plant

classifications for *T. septentrionalis* and 33 plant classifications for *M. turrifex*. There was

considerable overlap in plant taxa composition between the two species, with 21 plant

classifications present in both *T. septentrionalis* and *M. turrifex* colonies. 11 plant classifications

were identified only from *T. septentrionalis* colonies and 12 plant classifications were identified

only from *M. turrifex* colonies.

Fig 4. Phylogenetic tree and binary heat map of plant taxa identified from fungus garden

samples. Each branch end is linked to a row in the binary heat map, showing

presence/absence of each respective plant classification identified in the *M. turrifex* (Mt) and *T.*

septentrionalis (Ts) fungus garden samples. Cells with a black dot indicate the plant

classification was present in both colonies from that pair.

In *T. septentrionalis* colonies, the most frequently occurring plant classifications (occurred in

≥50% of *T. septentrionalis* samples) were: *Fagaceae*, *Magnoliopsida*, *Rubiaceae*, *Asteraceae*,

and *Fagales.* For *M. turrifex* colonies, the most frequently occurring plant classifications

(occurred in ≥50% of *M. turrifex* samples) were: *Fagaceae*, *Oxalis* (*Oxalidaceae*),

Magnoliopsida, *Rubiaceae*, *Fagales*, *Acalypha* (*Euphorbiaceae*), and *Poaceae*.

Plant Taxa Alpha Diversity

ANOVA analyses were used to determine if Faith's PD or species richness of fungal substrate

plant taxa differed significantly between colonies based on colony ant species or colony pair ID.

Trachymyrmex septentrionalis and *M. turrifex* fungus gardens did not have significantly different

Faith's PD ($F_{1,18}$ = 0.045, p = 0.834), nor did they have significantly different species richness

($F_{1,18}$ = 0.223, p = 0.643). There were, however, significant differences in Faith's PD ($F_{9,10}$ =

3.921, p = 0.022) and species richness ($F_{9,10}$ = 3.105, p = 0.046) between colony pairs.

Plant Taxa Beta Diversity

A PerMANOVA analysis was used to determine if the uUniFrac dissimilarity distances of fungal

substrate plant taxa were significantly different between colonies based on colony ant species

or colony pair ID. There was a significant difference found between colony ant species and the

uUniFrac dissimilarity distances of their plant taxa (p = 0.046) (Figure 5). There was also a

significant difference between colony pairs and the uUniFrac dissimilarity distances of their plant

taxa (p = 0.014).

Fig 5. NMDS plot showing the uUniFrac dissimilarity of plant taxa identified from fungus

garden samples. Colony pairs are connected by grey lines.

Arthropod Taxa Identified from Fungus Garden Samples

A total of six arthropod classifications were identified from *COI* sequences. All colony fungus

garden samples contained *Insecta* sequences. Only four samples contained sequences

identified to a lower taxonomic level than *Insecta*: *M. turrifex* colony 44's fungus garden

contained *Carabidae* (*Coleoptera*) and *Lepidoptera* sequences, *M. turrifex* colony 14's fungus

garden contained *Anomoea laticlavia* (*Coleoptera Chrysomelidae*) sequences, *M. turrifex* colony

20's fungus garden contained *Coleoptera* sequences, and *T. septentrionalis* colony 112's

fungus garden contained *Hemiptera* sequences.

Discussion

Dietary DNA techniques successfully identified plant and arthropod taxa present in the fungus

gardens of *T. septentrionalis* and *M. turrifex*. The dDNA analysis was particularly successful

with the *trnL* primer, as it identified a relatively large number of plant taxa, many of them to a

relatively high level of taxonomic resolution. These results could be particularly impactful

considering the literature deficit of specific plant taxa known to be foraged by non-leafcutting

fungus gardening ants. Only a handful of previous studies make any mention of specific foraged

plant taxa, with most only describing the general type of material the ants were collecting such

as flower petals or dead leaves [1,2,40-42,64–68].

Colony ant species appeared to be a significant factor in beta diversity metrics, but not alpha

diversity metrics, of fungal substrate plant taxa. Similarly, Colony pair appeared to be a

significant factor in both plant taxa beta diversity metrics and alpha diversity metrics. These

results suggest that diet preference is likely driven both by ant species-specific plant preference

and colony location-specific plant resource availability. The presence of dietary differences

between *T. septentrionalis* and *M. turrifex* colonies on our site appears similar to short-term

observations of dietary differences made on co-occurring *Atta* and *Acromyrmex* colonies

[33,34].

The plant taxa we identified from *T. septentrionalis* and *M. turrifex* fungus garden sequences

were consistent with the limited existing descriptions of forage material collected by these ant

species. One previous study observed *M. turrifex* collecting *Sorghum halepense* and *Solidago*

sp., both of whose families, *Poaceae* and *Asteraceae* respectively, were present in our samples

[40]. Another previous study suggested that *T. septentrionalis* collects *Quercus* (*Fagaceae*),

which was also present in our samples [41].

If we include descriptions of known forage material from other higher agriculture non-leafcutting

ants, we find additional overlap between our identified plant taxa and that of previous studies.

Lauraceae, *Rosaceae*, *Rubiaceae*, *Poaceae*, and *Euphorbiaceae* are all plant families which

were identified in our samples, and are known to be collected by or, in the case of lab colonies,

accepted by higher agriculture non-leafcutting ants [64,66]. Leafcutting species like *Atta texana*

are known to forage on *Pinus* (*Pinaceae*) needles, but our study is the first published record of a

non-leafcutting species collecting *Pinus* [69]. The ants may be collecting pieces of male pine

cones, as this has been observed in the field and laboratory during early spring (March-April)

(JN Seal, unpublished observations).

A notable drawback of only using dDNA analysis on sampled fungus gardens is that dDNA

analysis alone cannot determine the state plant material was in when it was collected by the

ants. Some higher agriculture fungus gardeners outside of *Atta* and *Acromyrmex*, like *T.*

septentrionalis, are known to cut fresh vegetative material to use as fungus garden substrate,

while others do not [2,41]. In our study, we could not determine whether the detected plant

sequences originated from the ants foraging fresh cut plant material, plant detritus, or frass. We

also could not determine which part of the plant they collected (i.e. leaves, flowers, or sticks).

The inability to detect which state plant material was collected in could obscure differences in

foraging behavior between these two ant species. For example, one ant species may forage on

Quercus (*Fagaceae*) catkins, while a second ant species may forage caterpillar frass that

contains *Quercus* leaves. Similarly, one ant species may cut soft *Quercus* vegetation, whereas

a second ant species might only take dead vegetation that has fallen from the tree. In all of

these examples the ant's fungus gardens would contain *Quercus* sequences, but dDNA

analysis would be unable to differentiate the state the material was collected in. Additionally,

since the number of sequencing reads cannot be reliably used as a quantitative measure of the

biomass of digested material, we cannot use dDNA to determine if colonies are foraging

different amounts of the same material [6].

While our study establishes dDNA analysis as a potential tool for identification of fungus garden

substrate, we also identified several limitations that should be accounted for in future studies

attempting to use similar techniques. One issue was the low taxonomic resolution of some plant

sequences. While we were able to successfully identify 44 plant classifications in the fungus

gardens using our selected *trnL* primer, unidentified *Magnoliopsida* had the sixth highest

absolute read abundance post-rarefaction, indicating that a number of plant species may be

present in the fungus gardens that our *trnL* primer was unable to identify to an informative

taxonomic level. In future studies this could be mitigated, at least in part, by using multiple plant

primers designed for different target regions [23,70,71]. Another issue was the misidentification

of plant sequences at the species level. While some identified plant species, like *Desmodium*

ciliare (*Fabaceae*), are known to occur on our study site; other species, such as *Salix hukaoana*

(*Salicaceae*), are not (though other species of *Salix* are certainly in the general vicinity). These

errors were likely the result of our classifier misidentifying sequences as related taxa due to an

incomplete reference database. Ideally, future studies could construct a region-specific

taxonomic library of plant sequences to be used when training their classifier [72]. Barcoding

insect species present in a study site would be more difficult, as they are more mobile and can

inhabit above both and below ground areas during different life stages.

Dietary DNA analysis alone is not capable of distinguishing *trnL* sequences from plant material

foraged directly by the ants and *trnL* sequences from insect frass foraged by the ants. Since

Trachymyrmex septentrionalis and *M. turrifex* both readily forage insect frass containing plant

material, plant DNA sequences from foraged frass were potentially detected by the *trnL* primer

we used [13,40,42]. We had hoped to use the *COI* primer results to gain insight into what

arthropod taxa the ants were collecting frass from, but the primer performed too poorly to draw

any meaningful conclusions. Other studies attempting to determine *Lepidoptera* species from

their frass found that amounts of dry frass less than 5 mg may be too small to detect insect DNA

sequences [73]. This insect sequence detectability cutoff was likely higher in our samples due to

the physical and enzymatic breakdown of foraged material occurring in the fungus gardens,

making it difficult to detect frass-producing-insect DNA [2]. Another potential confounding issue

could be a decrease in PCR efficiency due to our use of a 1-step PCR approach versus a 2-step

PCR approach for sequencing (See Methods Bohmann et al. (2022) for a discussion on the

topic) [74].

Many of the issues we discussed can be mitigated by taking the limitations of dDNA analysis

into account when designing future studies. Our recommendations for future studies include: 1)

using multiple plant primers to maximize taxonomic coverage, 2) constructing a region-specific

taxonomic library of plant sequences, 3) sampling at several different time points to allow

detection of temporal changes in fungal substrate plant composition, and 4) incorporating a

supplemental observational component to gain an understanding of what state forage material

is being collected in. Overall, we have demonstrated the efficacy of using dDNA analysis as a

tool for the taxonomic identification of forage material in fungus-gardening ants, and as a

possible tool to detect dietary differences between species. Dietary DNA analysis offers an

opportunity to improve our understanding of plant taxa foraged by non-leafcutting fungus

gardening ants, and, with appropriate supporting methods, can be applied to gain new insight

into their ecology and life history.

Acknowledgments

We would like to thank Christine Bays, Dillon Flowers, Gabriel McDanield, and Grant Rook, all

of whom took time out of their lives to help with colony excavation and sampling.

Data Availability

The raw sequences for all samples have been deposited with links to BioProject accession

number PRJNA1124954 in the NCBI BioProject database

(https://www.ncbi.nlm.nih.gov/bioproject/PRJNA1124954). The Qiime2, RESCRIPt, and R code

used during this project can be found at https://github.com/mrichardsperhatchv/FORAGE-

PREFERENCE-IN-TWO-GEOGRAPHICALLY-CO-OCCURRING-FUNGUS-GARDENING-

ANTS.

Supplemental

Fig S1. The F-measure of our classifiers being used on their own training dataset.

References

1. Leal IR, Oliveira PS. Foraging ecology of attine ants in a Neotropical savanna: seasonal use of fungal substrate in the cerrado vegetation of Brazil. Insectes sociaux. 2000;47: 376–382. doi:10.1007/PL00001734

2. De Fine Licht HH, Boomsma JJ. Forage collection, substrate preparation, and diet composition in fungus-growing ants. Ecological Entomology. 2010;35: 259–269. doi:10.1111/j.1365-2311.2010.01193.x

3. Solomon SE, Rabeling C, Sosa-Calvo J, Lopes CT, Rodrigues A, Vasconcelos HL, et al. The molecular phylogenetics of *Trachymyrmex Forel* ants and their fungal cultivars provide insights into the origin and coevolutionary history of 'higher-attine' ant agriculture. Systematic entomology. 2019;44: 939–956. doi:10.1111/syen.12370

4. Schultz TR, Brady SG. Major evolutionary transitions in ant agriculture. Proceedings of the National Academy of Sciences of the United States of America. 2008;105: 5435–5440. doi:10.1073/pnas.0711024105

5. De Fine Licht HH, Schiøtt M, Mueller UG, Boomsma JJ. Evolutionary transitions in enzyme

activity of ant fungus gardens. Evolution. 2010;64: 2055–2069. doi:10.1111/j.1558-
5646.2010.00948.x

6.   Nielsen JM, Clare EL, Hayden B, Brett MT, Kratina P. Diet tracing in ecology: Method
comparison and selection. Methods Ecol Evol. 2018;9: 278–291. doi:10.1111/2041-
210X.12869

7.   Sousa LL, Silva SM, Xavier R. DNA metabarcoding in diet studies: Unveiling ecological
aspects in aquatic and terrestrial ecosystems. Environmental DNA. 2019;1: 199–214.
doi:10.1002/edn3.27

8.   Wulff JA, Kjeldgaard MK, Eubanks MD, Sword GA. From the bellies of babes: a larval-
based approach to ant molecular gut content analysis. Insectes sociaux. 2021;68: 245–
253. doi:10.1007/s00040-021-00822-8

9.   Mintzer A. Foraging activity of the Mexican leafcutting ant *Atta mexicana* (F. Smith), in a
sonoran desert habitat (*Hymenoptera*, *Formicidae*). Insectes sociaux. 1979;26: 364–372.
doi:10.1007/BF02223555

10.   Berish CW. Leaf-Cutting Ants (*Atta cephalotes*) Select Nitrogen-Rich Forage. American
Midland Naturalist. 1986;115: 268. doi:10.2307/2425863

11.   Mintzer A. Diet of the Leafcutting Ant, *Atta Mexicana* (*Hymenoptera*: *Formicidae*), in a
Sonoran Desert Habitat. Journal of the Arizona-Nevada Academy of Science. 1995.

12.   Weber NA. Gardening ants the attines. American Philosophical Society. 1972; 146.

13.   Ando H, Mukai H, Komura T, Dewi T, Ando M, Isagi Y. Methodological trends and
perspectives of animal dietary studies by noninvasive fecal DNA metabarcoding.
Environmental DNA. 2020;2: 391–406. doi:10.1002/edn3.117

14.   Muilenburg VL, Goggin FL, Hebert SL, Jia L, Stephen FM. Ant predation on red oak borer
confirmed by field observation and molecular gut-content analysis. Agricultural and forest
entomology. 2008;10: 205–213. doi:10.1111/j.1461-9563.2008.00371.x

15.   Penn HJ, Chapman EG, Harwood JD. Overcoming PCR Inhibition During DNA-Based Gut
Content Analysis of Ants. Environmental Entomology. 2016;45: 1255–1261.
doi:10.1093/ee/nvw090

16.   Gadau J. DNA isolation from ants. Cold Spring Harbor Protocols. 2009;2009:
pdb.prot5245. doi:10.1101/pdb.prot5245

17.   Holldobler B, Wilson EO. The Ants. 1st ed. Cambridge, Mass: Belknap Press: An Imprint of
Harvard University Press; 1990.

18.   De Fine Licht HH, Boomsma JJ, Tunlid A. Symbiotic adaptations in the fungal cultivar of
leaf-cutting ants. Nature Communications. 2014;5: 5675. doi:10.1038/ncomms6675

19.   Seal JN, Schiøtt M, Mueller UG. Ant-fungus species combinations engineer physiological
activity of fungus gardens. Journal of Experimental Biology. 2014;217: 2540–2547.
doi:10.1242/jeb.098483

20.   Moller IE, De Fine Licht HH, Harholt J, Willats WGT, Boomsma JJ. The dynamics of plant
551   cell-wall polysaccharide decomposition in leaf-cutting ant fungus gardens. Plos One.
552   2011;6: e17506. doi:10.1371/journal.pone.0017506

553   21.   Burgar JM, Murray DC, Craig MD, Haile J, Houston J, Stokes V, et al. Who's for dinner?
554   High-throughput sequencing reveals bat dietary differentiation in a biodiversity hotspot
555   where prey taxonomy is largely undescribed. Mol Ecol. 2014;23: 3605–3617.
556   doi:10.1111/mec.12531

557   22.   Kartzinel TR, Chen PA, Coverdale TC, Erickson DL, Kress WJ, Kuzmina ML, et al. DNA
558   metabarcoding illuminates dietary niche partitioning by African large herbivores. Proc Natl
559   Acad Sci USA. 2015;112: 8019–8024. doi:10.1073/pnas.1503283112

560   23.   Soininen EM, Gauthier G, Bilodeau F, Berteaux D, Gielly L, Taberlet P, et al. Highly
561   overlapping winter diet in two sympatric lemming species revealed by DNA metabarcoding.
562   PLoS ONE. 2015;10: e0115335. doi:10.1371/journal.pone.0115335

563   24.   Sato JJ, Shimada T, Kyogoku D, Komura T, Uemura S, Saitoh T, et al. Dietary niche
564   partitioning between sympatric wood mouse species (*Muridae*: *Apodemus*) revealed by
565   DNA meta-barcoding analysis. J Mammal. 2018;99: 952–964.
566   doi:10.1093/jmammal/gyy063

567   25.   Smith JA, Thomas AC, Levi T, Wang Y, Wilmers CC. Human activity reduces niche
568   partitioning among three widespread mesocarnivores. Oikos. 2018 [cited 21 Oct 2023].
569   doi:10.1111/oik.04592

570   26.   Aizpurua O, Budinski I, Georgiakakis P, Gopalakrishnan S, Ibañez C, Mata V, et al.
571   Agriculture shapes the trophic niche of a bat preying on multiple pest arthropods across
572   Europe: Evidence from DNA metabarcoding. Mol Ecol. 2018;27: 815–825.
573   doi:10.1111/mec.14474

574   27.   Komura T, Ando H, Horikoshi K, Suzuki H, Isagi Y. DNA barcoding reveals seasonal shifts
575   in diet and consumption of deep-sea fishes in wedge-tailed shearwaters. PLoS ONE.
576   2018;13: e0195385. doi:10.1371/journal.pone.0195385

577   28.   Forsman AM, Hoenig BD, Gaspar SA, Fischer JD, Siegrist J, Fraser K. Evaluating the
578   impacts of metabarcoding primer selection on DNA characterization of diet in an aerial
579   insectivore, the Purple Martin. Ornithology. 2021 [cited 21 Oct 2023].
580   doi:10.1093/ornithology/ukab075

581   29.   Palumbo F, Squartini A, Barcaccia G, Macolino S, Pornaro C, Pindo M, et al. A multi-
582   kingdom metabarcoding study on cattle grazing Alpine pastures discloses intra-seasonal
583   shifts in plant selection and faecal microbiota. Sci Rep. 2021;11: 889. doi:10.1038/s41598-
584   020-79474-w

585   30.   Roffler GH, Allen JM, Massey A, Levi T. Metabarcoding of fecal DNA shows dietary
586   diversification in wolves substitutes for ungulates in an island archipelago. Ecosphere.
587   2021;12. doi:10.1002/ecs2.3297

588   31.   Brun L, Schneider J, Carrió EM, Dongre P, Taberlet P, van de Waal, et al. Focal vs. fecal:
589   Seasonal variation in the diet of wild vervet monkeys from observational and DNA

metabarcoding data. Ecol Evol. 2022;12: e9358. doi:10.1002/ece3.9358

32. Wang B, Li Y, Zhang G, Yang J, Deng C, Hu H, et al. Seasonal variations in the plant diet
of the Chinese Monal revealed by fecal DNA metabarcoding analysis. Avian Res. 2022;13:
100034. doi:10.1016/j.avrs.2022.100034

33. Wetterer J. Foraging Ecology of the Leaf-Cutting Ant Acromyrmex Octospinosus in a Costa
Rican Rain Forest. Psyche. 1991;98: 361–371. doi:10.1155/1991/46737

34. Kooij PW, Liberti J, Giampoudakis K, Schiøtt M, Boomsma JJ. Differences in forage-
acquisition and fungal enzyme activity contribute to niche segregation in Panamanian leaf-
cutting ants. Plos One. 2014;9: e94284. doi:10.1371/journal.pone.0094284

35. Seal JN, Brown L, Ontiveros C, Thiebaud J, Mueller UG. Gone to Texas: phylogeography
of two (*Hymenoptera*: *Formicidae*) species along the southeastern coastal plain of North
America. Biological Journal of the Linnean Society. 2015;114: 689–698.
doi:https://doi.org/10.1111/bij.12426

36. Senula SF, Scavetta JT, Banta JA, Mueller UG, Seal JN, Kellner K. Potential Distribution of
Six North American Higher-Attine Fungus-Farming Ant (*Hymenoptera*: *Formicidae*)
Species. Journal of Insect Science (Online). 2019;19. doi:10.1093/jisesa/iez118

37. Rabeling C, Cover SP, Johnson RA, Mueller, G U. A review of the North American species
of the fungus-gardening ant genus *Trachymyrmex (Hymenoptera*: *Formicidae*). Zenodo.
2007 [cited 7 Sep 2023]. doi:10.5281/zenodo.180014

38. Sanchez-Peña SR. Some Fungus-Growing Ants (*Hymenoptera*: *Formicidae*) from
Northeastern Mexico. Florida Entomologist. 2010;93: 501–504. doi:10.1653/024.093.0404

39. Wheeler WM. The fungus-growing ants of North America. 1907.

40. Waller DA. Foraging Behavior of *Trachymyrmex turrifex* Wheeler (*Formicidae*: *Attini*).
Source: The Southwestern Naturalist. 1989;34: 271–275.

41. Seal JN, Tschinkel WR. Complexity in an obligate mutualism: do fungus-gardening ants
know what makes their garden grow? Behavioral Ecology and Sociobiology. 2007;61:
1151–1160. doi:10.1007/s00265-006-0328-4

42. Seal JN, Tschinkel WR. Food limitation in the fungus-gardening ant, *Trachymyrmex
septentrionalis*. Ecological Entomology. 2008;33: 597–607. doi:10.1111/j.1365-
2311.2008.01010.x

43. Taberlet P, Coissac E, Pompanon F, Gielly L, Miquel C, Valentini A, et al. Power and
limitations of the chloroplast trnL (UAA) intron for plant DNA barcoding. Nucleic Acids Res.
2007;35: e14. doi:10.1093/nar/gkl938

44. Razgour O, Clare EL, Zeale MRK, Hanmer J, Schnell IB, Rasmussen M, et al. High-
throughput sequencing offers insight into mechanisms of resource partitioning in cryptic bat
species. Ecol Evol. 2011;1: 556–570. doi:10.1002/ece3.49

45. Galanti L, Shasha D, Gunsalus KC. Pheniqs 2.0: accurate, high-performance Bayesian

decoding and confidence estimation for combinatorial barcode indexing. BMC
Bioinformatics. 2021;22: 359. doi:10.1186/s12859-021-04267-5

46. Bolyen E, Rideout JR, Dillon MR, Bokulich NA, Abnet CC, Al-Ghalith GA, et al.
Reproducible, interactive, scalable and extensible microbiome data science using QIIME 2.
Nature Biotechnology. 2019;37: 852–857. doi:10.1038/s41587-019-0209-9

47. Callahan BJ, McMurdie PJ, Rosen MJ, Han AW, Johnson AJA, Holmes SP. DADA2: High-
resolution sample inference from Illumina amplicon data. Nature Methods. 2016;13: 581–
583. doi:10.1038/nmeth.3869

48. Robeson MS, O'Rourke DR, Kaehler BD, Ziemski M, Dillon MR, Foster JT, et al.
RESCRIPt: Reproducible sequence taxonomy reference database management. PLoS
Computational Biology. 2021;17: e1009581. doi:10.1371/journal.pcbi.1009581

49. Robeson M. Using RESCRIPt's "extract-seq-segments" to extract reference sequences
without PCR primer pairs. 2022. Available: https://forum.qiime2.org/t/using-rescripts-
extract-seq-segments-to-extract-reference-sequences-without-pcr-primer-pairs/23618

50. McMurdie PJ, Holmes S. phyloseq: an R package for reproducible interactive analysis and
graphics of microbiome census data. Plos One. 2013;8: e61217.
doi:10.1371/journal.pone.0061217

51. Bisanz JE. qiime2R: Importing QIIME2 artifacts and associated data into R sessions. 2018.
Available: https://github.com/jbisanz/{qiime2R}

52. Team RC. R: A Language and Environment for Statistical Computing. Vienna, Austria;
2023. Available: https://www.R-project.org/

53. Jin Y, Qian H. V.PhyloMaker2: An updated and enlarged R package that can generate
very large phylogenies for vascular plants. Plant Diversity. 2022;44: 335–339.
doi:10.1016/j.pld.2022.05.005

54. Jin Y, Qian H. U.PhyloMaker: An R package that can generate large phylogenetic trees for
plants and animals. Plant Diversity. 2023;45: 347–352. doi:10.1016/j.pld.2022.12.007

55. Paradis E, Schliep K. ape 5.0: an environment for modern phylogenetics and evolutionary
analyses in R. Schwartz R, editor. Bioinformatics. 2019;35: 526–528.
doi:10.1093/bioinformatics/bty633

56. Shetty S, Person P, Person P. biomeUtils: Utilities for microbiome analytics. 2024.

57. Deagle BE, Thomas AC, Shaffer AK, Trites AW, Jarman SN. Quantifying sequence
proportions in a DNA-based diet study using Ion Torrent amplicon sequencing: which
counts count? Mol Ecol Resour. 2013;13: 620–633. doi:10.1111/1755-0998.12103

58. Clare EL, Chain FJJ, Littlefair JE, Cristescu ME. The effects of parameter choice on
defining molecular operational taxonomic units and resulting ecological analyses of
metabarcoding data. Genome. 2016;59: 981–990. doi:10.1139/gen-2015-0184

59. Hambäck PA, Weingartner E, Dalén L, Wirta H, Roslin T. Spatial subsidies in spider diets

vary with shoreline structure: Complementary evidence from molecular diet analysis and
stable isotopes. Ecology and Evolution. 2016;6: 8431–8439. doi:10.1002/ece3.2536

60.　Oksanen J, Simpson GL, Blanchet FG, Kindt R, Legendre P, Minchin PR, et al. vegan:
Community Ecology Package. 2024. Available: https://vegandevs.github.io/vegan/

61.　Yu G, Smith DK, Zhu H, Guan Y, Lam TT. Ggtree: An R Package for Visualization and
Annotation of Phylogenetic Trees With Their Covariates and Other Associated Data.
Methods in Ecology and Evolution. 2016;8: 28–36. doi:10.1111/2041-210x.12628

62.　Wickham H, Averick M, Bryan J, Chang W, McGowan L, François R, et al. Welcome to the
Tidyverse. JOSS. 2019;4: 1686. doi:10.21105/joss.01686

63.　Slowikowski K. ggrepel: Automatically Position Non-Overlapping Text Labels with
"ggplot2." 2024. Available: https://{CRAN}.R-project.org/package=ggrepel

64.　Jaffé K, Villegas G. On the communication systems of the fungus-growing ant
trachymyrmex urichi. Insectes sociaux. 1985;32: 257–274. doi:10.1007/BF02224915

65.　Kaspari M. Removal of seeds from Neotropical frugivore droppings. Oecologia. 1993;95:
81–88. doi:10.1007/BF00649510

66.　Leal LR, Oliveira PS. Interactions between Fungus-Growing Ants (*Attini*), Fruits and Seeds
in Cerrado Vegetation in Southeast Brazil'. 1998 pp. 170–178.

67.　Pizo MA, Oliveira PS. The Use of Fruits and Seeds by Ants in the Atlantic Forest of
Southeast Brazil [1]. Biotropica. 2000;32: 851–861. doi:10.1111/j.1744-7429.2000.tb00623.x

68.　Moreira AA, Bueno FC, Oliveira CG, Diniz EA, Garcez DL, Campos MCG, et al. Foraging
activity of *Trachymyrmex fuscus* (*hymenoptera*: *formicidae*) in a degraded area.
Sociobiology. 2007;50: 409–417.

69.　Kulhavy D, Smith L, Ross W. Impact of the Texas Leaf-Cutting Ant (*Atta texana* (Buckley))
(Order *Hymenoptera*, Family *Formicidae*) on a Forested Landscape. 2001.

70.　Hibert F, Taberlet P, Chave J, Scotti-Saintagne C, Sabatier D, Richard-Hansen C.
Unveiling the diet of elusive rainforest herbivores in next generation sequencing era? The
tapir as a case study. PLoS ONE. 2013;8: e60799. doi:10.1371/journal.pone.0060799

71.　Clare EL. Molecular detection of trophic interactions: emerging trends, distinct advantages,
significant considerations and conservation applications. Evol Appl. 2014;7: 1144–1157.
doi:10.1111/eva.12225

72.　Furlan EM, Davis J, Duncan RP. Identifying error and accurately interpreting environmental
DNA metabarcoding results: A case study to detect vertebrates at arid zone waterholes.
Molecular Ecology Resources. 2020;20: 1259–1276. doi:10.1111/1755-0998.13170

73.　Rytkönen S, Vesterinen EJ, Westerduin C, Leviäkangas T, Vatka E, Mutanen M, et al.
From feces to data: A metabarcoding method for analyzing consumed and available prey
in a bird-insect food web. Ecol Evol. 2019;9: 631–639. doi:10.1002/ece3.4787

74. Bohmann K, Elbrecht V, Carøe C, Bista I, Leese F, Bunce M, et al. Strategies for sample
labelling and library preparation in DNA metabarcoding studies. Molecular ecology
resources. 2022;22: 1231-1246. doi:10.1111/1755-0998.13512
703