Social task and aggressiveness shape immunity gene expression levels in the head of highly eusocial bees

Leonardo Campana[1] (ORCID 0000-0002-6049-3420), Carlos Antônio Mendes Cardoso Júnior[2] (ORCID 0000-0002-5039-8358), Anete Pedro Lourenço[3](ORCID 0000-0001-6929-7277), Klaus Hartfelder[1,2,*](ORCID 0000-0001-7981-8427)

[1]Departamento de Genética, Faculdade de Medicina de Ribeirão Preto, Universidade de São Paulo, Ribeirão Preto, SP, Brazil.

[2]Departamento de Biologia Celular, Molecular e Bioagentes Patogênicos, Faculdade de Medicina de Ribeirão Preto, Universidade de São Paulo, Ribeirão Preto, SP, Brazil.

[3]Departamento de Ciências Biológicas, Faculdade de Ciências Biológicas e da Saúde, Universidade Federal dos Vales do Jequitinhonha e Mucuri, Diamantina, MG, Brazil.

*Corresponding author.

Klaus Hartfelder, Departamento de Biologia Celular, Molecular e Bioagentes Patogênicos, Faculdade de Medicina de Ribeirão Preto, Universidade de São Paulo, Avenida Bandeirantes 3900, 14049-900 Ribeirão Preto, SP, Brazil

Email: klaus@fmrp.usp.br

Running title: Aggressiveness and immunity

Word count text (incl. spaces): 8973

Number of Figures: 7

Supplementary material: 3 figures, 6 tables

BULLETED HIGHLIGHTS

- The expression of immune system genes increases with social task/age in the abdomen and head of highly eusocial bees.

- The expression of Toll and Imd pathway genes is upregulated in the heads of the more aggressive stingless bee species.

- In the more docile stingless bee species, immune gene expression is higher in the abdomen than in the head.

- Our findings suggest a possible involvement of immunity pathway signalling in the regulation of key social behaviours.

ABSTRACT

Living in crowded perennial nests, social insects encounter high risks of disease outbreaks

and predation on colony resources. The innate immune system plays an important barrier

against microbial pathogens, but its association with group defensive behaviours remains

unexplored. Here we investigated the expression of innate immune system genes in the

workers of three species of Brazilian stingless bees (*Scaptotrigona postica*,

Frieseomelitta varia, and *Melipona quadrifasciata*), which differ in worker reproductive

traits and colony defence strategies. First, we standardized a behavioural arena test to

assess the species-specific individual aggressiveness level of the workers. Then, we

quantified the transcript levels of four core genes of the Toll and Imd pathways in the

heads and abdomens for the three main life cycle stages of the workers (nurse, guard and

forager). For the heads we found a strong association between immunity gene expression

and aggressiveness, with an upregulation in the heads of guards and foragers in

comparison to their abdomen tissues of the aggressive species, *S. postica*. while the less

aggressive species, especially *M. quadrifasciata* workers, did not show such strong up-

regulation. Our study establishes a potential molecular link between immunity pathways

and complex social behaviours in the Brazilian stingless bees, which evolutionarily lost

the sting apparatus for the defence of their colonies.

Keywords: Stingless bee; colony defence; innate immunity; life history; *Apis mellifera,*

social behaviour.

BULLETED HIGHLIGHTS

• The expression of immune system genes increases with social task/age in the

abdomen and head of highly eusocial bees.

• The expression of Toll pathway genes is highly upregulated in the heads of the

more aggressive stingless bee species.

• In the more docile stingless bee species, immune gene expression is higher in the

abdomen than in the head.

• Our findings suggest a possible involvement of immunity pathway signalling in

the regulation of key social behaviours.

ABBREVIATIONS

AMP – Antimicrobial peptide

GLM – General Linear Model

GLMM – General Linear Mixed Model

NF-kB - Nuclear Factor-kappa B

RT-qPCR - Quantitative reverse transcription polymerase chain reaction

TLRs – Toll-like receptors

Vg - Vitellogenin

INTRODUCTION

Social bees, including the honeybee and the stingless bees, are territorial animals that build perennial nests where they store food resources (honey and pollen) and rear and protect the brood and the highly reproductive queen. Such social life presents two types of risks. On the one hand, the nutritious resources in the colony can attract a variety of predators and nest parasites, and on the other, the large number of individuals in these nests represents an elevated risk for the spreading of diseases (Abbot, 2022). Yet strikingly, the analysis of the honeybee genome (The Honeybee Genome Sequencing Consortium, 2006) indicated a reduction in the number of genes comprising the bees' immune response repertoire compared to other insects, such as *Drosophila melanogaster* and *Anopheles gambiae*. While this loss can be compensated by social behaviours (social immunity) that provide barriers to the spreading of diseases (Cremer et al., 2018), a reduction in the repertoire of immune system genes, however, was also observed in bee species with a solitary lifestyle (Barribeau et al., 2015), apparently representing an ancestral trait in the bees, and thus unrelated to the evolution of advanced sociality in this clade.

For the defence of their nests, social bees adopt different collective behaviours. In the honeybee, *Apis mellifera*, guard workers use the sting and also biting as the primary strategies against intruders (Abbot, 2022; Nouvian et al., 2016). In contrast, having evolutionarily lost their sting apparatus, the stingless bees (Meliponini) exhibit diverse defence strategies. Some launch mass attacks, biting invaders, some apply sticky resins on them, while others have nest entrances that are difficult to access (for reviews see Grüter, 2020; Roubik, 2023; Shanahan and Spivak, 2021). Also, guards can respond to invasion threats by attacking aggressively, releasing alarm pheromones, by forming a

corridor of hovering guards, or by simply blocking a narrow nest entrance (Kärcher and

Ratnieks, 2009).

Molecular signatures of the defensive (aggressive) behaviours against threats have

been mainly studied in the Western honeybee, *A. mellifera*, and a strong correlation was

found between the colony level of aggression and the frequency of certain alleles in the

population (Avalos et al., 2020). At the individual level, workers were found to exhibit

distinct gene expression profiles in their brains, differentiating more aggressive

individuals from more docile ones (Rittschof, 2017). Such individual tendencies in

aggressiveness are, however, masked by the prevailing behavioural disposition of the

colony at any given moment, indicating that the colony genetic composition may override

the individual workers' genetic background in the decision to place or not an attack

(Avalos et al., 2020). At the cellular level, a recent transcriptomics study (Traniello et al.,

2023) revealed genomic subnetworks associated with colony aggression across various

cell types and across brain regions involved in olfaction, vision, and multimodal sensory

integration.

Previous studies have also demonstrated a correlation between aggressiveness and

immune system activity. For instance, the highly aggressive Africanized hybrids of *A.*

mellifera present strong traits of resistance against ectoparasites compared to more docile

A. mellifera subspecies (Camazine, 1986). Also, relatively docile, healthy workers, as

well as workers raised in less aggressive colonies, presented a gene transcriptional profile

in the brain and in the abdominal fat body that was very similar to that of individuals

suffering from acute pathogen infection (Rittschof et al., 2019), indicating that low

aggression resembles a sickness-like behavioural state. Furthermore, transcriptional

regulatory network analyses demonstrated that the neuro-genomic behavioural state of

aggressive honeybee workers is characterized by high expression levels of the

transcription factor Dorsal (Chandrasekaran et al., 2011).

Dorsal, a Nuclear Factor-kappa B (NF-kB) protein in insects, is a transcription

factor that plays a key role in the Toll signalling pathway of the innate immune response

of insects (Hoffmann, 1995), similar to NF-kB in the Toll-like receptors (TLR) signalling

pathway in vertebrates (Imler and Hoffmann, 2001). Unlike vertebrates however,

invertebrates rely solely on innate immunity as their defence against pathogens. This

system consists of humoral and cellular responses mediated by haemocytes and

antimicrobial peptides (AMPs) (Danihlík et al., 2015; Janeway and Medzhitov, 2002;

Lourenço et al., 2018; Strand, 2008). In this context, the Toll signalling pathway plays a

vital role in the immune response of insects. When activated in response to recognition

proteins that detect specific components of the cell wall of fungi and Gram-positive

bacteria, the Toll pathway triggers the translocation of the transcription factor Dorsal to

the nucleus, leading to the transcription of AMP genes (Evans et al., 2006; Kingsolver et

al., 2013; Zhang et al., 2021).

Similarly, the Imd signalling pathway is triggered by the recognition of

peptidoglycans from Gram-negative bacteria, resulting in the activation of the NF-kB

transcription factor Relish and, consequently, the expression of specific AMPs (Zhai et

al., 2018). In bees, the Toll and Imd pathways are known to regulate the expression of

AMPs, such as Abaecin and Defensin-1 (Lourenço et al., 2013, 2018). These pathways

are, however, not restricted to immune functions. In fact, as indicated by the primary

name 'Dorsal', this gene was first identified in *Drosophila* embryonic mutant screens as

a key transcription factor in the determination of the dorsal-ventral embryonic axis

(Lynch and Roth, 2011).

Both, innate immunity and aggressive behaviours are aspects related to somatic

maintenance, and both are energetically costly and, thus, likely of relevance to trade-offs

in life histories (Flatt et al., 2013; Stearns, 1992). In general, reproductive activity and

immune responses frequently show a negative correlation in insects, a pattern observed

in both solitary species from orders such as Orthoptera, Diptera, Odonata, Hemiptera, and

Coleoptera, as well as in some social insects, such as certain ant species (Schwenke et al.,

2016). But how does this correlation look like in social insects, especially in the highly

eusocial ones, where colony (somatic) maintenance and reproductive activity are

dissociated into two morphologically distinct castes, the workers and the queen?

Studies on social insect immunity, colony defence strategy, reproductive activity,

and their associated gene networks have been mainly conducted using *A. mellifera*

workers as a model. However, the honeybees (Apini) represent a single genus (*Apis*)

consisting of less than a dozen species, and thus are a relatively small and rather

homogeneous group in the phylogeny of the bees (Apidae). In contrast, the stingless bees

(Meliponini) comprise over 500 recognized species distributed among 58 genera (Grüter,

2020; Melo, 2021). With a pantropical distribution, they exhibit a high level of biological

diversity in terms of nesting sites, reproductive strategies, and nest defence. Stingless

bees, like honeybees, are threatened by anthropogenic impacts, such as crop monoculture

and pesticide use. However, unlike honeybees, they thrive in greenhouse environments,

where they are increasingly recognized as efficient pollinators in high-value agriculture.

(Grüter, 2020).

Taking advantage of the stingless bees' biological variability, we set out in this

study to investigate the relationship between aggressiveness (defence behaviour) and the

immune system profile of the workers of three stingless bee species with distinct life

history configurations (Fig. 1). The workers of *Scaptotrigona postica* are very aggressive

in their nest defence and are able to lay eggs in the presence of the queen (Nogueira-Neto,

1997). The workers of *Melipona quadrifasciata* are also able to lay eggs in the presence

of the queen, but they are notably docile (Grüter, 2020; Nogueira-Neto, 1997), and the

workers of *Frieseomelitta varia,* while moderately defensive, are completely sterile

(Boleli et al., 1999; Nogueira-Neto, 1997). Hence, we considered that this contrast in

character combinations would allow us to address the question as to how life histories

and their respective trade-offs (Stearns, 1988), specifically those between the workers'

aggressiveness in the nest defence and their reproductive potential became configured

with respect to a costly system, such as the innate immune response. Specifically, while

on the one hand the investment in aggressiveness is likely very costly to the individual

worker, but benefits the colony fitness, worker reproduction on the other hand creates a

fitness conflict, primarily with the queen, but in the honey bee, also with the other workers

(Ratnieks et al., 2006).

Hence, the underlying hypotheses that we proposed to address were that: (a) even

in the absence of a specific immune challenge, the expression levels of immunity-related

genes increase with the age of the bees, reflecting their progression in social behaviours

(Wilson-Rich et al., 2008), and (b) the expression levels reflect the level of aggressiveness

and the reproductive biology of the workers. To do so, we adapted an arena assay that

allowed us to quantitatively assess the species-specific levels of aggressiveness for

individual workers, and we evaluated the expression levels of a set of four key genes of

the Toll and Imd pathways in the heads and abdomens for three important behavioural

stages in the workers' life cycle. Finally, we compared the results obtained for the three

stingless bees with those of a corresponding analysis that we performed on honeybee

workers.

MATERIAL AND METHODS

The arena behavioural assay: A quantitative tool for assessing the species-specific aggressiveness level of individual workers

To obtain information about the aggressiveness at the individual level for the workers of the three focal stingless bee species, we performed an arena assay based on the protocol of Dierick (2007) for *Drosophila melanogaster*, which has subsequently been applied to honeybees by Shpigler et al. (2017). This assay eliminates variables associated with the social colony context, including the presence of alarm pheromones elicited by nestmates, as well as environmental factors, such as ambient temperature, group size, diet, time, and territorial size in the bees' reaction against a perceived threat.

Workers of *Scaptotrigona postica* (Latreille, 1807), *Frieseomelitta varia* (Lepeletier, 1836), and *Melipona quadrifasciata* (Lepeletier, 1836) were collected from one single colony of each species, all maintained in the apiary at the Department of Genetics, Ribeirão Preto Campus, University of São Paulo, Brazil. From each colony of *S. postica* and *F. varia* (Lepeletier, 1836), 100 newly-emerged workers and 10 nurse-stage workers were collected and kept together in a single cage (8 x 11 x 13 cm) in an incubator at 34 °C for 14 days. For *M. quadrifasciata*, due to their larger body size, the number of newly emerged workers (N = 40) and nurse-stage workers (N = 4) in the cage was reduced. The cages were provided daily with pollen collected from the colonies of origin and with commercial *A. mellifera* honey diluted in water at 50%. The presence of nurse bees in the cages is important, as they feed the newly emerged workers during their first days of adult life.

After 14 days in the cages, the workers were submitted to a 15-hour food restriction period prior to the arena assays. This food restriction period proved to be essential, as, during a pilot experiment, we had noted, in the case of all three species, that

workers that were fed until the day of the assay displayed a disregard for the presence of

an intruder in the assay arena (Fig. S1). The 15-days-old workers were then individually

placed into 9 cm diameter Petri dishes lined with filter paper. After one minute of

acclimatization, an intruder, a newly emerged *Tetragonisca angustula* worker, was

introduced into each dish. The workers of the stingless bee species *T. angustula* are

smaller than those of our focal species, and not aggressive when they are still young. Each

Petri dish was filmed for 5 minutes, and the recordings were then analysed using Boris v.

8.11.3 software (Friard and Gamba, 2016). In total, we performed arena assays for 42 *S.*

postica workers, for 42 *F. varia* workers, and 19 *M. quadrifasciata* workers.

The following aggression parameters were considered: (i) Antennation (events of

touching the intruder with the antennae), (ii) frequency of attacking (percentage of tested

individuals that show any attack behaviour against the intruder during the observation

period), (iii) latency to attack (time until the first attacking event occurred), (iv) the

attacking index (total amount of time spent attacking the intruder during the five-minute

period), and (v) the pursuit index (total amount of time spent pursuing the intruder during

the five minutes).

Sampling of the worker bee life cycle stages for gene expression analysis

Workers of *S. postica*, *F. varia*, and *M. quadrifasciata* were obtained from the same

colonies as those used for the arena aggression assays. For each species, sampling was

done from three colonies. From each colony, seven workers were retrieved for each of

the three life cycle stages: nurse, guard, and forager. Sampling was done during the

Southern summer season, between November 2021 and March 2022, always between

9:00 and 10:00 am. Workers of the corresponding life cycle stages were also sampled

from a honeybee colony kept in the same apiary (*Apis mellifera*, Africanized hybrids) in

August 2022.

The sampling procedure was as follows: First, guards were collected, considering

that these were the bees that left the nest and attacked the collector's entomological sweep

net after three knocks with a hive chisel on the front board of the hive. Guards of *S. postica*

and *M. quadrifasciata* generally had an empty pollen basket (corbicula). In contrast, the

workers of *F. varia* that attacked the net all had their corbiculae (pollen baskets on the

hind legs) full of resin, as part of their colony defence strategy (Roubik, 2023). Next,

foragers were collected in front of the hives, when returning from the field with pollen-

filled corbiculae. Finally, the colonies were opened, and nurse-stage bees were collected

from the brood area. In all three species, nurse-stage bees can be identified by their

enlarged abdomen, due to their larval food-filled crop, and, their lighter body colour, as

they are still younger than the guards or foragers, and not yet fully pigmented (Nogueira-

Neto, 1997). Nurse bees are primarily engaged in the provisioning of newly constructed

brood cells with larval food. Fig. S2 shows the phenotypic characteristics of these life

cycle stages for each of the three species.

After sampling, the bees were briefly put on ice to diminish their motility and then

snap frozen in liquid nitrogen. The heads and abdomens were separately transferred into

Eppendorf vials containing 300 µL of TRIzol reagent (Invitrogen, Waltham, MA, USA)

and stored at -80 °C until RNA extraction. For the two species of smaller body size, *S.*

postica and *F. varia*, each sample was composed of a pool of heads or abdomens from

three individuals, while for the larger species *M. quadrifasciata* and *A. mellifera*, each

sample consisted of a single individual only. Pooling was necessary to obtain the RNA

yield required for gene expression analysis.

In this respect, we recognize as a limit of our study that we not separate between

age and social function. Separating these would require the setup of single-cohort

colonies, but such an experimental design is currently only available for honey bees

(Marco Antonio et al., 2008; Sullivan et al., 2000) and not for stingless bees. While in the

current study our emphasis was primarily on social task and not on age, the compiled

bionomics data available for stingless bees (Grüter, 2020) show that there is a clear age

sequence in social task performance.

RNA extraction and gene expression analysis

The head and abdominal tissues were macerated and homogenized with a pestle while

still in the TRIzol reagent before proceeding with the RNA extraction according to the

manufacturer's protocol. RNA extracts were treated with RNase-free DNase (Invitrogen)

to remove residual DNA. The quantity and quality of the RNA was checked

spectrophotometrically (NanoVue, GE Healthcare, Chicago, IL, USA). Reverse

transcription was performed using 1 μg of total RNA for cDNA synthesis, using the

Ultrascript 2.0 cDNA synthesis kit (PCRBiosystems, London, UK), following the

manufacturer's instructions.

The transcript levels of *Toll*, *Dorsal*, *Myd88* (Toll signalling pathway), and *Imd*

(Imd signalling pathway) were assessed through quantitative real-time PCR (RT-qPCR)

assays. Primer sequences for these genes were based on the respective genomic sequences

of *M. quadrifasciata* and *F. varia* (Table S1). The same primers were also tested and

successfully used to amplify the corresponding transcripts of *S. postica,* for which a

sequenced genome is not yet available.

Primer efficiencies were calculated as $E = 10^{[-1/\text{slope}]}$, using a 1:10 dilution series

of a cDNA pool containing equal amounts of RNA from all experimental groups and

species (Table S1). This pooling approach is a widely accepted method to estimate general primer performance across conditions. The primers for the endogenous control genes *Rpl32* and *Rps18* had previously been designed and validated for *F. varia and M. quadrifasciata* (Freitas et al. (2019). Additionally, as this was the first time these reference genes were used here also for *S. postica*, we confirmed their suitability by a series of additional efficiency tests using individual cDNA pools from each experimental group (head and abdomen tissues from both guards and foragers). The results show their high and consistent efficiencies (Table S1). For the normalization process, the Ct results of the two reference genes were combined by calculating their mean, and this average Ct value was used. The primer sequences for the corresponding *A. mellifera* immune system genes are based on the work by Evans et al. (2006), and those for the honeybee endogenous control gene *Rpl32* (*Rp49*) on Lourenço et al. (2008).

Amplification mixtures were set up in a total volume of 10 μL, containing 5 μl of the 2×qPCR SyGreen Mix kit (PCR Biosystems, London, UK), 1 μL of 10 × diluted cDNA, primers (forward and reverse) at a final concentration of 0.125 pmol, and ultrapure water completing the final volume. The assays were performed in a StepOnePlus real-time PCR system (Applied Biosystems, Waltham, MA, USA) with the following amplification protocol: 95 °C for 10 min, followed by 40 cycles at 95 °C for 15 s and 60 °C for 30 s. After these cycles, a melting curve analysis was performed to confirm the specificity of the amplification products. Assays were done in biological septuplicates (n = 7 per colony), and each sample was analysed in technical triplicates. The relative expression of the genes of interest was calculated by the $2^{-\Delta\Delta Ct}$ formula proposed by Livak and Schmittgen (2001).

To ensure the robustness of our analyses, we decided to remove the nurse bee samples of *S. postica* from the analyses, as for this group the reference genes exhibited

considerable variation compared to the guards and foragers (Fig. S3). Despite this

adjustment, it is still possible to compare gene expression profiles between species with

reproductive (*M. quadrifasciata*) and non-reproductive (*F. varia*) workers, while the *S.*

postica data were used to specifically discuss the relationship between the life history

traits aggression and immunity.

Statistical analysis

Statistical analyses were performed in the RStudio platform (R version 4.3.0). Data

obtained from the arena assays were submitted to Shapiro-Wilk and Kolmogorov-

Smirnov tests to check the adequacy for parametric testing. Levene's test was used to

verify homoscedasticity. As these conditions were not met, we used the Kruskal-Wallis

test (dplyr package) with Dunn's post-hoc test for multiple comparisons of three or more

groups. For the analysis of the gene expression data, we used a Generalized Linear Mixed

Model (GLMM) (lme4 package), considering "body compartment" and "life cycle stage"

as fixed factors and "colony" as random. For *Apis mellifera*, in the absence of colony

replicates, we utilized the GLM test (lme4 package) to identify differences between

means. The log link function was used to approximate the gene expression data to a

Gaussian distribution, which was checked by inspecting the residuals (DHARMa

package). Graphical representations of the data were generated with GraphPad Prism v.

7 (GraphPad Softwar, Boston, MA). An adjusted p-value < 0.05 was considered

significant in all analyses.

Ethics statement

For the behavioural assay, the bees were kept in social groups in cages of suitable size

and with *ad libitum* access to food. The 15-hours food restriction period had no visible

negative effect on the well-being of the bees. After the arena assay, the bees were

sacrificed by instant freezing in liquid nitrogen, as they could not be returned to their

colonies without being attacked there as intruders. For the gene expression analyses the

bees were briefly cooled on ice to diminish their motility before instantaneous freezing in

liquid nitrogen. This is a standard procedure to calm down before introducing them into

any experimental procedure.

The maintenance of the native stingless bee colonies in the apiary of the Ribeirão

Preto Campus of the University of São Paulo is authorized by the *Instituto Chico Mendes*

de Conservação de Biodiversidade – Ministério de Meio Ambiente, Brazil, under the

licence 81843-1 to the senior author.

RESULTS

Species-specific individual aggressiveness of stingless bee workers

The arena assays demonstrated that under standardised conditions (number of bees of

similar age, temperature, and diet) *S. postica* showed the highest scores in the aggression

indices, while *M. quadrifasciata* exhibited the lowest scores (Fig. 2, Table S2). The scores

for *F. varia* indicated intermediate levels of aggressiveness. *S. postica* was the species

with the highest frequency of individuals that attacked the intruder in the arena. This

species also was the one with the highest frequency of fatal attacks (Fig. 2A). In

comparison, the workers of *F. varia* and *M. quadrifasciata* had much lower attacking

frequencies. The behavioural profiles of the individual bees inside the arena were also

different for the three species. Already within the first minute after introducing the

intruder specimen, the workers of *S. postica* behaved agitated, walking around quickly,

while those of *F. varia* took short flights, during which they collided with the intruder or

the arena wall. *M. quadrifasciata* workers, in turn, appeared rather lethargic, with little

movement in the arena.

All workers of *S. postica* and *F. varia*, even the non-attacking individuals,

exhibited curiosity towards the intruder, with several events of observed antennation. In

contrast, the workers of *M. quadrifasciata* did not engage with the intruder, although they

were active and moving around the arena like the other species. (Fig. 2B). The only attack

event that we saw for an *M. quadrifasciata* worker was by an individual that behaved

very agitated already from the start of the observation period, heavily beating its wings

and vibrating its body.

The species with the shortest attack latency towards the intruder, that is, it attacked

the intruder faster than the other species, was *S. postica* (Fig. 2C: Kruskall-Wallis test: H

$= 21.5$, $P < 0.01$). *F. varia* and *M. quadrifasciata* showed longer attack latencies. The

attacking index, which is the time the individual remained attacking the intruder, was

higher for *S. postica* than for *F. varia* and *M. quadrifasciata* (Fig. 2D: Kruskal-Wallis

test: $H = 23.31$, $P < 0.001$), and among the three species, *S. postica* and *F. varia* workers

presented the highest number of pursuit events (Fig. 2E: Kruskal-Wallis test: $H = 6.094$,

$P < 0.5$). During these events, *F. varia* workers often pursued the intruder with opened

wings, a behavior not observed in *S. postica* workers. However, these short flights were

not counted as pursuit events; only pursuits that occurred while walking on the plate were

considered, to maintain the same standard for all three species. No pursuit event was

observed in the case of *M. quadrifasciata*. In these events, the *F. varia* workers used to

pursue the intruder with opened wings, a behaviour that was not seen in *S. postica*

workers. No pursuit event was observed in the case of *M. quadrifasciata*.

As the arena test permitted us to quantify the individual aggression levels of the

three species under a controlled experimental setup, our results provide evidence that *S.*

postica workers exhibit higher levels of aggression compared to the other species. *M.*

quadrifasciata workers are rather docile, and *F. varia* workers show an intermediate level

of aggressiveness. While this confirms the common impression one typically has when

working with colonies of these species, the arena test results can now provide quantitative

indices for the individuals' species-specific aggressiveness levels, independent of

eventually variable colony conditions, which cannot be easily standardized in the field.

Of note, we did not include *A. mellifera* workers in the arena test because their defensive

behaviour differs significantly from that of stingless bees. While stingless bees primarily

rely on biting as a means of defence, *A. mellifera* workers typically attempt to sting, rarely

engaging in biting. This fundamental difference in behaviour would require distinct

criteria for quantifying aggression, making direct comparisons between the species

problematic. Additionally, *A. mellifera* workers do not survive sufficiently long in

confinement due to their inability to defecate without flying, leading to physical

debilitation and mortality over time. In contrast, stingless bees can defecate within the

cage and even organize waste in specific areas, allowing for easier maintenance and

ensuring their survival in standardized experimental setups (See Discussion for details).

Body compartment and social task-specific levels of immunity gene expression in the

workers of stingless bees

The three stingless bee species presented distinct profiles of immunity gene expression

levels for the head and abdomen. For *S. postica*, the most aggressive of the three species,

we found that the workers showed elevated transcript levels of *Toll* (GLMM: $\chi^2 = 109.6$,

$P < 0.001$), *Myd88* (GLMM: $\chi^2 = 25.9$, $P < 0.001$), *Dorsal* (GLMM: $\chi^2 = 34.8$, $P < 0.001$),

and *Imd* (GLMM: $\chi^2 = 30.1$, $P < 0.001$) in the head when compared to the abdomen (Fig.

3, Table S3). For the *Myd88* gene, the guards' heads actually showed higher expression

when compared with the foragers' heads (GLMM: $t = -3.5$, $P = <0.001$). For *M.*

quadrifasciata, the most docile of the three species, the directionality was exactly the

opposite, with significantly higher levels for *Toll* (GLMM: $\chi^2 = 133.1$, $P < 0.001$), *Myd88*

(GLMM: $\chi^2 = 4.2$, $P = 0.03$), and *Dorsal* (GLMM: $\chi^2 = 11.5$, $P < 0.001$) in the abdomen

(Fig. 4, Table S4). *F. varia* was similar to *S. postica* regarding the expression of the genes

Myd88 (GLMM: $\chi^2 = 41.6$, $P < 0.001$) and *Imd* for guards and foragers (Fig. 5, Table S5:

GLMM: $\chi^2 = 19.7$, $P < 0.001$), but the difference between the two body compartments

was much less pronounced.

With respect to the life cycle stages, the guards and foragers of *F. varia* and *M.*

quadrifasciata in general showed a tendency towards higher levels of immunity gene

expression compared to the nurse-stage workers. The workers of *M. quadrifasciata*

showed differences in gene expression levels between the younger (nurse-stage) and the

older bees (foragers and guards) in both body compartments for the genes *Toll* (GLMM:

$\chi^2 = 133.1$, $P < 0.001$) and *Myd88* (Fig. 4: GLMM: $\chi^2 = 4.2$, $P < 0.05$). With respect to the

other two genes, an abdominal up-regulation of *Dorsal* (GLMM: $\chi^2 = 11.5$, $P < 0.001$)

was associated with the guard and forager stages, while *Imd* was higher expressed only

in the heads of guards compared to nurses (GLMM: $t = 3.2$; $P < 0.05$).

For *F. varia* workers, the heads showed significantly higher *Imd* expression in

both guards (GLMM: $t = 8.25$; $P < 0.01$) and foragers (GLMM: $t = 9.75$; $P < 0.01$)

compared to nurses. *Dorsal* and *Toll* are both higher expressed in foragers compared to

nurses (*Dorsal* GLMM: $t = 2.8$; $P < 0.05$; *Toll* GLMM: $t = 4.57$; $P < 0.01$) but not in

guards, while *Myd88* is higher expressed in guards compared to nurses (GLMM: $t = 3.12$;

$P < 0.01$), but not in foragers (Fig. 5; Table S5).

Notably, we found high levels of the immune gene transcript levels in the heads

of guards and foragers, especially in the more aggressive species, *S. postica*. Yet, given

that the abdominal fat body is the primary insect tissue for mounting an immunity

response (Li et al., 2019), we asked whether these high expression levels in the bees'

heads would in fact be related to an actual immune response. To address this, we assessed

the levels of *Abaecin* transcripts, as these should represent a direct readout for a canonical

immune response (Lourenço et al. 2018).

The *Abaecin* gene encodes a major antimicrobial peptide, and its expression is

driven by both the Toll and the Imd pathway in the honeybee (Lourenço et al., 2018). For

S. postica and *F. varia* we found no differences in the *Abaecin* expression levels between

head and abdomen (GLMM: $\chi^2 = 1.8$, $P = 0.17$; and $\chi^2 = 0.5$, $P=0.44$, respectively), while

M. quadrifasciata workers showed higher *Abaecin* expression in the abdomen than in the

head (Fig. 6: GLMM: $\chi^2 = 8.7$, $P < 0.003$). Furthermore, guards and foragers of both *S.*

postica and *M. quadrifasciata* had higher levels of *Abaecin* expression than nurses in both

body compartments, while for *F. varia* and *A. mellifera,* this was the case only for the

head.

The similarity in the expression levels for *Abaecin* with respect to social task

status across the four species indicates that, even in the absence of an immune challenge,

the immune system readiness level is higher in the older bees (guards and foragers),

possibly because they perform riskier tasks outside the colony in terms of predation threat

and pathogen exposure. Hence, while the expression patterns of the Toll and Imd pathway

genes showed pronounced differences for the head *vs.* abdominal tissues across the four

bee species, associated with their respective aggressiveness levels, the *Abaecin*

expression levels showed an age/task-related pattern that was common across the four

species. This discrepancy between the two signalling pathways and the actual immunity

readout (*Abaecin*) indicates a non-canonical function for the Toll and Imd pathways in

the heads of the more aggressive stingless bees.

Immunity gene expression in the workers of Apis mellifera

The association between aggressiveness and immune gene expression levels in the heads of the stingless bee workers that perform the higher risk tasks of guarding and foraging, thus raised the question as to whether this could be a conserved trait across the highly eusocial corbiculate bees (Apini and Meliponini). To address this, we assessed the transcript levels of the same innate immunity genes also in the honeybee, *A. mellifera* (Fig. 7, Table S6).

We found that only *MyD88* and *Imd* are higher expressed in the heads compared to the abdomen (GLM: $\chi^2 = 4.14$, $P = 0.04$; and $\chi^2 = 26.86$, $p < 0.001$, respectively). Also, for all tested genes, we observed an increase in gene expression in at least one tissue when comparing guards and foragers to nurse bees (Fig. 7; Table S6). Furthermore, like in *M. quadrifasciata*, the *Abaecin* expression was higher in the abdomen than in the head in all the life cycle stages considered in this study (GLM: $\chi^2 = 16.78$, $P < 0.001$). These findings for the notoriously aggressive Africanized honey bee provide additional support for a possibly non-canonical function of the Toll and Imd pathways, associated with the more aggressive Meliponini species tested in this work.

DISCUSSION

Based on the diversity in life history parameters seen in stingless bees, exemplified by the three focal species of this study, we intended to investigate the possible relationship between the bees' social tasks, their aggressiveness, and their immune system status against the background of the worker's reproductive biology (Fig. 1). The most striking finding was the upregulation in the expression of all four of the Toll and Imd pathway genes in the heads of the guards and foragers of *S. postica* in comparison to their abdominal tissues (Fig. 3), as this is the more aggressive among the three species that we

tested (Fig. 2). In the heads of *F. varia,* only two of these genes (*Myd88* and *Imd)* are

higher expressed in the heads of the workers (Fig. 5)*,* while in *M. quadrifasciata,* the

opposite is seen, with two genes (*Toll* and *Dorsa*l) actually higher expressed in the

abdomen (Fig. 4)*.* In *A. mellifera*, a possibly aggressiveness-related pattern was observed

for only two of the four genes (*Myd88* and *Imd*) in the heads of guards and foragers.

However, as age and task are often correlated in social insects, we acknowledge that these

differences may also reflect age-related variation rather than task-specific regulation.

Nonetheless, we interpret these findings as indicating a possibly non-canonical

(neo)functionalization of the immune system pathways in the context of colony defence

behaviours of highly eusocial bees.

Nest defence and aggression in stingless bees

The possible reasons for the evolutionary loss of the stinging apparatus in the Meliponini

are still an open question. Yet, the idea that the small size of the bees (miniaturization)

would reduce the efficiency of their sting is not a very convincing argument, considering

the continued presence of a sting in even smaller ants. Similarly problematic is the

concomitant reduction of their venom gland, in view of the fact that genes encoding

venom proteins were found conserved in stingless bee genomes (Koludarov et al., 2023).

Yet, what is well known among beekeepers, is that stingless bee species greatly vary in

their aggressiveness as part of their strategies of nest defence.

The results obtained here via the arena assay confirmed that *S. postica* workers*,*

at the individual level, exhibit a strong aggression response towards an intruder. This is

consistent with the fact that, among the three species that we analysed, *S. postica* also has

the largest nest entrance tunnel and the largest number of guards controlling the intense

flight traffic (Couvillon et al., 2008). Furthermore, when they attack people with bites,

the sensation is quite painful compared to bites from other stingless bees (Shackleton et al., 2015). Hence, bees of the genus *Scaptotrigona* have an efficient, aggression-based colony defence strategy. In fact, beekeepers often emphasize the precautions that are needed when working with colonies of this genus (Blochtein, 2008).

F. varia, the species that showed the intermediate level of aggressiveness in the arena test (Fig. 2), also showed attention towards the intruder. However, they did not attack with bites in most of the assay rounds. They have smaller mandibles than *S. postica*, and their main defence strategy is to apply sticky resin onto intruders (Shackleton et al., 2015). This strategy may not have much of an effect against vertebrate enemies, but it is very efficient against other arthropods (Couvillon et al., 2008; Grüter, 2020). In line with that, during the arena assays the individuals were seen to perform short flights, bouncing against the intruder inside the Petri dish. In the field, such collision flights would allow the *F. varia* workers to stick the resin from their corbiculae onto the potential intruder, causing immobilization of the latter (Roubik, 2023).

In contrast to the other two species, *M. quadrifasciata* workers were completely unaggressive in the arena assay. In fact, they were even less aggressive individually than what is observed when they are confronted with a potential intruder at the entrance of a colony in the field (Breed and Page, 1991; Nunes et al., 2014).

Pros and cons of the arena assay

The arena assay allowed us to quantitatively assess the aggressiveness level of individual stingless bee workers of three focal species, independent of variation in colony contexts. Yet, like in other behavioural tests on individual bees, certain adjustments were necessary. This included the necessity of introducing a 15-hour food restriction period prior to testing a bee's aggressiveness towards an intruder in the arena. Like in the proboscis extension

reflex (PER) assay (Bittermann et al., 1983; Smith and Burden, 2014) a food restriction

turned out to be necessary for the appropriate motivation of the bees to respond to the

stimulus. As previously shown in a different behavioural context, food restriction affects

the metabolism of *A. mellifera* workers, including their brain metabolism (Rittschof et al.,

2018). Furthermore, there is strong evidence that aggressive behaviours are associated

with significant metabolic changes in the brains of honeybees (Chandrasekaran et al.,

2015; Li-Byarlay et al., 2014), underlining the importance of taking into account the bees'

nutritional status when designing and standardising aggression tests.

With this in mind, there is the possibility that bees of larger size, like *M.*

quadrifasciata, may have larger energy reserves than the smaller *S. postica* and *F. varia*,

and potentially, this could make them less responsive in the arena assay. However, when

we extended the food restriction to 24 hours, this led to increased mortality. Hence, the

results of the arena assay performed under these standardised conditions confirmed the

field observations regarding the differences in the species' aggressiveness also at the level

of the individual workers.

Notably, we did not perform the arena test for the testing of honeybee workers,

because the protocol conditions that we used for the stingless bees were found not suitable

for the former. The main reason was that for the confinement in the cages, the stingless

bee workers received a diet that contained pollen. Yet, *A. mellifera* workers consuming

pollen hardly survive for more than 14 days in the cages because, different from the

stingless bees, they cannot defecate in confinement, without flying (Cole et al., 2021).

Furthermore, an arena aggression assay for *A. mellifera* workers was formerly designed

and performed with small groups of bees and not with individual bees (Nouvian et al.,

2016).

Elevated gene expression levels in the head of an aggressive stingless bee species indicate

a non-canonical function for innate immunity signalling pathways

To address the question as to whether and how the species-specific levels of

aggressiveness and the innate immune system of the stingless bee workers may be

intertwined in terms of life history strategies and trade-offs (i.e., reproduction vs.

immunity/somatic maintenance), we sampled workers according to their social task in the

colony. What we found is that in the more aggressive species, *S. postica*, the levels of

immune gene expression were upregulated in the head of the guards and foragers (Fig.

3). Since the abdominal fat body, but not the head, is the primary organ responsible for

the production of antimicrobial peptides (Li et al., 2019), this strong overexpression of

the Toll and Imd pathway genes in the bees' heads is clearly remarkable.

Toll and *Dorsal* genes (*NF-kB* in vertebrates) are present in the genomes of most

organisms, and the Toll/Toll-like receptors (TLR) signalling pathway is a very ancient

component in metazoan history (Leulier and Lemaitre, 2008). Actually, while it has been

hypothesized that the TLR immune functions involving Myd88 and NF-kB arose in the

Bilateria (Kim and Ausubel, 2005), all the core elements of Toll signalling are already

present in Cnidaria (Chapman et al., 2010). And even though the primary function of Toll

signalling is the recognition of pathogenic microorganisms, it has also acquired an

embryonic function in insects, determining the dorsoventral axis (Anderson et al., 1985).

Furthermore, in addition to its canonical function as a transmembrane receptor for

immune signalling, Toll may serve additional roles in the nervous system in both

vertebrates and invertebrates (Dresselhaus and Meffert, 2019).

Myd88, an adapter protein downstream to the transmembrane protein Toll (Horng

and Medzhitov, 2001), exhibited higher expression levels in the heads of *S. postica* guards

compared to foragers (Fig. 3). Interestingly, Myd88 is already known to play an important

regulatory role in mammalian neural and behavioural functions, including the regulation

of chronic stress responses (Hosoi et al., 2021). Furthermore, inhibition of this protein

promotes depression-like behaviour (Yao et al., 2023) and depression-like sleep disorders

(Choudhury et al., 2022). Since *Myd88* was found higher expressed in the aggressive

guard bees of *S. postica,* this molecular signature strengthens the hypotheses that ancestral

immunity proteins were co-opted to shape complex collective colony defence behaviours

in the highly eusocial stingless bees.

Dorsal (NF-kB), the main transcription factor of the Toll pathway, was also found

overexpressed in the heads of *S. postica* workers compared to their abdomens, whereas

in the docile *M. quadrifasciata* workers, this pattern is inverted. In mammals, NF-kB is

known to be involved in cognitive processes (Dresselhaus and Meffert, 2019). It is

constitutively activated in neurons from various regions of the central nervous system,

mainly in glutamatergic neurons (Bhakar et al., 2002; Kaltschmidt et al., 1994), and it

notably accumulates in the terminal regions of these neurons. Compelling evidence

indicates that the activation of NF-kB is induced by the neurotransmitter glutamate and

the influx of calcium ions (Ca^{2+}) (Lilienbaum and Israël, 2003; Meffert et al., 2003), and

upon activation, NF-kB undergoes a retrograde transport and becomes translocated to the

nucleus of the respective neurons (Meffert et al., 2003; Meffert and Baltimore, 2005;

Wellmann et al., 2001). In this way, it functions as a transducer of synaptic information

to the nucleus, where it assumes its role as a transcription factor (Mémet, 2006). In

addition, NF-κB is associated with memory consolidation and learning in vertebrates and

invertebrates, such as crustaceans and insects (Freudenthal and Romano, 2000; Mémet,

2006; Merlo et al., 2005). This response apparently occurs in neurons, but not in glial

cells, where NF-κB is strictly associated with the control of mammalian immune

responses and metabolism (Dresselhaus and Meffert, 2019). Based on our results obtained

for stingless bees, it is thus, reasonable to assume that in these bees, the Toll signalling

pathway, including its key transcription factor Dorsal, were co-opted to perform a

behavioural function in addition to its canonical immune system role.

With respect to *Imd* expression, the pattern was similar to *Dorsal* in *S. postica* in

terms of body compartment and social task specificity. However, aside from its immune

system function, little is known about potentially alternative roles of Imd signalling in

insects. Nonetheless, there are reports on its involvement in neurodegeneration, cell

death, and regulation of the gut microbiota (Hanson and Lemaitre, 2020; Zhai et al.,

2018). In this respect, our results may be the first ones to identify a link between the Imd

pathway and behavioural traits also in an invertebrate.

As emphasized, the expression patterns of all four immune signalling pathway

genes quantified in this study indicate an increase associated with the age-related social

task performed by the workers of the stingless bees, in both the head and the abdomen.

But how does this actually relate to the bees' immune system status? In *D.* melanogaster,

healthy and free of infection individuals showed a chronic increase with age in the

immune system activity, leading to an increased expression of antimicrobial peptides

(AMPs) in the brain, followed by marked neurodegeneration (Stuart et al., 2022).

Furthermore, a study by Kounatidis et al. (2017) showed that suppression of the Imd/NF-

κB signalling pathway in glial cells resulted in an extension of the flies' lifespan. This is

in line with data for *A. mellifera* (Lourenço et al., 2019), where nurse bees have much

lower expression levels of immunity genes and an increased lifespan.

In adult insects, AMPs are the most prominent products generated by the

activation of the Imd and Toll signalling pathways (Bulet and Stocklin, 2005), and

activation of the immune system in the brain leads to high AMP expression. This can

have collateral cytotoxic effects related to neurodegeneration, and consequently

behavioural changes, including sleep and memory dysregulation (Barajas-Azpeleta et al., 2018; Stączek et al., 2023). The brain of *A. mellifera* was previously seen to exhibit AMP overexpression upon infection with Deformed Wing Virus, a common bee virus, demonstrating the ability of the brain to elicit a strong immune response (Pizzorno et al., 2021). Studies with *D. melanogaster* in turn demonstrated that the production of AMPs may also be regulated by pathways other than Toll and Imd (Stączek et al., 2023). For instance, the transcription factor FoxO, which is a downstream regulator of the insulin signalling pathway, can activate the expression of AMPs in flies undergoing starvation, even in the absence of an infection or injury event (Becker et al., 2010). Therefore, the elevated *Abaecin* transcript levels that we found in the head of the older stingless bees may not be exclusively driven by the immune system pathways Toll and Imd, but also by pathways associated with their age-specific diets and, consequently, metabolic regulation. Nonetheless, in our previous study, where we compared head and abdominal expression of candidate genes for social task performance in stingless bees (de Souza and Hartfelder, 2023) we did not see any expression difference in the abdominal *tor* gene expression for *F. varia* and *M. quadrifasciata* nurses and foragers. In their heads, however, we saw an inverse expression pattern between the two species, indicating that this major nutrient-sensing pathway may be linked to the head immune gene expression profiles.

Finally, in this respect as already previously stated, we recognize as a limit of our study that we cannot not separate between age and social function. These two factors are strongly intertwined, with young bees preferentially performing tasks within the nest, while older ones take on guarding and foraging task, and these tasks are strongly associated with immune system functionality (Amdam et al., 2005; Lourenço et al., 2019).

Immune system gene expression and the life histories of social bees

At this point it is important to state that the question that we asked differs from the one

addressed in most studies on immune system functions in social insects. Specifically, we

did not ask how does the bees' immune system respond to a pathogen challenge, but

rather, whether the aggressiveness level of the workers, as part of the colony defence

against intruders and predators, is reflected in their (unchallenged) immune system status.

We addressed this question in the workers of three species of stingless bees that vary in

important life history traits, not only with respect to nest defence, but also with the

workers' reproductive status.

Different from the honeybees, whose workers are functionally sterile in the

presence of the queen, the workers of stingless bees differ considerably in their potential

of laying eggs (Hartfelder et al., 2006; Vollet-Neto et al., 2018). While *F. varia* workers

are completely sterile, those of *M. quadrifasciata* and *S. postica* lay trophic and also

reproductive eggs in the presence of a queen, thus contributing to the colonies'

reproductive output. A key reproductive protein in female insects is vitellogenin (Vg)

(Sappington and Raikhel, 1998), but interestingly, in the functionally sterile honeybee

workers, the major function of Vg is not oogenesis. Rather, together with juvenile

hormone, it regulates the behavioural transition from intranidal (nursing) to extranidal

(foraging) behaviour (Amdam and Omholt, 2003), and it is in this context that the high

Vg levels in the haemolymph of the nurses prevent the onset of immunosenescence

(Amdam et al., 2005). Moreover, Vg itself is a powerful antioxidant protein that plays

key roles in the regulation of the lifespan of the honeybee queens and workers (Corona et

al., 2007; Havukainen et al., 2013; Seehuus et al., 2006).

In a recent study on gene expression underlying the nurse-to-forager transition in

stingless bees (de Souza and Hartfelder, 2023) we found an exactly opposite expression

pattern for the *Vg* gene in two species, showing upregulated transcript levels in the heads

of *F. varia* foragers*,* which are sterile, contrasting with upregulated *Vg* levels in the

abdomen of *M. quadrifasciata* nurses, which can lay eggs. Now taken together with the

current data on immunity gene expression, this suggests that, contingent on the

reproductive status of the workers, there appears to be a trade-off in the investment

between the Vg-related and the Toll/Imd-related immunity status. This trade-off not only

extends beyond the honeybees (Lourenço et al., 2012), it also appears to be species-

specifically balanced against the aggressiveness level in the stingless bees, as shown in

the current study.

Conclusion

The high transcript levels of the Toll and Imd pathway genes in the heads of the workers

of the aggressive species *S. postica*, compared to those in the docile workers of *M.*

quadrifasciata, indicate a possible non-canonical involvement of the innate immune

system in behaviours related to nest-defence strategies of social bees. The nest is an

important social factor that protects the society and its individuals from stressful

situations and, inherently, shapes their life histories (Walton et al., 2014). Given the

intricate interplay between the innate immune system and stress responses documented

in insects (Štětina et al., 2019; Hao et al., 2023), including bees (McMenamin et al., 2020),

made us, thus, raise the hypothesis of a possible association between the patterns of

immune gene expression and the levels aggressiveness in the workers of social bees.

Of specific interest in this context is the recent proteomics study by McAfee et al.

(2024), who investigated a possibly caste-specific reproduction-immunity trade-off in the

bumble bee *Bombus impatiens*. While they found no evidence for such a trade-off in the

workers, they did so for the queens. However, the authors consider this to be due to

confounding effects resulting from the queens' complex life cycle, including a diapause

phase for overwintering, followed by a solitary nest foundation before a functional colony

gets established. They conclude that additional factors, such as worker age and

aggressiveness should be taken into consideration. It is this question what we now

addressed in the stingless bees. Not only do their queens spend their entire life in the

centre of a well-protected nest, except for a brief mating flight, it is also the species

richness of the Meliponini and their extraordinary variability in the biology of their

workers that makes them a valuable source for comparative studies on life history

strategies and functional genomics in social bees.

As next a step, extending this study to other stingless bee species that aggressively

defend their nests, or aggressively perform raids on other colonies, like *Lestrimelitta*

limao, will be important to understand whether this non-canonical pattern of immune

system gene expression represents a general trait in these highly eusocial bees.

Furthermore, the development of behavioural assays that allow the quantification of

aggressiveness levels, such as the arena assay, coupled with pharmacological

manipulation and/or RNAi functional assays, should allow mechanistic insights into the

importance of the Toll and Imd pathways in the sophisticated social behaviours of the

bees.

COMPETING INTERESTS

The authors declare no competing or financial interests.

DATA AVAILABILITY

All relevant data can be found within the article and its supplementary information. The

raw data is available from the following link: https://doi.org/10.5281/zenodo.13374927

REFERENCES

Abbot, P., 2022. Defense in social insects: Diversity, division of labor, and evolution.

Annual Review of Entomology 67, 407–436. https://doi.org/10.1146/annurev-ento-

082521-072638

Amdam, G.V., Omholt, S.W., 2003. The hive bee to forager transition in honeybee

colonies: The double repressor hypothesis. *Journal of Theoretical Biology* 223, 451–

464. https://doi.org/10.1016/S0022-5193(03)00121-8

Amdam, G. V., Aase, A.L.T.O., Seehuus, S.-C., Kim Fondrk, M., Norberg, K.,

Hartfelder, K., 2005. Social reversal of immunosenescence in honey bee workers.

Experimental Gerontolology 40, 939–947.

https://doi.org/10.1016/j.exger.2005.08.004

Anderson, K. V, Bokla, L., Nüsslein-Volhard, C., 1985. Establishment of dorsal-ventral

polarity in the Drosophila embryo: The induction of polarity by the Toll gene

product. *Cell* 42, 791–798. https://doi.org/https://doi.org/10.1016/0092-

8674(85)90275-2

Avalos, A., Fang, M., Pan, H., Ramirez Lluch, A., Lipka, A.E., Dave Zhao, S., Giray, T.,

Robinson, G.E., Zhang, G., Hudson, M.E., 2020. Genomic regions influencing

aggressive behavior in honey bees are defined by colony allele frequencies.

Proceedings of the National Academy of Sciences of the United States of America

117, 17135–17141. https://doi.org/10.5061/dryad.q573n5tg8

Barajas-Azpeleta, R., Wu, J., Gill, J., Welte, R., Seidel, C., McKinney, S., Dissel, S., Si,

K., 2018. Antimicrobial peptides modulate long-term memory. *PLoS Genetics* 14,

e1007440. https://doi.org/10.1371/journal.pgen.1007440

Barribeau, S.M., Sadd, B.M., du Plessis, L., Brown, M.J., Buechel, S.D., Cappelle, K.,

Carolan, J.C., Christiaens, O., Colgan, T.J., Erler, S., Evans, J., Helbing, S., Karaus,

E., Lattorff, H.M.G., Marxer, M., Meeus, I., Näpflin, K., Niu, J., Schmid-Hempel,

R., Smagghe, G., Waterhouse, R.M., Yu, N., Zdobnov, E.M., Schmid-Hempel, P.,

2015. A depauperate immune repertoire precedes evolution of sociality in bees.

Genome Biology 16, 83. https://doi.org/10.1186/s13059-015-0628-y

Becker, T., Loch, G., Beyer, M., Zinke, I., Aschenbrenner, A.C., Carrera, P., Inhester, T.,

Schultze, J.L., Hoch, M., 2010. FOXO-dependent regulation of innate immune

homeostasis. *Nature* 463, 369–373. https://doi.org/10.1038/nature08698

Bhakar, A.L., Tannis, L.-L., Zeindler, C., Russo, M.P., Jobin, C., Park, D.S., MacPherson,

S., Barker, P.A., 2002. Constitutive Nuclear Factor-κB activity is required for central

neuron survival. *The Journal of Neuroscience* 22, 8466.

https://doi.org/10.1523/JNEUROSCI.22-19-08466.2002

Bittermann, M.E., Menzel, R., Fietz, A., Schäfer, S., 1983. Classical conditioning of

proboscis extension in honeybees (*Apis mellifera*). *Journal of Comparative*

Psychology 97, 107-119.

Blochtein, B., 2008. *Manual de boas práticas para a criação e manejo racional de*

abelhas sem ferrão no RS: Guaraipo, manduri e tubuna. EDIPUCRS, Porto Alegre.

Boleli, I.C., Paulino-Simões, Z.L., Gentile Bitondi, M.M., 1999. Cell death in ovarioles

causes permanent sterility in *Frieseomelitta varia* worker bees. *Journal of*

Morphology 242, 271–282. https://doi.org/10.1002/(SICI)1097-4687(199912)242:3

Breed, M.D., Page, R.E., 1991. Intra- and interspecific nestmate recognition in *Melipona*

workers (Hymenoptera: Apidae). *Journal of Insect Behavior* 4, 463–469.

https://doi.org/10.1007/BF01049331

Bulet, P., Stocklin, R., 2005. Insect antimicrobial peptides: Structures, properties and

gene regulation. *Protein & Peptide Letters* 12, 3–11.

https://doi.org/10.2174/0929866053406011

Camazine, S., 1986. Differential reproduction of the mite, *Varroa jacobso*ni

(Mesostigmata: Varroidae), on Africanized and European honey bees

(Hymenoptera: Apidae). *Annals of the Entomological Society of America* 79, 801–

803. https://doi.org/10.1093/aesa/79.5.801

Chandrasekaran, S., Ament, S.A., Eddy, J.A., Rodriguez-Zas, S.L., Schatz, B.R., Price,

N.D., Robinson, G.E., 2011. Behavior-specific changes in transcriptional modules

lead to distinct and predictable neurogenomic states. *Proceedings of the National*

Academy of Sciences of the United States of America 108, 18020–18025.

https://doi.org/10.1073/pnas.1114093108

Chandrasekaran, S., Rittschof, C.C., Djukovic, D., Gu, H., Raftery, D., Price, N.D.,

Robinson, G.E., 2015. Aggression is associated with aerobic glycolysis in the honey

bee brain1. *Genes, Brain and Behavior* 14, 158–166.

https://doi.org/10.1111/gbb.12201

Chapman, J.A., Kirkness, E.F., Simakov, O., Hampson, S.E., Mitros, T., Weinmaier, T.,

Rattei, T., Balasubramanian, P.G., Borman, J., Busam, D., Disbennett, K.,

Pfannkoch, C., Sumin, N., Sutton, G.G., Viswanathan, L.D., Walenz, B., Goodstein,

D.M., Hellsten, U., Kawashima, T., Prochnik, S.E., Putnam, N.H., Shu, S.,

Blumberg, B., Dana, C.E., Gee, L., Kibler, D.F., Law, L., Lindgens, D., Martinez,

D.E., Peng, J., Wigge, P.A., Bertulat, B., Guder, C., Nakamura, Y., Ozbek, S.,

Watanabe, H., Khalturin, K., Hemmrich, G., Franke, A., Augustin, R., Fraune, S.,

Hayakawa, E., Hayakawa, S., Hirose, M., Hwang, J.S., Ikeo, K., Nishimiya-

Fujisawa, C., Ogura, A., Takahashi, T., Steinmetz, P.R.H., Zhang, X., Aufschnaiter,

R., Eder, M.-K., Gorny, A.-K., Salvenmoser, W., Heimberg, A.M., Wheeler, B.M.,

Peterson, K.J., Böttger, A., Tischler, P., Wolf, A., Gojobori, T., Remington, K.A.,

Strausberg, R.L., Venter, J.C., Technau, U., Hobmayer, B., Bosch, T.C.G., Holstein,

T.W., Fujisawa, T., Bode, H.R., David, C.N., Rokhsar, D.S., Steele, R.E., 2010. The

dynamic   genome   of   *Hydra*.   *Nature*   464,   592–596.

https://doi.org/10.1038/nature08830

Choudhury, M.E., Mikami, K., Nakanishi, Y., Matsuura, T., Utsunomiya, R., Yano, H.,

Kubo, M., Ando, R., Iwanami, J., Yamashita, M., Nagai, M., Tanaka, J., 2022.

Insomnia and depressive behavior of MyD88-deficient mice: Relationships with

altered   microglial   functions.   *Journal of Neuroimmunology*   363,   577794.

https://doi.org/https://doi.org/10.1016/j.jneuroim.2021.577794

Cole, M.E., Ceja-Navarro, J.A., Mikaelyan, A., 2021. The power of poop: Defecation

behaviors   and   social   hygiene   in   insects.   *PLoS Pathogens*   17,   e1009964.

https://doi.org/10.1371/journal.ppat.1009964

Corona, M., Velarde, R.A., Remolina, S., Moran-Lauter, A., Wang, Y., Hughes, K.A.,

Robinson, G.E., 2007. Vitellogenin, juvenile hormone, insulin signaling, and queen

honey bee longevity. *Proceedings of the National Academy of Sciences of the United*

States of America 104, 7128–7133. https://doi.org/10.1073/pnas.0701909104

Couvillon, M.J., Wenseleers, T., Imperatriz-Fonseca, V.L., Nogueira-Neto, P., Ratnieks,

F.L.W., 2008. Comparative study in stingless bees (Meliponini) demonstrates that

nest entrance size predicts traffic and defensivity. *Journal of Evolutionary Biology*

21, 194–201. https://doi.org/10.1111/j.1420-9101.2007.01457.x

Cremer, S., Pull, C.D., Fürst, M.A., 2018. Social immunity: Emergence and evolution of

colony-level   disease   protection.   *Annual Review of Entomology*   63,   105–123.

https://doi.org/10.1146/annurev-ento-020117-043110

Danihlík, J., Aronstein, K., Petřivalský, M., 2015. Antimicrobial peptides: A key

component of honey bee innate immunity. *Journal of Apicultural Research*

https://doi.org/10.1080/00218839.2015.1109919

de Souza, L.D.R., Hartfelder, K., 2023. Reproductive potential shapes the expression of

nurse-to-forager transition genes in the workers of stingless bees (Meliponini).

Apidologie 54, 41. https://doi.org/10.1007/s13592-023-01021-z

Dierick, H.A., 2007. A method for quantifying aggression in male *Drosophila*

melanogaster.   *Nature*   *Protocols*   2,   2712–2718.

https://doi.org/10.1038/nprot.2007.404

Dresselhaus, E.C., Meffert, M.K., 2019. Cellular specificity of NF-κB function in the

nervous   system.   *Frontiers*   *in*   *Immunology*   10,   1043.

https://doi.org/10.3389/fimmu.2019.01043

Evans, J.D., Aronstein, K., Chen, Y.P., Hetru, C., Imler, J.-L., Jiang, H., Kanost, M.,

Thompson, G.J., Zou, Z., Hultmark, D., 2006. Immune pathways and defence

mechanisms in honey bees *Apis mellifera*. *Insect Molecular Biology* 15, 645–656.

https://doi.org/10.1111/j.1365-2583.2006.00682.x

Flatt, T., Amdam, G. V., Kirkwood, T.B.L., Omholt, S.W., 2013. Life-history evolution

and the polyphenic regulation of somatic maintenance and survival. *Quarterly*

Review of Biology 88, 185–218. https://doi.org/10.1086/671484

Freitas, F.C.P., Depintor, T.S., Agostini, L.T., Luna-Lucena, D., Nunes, F.M.F., Bitondi,

M.M.G., Simões, Z.L.P., Lourenço, A.P., 2019. Evaluation of reference genes for

gene expression analysis by real-time quantitative PCR (qPCR) in three stingless

bee species (Hymenoptera: Apidae: Meliponini). *Scientific Reports* 9, 17692.

https://doi.org/10.1038/s41598-019-53544-0

Freudenthal, R., Romano, A., 2000. Participation of Rel/NF-κB transcription factors in

long-term memory in the crab *Chasmagnathus*. *Brain Research* 855, 274–281.

https://doi.org/10.1016/S0006-8993(99)02358-6

Friard, O., Gamba, M., 2016. BORIS: a free, versatile open-source event-logging

software for video/audio coding and live observations. *Methods in Ecology and*

Evolution 7, 1325–1330. https://doi.org/10.1111/2041-210X.12584

Grüter, C., 2020. *Stingless Bees*, *Their Behaviour, Ecology and Evolution.* Springer

International Publishing, Cham. https://doi.org/10.1007/978-3-030-60090-7

Hanson, M.A., Lemaitre, B., 2020. New insights *on Drosophila* antimicrobial peptide

function in host defense and beyond. *Current Opinion in Immunology* 62, 22–30

https://doi.org/10.1016/j.coi.2019.11.008

Hao, Z.H., Lu, Q.Y., Zhou, Y.Y., Liang, Y.T., Gao, Y., Ma, H.Y., Xu, Y.S., Wang, H.B.,

2023. Molecular characterization of MyD88 as a potential biomarker for pesticide-

induced stress in *Bombyx mori*. *Pesticide Biochemistry and Physiology* 196, 105610.

https://doi.org/10.1016/j.pestbp.2023.105610.

Hartfelder, K., Makert, G.R., Judice, C.C., Pereira, G.A.G., Santana, W.C., Dallacqua,

R., Bitondi, M.M.G., 2006. Physiological and genetic mechanisms underlying caste

development, reproduction and division of labor in stingless bees. *Apidologie* 37,

144–163. https://doi.org/10.1051/apido:2006013

Havukainen, H., Munch, D., Baumann, A., Zhong, S., Halskau, O., Krogsgaard, M.,

Amdam, G. V, 2013. Vitellogenin recognizes cell damage through membrane

binding and shields living cells from reactive oxygen species. *Journal of Biological*

Chemistry 288, 28369–28381. https://doi.org/10.1074/jbc.M113.465021

Hoffmann, J.A., 1995. Innate immunity of insects. *Current Opinion in Immunology* 7, 4–

10. https://doi.org/https://doi.org/10.1016/0952-7915(95)80022-0

Horng, T., Medzhitov, R., 2001. *Drosophila* MyD88 is an adapter in the Toll signaling

pathway. *Proceedings of the National Academy of Sciences of the United States of*

America 98, 12654–12658. https://doi.org/10.1073/pnas.231471798

Hosoi, T., Yamawaki, Y., Kimura, H., Honda, S., Ozawa, K., 2021. Possible involvement

of MyD88 in regulating stress response in mice. *Frontiers in Neuroscience* 15,

621446. https://doi.org/10.3389/fnins.2021.621446

Imler, J.-L., Hoffmann, J.A., 2001. Toll receptors in innate immunity. *Trends in Cell*

Biology 11, 304–311. https://doi.org/10.1016/S0962-8924(01)02004-9

Janeway, C.A., Medzhitov, R., 2002. Innate immune recognition. *Annual Review of*

Immunology https://doi.org/10.1146/annurev.immunol.20.083001.084359

Kaltschmidt, C., Kaltschmidt, B., Neumann, H., Wekerle, H., Baeuerle, P.A., 1994.

Constitutive NF-kB activity in neurons. *Molecular and Cellular Biology* 14, 3981–

3992. https://doi.org/10.1128/mcb.14.6.3981-3992.1994

Kärcher, M.H., Ratnieks, F.L.W., 2009. Standing and hovering guards of the stingless

bee *Tetragonisca angustula* complement each other in entrance guarding and

intruder recognition. *Journal of Apicultural Research* 48, 209–214.

https://doi.org/10.3896/IBRA.1.48.3.10

Kim, D.H., Ausubel, F.M., 2005. Evolutionary perspectives on innate immunity from the

study of *Caenorhabditis elegans*. *Current Opinion in Immunology* 17, 4–10.

https://doi.org/10.1016/j.coi.2004.11.007

Kingsolver, M.B., Huang, Z., Hardy, R.W., 2013. Insect antiviral innate immunity:

Pathways, effectors, and connections. *Journal of Molecular Biolog*y 425, 4921–

4936. https://doi.org/10.1016/j.jmb.2013.10.006

Koludarov, I., Velasque, M., Senoner, T., Timm, T., Greve, C., Hamadou, A. Ben, Gupta,

D.K., Lochnit, G., Heinzinger, M., Vilcinskas, A., Gloag, R., Harpur, B.A.,

Podsiadlowski, L., Rost, B., Jackson, T.N.W., Dutertre, S., Stolle, E., von Reumont,

B.M., 2023. Prevalent bee venom genes evolved before the aculeate stinger and

eusociality. *BMC Biology* 21, 229. https://doi.org/10.1186/s12915-023-01656-5

Kounatidis, I., Chtarbanova, S., Cao, Y., Hayne, M., Jayanth, D., Ganetzky, B.,

Ligoxygakis, P., 2017. NF-κB immunity in the brain determines fly lifespan in

healthy aging and age-related neurodegeneration. *Cell Reports* 19, 836–848.

https://doi.org/10.1016/j.celrep.2017.04.007

Leulier, F., Lemaitre, B., 2008. Toll-like receptors - taking an evolutionary approach.

Nature Reviews Genetics https://doi.org/10.1038/nrg2303

Li, S., Yu, X., Feng, Q., 2019. Fat body biology in the last decade. *Annual Review of*

Entomology 64, 315–333. https://doi.org/10.1146/annurev-ento-011118-112007

Li-Byarlay, H., Rittschof, C.C., Massey, J.H., Pittendrigh, B.R., Robinson, G.E., 2014.

Socially responsive effects of brain oxidative metabolism on aggression.

Proceedings of the National Academy of Sciences of the United States of America

111, 12533–12537. https://doi.org/10.1073/pnas.1412306111

Lilienbaum, A., Israël, A., 2003. From calcium to NF-κB signaling pathways in neurons.

Molecular   and   Cellular   Biology   23,   2680–2698.

https://doi.org/10.1128/MCB.23.8.2680-2698.2003

Livak, K.J., Schmittgen, T.D., 2001. Analysis of relative gene expression data using Real-

Time quantitative PCR and the $2^{-\Delta\Delta CT}$ method. *Methods* 25, 402–408.

Lourenço, A.P., Mackert, A., Cristino, A.D.S., Simões, Z.L.P., 2008. Validation of

reference genes for gene expression studies in the honey bee, *Apis mellifera*, by

quantitative   real-time   RT-PCR.   *Apidologie*   39,   372–385.

https://doi.org/10.1051/apido:2008015

Lourenco, A.P., Martins, J.R., Guidugli-Lazzarini, K.R., Macedo, L.M., Bitondi, M.M.,

Simoes, Z.L., 2012. Potential costs of bacterial infection on storage protein gene

expression and reproduction in queenless *Apis mellifera* worker bees on distinct

dietary regimes. *Journal of Insect Physiology* 58, 1217–1225.

https://doi.org/10.1016/j.jinsphys.2012.06.006

Lourenço, A.P., Guidugli-Lazzarini, K.R., Freitas, F.C.P., Bitondi, M.M.G., Simões,

Z.L.P., 2013. Bacterial infection activates the immune system response and

dysregulates microRNA expression in honey bees. *Insect Biochemistry and*

Molecular Biology 43, 474–482. https://doi.org/10.1016/j.ibmb.2013.03.001

Lourenço, A.P., Florecki, M.M., Simões, Z.L.P., Evans, J.D., 2018. Silencing of *Apis*

mellifera dorsal genes reveals their role in expression of the antimicrobial peptide

defensin-1. *Insect Molecular Biology* 27, 577–589.

https://doi.org/10.1111/imb.12498

Lourenço, A.P., Martins, J.R., Torres, F.A.S., Mackert, A., Aguiar, L.R., Hartfelder, K.,

Bitondi, M.M.G., Simões, Z.L.P., 2019. Immunosenescence in honey bees (*Apis*

mellifera L.) is caused by intrinsic senescence and behavioral physiology.

Experimental Gerontology 119, 174–183.

https://doi.org/10.1016/j.exger.2019.02.005

Lynch, J.A., Roth, S., 2011. The evolution of dorsal-ventral patterning mechanisms in

insects. *Genes & Development* 25, 107–118. https://doi.org/10.1101/gad.2010711

McAfee, A., Chapman, A., Bao, G., Tarpy, D.R., Foster, L.J., 2024. Investigating trade-

offs between ovary activation and immune protein expression in bumble bee

(*Bombus impatiens*) workers and queens. *Proceedings of the Royal Society B:*

Biological Sciences 291, 20232463. https://doi.org/10.1098/rspb.2023.2463

McMenamin, A.J., Daughenbaugh, K.F., Flenniken, M.L., 2020. The heat shock response

in the Western honey bee (*Apis mellifera*) is antiviral. *Viruses (Basel)* 12, 245.

htpps://doi.org/.10.3390/v12020245

Meffert, M.K., Baltimore, D., 2005. Physiological functions for brain NF-κB. *Trends in*

Neuroscience 28, 37–43. https://doi.org/10.1016/j.tins.2004.11.002

Meffert, M.K., Chang, J.M., Wiltgen, B.J., Fanselow, M.S., Baltimore, D., 2003. NF-κB

functions in synaptic signaling and behavior. *Nature Neuroscience* 6, 1072–1078.

https://doi.org/10.1038/nn1110

Melo, G.A.R., 2021. Stingless Bees (Meliponini), in: Starr C (ed) *Encyclopedia of Social*

Insects.   Springer   International   Publishing,   Cham,   pp.   883–900.

https://doi.org/10.1007/978-3-030-28102-1_117

Mémet, S., 2006. NF-κB functions in the nervous system: From development to disease.

Biochemical   Pharmacology   72,   1180–1195.

https://doi.org/10.1016/j.bcp.2006.09.003

Merlo, E., Freudenthal, R., Maldonado, H., Romano, A., 2005. Activation of the

transcription factor NF-κB by retrieval is required for long-term memory

reconsolidation. L*earning & Memory* 12, 23–29. https://doi.org/10.1101/lm.82705

Nogueira-Neto, P., 1997. *Vida e Criação de Abelhas Indígenas Sem Ferrão*. Nogueirapis,

São Paulo.

Nouvian, M., Reinhard, J., Giurfa, M., 2016. The defensive response of the honeybee

Apis   mellifera.   Journal   of   Experimental   Biology   219,   3505–3517.

https://doi.org/10.1242/jeb.143016

Nunes, T.M., Von Zuben, L.G., Costa, L., Venturieri, G.C., 2014. Defensive repertoire

of the stingless bee *Melipona flavolineata* Friese (Hymenoptera: Apidae).

Sociobiology 61, 541–546. https://doi.org/10.13102/sociobiology.v61i4.541-546

Pizzorno, M.C., Field, K., Kobokovich, A.L., Martin, P.L., Gupta, R.A., Mammone, R.,

Rovnyak, D., Capaldi, E.A., 2021. Transcriptomic responses of the honey bee brain

to infection with deformed wing virus. *Viruses (Basel)* 13, 287.

https://doi.org/10.3390/v13020287

Ratnieks, F.L., Foster, K.R., Wenseleers, T., 2006. Conflict resolution in insect societies.

Annual Review of Entomology 51, 581-608. https://doi:

10.1146/annurev.ento.51.110104.151003

Rittschof, C.C., 2017. Sequential social experiences interact to modulate aggression but

not brain gene expression in the honey bee (*Apis mellifera*). *Frontiers in Zoology*

14, 16. https://doi.org/10.1186/s12983-017-0199-8

Rittschof, C.C., Vekaria, H.J., Palmer, J.H., Sullivan, P.G., 2018. Brain mitochondrial

bioenergetics change with rapid and prolonged shifts in aggression in the honey bee,

Apis mellifera. Journal of Experimental Biology 221, jeb176917.

https://doi.org/10.1242/jeb.176917

Rittschof, C.C., Rubin, B.E.R., Palmer, J.H., 2019. The transcriptomic signature of low

aggression in honey bees resembles a response to infection. *BMC Genomics* 20,

1029. https://doi.org/10.1186/s12864-019-6417-3

Roubik, D.W., 2023. Stingless bee (Apidae: Apinae: Meliponini) ecology. *Annual Review*

of Entomology 68, 231–256. https://doi.org/10.1146/annurev-ento-120120-103938

Sappington, T.W., S. Raikhel, A., 1998. Molecular characteristics of insect vitellogenins

and vitellogenin receptors. *Insect Biochemistry and Molecular Biology* 28, 277–300.

https://doi.org/10.1016/S0965-1748(97)00110-0

Schwenke, R.A., Lazzaro, B.P., Wolfner, M.F., 2016. Reproduction-immunity trade-offs

in insects. *Annual Review of Entomology* 61, 239–256.

https://doi.org/10.1146/annurev-ento-010715-023924

Seehuus, S.C., Norberg, K., Gimsa, U., Krekling, T., Amdam, G., 2006. Reproductive

protein protects functionally sterile honey bee workers from oxidative stress.

Proceedings of the National Academy of Sciences of the United States of America

103, 962–967. https://doi.org/10.1073/pnas.0502681103

Sétima, T., Poupardin. R., Moos, M., Šimek, P., Šmilauer, P., Koštál, V., 2019. Larvae of

Drosophila melanogaster exhibit transcriptional activation of immune response

pathways and antimicrobial peptides during recovery from supercooling stress.

Insect Biochemistry and Molecular Biology 105, 60-68.

htpps://doi.org/10.1016/j.ibmb.2019.01.006

Shackleton, K., Toufailia, H. Al, Balfour, N.J., Nascimento, F.S., Alves, D.A., Ratnieks,

F.L.W., 2015. Appetite for self-destruction: Suicidal biting as a nest defense strategy

in *Trigona* stingless bees. *Behavioral Ecology and Sociobiology* 69, 273–281.

https://doi.org/10.1007/s00265-014-1840-6

Shanahan, M., Spivak, M., 2021. Resin use by stingless bees: A review. *Insects* 12, 719.

https://doi.org/10.3390/insects12080719

Shpigler, H.Y., Saul, M.C., Corona, F., Block, L., Cash Ahmed, A., Zhao, S.D.,

Robinson, G.E., 2017. Deep evolutionary conservation of autism-related genes.

Proceedings of the National Academy of Sciences of the United States of America

114, 9653–9658. https://doi.org/10.1073/pnas.1708127114

Smith, B.H., Burden, C.M., 2014. A proboscis extension response protocol for

investigating behavioral plasticity in insects: application to basic, biomedical, and

agricultral research. *Journal of Visual Experiments* 91, e51057.

https://doi.org/10.3791/51057

Stączek, S., Cytryńska, M., Zdybicka-Barabas, A., 2023. Unraveling the role of

antimicrobial peptides in insects. *International Journal of Molecular Science* 24,

5753. https://doi.org/10.3390/ijms24065753

Stearns, S., 1992. *The Evolution of Life Histories*. Oxford University Press, Oxford.

Strand, M.R., 2008. The insect cellular immune response. *Insect Science*

https://doi.org/10.1111/j.1744-7917.2008.00183.x

Stuart, B.A.R., Franitza, A.L., Lezi, E., 2022. Regulatory roles of antimicrobial peptides

in the nervous system: Implications for neuronal aging. *Frontiers in Cellular*

Neuroscience 16, 843790. https://doi.org/10.3389/fncel.2022.843790

The Honeybee Genome Sequencing Consortium, 2006. Insights into social insects from

the genome of the honeybee *Apis mellifera*. *Nature* 443, 931–949.

https://doi.org/10.1038/nature05260

Traniello, I.M., Bukhari, S.A., Dibaeinia, P., Serrano, G., Avalos, A., Ahmed, A.C.,

Sankey, A.L., Hernaez, M., Sinha, S., Zhao, S.D., Catchen, J., Robinson, G.E., 2023.

Single-cell dissection of aggression in honeybee colonies. *Nature Ecology and*

Evolution 7,1232-1244. https://doi.org/10.1038/s41559-023-02090-0

Vollet-Neto, A., Koffler, S., dos Santos, C.F., Menezes, C., Nunes, F.M.F., Hartfelder,

K., Imperatriz-Fonseca, V.L., Alves, D.A., 2018. Recent advances in reproductive

biology of stingless bees. *Insectes Sociaux* 65, 201–212.

https://doi.org/10.1007/s00040-018-0607-x

Walton, A., Herman, J.J., Rueppell, O., 2024. Social life results in social stress protection:

a novel concept to explain individual life-history patterns in social insects.

Biological Reviews 99, 1444–1457. htpps://doi.org/10.1111/brv.13074

Wellmann, H., Kaltschmidt, B., Kaltschmidt, C., 2001. Retrograde transport of

transcription factor NF-κB in living neurons. *Journal of Biological Chemistry* 276,

11821–11829. https://doi.org/https://doi.org/10.1074/jbc.M009253200

Wilson-Rich, N., Dres, S.T., Starks, P.T., 2008. The ontogeny of immunity: development

of innate immune strength in the honey bee (*Apis mellifera*). *Journal of Insect*

Physiology 54, 1392-1399. https://doi: 10.1016/j.jinsphys.2008.07.016

Yao, X.P., Ye, J., Feng, T., Jiang, F.C., Zhou, P., Wang, F., Chen, J.G., Wu, P.F., 2023.

Adaptor protein MyD88 confers the susceptibility to stress via amplifying immune

danger   signals.   *Brain   Behavior   and   Immunity*   108,   204–220.

https://doi.org/10.1016/j.bbi.2022.12.007

Zhai, Z., Huang, X., Yin, Y., 2018. Beyond immunity: The Imd pathway as a coordinator

of host defense, organismal physiology and behavior. *Developmental &*

Comparative Immunology 83, 51–59. https://doi.org/10.1016/j.dci.2017.11.008

Zhang, W., Tettamanti, G., Bassal, T., Heryanto, C., Eleftherianos, I., Mohamed, A.,

2021. Regulators and signalling in insect antimicrobial innate immunity: Functional

molecules   and   cellular   pathways.   *Cell   Signaling*   83,   110003.

https://doi.org/10.1016/j.cellsig.2021.110003

1046

1047

1048

Figure 1 – The stingless bee species investigated in this study and their major life history traits. Images are credited to Dr. Cristiano Menezes.

Figure 2 – Arena test results on aggressiveness of individual workers of three stingless bee species. (A) Attack frequency and fatal attack frequencies for *S. postica*, *F. varia*, and *M. quadrifasciata*. The dark red colour represents the percentage of fatal attackers, i.e., those that killed the intruder, the red colour represents the percentage of attackers that did not kill the intruder, and the grey colour represents the percentage of bees that did not attack the intruder. (B) Number of antennation events: *S. postica* and *F. varia* exhibit have higher number of antennation events when compared to *M. quadrifasciata* (Kruskal-Wallis: $H = 20.02$, $P < 0.001$). (C) Latency to attack: *S. postica* exhibits the shortest latency to attack compared to *F. varia* and *M. quadrifasciata,* with no significant differences among the latter two species (Kruskal-Wallis: $H = 21.05$, $P < 0.001$). (D) Attacking index: *S. postica* has a higher attacking index than *F. varia* and *M. quadrifasciata* (Kruskal-Wallis: $H = 23.31$, $P < 0.001$). (E) Pursuit index: All three species showed differences in the chasing index (Kruskal-Wallis: $H = 6.09$, $P < 0.05$); however, pairwise comparisons revealed no significant differences. Orange: *S. postica* (Sp), Yellow: *F. varia* (Fv), Black: *M. quadrifasciata* (Mq). Different letters indicate statistically significant differences among the three species. The error bars represent the standard error of the mean (SEM).

Figure 3 – Relative expression levels of four immune system genes in the heads and abdomens of *Scaptotrigona postica* guards and foragers. Bar plots representing the transcript levels for (A) *Toll,* (B) *Myd88*, (C) *Dorsal,* and (D) *Imd*. Blue: Guard (G) and

Green: Forager (F). Different letters indicate statistically significantly differences among

the three live cycle stages within each body compartment. Asterisks indicate statistical

differences between the two body compartments (GLMM). Symbols (***) $P < 0.0001$,

(**) $P < 0.001$, (*) $P < 0.05$, (n.s.) non-significant. Sample size per colony, $N = 7$; per life

cycle stage within each body compartment, $N = 21$ (3 colonies). The error bars represent

the standard error of the mean (SEM).

1080

Figure 4 – Relative expression levels of four immune system genes in the heads and

abdomens of *Melipona quadrifasciata* nurses, guards, and foragers. Bar plots

representing the transcript levels for (A) *Toll,* (B) *Myd88*, (C) *Dorsal,* and (D) *Imd.*

Purple: Nurse (N), Blue: Guard (G), Green: Forager (F). Different letters indicate

statistically significantly differences among the three live cycle stages within each body

compartment. Asterisks indicate statistical differences between the two body

compartments (GLMM). Symbols (***) $P < 0.0001$, (**) $P < 0.001$, (*) $P < 0.05$, (n.s.)

non-significant. Sample size per colony, $N = 7$; per life stage within each body

compartment, $N = 21$ (3 colonies). The error bars represent the standard error of the mean

(SEM).

1091

Figure 5 – Relative expression levels of four immune system genes in the heads and

abdomens of *Frieseomelitta varia* nurses, guards, and foragers. Bar plots representing

the transcript levels for (A) *Toll,* (B) *Myd88*, (C) *Dorsal,* and (D) *Imd.* Purple: Nurse (N),

Blue: Guard (G), Green: Forager (F). Different letters indicate statistically significantly

differences among the three live cycle stages within each body compartment. Asterisks

indicate statistical differences between the two body compartments (GLMM). Symbols

(***) $P < 0.0001$, (**) $P < 0.001$, (*) $P < 0.05$, (n.s.) non-significant. Sample size per

colony, $N = 7$; per life stage within each body compartment, $N = 21$ (3 colonies). The

error bars represent the standard error of the mean (SEM).

1101

Figure 6 – *Abaecin* gene expression in the heads and abdomens of the workers of stingless

bees. Bar plots for *Abaecin* gene expression of *S. postica* (upper left) and *F. varia* (upper

right) showing no differences between the two body compartments. *M. quadrifasciata*

workers (lower left) and *A. mellifera* (lower right) have higher *Abaecin* gene expression

levels in the abdomen. Purple: Nurse (N), Blue: Guard (G), Green: Forager (F). Different

letters indicate significant differences within each body compartment. Asterisks indicate

statistical differences between the two body compartments (GLMM for *S. postica*, *F.*

varia, and *M. quadrifasciata*; GLM for *A. mellifera*). Symbols (***) $P < 0.0001$, (**) P

< 0.001, (*) $P < 0.05$, (n.s.) non-significant. Sample size per colony, $N = 7$; per life stage

within each body compartment, $N = 21$ (3 colonies for each stingless bee species); $N = 7$

for *A. mellifera* (one colony). The error bars represent the standard error of the mean

(SEM).

1114

Figure 7 – Relative expression levels of four immune system genes in the heads and

abdomens of honeybee, *Apis mellifera,* workers. Bar plots representing the transcript

levels for (A) *Toll,* (B) *Myd88,* (C) *Dorsal,* and (D) *Imd*. Purple: Nurse (N), Blue: Guard

(G), Green: Forager (F). Different letters indicate statistically significantly differences

among the three live cycle stages within each body compartment. Asterisks indicate

statistical differences between the two body compartments (GLM). Symbols (***) $P <$

0.0001, (**) $P < 0.001$, (*) $P < 0.05$, (n.s.) non-significant. Sample size per life stage

within each body compartment, $N = 7$. The error bars represent the standard error of the

mean (SEM).

SUPPLEMENTARY MATERIAL

Figure S1 – Effect of dietary restriction on the number of attacking and non-attacking workers.

Figure S2 – Images illustrating the phenotypic characteristics of the workers of the three stingless bee species used in this study

Figure S3 – Ct values of the two reference genes (*Rps18* and *Rpl32*) across behavioral subcastes and tissues in the *S. postica*, *F. varia* e *M. quadrifasciata*.

Table S1 - Primers used in the RT-qPCR assays.

Table S2 - Indices of aggressiveness for individual workers of stingless bees tested in arena assays.

Table S3 - Statistical details of the GLMM test used to analyse the expression of immune pathways genes in *S. postica*.

Table S4 - Statistical details of the GLMM test used to analyse the expression of immune pathways genes in *M. quadrifasciata*.

Table S5 - Statistical details of the GLMM test used to analyse the expression of immune pathways genes in *F. varia*.

1149   **Table S6 -** Statistical details of the GLM test used to analyse the expression of immune

1150   pathways genes in *A. mellifera.*

1151

ABSTRACT

Living in crowded perennial nests, social insects encounter high risks of disease outbreaks

and predation on colony resources. The innate immune system plays an important barrier

against microbial pathogens, but its association with group defensive behaviours remains

unexplored. Here we investigated the expression of innate immune system genes in the

workers of three species of Brazilian stingless bees (*Scaptotrigona postica*,

Frieseomelitta varia, and *Melipona quadrifasciata*), which differ in worker reproductive

traits and colony defence strategies. First, we standardized a behavioural arena test to

assess the species-specific individual aggressiveness level of the workers. Then, we

quantified the transcript levels of four core genes of the Toll and Imd pathways in the

heads and abdomens for the three main life cycle stages of the workers (nurse, guard and

forager). For the heads we found a strong association between immunity gene expression

and aggressiveness, with an upregulation in the heads of guards and foragers in

comparison to their abdomen tissues of the aggressive species, *S. postica*. while the less

aggressive species, especially *M. quadrifasciata* workers, did not show such strong up-

regulation. Our study establishes a potential molecular link between immunity pathways

and complex social behaviours in the Brazilian stingless bees, which evolutionarily lost

the sting apparatus for the defence of their colonies.

Keywords: Stingless bee; colony defence; innate immunity; life history; *Apis mellifera,*

social behaviour.

BULLETED HIGHLIGHTS

• The expression of immune system genes increases with social task/age in the

abdomen and head of highly eusocial bees.

• The expression of Toll pathway genes is highly upregulated in the heads of the

more aggressive stingless bee species.

• In the more docile stingless bee species, immune gene expression is higher in the

abdomen than in the head.

• Our findings suggest a possible involvement of immunity pathway signalling in

the regulation of key social behaviours.

ABBREVIATIONS

AMP – Antimicrobial peptide

GLM – General Linear Model

GLMM – General Linear Mixed Model

NF-kB - Nuclear Factor-kappa B

RT-qPCR - Quantitative reverse transcription polymerase chain reaction

TLRs – Toll-like receptors

Vg - Vitellogenin

INTRODUCTION

Social bees, including the honeybee and the stingless bees, are territorial animals that build perennial nests where they store food resources (honey and pollen) and rear and protect the brood and the highly reproductive queen. Such social life presents two types of risks. On the one hand, the nutritious resources in the colony can attract a variety of predators and nest parasites, and on the other, the large number of individuals in these nests represents an elevated risk for the spreading of diseases (Abbot, 2022). Yet strikingly, the analysis of the honeybee genome (The Honeybee Genome Sequencing Consortium, 2006) indicated a reduction in the number of genes comprising the bees' immune response repertoire compared to other insects, such as *Drosophila melanogaster* and *Anopheles gambiae*. While this loss can be compensated by social behaviours (social immunity) that provide barriers to the spreading of diseases (Cremer et al., 2018), a reduction in the repertoire of immune system genes, however, was also observed in bee species with a solitary lifestyle (Barribeau et al., 2015), apparently representing an ancestral trait in the bees, and thus unrelated to the evolution of advanced sociality in this clade.

For the defence of their nests, social bees adopt different collective behaviours. In the honeybee, *Apis mellifera*, guard workers use the sting and also biting as the primary strategies against intruders (Abbot, 2022; Nouvian et al., 2016). In contrast, having evolutionarily lost their sting apparatus, the stingless bees (Meliponini) exhibit diverse defence strategies. Some launch mass attacks, biting invaders, some apply sticky resins on them, while others have nest entrances that are difficult to access (for reviews see Grüter, 2020; Roubik, 2023; Shanahan and Spivak, 2021). Also, guards can respond to invasion threats by attacking aggressively, releasing alarm pheromones, by forming a

corridor of hovering guards, or by simply blocking a narrow nest entrance (Kärcher and

Ratnieks, 2009).

Molecular signatures of the defensive (aggressive) behaviours against threats have

been mainly studied in the Western honeybee, *A. mellifera*, and a strong correlation was

found between the colony level of aggression and the frequency of certain alleles in the

population (Avalos et al., 2020). At the individual level, workers were found to exhibit

distinct gene expression profiles in their brains, differentiating more aggressive

individuals from more docile ones (Rittschof, 2017). Such individual tendencies in

aggressiveness are, however, masked by the prevailing behavioural disposition of the

colony at any given moment, indicating that the colony genetic composition may override

the individual workers' genetic background in the decision to place or not an attack

(Avalos et al., 2020). At the cellular level, a recent transcriptomics study (Traniello et al.,

2023) revealed genomic subnetworks associated with colony aggression across various

cell types and across brain regions involved in olfaction, vision, and multimodal sensory

integration.

Previous studies have also demonstrated a correlation between aggressiveness and

immune system activity. For instance, the highly aggressive Africanized hybrids of *A.*

mellifera present strong traits of resistance against ectoparasites compared to more docile

A. mellifera subspecies (Camazine, 1986). Also, relatively docile, healthy workers, as

well as workers raised in less aggressive colonies, presented a gene transcriptional profile

in the brain and in the abdominal fat body that was very similar to that of individuals

suffering from acute pathogen infection (Rittschof et al., 2019), indicating that low

aggression resembles a sickness-like behavioural state. Furthermore, transcriptional

regulatory network analyses demonstrated that the neuro-genomic behavioural state of

aggressive honeybee workers is characterized by high expression levels of the

transcription factor Dorsal (Chandrasekaran et al., 2011).

Dorsal, a Nuclear Factor-kappa B (NF-kB) protein in insects, is a transcription

factor that plays a key role in the Toll signalling pathway of the innate immune response

of insects (Hoffmann, 1995), similar to NF-kB in the Toll-like receptors (TLR) signalling

pathway in vertebrates (Imler and Hoffmann, 2001). Unlike vertebrates however,

invertebrates rely solely on innate immunity as their defence against pathogens. This

system consists of humoral and cellular responses mediated by haemocytes and

antimicrobial peptides (AMPs) (Danihlík et al., 2015; Janeway and Medzhitov, 2002;

Lourenço et al., 2018; Strand, 2008). In this context, the Toll signalling pathway plays a

vital role in the immune response of insects. When activated in response to recognition

proteins that detect specific components of the cell wall of fungi and Gram-positive

bacteria, the Toll pathway triggers the translocation of the transcription factor Dorsal to

the nucleus, leading to the transcription of AMP genes (Evans et al., 2006; Kingsolver et

al., 2013; Zhang et al., 2021).

Similarly, the Imd signalling pathway is triggered by the recognition of

peptidoglycans from Gram-negative bacteria, resulting in the activation of the NF-kB

transcription factor Relish and, consequently, the expression of specific AMPs (Zhai et

al., 2018). In bees, the Toll and Imd pathways are known to regulate the expression of

AMPs, such as Abaecin and Defensin-1 (Lourenço et al., 2013, 2018). These pathways

are, however, not restricted to immune functions. In fact, as indicated by the primary

name 'Dorsal', this gene was first identified in *Drosophila* embryonic mutant screens as

a key transcription factor in the determination of the dorsal-ventral embryonic axis

(Lynch and Roth, 2011).

Both, innate immunity and aggressive behaviours are aspects related to somatic maintenance, and both are energetically costly and, thus, likely of relevance to trade-offs in life histories (Flatt et al., 2013; Stearns, 1992). In general, reproductive activity and immune responses frequently show a negative correlation in insects, a pattern observed in both solitary species from orders such as Orthoptera, Diptera, Odonata, Hemiptera, and Coleoptera, as well as in some social insects, such as certain ant species (Schwenke et al., 2016). But how does this correlation look like in social insects, especially in the highly eusocial ones, where colony (somatic) maintenance and reproductive activity are dissociated into two morphologically distinct castes, the workers and the queen?

Studies on social insect immunity, colony defence strategy, reproductive activity, and their associated gene networks have been mainly conducted using *A. mellifera* workers as a model. However, the honeybees (Apini) represent a single genus (*Apis*) consisting of less than a dozen species, and thus are a relatively small and rather homogeneous group in the phylogeny of the bees (Apidae). In contrast, the stingless bees (Meliponini) comprise over 500 recognized species distributed among 58 genera (Grüter, 2020; Melo, 2021). With a pantropical distribution, they exhibit a high level of biological diversity in terms of nesting sites, reproductive strategies, and nest defence. Stingless bees, like honeybees, are threatened by anthropogenic impacts, such as crop monoculture and pesticide use. However, unlike honeybees, they thrive in greenhouse environments, where they are increasingly recognized as efficient pollinators in high-value agriculture. (Grüter, 2020).

Taking advantage of the stingless bees' biological variability, we set out in this study to investigate the relationship between aggressiveness (defence behaviour) and the immune system profile of the workers of three stingless bee species with distinct life history configurations (Fig. 1). The workers of *Scaptotrigona postica* are very aggressive

in their nest defence and are able to lay eggs in the presence of the queen (Nogueira-Neto,

1997). The workers of *Melipona quadrifasciata* are also able to lay eggs in the presence

of the queen, but they are notably docile (Grüter, 2020; Nogueira-Neto, 1997), and the

workers of *Frieseomelitta varia,* while moderately defensive, are completely sterile

(Boleli et al., 1999; Nogueira-Neto, 1997). Hence, we considered that this contrast in

character combinations would allow us to address the question as to how life histories

and their respective trade-offs (Stearns, 1988), specifically those between the workers'

aggressiveness in the nest defence and their reproductive potential became configured

with respect to a costly system, such as the innate immune response. Specifically, while

on the one hand the investment in aggressiveness is likely very costly to the individual

worker, but benefits the colony fitness, worker reproduction on the other hand creates a

fitness conflict, primarily with the queen, but in the honey bee, also with the other workers

(Ratnieks et al., 2006).

Hence, the underlying hypotheses that we proposed to address were that: (a) even

in the absence of a specific immune challenge, the expression levels of immunity-related

genes increase with the age of the bees, reflecting their progression in social behaviours

(Wilson-Rich et al., 2008), and (b) the expression levels reflect the level of aggressiveness

and the reproductive biology of the workers. To do so, we adapted an arena assay that

allowed us to quantitatively assess the species-specific levels of aggressiveness for

individual workers, and we evaluated the expression levels of a set of four key genes of

the Toll and Imd pathways in the heads and abdomens for three important behavioural

stages in the workers' life cycle. Finally, we compared the results obtained for the three

stingless bees with those of a corresponding analysis that we performed on honeybee

workers.

MATERIAL AND METHODS

The arena behavioural assay: A quantitative tool for assessing the species-specific

aggressiveness level of individual workers

To obtain information about the aggressiveness at the individual level for the workers of

the three focal stingless bee species, we performed an arena assay based on the protocol

of Dierick (2007) for *Drosophila melanogaster*, which has subsequently been applied to

honeybees by Shpigler et al. (2017). This assay eliminates variables associated with the

social colony context, including the presence of alarm pheromones elicited by nestmates,

as well as environmental factors, such as ambient temperature, group size, diet, time, and

territorial size in the bees' reaction against a perceived threat.

Workers of *Scaptotrigona postica* (Latreille, 1807), *Frieseomelitta varia*

(Lepeletier, 1836), and *Melipona quadrifasciata* (Lepeletier, 1836) were collected from

one single colony of each species, all maintained in the apiary at the Department of

Genetics, Ribeirão Preto Campus, University of São Paulo, Brazil. From each colony of

S. postica and *F. varia* (Lepeletier, 1836), 100 newly-emerged workers and 10 nurse-

stage workers were collected and kept together in a single cage (8 x 11 x 13 cm) in an

incubator at 34 °C for 14 days. For *M. quadrifasciata*, due to their larger body size, the

number of newly emerged workers (N = 40) and nurse-stage workers (N = 4) in the cage

was reduced. The cages were provided daily with pollen collected from the colonies of

origin and with commercial *A. mellifera* honey diluted in water at 50%. The presence of

nurse bees in the cages is important, as they feed the newly emerged workers during their

first days of adult life.

After 14 days in the cages, the workers were submitted to a 15-hour food

restriction period prior to the arena assays. This food restriction period proved to be

essential, as, during a pilot experiment, we had noted, in the case of all three species, that

workers that were fed until the day of the assay displayed a disregard for the presence of

an intruder in the assay arena (Fig. S1). The 15-days-old workers were then individually

placed into 9 cm diameter Petri dishes lined with filter paper. After one minute of

acclimatization, an intruder, a newly emerged *Tetragonisca angustula* worker, was

introduced into each dish. The workers of the stingless bee species *T. angustula* are

smaller than those of our focal species, and not aggressive when they are still young. Each

Petri dish was filmed for 5 minutes, and the recordings were then analysed using Boris v.

8.11.3 software (Friard and Gamba, 2016). In total, we performed arena assays for 42 *S.*

postica workers, for 42 *F. varia* workers, and 19 *M. quadrifasciata* workers.

The following aggression parameters were considered: (i) Antennation (events of

touching the intruder with the antennae), (ii) frequency of attacking (percentage of tested

individuals that show any attack behaviour against the intruder during the observation

period), (iii) latency to attack (time until the first attacking event occurred), (iv) the

attacking index (total amount of time spent attacking the intruder during the five-minute

period), and (v) the pursuit index (total amount of time spent pursuing the intruder during

the five minutes).

Sampling of the worker bee life cycle stages for gene expression analysis

Workers of *S. postica*, *F. varia*, and *M. quadrifasciata* were obtained from the same

colonies as those used for the arena aggression assays. For each species, sampling was

done from three colonies. From each colony, seven workers were retrieved for each of

the three life cycle stages: nurse, guard, and forager. Sampling was done during the

Southern summer season, between November 2021 and March 2022, always between

9:00 and 10:00 am. Workers of the corresponding life cycle stages were also sampled

from a honeybee colony kept in the same apiary (*Apis mellifera*, Africanized hybrids) in

August 2022.

The sampling procedure was as follows: First, guards were collected, considering

that these were the bees that left the nest and attacked the collector's entomological sweep

net after three knocks with a hive chisel on the front board of the hive. Guards of *S. postica*

and *M. quadrifasciata* generally had an empty pollen basket (corbicula). In contrast, the

workers of *F. varia* that attacked the net all had their corbiculae (pollen baskets on the

hind legs) full of resin, as part of their colony defence strategy (Roubik, 2023). Next,

foragers were collected in front of the hives, when returning from the field with pollen-

filled corbiculae. Finally, the colonies were opened, and nurse-stage bees were collected

from the brood area. In all three species, nurse-stage bees can be identified by their

enlarged abdomen, due to their larval food-filled crop, and, their lighter body colour, as

they are still younger than the guards or foragers, and not yet fully pigmented (Nogueira-

Neto, 1997). Nurse bees are primarily engaged in the provisioning of newly constructed

brood cells with larval food. Fig. S2 shows the phenotypic characteristics of these life

cycle stages for each of the three species.

After sampling, the bees were briefly put on ice to diminish their motility and then

snap frozen in liquid nitrogen. The heads and abdomens were separately transferred into

Eppendorf vials containing 300 μL of TRIzol reagent (Invitrogen, Waltham, MA, USA)

and stored at -80 °C until RNA extraction. For the two species of smaller body size, *S.*

postica and *F. varia*, each sample was composed of a pool of heads or abdomens from

three individuals, while for the larger species *M. quadrifasciata* and *A. mellifera*, each

sample consisted of a single individual only. Pooling was necessary to obtain the RNA

yield required for gene expression analysis.

In this respect, we recognize as a limit of our study that we not separate between

age and social function. Separating these would require the setup of single-cohort

colonies, but such an experimental design is currently only available for honey bees

(Marco Antonio et al., 2008; Sullivan et al., 2000) and not for stingless bees. While in the

current study our emphasis was primarily on social task and not on age, the compiled

bionomics data available for stingless bees (Grüter, 2020) show that there is a clear age

sequence in social task performance.

RNA extraction and gene expression analysis

The head and abdominal tissues were macerated and homogenized with a pestle while

still in the TRIzol reagent before proceeding with the RNA extraction according to the

manufacturer's protocol. RNA extracts were treated with RNase-free DNase (Invitrogen)

to remove residual DNA. The quantity and quality of the RNA was checked

spectrophotometrically (NanoVue, GE Healthcare, Chicago, IL, USA). Reverse

transcription was performed using 1 μg of total RNA for cDNA synthesis, using the

Ultrascript 2.0 cDNA synthesis kit (PCRBiosystems, London, UK), following the

manufacturer's instructions.

The transcript levels of *Toll*, *Dorsal*, *Myd88* (Toll signalling pathway), and *Imd*

(Imd signalling pathway) were assessed through quantitative real-time PCR (RT-qPCR)

assays. Primer sequences for these genes were based on the respective genomic sequences

of *M. quadrifasciata* and *F. varia* (Table S1). The same primers were also tested and

successfully used to amplify the corresponding transcripts of *S. postica,* for which a

sequenced genome is not yet available.

Primer efficiencies were calculated as $E = 10^{[-1/slope]}$, using a 1:10 dilution series

of a cDNA pool containing equal amounts of RNA from all experimental groups and

species (Table S1). This pooling approach is a widely accepted method to estimate general primer performance across conditions. The primers for the endogenous control genes *Rpl32* and *Rps18* had previously been designed and validated for *F. varia and M. quadrifasciata* (Freitas et al. (2019). Additionally, as this was the first time these reference genes were used here also for *S. postica*, we confirmed their suitability by a series of additional efficiency tests using individual cDNA pools from each experimental group (head and abdomen tissues from both guards and foragers). The results show their high and consistent efficiencies (Table S1). For the normalization process, the Ct results of the two reference genes were combined by calculating their mean, and this average Ct value was used. The primer sequences for the corresponding *A. mellifera* immune system genes are based on the work by Evans et al. (2006), and those for the honeybee endogenous control gene *Rpl32* (*Rp49*) on Lourenço et al. (2008).

Amplification mixtures were set up in a total volume of 10 μL, containing 5 μl of the 2×qPCR SyGreen Mix kit (PCR Biosystems, London, UK), 1 μL of $10 \times$ diluted cDNA, primers (forward and reverse) at a final concentration of 0.125 pmol, and ultrapure water completing the final volume. The assays were performed in a StepOnePlus real-time PCR system (Applied Biosystems, Waltham, MA, USA) with the following amplification protocol: 95 °C for 10 min, followed by 40 cycles at 95 °C for 15 s and 60 °C for 30 s. After these cycles, a melting curve analysis was performed to confirm the specificity of the amplification products. Assays were done in biological septuplicates (n = 7 per colony), and each sample was analysed in technical triplicates. The relative expression of the genes of interest was calculated by the $2^{-\Delta\Delta Ct}$ formula proposed by Livak and Schmittgen (2001).

To ensure the robustness of our analyses, we decided to remove the nurse bee samples of *S. postica* from the analyses, as for this group the reference genes exhibited

considerable variation compared to the guards and foragers (Fig. S3). Despite this

adjustment, it is still possible to compare gene expression profiles between species with

reproductive (*M. quadrifasciata*) and non-reproductive (*F. varia*) workers, while the *S.*

postica data were used to specifically discuss the relationship between the life history

traits aggression and immunity.

Statistical analysis

Statistical analyses were performed in the RStudio platform (R version 4.3.0). Data

obtained from the arena assays were submitted to Shapiro-Wilk and Kolmogorov-

Smirnov tests to check the adequacy for parametric testing. Levene's test was used to

verify homoscedasticity. As these conditions were not met, we used the Kruskal-Wallis

test (dplyr package) with Dunn's post-hoc test for multiple comparisons of three or more

groups. For the analysis of the gene expression data, we used a Generalized Linear Mixed

Model (GLMM) (lme4 package), considering "body compartment" and "life cycle stage"

as fixed factors and "colony" as random. For *Apis mellifera*, in the absence of colony

replicates, we utilized the GLM test (lme4 package) to identify differences between

means. The log link function was used to approximate the gene expression data to a

Gaussian distribution, which was checked by inspecting the residuals (DHARMa

package). Graphical representations of the data were generated with GraphPad Prism v.

7 (GraphPad Softwar, Boston, MA). An adjusted *p*-value < 0.05 was considered

significant in all analyses.

Ethics statement

For the behavioural assay, the bees were kept in social groups in cages of suitable size

and with *ad libitum* access to food. The 15-hours food restriction period had no visible

negative effect on the well-being of the bees. After the arena assay, the bees were

sacrificed by instant freezing in liquid nitrogen, as they could not be returned to their

colonies without being attacked there as intruders. For the gene expression analyses the

bees were briefly cooled on ice to diminish their motility before instantaneous freezing in

liquid nitrogen. This is a standard procedure to calm down before introducing them into

any experimental procedure.

The maintenance of the native stingless bee colonies in the apiary of the Ribeirão

Preto Campus of the University of São Paulo is authorized by the *Instituto Chico Mendes*

de Conservação de Biodiversidade – Ministério de Meio Ambiente, Brazil, under the

licence 81843-1 to the senior author.

RESULTS

Species-specific individual aggressiveness of stingless bee workers

The arena assays demonstrated that under standardised conditions (number of bees of

similar age, temperature, and diet) *S. postica* showed the highest scores in the aggression

indices, while *M. quadrifasciata* exhibited the lowest scores (Fig. 2, Table S2). The scores

for *F. varia* indicated intermediate levels of aggressiveness. *S. postica* was the species

with the highest frequency of individuals that attacked the intruder in the arena. This

species also was the one with the highest frequency of fatal attacks (Fig. 2A). In

comparison, the workers of *F. varia* and *M. quadrifasciata* had much lower attacking

frequencies. The behavioural profiles of the individual bees inside the arena were also

different for the three species. Already within the first minute after introducing the

intruder specimen, the workers of *S. postica* behaved agitated, walking around quickly,

while those of *F. varia* took short flights, during which they collided with the intruder or

the arena wall. *M. quadrifasciata* workers, in turn, appeared rather lethargic, with little

movement in the arena.

All workers of *S. postica* and *F. varia*, even the non-attacking individuals,

exhibited curiosity towards the intruder, with several events of observed antennation. In

contrast, the workers of *M. quadrifasciata* did not engage with the intruder, although they

were active and moving around the arena like the other species. (Fig. 2B). The only attack

event that we saw for an *M. quadrifasciata* worker was by an individual that behaved

very agitated already from the start of the observation period, heavily beating its wings

and vibrating its body.

The species with the shortest attack latency towards the intruder, that is, it attacked

the intruder faster than the other species, was *S. postica* (Fig. 2C: Kruskall-Wallis test: H

$= 21.5$, $P < 0.01$). *F. varia* and *M. quadrifasciata* showed longer attack latencies. The

attacking index, which is the time the individual remained attacking the intruder, was

higher for *S. postica* than for *F. varia* and *M. quadrifasciata* (Fig. 2D: Kruskal-Wallis

test: $H = 23.31$, $P < 0.001$), and among the three species, *S. postica* and *F. varia* workers

presented the highest number of pursuit events (Fig. 2E: Kruskal-Wallis test: $H = 6.094$,

$P < 0.5$). During these events, *F. varia* workers often pursued the intruder with opened

wings, a behavior not observed in *S. postica* workers. However, these short flights were

not counted as pursuit events; only pursuits that occurred while walking on the plate were

considered, to maintain the same standard for all three species. No pursuit event was

observed in the case of *M. quadrifasciata*. In these events, the *F. varia* workers used to

pursue the intruder with opened wings, a behaviour that was not seen in *S. postica*

workers. No pursuit event was observed in the case of *M. quadrifasciata*.

As the arena test permitted us to quantify the individual aggression levels of the

three species under a controlled experimental setup, our results provide evidence that *S.*

postica workers exhibit higher levels of aggression compared to the other species. *M.*

quadrifasciata workers are rather docile, and *F. varia* workers show an intermediate level

of aggressiveness. While this confirms the common impression one typically has when

working with colonies of these species, the arena test results can now provide quantitative

indices for the individuals' species-specific aggressiveness levels, independent of

eventually variable colony conditions, which cannot be easily standardized in the field.

Of note, we did not include *A. mellifera* workers in the arena test because their defensive

behaviour differs significantly from that of stingless bees. While stingless bees primarily

rely on biting as a means of defence, *A. mellifera* workers typically attempt to sting, rarely

engaging in biting. This fundamental difference in behaviour would require distinct

criteria for quantifying aggression, making direct comparisons between the species

problematic. Additionally, *A. mellifera* workers do not survive sufficiently long in

confinement due to their inability to defecate without flying, leading to physical

debilitation and mortality over time. In contrast, stingless bees can defecate within the

cage and even organize waste in specific areas, allowing for easier maintenance and

ensuring their survival in standardized experimental setups (See Discussion for details).

Body compartment and social task-specific levels of immunity gene expression in the

workers of stingless bees

The three stingless bee species presented distinct profiles of immunity gene expression

levels for the head and abdomen. For *S. postica*, the most aggressive of the three species,

we found that the workers showed elevated transcript levels of *Toll* (GLMM: $\chi^2 = 109.6$,

$P < 0.001$), *Myd88* (GLMM: $\chi^2 = 25.9$, $P < 0.001$), *Dorsal* (GLMM: $\chi^2 = 34.8$, $P < 0.001$),

and *Imd* (GLMM: $\chi^2 = 30.1$, $P < 0.001$) in the head when compared to the abdomen (Fig.

3, Table S3). For the *Myd88* gene, the guards' heads actually showed higher expression

when compared with the foragers' heads (GLMM: $t = $ -3.5, $P = < 0.001$). For *M.*

quadrifasciata, the most docile of the three species, the directionality was exactly the

opposite, with significantly higher levels for *Toll* (GLMM: $\chi^2 = 133.1$, $P < 0.001$), *Myd88*

(GLMM: $\chi^2 = 4.2$, $P = 0.03$), and *Dorsal* (GLMM: $\chi^2 = 11.5$, $P < 0.001$) in the abdomen

(Fig. 4, Table S4). *F. varia* was similar to *S. postica* regarding the expression of the genes

Myd88 (GLMM: $\chi^2 = 41.6$, $P < 0.001$) and *Imd* for guards and foragers (Fig. 5, Table S5:

GLMM: $\chi^2 = 19.7$, $P < 0.001$), but the difference between the two body compartments

was much less pronounced.

With respect to the life cycle stages, the guards and foragers of *F. varia* and *M.*

quadrifasciata in general showed a tendency towards higher levels of immunity gene

expression compared to the nurse-stage workers. The workers of *M. quadrifasciata*

showed differences in gene expression levels between the younger (nurse-stage) and the

older bees (foragers and guards) in both body compartments for the genes *Toll* (GLMM:

$\chi^2 = 133.1$, $P < 0.001$) and *Myd88* (Fig. 4: GLMM: $\chi^2 = 4.2$, $P < 0.05$). With respect to the

other two genes, an abdominal up-regulation of *Dorsal* (GLMM: $\chi^2 = 11.5$, $P < 0.001$)

was associated with the guard and forager stages, while *Imd* was higher expressed only

in the heads of guards compared to nurses (GLMM: $t = 3.2$; $P < 0.05$).

For *F. varia* workers, the heads showed significantly higher *Imd* expression in

both guards (GLMM: $t = 8.25$; $P < 0.01$) and foragers (GLMM: $t = 9.75$; $P < 0.01$)

compared to nurses. *Dorsal* and *Toll* are both higher expressed in foragers compared to

nurses (*Dorsal* GLMM: $t = 2.8$; $P < 0.05$; *Toll* GLMM: $t = 4.57$; $P < 0.01$) but not in

guards, while *Myd88* is higher expressed in guards compared to nurses (GLMM: $t = 3.12$;

$P < 0.01$), but not in foragers (Fig. 5; Table S5).

Notably, we found high levels of the immune gene transcript levels in the heads

of guards and foragers, especially in the more aggressive species, *S. postica*. Yet, given

that the abdominal fat body is the primary insect tissue for mounting an immunity

response (Li et al., 2019), we asked whether these high expression levels in the bees'

heads would in fact be related to an actual immune response. To address this, we assessed

the levels of *Abaecin* transcripts, as these should represent a direct readout for a canonical

immune response (Lourenço et al. 2018).

The *Abaecin* gene encodes a major antimicrobial peptide, and its expression is

driven by both the Toll and the Imd pathway in the honeybee (Lourenço et al., 2018). For

S. postica and *F. varia* we found no differences in the *Abaecin* expression levels between

head and abdomen (GLMM: $\chi^2 = 1.8$, $P = 0.17$; and $\chi^2 = 0.5$, $P = 0.44$, respectively), while

M. quadrifasciata workers showed higher *Abaecin* expression in the abdomen than in the

head (Fig. 6: GLMM: $\chi^2 = 8.7$, $P < 0.003$). Furthermore, guards and foragers of both *S.*

postica and *M. quadrifasciata* had higher levels of *Abaecin* expression than nurses in both

body compartments, while for *F. varia* and *A. mellifera,* this was the case only for the

head.

The similarity in the expression levels for *Abaecin* with respect to social task

status across the four species indicates that, even in the absence of an immune challenge,

the immune system readiness level is higher in the older bees (guards and foragers),

possibly because they perform riskier tasks outside the colony in terms of predation threat

and pathogen exposure. Hence, while the expression patterns of the Toll and Imd pathway

genes showed pronounced differences for the head *vs.* abdominal tissues across the four

bee species, associated with their respective aggressiveness levels, the *Abaecin*

expression levels showed an age/task-related pattern that was common across the four

species. This discrepancy between the two signalling pathways and the actual immunity

readout (*Abaecin*) indicates a non-canonical function for the Toll and Imd pathways in

the heads of the more aggressive stingless bees.

Immunity gene expression in the workers of Apis mellifera

The association between aggressiveness and immune gene expression levels in

the heads of the stingless bee workers that perform the higher risk tasks of guarding and

foraging, thus raised the question as to whether this could be a conserved trait across the

highly eusocial corbiculate bees (Apini and Meliponini). To address this, we assessed the

transcript levels of the same innate immunity genes also in the honeybee, *A. mellifera*

(Fig. 7, Table S6).

We found that only *MyD88* and *Imd* are higher expressed in the heads compared

to the abdomen (GLM: $\chi^2 = 4.14$, $P = 0.04$; and $\chi^2 = 26.86$, p < 0.001, respectively). Also,

for all tested genes, we observed an increase in gene expression in at least one tissue when

comparing guards and foragers to nurse bees (Fig. 7; Table S6). Furthermore, like in *M.*

quadrifasciata, the *Abaecin* expression was higher in the abdomen than in the head in all

the life cycle stages considered in this study (GLM: $\chi^2 = 16.78$, $P < 0.001$). These findings

for the notoriously aggressive Africanized honey bee provide additional support for a

possibly non-canonical function of the Toll and Imd pathways, associated with the more

aggressive Meliponini species tested in this work.

DISCUSSION

Based on the diversity in life history parameters seen in stingless bees, exemplified by

the three focal species of this study, we intended to investigate the possible relationship

between the bees' social tasks, their aggressiveness, and their immune system status

against the background of the worker's reproductive biology (Fig. 1). The most striking

finding was the upregulation in the expression of all four of the Toll and Imd pathway

genes in the heads of the guards and foragers of *S. postica* in comparison to their

abdominal tissues (Fig. 3), as this is the more aggressive among the three species that we

tested (Fig. 2). In the heads of *F. varia,* only two of these genes (*Myd88* and *Imd)* are

higher expressed in the heads of the workers (Fig. 5)*,* while in *M. quadrifasciata,* the

opposite is seen, with two genes (*Toll* and *Dorsa*l) actually higher expressed in the

abdomen (Fig. 4)*.* In *A. mellifera*, a possibly aggressiveness-related pattern was observed

for only two of the four genes (*Myd88* and *Imd*) in the heads of guards and foragers.

However, as age and task are often correlated in social insects, we acknowledge that these

differences may also reflect age-related variation rather than task-specific regulation.

Nonetheless, we interpret these findings as indicating a possibly non-canonical

(neo)functionalization of the immune system pathways in the context of colony defence

behaviours of highly eusocial bees.

Nest defence and aggression in stingless bees

The possible reasons for the evolutionary loss of the stinging apparatus in the Meliponini

are still an open question. Yet, the idea that the small size of the bees (miniaturization)

would reduce the efficiency of their sting is not a very convincing argument, considering

the continued presence of a sting in even smaller ants. Similarly problematic is the

concomitant reduction of their venom gland, in view of the fact that genes encoding

venom proteins were found conserved in stingless bee genomes (Koludarov et al., 2023).

Yet, what is well known among beekeepers, is that stingless bee species greatly vary in

their aggressiveness as part of their strategies of nest defence.

The results obtained here via the arena assay confirmed that *S. postica* workers*,*

at the individual level, exhibit a strong aggression response towards an intruder. This is

consistent with the fact that, among the three species that we analysed, *S. postica* also has

the largest nest entrance tunnel and the largest number of guards controlling the intense

flight traffic (Couvillon et al., 2008). Furthermore, when they attack people with bites,

the sensation is quite painful compared to bites from other stingless bees (Shackleton et

al., 2015). Hence, bees of the genus *Scaptotrigona* have an efficient, aggression-based

colony defence strategy. In fact, beekeepers often emphasize the precautions that are

needed when working with colonies of this genus (Blochtein, 2008).

F. varia, the species that showed the intermediate level of aggressiveness in the

arena test (Fig. 2), also showed attention towards the intruder. However, they did not

attack with bites in most of the assay rounds. They have smaller mandibles than *S. postica*,

and their main defence strategy is to apply sticky resin onto intruders (Shackleton et al.,

2015). This strategy may not have much of an effect against vertebrate enemies, but it is

very efficient against other arthropods (Couvillon et al., 2008; Grüter, 2020). In line with

that, during the arena assays the individuals were seen to perform short flights, bouncing

against the intruder inside the Petri dish. In the field, such collision flights would allow

the *F. varia* workers to stick the resin from their corbiculae onto the potential intruder,

causing immobilization of the latter (Roubik, 2023).

In contrast to the other two species, *M. quadrifasciata* workers were completely

unaggressive in the arena assay. In fact, they were even less aggressive individually than

what is observed when they are confronted with a potential intruder at the entrance of a

colony in the field (Breed and Page, 1991; Nunes et al., 2014).

Pros and cons of the arena assay

The arena assay allowed us to quantitatively assess the aggressiveness level of individual

stingless bee workers of three focal species, independent of variation in colony contexts.

Yet, like in other behavioural tests on individual bees, certain adjustments were necessary.

This included the necessity of introducing a 15-hour food restriction period prior to testing

a bee's aggressiveness towards an intruder in the arena. Like in the proboscis extension

reflex (PER) assay (Bittermann et al., 1983; Smith and Burden, 2014) a food restriction

turned out to be necessary for the appropriate motivation of the bees to respond to the

stimulus. As previously shown in a different behavioural context, food restriction affects

the metabolism of *A. mellifera* workers, including their brain metabolism (Rittschof et al.,

2018). Furthermore, there is strong evidence that aggressive behaviours are associated

with significant metabolic changes in the brains of honeybees (Chandrasekaran et al.,

2015; Li-Byarlay et al., 2014), underlining the importance of taking into account the bees'

nutritional status when designing and standardising aggression tests.

With this in mind, there is the possibility that bees of larger size, like *M.*

quadrifasciata, may have larger energy reserves than the smaller *S. postica* and *F. varia*,

and potentially, this could make them less responsive in the arena assay. However, when

we extended the food restriction to 24 hours, this led to increased mortality. Hence, the

results of the arena assay performed under these standardised conditions confirmed the

field observations regarding the differences in the species' aggressiveness also at the level

of the individual workers.

Notably, we did not perform the arena test for the testing of honeybee workers,

because the protocol conditions that we used for the stingless bees were found not suitable

for the former. The main reason was that for the confinement in the cages, the stingless

bee workers received a diet that contained pollen. Yet, *A. mellifera* workers consuming

pollen hardly survive for more than 14 days in the cages because, different from the

stingless bees, they cannot defecate in confinement, without flying (Cole et al., 2021).

Furthermore, an arena aggression assay for *A. mellifera* workers was formerly designed

and performed with small groups of bees and not with individual bees (Nouvian et al.,

2016).

Elevated gene expression levels in the head of an aggressive stingless bee species indicate

a non-canonical function for innate immunity signalling pathways

To address the question as to whether and how the species-specific levels of

aggressiveness and the innate immune system of the stingless bee workers may be

intertwined in terms of life history strategies and trade-offs (i.e., reproduction vs.

immunity/somatic maintenance), we sampled workers according to their social task in the

colony. What we found is that in the more aggressive species, *S. postica*, the levels of

immune gene expression were upregulated in the head of the guards and foragers (Fig.

3). Since the abdominal fat body, but not the head, is the primary organ responsible for

the production of antimicrobial peptides (Li et al., 2019), this strong overexpression of

the Toll and Imd pathway genes in the bees' heads is clearly remarkable.

549   *Toll* and *Dorsal* genes (*NF-kB* in vertebrates) are present in the genomes of most

organisms, and the Toll/Toll-like receptors (TLR) signalling pathway is a very ancient

component in metazoan history (Leulier and Lemaitre, 2008). Actually, while it has been

hypothesized that the TLR immune functions involving Myd88 and NF-kB arose in the

Bilateria (Kim and Ausubel, 2005), all the core elements of Toll signalling are already

present in Cnidaria (Chapman et al., 2010). And even though the primary function of Toll

signalling is the recognition of pathogenic microorganisms, it has also acquired an

embryonic function in insects, determining the dorsoventral axis (Anderson et al., 1985).

Furthermore, in addition to its canonical function as a transmembrane receptor for

immune signalling, Toll may serve additional roles in the nervous system in both

vertebrates and invertebrates (Dresselhaus and Meffert, 2019).

560   Myd88, an adapter protein downstream to the transmembrane protein Toll (Horng

and Medzhitov, 2001), exhibited higher expression levels in the heads of *S. postica* guards

compared to foragers (Fig. 3). Interestingly, Myd88 is already known to play an important

regulatory role in mammalian neural and behavioural functions, including the regulation

of chronic stress responses (Hosoi et al., 2021). Furthermore, inhibition of this protein

promotes depression-like behaviour (Yao et al., 2023) and depression-like sleep disorders

(Choudhury et al., 2022). Since *Myd88* was found higher expressed in the aggressive

guard bees of *S. postica,* this molecular signature strengthens the hypotheses that ancestral

immunity proteins were co-opted to shape complex collective colony defence behaviours

in the highly eusocial stingless bees.

Dorsal (NF-kB), the main transcription factor of the Toll pathway, was also found

overexpressed in the heads of *S. postica* workers compared to their abdomens, whereas

in the docile *M. quadrifasciata* workers, this pattern is inverted. In mammals, NF-kB is

known to be involved in cognitive processes (Dresselhaus and Meffert, 2019). It is

constitutively activated in neurons from various regions of the central nervous system,

mainly in glutamatergic neurons (Bhakar et al., 2002; Kaltschmidt et al., 1994), and it

notably accumulates in the terminal regions of these neurons. Compelling evidence

indicates that the activation of NF-kB is induced by the neurotransmitter glutamate and

the influx of calcium ions (Ca^{2+}) (Lilienbaum and Israël, 2003; Meffert et al., 2003), and

upon activation, NF-kB undergoes a retrograde transport and becomes translocated to the

nucleus of the respective neurons (Meffert et al., 2003; Meffert and Baltimore, 2005;

Wellmann et al., 2001). In this way, it functions as a transducer of synaptic information

to the nucleus, where it assumes its role as a transcription factor (Mémet, 2006). In

addition, NF-κB is associated with memory consolidation and learning in vertebrates and

invertebrates, such as crustaceans and insects (Freudenthal and Romano, 2000; Mémet,

2006; Merlo et al., 2005). This response apparently occurs in neurons, but not in glial

cells, where NF-κB is strictly associated with the control of mammalian immune

responses and metabolism (Dresselhaus and Meffert, 2019). Based on our results obtained

for stingless bees, it is thus, reasonable to assume that in these bees, the Toll signalling

pathway, including its key transcription factor Dorsal, were co-opted to perform a

behavioural function in addition to its canonical immune system role.

With respect to *Imd* expression, the pattern was similar to *Dorsal* in *S. postica* in

terms of body compartment and social task specificity. However, aside from its immune

system function, little is known about potentially alternative roles of Imd signalling in

insects. Nonetheless, there are reports on its involvement in neurodegeneration, cell

death, and regulation of the gut microbiota (Hanson and Lemaitre, 2020; Zhai et al.,

2018). In this respect, our results may be the first ones to identify a link between the Imd

pathway and behavioural traits also in an invertebrate.

As emphasized, the expression patterns of all four immune signalling pathway

genes quantified in this study indicate an increase associated with the age-related social

task performed by the workers of the stingless bees, in both the head and the abdomen.

But how does this actually relate to the bees' immune system status? In *D.* melanogaster,

healthy and free of infection individuals showed a chronic increase with age in the

immune system activity, leading to an increased expression of antimicrobial peptides

(AMPs) in the brain, followed by marked neurodegeneration (Stuart et al., 2022).

Furthermore, a study by Kounatidis et al. (2017) showed that suppression of the Imd/NF-

κB signalling pathway in glial cells resulted in an extension of the flies' lifespan. This is

in line with data for *A. mellifera* (Lourenço et al., 2019), where nurse bees have much

lower expression levels of immunity genes and an increased lifespan.

In adult insects, AMPs are the most prominent products generated by the

activation of the Imd and Toll signalling pathways (Bulet and Stocklin, 2005), and

activation of the immune system in the brain leads to high AMP expression. This can

have collateral cytotoxic effects related to neurodegeneration, and consequently

behavioural changes, including sleep and memory dysregulation (Barajas-Azpeleta et al.,

2018; Stączek et al., 2023). The brain of *A. mellifera* was previously seen to exhibit AMP

overexpression upon infection with Deformed Wing Virus, a common bee virus,

demonstrating the ability of the brain to elicit a strong immune response (Pizzorno et al.,

2021). Studies with *D. melanogaster* in turn demonstrated that the production of AMPs

may also be regulated by pathways other than Toll and Imd (Stączek et al., 2023). For

instance, the transcription factor FoxO, which is a downstream regulator of the insulin

signalling pathway, can activate the expression of AMPs in flies undergoing starvation,

even in the absence of an infection or injury event (Becker et al., 2010). Therefore, the

elevated *Abaecin* transcript levels that we found in the head of the older stingless bees

may not be exclusively driven by the immune system pathways Toll and Imd, but also by

pathways associated with their age-specific diets and, consequently, metabolic regulation.

Nonetheless, in our previous study, where we compared head and abdominal expression

of candidate genes for social task performance in stingless bees (de Souza and Hartfelder,

2023) we did not see any expression difference in the abdominal *tor* gene expression for

F. varia and *M. quadrifasciata* nurses and foragers. In their heads, however, we saw an

inverse expression pattern between the two species, indicating that this major nutrient-

sensing pathway may be linked to the head immune gene expression profiles.

Finally, in this respect as already previously stated, we recognize as a limit of our

study that we cannot not separate between age and social function. These two factors are

strongly intertwined, with young bees preferentially performing tasks within the nest,

while older ones take on guarding and foraging task, and these tasks are strongly

associated with immune system functionality (Amdam et al., 2005; Lourenço et al.,

2019).

Immune system gene expression and the life histories of social bees

At this point it is important to state that the question that we asked differs from the one

addressed in most studies on immune system functions in social insects. Specifically, we

did not ask how does the bees' immune system respond to a pathogen challenge, but

rather, whether the aggressiveness level of the workers, as part of the colony defence

against intruders and predators, is reflected in their (unchallenged) immune system status.

We addressed this question in the workers of three species of stingless bees that vary in

important life history traits, not only with respect to nest defence, but also with the

workers' reproductive status.

Different from the honeybees, whose workers are functionally sterile in the

presence of the queen, the workers of stingless bees differ considerably in their potential

of laying eggs (Hartfelder et al., 2006; Vollet-Neto et al., 2018). While *F. varia* workers

are completely sterile, those of *M. quadrifasciata* and *S. postica* lay trophic and also

reproductive eggs in the presence of a queen, thus contributing to the colonies'

reproductive output. A key reproductive protein in female insects is vitellogenin (Vg)

(Sappington and Raikhel, 1998), but interestingly, in the functionally sterile honeybee

workers, the major function of Vg is not oogenesis. Rather, together with juvenile

hormone, it regulates the behavioural transition from intranidal (nursing) to extranidal

(foraging) behaviour (Amdam and Omholt, 2003), and it is in this context that the high

Vg levels in the haemolymph of the nurses prevent the onset of immunosenescence

(Amdam et al., 2005). Moreover, Vg itself is a powerful antioxidant protein that plays

key roles in the regulation of the lifespan of the honeybee queens and workers (Corona et

al., 2007; Havukainen et al., 2013; Seehuus et al., 2006).

In a recent study on gene expression underlying the nurse-to-forager transition in

stingless bees (de Souza and Hartfelder, 2023) we found an exactly opposite expression

pattern for the *Vg* gene in two species, showing upregulated transcript levels in the heads

of *F. varia* foragers, which are sterile, contrasting with upregulated *Vg* levels in the

abdomen of *M. quadrifasciata* nurses, which can lay eggs. Now taken together with the

current data on immunity gene expression, this suggests that, contingent on the

reproductive status of the workers, there appears to be a trade-off in the investment

between the Vg-related and the Toll/Imd-related immunity status. This trade-off not only

extends beyond the honeybees (Lourenço et al., 2012), it also appears to be species-

specifically balanced against the aggressiveness level in the stingless bees, as shown in

the current study.

Conclusion

The high transcript levels of the Toll and Imd pathway genes in the heads of the workers

of the aggressive species *S. postica*, compared to those in the docile workers of *M.*

quadrifasciata, indicate a possible non-canonical involvement of the innate immune

system in behaviours related to nest-defence strategies of social bees. The nest is an

important social factor that protects the society and its individuals from stressful

situations and, inherently, shapes their life histories (Walton et al., 2014). Given the

intricate interplay between the innate immune system and stress responses documented

in insects (Štětina et al., 2019; Hao et al., 2023), including bees (McMenamin et al., 2020),

made us, thus, raise the hypothesis of a possible association between the patterns of

immune gene expression and the levels of aggressiveness in the workers of social bees.

Of specific interest in this context is the recent proteomics study by McAfee et al.

(2024), who investigated a possibly caste-specific reproduction-immunity trade-off in the

bumble bee *Bombus impatiens*. While they found no evidence for such a trade-off in the

workers, they did so for the queens. However, the authors consider this to be due to

confounding effects resulting from the queens' complex life cycle, including a diapause

phase for overwintering, followed by a solitary nest foundation before a functional colony

gets established. They conclude that additional factors, such as worker age and

aggressiveness should be taken into consideration. It is this question what we now

addressed in the stingless bees. Not only do their queens spend their entire life in the

centre of a well-protected nest, except for a brief mating flight, it is also the species

richness of the Meliponini and their extraordinary variability in the biology of their

workers that makes them a valuable source for comparative studies on life history

strategies and functional genomics in social bees.

As next a step, extending this study to other stingless bee species that aggressively

defend their nests, or aggressively perform raids on other colonies, like *Lestrimelitta*

limao, will be important to understand whether this non-canonical pattern of immune

system gene expression represents a general trait in these highly eusocial bees.

Furthermore, the development of behavioural assays that allow the quantification of

aggressiveness levels, such as the arena assay, coupled with pharmacological

manipulation and/or RNAi functional assays, should allow mechanistic insights into the

importance of the Toll and Imd pathways in the sophisticated social behaviours of the

bees.

COMPETING INTERESTS

The authors declare no competing or financial interests.

DATA AVAILABILITY

All relevant data can be found within the article and its supplementary information. The

raw data is available from the following link: https://doi.org/10.5281/zenodo.13374927

REFERENCES

Abbot, P., 2022. Defense in social insects: Diversity, division of labor, and evolution.

Annual Review of Entomology 67, 407–436. https://doi.org/10.1146/annurev-ento-

082521-072638

Amdam, G.V., Omholt, S.W., 2003. The hive bee to forager transition in honeybee

colonies: The double repressor hypothesis. *Journal of Theoretical Biology* 223, 451–

464. https://doi.org/10.1016/S0022-5193(03)00121-8

Amdam, G. V., Aase, A.L.T.O., Seehuus, S.-C., Kim Fondrk, M., Norberg, K.,

Hartfelder, K., 2005. Social reversal of immunosenescence in honey bee workers.

Experimental Gerontololology 40, 939–947.

https://doi.org/10.1016/j.exger.2005.08.004

Anderson, K. V, Bokla, L., Nüsslein-Volhard, C., 1985. Establishment of dorsal-ventral

polarity in the Drosophila embryo: The induction of polarity by the Toll gene

product. *Cell* 42, 791–798. https://doi.org/https://doi.org/10.1016/0092-

8674(85)90275-2

Avalos, A., Fang, M., Pan, H., Ramirez Lluch, A., Lipka, A.E., Dave Zhao, S., Giray, T.,

Robinson, G.E., Zhang, G., Hudson, M.E., 2020. Genomic regions influencing

aggressive behavior in honey bees are defined by colony allele frequencies.

Proceedings of the National Academy of Sciences of the United States of America

117, 17135–17141. https://doi.org/10.5061/dryad.q573n5tg8

Barajas-Azpeleta, R., Wu, J., Gill, J., Welte, R., Seidel, C., McKinney, S., Dissel, S., Si,

K., 2018. Antimicrobial peptides modulate long-term memory. *PLoS Genetics* 14,

e1007440. https://doi.org/10.1371/journal.pgen.1007440

Barribeau, S.M., Sadd, B.M., du Plessis, L., Brown, M.J., Buechel, S.D., Cappelle, K.,

Carolan, J.C., Christiaens, O., Colgan, T.J., Erler, S., Evans, J., Helbing, S., Karaus,

E., Lattorff, H.M.G., Marxer, M., Meeus, I., Näpflin, K., Niu, J., Schmid-Hempel,

R., Smagghe, G., Waterhouse, R.M., Yu, N., Zdobnov, E.M., Schmid-Hempel, P.,

2015. A depauperate immune repertoire precedes evolution of sociality in bees.

Genome Biology 16, 83. https://doi.org/10.1186/s13059-015-0628-y

Becker, T., Loch, G., Beyer, M., Zinke, I., Aschenbrenner, A.C., Carrera, P., Inhester, T.,

Schultze, J.L., Hoch, M., 2010. FOXO-dependent regulation of innate immune

homeostasis. *Nature* 463, 369–373. https://doi.org/10.1038/nature08698

Bhakar, A.L., Tannis, L.-L., Zeindler, C., Russo, M.P., Jobin, C., Park, D.S., MacPherson,

S., Barker, P.A., 2002. Constitutive Nuclear Factor-κB activity is required for central

neuron survival. *The Journal of Neuroscience* 22, 8466.

https://doi.org/10.1523/JNEUROSCI.22-19-08466.2002

Bittermann, M.E., Menzel, R., Fietz, A., Schäfer, S., 1983. Classical conditioning of

proboscis extension in honeybees (*Apis mellifera*). *Journal of Comparative*

Psychology 97, 107-119.

Blochtein, B., 2008. *Manual de boas práticas para a criação e manejo racional de*

abelhas sem ferrão no RS: Guaraipo, manduri e tubuna. EDIPUCRS, Porto Alegre.

Boleli, I.C., Paulino-Simões, Z.L., Gentile Bitondi, M.M., 1999. Cell death in ovarioles

causes permanent sterility in *Frieseomelitta varia* worker bees. *Journal of*

Morphology 242, 271–282. https://doi.org/10.1002/(SICI)1097-4687(199912)242:3

Breed, M.D., Page, R.E., 1991. Intra- and interspecific nestmate recognition in *Melipona*

workers (Hymenoptera: Apidae). *Journal of Insect Behavior* 4, 463–469.

https://doi.org/10.1007/BF01049331

Bulet, P., Stocklin, R., 2005. Insect antimicrobial peptides: Structures, properties and

gene   regulation.   *Protein   &   Peptide   Letters*   12,   3–11.

https://doi.org/10.2174/0929866053406011

Camazine, S., 1986. Differential reproduction of the mite, *Varroa jacobso*ni

(Mesostigmata: Varroidae), on Africanized and European honey bees

(Hymenoptera: Apidae). *Annals of the Entomological Society of America* 79, 801–

803. https://doi.org/10.1093/aesa/79.5.801

Chandrasekaran, S., Ament, S.A., Eddy, J.A., Rodriguez-Zas, S.L., Schatz, B.R., Price,

N.D., Robinson, G.E., 2011. Behavior-specific changes in transcriptional modules

lead to distinct and predictable neurogenomic states. *Proceedings of the National*

Academy of Sciences of the United States of America 108, 18020–18025.

https://doi.org/10.1073/pnas.1114093108

Chandrasekaran, S., Rittschof, C.C., Djukovic, D., Gu, H., Raftery, D., Price, N.D.,

Robinson, G.E., 2015. Aggression is associated with aerobic glycolysis in the honey

bee   brain1.   *Genes,   Brain   and   Behavior*   14,   158–166.

https://doi.org/10.1111/gbb.12201

Chapman, J.A., Kirkness, E.F., Simakov, O., Hampson, S.E., Mitros, T., Weinmaier, T.,

Rattei, T., Balasubramanian, P.G., Borman, J., Busam, D., Disbennett, K.,

Pfannkoch, C., Sumin, N., Sutton, G.G., Viswanathan, L.D., Walenz, B., Goodstein,

D.M., Hellsten, U., Kawashima, T., Prochnik, S.E., Putnam, N.H., Shu, S.,

Blumberg, B., Dana, C.E., Gee, L., Kibler, D.F., Law, L., Lindgens, D., Martinez,

D.E., Peng, J., Wigge, P.A., Bertulat, B., Guder, C., Nakamura, Y., Ozbek, S.,

Watanabe, H., Khalturin, K., Hemmrich, G., Franke, A., Augustin, R., Fraune, S.,

Hayakawa, E., Hayakawa, S., Hirose, M., Hwang, J.S., Ikeo, K., Nishimiya-

Fujisawa, C., Ogura, A., Takahashi, T., Steinmetz, P.R.H., Zhang, X., Aufschnaiter,

786 R., Eder, M.-K., Gorny, A.-K., Salvenmoser, W., Heimberg, A.M., Wheeler, B.M.,

787 Peterson, K.J., Böttger, A., Tischler, P., Wolf, A., Gojobori, T., Remington, K.A.,

788 Strausberg, R.L., Venter, J.C., Technau, U., Hobmayer, B., Bosch, T.C.G., Holstein,

789 T.W., Fujisawa, T., Bode, H.R., David, C.N., Rokhsar, D.S., Steele, R.E., 2010. The

790 dynamic genome of *Hydra*. *Nature* 464, 592–596.

791 https://doi.org/10.1038/nature08830

792 Choudhury, M.E., Mikami, K., Nakanishi, Y., Matsuura, T., Utsunomiya, R., Yano, H.,

793 Kubo, M., Ando, R., Iwanami, J., Yamashita, M., Nagai, M., Tanaka, J., 2022.

794 Insomnia and depressive behavior of MyD88-deficient mice: Relationships with

795 altered microglial functions. *Journal of Neuroimmunology* 363, 577794.

796 https://doi.org/https://doi.org/10.1016/j.jneuroim.2021.577794

797 Cole, M.E., Ceja-Navarro, J.A., Mikaelyan, A., 2021. The power of poop: Defecation

798 behaviors and social hygiene in insects. *PLoS Pathogens* 17, e1009964.

799 https://doi.org/10.1371/journal.ppat.1009964

800 Corona, M., Velarde, R.A., Remolina, S., Moran-Lauter, A., Wang, Y., Hughes, K.A.,

801 Robinson, G.E., 2007. Vitellogenin, juvenile hormone, insulin signaling, and queen

802 honey bee longevity. *Proceedings of the National Academy of Sciences of the United*

803 *States of America* 104, 7128–7133. https://doi.org/10.1073/pnas.0701909104

804 Couvillon, M.J., Wenseleers, T., Imperatriz-Fonseca, V.L., Nogueira-Neto, P., Ratnieks,

805 F.L.W., 2008. Comparative study in stingless bees (Meliponini) demonstrates that

806 nest entrance size predicts traffic and defensivity. *Journal of Evolutionary Biology*

807 21, 194–201. https://doi.org/10.1111/j.1420-9101.2007.01457.x

808 Cremer, S., Pull, C.D., Fürst, M.A., 2018. Social immunity: Emergence and evolution of

809 colony-level disease protection. *Annual Review of Entomology* 63, 105–123.

810 https://doi.org/10.1146/annurev-ento-020117-043110

Danihlík, J., Aronstein, K., Petřivalský, M., 2015. Antimicrobial peptides: A key

component of honey bee innate immunity. *Journal of Apicultural Research*

https://doi.org/10.1080/00218839.2015.1109919

de Souza, L.D.R., Hartfelder, K., 2023. Reproductive potential shapes the expression of

nurse-to-forager transition genes in the workers of stingless bees (Meliponini).

Apidologie 54, 41. https://doi.org/10.1007/s13592-023-01021-z

Dierick, H.A., 2007. A method for quantifying aggression in male *Drosophila*

melanogaster.   *Nature*   *Protocols*   2,   2712–2718.

https://doi.org/10.1038/nprot.2007.404

Dresselhaus, E.C., Meffert, M.K., 2019. Cellular specificity of NF-κB function in the

nervous   system.   *Frontiers*   *in*   *Immunology*   10,   1043.

https://doi.org/10.3389/fimmu.2019.01043

Evans, J.D., Aronstein, K., Chen, Y.P., Hetru, C., Imler, J.-L., Jiang, H., Kanost, M.,

Thompson, G.J., Zou, Z., Hultmark, D., 2006. Immune pathways and defence

mechanisms in honey bees *Apis mellifera*. *Insect Molecular Biology* 15, 645–656.

https://doi.org/10.1111/j.1365-2583.2006.00682.x

Flatt, T., Amdam, G. V., Kirkwood, T.B.L., Omholt, S.W., 2013. Life-history evolution

and the polyphenic regulation of somatic maintenance and survival. *Quarterly*

Review of Biology 88, 185–218. https://doi.org/10.1086/671484

Freitas, F.C.P., Depintor, T.S., Agostini, L.T., Luna-Lucena, D., Nunes, F.M.F., Bitondi,

M.M.G., Simões, Z.L.P., Lourenço, A.P., 2019. Evaluation of reference genes for

gene expression analysis by real-time quantitative PCR (qPCR) in three stingless

bee species (Hymenoptera: Apidae: Meliponini). *Scientific Reports* 9, 17692.

https://doi.org/10.1038/s41598-019-53544-0

Freudenthal, R., Romano, A., 2000. Participation of Rel/NF-κB transcription factors in

long-term memory in the crab *Chasmagnathus*. *Brain Research* 855, 274–281.

https://doi.org/10.1016/S0006-8993(99)02358-6

Friard, O., Gamba, M., 2016. BORIS: a free, versatile open-source event-logging

software for video/audio coding and live observations. *Methods in Ecology and*

Evolution 7, 1325–1330. https://doi.org/10.1111/2041-210X.12584

Grüter, C., 2020. *Stingless Bees*, *Their Behaviour, Ecology and Evolution.* Springer

International Publishing, Cham. https://doi.org/10.1007/978-3-030-60090-7

Hanson, M.A., Lemaitre, B., 2020. New insights *on Drosophila* antimicrobial peptide

function in host defense and beyond. *Current Opinion in Immunology* 62, 22–30

https://doi.org/10.1016/j.coi.2019.11.008

Hao, Z.H., Lu, Q.Y., Zhou, Y.Y., Liang, Y.T., Gao, Y., Ma, H.Y., Xu, Y.S., Wang, H.B.,

2023. Molecular characterization of MyD88 as a potential biomarker for pesticide-

induced stress in *Bombyx mori*. *Pesticide Biochemistry and Physiology* 196, 105610.

https://doi.org/10.1016/j.pestbp.2023.105610.

Hartfelder, K., Makert, G.R., Judice, C.C., Pereira, G.A.G., Santana, W.C., Dallacqua,

R., Bitondi, M.M.G., 2006. Physiological and genetic mechanisms underlying caste

development, reproduction and division of labor in stingless bees. *Apidologie* 37,

144–163. https://doi.org/10.1051/apido:2006013

Havukainen, H., Munch, D., Baumann, A., Zhong, S., Halskau, O., Krogsgaard, M.,

Amdam, G. V, 2013. Vitellogenin recognizes cell damage through membrane

binding and shields living cells from reactive oxygen species. *Journal of Biological*

Chemistry 288, 28369–28381. https://doi.org/10.1074/jbc.M113.465021

Hoffmann, J.A., 1995. Innate immunity of insects. *Current Opinion in Immunology* 7, 4–

10. https://doi.org/https://doi.org/10.1016/0952-7915(95)80022-0

Horng, T., Medzhitov, R., 2001. *Drosophila* MyD88 is an adapter in the Toll signaling

pathway. *Proceedings of the National Academy of Sciences of the United States of*

America 98, 12654–12658. https://doi.org/10.1073/pnas.231471798

Hosoi, T., Yamawaki, Y., Kimura, H., Honda, S., Ozawa, K., 2021. Possible involvement

of MyD88 in regulating stress response in mice. *Frontiers in Neuroscience* 15,

621446. https://doi.org/10.3389/fnins.2021.621446

Imler, J.-L., Hoffmann, J.A., 2001. Toll receptors in innate immunity. *Trends in Cell*

Biology 11, 304–311. https://doi.org/10.1016/S0962-8924(01)02004-9

Janeway, C.A., Medzhitov, R., 2002. Innate immune recognition. *Annual Review of*

Immunology https://doi.org/10.1146/annurev.immunol.20.083001.084359

Kaltschmidt, C., Kaltschmidt, B., Neumann, H., Wekerle, H., Baeuerle, P.A., 1994.

Constitutive NF-kB activity in neurons. *Molecular and Cellular Biology* 14, 3981–

3992. https://doi.org/10.1128/mcb.14.6.3981-3992.1994

Kärcher, M.H., Ratnieks, F.L.W., 2009. Standing and hovering guards of the stingless

bee *Tetragonisca angustula* complement each other in entrance guarding and

intruder   recognition.   *Journal   of   Apicultural   Research*   48,   209–214.

https://doi.org/10.3896/IBRA.1.48.3.10

Kim, D.H., Ausubel, F.M., 2005. Evolutionary perspectives on innate immunity from the

study of *Caenorhabditis elegans*. *Current Opinion in Immunology* 17, 4–10.

https://doi.org/10.1016/j.coi.2004.11.007

Kingsolver, M.B., Huang, Z., Hardy, R.W., 2013. Insect antiviral innate immunity:

Pathways, effectors, and connections. *Journal of Molecular Biolog*y 425, 4921–

4936. https://doi.org/10.1016/j.jmb.2013.10.006

Koludarov, I., Velasque, M., Senoner, T., Timm, T., Greve, C., Hamadou, A. Ben, Gupta,

D.K., Lochnit, G., Heinzinger, M., Vilcinskas, A., Gloag, R., Harpur, B.A.,

Podsiadlowski, L., Rost, B., Jackson, T.N.W., Dutertre, S., Stolle, E., von Reumont,

B.M., 2023. Prevalent bee venom genes evolved before the aculeate stinger and

eusociality. *BMC Biology* 21, 229. https://doi.org/10.1186/s12915-023-01656-5

Kounatidis, I., Chtarbanova, S., Cao, Y., Hayne, M., Jayanth, D., Ganetzky, B.,

Ligoxygakis, P., 2017. NF-κB immunity in the brain determines fly lifespan in

healthy aging and age-related neurodegeneration. *Cell Reports* 19, 836–848.

https://doi.org/10.1016/j.celrep.2017.04.007

Leulier, F., Lemaitre, B., 2008. Toll-like receptors - taking an evolutionary approach.

Nature Reviews Genetics https://doi.org/10.1038/nrg2303

Li, S., Yu, X., Feng, Q., 2019. Fat body biology in the last decade. *Annual Review of*

Entomology 64, 315–333. https://doi.org/10.1146/annurev-ento-011118-112007

Li-Byarlay, H., Rittschof, C.C., Massey, J.H., Pittendrigh, B.R., Robinson, G.E., 2014.

Socially responsive effects of brain oxidative metabolism on aggression.

Proceedings of the National Academy of Sciences of the United States of America

111, 12533–12537. https://doi.org/10.1073/pnas.1412306111

Lilienbaum, A., Israël, A., 2003. From calcium to NF-κB signaling pathways in neurons.

Molecular and Cellular Biology 23, 2680–2698.

https://doi.org/10.1128/MCB.23.8.2680-2698.2003

Livak, K.J., Schmittgen, T.D., 2001. Analysis of relative gene expression data using Real-

Time quantitative PCR and the $2^{-\Delta\Delta CT}$ method. *Methods* 25, 402–408.

Lourenço, A.P., Mackert, A., Cristino, A.D.S., Simões, Z.L.P., 2008. Validation of

reference genes for gene expression studies in the honey bee, *Apis mellifera*, by

quantitative real-time RT-PCR. *Apidologie* 39, 372–385.

https://doi.org/10.1051/apido:2008015

Lourenco, A.P., Martins, J.R., Guidugli-Lazzarini, K.R., Macedo, L.M., Bitondi, M.M.,
Simoes, Z.L., 2012. Potential costs of bacterial infection on storage protein gene
expression and reproduction in queenless *Apis mellifera* worker bees on distinct
dietary regimes. *Journal of Insect Physiology* 58, 1217–1225.
https://doi.org/10.1016/j.jinsphys.2012.06.006

Lourenço, A.P., Guidugli-Lazzarini, K.R., Freitas, F.C.P., Bitondi, M.M.G., Simões,
Z.L.P., 2013. Bacterial infection activates the immune system response and
dysregulates microRNA expression in honey bees. *Insect Biochemistry and*
Molecular Biology 43, 474–482. https://doi.org/10.1016/j.ibmb.2013.03.001

Lourenço, A.P., Florecki, M.M., Simões, Z.L.P., Evans, J.D., 2018. Silencing of *Apis*
mellifera dorsal genes reveals their role in expression of the antimicrobial peptide
defensin-1. *Insect Molecular Biology* 27, 577–589.
https://doi.org/10.1111/imb.12498

Lourenço, A.P., Martins, J.R., Torres, F.A.S., Mackert, A., Aguiar, L.R., Hartfelder, K.,
Bitondi, M.M.G., Simões, Z.L.P., 2019. Immunosenescence in honey bees (*Apis*
mellifera L.) is caused by intrinsic senescence and behavioral physiology.
Experimental Gerontology 119, 174–183.
https://doi.org/10.1016/j.exger.2019.02.005

Lynch, J.A., Roth, S., 2011. The evolution of dorsal-ventral patterning mechanisms in
insects. *Genes & Development* 25, 107–118. https://doi.org/10.1101/gad.2010711

McAfee, A., Chapman, A., Bao, G., Tarpy, D.R., Foster, L.J., 2024. Investigating trade-
offs between ovary activation and immune protein expression in bumble bee
(*Bombus impatiens*) workers and queens. *Proceedings of the Royal Society B:*
Biological Sciences 291, 20232463. https://doi.org/10.1098/rspb.2023.2463

McMenamin, A.J., Daughenbaugh, K.F., Flenniken, M.L., 2020. The heat shock response

in the Western honey bee (*Apis mellifera*) is antiviral. *Viruses (Basel)* 12, 245.

htpps://doi.org/.10.3390/v12020245

Meffert, M.K., Baltimore, D., 2005. Physiological functions for brain NF-κB. *Trends in*

Neuroscience 28, 37–43. https://doi.org/10.1016/j.tins.2004.11.002

Meffert, M.K., Chang, J.M., Wiltgen, B.J., Fanselow, M.S., Baltimore, D., 2003. NF-κB

functions in synaptic signaling and behavior. *Nature Neuroscience* 6, 1072–1078.

https://doi.org/10.1038/nn1110

Melo, G.A.R., 2021. Stingless Bees (Meliponini), in: Starr C (ed) *Encyclopedia of Social*

Insects.   Springer   International   Publishing,   Cham,   pp.   883–900.

https://doi.org/10.1007/978-3-030-28102-1_117

Mémet, S., 2006. NF-κB functions in the nervous system: From development to disease.

Biochemical   Pharmacology   72,   1180–1195.

https://doi.org/10.1016/j.bcp.2006.09.003

Merlo, E., Freudenthal, R., Maldonado, H., Romano, A., 2005. Activation of the

transcription factor NF-κB by retrieval is required for long-term memory

reconsolidation. L*earning & Memory* 12, 23–29. https://doi.org/10.1101/lm.82705

Nogueira-Neto, P., 1997. *Vida e Criação de Abelhas Indígenas Sem Ferrão*. Nogueirapis,

São Paulo.

Nouvian, M., Reinhard, J., Giurfa, M., 2016. The defensive response of the honeybee

Apis   mellifera.   Journal   of   Experimental   Biology   219,   3505–3517.

https://doi.org/10.1242/jeb.143016

Nunes, T.M., Von Zuben, L.G., Costa, L., Venturieri, G.C., 2014. Defensive repertoire

of the stingless bee *Melipona flavolineata* Friese (Hymenoptera: Apidae).

Sociobiology 61, 541–546. https://doi.org/10.13102/sociobiology.v61i4.541-546

Pizzorno, M.C., Field, K., Kobokovich, A.L., Martin, P.L., Gupta, R.A., Mammone, R.,

Rovnyak, D., Capaldi, E.A., 2021. Transcriptomic responses of the honey bee brain

to infection with deformed wing virus. *Viruses (Basel)* 13, 287.

https://doi.org/10.3390/v13020287

Ratnieks, F.L., Foster, K.R., Wenseleers, T., 2006. Conflict resolution in insect societies.

Annual Review of Entomology 51, 581-608. https://doi:

10.1146/annurev.ento.51.110104.151003

Rittschof, C.C., 2017. Sequential social experiences interact to modulate aggression but

not brain gene expression in the honey bee (*Apis mellifera*). *Frontiers in Zoology*

14, 16. https://doi.org/10.1186/s12983-017-0199-8

Rittschof, C.C., Vekaria, H.J., Palmer, J.H., Sullivan, P.G., 2018. Brain mitochondrial

bioenergetics change with rapid and prolonged shifts in aggression in the honey bee,

Apis mellifera. *Journal of Experimental Biology* 221, jeb176917.

https://doi.org/10.1242/jeb.176917

Rittschof, C.C., Rubin, B.E.R., Palmer, J.H., 2019. The transcriptomic signature of low

aggression in honey bees resembles a response to infection. *BMC Genomics* 20,

1029. https://doi.org/10.1186/s12864-019-6417-3

Roubik, D.W., 2023. Stingless bee (Apidae: Apinae: Meliponini) ecology. *Annual Review*

of Entomology 68, 231–256. https://doi.org/10.1146/annurev-ento-120120-103938

Sappington, T.W., S. Raikhel, A., 1998. Molecular characteristics of insect vitellogenins

and vitellogenin receptors. *Insect Biochemistry and Molecular Biology* 28, 277–300.

https://doi.org/10.1016/S0965-1748(97)00110-0

Schwenke, R.A., Lazzaro, B.P., Wolfner, M.F., 2016. Reproduction-immunity trade-offs

in insects. *Annual Review of Entomology* 61, 239–256.

https://doi.org/10.1146/annurev-ento-010715-023924

983 Seehuus, S.C., Norberg, K., Gimsa, U., Krekling, T., Amdam, G., 2006. Reproductive

984   protein protects functionally sterile honey bee workers from oxidative stress.

985   *Proceedings of the National Academy of Sciences of the United States of America*

986   103, 962–967. https://doi.org/10.1073/pnas.0502681103

987 Sétima, T., Poupardin. R., Moos, M., Šimek, P., Šmilauer, P., Koštál, V., 2019. Larvae of

988   *Drosophila melanogaster* exhibit transcriptional activation of immune response

989   pathways and antimicrobial peptides during recovery from supercooling stress.

990   *Insect Biochemistry and Molecular Biology* 105, 60-68.

991   htpps://doi.org/10.1016/j.ibmb.2019.01.006

992 Shackleton, K., Toufailia, H. Al, Balfour, N.J., Nascimento, F.S., Alves, D.A., Ratnieks,

993   F.L.W., 2015. Appetite for self-destruction: Suicidal biting as a nest defense strategy

994   in *Trigona* stingless bees. *Behavioral Ecology and Sociobiology* 69, 273–281.

995   https://doi.org/10.1007/s00265-014-1840-6

996 Shanahan, M., Spivak, M., 2021. Resin use by stingless bees: A review. *Insects* 12, 719.

997   https://doi.org/10.3390/insects12080719

998 Shpigler, H.Y., Saul, M.C., Corona, F., Block, L., Cash Ahmed, A., Zhao, S.D.,

999   Robinson, G.E., 2017. Deep evolutionary conservation of autism-related genes.

1000   *Proceedings of the National Academy of Sciences of the United States of America*

1001   114, 9653–9658. https://doi.org/10.1073/pnas.1708127114

1002 Smith, B.H., Burden, C.M., 2014. A proboscis extension response protocol for

1003   investigating behavioral plasticity in insects: application to basic, biomedical, and

1004   agricultral research. *Journal of Visual Experiments* 91, e51057.

1005   https://doi.org/10.3791/51057

Stączek, S., Cytryńska, M., Zdybicka-Barabas, A., 2023. Unraveling the role of antimicrobial peptides in insects. *International Journal of Molecular Science* 24, 5753. https://doi.org/10.3390/ijms24065753

Stearns, S., 1992. *The Evolution of Life Histories*. Oxford University Press, Oxford.

Strand, M.R., 2008. The insect cellular immune response. *Insect Science* https://doi.org/10.1111/j.1744-7917.2008.00183.x

Stuart, B.A.R., Franitza, A.L., Lezi, E., 2022. Regulatory roles of antimicrobial peptides in the nervous system: Implications for neuronal aging. *Frontiers in Cellular Neuroscience* 16, 843790. https://doi.org/10.3389/fncel.2022.843790

The Honeybee Genome Sequencing Consortium, 2006. Insights into social insects from the genome of the honeybee *Apis mellifera*. *Nature* 443, 931–949. https://doi.org/10.1038/nature05260

Traniello, I.M., Bukhari, S.A., Dibaeinia, P., Serrano, G., Avalos, A., Ahmed, A.C., Sankey, A.L., Hernaez, M., Sinha, S., Zhao, S.D., Catchen, J., Robinson, G.E., 2023. Single-cell dissection of aggression in honeybee colonies. *Nature Ecology and Evolution* 7,1232-1244. https://doi.org/10.1038/s41559-023-02090-0

Vollet-Neto, A., Koffler, S., dos Santos, C.F., Menezes, C., Nunes, F.M.F., Hartfelder, K., Imperatriz-Fonseca, V.L., Alves, D.A., 2018. Recent advances in reproductive biology of stingless bees. *Insectes Sociaux* 65, 201–212. https://doi.org/10.1007/s00040-018-0607-x

Walton, A., Herman, J.J., Rueppell, O., 2024. Social life results in social stress protection: a novel concept to explain individual life-history patterns in social insects. *Biological Reviews* 99, 1444–1457. htpps://doi.org/10.1111/brv.13074

Wellmann, H., Kaltschmidt, B., Kaltschmidt, C., 2001. Retrograde transport of

transcription factor NF-κB in living neurons. *Journal of Biological Chemistry* 276,

11821–11829. https://doi.org/https://doi.org/10.1074/jbc.M009253200

Wilson-Rich, N., Dres, S.T., Starks, P.T., 2008. The ontogeny of immunity: development

of innate immune strength in the honey bee (*Apis mellifera*). *Journal of Insect*

Physiology 54, 1392-1399. https://doi: 10.1016/j.jinsphys.2008.07.016

Yao, X.P., Ye, J., Feng, T., Jiang, F.C., Zhou, P., Wang, F., Chen, J.G., Wu, P.F., 2023.

Adaptor protein MyD88 confers the susceptibility to stress via amplifying immune

danger signals. *Brain Behavior and Immunity* 108, 204–220.

https://doi.org/10.1016/j.bbi.2022.12.007

Zhai, Z., Huang, X., Yin, Y., 2018. Beyond immunity: The Imd pathway as a coordinator

of host defense, organismal physiology and behavior. *Developmental &*

Comparative Immunology 83, 51–59. https://doi.org/10.1016/j.dci.2017.11.008

Zhang, W., Tettamanti, G., Bassal, T., Heryanto, C., Eleftherianos, I., Mohamed, A.,

2021. Regulators and signalling in insect antimicrobial innate immunity: Functional

molecules and cellular pathways. *Cell Signaling* 83, 110003.

https://doi.org/10.1016/j.cellsig.2021.110003

1046

1047

1048

Figure 1 – The stingless bee species investigated in this study and their major life history traits. Images are credited to Dr. Cristiano Menezes.

Figure 2 – Arena test results on aggressiveness of individual workers of three stingless bee species. (A) Attack frequency and fatal attack frequencies for *S. postica*, *F. varia*, and *M. quadrifasciata*. The dark red colour represents the percentage of fatal attackers, i.e., those that killed the intruder, the red colour represents the percentage of attackers that did not kill the intruder, and the grey colour represents the percentage of bees that did not attack the intruder. (B) Number of antennation events: *S. postica* and *F. varia* exhibit have higher number of antennation events when compared to *M. quadrifasciata* (Kruskal-Wallis: $H = 20.02$, $P < 0.001$). (C) Latency to attack: *S. postica* exhibits the shortest latency to attack compared to *F. varia* and *M. quadrifasciata,* with no significant differences among the latter two species (Kruskal-Wallis: $H = 21.05$, $P < 0.001$). (D) Attacking index: *S. postica* has a higher attacking index than *F. varia* and *M. quadrifasciata* (Kruskal-Wallis: $H = 23.31$, $P < 0.001$). (E) Pursuit index: All three species showed differences in the chasing index (Kruskal-Wallis: $H = 6.09$, $P < 0.05$); however, pairwise comparisons revealed no significant differences. Orange: *S. postica* (Sp), Yellow: *F. varia* (Fv), Black: *M. quadrifasciata* (Mq). Different letters indicate statistically significant differences among the three species. The error bars represent the standard error of the mean (SEM).

Figure 3 – Relative expression levels of four immune system genes in the heads and abdomens of *Scaptotrigona postica* guards and foragers. Bar plots representing the transcript levels for (A) *Toll,* (B) *Myd88*, (C) *Dorsal,* and (D) *Imd*. Blue: Guard (G) and

Green: Forager (F). Different letters indicate statistically significantly differences among

the three live cycle stages within each body compartment. Asterisks indicate statistical

differences between the two body compartments (GLMM). Symbols (***) $P < 0.0001$,

(**) $P < 0.001$, (*) $P < 0.05$, (n.s.) non-significant. Sample size per colony, $N = 7$; per life

cycle stage within each body compartment, $N = 21$ (3 colonies). The error bars represent

the standard error of the mean (SEM).

1080

Figure 4 – Relative expression levels of four immune system genes in the heads and

abdomens of *Melipona quadrifasciata* nurses, guards, and foragers. Bar plots

representing the transcript levels for (A) *Toll,* (B) *Myd88*, (C) *Dorsal,* and (D) *Imd.*

Purple: Nurse (N), Blue: Guard (G), Green: Forager (F). Different letters indicate

statistically significantly differences among the three live cycle stages within each body

compartment. Asterisks indicate statistical differences between the two body

compartments (GLMM). Symbols (***) $P < 0.0001$, (**) $P < 0.001$, (*) $P < 0.05$, (n.s.)

non-significant. Sample size per colony, $N = 7$; per life stage within each body

compartment, $N = 21$ (3 colonies). The error bars represent the standard error of the mean

(SEM).

1091

Figure 5 – Relative expression levels of four immune system genes in the heads and

abdomens of *Frieseomelitta varia* nurses, guards, and foragers. Bar plots representing

the transcript levels for (A) *Toll,* (B) *Myd88*, (C) *Dorsal,* and (D) *Imd.* Purple: Nurse (N),

Blue: Guard (G), Green: Forager (F). Different letters indicate statistically significantly

differences among the three live cycle stages within each body compartment. Asterisks

indicate statistical differences between the two body compartments (GLMM). Symbols

(***) $P < 0.0001$, (**) $P < 0.001$, (*) $P < 0.05$, (n.s.) non-significant. Sample size per

colony, $N = 7$; per life stage within each body compartment, $N = 21$ (3 colonies). The

error bars represent the standard error of the mean (SEM).

1101

Figure 6 – *Abaecin* gene expression in the heads and abdomens of the workers of stingless

bees. Bar plots for *Abaecin* gene expression of *S. postica* (upper left) and *F. varia* (upper

right) showing no differences between the two body compartments. *M. quadrifasciata*

workers (lower left) and *A. mellifera* (lower right) have higher *Abaecin* gene expression

levels in the abdomen. Purple: Nurse (N), Blue: Guard (G), Green: Forager (F). Different

letters indicate significant differences within each body compartment. Asterisks indicate

statistical differences between the two body compartments (GLMM for *S. postica*, *F.*

varia, and *M. quadrifasciata*; GLM for *A. mellifera*). Symbols (***) $P < 0.0001$, (**) P

< 0.001, (*) $P < 0.05$, (n.s.) non-significant. Sample size per colony, $N = 7$; per life stage

within each body compartment, $N = 21$ (3 colonies for each stingless bee species); $N = 7$

for *A. mellifera* (one colony). The error bars represent the standard error of the mean

(SEM).

1114

Figure 7 – Relative expression levels of four immune system genes in the heads and

abdomens of honeybee, *Apis mellifera,* workers. Bar plots representing the transcript

levels for (A) *Toll,* (B) *Myd88,* (C) *Dorsal,* and (D) *Imd*. Purple: Nurse (N), Blue: Guard

(G), Green: Forager (F). Different letters indicate statistically significantly differences

among the three live cycle stages within each body compartment. Asterisks indicate

statistical differences between the two body compartments (GLM). Symbols (***) $P <$

0.0001, (**) $P < 0.001$, (*) $P < 0.05$, (n.s.) non-significant. Sample size per life stage

within each body compartment, $N = 7$. The error bars represent the standard error of the

mean (SEM).

 SUPPLEMENTARY MATERIAL

1125

1126 **Figure S1** – Effect of dietary restriction on the number of attacking and non-attacking

1127 workers.

1128

1129 **Figure S2 –** Images illustrating the phenotypic characteristics of the workers of the three

1130 stingless bee species used in this study

1131

1132 **Figure S3** – Ct values of the two reference genes (*Rps18* and *Rpl32*) across behavioral

1133 subcastes and tissues in the *S. postica*, *F. varia* e *M. quadrifasciata*.

1134

1135 **Table S1 -** Primers used in the RT-qPCR assays.

1136

1137 **Table S2 -** Indices of aggressiveness for individual workers of stingless bees tested in

1138 arena assays.

1139

1140 **Table S3 -** Statistical details of the GLMM test used to analyse the expression of immune

1141 pathways genes in *S. postica*.

1142

1143 **Table S4 -** Statistical details of the GLMM test used to analyse the expression of immune

1144 pathways genes in *M. quadrifasciata*.

1145

1146 **Table S5 -** Statistical details of the GLMM test used to analyse the expression of immune

1147 pathways genes in *F. varia*.

1148

1149   **Table S6 -** Statistical details of the GLM test used to analyse the expression of immune

1150   pathways genes in *A. mellifera.*

1151

ACKNOWLEDGMENTS

We thank Jairo de Souza and Luiz Roberto Aguiar for helping with the maintenance of the bee colonies in the meliponary and apiary of the Ribeirão Preto Campus of the University of São Paulo.

FUNDING

This work was supported by the São Paulo Research Foundation (FAPESP 2020/13296-9 to KH, 2020/08524-2 to CAM), the National Council for Scientific and Technological Development CNPq (302209/2022-0), and the Coordination for the Improvement of Higher Education Personnel (CAPES – Brazil Finance code 001).

STINGLESS BEES FEATURES			
Species	*S. postica*	*F. varia*	*M. quadrifasciata*
Worker size	~ 6 mm	~ 5,5 mm	~ 10 mm
Colony size	6000 to 100000 workers	800 to 1600 workers	300 to 1500 workers
Worker reproduction	Workers can lay eggs	Sterile workers	Workers can lay eggs
Colony defense	Mass attacking/Biting	Sticking resin/Biting	Blocking entrance