Adaptive decision-making by ants in response to past, imminent, and predicted adversity

AUTHOR

Yusuke Notomi[1,2], Shigeto Dobata[1], Tomoki Kazawa[2], So Maezawa[3], Shigehiro Namiki[2],

Ryohei Kanzaki[2], Stephan Shuichi Haupt[2]*

[1]Department of General Systems Studies, Graduate School of Arts and Sciences, The University

of Tokyo, 3-8-1, Komaba, Meguro-ku, Tokyo, 153-8902, Japan.

[2]Research Center for Advanced Science and Technology, The University of Tokyo, 4-6-1

Komaba, Meguro-ku, Tokyo 153-8904, Japan.

[3]Department of Applied Biological Science, Faculty of Science and Technology, Tokyo

University of Science, 2641 Yamazaki, Noda-shi, Chiba 278-8510, Japan.

*Corresponding author

CONTACT INFORMATION

Yusuke NOTOMI: notomiant@gmail.com

Stephan Shuichi HAUPT: haupt@brain.imi.i.u-tokyo.ac.jp

KEY WORDS

Visual orientation, Swimming, Locomotor pattern, Substrate condition, Internal state, Scototaxis

ABSTRACT

Many animals exhibit innate behaviours, which are often interpreted as hard-wired, reflex-like

responses, particularly in insects. Among these behaviours, beacon-aiming—an approach

towards dark areas or objects—is observed in many animals; however, its functional

significance remains unclear, and some ant species do not exhibit it. Here we show that in one

such species, *Camponotus japonicus,* the behaviour was triggered only under adverse substrate

conditions, such as during the crossing of liquid-covered surfaces, regardless of the locomotor

patterns like walking or swimming, or the mere presence of water, or when walking upside-

down. Once initiated, beacon-aiming persists even under normal substrate conditions, as

demonstrated by ants transitioning from water-covered to dry substrates suitable for

comfortable walking. This behavioural flexibility indicates that the innate behaviour is not hard-

wired but modulated by internal states. Furthermore, changes in internal states may serve

adaptive decision-making, potentially allowing ants to prepare for future adverse conditions.

The isolated ants on a water-surrounded platform gradually established an attraction to the

direction of the beacon before ultimately swimming towards it. These findings suggest that

beacon-aiming is regulated by internal states, especially anxiety-like states formed in response

to past, imminent, and predicted adverse substrate conditions in ants.

1. INTRODUCTION

Decision-making processes in animals are highly intricate, providing flexible behavioural

outputs that adapt to environmental and behavioural contexts [1,2]. While larger brains enable

advanced cognitive processing, they also bring about significant metabolic costs arising from

development and operation [3,4]. Importantly, cognitive function is not strictly dependent on

brain size [5]. Even insects with small brains and limited computational resources [6,7] are

capable of complex cognitive processing [8,9], such as path integration [10,11], tool use

(reviewed in [12]), and multi-modal cue integration [13]. Such processes are often governed by

learning, which involves plasticity in neural circuits [14]. Learning allows some forms of

inference, even in insects [15,16]. Additionally, insect behaviour can be temporarily promoted

or suppressed by physiological factors, such as the perception of threats during courtship [17]

and winner and loser effects [18].

Such transient behavioural changes are not limited to learned actions but also occur in innate

behaviours [19,20], which are typically considered relatively fixed [21]. However, the neural

mechanisms underlying the dynamic modulation of innate behaviours remain poorly

understood [22,23]. This gap partly arises from the paucity of simple experimental paradigms

to manipulate decision-making processes involved in innate behaviours, which are defined by

their stability and induction through simple stimuli [24,25].

In this study, we focus on beacon-aiming [26], a behaviour consisting of a spontaneous

orientation towards visually conspicuous dark objects. In ethological contexts, this behaviour

is classified as scototaxis and has been observed across almost all insect orders (reviewed in

[27]) and other invertebrates [28,29]. Beacon-aiming has also been reported in insects that fall

onto water surfaces [30,31]. While many species experience difficulties upon falling into water

[31,32], some are capable of coordinated locomotion across water surfaces [33,34].

In ants, beacon-aiming on land or water has been documented in various species [35,36] and

is typically interpreted as an innate response, independent of age, feeding state, or caste

[27,37,38]. However, a comparative study found that some ant species do not exhibit beacon-

aiming on plastic surfaces of experimental arenas [27].

In the present study, we departed from the hypothesis that poor substrate engagement

impairing locomotion could be a trigger for beacon-aiming. Choosing a species that did not

perform beacon-aiming in previous tests, the partially arboreal *Camponotus japonicus* [39]

capable of climbing extremely smooth vertical walls, we show that it does perform beacon-

aiming when substrate conditions, such as immersion in water, are adverse. We then introduce

experimental paradigms in which an ant has to transition from one substrate to another, i.e. from

water to dry substrate and vice-versa. This set-up allows us to probe the decision-making

dynamics in ants before, during, and after transition between substrates.

2. MATERIALS AND METHODS

(a) Animals

Foragers of the Japanese carpenter ant (*Camponotus japonicus* Mayr, 1866) were collected from

Komaba I, II campus of the University of Tokyo (35.661°N, 139.684°E; 35.662°N, 139.678°E)

and were individually isolated in 50 ml centrifuge tubes. Prior to the experiments, they were

acclimatised under laboratory conditions with light exposure for at least one hour. Only

physically intact (no visible damage to legs and antennae) ants were selected for the

experiments.

(b) General procedures

All experiments were conducted inside an opaque cylindrical arena (radius, approximately 30

cm). A vertical black rectangle (beacon) measuring 20° horizontally and 45° vertically from the

effective centre of the floor was placed on the inner wall of the arena, based on prior studies

[27,40,41]. To minimise the potential influence of external cues, the beacon was placed in one

of three possible positions, each separated by 120°, with trials evenly distributed across these

positions [27]. Each trial began with the ant placed at the centre of the floor. The behavioural

sequence was recorded using an ILME–FX3 camera (Sony) equipped with a 24–105mm F4 DG

OS HSM lens (Sigma) at 59.94 fps, until the ants reached the edge of the circular area used for

analysis (arena edge), as defined for each experiment.

(c) Experimental procedures

(i) Influence of substrate conditions, locomotor patterns and water reception on beacon-aiming

To investigate whether *C. japonicus* exhibits different orientation responses on land and water surfaces, two conditions were prepared: 'Land-walking' – walking on a dry substrate (figure 1a corresponding to [27]); and 'Water-swimming' – forced swimming in the same arena filled with water (figure 1b corresponding to [31]). To exclude the possibility that the observed behaviour is specifically associated with locomotion unique to swimming, we provided the ants with a 'Water-wading' condition where a shallow water depth allowed the ants to walk (wade) using a walking locomotor pattern (figure 1b). The behavioural sequence was recorded until the ant reached a defined arena edge (radius, 20 cm).

Furthermore, the involvement of water reception in beacon-aiming was assessed by replacing water with two kinds of liquids (figure 1c). 'Glycerol-swimming' involved swimming in a mixture of 80% (v/v) glycerol and 20% (v/v) 2-propanol, which had a dynamic viscosity of 167.1 ± 2.1 cSt (average of three measurements at 24.0°C). In the 'Tylose-swimming' used a 1.0% (w/w) tylose (H 10000 P2, SE Tylose, Wiesbaden, Germany) aqueous solution adjusted to closely match the dynamic viscosity of the glycerol-propanol mixture (166.5 ± 1.8 cSt, average of three measurements at 23.7°C, respectively). A transparent glass Petri dish (radius, 10 cm), placed at the centre of the arena, was filled with liquid. The behavioural sequence was recorded until the ant reached a defined arena edge (radius, 9 cm).

In addition, as an adverse substrate condition that does not employ liquids at all, we forced ants to walk upside-down ('Upside-down-walking') on sandpaper (#2000). Under these conditions, 32/77 ants dropped from the substrate, demonstrating it is difficult to negotiate for

them. The edge of the arena (radius, 9 cm) was defined in alignment with the dimensions of the

sandpaper (23 cm × 28 cm).

(ii) After-effects of immersion after a water-to-land transition

Building on the results of experiment (i), we examined whether the directional decision

persisted after a transition from water to a dry substrate (equivalent to aquathlon, figure 1d).

The experiment was conducted under three conditions: 'Beacon Fixed' – the beacon remained

stationary on the arena wall; 'Beacon Displaced' – the beacon was rapidly shifted by 60° upon

the transition of the ants from water to dry substrate; and 'Walking-without-transition' – ants

walked on a dry arena without a pool to account for differences in floor height and texture

compared to the standard arena used in experiment (i). The behavioural sequence was recorded

until they reached a defined arena edge (radius, 20 cm), passing through a defined pool edge

(radius, 8 cm).

(iii) Decision-making process for beacon-aiming before a land-to-water transition

In preliminary experiments, we observed that workers of *C. japonicus*, once placed on a small,

dry platform surrounded by water (referred to as an 'island'), spontaneously left the island by

swimming after a period of exploration on the platform (which would correspond to reverse

aquathlon). Leveraging this behaviour, we investigated whether the ants orient towards a

beacon prior to starting to swim, thereby examining whether the decision-making process

occurs independently of immersion.

A transparent glass cup was inverted to create an island (radius, 2.75 cm) at the centre of the

arena, surrounded by water. To prevent ants from inadvertently leaving the island due to their

initial momentum, they were first confined within a Fluon-coated cylinder on the island. The

cylinder was removed to begin the trial.

The behavioural sequence was classified into four distinct phases:

1. Initial Phase (prior to first water contact): This phase lasted until the mandibles crossed the

defined edge of the island (radius, 2.75 cm).

2. Contact Phase (before first immersion, after first water contact): Starting at the end of the

Initial Phase, this phase concluded when the mandibles crossed the defined immersion edge,

represented by a circle with a radius equal to the island's radius plus two-thirds of the ant's

body length. In this phase, ants contact water mainly with their antennae and forelegs during

exploration (defined as water contacts).

3. Immersion Phase (before initiating swimming, after first immersion): This phase spanned

from the end of the Contact Phase to the moment when the ant lost mechanical contact with

the island. In this phase, ants protrude much more over the boundary, apparently exploring

beyond for invisible gaps. This is defined as water immersion, but water contacts continue

also during this phase.

4. Swimming Phase: Beginning at the end of the Immersion Phase, this phase ended when

the ant reached the defined arena edge (radius, 20 cm).

(d) Analysis

Trajectories of the ants were obtained using DeepLabCut 2.3.8 [42], except for Upside-down-

walking, where UMATracker Release-15 [43] was instead used. The direction of the vector

connecting the centre of the arena to the position where the ant reached the arena edge was

defined as the final bearing of each trial. Similarly, the final bearings at other defined edges and

the direction of the body axis were also calculated.

Statistical analyses were conducted as previously described [27] using Python

(https://www.python.org) and the package 'circular' (version: 0.4–93) implemented in R [44].

Circular box-and-whisker plots were generated using a modified version of R code from

Buttarazzi et al (2018 [45]). The Rayleigh test was used to test for uniformity of circular

distributions. Differences in circular distributions among experimental groups were evaluated

using the Mardia–Watson–Wheeler test. For comparisons made for non-circular data groups,

the Mann–Whitney U test was used. All statistical tests were two-tailed and Bonferroni

correction applied where necessary.

Further details of MATERIALS AND METHODS are provided in the electronic

supplementary material, text S1.

3. RESULTS

(a) Experiment (i): Influence of substrate conditions, locomotor patterns and water

reception on beacon-aiming

Under 'Land-walking' conditions, ant trajectories were relatively straight (figure 2, electronic

supplementary material, figure S1); however, their orientation appeared to be in random

directions (figure 2a). This was reflected by a wide interquartile range spanning more than 90

degrees in the circular boxplot of the final bearings, indicating no clear directional preference

(Rayleigh test: $\overline{R} = 0.19$, $P = 0.140$, $N = 57$).

Under 'Water-swimming' conditions, except for occasional difficulties, ants adopted a

distinctive swimming posture with their antennae raised above the water and a locomotor

pattern characterised by extended hind legs. During swimming, most trajectories meandered

slightly but generally oriented towards the beacon or its vicinity (figure 2b). The interquartile

range of the circular boxplot of the final bearings of Water-swimming conditions overlapped

with the beacon range and was relatively narrow, indicating a preference for the centre of the

beacon (Rayleigh test: $\overline{R}$ = 0.95, $P < 0.001$, $N = 57$; mean direction: 1.3°, 95% confidence

interval: [−1.8°, 4.4°]). The final bearing distribution was significantly different from Land-

walking (Mardia–Watson–Wheeler test: $W_{(2)} = 73.01$, $P < 0.001$).

Under 'Water-wading' conditions, the distribution of final bearings (Rayleigh test: $\overline{R}$ = 0.97,

$P < 0.001$, $N = 57$; mean direction: −2.1°, 95% confidence interval: [−5.9°, 1.6°], figure 2c)

was similar to that in Water-swimming. The final bearing distribution was significantly different

from Land-walking (Mardia–Watson–Wheeler test: $W_{(2)}$ = 69.28, $P < 0.001$) but not

significantly different from Water-swimming (Mardia–Watson–Wheeler test: $W_{(2)} = 3.12$, $P =$

0.629).

When introduced into a glycerol-propanol mixture or an aqueous solution of tylose, the

smooth paddling motion of ants was hindered by the high viscosity of these liquids. Under

'Glycerol-swimming' conditions, they swam towards the beacon across the liquid surface,

although their locomotion was impaired (Glycerol-swimming, Rayleigh test: $\overline{R}$ = 0.95, $P <$

0.001, $N = 37$; mean direction: 1.5°, confidence interval: [−4.2°, 6.9°], figure 2d). Under

'Tylose-swimming' conditions, ants also swam towards the beacon (Rayleigh test: $\overline{R}$ = 0.96,

$P < 0.001$, $N = 49$; mean direction: 1.0°, confidence interval: [−3.3°, 5.2°], figure 2e). No

significant difference was observed in the distribution of final bearings between swimming

across these two liquids (Mardia–Watson–Wheeler test: Glycerol-swimming and Tylose-

swimming: $W_{(2)} = 2.66$, $P = 0.265$). These results indicate that the presence of water is not

required for beacon-aiming.

In addition, ants forced to perform upside-down walking on sandpaper also significantly

oriented towards the beacon direction ('Upside-down-walking', Rayleigh test: $\overline{R}$ = 0.72, $P <$

0.001, $N = 45$; mean direction: 10.0°, confidence interval: [−2.7°, 24.1°], figure 2f). Therefore,

liquid immersion is not necessary to induce beacon-aiming, but it is considerably more

convenient in experimental situations.

(b) Experiment (ii): Persistence of beacon-aiming decision made after water-to-land

transition

Swimming in a pool placed at the centre of the arena induced beacon-aiming, shown as the

distributions of the final bearings on the pool edge under all conditions ('Beacon Fixed',

'Beacon Displaced': Rayleigh test: $\overline{R}$ = 0.85, 0.84, both $P < 0.001$, $N = 33, 33$, mean direction:

1.0°, 0.0°, confidence interval: [−8.2°, 9.2°], [−7.9°, 7.3°], figure 3b, c, respectively).

Under Beacon Fixed conditions, beacon-aiming persisted after the transition from water to

land (Rayleigh test: $\overline{R}$ = 0.78, $P < 0.001$, $N = 33$, mean direction: −2.7°, confidence interval:

[−10.5°, 5.3°], figure 3d). No significant difference was observed in the distributions of final

bearings at the pool edge and arena edge (Mardia–Watson–Wheeler test: $W_{(2)} = 0.27$, $P = 0.875$).

Under 'Walking-without-transition' conditions on the same dry polystyrene substrate, no

orientation towards the beacon was observed (Rayleigh test: $\overline{R}$ = 0.28, $P = 0.145$, $N = 33$,

figure 3a, electronic supplementary material, figure S2). The distribution of final bearings at

arena edge in the water to land transition experiment under Beacon Fixed conditions was

significantly different from Walking-without-transition (Mardia–Watson–Wheeler test: $W_{(2)} =$

25.74, $P < 0.001$, $N = 33, 33$).

Under Beacon Displaced conditions, the beacon-aiming persisted and the directional decision

was updated even after the transition from water to land, as indicated by the mean direction

shifting towards the new beacon position, which was displaced by 60° (Rayleigh test: $\overline{R}$ = 0.61,

$P < 0.01$, $N = 33$, mean direction: 31.5°, confidence interval: [12.1°, 48.6°], figure 3e). This

shift resulted in a significant difference between the distributions of final bearings at the pool

edge and the arena edge (Watson−Wheeler test: $W_{(2)} = 30.58$, $P < 0.001$) as well as between the

distributions of final bearings at arena edge under Beacon Fixed and Beacon Displaced

conditions (Mardia–Watson–Wheeler test: $W_{(2)} = 30.69$, $P < 0.001$). Final bearings in the

Beacon Displaced condition exhibited a somewhat variable partial shift to the new beacon

position such that no significant difference was found in comparison to the Walking-without-

transition condition (Mardia–Watson–Wheeler test: $W_{(2)} = 7.72$, $P = 0.021$, $N = 33, 33$, with the

Bonferroni correction accounted for multiple comparisons among three conditions). However,

the body axis direction was significantly oriented towards the new beacon position (Rayleigh

test: $\overline{R} = 0.50$, $P < 0.001$, mean direction: 68.6°, confidence interval: [45.1°, 92.7°], figure 3f).

Once the ants reached the shore, they did not re-enter the water.

(c) Experiment (iii): Decision-making process of beacon-aiming before a land-to-water

transition

Ants, after released onto an island surrounded by water, showed distinct phases of behaviour.

During the 'Initial Phase', the ants contacted water without a directional bias (Rayleigh test: $\overline{R}$

$= 0.05$, $P = 0.917$, $N = 33$, figure 4a, b). However, a significant directional bias was evident in

the distribution of all water contacts during the 'Contact Phase' (Rayleigh test: $\overline{R} = 0.27$, $P <$

0.001, $N = 282$, mean direction: 23.2°, confidence interval: [6.3°, 39.8°], figure 4a, c, d,

electronic supplementary material, figure S3a), which is interpreted as a preference for the right

side of the beacon (10°). Similarly, the distribution of the first immersions is significantly biased

to the right edge of the beacon (Rayleigh test: $\overline{R} = 0.34$, $P = 0.033$, $N = 33$, mean direction:

6.0°, confidence interval: [−33.5°, 53.0°], figure 4c), and the distribution of all water

immersions is also significantly biased to the right edge of the beacon (Rayleigh test: $\overline{R} = 0.66$,

$P < 0.001$, $N = 33$, mean direction: 13.2°, confidence interval: [10.5°, 15.7°], figure 4a, e, f).

After several water immersions (electronic supplementary material, figure S3b), the ants

exhibited 'pre-swimming behaviour'. Some behaviours classified as immersions were

identified as pre-swimming behaviour. This behaviour, characterised by paddling movements

while maintaining contact with the shore using at least one hind leg, displayed intermediate

characteristics between walking and swimming, complicating quantitative analysis. Most ants

began swimming after clear pre-swimming behaviour, showing a directional bias towards the

beacon (Rayleigh test: $\overline{R}$ = 0.94, $P < 0.001$, $N = 33$, mean direction: 7.0°, confidence interval:

[6.1°, 14.8°], figure 4a, e). During the 'Swimming Phase', ants finally decided orienting

towards the centre of the beacon (Rayleigh test: $\overline{R}$ = 0.99, $P < 0.001$, $N = 33$, mean direction:

1.4°, confidence interval: [-1.0°/3.9°], figure 4a, g).

These results suggest that while the ants showed no interest in the beacon during the Initial

Phase, they exhibited a preference for the sides or edges of the beacon during the Contact Phase,

eventually aligning themselves with the centre of the beacon after swimming.

4. Discussion

Animals act autonomously in diverse and complex natural environments. The elements defining

their behaviour have been classified into 1) reflexes, fixed action patterns, and innate

behaviours which can be seen as characteristics of an automatic machine, the extensions of such

a machine, or a combination of both, and 2) learning, decision-making, and related processes

which are key elements enabling flexible adaptation to diverse natural environments. The

primary focus of the research aimed at understanding flexibility and adaptability has been on

the latter.

While some ant species are known to orient towards a visual landmark on land [26,35,37],

recent findings have revealed that *C. japonicus* does not exhibit such orientation under

terrestrial conditions [27], except in cases where the landmark is associated with appetitive

learning [46]. Our research showed that the attraction to the beacon in this species when

crossing water surfaces was triggered by immersion rather than by reception of water itself or

the use of swimming-specific locomotor patterns (experiment (i)). Changes in orientation

decisions depending on substrate conditions are known from a few intertidal species in specific

ecological contexts [47,48]. When forced to walk upside-down under danger of dropping, the

ants performed beacon-aiming too. Insects are known to internally monitor their walking

movements, detecting reductions in walking speed and/or abnormal leg rotation rates [49].

Therefore, they are sensitive to the state of contact with the substrate. This is especially true for

arboreal arthropods, as dropping from arboreal environments can induce isolation from familiar

environments and/or falling onto water surfaces. Unfamiliar environments pose difficulties for

navigation, which is a significant problem, particularly for central place foragers [50,51].

Furthermore, movements across water surfaces pose risks of drowning or predation, especially

for small animals [52–54]. Thus, adverse substrate conditions or even adverse conditions in

general are candidates for inducing beacon-aiming. This may apply to other arboreal arthropods

or even a multitude of terrestrial animals.

Context-dependent decision-making has been studied in diverse animal species and in

various contexts [55,56]. Although innate behaviours are generally considered to be hard-wired

[37,57], a certain degree of context-dependence has been reported [19,58]. However, it remains

to be explained whether these context-dependent responses are similar to program behaviours

triggered directly by a specific sensory input [59] or are instead regulated by transient changes

in internal state [60,61].

Ants having exhibited beacon-aiming in the pool (experiment (ii)) continued to orient

towards the beacon even after beginning to walk. This result contrasts with the absence of

beacon-aiming without prior swimming experience. Most ants re-oriented themselves towards

the displaced beacon, suggesting that their behaviour was not merely maintaining directional

consistency or a direct sensory behaviour; rather, it should be regarded as a continuation of an

internal state [23] that induces beacon-aiming. Internal states provide specialized consistency

to behavioural tendencies in animals across similar behavioural contexts, such as hunger [62],

thirst [63], social isolation [64], and focusing on courtship [17]. The temporal persistence of

internal states has been observed in other arthropods [65,66].

Ants having been introduced onto the island (experiment (iii)) made initial contacts with the

surrounding water in various directions and then began to access boundary areas in different

directions. Thorough scanning in unfamiliar visual environments in ants aligns with previous

observations [67]. It is noteworthy that the orientation of the ants towards the periphery of the

beacon was induced prior to the experience of immersion. Furthermore, even after repeated

water contacts and subsequent beacon-aiming, the ants often continued to explore other

directions thoroughly (electronic supplementary material, figure S3b). This observation

suggests that beacon-aiming in ants is not a binary response but rather a behaviour shaped

gradually and most likely caused by the formation of an anxiety-like state [68]. In summary,

the results of our study demonstrate that beacon-aiming in the ant *C. japonicus* elicited by

adverse substrate conditions can be interpreted as an anxiety-like behaviour.

Anxiety-like behaviour has been studied primarily in rodents [69] and fish [70], as well as in

some arthropods [71,72], including insects [73], yet detailed research remains scarce for non-

model species. The dynamic changes in orientation decisions observed in *C. japonicus* on an

island, which might be viewed as a reverse analogue of the Morris water maze, illustrate the

process of anxiety-like state formation. To the best of our knowledge, spontaneous escape

behaviours from a platform have been studied only in a few invertebrates [74,75].

A commonly used experimental paradigm in studies of anxiety is the light-dark test, which

induces a preference for darkness (scototaxis) in animals. The simplest explanation for the

ecological significance of this preference is that the bright, open areas are innately aversive to many animals, whereas darker spaces are relatively predictable as more secure under predation risks [76,77], making them valuable to approach or remain in. Scototaxis has been observed in rodents [78,79], fishes [80], insects [81,82], crustaceans [83,84], arachnids [85], echinoderms [86], and molluscs [29,87]. Conversely, this innate preference can be altered depending on habitat [88], developmental stage [89], and time of day [88].

The relationship between beacon-aiming and dark preference in general remains to be explained. For instance, tendencies such as preference for an area which includes a number of dark spots over overall darkness [22], back-and-forth movement between black stripes in Buridan's paradigm [90], and attraction to a bright beacon against a dark background during swimming [40] have been observed. If these behaviours are included in beacon-aiming, the relationship between scototaxis as an anxiety-like behaviour and beacon-aiming would have to be reconsidered.

Is beacon-aiming simply a result of an internal state leading to this behaviour or is there a more sophisticated cognitive process involved to decide to head for the beacon? This is something we cannot decide at this point. Interestingly, beacon-aiming on the island (experiment (iii)) is adapted to handle the adversity regardless of the actual mechanism that can be the result of a change in internal state [72] or cognitive processing [91]. Examples of responses to future adversity include ants avoiding similar situations by generalizing based on their experiences even after a single emergency encounter [92], and grasshoppers that, under chronic predation risk, learn to escape faster and farther [93].

Various substances are known to play a major role in forming and maintaining internal states [18,65], but knowledge about the mechanisms regulating innate behaviour remains limited [19,23]. Brain areas known to be involved in beacon-aiming are the central complex, which is necessary for turning during visual following in the context of beacon-aiming [94], and the

360   mushroom bodies, responsible for allowing navigation relative to a beacon by learning rather

361   than aiming for it [95]. The activation of neurons expressing dNPF (*Drosophila* neuropeptide

362   F) [96] or octopaminergic neurons [97] can induce beacon-aiming towards small objects that

363   *Drosophila* otherwise avoid [41]. Whether such mechanisms are related to state changes that

364   result in the behaviours we observe around transitions between substrates remains to be seen.

5. Conclusion

A simple and supposedly hard-wired orientation response, beacon-aiming, was shown to be

induced by adverse substrate conditions in an ant species. Using paradigms with transitions

between different substrates enabled us to observe dynamical aspects of the decisions for

orientation. As both after-effects and pre-emptive effects could be shown, we conclude that

substrate conditions adverse to locomotion are a factor altering the internal state of the ants,

leading to an anxiety-like state which promotes the orientation towards beacons. Similarities to

anxiety-like behaviours in other animals open up the interesting question whether similar

behavioural strategies are used by a wide range of animal species.

Reference

1. Dolan RJ. 2002 Emotion, cognition, and behavior. *Science* **298**, 1191–1194.
(doi:10.1126/science.1076358)

2. Webb B. 2012 Cognition in insects. *Philos Trans R Soc Lond B Biol Sci* **367**, 2715–2722.
(doi:10.1098/rstb.2012.0218)

3. Aiello LC, Wheeler P. 1995 The expensive-tissue hypothesis: the brain and the digestive
system in human and primate evolution. *Curr. Anthropol.* **36**, 199–221.

4. van der Woude E, Groothuis J, Smid HM. 2019 No gains for bigger brains: Functional and
neuroanatomical consequences of relative brain size in a parasitic wasp. *J. Exp. Biol.* **32**, 694–
705. (doi:10.1111/jeb.13450)

5. Chittka L, Niven J. 2009 Are bigger brains better? *Curr. Biol.* **19**, R995–R1008.
(doi:10.1016/j.cub.2009.08.023)

6. Abbott LF *et al.* 2020 The mind of a mouse. *Cell* **182**, 1372–1376.
(doi:10.1016/j.cell.2020.08.010)

7. Raji JI, Potter CJ. 2021 The number of neurons in *Drosophila* and mosquito brains. *PLOS*
ONE **16**, e0250381. (doi:10.1371/journal.pone.0250381)

8. Chittka L, Skorupski P. 2011 Information processing in miniature brains. *Proc. R. Soc. B.* **278**,
885–888. (doi:10.1098/rspb.2010.2699)

9. Hulse BK *et al.* 2021 A connectome of the *Drosophila* central complex reveals network motifs
suitable for flexible navigation and context-dependent action selection. *Elife* **10**, e66039.
(doi:10.7554/eLife.66039)

10. Müller M, Wehner R. 1988 Path integration in desert ants, *Cataglyphis fortis*. *Proc. Natl.*
Acad. Sci. U. S. A. **85**, 5287–5290. (doi:10.1073/pnas.85.14.5287)

11. Wittlinger M, Wehner R, Wolf H. 2006 The ant odometer: stepping on stilts and stumps.
Science **312**, 1965–1967. (doi:10.1126/science.1126912)

12. Tomer J. Czaczkes. 2022 Advanced cognition in ants. *Myrmecol. News* **32**, 51–64.
(doi:10.25849/MYRMECOL.NEWS_032:051)

13. Buehlmann C, Mangan M, Graham P. 2020 Multimodal interactions in insect navigation.
Anim. Cogn. **23**, 1129–1141. (doi:10.1007/s10071-020-01383-2)

14. Rössler W. 2019 Neuroplasticity in desert ants (Hymenoptera: Formicidae) – importance for
the ontogeny of navigation. *Myrmecol. News* **29**, 61–69.
(doi:10.25849/myrmecol.news_029:001)

15. Martin-Ordas G. 2024 Inferential reasoning abilities in wild-caught bumblebees. *Biol. Lett.*
20, 20230561. (doi:10.1098/rsbl.2023.0561)

16. Tibbetts EA, Agudelo J, Pandit S, Riojas J. 2019 Transitive inference in *Polistes* paper wasps.
Biol. Lett. **15**, 20190015. (doi:10.1098/rsbl.2019.0015)

17. Cazalé-Debat L *et al.* 2024 Mating proximity blinds threat perception. *Nature* **634**, 635–643.
(doi:10.1038/s41586-024-07890-3)

18. Stevenson PA, Rillich J. 2016 Controlling the decision to fight or flee: the roles of biogenic
amines and nitric oxide in the cricket † . *Curr. Zool.* **62**, 265–275. (doi:10.1093/cz/zow028)

19. Gorostiza EA, Colomb J, Brembs B. 2016 A decision underlies phototaxis in an insect. *Open*
Biol. **6**, 160229. (doi:10.1098/rsob.160229)

20. McEwen RS. 1918 The reactions to light and to gravity in Drosophila and its mutants. *Journal*
of Experimental Zoology **25**, 49–106. (doi:10.1002/jez.1400250103)

21. Jander R. 1963 Insect orientation. *Annu. Rev. Entomol.* **8**, 95–114.
(doi:10.1146/annurev.en.08.010163.000523)

22. Bengochea M, Sitt JD, Izard V, Preat T, Cohen L, Hassan BA. 2023 Numerical discrimination
in *Drosophila melanogaster*. *Cell Rep.* **42**, 112772. (doi:10.1016/j.celrep.2023.112772)

23. Grunwald Kadow IC. 2019 State-dependent plasticity of innate behavior in fruit flies. *Curr.*
Opin. Neurobiol. **54**, 60–65. (doi:10.1016/j.conb.2018.08.014)

24. Bourin M, Hascoët M. 2003 The mouse light/dark box test. *Eur. J. Pharmacol.* **463**, 55–65.
(doi:10.1016/S0014-2999(03)01274-3)

25. Crawley JN, Marangos PJ, Paul SM, Skolnick P, Goodwin FK. 1981 Interaction between
purine and benzodiazepine: inosine reverses diazepam-induced stimulation of mouse
exploratory behavior. *Science* **211**, 725–727. (doi:10.1126/science.6256859)

26. Graham P, Fauria K, Collett TS. 2003 The influence of beacon-aiming on the routes of wood
ants. *J. Exp. Biol.* **206**, 535–541. (doi:10.1242/jeb.00115)

27. Notomi Y, Kazawa T, Maezawa S, Kanzaki R, Haupt SS. 2022 Use of visual information by ant
species occurring in similar urban anthropogenic environments. *Zool. Sci.* **39**, 529–544.
(doi:10.2108/zs220035)

28. Chiussi R. 2003 Orientation and shape discrimination in juveniles and adults of the mangrove
crab *Aratus pisonii* (H. Milne Edwards, 1837): effect of predator and chemical cues. *Mar.*
Freshw. Behav. Physiol. **36**, 41–50. (doi:10.1080/10236240290031548)

29. Zanforlin M. 1976 Observations on the visual perception of the snail *Euparipha pisana*
(Müller). *Boll. Zool.* **43**, 303–315. (doi:10.1080/11250007609430147)

30. Beugnon G. 1986 Learned orientation in landward swimming in the cricket *Pteronemobius*
Lineolatus. Behav. Process **12**, 215–226. (doi:10.1016/0376-6357(86)90037-9)

31. Yanoviak SP, Frederick DN. 2014 Water surface locomotion in tropical canopy ants. *J. Exp.*
Biol. **217**, 2163–2170. (doi:10.1242/jeb.101600)

32. Miyamoto S. 1955 On a special mode of locomotion utilizing surface tension at the water-edge
in some semi-aquatic insects. *Kontyu* **23**.

33. Bush JWM, Hu DL. 2006 Walking on water: Biolocomotion at the Interface. *Annu. Rev. Fluid*
Mech. **38**, 339–369. (doi:10.1146/annurev.fluid.38.050304.092157)

34. DuBois MB, Jander R. 1985 Leg coordination and swimming in an ant, *Camponotus*
americanus. Physiol. Entomol. **10**, 267–270. (doi:10.1111/j.1365-3032.1985.tb00046.x)

35. Macquart D, Garnier L, Combe M, Beugnon G. 2006 Ant navigation en route to the goal:
signature routes facilitate way-finding of *Gigantiops destructor. J. Comp. Physiol. A* **192**, 221–
234. (doi:10.1007/s00359-005-0064-7)

36. Schultheiss P, Guénard B. 2021 Kinematic study of six mangrove ant species (Hymenoptera:
Formicidae) reveals different swimming styles and abilities. *Myrmecol. News* **31**, 217–224.
(doi:10.25849/MYRMECOL.NEWS_031:217)

37. Buehlmann C, Graham P. 2022 Innate visual attraction in wood ants is a hardwired behavior
seen across different motivational and ecological contexts. *Insect. Soc.* **69**, 271–277.
(doi:10.1007/s00040-022-00867-3)

38. Jander R. 1957 Die optische Richtungsorientierung der Roten Waldameise (*Formica rufa* L.).
Z. Vergl. Physiol. **40**, 162–238. (doi:10.1007/BF00297947)

39. Fukushi T, Sato J, Kusaka Y, Konno R, Sasaki M, Takahashi H. 1999 Visual orientation in
outbound and homing travels in the ant, *Camponotus japonicus. Bull. Miyagi Univ. Educ.* **34**,
61–71.

40. Gora EM, Gripshover N, Yanoviak SP. 2016 Orientation at the water surface by the carpenter
ant *Camponotus pennsylvanicus* (DE GEER, 1773) (Hymenoptera: Formicidae). *Myrmecol.*
News **23**, 33–39. (doi:https://doi.org/10.25849/myrmecol.news_023:033)

41. Maimon G, Straw AD, Dickinson MH. 2008 A simple vision-based algorithm for decision
making in flying *Drosophila. Curr. Biol.* **18**, 464–470. (doi:10.1016/j.cub.2008.02.054)

42. Mathis A, Mamidanna P, Cury KM, Abe T, Murthy VN, Mathis MW, Bethge M. 2018
DeepLabCut: markerless pose estimation of user-defined body parts with deep learning. *Nat.*
Neurosci. **21**, 1281–1289. (doi:10.1038/s41593-018-0209-y)

43. Yamanaka O, Takeuchi R. 2018 UMATracker: an intuitive image-based tracking platform. *J.*
Exp. Biol. **221**, jeb182469. (doi:10.1242/jeb.182469)

44. R Core Team. 2021 R: A language and environment for statistical computing. R Foundation
for Statistical Computing, Vienna, Austria. See https://www.r-project.org/ (accessed on 16
July 2022).

45. Buttarazzi D, Pandolfo G, Porzio GC. 2018 A boxplot for circular data. *Biometrics* **74**, 1492–
1501. (doi:10.1111/biom.12889)

46. Sakiyama T, Suda K. 2023 Movement during the acquisition of a visual landmark may be
necessary for rapid learning in ants. *J. Comp. Physiol. A* (doi:10.1007/s00359-023-01652-8)

47. Avellanal MH, Jaramillo E, Naylor E, Kennedy F. 2000 Orientation of *Phalerisida maculata*
Kulzer (Coleoptera, Tenebrionidae) in sandy beaches of the Chilean coast: Orientation of
Phalerisida maculata in sandy beaches. *J. Exp. Mar. Biol. Ecol.* **247**, 153–167.
(doi:10.1016/S0022-0981(99)00185-9)

48. Ugolini A, Scapini F, Pardi L. 1986 Interaction between solar orientation and landscape
visibility in *Talitrus saltator* (Crustacea: Amphipoda). *Mar. Biol.* **90**, 449–460.
(doi:10.1007/BF00428568)

49. Fujiwara T, Cruz TL, Bohnslav JP, Chiappe ME. 2017 A faithful internal representation of
walking movements in the *Drosophila* visual system. *Nat. Neurosci.* **20**, 72–81.
(doi:10.1038/nn.4435)

50. Sakiyama T. 2019 Emergence of a complex movement pattern in an unfamiliar food place by
foraging ants. *J. Comp. Physiol. A* **205**, 61–66. (doi:10.1007/s00359-018-1303-z)

51. Schwarz S, Wystrach A, Cheng K. 2017 Ants' navigation in an unfamiliar environment is
influenced by their experience of a familiar route. *Sci. Rep.* **7**, 14161. (doi:10.1038/s41598-
017-14036-1)

52. Adis J. 1982 Eco-Entomological observations from the Amazon: III. How do leafcutting ants of
inundation forests survive flooding? *Acta Amaz.* **12**, 839–840. (doi:10.1590/1809-
43921982124839)

53. Small GE, Torres PJ, Schweizer LM, Duff JH, Pringle CM. 2013 Importance of terrestrial
arthropods as subsidies in lowland Neotropical rain forest stream ecosystems. *Biotropica* **45**,
80–87. (doi:10.1111/j.1744-7429.2012.00896.x)

54. Yanoviak SP, Munk Y, Dudley R. 2011 Evolution and ecology of directed aerial descent in
arboreal ants. *Integr. Comp. Biol.* **51**, 944–956. (doi:10.1093/icb/icr006)

55. Barbasch TA, Abuwa VI, Carswell B, Bell AM. 2024 Managing the tradeoff between
reproduction and survival requires flexibility in behaviour and gene regulation in three-spined
stickleback. *Proc. R. Soc. B.* **291**, 20242296. (doi:10.1098/rspb.2024.2296)

56. Wright TF, Eberhard JR, Hobson EA, Avery ML, Russello MA. 2010 Behavioral flexibility and
species invasions: the adaptive flexibility hypothesis. *Ethol. Ecol. Evol.* **22**, 393–404.
(doi:10.1080/03949370.2010.505580)

57. Perry CJ, Chittka L. 2019 How foresight might support the behavioral flexibility of arthropods.
Curr. Opin. Neurobiol. **54**, 171–177. (doi:10.1016/j.conb.2018.10.014)

58. Cheng KY, Frye MA. 2021 Odour boosts visual object approach in flies. *Biol. Lett.* **17**,
20200770. (doi:10.1098/rsbl.2020.0770)

59. Kanzaki R, Sugi N, Shibuya T. 1992 Selfgenerated zigzag turning of *Bombyx mori* males
during pheromone-mediated upwind walking. *Zoological Science* **9**, 515–527.
(doi:10.34425/zs000983)

60. Deakin A, Mendl M, Browne WJ, Paul ES, Hodge JJL. 2018 State-dependent judgement bias
in *Drosophila*: evidence for evolutionarily primitive affective processes. *Biol. Lett.* **14**,
20170779. (doi:10.1098/rsbl.2017.0779)

61. Iwasaki M, Delago A, Nishino H, Aonuma H. 2006 Effects of previous experience on the
agonistic behaviour of male crickets, *Gryllus bimaculatus*. *Zool. Sci.* **23**, 863–872.
(doi:10.2108/zsj.23.863)

62. Navas CA, Agudelo-Cantero GA, Loeschcke V. 2022 Thermal boldness: Volunteer exploration
of extreme temperatures in fruit flies. *J. Insect Physiol.* **136**, 104330.
(doi:10.1016/j.jinsphys.2021.104330)

63. Lin S, Owald D, Chandra V, Talbot C, Huetteroth W, Waddell S. 2014 Neural correlates of
water reward in thirsty *Drosophila. Nat. Neurosci.* **17**, 1536–1542. (doi:10.1038/nn.3827)

64. Hoffmann AA. 1990 The influence of age and experience with conspecifics on territorial
behavior in *Drosophila melanogaster. J. Insect Behav.* **3**, 1–12. (doi:10.1007/BF01049191)

65. Iwasaki M, Nishino H, Delago A, Aonuma H. 2007 Effects of NO/cGMP signaling on
behavioral changes in subordinate male crickets, *Gryllus bimaculatus. Zool. Sci.* **24**, 860–868.
(doi:10.2108/zsj.24.860)

66. Stevenson PA, Rillich J. 2013 Isolation associated aggression – a consequence of recovery from
defeat in a territorial animal. *PLOS ONE* **8**, e74965. (doi:10.1371/journal.pone.0074965)

67. Wystrach A, Philippides A, Aurejac A, Cheng K, Graham P. 2014 Visual scanning behaviours
and their role in the navigation of the Australian desert ant *Melophorus bagoti. J. Comp.*
Physiol. A **200**, 615–626. (doi:10.1007/s00359-014-0900-8)

68. Guitton MJ, Dudai Y. 2004 Anxiety-like state associates with taste to produce conditioned
taste aversion. *Biol. Psychiatry* **56**, 901–904. (doi:10.1016/j.biopsych.2004.08.024)

69. Belzung C, Griebel G. 2001 Measuring normal and pathological anxiety-like behaviour in
mice: a review. *Behavioural Brain Research* **125**, 141–149. (doi:10.1016/S0166-
4328(01)00291-1)

70. Stewart A, Gaikwad S, Kyzar E, Green J, Roth A, Kalueff AV. 2012 Modeling anxiety using
adult zebrafish: A conceptual review. *Neuropharmacology* **62**, 135–143.
(doi:10.1016/j.neuropharm.2011.07.037)

71. Fossat P, Bacqué-Cazenave J, De Deurwaerdère P, Delbecque J-P, Cattaert D. 2014 Anxiety-
like behavior in crayfish is controlled by serotonin. *Science* **344**, 1293–1297.
(doi:10.1126/science.1248811)

72. Perrot-Minnot M-J, Banchetry L, Cézilly F. 2017 Anxiety-like behaviour increases safety from
fish predation in an amphipod crustacea. *R. Soc. Open Sci.* **4**, 171558.
(doi:10.1098/rsos.171558)

73. Mohammad F, Aryal S, Ho J, Stewart JC, Norman NA, Tan TL, Eisaka A, Claridge-Chang A.
2016 Ancient Anxiety Pathways Influence *Drosophila* Defense Behaviors. *Curr. Biol.* **26**, 981–
986. (doi:10.1016/j.cub.2016.02.031)

74. Yamaguchi T. 2017 Behavior of *Formica japonica* (Hymenoptera: Formicidae) dropped on an
artificial small "island". *ARI, J. Myrmecol. Soc. Japan* **38**, 31–44.

75. Yamaguchi T. 2018 Effects of photoenvironments on the behavior of *Formica japonica*
(Hymenoptera: Formicidae) dropped on an artificial small "island". *ARI, J. Myrmecol. Soc.*
Japan **39**, 37–59.

76. Culp JM, Glozier NE, Scrimgeour GJ. 1991 Reduction of predation risk under the cover of
darkness: Avoidance responses of mayfly larvae to a benthic fish. *Oecologia* **86**, 163–169.
(doi:10.1007/BF00317527)

77. Kotler BP, Brown JS, Hasson O. 1991 Factors affecting gerbil foraging behavior and rates of
owl predation. *Ecology* **72**, 2249–2260. (doi:10.2307/1941575)

78. Hascoët M, Bourin M, Nic Dhonnchadha BrídÁ. 2001 The mouse ligth-dark paradigm: A
review. *Prog. Neuropsychopharmacol. Biol. Psychiatry.* **25**, 141–166. (doi:10.1016/S0278-
5846(00)00151-2)

79. Crawley J, Goodwin FK. 1980 Preliminary report of a simple animal behavior model for the
anxiolytic effects of benzodiazepines. *Pharmacol. Biochem. Behav.* **13**, 167–170.
(doi:10.1016/0091-3057(80)90067-2)

80. Maximino C, Marques de Brito T, Dias CAG de M, Gouveia A, Morato S. 2010 Scototaxis as
anxiety-like behavior in fish. *Nat Protoc* **5**, 209–216. (doi:10.1038/nprot.2009.225)

81. Colombini I, Chelazzi L, Scapini F. 1994 Solar and landscape cues as orientation mechanisms
in the beach-dwelling beetle *Eurynebria complanata* (Coleoptera, Carabidae). *Mar. Biol.* **118**,
425–432. (doi:10.1007/BF00350299)

82. Kral K. 2019 Visually guided search behavior during walking in insects with different habitat
utilization strategies. *J. Insect Behav.* **32**, 290–305. (doi:10.1007/s10905-019-09735-8)

83. Cohen JH, Putts MR. 2013 Polarotaxis and scototaxis in the supratidal amphipod
Platorchestia platensis. J. Comp. Physiol. A **199**, 669–680. (doi:10.1007/s00359-013-0825-7)

84. Huang H-D, Rittschof D, Jeng M-S. 2005 Visual orientation of the symbiotic snapping shrimp
Synalpheus demani. J. Exp. Mar. Biol. Ecol. **326**, 56–66. (doi:10.1016/j.jembe.2005.05.008)

85. Kopp K, Gothe R. 1995 *Hyalomma truncatum* (Acari; Ixodidae): investigations into the
scototaxis of unfed adult ticks. *Exp. Appl. Acarol.* **19**, 53–64. (doi:10.1007/BF00051937)

86. Blevins E, Johnsen S. 2004 Spatial vision in the echinoid genus *Echinometra. J. Exp. Biol.* **207**,
4249–4253. (doi:10.1242/jeb.01286)

87. Bisazza A, Gatto E. 2021 Continuous versus discrete quantity discrimination in dune snail
(Mollusca: Gastropoda) seeking thermal refuges. *Sci. Rep.* **11**, 3757. (doi:10.1038/s41598-
021-82249-6)

88. Mezzetti MC, Nardi M, Scapini F. 1997 Variation and ontogeny of the response to visual
stimuli in Mediterranean populations of *Talitrus saltator* (Montagu) (Crustacea-Amphipoda).
J. Exp. Mar. Biol. Ecol. **211**, 129–142. (doi:10.1016/S0022-0981(96)02718-9)

89. Medina JM, Tankersley RA. 2010 Orientation of larval and juvenile horseshoe crabs *Limulus
polyphemus* to visual cues: Effects of chemical odors. *Curr. Zool.* **56**, 618–633.
(doi:10.1093/czoolo/56.5.618)

90. Han R, Wei T-M, Tseng S-C, Lo C-C. 2021 Characterizing approach behavior of *Drosophila
melanogaster* in Buridan's paradigm. *PLOS ONE* **16**, e0245990.
(doi:10.1371/journal.pone.0245990)

91. Suddendorf T, Busby J. 2003 Mental time travel in animals? *Trends Cogn. Sci.* **7**, 391–396.
(doi:10.1016/S1364-6613(03)00187-6)

92. Hollis KL, McNew K, Sosa T, Harrsch FA, Nowbahari E. 2017 Natural aversive learning in
Tetramorium ants reveals ability to form a generalizable memory of predators' pit traps.
Behav. Process **139**, 19–25. (doi:10.1016/j.beproc.2017.03.003)

93. Hawlena D, Kress H, Dufresne ER, Schmitz OJ. 2011 Grasshoppers alter jumping
biomechanics to enhance escape performance under chronic risk of spider predation. *Funct.
Ecol.* **25**, 279–288. (doi:10.1111/j.1365-2435.2010.01767.x)

94. Buehlmann C, Dell-Cronin S, Pathirannahelage AD, Goulard R, Webb B, Niven JE, Graham
P. 2023 Impact of central complex lesions on innate and learnt visual navigation in ants. *J.
Comp. Physiol. A* **209**, 737–746. (doi:10.1007/s00359-023-01613-1)

95. Buehlmann C, Wozniak B, Goulard R, Webb B, Graham P, Niven JE. 2020 Mushroom bodies
are required for learned visual navigation, but not for innate visual behavior, in ants. *Curr.
Biol.* **30**, 3438-3443.e2. (doi:10.1016/j.cub.2020.07.013)

96. Grabowska MJ, Steeves J, Alpay J, Van de Poll M, Ertekin D, van Swinderen B. 2018 Innate
visual preferences and behavioral flexibility in *Drosophila*. *J. Exp. Biol.* **221**, jeb185918.
(doi:10.1242/jeb.185918)

97. Cheng KY, Colbath RA, Frye MA. 2019 Olfactory and neuromodulatory signals reverse visual
object avoidance to approach in *Drosophila*. *Curr. Biol.* **29**, 2058-2065.e2.
(doi:10.1016/j.cub.2019.05.010)

(a) Land-walking

(b) Water-swimming / Water-wading

(c) Glycerol-swimming / Tylose-swimming

(d) Beacon Fixed / Beacon Moving   Walking-without-transition

(e) Island

Figure 1. All experiments were conducted in an arena with a black beacon (height: 45°, width: 20°, measured from the effective centre of the floor) placed against the wall. Arrows and cross marks represent the overall orientation trend. (a) Basic arena used for Land-walking. (b) Arena filled with water for swimming (Water-swimming) or with shallow water for wading (Water-wading). (c) Arena, same as (a), with a transparent dish filled with either a 80% (v/v) glycerol and 20% (v/v) 2-propanol mixture (Glycerol-swimming) or a 1.0% (w/w) tylose aqueous solution (Tylose-swimming). (d) Arena fitted with extruded polystyrene foam insets to enable the transition from swimming in water to walking on a dry substrate. One side of the inset contained a central pool that could be filled with water (for the conditions Beacon Fixed, Beacon Displaced), while the other side served as a control (for condition Walking-without-transition), providing only the uniform dry substrate. The final position of the beacon, which was displaced by 60° under the Beacon Displaced condition, is also illustrated (dashed outline). (e) Arena, same as (b), with a transparent cup in the centre and the water level matched to the base of the cup (Island), effectively creating a central platform surrounded by water ("island").

Figure 2 Trajectories and circular box-and-whisker plots of final bearings are shown under the following conditions: (a) Land-walking (walking on dry substrate, $N = 57$); (b) Water-swimming (swimming in water, $N = 57$); (c) Water-wading (walking in shallow water, $N = 57$); (d) Glycerol-swimming (swimming in an 80% (v/v) glycerol and 20% (v/v) 2-propanol mixture, $N = 47$); and (e) Tylose-swimming (swimming in a 1.0% (w/w) tylose aqueous solution, $N = 59$); (f) Upside-down-walking ($N = 45$). The beacon range is represented as a black arc segment on the left side of each plot. Individual trajectories correspond to single trials. Circular box-and-whisker plots show the median, first quartile, third quartile (coloured interquartile range), minimum, maximum (whiskers), and outliers (black-filled circles) of the final bearing distributions.

(a) Walking-without-transition

(b) Beacon Fixed (pool edge)

(c) Beacon Displaced (pool edge)

(d) Beacon Fixed

(e) Beacon Displaced

(f) Beacon Displaced (body axis)

Figure 3. (a–e) Trajectories and circular box-and-whisker plots of final bearings are shown under the following conditions: Walking-without-transition condition, until ants reached the arena edge (a, N=33); Beacon Fixed condition, until ants reached the pool edge (b, $N = 33$); Beacon Displaced condition, until ants reached the pool edge (c, $N = 33$); Beacon Fixed condition, until ants reached the arena edge (d, $N = 33$); and Beacon Displaced condition, until ants reached the arena edge (e, $N = 33$), following the format of figure 2. For the condition where the beacon was displaced by 60° (e), an arrow indicates the direction of the beacon shift. The initial beacon range is shown in grey, the final beacon range after displacement in black. (f) Circular box-and-whisker plots illustrate the distribution of the body axis direction (from abdomen to head) at the arena edge under Beacon Displaced condition, as described in (e).

(a)

Initial water contact

First immersion

Start of swimming

End of swimming

(b) Initial Phase

2 cm

(c) Contact Phase

2 cm

(d)

90°

0°   ±180°

Before 1st immersion

-90°

Direction of water contacts

(e) Immersion Phase

2 cm

(f)

90°

0°   ±180°

-90°

Direction of immersions

(g) Swimming Phase

10 cm

Figure 4. (a) Representative frame images capturing key moments: Initial water contact, First immersion, Start of swimming, and End of swimming. (b, c) Trajectories during the Initial Phase (b) and Contact Phase (c) are displayed along with circular box-and-whisker plots indicating the directions of initial water contacts for all ants (N = 33), following the format of Figure 2. A representative trajectory is highlighted in magenta, and the water contact positions during the Contact Phase are marked with light yellow stars. Island edges are shown as black dotted circles, and the trajectory scales are located at the bottom right of each plot. (d) A circular box-and-whisker plot illustrates the directional distribution of all water contacts made by ants before the first immersion, formatted as in Figure 3. (e) Trajectories during the Immersion Phase are presented with circular box-and-whisker plots of initial water contacts, consistent with (b). A representative trajectory (magenta) is included, and positions of all immersions are indicated by light blue stars. (f) A circular box-and-whisker plot shows the distribution of water immersion directions before the ants started swimming, as in (d). (g) Trajectories during the Swimming Phase are depicted with circular box-and-whisker plots showing the final swimming bearings. The format is consistent with (b), and the representative trajectory (magenta) is included.