An in-depth dataset of northwestern European arthropod life histories and ecological traits

Garben Logghe[1,2], Femke Batsleer[1], Dirk Maes[2,3], Tristan Permentier[1], Matty P. Berg[4,5], Dimitri Brosens[2], Stijn Cooleman, Pallieter De Smedt[6], Jonas Hagge[7,8], Jorg Lambrechts[9], Marc Pollet[2,10], Fons Verheyde[11], Dries Bonte[1]

Ghent University, Department of Biology, Terrestrial Ecology Unit, Karel Lodewijk Ledeganckstraat 35, 9000 Ghent, Belgium

Research Institute for Nature and Forest (INBO), Herman Teirlinckgebouw, Havenlaan 88 Bus 73, 1000 Brussels, Belgium

Radboud Institute for Biological and Environmental Sciences (RIBES), Radboud University, PO Box 9010, NL-6500 GL Nijmegen, The Netherlands

Amsterdam Institute for Life and Environment (A-LIFE), Section Ecology and Evolution, Vrije Universiteit Amsterdam, De Boelelaan 1108, 1081 HZ Amsterdam, The Netherlands

Groningen Institute for Evolutionary Life Sciences (GELIFES), University of Groningen Nijenborgh 7 / 9747 AG Groningen, The Netherlands

Ghent University, Department of Environment, Forest & Nature Lab, Geraardsbergsesteenweg 267, 9090 Gontrode, Belgium

Northwest German Forest Research Institute, Forest Nature Conservation, Prof.-Oelkers-Str. 6, 34346 Hann. Münden, Germany

University of Göttingen, Forest Nature Conservation, Büsgenweg 3, 37077 Göttingen, Germany

9 Natuurpunt Studie, Coxiestraat 11, 2800 Mechelen, Belgium

10 Operational Directory Taxonomy and Phylogeny, Entomology, Royal Belgian Institute of Natural

Sciences (RBINS), Vautierstraat 29, 1000 Brussels, Belgium

11 Flanders Marine Institute, Jacobsenstraat 1, 8400 Oostende, Belgium

Corresponding author: G. Logghe, Ghent University, Department of Biology, Terrestrial Ecology Unit,

K. L. Ledeganckstraat 35, 9000 Ghent, Belgium. E-mail: garben.logghe@ugent.be

Abstract

In response to the ongoing biodiversity crisis among arthropods, it is essential to implement efficient conservation strategies to safeguard both species diversity and the vital ecosystem services they provide. Developing such strategies requires reliable predictive models that can identify the species that are the most vulnerable to current and future threats, including those posed by climate and land-use change. Species life histories are central to these models, as they influence both population dynamics and spread rates. To support this effort, we compiled a dataset with key traits for arthropods based on several literature sources and expert knowledge. The dataset contains data on body size, life history, thermal niche and ecology for 4874 northwestern European species across 10 different orders. By gathering these essential trait data, we aim to create a robust foundation for predicting species vulnerability and anticipating shifts in arthropod communities in response to global change.

Introduction

Arthropods are an exceptionally species-rich taxon that provide essential ecosystem services such as pollination or nutrient cycling (Stork, 2018; Yang & Gratton, 2014). Despite their critical importance, arthropods remain noticeably understudied in conservation research compared to vertebrates (Clark & May, 2002). This oversight is particularly concerning as these vital organisms are currently under threat from a widespread crisis referred to as "a death by a thousand cuts" (Wagner et al., 2021). The problem stems from a combination of global threats, including climate extremes, pollution, eutrophication, invasive species and urbanization, which have collectively led to significant declines in both the abundance and species richness of arthropods (Harvey et al., 2023; Wagner, 2020). The situation is expected to worsen with impending global change, putting arthropods at even greater risk (Hallmann et al., 2017; Seibold et al., 2019; Soroye et al., 2020).

A key strategy to increase arthropod community resilience against global change involves maintaining biotic complexity (Samways et al., 2020). This can be achieved locally by conserving both at-risk habitats and microhabitat heterogeneity (Toonen et al., 2022). On a regional scale, it requires restoring habitat quality and connectivity, enabling species with restricted environmental tolerances to track their optimal niches under global change (Travis et al., 2013). However, predicting which species will be most vulnerable and where conservation measures should be prioritized remains challenging. Existing predictive models are mostly based on correlative species distribution projections but neglect biological mechanisms such as demography, dispersal and interactions between species. Including these mechanisms is crucial for producing accurate forecasts, as merely extrapolating correlations between species' range and climate is insufficient (Urban et al., 2016).

Implementing these biological mechanisms in predictive models requires detailed information on species' demographic and ecological traits (Comte et al., 2024; Urban et al., 2016). Compiling such trait data for arthropods is challenging for two main reasons: many arthropod groups are less studied compared to vertebrates (Leather, 2013) or plants (Kleyer et al., 2008), and their high species richness, small sizes and hidden life styles make it difficult to quantify key traits. Furthermore, available trait data are mostly scattered across literature, with non-standardized measurements that hinder cross-taxon comparisons (Moretti et al., 2017).

To address this gap, we aimed to compile relevant trait data needed for predictive demographic models of arthropod diversity under global change pressures. We synthesised data on life history traits such as fecundity, dispersal ability and development time from various literature sources and expert judgement. Additionally, we included information on readily available ecological traits like habitat, feeding guild and thermal niche. With the exception of genetic adaption, our goal is to provide data on

72   all necessary ecological components to construct reliable predictive models for forecasting the impact
73   of global change on arthropod species (Urban et al., 2016).

General description

Purpose: This dataset summarizes life history and ecological traits for 4874 European arthropod
species, aiming to construct a comprehensive dataset that could be incorporated in predictive models.
The selection of traits was guided by four of the six mechanisms identified by Urban et al. (2016) that
determine biological responses to climate change: dispersal, demography, species interactions and
physiological processes. We are convinced that most of these mechanisms can also be integrated into
models assessing the effects of other threats, such as land-use change. The first mechanism is
dispersal, defined as the movement of a species from its place of birth to their location of reproduction
(Clobert et al., 2009). The dispersal ability of a species is crucial for its ability to track suitable niches in
new environments, making it a key component of this dataset (Travis et al., 2013). A second important
mechanism is demography, determining population growth rates (Altermatt, 2010; Bommarco, 2001).
It is here represented by the following life history traits: fecundity, development time, voltinism and
lifespan. Overwintering stage was also included due to its strong connection with climate change-
induced phenological shifts, which can significantly alter population dynamics (Marshall et al., 2020).
A third mechanism is species interactions, acknowledging that species do not exist in isolation.
Although it was not feasible to include direct measures of competition (e.g. functional responses) in
this dataset, we did incorporate feeding guild and trophic range, which can be relevant in the context
of global change (Burghardt & Tallamy, 2013; Mitchell & Litt, 2016). The fourth type of mechanisms is
physiological processes, which are crucial to understand population change under climate change. To
address this, we included realised thermal niches to indicate a species' vulnerability to shifts in
temperature regimes. Additionally, diurnality was included, as it may influence species' responses to
climate change (Levy et al., 2019). The final two mechanisms identified by Urban et al. (2016) –
environment and evolution – were not included in this dataset due to practical constraints.
Environmental factors are largely species-independent and should be sourced from other datasets.
Nevertheless, we included habitat information, as it is essential for assessing the responses of specific
communities. The evolution mechanism, which involves genetic variation, was omitted due to the lack
of available data for most arthropod species. In addition to traits related to these six mechanisms, we
included data on body size, overwintering stage and diurnality. Body size is often considered a "master
trait" because it correlates with many other traits related to several of the key mechanisms (White et
al., 2007; Woodward et al., 2005). Body size can determine maximal population densities (Currie, 1993;
White et al., 2007), movement capacities (Hillaert et al., 2018; Logghe et al., 2024), fecundity (Berger
et al., 2008) and thermal limits (Klockmann et al., 2017). The inclusion of body size in our data

consequently allows it to be used as a correlated trait and proxy for potential missing data on specific
mechanisms. With this selection of traits, we aim to facilitate the implementation of biological
mechanisms in predictive models focused on arthropods, thereby enhancing our ability to forecast the
impact of global change on these animals.

Sampling methods

Sampling description: Initially, we conducted an extensive search for literature sources that describe
life history and/or ecological traits of arthropods. Our goal was to include as many arthropod orders
as possible. However, we soon realized that to obtain sufficient data per taxon, we needed to focus on
10 specific orders (see Taxonomic coverage). The data were compiled in separate Excel sheets for each
order. Each source was recorded on a separate line in the dataset, regardless of whether the species
had already been included. This approach allowed us to consider multiple sources when estimating
trait values for different species. Later, this information was merged into a single dataset (see below).
We consulted 83 different literature sources (see Supplement 1 for a full list), including scientific
papers, books and websites (Figure 1). Initially, the search for data was focused on existing datasets or
books with extensive descriptions of specific groups, aiming to find sources with the most
comprehensive information. Subsequently, we targeted specific papers on individual species or small
groups to fill gaps in the dataset.

In the next step, several authors contributed personal data, measurements and expert judgements on
specific arthropod groups to address noticeable gaps in the dataset. These data points were added as
separate rows to the raw datasets, with the contributing author's name cited as the source. One of the
most significant contributions from these experts was the estimation of dispersal ability, defined as a
species' potential to move from its birthplace to new areas. This estimation was invaluable, as dispersal
data is often difficult to obtain from literature or simply unmeasured.

Additionally, we estimated the realised thermal niche for each species in the datasets. This was
accomplished by overlaying species distribution data derived from GBIF (GBIF.org, 2023) with
WorldClim climate data (Fick & Hijmans, 2017) in QGIS (QGIS Development Team, 2023). First, an R
script was developed to automate the retrieval and processing of species occurrence from GBIF. The
script utilizes the "rgbif" package (Chamberlain et al., 2024) to interface directly with the GBIF API,
allowing for large-scale download of biodiversity data for multiple species. Specifically, the script
matches a list of species names to their corresponding "speciesKey" in the GBIF taxonomic backbone.
Once matched, the script initiates a data download using the "occ_download" function, filtering the
data to include only records with geographic coordinates and excluding records with geospatial issues,
null observations, preserved specimens, and fossils. After submitting the download request, GBIF

provides a unique download key, which is used by the script to retrieve the dataset once it is available
using the "occ_download_get" function. The data is imported into an R dataframe, where potential
issues with name matching or data availability can be checked. The script then creates individual CSV
files for each species, containing species names and geographic coordinates. For the next step, a
Python (Van Rossum & Drake Jr., 1995) script was developed to run in QGIS. After loading a raster with
annual mean temperatures (BIO1) derived from WorldClim, the script processes each CSV file to derive
distribution data. The script creates a buffer around each species observation point (0.2 degrees) to
expand it to a small area, ensuring that temperature data is averaged over a meaningful geographic
range rather than at a single point. The zonal statistics tool is then used to calculate temperature
statistics (minimum, maximum, and mean) within each buffer area by extracting raster values (annual
mean temperature) that overlap with the buffers. The resulting data is stored in a CSV file containing
the three temperature measures, with one row per assessed species. Finally, temperature range was
calculated by subtracting the minimum temperature from the maximum.

In the final step, we aimed to create a centralized dataset that consolidated all species traits into a
single, averaged value per trait for each species. This involved merging the raw datasets with the data
on temperature niches and then summarizing each trait per species using R. For continuous traits, such
as body size, we averaged all available values from different sources into a single value per species.
When sources provided separate values for males and females, we first averaged sex-specific values
and then combined them with values from other sources to derive the final species value. For
categorical variables, we adopted the value representing the most inclusive or highest category. For
example, if a species was described as both omnivorous and carnivorous by different sources, we
classified it as omnivorous in the final dataset. Similarly, if one source indicated that a species
overwinters as larva and another source did not, we classified the species as overwintering in the larval
stage in the final dataset. Averaging dispersal ability was particularly challenging due to the significant
variation in dispersal modes and studied proxies (from morphology to behaviour). For example,
dispersal ability in spiders is often experimentally assessed by ballooning propensity, whereas in
beetles, wing load is the most prevalent measure. Different proxies of dispersal, and metrics of
dispersal ability make it further impossible to directly average these values across and within taxa. For
example, one source might use a scale of 1-9 for estimating the dispersal ability of a butterfly, while
another might use a scale of 1-10. To address this, we rescaled the dispersal ability values to a relative
scale between 0.1 and 1 within each order. For example, if a butterfly received the highest dispersal
score from a particular source (e.g. 9 on a 1-9 scale), it was assigned a value of 1. Conversely, the
species with the lowest dispersal score (e.g. 1 on a 1-9 scale) would receive a value of 0.1. This process
was applied separately to each order, ensuring that every group included species with dispersal values

 ranging from 0.1 to 1. By rescaling and averaging dispersal ability within each order, we aimed to create

 relative dispersal values that could be meaningfully compared across different arthropod orders.

 Figure 1: Overview of the proportional contribution of various data sources to each order. Absolute species counts are depicted

 by the white numbers within the graph. Total number of species per order are below order names between brackets.

Geographic coverage

 Currently, the dataset is restricted to species that are native to northwest Europe, which includes

 Belgium, Luxembourg, the Netherlands, northern France, United Kingdom and western Germany.

 Introduced species were mostly avoided, but there are some records of southern European species

 that have been established for a long time (e.g. *Pholcus phalangioides, Porcellionides pruinosus*).

Taxonomic coverage

 The dataset currently covers 4874 arthropod species from 10 distinct orders (Figure 2). The dataset is

 restricted to species with a terrestrial adult stage, which means that groups with aquatic larvae (such

 as Odonata) could still be incorporated into this dataset. Both the orders and species included were

selected based on the availability of relevant trait data. While we aimed to be as inclusive as possible, the representation of orders and families within the dataset varies. Smaller orders like Isopoda (9 families), Odonata (9 families), Orthoptera (6 families) and Opiliones (5 families) are relatively well-covered and nearly complete. In contrast, for larger orders such as Araneae (31 families), Hemiptera (32 families) and Hymenoptera (29 families), the dataset includes about 50% of the species. Coleoptera (75 families) and Diptera (6 families) are relatively less-well represented due to a high number of species and very limited availability of trait data. Within Diptera, most data are limited to the families Syrphidae and Dolichopodidae. Lepidoptera (25 families) also appear rather incomplete when considering all species within the sampled region. However, the dataset focuses on macrolepidoptera, which are well-covered compared to the total number of species in that group.

Figure 2: Overview of the arthropod orders that were included in the dataset. The white numbers represent the total number of species included for each order. All arthropod icons originate from Noun Project (CC BY-NC-ND 2.0). From left to right: Beetle by Rachel Siao, Moth by parkjisun, Cicada by Alejandro Capellan, wasp by parkjisun, fly by Hermine Blanquart, Spider by Matthew Davis, Dragonfly by Hermine Blanquart, Cricket by Ed Harrison, isopod by Pham Thanh Lôc and Spider by Miroslava.

200 No data have been included for taxa with lower taxonomic resolution than species level. For each
201 species, other taxonomic ranks including kingdom, phylum, order, suborder and family were added to
202 the dataset to provide fully detailed information on taxonomic position.

Traits coverage

204 The dataset comprises 28 traits describing ecological and life history traits of the assessed arthropod
205 species (Table 1). These traits were selected based on their significance in predictive modelling
206 concerning the future effects of global change on arthropods (see above for more details).

207 *Table 1: Definition of the 28 traits included in the dataset, based on (Moretti et al., 2017). In several instances, these traits*
208 *constitute subcategories of broader traits. In those cases, only the subcategories have dedicated columns in the dataset.*

Trait category	Subcategory	Definition and description of the trait
Body size		Body length of the adult (in millimetres)
Fecundity		Number of offspring produced by a single female
Development time		Time between laying of the egg and eclosion of the adult (in days)
Lifespan		Amount of time an adult individual lives, from emergence from last instar until death (in days)
Voltinism		Number of generations an organism completes in a single year
	Voltinism mean	Mean number of generations per year; calculated by averaging all possible scenarios (e.g. a species that can have 1 or 2 generations per year would show a value of 1.5)
	Voltinism max	Maximum number of generations per year; partivoltine: new generation emerges every two years univoltine: a single generation per year bivoltine: two generations per year multivoltine: three or more generations per year
Overwintering		Life stage in which the species overwinters (categorical)
	Overwintering egg	0: species never overwinters as egg 1: species overwinters as egg
	Overwintering larva	0: species never overwinters as larva 1: species overwinters as larva
	Overwintering pupa	0: species never overwinters as pupa 1: species overwinters as pupa
	Overwintering adult	0: species never overwinters as adult

		1: species overwinters as adult
Diurnality		diurnal: species is primarily active by day
		cathemeral: species is active by both day and night
		nocturnal: species is primarily active by night
Feeding guild		Expresses the main diet of the species (categorical)
	Feeding guild larva	herbivore: larva feeds on plant material (leaves, stem, roots)
		palynivore: larva feeds on pollen
		omnivore: larva feeds on both plants and animals
		fungivore: larva feeds on fungi
		carnivore: larva feeds on other animals
		detritivore: larva feeds on decaying organic material
	Feeding guild adult	non-eating: adult does not eat
		herbivore: adult feeds on plant material (leaves, stem, roots)
		pollinator: adult feeds on pollen or nectar
		omnivore: adult feeds on both plants and animals
		fungivore: adult feeds on fungi
		carnivore: adult feeds on other animals
		detritivore: adult feeds on decaying organic material
Trophic range		Expresses the dietary breadth of the species (categorical)
	Trophic range larva	monophagous: larva feeds on a single (or a very few related) species
		oligophagous: larva feeds on multiple species within the same taxon
		polyphagous: larva feeds on species from multiple families or orders
	Trophic range adult	monophagous: adult feeds on a single (or a very few related) species
		oligophagous: adult feeds on multiple species within the same taxon
		polyphagous: adult feeds on species from multiple families or orders
Habitat		Expresses the main habitat types where the species feeds and reproduces (categorical)
	Habitat urban	0: the species is not found in urbanised environments
		1: the species occurs in urbanised environments (including indoors, parks and gardens)
	Habitat agriculture	0: the species does not occur in agricultural environments
		1: the species occurs in agricultural environments (including orchards and vineyards)

	Habitat forest	0: the species does not occur in forests 1: the species occurs in forests (including broadleaved forest, mixed forest, coniferous forest and forest edges)
	Habitat grassland	0: the species does not occur in semi-natural to natural grasslands 1: the species occurs in semi-natural to natural grasslands (ranging from extensive meadows to sparsely vegetated chalk grasslands)
	Habitat heathland	0: the species does not occur in heathlands 1: the species occurs in heathlands (including moorlands and bogs)
	Habitat dunes	0: the species does not occur in coastal dunes 1: the species occurs in coastal dunes
	Habitat fresh marshes	0: the species does not occur in freshwater marshes 1: the species occurs in freshwater marshes (including floodplains)
	Habitat salt marshes	0: the species does not occur in salt water marshes 1: the species occurs in salt water marshes (including tidal flats)
Dispersal ability		Expresses the potential of a species to disperse, to move a certain distance from its place of birth (on a relative scale from 0.1 to 1)
Thermal niche		Expresses the thermal niche of the species on a large spatial scale (in °C); values indicate whether the species occurs in colder or warmer climates.
	Thermal mean	Mean average annual temperature within the species range
	Thermal minimum	Minimum average annual temperature within the species range
	Thermal maximum	Maximum average annual temperature within the species range
	Thermal range	Difference between Thermal minimum and Thermal maximum

Data coverage of traits

The amount of available data differs substantially between traits (Figure 3). Data on body size, habitat and distribution (used for calculating thermal niche) were readily available for a wide range of species. However, data on demographic traits like development time, fecundity and lifespan were much more difficult to obtain. Most sources assessing demographic traits tend to focus on a few closely related species, causing a noticeable lack of data availability. It must be noted that even though it seems that dispersal ability has a lot of entries in the dataset, these datapoints are mostly based on proxies or expert judgement. Data on actual dispersal distances of arthropods is extremely rare, mainly due to the inherent challenges that come with tracking these small organisms (Batsleer et al., 2020).

Figure 3: Overview of the number of species for which data on a given trait is available. The trait categories with subcategories (Habitat, Thermal niche, Overwintering & Voltinism; see table 1) have the same number of species for each subcategory for which trait data is available.

Data resources

The final dataset was published on GBIF: https://doi.org/10.15468/75g5z9, raw datasets and code will be available from Zenodo: 10.5281/zenodo.13379714.

Additional information

Limitations

While we aimed to be as complete as possible, this dataset has some important limitations. First, we did not account for intraspecific variation in traits, opting instead to use averaged values for each trait. Similarly, we did not consider potential variation in average trait values across a species' range. This decision was made because many sources report only average values, making data on trait variation difficult to obtain. Furthermore, we acknowledge there are significant limitations to our calculation of thermal niche. These data should not be used as precise thermal limits but rather as an indication whether the species is more adapted to warmer or cooler climates. While data on microclimates or physiological thermal limits would offer a more accurate estimate of thermal tolerance, they are much harder to obtain and less relevant for analysing patterns at large spatial scales and across many taxa. It is also important to note that this method calculates realised thermal niche, meaning that species might have broader thermal niches than observed but can be constricted in their distribution range by, for instance, species interactions or fragmentation (Sax et al., 2013). The accuracy of these calculations can be somewhat compromised for species with few records in GBIF. Nevertheless, we found that the values generally align well for some groups with climate niches reported in previous studies (see Supplement 2). Finally, while we present an extensive dataset covering more than 4000 species, we acknowledge that this represents only a small fraction of the total arthropod diversity in northwestern Europe. This limitation is largely due to the lack of available data for many arthropod groups, as research tends to be biased toward large and more charismatic species (Cardoso, 2012).

Acknowledgements

The authors would like to thank the following people for providing and/or recommending literature: Diana Bowler, Wouter Dekoninck, Frederik Hendrickx, Thomas Merckx, Thomas Parmentier, Arno Thomaes, Elias Van Den Broeck, Luc Vanhercke and Pieter Vantieghem. Special thanks to Uche Osajie, who tried out the method for calculating thermal niches in QGIS and contrasted the results with available literature data.

Author contributions

Garben Logghe, Femke Batsleer, Dirk Maes and Dries Bonte devised the content and structure of the dataset. Garben Logghe collected and synthesised the data. Tristan Permentier and Garben Logghe developed the method for calculating thermal niches. Dimitri Brosens and Stijn Cooleman coordinated the data publishing process. The following authors provided personal data and/or expert judgement for specific taxa: Matty Berg (Isopoda), Pallieter De Smedt (Isopoda), Dirk Maes (Lepidoptera), Jonas Hagge (xylophagous Coleoptera), Jorg Lambrechts (Orthoptera), Marc Pollet (Dolichopodidae) and Fons Verheyde (Hymenoptera). Garben Logghe wrote the first draft of the manuscript. All authors contributed to a first revision of the manuscript.

Funding

Garben Logghe was funded by FWO (grantnr: 1130223N).

References

Altermatt, F. (2010). Climatic warming increases voltinism in European butterflies and moths. *Proceedings of the Royal Society B: Biological Sciences*, *277*(1685), 1281–1287. https://doi.org/10.1098/rspb.2009.1910

Batsleer, F., Bonte, D., Dekeukeleire, D., Goossens, S., Poelmans, W., Van der Cruyssen, E., Maes, D., & Vandegehuchte, M. L. (2020). The neglected impact of tracking devices on terrestrial arthropods. *Methods in Ecology and Evolution*, *11*(3), 350–361. https://doi.org/10.1111/2041-210X.13356

Berger, D., Walters, R., & Gotthard, K. (2008). What limits insect fecundity? Body size- and temperature-dependent egg maturation and oviposition in a butterfly. *Functional Ecology*, *22*(3), 523–529. https://doi.org/10.1111/j.1365-2435.2008.01392.x

Bommarco, R. (2001). Using matrix models to explore the influence of temperature on population growth of arthropod pests. *Agricultural and Forest Entomology*, *3*(4), 275–283. https://doi.org/10.1046/j.1461-9555.2001.00114.x

Bowler, D. E., Hof, C., Haase, P., Kröncke, I., Schweiger, O., Adrian, R., Baert, L., Bauer, H.-G., Blick, T., Brooker, R. W., Dekoninck, W., Domisch, S., Eckmann, R., Hendrickx, F., Hickler, T., Klotz, S.,

Kraberg, A., Kühn, I., Matesanz, S., … Böhning-Gaese, K. (2017). Cross-realm assessment of

climate change impacts on species' abundance trends. *Nature Ecology & Evolution*, *1*(3),

0067. https://doi.org/10.1038/s41559-016-0067

Burghardt, K. T., & Tallamy, D. W. (2013). Plant origin asymmetrically impacts feeding guilds and life

stages driving community structure of herbivorous arthropods. *Diversity and Distributions*,

19(12), 1553–1565. https://doi.org/10.1111/ddi.12122

Cardoso, P. (2012). Habitats Directive species lists: Urgent need of revision. *Insect Conservation and*

Diversity, *5*(2), 169–174. https://doi.org/10.1111/j.1752-4598.2011.00140.x

Chamberlain, S., Barve, V., Mcglinn, D., Oldoni, D., Desmet, P., Geffert, L., & Ram, K. (2024). *rgbif:*

Interface to the Global Biodiversity Information Facility API (Version 3.7.9) [Computer

software]. https://CRAN.R-project.org/package=rgbif

Clark, J. A., & May, R. M. (2002). Taxonomic bias in conservation research. *Science, 297*(5579), 191–

192.

Clobert, J., Le Galliard, J.-F., Cote, J., Meylan, S., & Massot, M. (2009). Informed dispersal,

heterogeneity in animal dispersal syndromes and the dynamics of spatially structured

populations. *Ecology Letters*, *12*(3), 197–209. https://doi.org/10.1111/j.1461-

0248.2008.01267.x

Comte, L., Bertrand, R., Diamond, S., Lancaster, L. T., Pinsky, M. L., Scheffers, B. R., Baecher, J. A.,

Chen, C., Lawlor, J. A., Moore, N. A., Oliveira, B. F., Murienne, J., Rolland, J., Rubenstein, M.

A., Sunday, J., Thompson, L. M., Villalobos, F., Weiskopf, S. R., & Lenoir, J. (2024). Bringing

traits back into the equation: A roadmap to understand species redistribution. *Global Change*

Biology, 30, e17271. https://doi.org/10.1111/gcb.17271

Currie, D. J. (1993). What Shape Is the Relationship between Body Size and Population Density?

Oikos, *66*(2), 353. https://doi.org/10.2307/3544825

Fick, S. E., & Hijmans, R. J. (2017). WorldClim 2: New 1-km spatial resolution climate surfaces for

global land areas. *International Journal of Climatology*, *37*(12), 4302–4315.

https://doi.org/10.1002/joc.5086

GBIF.org. (2023). *GBIF Occurence Download* [Computer software].

https://doi.org/10.15468/dl.e7kudt

Hallmann, C. A., Sorg, M., Jongejans, E., Siepel, H., Hofland, N., Schwan, H., Stenmans, W., Müller, A.,

Sumser, H., Hörren, T., Goulson, D., & de Kroon, H. (2017). More than 75 percent decline

over 27 years in total flying insect biomass in protected areas. *PLOS ONE*, *12*(10), e0185809.

https://doi.org/10.1371/journal.pone.0185809

Harvey, J. A., Tougeron, K., Gols, R., Heinen, R., Abarca, M., Abram, P. K., Basset, Y., Berg, M., Boggs,

C., Brodeur, J., Cardoso, P., de Boer, J. G., De Snoo, G. R., Deacon, C., Dell, J. E., Desneux, N.,

Dillon, M. E., Duffy, G. A., Dyer, L. A., … Chown, S. L. (2023). Scientists' warning on climate

change and insects. *Ecological Monographs*, *93*(1), e1553. https://doi.org/10.1002/ecm.1553

Hillaert, J., Hovestadt, T., Vandegehuchte, M. L., & Bonte, D. (2018). Size-dependent movement

explains why bigger is better in fragmented landscapes. *Ecology and Evolution*, *8*(22), 10754–

10767. https://doi.org/10.1002/ece3.4524

Kleyer, M., Bekker, R. M., Knevel, I. C., Bakker, J. P., Thompson, K., Sonnenschein, M., Poschlod, P.,

van Groenendael, J. M., Klimes, L., Klimesova, J., Klotz, S., Rusch, G. M., Hermy, M., Adriaens,

D., Boedeltje, G., Bossuyt, B., Dannemann, A., Endels, P., Götzenberger, L., … Peco, B. (2008).

The LEDA Traitbase: A database of life-history traits of the Northwest European flora. *Journal*

of Ecology, 96, 1266–1274. https://doi.org/10.1111/j.1365-2745.2008.01430.x

Klockmann, M., Günter, F., & Fischer, K. (2017). Heat resistance throughout ontogeny: Body size

constrains thermal tolerance. *Global Change Biology*, *23*(2), 686–696.

https://doi.org/10.1111/gcb.13407

Leather, S. R. (2013). Institutional vertebratism hampers insect conservation generally; not just

saproxylic beetle conservation. *Animal Conservation*, *16*(4), 379–380.

https://doi.org/10.1111/acv.12068

Levy, O., Dayan, T., Porter, W. P., & Kronfeld-Schor, N. (2019). Time and ecological resilience: Can

diurnal animals compensate for climate change by shifting to nocturnal activity? *Ecological*

Monographs, *89*(1), e01334. https://doi.org/10.1002/ecm.1334

Logghe, G., Taelman, C., Van Hecke, F., Batsleer, F., Maes, D., & Bonte, D. (2024). Unravelling

arthropod movement in natural landscapes: Small-scale effects of body size and weather

conditions. *Journal of Animal Ecology*, *23*(9), 1365–1379. https://doi.org/10.1111/1365-

2656.14161

Marshall, K. E., Gotthard, K., & Williams, C. M. (2020). Evolutionary impacts of winter climate change

on insects. *Current Opinion in Insect Science*, *41*, 54–62.

https://doi.org/10.1016/j.cois.2020.06.003

Mitchell, A. B., & Litt, A. R. (2016). Nonnative plant shifts functional groups of arthropods following

drought. *Biological Invasions*, *18*(5), 1351–1361. https://doi.org/10.1007/s10530-016-1072-y

Moretti, M., Dias, A. T. C., Bello, F., Altermatt, F., Chown, S. L., Azcárate, F. M., Bell, J. R., Fournier, B.,

Hedde, M., Hortal, J., Ibanez, S., Öckinger, E., Sousa, J. P., Ellers, J., & Berg, M. P. (2017).

Handbook of protocols for standardized measurement of terrestrial invertebrate functional

traits. *Functional Ecology*, *31*(3), 558–567. https://doi.org/10.1111/1365-2435.12776

Osajie, U. (2021). *Integratie van de temperatuurniche in de levensgeschiedenis van enkele Arthropoda*

en de Global Biodiversity Information Facility (GBIF) als bron voor temperatuurdata [Bachelor

thesis]. Ghent University.

QGIS Development Team. (2023). *QGIS Geographic Information System* [Computer software]. Open

Source Geospatial Foundation Project. http://qgis.osgeo.org

Samways, M. J., Barton, P. S., Birkhofer, K., Chichorro, F., Deacon, C., Fartmann, T., Fukushima, C. S.,

Gaigher, R., Habel, J. C., Hallmann, C. A., Hill, M. J., Hochkirch, A., Kaila, L., Kwak, M. L., Maes,

D., Mammola, S., Noriega, J. A., Orfinger, A. B., Pedraza, F., … Cardoso, P. (2020). Solutions

for humanity on how to conserve insects. *Biological Conservation*, *242*, 108427.

https://doi.org/10.1016/j.biocon.2020.108427

Sax, D. F., Early, R., & Bellemare, J. (2013). Niche syndromes, species extinction risks, and

management under climate change. *Trends in Ecology & Evolution*, *28*(9), 517–523.

https://doi.org/10.1016/j.tree.2013.05.010

Schweiger, O., Harpke, A., Wiemers, M., & Settele, J. (2014). CLIMBER: Climatic niche characteristics

of the butterflies in Europe. *ZooKeys*, *367*, 65–84. https://doi.org/10.3897/zookeys.367.6185

Seibold, S., Gossner, M. M., Simons, N. K., Blüthgen, N., Müller, J., Ambarlı, D., Ammer, C., Bauhus, J.,

Fischer, M., Habel, J. C., Linsenmair, K. E., Nauss, T., Penone, C., Prati, D., Schall, P., Schulze,

E.-D., Vogt, J., Wöllauer, S., & Weisser, W. W. (2019). Arthropod decline in grasslands and

forests is associated with landscape-level drivers. *Nature*, *574*(7780), 671–674.

https://doi.org/10.1038/s41586-019-1684-3

Soroye, P., Newbold, T., & Kerr, J. (2020). Climate change contributes to widespread declines among

bumble bees across continents. *Science*, *367*(6478), 685–688.

https://doi.org/10.1126/science.aax8591

Stork, N. E. (2018). How Many Species of Insects and Other Terrestrial Arthropods Are There on

Earth? *Annual Review of Entomology*, *63*(1), 31–45. https://doi.org/10.1146/annurev-ento-

020117-043348

Toonen, M., Van Den Berge, S., De Frenne, P., & Vandekerkhove, K. (2022). Klimaatadaptatie en

natuurbeheer. In *Van Uytvanck, J. et al. Natuurbeheer: Praktijk en wetenschap hand in hand*

(pp. 451–468). Sterck & De Vreese, Gorredijk.

Travis, J. M. J., Delgado, M., Bocedi, G., Baguette, M., Bartoń, K., Bonte, D., Boulangeat, I., Hodgson, J.

A., Kubisch, A., Penteriani, V., Saastamoinen, M., Stevens, V. M., & Bullock, J. M. (2013).

Dispersal and species' responses to climate change. *Oikos*, *122*(11), 1532–1540.

https://doi.org/10.1111/j.1600-0706.2013.00399.x

Urban, M. C., Bocedi, G., Hendry, A. P., Mihoub, J.-B., Pe'er, G., Singer, A., Bridle, J. R., Crozier, L. G.,

De Meester, L., Godsoe, W., Gonzalez, A., Hellmann, J. J., Holt, R. D., Huth, A., Johst, K., Krug,

C. B., Leadley, P. W., Palmer, S. C. F., Pantel, J. H., … Travis, J. M. J. (2016). Improving the

forecast for biodiversity under climate change. *Science*, *353*(6304), aad8466.

https://doi.org/10.1126/science.aad8466

Van Rossum, G., & Drake Jr, F. L. (1995). *Python reference manual*. Centrum voor Wiskunde en

Informatica, Amsterdam.

Wagner, D. L. (2020). Insect Declines in the Anthropocene. *Annual Review of Entomology, 65*, 457–

480. https://doi/10.1146/annurev-ento-011019-025151

Wagner, D. L., Grames, E. M., Forister, M. L., Berenbaum, M. R., & Stopak, D. (2021). Insect decline in

the Anthropocene: Death by a thousand cuts. *Proceedings of the National Academy of*

Sciences, *118*(2), e2023989118. https://doi.org/10.1073/pnas.2023989118

White, E. P., Ernest, S. K. M., Kerkhoff, A. J., & Enquist, B. J. (2007). Relationships between body size

and abundance in ecology. *Trends in Ecology & Evolution*, *22*(6), 323–330.

https://doi.org/10.1016/j.tree.2007.03.007

Woodward, G., Ebenman, B., Emmerson, M., Montoya, J., Olesen, J., Valido, A., & Warren, P. (2005).

Body size in ecological networks. *Trends in Ecology & Evolution*, *20*(7), 402–409.

https://doi.org/10.1016/j.tree.2005.04.005

Yang, L. H., & Gratton, C. (2014). Insects as drivers of ecosystem processes. *Current Opinion in Insect*

Science, *2*, 26–32. https://doi.org/10.1016/j.cois.2014.06.004