Loss of olfaction reduces caterpillar performance and increases susceptibility to a

natural enemy

Qi Wang [1], Yufei Jia [1], Hans M. Smid [1], Berhane T. Weldegergis [1], Liana O. Greenberg [2],

Maarten Jongsma [3], Marcel Dicke [1], Alexander Haverkamp [1,4]*

[1] Laboratory of Entomology, Wageningen University & Research, Wageningen, the Netherlands

[2] Biosystematics Group, Wageningen University & Research, Wageningen, the Netherlands

[3] Business Unit Bioscience, Wageningen University & Research, Wageningen, the Netherlands

[4] Lead contact

* Correspondence: alexander.haverkamp@wur.nl (A.H.)

Abstract

Insect herbivores such as caterpillars, are under strong selection pressure from natural enemies, especially parasitoid wasps. Although the role of olfaction in host-plant seeking has been investigated in great detail in parasitoids and adult lepidopteran, the caterpillar olfactory system and its significance in tri-trophic interactions remains poorly understood. In this study, we investigated the olfactory system of *Pieris brassicae* caterpillars and the importance of olfactory information in the interactions among this herbivore, its host-plant *Brassica oleracea* and its primary natural enemy *Cotesia glomerata*. To examine the role of olfaction, we utilized CRISPR/Cas9 to knock out the odorant receptor co-receptor (*Orco*). This knockout (KO) impaired olfactory detection and primary processing in the brain. *Orco* KO caterpillars exhibited reduced weight and lost preference for their optimal food plant. Interestingly, the KO caterpillars also experienced reduced weight when challenged by the parasitoid *C. glomerata* whose ovipositor had been removed, and the mortality of the KO caterpillars under the attack of unmanipulated parasitoids increased. We then investigated the behavior of *P. brassicae* caterpillars in response to volatiles from plants attacked by conspecific caterpillars and volatiles from plants on which the caterpillars were themselves attacked by *C. glomerata*. After analyzing the volatile compounds involved in these interactions, we concluded that olfactory information enables caterpillars to locate suitable food sources more efficiently as well as selecting enemy-free spaces. Our results reveal the crucial role of olfaction in caterpillar feeding and natural enemy avoidance, highlighting the significance of chemoreceptor genes in shaping ecological interactions.

Keywords: insect larvae, natural enemy, olfaction, multitrophic interactions, Orco, host-plant selection

Introduction

Chemoreception plays a critical role in the insect life cycle. For instance, insects exploit chemical cues to locate food plants, to find and select mates, to evaluate oviposition sites, and to avoid natural enemies (Dicke & Grostal, 2001). Olfaction, a pivotal component of chemoreception, involves multiple processes in insects and has been extensively investigated in many species (Haverkamp et al., 2018), including its role in the interactions with conspecifics and natural enemies (Kannan et al., 2022). Thus far, the study of olfaction in insects has largely centered upon imagos, the final life stage of insects. However, many insects are holometabolous and spend a significant portion of their lives as larvae, during this life stage chemical information likely play an essential role in host-plant choice and natural-enemy avoidance. Nonetheless, the olfactory system in larval herbivores remains largely unexplored.

Tri-trophic interactions among plants, herbivores and natural enemies have been widely investigated in the last few decades, and chemoreception is considered to play an important role in these systems (Turlings & Erb, 2018; Vet & Dicke, 1992; Wang, 2023). In particular, abundant research has demonstrated that olfactory information, such as plant volatiles, are crucial in the coevolution of herbivores and natural enemies (Dicke & Baldwin, 2010; Haverkamp & Smid, 2020; Turlings & Erb, 2018). In addition to plant volatiles, chemicals emitted by the insects themselves also play an important role (Ebrahim et al., 2015; Venugopal et al., 2020). Under the strong selection pressure from natural enemies, herbivores have developed several integrated avoidance strategies based on the detection of chemical cues. *Drosophila melanogaster* fruit flies, for example, prefer to lay eggs on citrus fruits because the volatiles emitted by citrus are repellent to the endoparasitoid *Leptopilina boulardi*, a natural enemy of fruit fly larvae (Dweck et al., 2013). Similarly, brown planthoppers deposit their eggs on rice plants infested by the striped stem borer, which is repellent to the egg parasitoid *Anagrus nilaparvatae* (Hu et al., 2020).

Insect olfaction is dominated by two classes of olfactory receptor genes, odorant receptors (ORs) and ionotropic receptors (IRs). ORs detect the majority of volatiles in the insect habitat. In all ORs the odorant receptor coreceptor (Orco) is required to form a functional ion channel in the membrane of the odorant receptor neurons (ORNs) (Jones et al., 2005; Sato et al., 2008; Wicher et al., 2008). Previous research indicates that silencing the *Orco* gene can lead to a severe olfaction deficiency, causing a loss of social behavior, impaired mating and significantly reduced foraging behavior (Fandino et al., 2019; Trible et al., 2017; Yan et al., 2017). Previous reports have also clearly shown that adult insect behavior

and neuronal development were greatly affected when *Orco* is knocked out (Trible et al., 2017; Yan et al., 2017). Yet, potential changes in olfaction-deficient larvae remain largely unexplored and little is known about how olfaction contributes to the co-adaptation of larvae and their natural enemies, especially in the larvae of non-model insects such as lepidopteran caterpillars.

Pieris brassicae is a very common butterfly species that causes substantial agricultural production loss across Europe and the Indian sub-continent (Feltwell, 1982; Hasan & Ansari, 2010; Hasan & Ansari, 2011). The butterflies utilize chemoreception to evaluate oviposition sites and host-plants (van Loon et al., 1992a; van Loon et al., 1992b; Wang et al., 2023a). The caterpillars also have a well-developed olfactory system (Wang et al., 2023b) and chemical communication has been demonstrated to be involved in the interaction between herbivores and their natural enemies (Dicke & Baldwin, 2010; Stam et al., 2014). Natural enemies such as parasitoid wasps are highly attracted to volatiles from cabbage plants infested by *P. brassicae* caterpillars (Mattiacci et al., 1994). *P. brassicae* caterpillars are thus under strong selection pressure from natural enemies, though caterpillars can defend themselves by spitting at natural enemies while being attacked.

To gain comprehensive insight into the role of olfaction in *P. brassicae*, and to understand the role of olfaction in their coevolution with natural enemies, we employed a well-established research system utilizing *Brassica oleracea* as host-plants, *P. brassicae* caterpillars as herbivores and *C. glomerata* parasitoid wasps as natural enemies. In this study, we knocked out *Orco* by CRISPR/Cas9 in *P. brassicae* to investigate the influence of odor-guided behavior by this primary herbivore on the next trophic levels. The knockout was verified in caterpillars by staining Orco and glomeruli in larval antennae and the larval antennal center (LAC) respectively and was further confirmed in butterflies by electrophysiological investigations. We also examined the egg-laying behavior of KO butterflies, which is the basis of establishing homozygous KO insect colony for our subsequent studies. The development of caterpillars on cabbage plants was evaluated, and the performance of caterpillars under the threat of natural enemies was studied to investigate the ecological significance of larval olfaction in interacting with their natural enemies. The larval host-plant seeking behaviors were further evaluated to determine the importance of olfaction in locating food sources. Volatiles emitted by caterpillars and parasitoids were analyzed to identify candidate chemicals that might play a role in this interaction. Our study provides further knowledge on the olfactory system in insect herbivores and might help to clarify the ecological significance of olfaction in the interaction with their natural enemies.

Results

Orco knockout and neuronal deficiency

Through screening all butterflies that were reared from injected eggs using PCR and sequencing, we identified an *Orco* mutant line in *P. brassicae*. This mutant line exhibited an 11 bp insertion and a four bp deletion in the second exon, resulting in a shifted reading frame (Fig 1A). The Orco amino acid sequence showed a significant change mediated by an early stop codon (Fig 1B). These results indicate that the knockout (KO) of the *Orco* gene was successful.

To further validate the KO in *P. brassicae* and investigate the neuronal mechanism underlying olfactory deficiency in KO caterpillars, we performed staining of Orco in third-instar larvae (L3) using a lepidopteran specific Orco antibody, which has been validated previously (Fandino et al., 2019; Nolte et al., 2016; Wang et al., 2023b). In WT L3 caterpillar antennae, a total of 34 ORNs were successfully identified (Wang et al., 2023b) (Fig 1C). However, no Orco-positive ORNs were found in the antennae of KO caterpillars (Fig 1D). Similarly, we detected an Orco-positive neuron in the maxillary palps of L3 WT caterpillars (Fig 1E), but no Orco-positive signal was detected in the maxillary palps of KO caterpillars (Fig 1F). By tracing the ORN axons to their glomeruli in the brain, we observed some differences between WT and *Orco* KO caterpillars. In the larval antennal center (LAC), glomeruli displayed well-defined shapes with clear boundaries in the WT LAC (Fig 1G). However, in the KO LAC, the glomeruli were less clearly separated (Fig 1H). Only 7-12 glomeruli were identified in KO LAC, which was a significant reduction compared to WT LAC (P = 0.0066, Wilcoxon rank-sum test) (Fig 1I).

Butterfly electrophysiological response to chemical compounds and oviposition

We then conducted additional tests comparing the antennal electrophysiological responses of male and female butterflies from both genotypes using electroantennography (EAG), to further confirm the KO of *Orco* in butterflies. *Orco* KO butterflies of both sexes displayed a complete loss of responses to the tested chemicals with the exception of acetic acid, n-caproic acid and hexanal, where no significant

difference was detected between KO and WT butterflies. Reduced responses were recorded in KO butterflies to 1-nonanal, (Z)-2-pentenal and (E)-2-hexenal. Besides, WT male and female butterflies responded similarly to most of the volatile compounds with a few exceptions where females showed slight responses to 1-methoxy-2-propanol oil ($P < 0.0001$, Kruskal-Wallis test) and methyl ITC compared to paraffin oil ($P < 0.0001$, Student's t-test) while males did not. Moreover, *Orco* KO males did not show significantly different responses to 3-methylbutanal and 3-pentanone compared to WT males. *Orco* KO females did not show significantly different responses to methyl ITC, 2,3-butanedione and dimethyl disulphide compared to WT females (Fig 2A and 2B).

To investigate the role of olfaction in butterfly mating and oviposition, and evaluate the feasibility of establishing homozygous *Orco* KO colony for subsequent larval studies, pairs of newly emerged butterflies were collected and reared in a cage and allowed to mate freely. Upon dissection, it was found that KO butterflies mated fewer times than WT butterflies ($P = 0.0199$, Wilcoxon rank-sum test), since the majority of the KO butterflies (9 out of 14) did not mate during the experiment (Fig 2C). Out of the five KO butterflies that did mate, all but one mated singly. Of the wildtype butterflies, all but 3 of 11 mated, with most mating twice and one individual even thrice. Female KO butterflies also deposited fewer eggs than WT butterflies during the experiment ($P < 0.0001$, GLM negative binominal) (Fig 2D). To determine the hatching rate, the number of newly hatched caterpillars of both genotypes were counted. It was found that virgin butterflies of both genotypes did not deposit any fertilized eggs, while mated butterflies exhibited similar fecundity with comparable hatching rates between the two genotypes without any significant difference (S1 Fig).

Caterpillar performance on cabbage plants

To assess the ecological significance of knocking out *Orco* in *P. brassicae* caterpillars, we first conducted a comparison of caterpillar growth between the two genotypes on their optimal host-plant. The results show that in the absence of parasitoids, WT caterpillars exhibited better performance, gaining more weight compared to KO caterpillars ($P = 0.0268$, Student's t-test) (Fig 3A). However, there is no developmental difference while being reared in a Petri dish in which a group of caterpillars were supplied with cut out leaf discs (S2 Fig). Subsequently, to determine if the differences in feeding and survival are dependent on food-plant location, a two-choice assay was conducted with cabbage leaf discs, tomato leaf discs and green paper discs in a glass Petri dish to investigate the role of olfaction in host-plant selection

(Fig 3B). Both caterpillar genotypes chose the cabbage leaf disc significantly more often than the green paper discs (P = 0.0016 for KO; P < 0.0001 for WT, Chi-square test), and no difference was detected between the two genotypes (P = 0.1460, Chi-square test). Similarly, when the cabbage leaf disc was replaced with a tomato leaf disc, both genotypes preferred the tomato leaf disc over the paper disc (P = 0.0143 for KO; P = 0.0009 for WT, Chi-square test) and no difference was found between the two genotypes (P = 0.5477, Chi-square test). However, when caterpillars were provided with both a cabbage leaf disc and a tomato leaf disc, a contrasting preference was observed between the two genotypes (P = 0.0004, Chi-square test). KO caterpillars significantly preferred the tomato leaf disc (P = 0.0466, Chi-square test), while WT caterpillars preferred the cabbage leaf disc (P = 0.0028, Chi-square test).

We then aimed to investigate if caterpillars could locate the optimal host-plant when supplied with differently treated plants using a Y-tube olfactometer. We compared the behaviors of both WT and KO caterpillars in response to clean air, a healthy plant and a caterpillar-infested plant (Fig 3C). Both WT and KO caterpillars tended to prefer a healthy cabbage plant to a clean air control, however WT caterpillars were significantly attracted by a healthy plant (P = 0.0164, Chi-square test) while the preference for KO caterpillars was not significant (P = 0.2207, Chi-square test). Subsequently, we challenged the caterpillars with a healthy plant and an infested plant and found that WT caterpillars prefer a healthy plant to an infested plant (P = 0.0495, Chi-square test), while KO caterpillars did not show any preference between the two provided plants (P = 0.7055, Chi-square test). We further compared larval host-plant seeking behaviors with the presence of natural enemies on caterpillar-infested plants (Fig 3D). WT caterpillars showed a significant preference for infested plants without parasitoid wasps *C. glomerata* (P = 0.0067, Chi-square test), while KO caterpillars did not show any preference for either odor source (P = 0.7055, Chi-square test).

Next we tested whether these differences in the foraging behavior also lead to differences in the development and survival of *P. brassicae* caterpillars when exposed to their natural enemies. We first evaluated caterpillar performance under the potential threat of natural enemies by exposing them to female parasitoids from which the ovipositor had been removed, so that these parasitoids could attack, but not kill the caterpillars (Fig 3E). The results show a significantly higher weight in WT caterpillars than in KO caterpillars (P = 0.0088, Student's t-test). To further explore the effect of knocking out *Orco* on the interaction of the caterpillars with their natural enemies, we compared the survival rates of caterpillars when exposed to two mated unmanipulated female *C. glomerata* wasps. The findings show

181   that KO caterpillars had significantly lower survival rates compared to WT caterpillars ($P = 0.0120$, GLM

182   beta-binomial) (Fig 3F).

Volatile headspace composition

Finally, we investigated if caterpillar- and parasitoid-derived chemicals are involved in the

behavioral responses. We identified a total of 45 chemical compounds among the five treatments

belonging to seven classes, including alcohols, aldehydes, aromatics, ketones, nitrogen and/or sulfur-

containing chemicals, terpenoids and others (S3 Table). The contribution of identified chemicals towards

separating different treatment groups was analyzed using principal component analysis (PCA). This

analysis showed a significant separation of most groups with PC 1 carrying 65.4 % of the variance ($P =$

0.001, PERMANOVA). *Pieris brassicae* caterpillars (Pb), *P. brassicae* caterpillar spit (Pb-S), *C.*

glomerata female wasps (Cg) and caterpillar frass (Pb-Fr) treatment groups were clearly separated (Pb

vs. Cg, Pb vs. Pb-S, Pb vs. Pb-Fr, Pb-S vs. Cg and Pb-Fr vs. Cg, P= 0.0013; Pb-S vs. Pb-Fr, P= 0.0022

pairwise PERMANOVA, Fig 4A), whereas the identified and measured volatile blend from *P. brassicae*

caterpillars (Pb) was similar to that of the *P. brassicae* caterpillars-*C. glomerata* parasitoids combination

(Pb-Cg) ($P = 0.0330$, pairwise PERMANOVA). The amount of volatiles among the five treatments was

also compared. Most of the identified compounds were found in significantly higher amounts in Pb-S

and/or Pb-Fr compared to the other treatments (Fig 4B, S3 Table). We also found that some chemicals

are strongly correlated. The identified chemicals are classified into several subgroups in the hierarchical

clustering heatmap according to the abundance among treatments. For instance, the seven chemicals (*Z*)-

3-hexen-1-ol, 1-penten-3-ol, dimethyl disulfide, methyl (methylthio) methyl sulfide, dimethyl trisulfide,

2-methylbutanal and 3-methylbutanal formed a clade. Similarly, the six chemicals, 3-methylbutanoic

acid, 1-pentanol, benzyl cyanide, 3-(methylthio) propanal, (*Z*)-2-hexen-1-ol and phenylacetaldehyde

formed another clade. The clustering analysis demonstrated that chemicals within the same clade are

highly correlated. The multivariate data analysis (MVDA) also showed similar clustering when

comparing the different treatments (S3-S7 Fig).

Based on the results of our chemical analysis, we selected 11 candidate chemicals (1-hexanol, 1-

methoxy-2-propanol, 1-penten-3-ol, 2,3-butanedione, 3-methylbutanal, benzyl cyanide, dimethyl

disulfide, pentanal, (*E*)-2-hexen-1-ol, (*Z*)-2-penten-1-ol and (*Z*)-3-hexen-1-ol) with high variable

importance in the projection (VIP) scores (> 1.0) (S3-S8 Table), and another four common plant volatiles

(linalool, (*Z*)-3-hexenyl acetate, hexanal and (*E*)-2-hexenal). These odorants were chosen to test the behavioral response of L3 caterpillars. Using a custom-made multiple channel setup (S7 Fig), we conducted behavioral experiments with the caterpillars. The results showed that WT caterpillars displayed a preference for two plant chemicals, namely (*Z*)-3-hexen-1-ol and (*E*)-2-hexenal over paraffin oil (P = 0.0482 and P = 0.0321) (Fig 4D). In contrast, KO caterpillars did not exhibit any significant attraction or avoidance behavior towards the tested chemicals (Fig 4E). Additionally, we found that WT caterpillars were more active in the arena, moving around (S8 Fig), while KO caterpillars were less active in the arena, often remaining at the same position within the channels (S9 Fig).

Discussion

Chemical communication plays a significant role in the interaction between plants, herbivores and their natural enemies. Infested host-plants emit herbivore-induced plant volatiles (HIPVs) and thereby attract natural enemies such as parasitoid wasps (Mattiacci et al., 1994). While the role of adult herbivores' olfaction in mediating these tri-trophic interactions has been investigated, the significance of caterpillar olfaction in tri-trophic interactions of plants, herbivorous and their natural enemies has remained largely underexplored. In this study, we severely impaired the olfaction of *P. brassicae* by knocking out *Orco* by CRISPR/Cas9 to investigated the ecological importance of olfaction in host-plant choice and enemy avoidance by *Pieris brassicae* larvae. The knockout of olfaction was successful, as we did not find Orco-proteins in the antennae of *Orco* KO caterpillars and showed an altered brain structure in the *Orco* KO caterpillars in comparison to WT caterpillars. Furthermore, the near complete lack of response to odors by *Orco* KO butterfly antennae compared to wildtype confirms that olfaction was largely lost. Intriguingly, although mating and oviposition were significantly reduced, KO butterflies still reproduce some homozygous offspring. After this characterization of our mutant line, we aimed to test the role of larval olfaction in a multitrophic framework. We found that the loss of olfaction had a significant effect on the foraging behavior of the caterpillars, which exhibited a reduced growth rate, a greater vulnerability to the parasitoid wasp *C. glomerata*.

The perception of volatile chemicals is mediated by the olfactory signal transduction cascade in the insect antennae and the further processing of information in the brain. In the *P. brassicae* KO caterpillar antennae, no Orco protein was detected, and chemical signals can therefore not be converted into electrical signals (Sato et al., 2008; Wicher et al., 2008). Interestingly, the malfunctional Orco decreased

the number of glomeruli in the LAC, indicating that Orco is likely to play a role in the glomerular

organization of the LAC (Fig 1C-I). Similar changes have been reported for adult insects of a few

hymenopteran species: the ants *Harpegnathos saltator* and *Ooceraea biroi* and the honeybee *Apis*

mellifera, where the number and total size of glomeruli decreased while the volume of individual

glomeruli increased in *Orco* KO mutants (Chen et al., 2021; Trible et al., 2017; Yan et al., 2017). In

contrast, the antennal lobe morphology of *D. melanogaster* was not significantly changed when *Orco*

was knocked out (Larsson et al., 2004). In Lepidoptera, the size and number of glomeruli in *Helicoverpa*

armigera were found to be comparable between WT and KO adult moths without any significant

difference (Fan et al., 2022). However, silencing the pheromone receptor, *SlitOR5* led to a decreased size

of a glomerulus in the adult moth *Spodoptera littoralis* and knocking out *Orco* also led to a reduced size

of OR-related glomeruli in adult *Manduca sexta* (Fandino et al., 2019; Koutroumpa et al., 2022). The

loss of olfactory glomeruli in the caterpillar might, therefore, either indicate that the formation of

glomeruli in the embryo is mediated by a different mechanism than during pupation (Oland & Tolbert,

2011), or that the loss of neuronal activity reduces axonal innervations and thereby the size of the

glomeruli to an extent that they could no longer be visualized by our method. However, the mechanism

of glomerular formation and the roles that olfactory genes play in this process needs to be further

investigated in both immature and adult insects (Williams et al., 2022).

The role of olfaction in the behavioral ecology of adult insects has already been studied in quite

some detail, and silencing *Orco* has been an important tool for unravelling the importance of chemical

communication in adults of various insect species. For instance, in *Orco* KO adults of several insect

species, there was a significant reduction in the electrophysiological responses to volatile chemicals such

as alcohols and esters (Fandino et al., 2019; Sun et al., 2020). Consequently, *Orco* mutants exhibited

notable changes in host-plant location and severely reduced fecundity due to impaired pheromone

perception and mating, which also resulted in significantly reduced egg-laying behavior (Fan et al., 2022;

Fandino et al., 2019; Liu et al., 2023). Similar to other species, *Orco* KO *P. brassicae* butterflies exhibited

a loss of electrophysiological responses to most plant volatiles we tested (Fan et al., 2022; Sun et al.,

2020) (Fig 2A and 2B), and the mating behavior of *P. brassicae Orco* KO butterflies was also disrupted

(Fig 2C). However, we observed some differences: *P. brassicae Orco* KO butterflies were still able to

mate to a limited degree (Fig 2D). Whereas *Orco* KO adults in some other insect species did not exhibit

any mating behavior (Fan et al., 2022; Fandino et al., 2019). This finding suggests that additional sensory

systems beyond olfaction, such as vision, may be involved in the mating processes of this day-active butterfly (Carpenter & Sparks, 1982; Obara et al., 2008). Interestingly, while many *P. brassicae* butterflies with impaired olfaction failed to mate, several KO butterflies still mate and deposit fertilized eggs when the oviposition substrate was supplied in our laboratory setting. The mated *Orco* KO butterflies laid a comparable number of fertilized eggs on cabbage plants to WT butterflies which is in line with the results from different moth species (Fan et al., 2022; Fandino et al., 2019; Sun et al., 2023), suggesting oviposition decisions are mostly guided by non-volatile compounds, when no host-plant location is required (van Loon et al., 1992a). Furthermore, the hatching rate of eggs laid by mated *Orco* KO butterflies was similar to that of WT butterflies (S1 Fig), indicating that the reduced fecundity in these insects is primarily due to impaired mating behavior and reduced mating frequency, which may result from a lack of pheromonal communication.

When directly placed on their host-plant, we found that WT caterpillars gained more weight than *Orco* KO caterpillars (Fig 3A). We speculate that olfaction is needed to evaluate plant tissues and to select different leaves or leaf parts in order to feed efficiently, even when the caterpillar is already on the plant. Secondary metabolites are often unevenly distributed even within a single leaf, and these concentration differences have been shown to influence the performance of different caterpillars (Kester et al., 2002; Yuan et al., 2022). It is conceivable that these localized differences in plant secondary metabolites are also detectable by the herbivore through its olfactory system (Hanson & Dethier, 1972). The difference between WT and KO caterpillars was amplified in the presence of disarmed parasitoids, which were able to attack but not injure the caterpillars, indicating that KO caterpillars were more susceptible to harassment by their natural enemies (Fig 3E). *Pieris brassicae* caterpillars are most vulnerable to *C. glomerata* during the first two larval instars. Due to their reduced growth rate, KO caterpillars spent a longer time in these vulnerable stages and, therefore, in a natural setting would likely have suffered more strongly from parasitoid attack than WT caterpillars (Brodeur et al., 1996; Haverkamp & Smid, 2020). Similarly, when the caterpillars were exposed to unmanipulated parasitoids, the KO caterpillars were more susceptible and exhibited a higher mortality rate, demonstrating again that olfaction is crucial in larvae to survive under the pressure of natural enemies (Fig 3F). Subsequently, we hypothesized that the differences in weight gain and survival rate between WT and KO caterpillars are due to different sensitivity to plant volatiles. By locating an available food source efficiently, WT caterpillars can sooner reach a body size in which they are no longer vulnerable to attack by *C. glomerata*

(Brodeur et al., 1996). The higher mortality rate of KO caterpillars is, therefore, at least partly, a result of a reduced foraging efficiency caused by the lack of olfactory information, which in turn led to an extended period in which the caterpillars were vulnerable to attack by their natural enemies.

Searching for host-plants suitable for feeding and free of natural enemies is largely accomplished via the olfactory system of herbivorous insects (Carrasco et al., 2015; Hu et al., 2020). Without functional ORs, *P. brassicae* caterpillars could still discriminate plant tissues from paper discs; however, *Orco* KO caterpillars preferred the non-host tomato plant over their natural food plant (Fig 3B). We surmise that OR-mediated olfaction is required in caterpillars to locate and identify their host-plants, and losing this part of their olfactory receptor repertoire would make it challenging to distinguish host from non-host-plants (Liu et al., 2023). Nonetheless, *Orco* KO caterpillars still retain ORNs expressing ionotropic receptors, which commonly detect acids and amines that have been found to elicit aversion in insect herbivores (Zhang et al., 2019). Fatty acids are present at the edge of both artificially damaged and caterpillar-infested cabbage leaves (Horikoshi et al., 1997), which, in the absence of ORs detecting attractive plant compounds, may have caused avoidance behavior in the KO caterpillar to their natural host-plant. Interestingly, the parasitoid *C. glomerata* also uses the fatty acids produced by the plant to locate *Pieris* caterpillars, and it has been argued that herbivores might avoid these compounds to escape from their natural enemies (Horikoshi et al., 1997; Zhang et al., 2019).

We found that WT caterpillars prefer caterpillar-infested plants free of wasps over caterpillar-infested plants with wasps, which suggests that caterpillars are able to detect certain volatiles that are derived from the interaction with caterpillars and wasps. By analyzing headspace samples of caterpillars, parasitoids as well as the interaction of caterpillars and parasitoids, we successfully identified some chemical compounds which were already known plant volatiles, such as alcohols and aldehydes. However, we did not find any compounds that were specifically abundant in Pb-Cg or Cg. This suggests that a volatile signal serving as a "danger signal" directly emitted by the wasps or caterpillars might not exist, or it might not be detectable with these methods (Fig 4B). Alternatively, indirect cues such as the volatiles emitted from the caterpillar spit during their defensive behavior might have triggered the avoidance behavior of the WT caterpillars to infested plants with natural enemies. When we tested the most abundant compounds in a two-choice behavioral assay against a solvent control, we found that *P. brassicae* WT larvae showed a preference to the plant volatiles (*Z*)-3-hexen-1-ol and (*E*)-2-hexenal (Fig 4D), which are known HIPVs. Previous studies on other species have demonstrated that caterpillars are

capable of detecting different HIPVs (de Fouchier et al., 2018; Di et al., 2017). For caterpillars, HIPVs can serve as indicators for the location of potential food plants and the presence of conspecifics (Di et al., 2017; Zhang et al., 2019), but they also attract natural enemies and enhance the risk of being attacked (Bernays, 1997; Ngumbi & Fadamiro, 2012; Yang et al., 2016). Most HIPVs identified in this study were found abundantly in caterpillar saliva that caterpillars spit onto parasitoids to defend against attack. In our study, caterpillars were attracted by HIPVs emitted by plants and by individual HIPVs (Fig 4D). We speculate that caterpillars use low to intermediate concentrations of HIPVs to find suitable host-plants and high concentrations to avoid sites where other caterpillars are under attack by their natural enemies. Therefore, perceiving plant leaf volatiles and approaching the food plants accordingly will increase the herbivores' feeding efficiency and enhance survival chances. In our study volatile cues of wasps were only found in very low quantities and these compounds hardly elicited any behavioral response in the caterpillars, however these chemicals might still play a role in the interaction between the caterpillars and wasps at a close range (Ebrahim et al., 2015).

Larval insects are under a strong selection pressure to the detect optimal food sources and enemy free spaces. Our study use a CRISPR/Cas9 knockout mutant line to investigate the significance of olfactory information for the survival and development of *P. brassicae* caterpillars. We provided novel insights into the tri-trophic interactions of plants, herbivores and their natural enemies from the perspective of a caterpillar and highlighted the role of olfaction for both foraging and escaping natural enemies.

Acknowledgements

We appreciate Janneke Bloem for helping with genome editing, and the rearing team at the Laboratory of Entomology for providing insects for our experiments in this study.

Author contributions

Conceptualization, Q.W., M.D. and A.H.; Methodology, Q.W., M.J., A.H.; Investigation, Q.W., Y.J.,

355 H.S., B.T.W., L.O.G.; Writing - Original Draf, Q.W., A.H.; Writing - Review & Editing, Q.W., L.O.G.,

356 M.D., A.H.; Funding Acquisition, M.D., A.H.; Resources: Q.W., M.D., A.H.; Supervision, M.D. and A.H.

Declaration of interests

358 The authors declare they have no competing interests.

Data availability

360 All data sets and scripts related to this study are available at the 4TU.ResearchData repository:

361 https://doi.org/10.4121/610e707d-b10a-467e-8afe-ead477827914

Materials and methods

Plant and insect rearing

Cabbage plants (*Brassica oleracea* var. gemmifera cv. Cobelius; Brussels sprouts) and tomato plants (*Solanum lycopersicum* cv. moneymaker) were planted in individual pots and grown for four weeks after germination before being used in the experiments or to feed the caterpillars. The cabbage plants were reared under glasshouse conditions at 22 ± 3 °C, RH 50-80 % and 16 h light: 8 h dark cycles. *Pieris brassicae* caterpillars and butterflies as well as *C. glomerata* wasps were collected from laboratory colonies maintained at the Laboratory of Entomology, Wageningen University & Research, the Netherlands. The caterpillars were reared on Brussels sprouts plants at 22 ± 3 °C, RH 25-35 % with 14 h light: 10 h dark cycles. The adults were supplied with 10 % sugar water as nutrient under the same environmental conditions. After successful gene editing, KO insects were transferred and reared in a glass compartment under conditions at 25 ± 3 °C, RH 25-35 %. Several KO caterpillars and butterflies from each generation were randomly collected for genotype screening to ensure that homozygosity remains. *Cotesia glomerata* adults were fed with organic honey, *P. brassicae* second-instar larvae (L2) were supplied to the parasitoid wasps for parasitization to maintain the colony. The wasp cocoons were collected in Petri dishes and incubated at 22 °C, RH 50-70 % with the same photoperiod as that of caterpillars. All the subsequent comparisons between WT and KO in this study were conducted in the glass compartment under the rearing condition of KO insects unless specified.

Orco knockout

The second exon of *Orco* was targeted to knock out the gene. The sgRNA was designed with Geneious Prime 2021 (Geneious, New Zealand) by searching N20 target + NGG PAM sequences in the second exon, sgRNA primers were selected according to activity scorings (Doench et al., 2016). The designated sgRNA off-target regions were evaluated by searching against the genome by est2genome function with a loose 50-score cut-off of using Exonerate 2.0 (Slater & Birney, 2005) and CHOPCHOP v3 (Labun et al., 2019). The sgRNAs that had no off-target sites in coding areas were selected. sgRNA templates were prepared by mixtures of 50 µL Q5 high-fidelity 2 × master mix (NEB, USA), 5 µL of the

following   forward   primers:

sgRNA1: ATTTAGGTGACACTATACATGTCAACTCTACTTAGGAGTTTTAGAGCTAGAAATA

GCAAG; sgRNA2: ATTTAGGTGACACTATACGATGAAGTAAACGAACTCAGTTTTAGAGCTA

GAAATAGCAAG; 5 µL constant reverse primer: AAAAGCACCGACTCGGTGCCACTTTTTCA

AGTTGATAACGGACTAGCCTTATTTTAACTTGCTATTTCTAGCTCTAAAAC and 40 µL H_2O. The

mixture was incubated at 98 °C for 2 min; followed by 38 cycles of 98 °C for 20 s, 65 °C for 10 s and

72 °C for 10 s; finalized at 72 °C for 5 min. The PCR products were checked by 1.0 % gel electrophoresis

and purified by QIAGEN PCR purification kit (QIAGEN, Germany) according to the manufacturer's

instructions. sgRNA was synthesized by a reaction system of 2 µL 10 × SP6 enzyme mix (Invitrogen,

USA), 2 µL 10 × SP6 reaction buffer, 2 µL each of ATP, UTP, GTP and CTP, 200 ug sgRNA template

and add RNase-free H_2O up to 20 µL. The mixture was incubated at 37 °C for 4 hours, followed by

adding 1 µL Turbo DNase and incubating at 37 °C for 15 min. The reaction products were purified by

Monarch RNA cleanup kit (NEB, USA). The concentration of sgRNA was determined by DeNovix

(DeNovix, USA). sgRNA was mixed with Cas9 protein (NEB, USA) and incubated at 25 °C for 10 min.

Newly laid eggs, not older than half an hour, were collected from our laboratory colony and were

fixed on glass slides. The sgRNA/Cas9 mixture was colored with 1 µL food dye to stain the eggs as an

indicator that injection with a Femo-jet (Eppendorf, Germany) was successful. The injected eggs were

incubated at 25 °C for 4 to 5 days until hatching. The caterpillars were reared until the emergence of the

butterflies, a leg of each butterfly was cut for genotyping. gDNA of the leg samples was extracted with

MyTaq Extract-PCR Kit (Bioline, Germany). Mutant screening was performed by PCR with 15 µL 2 ×

MyTaq HS Red mix, 2 µL forward primer (TCTGGCTTCGGTATTACATTTC), 2 µL reverse primer

(CTTTTATGGCGTGTTTTATTTG), 2 µL gDNA template and 9 µL H_2O. The PCR products were

checked by gel electrophoresis and sequenced with the same primers by Eurofins (Eurofins, the

Netherlands).

ORN staining and axon tracing

414   L3 caterpillars (n = 6 for WT and n = 5 for KO) were anesthetized on ice and decapitated. Heads

415 were cut along the midline with micro-scissors to facilitate penetration of the fixative and antibodies.

416 The collected heads were immersed in freshly prepared 4 % formaldehyde in 0.1 M phosphate buffer at

417 pH 7.3 and fixed overnight at 4 °C. The fixed samples were rinsed in 70 % ethanol, followed by

incubation in a mixture of 96 % ethanol: 30 % hydrogen peroxide at 1:1 ratio for 7 days at 4 °C. The

incubated samples were then dehydrated in a graded series of ethanol, degreased in xylene for two mins,

rehydrated and incubated in 0.1 M phosphate buffered saline (Oxoid, Dulbecco A) with 0.5% triton X-

100 (PBS-T) followed by four washes in PBS-T for 15 min each. Preincubation of head samples was

performed in 10 % normal goat serum (Bio-connect services, The Netherlands) in PBS-T (PBS-T-NGS)

for 1 hour, and then incubated in rabbit anti-Orco antibody which specificity was tested previously (Nolte

et al., 2016) diluted in 1: 500 in PBS-T-NGS for three days at 4 °C, followed by 6 washes in PBS-T for

30 min each. Samples were subsequently incubated in goat anti-rabbit conjugated to Alexa fluor 488

(Thermo Fisher Scientific, USA) diluted 1:100 and TO-PRO-3 iodide (Thermo Fisher Scientific, USA)

diluted 1:1000 in PBS-T-NGS for two days at 4 °C. Heads were washed two times in PBS-T for 30 mins

and once overnight at 4 °C, following with two 1-hour PBS washes at room temperature. The samples

were then dehydrated with a graded series of ethanol, cleared in xylene and mounted in DPX. Confocal

microscopy was performed with a Leica Stellaris 5- DIM 8 confocal microscope (Leica, Germany), using

a 63× oil-immersion plan APO objective NA1.4. and a spectrally flexible white light laser for excitation

of the two fluorophores using the pre-sets for Alexa fluor 488 and To-Pro-3 iodide.

L4 caterpillars of similar head size were anesthetized on ice and immobilized in clay. The tip of each

antenna was peeled, and a layer of petroleum jelly (Vaseline) was applied around the antenna. A 0.5 μL

drop of 2.5 % biotin-dextran solution was added to the cut and covered with petroleum jelly, then

incubated at 4 °C overnight. Brains were dissected in PBS, fixed in 4 % formaldehyde in PBS at 4 °C

overnight, dehydrated in xylene and rehydrated in PBS-T, followed by three washes in PBS-T for 2 hours

each. Brains were then incubated in 10 % NGS in PBS-T for 1 hour, followed by incubation in 1:200

streptavidin 488 and 1:1000 To-Pro iodide in PBS-T at 4 °C for three days. After further rinsing in PBS-

T overnight, brains were mounted in DPX. Whole-mount samples were scanned using confocal

microscopy with settings similar to those used for ORN staining.

Electroantennography (EAG)

A panel of 54 chemicals (S1 Table) including esters, alcohols, aldehydes, alkenes, isothiocyanates

(ITCs), heterocyclics and nitriles were selected from the literature (Bourne et al., 2023; Liu et al., 2020;

van Loon et al., 1992b; Zhu et al., 2015) and our current volatile identification data, to investigate the

EAG responses of both KO and wildtype (WT) butterfly antennae (n =15-18) of both sexes. Filter papers

loaded with 10 µL of selected chemicals, diluted in paraffin oil (10^{-2} v/v), were inserted into Pasteur

pipettes, clean filter paper was employed as negative control. The left antennae of butterflies were excised

at the basal end and the distal tip was also removed for better conductivity. Antennae were placed into

glass capillaries filled with EAG Ringer solution. Glass capillaries were placed on Ag-AgCl wires

connected to a ground electrode and a 10× high-impedance DC amplifier (Ockenfels Syntech, Germany).

The electrical signals were transformed using an IDAC-232 (Ockenfels Syntech, Germany) analog-

digital converter connected to a personal computer. Signals were finally recorded using the software

EAG Pro (Ockenfels Syntech, Germany). The first puff of each chemical was abandoned before the first

round of testing. Antennae were exposed to the chemicals in random order and were given intervals of

at least 20 s for recovering until the baseline stabilized.

Female oviposition behavior

A mating pair of newly emerged butterflies (n = 11 for WT and n = 14 for KO) were put in a cage

and supplied with a four-week-old Brussels sprouts cabbage plant as oviposition substrate and sugar

water as nutrients. The numbers of eggs were counted on a daily basis for seven days. The female

butterflies were then collected and dissected to compare mating frequency between WT and KO females.

Mating frequency was determined by the number of spermatophores dissected from the females'

abdomens. The eggs remained on the plants for another four to five days to allow hatching. The number

of caterpillars hatched enabled us to compare the number of fertilized eggs and hatching rate between

the two butterfly genotypes.

Caterpillar performance on plants

Caterpillar performance on cabbage plants was evaluated by three different treatments:

(1) For each genotype, a group of ten L1 caterpillars (n = 17) was placed on one leaf of a four-week-

old cabbage plant. Caterpillars were reared for ten days, and more plants were provided when needed.

The caterpillars were then collected for weighing to compare the growth between the two genotypes.

(2) The growth of both genotypes of caterpillars (n = 8) was further compared by rearing in the

presence of *C. glomerata* parasitoid wasps from which the ovipositor had been removed (hereafter:

"disarmed"). Groups of ten L1 caterpillars were placed on individual four-week-old cabbage plants as

described above. The parasitoid wasps were anesthetized on a CO_2 plate and the ovipositor was removed with fine tweezers to ensure that the parasitoid wasps can interact with the caterpillars but cannot parasitize them. Two disarmed parasitoid wasps were released in each rearing cage 24 hours after the caterpillars had been placed on the plants. The disarmed parasitoid wasps were replaced by newly disarmed parasitoids every day, the caterpillars were reared on the plants for ten days and then collected for weighing.

(3) To compare the survival rate of caterpillars (n = 7 groups of ten caterpillars) on the plants under the threat of *C. glomerata* parasitoid wasps, a group of ten L1 caterpillars was placed on a leaf of a four-week-old cabbage plant two days prior to placing two female parasitoid wasps. The parasitoid wasps were replaced every day to ensure that caterpillars were always exposed to natural enemies. The caterpillars were collected after ten days to count the number that survived; the collected caterpillars were further reared until they pupated or *C. glomerata* larvae emerged.

Larval host-plant seeking behavior

Caterpillar (n = 54-73) food-plant seeking behavior was evaluated in a greenhouse under the same conditions in which the caterpillars were reared. Plant leaf discs or green paper discs (Trophée 1224 "Forest Green", Clairefontaine, France), with 12 mm diameter were placed at opposite positions along the edge of a 188 mm-diameter Petri dish. A single L3 caterpillar was then placed at the center of the arena. Each caterpillar was observed for 10 min, the first choice that the caterpillars made and the time spent were recorded. The caterpillars were provided with a choice between either (i) a cabbage leaf disc and a paper disc, (ii) a tomato leaf disc and a paper disc, or (iii) a cabbage leaf disc and a tomato leaf disc. A choice was recorded when caterpillars began feeding on the cabbage leaf disc or contacted the tomato leaf / green paper discs. New discs were used for every caterpillar and the positions of discs were exchanged after every ten tests.

To further test the host-plant locating behavior of L3 caterpillars, we investigated the behaviors of caterpillars in a Y-tube olfactometer (diameter 35 mm) with a built-in black Y-shaped wire, allowing caterpillars to crawl along the wire and make a choice smoothly in the laboratory. Each arm was provided with a 1.1 L/min inlet airflow which had been purified by charcoal and humidified by water before entering the treatment jars. Healthy plants were untreated, infested plants were infested with 30 early L3 caterpillars for 24 hours and with caterpillars on plants while testing. Differently treated plants were

503   placed in jars to test caterpillar behaviors. The caterpillars had been starved for approximately 5 hours

504   before testing. Each caterpillar was tested and observed for 5 min, caterpillar behavior was monitored by

505   a camera. The position of inlet airflow was changed every five tests to avoid position bias.

Volatile analysis

Volatiles were collected from L3 *P. brassicae* caterpillars, three-day-old female *C. glomerata* wasps,

P. brassicae caterpillars and *C. glomerata* wasps together, caterpillar frass collected from L3 caterpillars

and caterpillar saliva also collected from L3 caterpillars using glass capillaries following Mattiacci et al.

(Mattiacci et al., 1994). Fresh frass and saliva samples were stored in a freezer at -20 °C and thawed

overnight at room temperature before volatile collection. Headspace collection of volatiles was

performed from five treatments: 20 female wasps (Cg) in a 10 mL glass vial (n =12); 20 L3 caterpillars

(Pb) in a 10 mL glass vial (n =10); 20 L3 caterpillars and 2 female wasps (Pb-Cg) in a 10 mL glass vial

(n =14); 100 mg caterpillar frass (Pb-Fr) in a 1.5 mL glass vial (n =10); and 50 µL caterpillar saliva (Pb-

S) in a 1.5 mL glass vial (n =12). Clean empty glass vials (10 mL and 1.5 mL) were used as a negative

control. Volatiles were collected through dynamic headspace sampling using Tenax TA adsorbent

material (20/35 mesh; Camsco, USA). Synthetic air (Air Synthetic 4.0 Monitoring from Linde Gas, the

Netherlands) at 110 mL min^{-1} to the samples in 10 mL vials and 55 mL min^{-1} to the samples in 1.5 mL

vials were constantly supplied as a carrier for the volatiles, while simultaneously the volatiles were

trapped by drawing air at 100 and 50 mL min^{-1}, respectively, through a stainless-steel tube filled with

200 mg Tenax TA for 2 h.

The collected volatiles were thermally released from the Tenax TA adsorbent using an Ultra 50:50

thermal desorption unit (Markes, UK) at 250 °C for 10 min under a helium flow of 20 mL min^{-1}, while

simultaneously re-collecting the volatiles in a thermally cooled universal solvent trap: Unity (Markes,

UK) at 0 °C. Once the desorption process was completed, the volatile compounds were released from

the cold trap by ballistic heating at 40°C s^{-1} to 280 °C, which was then kept for 10 min, while all the

volatiles were transferred to a ZB-5MS analytical column (30 mL × 0.25 mm ID × 1 mm F.T. with 10 m

built-in guard column (Phenomenex, USA), placed inside the oven of a Thermo Trace GC Ultra (Thermo

Fisher Scientific, USA), for separation of volatiles. The GC oven temperature was initially held at 40 °C

for 2 min and was immediately raised at 6 °C min^{-1} to a final temperature of 280°C, where it was kept

for 4 min under a constant helium flow of 1 mL min^{-1}. For the detection of volatiles, a Thermo Trace

DSQ quadrupole mass spectrometer (Thermo Fisher Scientific, USA) coupled to the GC was operated

in an electron impact ionization (EI) mode at 70 eV in a fullscan mode with a mass range of 35 - 400 amu

at 4.70 scans s^{-1}. The MS transfer line and ion source were set at 275 and 250 °C, respectively.

Automated baseline correction, peak selection (S/N > 3) and alignments of all extracted mass signals

of the raw data were processed following an untargeted metabolomic workflow using MetAlign software,

producing detailed information on the relative abundance of mass signals representing the available

metabolites (Lommen, 2009). This was followed by the reconstruction of the extracted mass features into

potential compounds using the MSClust software through data reduction by means of unsupervised

clustering and extraction of putative metabolite mass spectra (Tikunov et al., 2012). Tentative

identification of volatile metabolites was based on a comparison of the reconstructed mass spectra with

those in the NIST 2008 and Wageningen Mass Spectral Database of Natural Products MS libraries, as

well as experimentally obtained linear retention indices (LRIs).

Caterpillar response to individual chemical compounds

L3 caterpillars (n = 31-50 for each compound, each genotype) were tested in a custom-made set-up

consisting of 25 separate arenas in the laboratory. Single caterpillars were placed in the central well of

the individual arenas before the test. Caterpillars were given a choice between a selected volatile

chemical of interest (S2 Table) and paraffin oil. Input odors were delivered at 1.1 L min^{-1} for both

chemicals and solvent controls, the odors were sucked out at 2.2 L min^{-1} at the same time from the center.

The arena was cleaned by ethanol and dried by high-pressure air. The position of chemicals was

exchanged after each test. The behavior of caterpillars was recorded for 10 min at 100 frames per minute

using a Canon TV lens JF16mm (Canon, Japan). Behavioral data was extracted by EthoVision XT 11.5

software (Noldus, The Netherlands). Data were removed from the dataset when the caterpillar stayed in

the arena for less than 10 seconds. All tests were performed at room temperature.

Statistical analysis

556   Unless stated otherwise, statistical tests were performed in R (version 4.4.0; R Core Team, 2016) in

557 combination with RStudio (Boston, USA). Data sets were tested for normal distribution using a Shapiro–

558 Wilk test and for equal variances using Levene's test. The difference of glomerular number between

wildtype and *Orco* knockout caterpillars was tested with a Wilcoxon rank-sum test. Electroantennographic responses between the two genotypes were tested statistically by using the Student's t-test when the values were normally distributed and had homogenous variance or tested by using the Kruskal-Wallis rank-sum test when the requirements for the t-test were not met. The difference in butterfly mating frequency was analyzed by using the Wilcoxon rank-sum test; differences in oviposition dynamics were tested with a generalized linear model (GLM) with a negative binominal distribution. The development differences between WT and KO caterpillars on full plants and under the thread of disarmed parasitoids were tested using a Student's t-test, and survival rate difference between the two genotypes was tested using a GLM with a beta-binomial distribution. Caterpillar host-plant selection behaviors in Petri dishes and in the Y-tube were analyzed with a Chi-square test. Differences of the relative time spent by the caterpillars between the control side and the odor side of the multichannel arenas were compared by a non-parametric Wilcoxon rank-sum test. Principal component analysis (PCA) of volatile blends was achieved by MetaboAnalyst (Pang et al., 2024). Data were normalized by logarithmic transformation and scaled in auto mode. Ward's method and Euclidean distance were employed in the hierarchical clustering heatmap by default settings without clustering samples. In addition, the abundance of volatiles as peak heights was imported into SIMCA-P 17 statistical software (Umetrics, Sweden), the data was analyzed by multivariate data analysis (MVDA). Supervised orthogonal partial least-squares discriminant analysis (OPLS-DA) was employed to compare and correlate treatment groups.

References

Bernays, E. A. (1997). Feeding by lepidopteran larvae is dangerous. *Ecological Entomology*, *22*(1), 121-123. https://doi.org/10.1046/j.1365-2311.1997.00042.x

Bourne, M. E., Gloder, G., Weldegergis, B. T., Slingerland, M., Ceribelli, A., Crauwels, S., Lievens, B., Jacquemyn, H., Dicke, M., & Poelman, E. H. (2023). Parasitism causes changes in caterpillar odours and associated bacterial communities with consequences for host-location by a hyperparasitoid. *PLoS Pathogens*, *19*(3), e1011262. https://doi.org/10.1371/journal.ppat.1011262

Brodeur, J., Geervliet, J. B. F., & Vet, L. E. M. (1996). The role of host species, age and defensive behaviour on ovipositional decisions in a solitary specialist and gregarious generalist parasitoid (Cotesia species). *Entomologia Experimentalis et Applicata*, *81*(2), 125-132. https://doi.org/https://doi.org/10.1111/j.1570-7458.1996.tb02024.x

Carpenter, J. E., & Sparks, A. N. (1982). Effects of vision on mating behavior of the male corn earworm. *Journal of Economic Entomology*, *75*(2), 248-250.

Carrasco, D., Larsson, M. C., & Anderson, P. (2015). Insect host plant selection in complex environments. *Current opinion in insect science*, *8*, 1-7. https://doi.org/10.1016/j.cois.2015.01.014

Chen, Z., Traniello, I. M., Rana, S., Cash-Ahmed, A. C., Sankey, A. L., Yang, C., & Robinson, G. E. (2021). Neurodevelopmental and transcriptomic effects of CRISPR/Cas9-induced somatic *orco* mutation in honey bees. *Journal of Neurogenetics*, *35*(3), 320-332. https://doi.org/10.1080/01677063.2021.1887173

de Fouchier, A., Sun, X., Caballero-Vidal, G., Travaillard, S., Jacquin-Joly, E., & Montagne, N. (2018). Behavioral effect of plant volatiles binding to *Spodoptera littoralis* larval odorant receptors. *Frontiers in Behavioral Neuroscience*, *12*, 264. https://doi.org/10.3389/fnbeh.2018.00264

Di, C., Ning, C., Huang, L. Q., & Wang, C. Z. (2017). Design of larval chemical attractants based on odorant response spectra of odorant receptors in the cotton bollworm. *Insect Biochemistry and Molecular Biology*, *84*, 48-62. https://doi.org/10.1016/j.ibmb.2017.03.007

Dicke, M., & Baldwin, I. T. (2010). The evolutionary context for herbivore-induced plant volatiles: beyond the 'cry for help'. *Trends in Plant Science*, *15*(3), 167-175. https://doi.org/10.1016/j.tplants.2009.12.002

Dicke, M., & Grostal, P. (2001). Chemical detection of natural enemies by arthropods: an ecological perspective. *Annual Review of Ecology and Systematics*, *32*, 1-23. https://doi.org/10.1146/annurev.ecolsys.32.081501.113951

Doench, J. G., Fusi, N., Sullender, M., Hegde, M., Vaimberg, E. W., Donovan, K. F., Smith, I., Tothova, Z., Wilen, C., Orchard, R., Virgin, H. W., Listgarten, J., & Root, D. E. (2016). Optimized

sgRNA design to maximize activity and minimize off-target effects of CRISPR-Cas9. *Nature*

Biotechnology, *34*(2), 184-191. https://doi.org/10.1038/nbt.3437

Dweck, H. K., Ebrahim, S. A., Kromann, S., Bown, D., Hillbur, Y., Sachse, S., Hansson, B. S., &

Stensmyr, M. C. (2013). Olfactory preference for egg laying on citrus substrates in

Drosophila. *Current Biology*, *23*(24), 2472-2480. https://doi.org/10.1016/j.cub.2013.10.047

Ebrahim, S. A., Dweck, H. K., Stokl, J., Hofferberth, J. E., Trona, F., Weniger, K., Rybak, J., Seki, Y.,

Stensmyr, M. C., Sachse, S., Hansson, B. S., & Knaden, M. (2015). *Drosophila* avoids

parasitoids by sensing their semiochemicals via a dedicated olfactory circuit. *PLoS Biology*,

13(12), e1002318. https://doi.org/10.1371/journal.pbio.1002318

Fan, X. B., Mo, B. T., Li, G. C., Huang, L. Q., Guo, H., Gong, X. L., & Wang, C. Z. (2022).

Mutagenesis of the odorant receptor co-receptor (*Orco*) reveals severe olfactory defects in the

crop pest moth *Helicoverpa armigera*. *BMC Biology*, *20*(1), 214.

https://doi.org/10.1186/s12915-022-01411-2

Fandino, R. A., Haverkamp, A., Bisch-Knaden, S., Zhang, J., Bucks, S., Nguyen, T. A. T., Schroder, K.,

Werckenthin, A., Rybak, J., Stengl, M., Knaden, M., Hansson, B. S., & Große-Wilde, E.

(2019). Mutagenesis of odorant coreceptor *Orco* fully disrupts foraging but not oviposition

behaviors in the hawkmoth *Manduca sexta*. *Proceedings of the National Academy of Sciences*

of the United States of America, *116*(31), 15677-15685.

https://doi.org/10.1073/pnas.1902089116

Feltwell, J. (1982). *Large white butterfly: the biology, biochemistry and physiology of Pieris brassicae*

(Linnaeus). Dr W. Junk Publishers.

Hanson, F. E., & Dethier, V. G. (1972). Role of gustation and olfaction in food plant discrimination in

the tobacco hornworm, *Manduca sexta*. *Journal of Insect Physiology*, *19*(5), 1019-1031.

Hasan, F., & Ansari, M. S. (2010). Effect of different cole crops on the biological parameters of *Pieris*

brassicae (L.) (Lepidoptera: Pieridae) under laboratory conditions. *Journal of Crop Science*

and Biotechnology, *13*(3), 195-202. https://doi.org/10.1007/s12892-010-0025-2

Hasan, F., & Ansari, M. S. (2011). Population growth of *Pieris brassicae* (L.) (Lepidoptera: Pieridae)

on different cole crops under laboratory conditions. *Journal of Pest Science*, *84*(2), 179-186.

https://doi.org/10.1007/s10340-010-0339-9

Haverkamp, A., Hansson, B. S., & Knaden, M. (2018). Combinatorial codes and labeled lines: how

insects use olfactory cues to find and judge food, mates, and oviposition sites in complex

environments. *Frontiers in physiology*, *9*, 49. https://doi.org/10.3389/fphys.2018.00049

Haverkamp, A., & Smid, H. M. (2020). A neuronal arms race: the role of learning in parasitoid-host

interactions. *Current opinion in insect science*, *42*, 47-54.

https://doi.org/10.1016/j.cois.2020.09.003

Horikoshi, M., Takabayashi, J., Yano, S., Yamaoka, R., Ohsaki, N., & Sato, Y. (1997). *Cotesia glomerata* female wasps use fatty acids from plant–herbivore complex in host searching. *Journal of Chemical Ecology*, *23*, 1505-1515.

Hu, X., Su, S., Liu, Q., Jiao, Y., Peng, Y., Li, Y., & Turlings, T. C. (2020). Caterpillar-induced rice volatiles provide enemy-free space for the offspring of the brown planthopper. *eLife*, *9*, e55421. https://doi.org/10.7554/eLife.55421

Jones, W. D., Nguyen, T. A., Kloss, B., Lee, K. J., & Vosshall, L. B. (2005). Functional conservation of an insect odorant receptor gene across 250 million years of evolution. *Current Biology*, *15*(4), R119-121. https://doi.org/10.1016/j.cub.2005.02.007

Kannan, K., Galizia, C. G., & Nouvian, M. (2022). Olfactory strategies in the defensive behaviour of insects. *Insects*, *13*(5), 470. https://doi.org/10.3390/insects13050470

Kester, K. M., Peterson, S. C., Hanson, F., Jackson, D. M., & Severson, R. F. (2002). The roles of nicotine and natural enemies in determining larval feeding site distributions of *Manduca sexta* L. and *Manduca quinquemaculata* (Haworth) on tobacco. *Chemoecology*, *12*, 1-10.

Koutroumpa, F., Monsempes, C., Anton, S., Francois, M. C., Montagne, N., & Jacquin-Joly, E. (2022). Pheromone receptor knock-out affects pheromone detection and brain structure in a moth. *Biomolecules*, *12*(3), 341. https://doi.org/10.3390/biom12030341

Labun, K., Montague, T. G., Krause, M., Torres Cleuren, Y. N., Tjeldnes, H., & Valen, E. (2019). CHOPCHOP v3: expanding the CRISPR web toolbox beyond genome editing. *Nucleic Acids Research*, *47*(W1), W171-W174. https://doi.org/10.1093/nar/gkz365

Larsson, M. C., Domingos, A. I., Jones, W. D., Chiappe, M. E., Amrein, H., & Vosshall, L. B. (2004). *Or83b* encodes a broadly expressed odorant receptor essential for *Drosophila* olfaction. *Neuron*, *43*(5), 703-714. https://doi.org/10.1016/j.neuron.2004.08.019

Liu, X. L., Zhang, J., Yan, Q., Miao, C. L., Han, W. K., Hou, W., Yang, K., Hansson, B. S., Peng, Y. C., Guo, J. M., Xu, H., Wang, C. Z., Dong, S. L., & Knaden, M. (2020). The molecular basis of host selection in a crucifer-specialized moth. *Current Biology*, *30*(22), 4476-4482 e4475. https://doi.org/10.1016/j.cub.2020.08.047

Liu, X. Y., Yan, R., Chen, S. J., Zhang, J. L., & Xu, H. J. (2023). Orco mutagenesis causes deficiencies in olfactory sensitivity and fertility in the migratory brown planthopper, *Nilaparvata lugens*. *Pest Management Science*, *79*(3), 1030-1039. https://doi.org/10.1002/ps.7286

Lommen, A. (2009). MetAlign: interface-driven, versatile metabolomics tool for hyphenated full-scan mass spectrometry data preprocessing. *Analytical Chemistry*, *81*(8), 3079-3086.

Mattiacci, L., Dicke, M., & Posthumus, M. A. (1994). Induction of parasitoid attracting synomone in brussels sprouts plants by feeding of *Pieris brassicae* larvae: role of mechanical damage and herbivore elicitor. *Journal of Chemical Ecology*, *20*(9), 2229-2247.

Ngumbi, E., & Fadamiro, H. (2012). Species and sexual differences in behavioural responses of a specialist and generalist parasitoid species to host-related volatiles. *Bulletin of Entomological Research*, *102*(6), 710-718. https://doi.org/10.1017/S0007485312000326

Nolte, A., Gawalek, P., Koerte, S., Wei, H., Schumann, R., Werckenthin, A., Krieger, J., & Stengl, M. (2016). No evidence for ionotropic pheromone transduction in the hawkmoth *Manduca sexta*. *PLoS One*, *11*(11), e0166060. https://doi.org/10.1371/journal.pone.0166060

Obara, Y., Koshitaka, H., & Arikawa, K. (2008). Better mate in the shade: enhancement of male mating behaviour in the cabbage butterfly, *Pieris rapae crucivora*, in a UV-rich environment. *The Journal of Experimental Biology*, *211*(Pt 23), 3698-3702. https://doi.org/10.1242/jeb.021980

Oland, L. A., & Tolbert, L. P. (2011). Roles of glial cells in neural circuit formation: insights from research in insects. *Glia*, *59*(9), 1273-1295. https://doi.org/10.1002/glia.21096

Pang, Z., Lu, Y., Zhou, G., Hui, F., Xu, L., Viau, C., Spigelman, A. F., MacDonald, P. E., Wishart, D. S., Li, S., & Xia, J. (2024). MetaboAnalyst 6.0: towards a unified platform for metabolomics data processing, analysis and interpretation. *Nucleic Acids Research*, *52*(W1), W398-W406. https://doi.org/10.1093/nar/gkae253

Sato, K., Pellegrino, M., Nakagawa, T., Nakagawa, T., Vosshall, L. B., & Touhara, K. (2008). Insect olfactory receptors are heteromeric ligand-gated ion channels. *Nature*, *452*(7190), 1002-1006. https://doi.org/10.1038/nature06850

Slater, G. S., & Birney, E. (2005). Automated generation of heuristics for biological sequence comparison. *BMC Bioinformatics*, *6*, 31. https://doi.org/10.1186/1471-2105-6-31

Stam, J. M., Kroes, A., Li, Y., Gols, R., van Loon, J. J. A., Poelman, E. H., & Dicke, M. (2014). Plant interactions with multiple insect herbivores: from community to genes. *Annual Review of Plant Biology*, *65*, 689-713. https://doi.org/10.1146/annurev-arplant-050213-035937

Sun, H., Bu, L. A., Su, S. C., Guo, D., Gao, C. F., & Wu, S. F. (2023). Knockout of the odorant receptor co-receptor, orco, impairs feeding, mating and egg-laying behavior in the fall armyworm *Spodoptera frugiperda*. *Insect Biochemistry and Molecular Biology*, *152*, 103889. https://doi.org/10.1016/j.ibmb.2022.103889

Sun, H., Liu, F., Ye, Z., Baker, A., & Zwiebel, L. J. (2020). Mutagenesis of the orco odorant receptor co-receptor impairs olfactory function in the malaria vector *Anopheles coluzzii*. *Insect Biochemistry and Molecular Biology*, *127*, 103497. https://doi.org/10.1016/j.ibmb.2020.103497

Tikunov, Y. M., Laptenok, S., Hall, R. D., Bovy, A., & de Vos, R. C. (2012). MSClust: a tool for unsupervised mass spectra extraction of chromatography-mass spectrometry ion-wise aligned data. *Metabolomics*, *8*(4), 714-718. https://doi.org/10.1007/s11306-011-0368-2

Trible, W., Olivos-Cisneros, L., McKenzie, S. K., Saragosti, J., Chang, N. C., Matthews, B. J., Oxley, P. R., & Kronauer, D. J. C. (2017). orco Mutagenesis Causes Loss of Antennal Lobe Glomeruli

and Impaired Social Behavior in Ants. *Cell*, *170*(4), 727-735 e710. https://doi.org/10.1016/j.cell.2017.07.001

Turlings, T. C. J., & Erb, M. (2018). Tritrophic interactions mediated by herbivore-induced plant volatiles: mechanisms, ecological relevance, and application potential. *Annual Review of Entomology*, *63*, 433-452. https://doi.org/10.1146/annurev-ento-020117-043507

van Loon, J. J. A., Blaakmeer, A., Griepink, F. C., van Beek, T. A., Schoonhoven, L. M., & de Groot, A. (1992a). Leaf surface compound from *Brassica oleracea* (Cruciferae) induces oviposition by *Pieris brassicae* (Lepidoptera: Pieridae). *Chemoecology*, *3*, 39-44.

van Loon, J. J. A., Frentz, W. H., & van Eeuwijk, F. A. (1992b). Electroantennogram responses to plant volatiles in two species of *Pieris* butterflies. *Entomologia Experimentalis et Applicata*, *62*(3), 253-260.

Venugopal, U., Jayanthi, P. D. K., KUMAR, P., Jagadeesh, K. S., & Mohan, K. M. (2020). Behavioural response of specific larval endoparasitoid, *Apanteles machaeralis* (Wilkinson) to volatile cues from its host insect, *Diaphania indica* (Saunders) and the host plant (*Cucumis sativus* L.). *Journal of Biological Control*, *34*(2), 132-139. https://doi.org/10.18311/jbc/2020/25359

Vet, L. E., & Dicke, M. (1992). Ecology of infochemical use by natural enemies in a tritrophic context. *Annual Review of Entomology*, *37*(1), 141-172. https://doi.org/10.1146/annurev.en.37.010192.001041

Wang, Q. (2023). *Odorant receptors in larvae and adults of a butterfly species: from identification to ecological significance* (Publication Number 31005520) [Ph.D., Wageningen University and Research]. Netherlands. https://library.wur.nl/WebQuery/rduser/ezproxy-login?url=https://www.proquest.com/dissertations-theses/odorant-receptors-larvae-adults-butterfly-species/docview/3059425559/se-2?accountid=27871

Wang, Q., Dicke, M., & Haverkamp, A. (2023a). Sympatric *Pieris* butterfly species exhibit a high conservation of chemoreceptors. *Frontiers in Cellular Neuroscience*, *17*, 1155405. https://doi.org/10.3389/fncel.2023.1155405

Wang, Q., Smid, H. M., Dicke, M., & Haverkamp, A. (2023b). The olfactory system of *Pieris brassicae* caterpillars: from receptors to glomeruli. *Insect Science*. https://doi.org/10.1111/1744-7917.13304

Wicher, D., Schafer, R., Bauernfeind, R., Stensmyr, M. C., Heller, R., Heinemann, S. H., & Hansson, B. S. (2008). *Drosophila* odorant receptors are both ligand-gated and cyclic-nucleotide-activated cation channels. *Nature*, *452*(7190), 1007-1011. https://doi.org/10.1038/nature06861

Williams, A. T., Verhulst, E. C., & Haverkamp, A. (2022). A unique sense of smell: development and evolution of a sexually dimorphic antennal lobe – a review. *Entomologia Experimentalis et Applicata*, *170*(4), 303-318. https://doi.org/https://doi.org/10.1111/eea.13145

Yan, H., Opachaloemphan, C., Mancini, G., Yang, H., Gallitto, M., Mlejnek, J., Leibholz, A., Haight,
K., Ghaninia, M., Huo, L., Perry, M., Slone, J., Zhou, X., Traficante, M., Penick, C. A.,
Dolezal, K., Gokhale, K., Stevens, K., Fetter-Pruneda, I., . . . Desplan, C. (2017). An
Engineered orco Mutation Produces Aberrant Social Behavior and Defective Neural
Development in Ants. *Cell*, *170*(4), 736-747 e739. https://doi.org/10.1016/j.cell.2017.06.051

Yang, G., Zhang, Y. N., Gurr, G. M., Vasseur, L., & You, M. S. (2016). Electroantennogram and
behavioral responses of *Cotesia plutellae* to plant volatiles. *Insect Science*, *23*(2), 245-252.
https://doi.org/10.1111/1744-7917.12308

Yuan, G. G., Zhao, L. C., Du, Y. W., Yu, H., Shi, X. B., Chen, W. C., & Chen, G. (2022). Repellence or
attraction: secondary metabolites in pepper mediate attraction and defense against Spodoptera
litura. *Pest Management Science*, *78*(11), 4859-4870. https://doi.org/10.1002/ps.7107

Zhang, J., Bisch-Knaden, S., Fandino, R. A., Yan, S., Obiero, G. F., Große-Wilde, E., Hansson, B. S., &
Knaden, M. (2019). The olfactory coreceptor IR8a governs larval feces-mediated competition
avoidance in a hawkmoth. *Proceedings of the National Academy of Sciences of the United*
States of America, *116*(43), 21828-21833. https://doi.org/10.1073/pnas.1913485116

Zhu, F., Broekgaarden, C., Weldegergis, B. T., Harvey, J. A., Vosman, B., Dicke, M., & Poelman, E. H.
(2015). Parasitism overrides herbivore identity allowing hyperparasitoids to locate their
parasitoid host using herbivore-induced plant volatiles. *Molecular Ecology*, *24*(11), 2886-
2899. https://doi.org/10.1111/mec.13164

Figures and figure legends

Fig 1. *Orco* knockout by CRISPR/Cas9 and verification in *Pieris brassicae*. (A), gene structure of *Orco*. Yellow blocks indicate exons and blue block indicates target area in the second exon. Black solid line indicates intron. Black arrows indicate the designated cleavage site of the Cas9 protein. The target knockout area is magnified to show the sequence. Blue letters indicate base pairs in the second exon segment, purple letters indicate mutation sites. Protospacer adjacent motif (PAM) sequences are in blue bold. (B), predicted transmembrane structure of Orco. The left and right panels represent wildtype Orco and mutated Orco transmembrane domains, respectively. Orange blocks indicate ORN membrane, extracellular (Extra) and intracellular (Intra) are shown. White circles indicate amino acids of Orco, blue circles on the left panel indicate the target mutation area. Cyan circles and the brown circle on the right panel indicate the mutated area and the early stop codon, respectively. Orco-positive ORNs and larval antennal center. *P. brassicae* larval antennae and larval antennal center (LAC) staining. (C), Odorant receptor neurons (ORNs) were stained (green cells, several are indicated by white arrows) in wildtype (WT) larval antennae. (D), ORNs were not stained in *Orco* knockout (KO) larval antennae. (E), An ORN was stained (green cell, indicated by white arrow) in WT larval palps. (F), ORNs were not stained in KO

larval palps. (G), glomeruli in the LAC of a WT brain. (H), glomeruli in the LAC of a KO brain. Glomeruli are indicated by white dashed circles in (G) and (H). (I), number of glomeruli counted in WT caterpillar brain (orange) and *Orco* KO caterpillar (blue) brain. Significant difference was detected by Wilcoxon rank-sum test.

Fig 2. *Pieris brassicae* butterfly electrophysiological responses to chemical compounds and mating and oviposition dynamics. (A), electroantennogram (EAG) response of male butterflies (n = 18 for both WT and KO). (B), EAG response of female butterflies (n = 15 for WT and n = 17 for KO). Left panels represent wildtype (WT) butterfly EAG responses, and right panels represent are knockout (KO) butterfly EAG responses. Significant differences between WT and KO butterflies were identified using Student's t-test when the data was normally distributed and the variances were equal or using a Kruskal-Wallis rank-sum test when these criteria did not apply. Significant difference levels are indicated by asterisks,

ns (P > 0.05), * (0.01 < P < 0.05), ** (0.001 < P < 0.01), *** (P < 0.001), - (no response recorded from

WT butterfly antennae). Butterfly antennal responses (mV) to the tested chemicals are indicated by a

color scale from navy (0 mV) via yellow to red (1.1 mV). (C), mating frequency of *P. brassicae* butterflies.

The boxplots indicate the number of spermatophores in WT female butterflies (n = 11) and in KO

butterflies (*Orco*^(-/-)) (n = 14), the difference was tested by using Wilcoxon rank-sum test. (D), *P. brassicae*

wildtype (WT) and *Orco*^(-/-) knockout butterfly oviposition dynamics. The orange line indicates the number

of eggs per female (mean ± SE) laid by wildtype butterflies (n = 11), the blue line indicates the number

of eggs laid by *Orco* knockout butterflies (n = 14). Differences were analyzed using a GLM with negative

binominal distribution, difference test results are shown in the line chart.

Fig 3. *Pieris brassicae* caterpillar growth and foraging behavior. (A), *P. brassicae* caterpillar growth on cabbage plants, y axis shows weight (g) of WT and KO caterpillars after 10 days of feeding , n = 17. (B), caterpillar behavioral choices in a Petri dish. The numbers of caterpillars that chose one of the two discs are shown in the respective bars (n = 54 -73). Schematic drawing shows behavioral setup. Petri dish diameter was 188 mm, disc diameters were 13 mm. Discs in the Petri dish represent cabbage, paper and tomato leaf discs. Grey bars indicate paper disc choices, light green indicates cabbage leaf disc choices and orange indicates choices for tomato leaf discs. (C), caterpillar behavioral choices in Y-tube olfactometer without parasitoid wasps. Schematic drawing shows the Y-tube olfactometer. The dashed line indicates a black metal Y wire in the center of glass Y-tube olfactometer. The main arm of the Y-olfactometer is 200 mm length, the lateral arms are 275 mm length, the angle between the lateral arms is 80 °. Light green, dark green and grey bars represent choices for healthy plant, infested plant and no plant respectively. Healthy plant, plants were not treated; no plant, an empty jar without any insect or plant; infested plant, plants were infested by early L3 caterpillars. (D), caterpillar behavioral choices in Y-tube olfactometer with parasitoid wasps. Different treatments are in different colors. Dark green, infested plant - wasps; magenta, infested plant + wasps. Significant differences were tested between wildtype (WT) and knockout (KO) caterpillars or between two discs by Chi-square test in panel B-D, P-values are shown on the right side of each bar. (E), *P. brassicae* caterpillar growth when exposed to disarmed *C. glomerata* female parasitoids, y axis shows weight (g) of caterpillars after 10 days of feeding, n = 8. Schematic drawing shows disarmed female *C. glomerata* (ovipositor removed). (F), *P. brassicae* caterpillar survival rate when exposed to healthy *C. glomerata* female wasps, n = 7. Significant differences in development and survival rate were assessed using a one-tailed Student's t-test, P values are indicated above boxplots. Schematic drawing shows a healthy female *C. glomerata* (unmanipulated). In both panels orange boxplots indicate wildtype caterpillars (WT), blue boxplots indicate *Orco* knockout caterpillars (KO).

Fig 4. Overview of caterpillar- (*Pieris brassicae*) and parasitoid wasp- (*Cotesia glomerata*) associated volatile compounds. (A), PCA (Principal Component Analysis) two-dimensional score plot of five treatment groups: Cg, *Cotesia glomerata* female parasitoid wasps (n = 12); Pb, *Pieris brassicae* caterpillars (n = 10); Pb-Cg, *P. brassicae* caterpillars in the presence of *C. glomerata* female parasitoid wasps (n = 14); Pb-Fr, *P. brassicae* caterpillar frass (n = 10); and Pb-S, *P. brassicae* spit (n = 12), based on their volatile blend composition. (B), hierarchical clustering heatmap showing the abundance of each identified volatile compound of each treatment. Clustering chemicals in the heatmap indicate higher correlation. (C), schematic drawing of the custom-designed multi-channel arena. (D) wildtype (WT)

caterpillar behavioral preference to the tested 15 odorants. (E), *Orco* knockout (KO) caterpillar

behavioral preference to the tested 15 chemicals. Unfilled bars represent tests that exhibit no difference

between paraffin oil and the odorant of interest. Blue bars indicate the cumulative duration ratio that

caterpillars stayed in the paraffin oil zone, dark orange bars indicate the cumulative duration ratio that

caterpillars stayed in the odorant zone. Relative time spent in odorant zone = Cumulative duration in a

specific zone / Cumulative duration in the arena. Error bars indicate standard errors. In (D) and (E),

differences were tested between the two zones using a Wilcoxon rank-sum test, P values are presented

above bars (n = 31-50). For each comparison, left bar represents paraffin oil and right bar represents

chemical compound.