Chemosynthesis enhances carbon fixation in an active microbialite ecosystem

Francesco Ricci[1,2,3], Pok Man Leung[1,2,*], Tess Hutchinson[1,2], Thanh Nguyen-Dinh[1,2], Ashleigh v.S. Hood[3], Vinícius W. Salazar[4], Vera Eate[5], Wei Wen Wong[5], Perran L.M. Cook[5], Chris Greening[1,2] & Harry McClelland[3,6,*]

1. Department of Microbiology, Biomedicine Discovery Institute, Monash University, Melbourne, VIC, Australia
2. Securing Antarctica's Environmental Future, Monash University, Melbourne, VIC, Australia
3. School of Geography, Earth and Atmospheric Sciences, University of Melbourne, Parkville, VIC, Australia
4. Melbourne Bioinformatics, University of Melbourne, Parkville, VIC, Australia
5. Water Studies Centre, School of Chemistry, Monash University, Clayton, VIC, Australia
6. Department of Structural and Molecular Biology, University College London, London, WC1E6BT, UK

*Corresponding authors: francesco.ricci@monash.edu, bob.leung@monash.edu, and h.mcclelland@ucl.ac.uk

Abstract

Microbialites—carbonate structures formed under the influence of microbial action—are the earliest macroscopic evidence of life. For three billion years, the microbial mat communities responsible for these structures fundamentally shaped Earth's biogeochemical cycles. In photosynthetic microbial communities, light energy ultimately drives primary production and the ensuing cascade of daisy-chained metabolisms. However, reduced compounds such as atmospheric trace gases and those released as metabolic byproducts in deeper, anaerobic regions of the mat, could also fuel chemosynthetic processes. Here, we investigated the intricate metabolic synergies that sustain microbialite community nutrient webs. We recovered 331 genomes spanning 40 bacterial and archaeal phyla, revealing a staggering diversity fuelled by the biogeochemistry of these ecosystems. While phototrophy is an important metabolism encoded by 17% of the genomes, over half

encode enzymes to harness energy from reduced compounds and 12% co-encode
carbon fixation pathways, using sulfide and hydrogen as major electron donors.
Consistent with these genomic predictions, we experimentally demonstrated that
microbialite communities oxidise ferrous iron, ammonia, sulfide and gas substrates
aerobically and anaerobically. Furthermore, ^{14}C-fixation assays revealed that
chemosynthesis contributes significantly to carbon fixation alongside photosynthesis.
Chemosynthesis in microbialite communities represents a complex interplay of
metabolic synergies and continuous nutrient cycling, which decouples community
carbon fixation from the diurnal cycle. As a result, this process mitigates the loss of
organic carbon from respiration, thus enhancing the net productivity of these highly
efficient ecosystems.
49
50
51
52

Introduction

Complex microbial communities in aquatic environments can form carbonate
structures named microbialites. With the earliest fossils dating to around three and a
half billion years[1,2], these structures are representative of the most ancient and
persistent microbial ecosystems in Earth's history. Over Earth's long history, dense,
benthic microbial communities played a pivotal role in shaping the composition of
Earth's atmosphere[3,4]. Notably, they contributed to the rise in atmospheric oxygen
following the evolution of oxygenic photosynthesis[3,5], which amplified global
biological productivity by 100-1000 fold[6]. Much of this new productivity first occured
in microbial mats that provided favourable conditions for photosynthetic communities
to thrive[7–9]. Additionally, mat communities mediated large fluxes of reduced gases
such as carbon monoxide (CO), hydrogen (H_2), and methane (CH_4) into the
atmosphere[6]. Today, living microbialites are generally confined to extreme
environments such as hypersaline lakes[10–14], where grazing metazoans and plants
are typically absent[3]. However, modern stratified microbial communities have a
structure resembling those from the Cambrian[15] and Archaean[5], positioning these
communities as natural laboratories for unravelling the functional processes
underlying these ancient systems.
71
Microbialites are hotspots of microbial diversity[16–18], yet the key processes that have
sustained the ecosystems associated with these structures remain enigmatic. Light
is thought to have had a central role throughout their history, initially sustaining
anoxygenic phototrophs and later cyanobacteria and algae, driving organic carbon
production, carbonate precipitation, and elemental cycling[5,6,19,20]. The resultant
photosynthetic end-products shape aerobic niches and fuel complex microbial
networks[3], including by supporting the activity of aerobic heterotrophs[21–23].
Anaerobes dominate beneath the surface of microbial mats, including fermenters,

sulfate-reducing bacteria (SRB), and methanogens that catabolise organic matter
and gases[6,24,25]. SRB use organic (e.g. acetate, sugars) and inorganic electron
donors (e.g. H_2) while releasing H_2S. Chemolithoautotrophs can exploit inorganic
compounds, encoding genes mediating the oxidation of H_2, CO, and various sulfur
compounds[26–28], though their ecological and biogeochemical roles within these
ecosystems remain poorly understood. While photosynthetic carbon fixation is
central to biomass production in these ecosystems[3,29], this is not the sole process
contributing to their primary production. Most notably, the abundant anoxic pockets
within microbialites foster ecological niches well-suited for anaerobic autotrophs
utilising the Wood–Ljungdahl pathway (WLP)[30–32]. Although gene- and genome-
resolved studies have provided insights into the functional potential of microbialite
communities, direct activity-based evidence of the complex metabolic interactions
that sustain these ecosystems remains limited.
93
Here we aim to shed light on the primary production and elementary cycling
processes sustaining the biodiversity of calcyfing microbial mats using living
microbialites from West Basin Lake, Victoria, Australia. While it is recognised that
microbialite communities have the potential for several photosynthetic and
chemosynthetic carbon fixation pathways, we still lack a comprehensive
understanding of their mediators, the role of various electron donors in driving these
processes, and their relative contribution to primary production. The classic view
positions photosynthesis as a leading process driving biological productivity[3,6,33,34].
However, microbialite communities produce substantial amounts of reduced
compounds such as H_2, H_2S, CO and CH_4[6,25], which could fuel diverse
chemosynthetic processes. Using living microbialite ecosystems, we integrate
genome-resolved metagenomics with high-resolution metabarcoding,
biogeochemistry, modelling and phylogenetic analysis to quantify the key
metabolisms and their contributions in supporting the nutrient web in microbialite
communities.
109
110

Results & Discussion

Taxonomically and metabolically diverse microbes control a complex nutrient web in living microbialites

115
116 We deeply sequenced nine metagenomes across three microbialite communities
117 from West Basin Lake in Victoria, Australia. This effort resulted in the recovery of
118 331 medium- to high-quality metagenome-assembled genomes (MAGs) dereplicated

at the species level using a 95% ANI threshold. These MAGs depict a remarkable diversity spanning 40 bacterial and archaeal phyla, including the first genomes from a microbialite community for the phyla JAHJDO01 and UBP6 (Fig. 1). Many of these microbes represent elusive and rare lineages, such as Cloacimonadota, Krumholzibacteriota, Hydrogenedentota, Omnitrophota, and Sumerlaeota. Representatives of these phyla have been previously found in extreme environments including an Antarctic lake[35], deep-sea trenches[36] and geothermal springs[37]. The most abundant phyla were Proteobacteria and Bacteroidota, represented by 81 and 56 MAGs respectively (Fig. 1). Cyanobacteria are considered key members of these ecosystems, though we only retrieved one medium-quality MAG of *Halothece* (total rel. abundance 3.11%) and 18 MAGs capable of anoxygenic chlorophototrophy (Fig. 1). Millimeter-scale 16S rRNA gene community profiling detected 30 cyanobacterial sequences, which were present at considerably different relative abundance between the analysed samples (sample C: 11.2% ± 7.6; sample D: 0.27% ± 0.35) (Supp. Data 1). Archaea are also significant members, represented by 18 MAGs spanning Asgardarchaeota, Halobacteriota, Iainarchaeota, Nanoarchaeota, Thermoplasmatota and Thermoproteota (Fig. 1). Eukaryotes have been found in most modern microbial communities[38–40] and accordingly we identified several major members of diatoms and red and green algae (Supp. Data 1).

Screening of key metabolic pathways in the MAGs revealed extensive metabolic diversity within West Basin Lake microbialite communities (Fig. 1), potentially underpinning these diverse microbial assemblages. Many microbes appeared to be facultative anaerobes that can input electrons from organic carbon to aerobic and anaerobic respiratory chains with 63% of genomes encoding at least one terminal oxidase (Supp. Data 2). The widespread capacity to oxidise various inorganic compounds (53.2% of the genomes), including hydrogen, sulfide, and carbon monoxide, for supplemental energy likely allows more efficient use of organic carbon for anabolism than catabolism (Supp. Data 2). Furthermore, our data suggest that at West Basin Lake, most organic carbon is produced *in situ*. We identified multiple lines of evidence supporting this hypothesis. First, the lake is not connected to any inlet and lacks aquatic vegetation. Second, we measured moderate concentrations of organic carbon in the water, averaging 4.83 mg/L. Notably, organic carbon abundance increases on average from 3.09 mg/L in shallow to 5.60 mg/L in deeper layers within the microbialites (Supp. Data 3). Third, 13.6% of the generated MAGs encode genes for carbon fixation (Fig. 1). Specifically, 19 MAGs encoded the WLP, whereas 15 MAGs co-encoded the CBB cycle with several other energy harvesting genes (Fig. 1; Supp. Data 2). Interestingly, some MAGs possess the genetic machinery to utilise multiple energy sources. For instance, a *Thiohalophilus* MAG encodes the CBB cycle with uptake [NiFe]-hydrogenases, sulfide quinone oxidoreductase, thiosulfohydrolase, reverse dissimilatory sulfate reductase and iron oxidising cytochrome (Fig. 1). We also found six MAGs encoding the reductive tricarboxylic acid cycle (rTCA), three of which belonged to the enigmatic group 3 [NiFe]-hydrogenase-encoding Thermoplasmatota class E2 (Fig. 1). Lastly,

phototrophy based carbon-fixation via the CBB cycle was mediated by members of
the Cyanobacteria, Proteobacteria and Gemmatimonadota (Supp. Data 2) as well as
microalgae in the phyla Cercozoa, Chlorophyta, Gyrista and Rhodophyta (Supp.
Data 1).

To gain a comprehensive, contextual understanding of the metabolic traits
supporting energy conservation and primary production in microbialite communities,
we conducted a comparative gene-centric analysis of West Basin Lake
metagenomes alongside previously sequenced datasets, including thrombolite,
oncolite, and stromatolite communities from five global sites (Fig. 1). The capacity for
carbon fixation was consistently high across microbialites worldwide. In West Basin
Lake, the CBB (*rbcL* av. 19.6%) and the WLP (*acsB* av. 9.92%) were predominant.
In the global datasets, the CBB was also the major pathway (av. 32.5 ± 17.0%),
while the WLP and 3-hydroxypropionate cycle showed occasional dominance in
specific samples (Fig. 1). The potential to utilise inorganic substrates exhibited
significant parallels across sites. Sulfide and H_2 appear to be dominant energy
sources, reflected by the widespread oxidation capacity (*sqr*: West Basin Lake av.
60.9%, global 40.1 ± 20.9%; Group 1-3 NiFe hydrogenase: West Basin Lake av.
24.4%, global 22.2 ± 16.0%). The capacity for CO and arsenite oxidation–ancient
metabolic traits–was also substantial across all microbial communities (*cooS & coxL*:
West Basin Lake av. 17.7%, global 11.2 ± 11.7%; *aro*: West Basin Lake av. 11.5%,
global 10.1 ± 7.3%). Conversely, photosynthesis genes showed some variation.
West Basin Lake had moderate capacity for chlorophylls and bacteriochlorophylls
mediated phototrophy (*psaA, psbA:* av. 21.9%) and microbial rhodopsins mediated
phototrophy (*rho:* av. 30.5%), whereas the global datasets exhibited broader ranges
(*psaA, psbA:* 12.6–141.3%; *rho:* 13.1–113.5%). This variability most likely
underscores the influence of local environmental conditions on energy acquisition
strategies. Overall, our findings demonstrate that the genetic potential for energy
conservation and carbon fixation in West Basin Lake microbialites is broadly
representative of other microbial communities forming microbialites that have been
studied globally. The high metabolic flexibility observed in these ecosystems enables
them to act as efficient engines of biological productivity, supporting diverse and
dynamic microbial communities.

195
196

197 Tight microbial interactions support intricate biogeochemical 198 cycles

Consistent with previous work[13,18,28,41,42], our findings reveal that microbialite
communities have remarkable metabolic diversity, and we sought to resolve the

complex molecular interactions underpinning such diversity. Millimeter-scale
community analysis revealed high microbial richness throughout the entire
microbialite structure (Supp. Fig. 1). While some taxa such as fermenters in the
order Clostridiales were confined to deeper anoxic niches (grey layer - Supp. Data
1), many other microbes were distributed across the whole microbialite structure
(Supp. Data 1), most notably Cyanobacteriales and the sulfur-oxidiser
Campylobacterales (Supp. Data 1). Similarly, members of the obligately anaerobic
order Spirochaetales were ubiquitous (Supp. Data 1). These data suggest that the
spatiotemporal physicochemical shifts such as the expansion of oxic pockets during
the day foster tight metabolic interaction that allow diverse functional guilds to
coexist and meet their physiological needs throughout the microbialite structure.

We developed a model to elucidate the array of molecular exchanges occurring
within microbialite communities (Fig. 2). Molecular hydrogen (H_2) emerges as a
central molecule. In these communities, H_2 is primarily produced through
fermentation of photosynthetically- and chemosynthetically-derived organic carbon
via diverse group 3 [NiFe]-hydrogenases and [FeFe]-hydrogenases, encoded by 90
and 55 MAGs respectively (Fig. 2). The genetic potential for other fermentation
pathways in the community is high, with many MAGs encoding marker genes
associated with acetate (*acdA, ack, pta*), formate (*fdhA, fdoG, pflD*), and lactate (*ldh*;
Supp. Data 5) fermentation. Diazotrophic Cyanobacteria and members of six other
phyla (16 MAGs) may also contribute to H_2 release as an obligate by-product of the
nitrogenase reaction. Upon diffusion into aerobic niches, H_2 is readily utilised by the
abundant community of gas oxidisers spanning 25 phyla (106 MAGs; Fig. 2), which
may utilise the electrons derived through this process for carbon fixation (Supp. Data
2). Anaerobic processes likely recycle much of the H_2, including through coupling to
denitrification (109 MAGs) and dissimilatory nitrate reduction to ammonium (55
MAGs; Fig. 2). Hydrogenotrophic acetogens were found across each microbialite
sample, albeit at low relative abundance (av. 1.2% of mapped reads). H_2 also fuels
methanogenesis and sulfate reduction to varying degrees, leading to the production
of methane and sulfide (Fig. 2). Aerobic and anaerobic CO oxidation are important
processes within microbialite communities (34 MAGs; Fig. 2), capable of supporting
energy conservation and carbon fixation in eight phyla (Supp. Data 2). A total of 78
MAGs, representing chemolithotrophic and photolithotrophic microorganisms across
nine phyla, possess the potential to oxidize sulfide under both aerobic and anaerobic
conditions (Fig. 2; Supp. Data 2). This capability likely provides a substantial
ecological advantage, facilitating their persistence in environments characterised by
dynamic redox fluctuations and intense resource competition.

239
240

Microbialite communities mediate broad aerobic and anaerobic elemental cycling

To validate our biogeochemical predictions, we performed *ex situ* microcosm experiments. The potential for oxygenic photosynthesis was assessed using chemical imaging that simultaneously visualises and quantifies O_2 concentrations in natural samples[43,44]. Algal and cyanobacterial populations, abundant in the microbialite communities, exhibited high activity, producing substantial O_2 when exposed to light intensities of ~450 µmol m^{-2} s^{-1}, mimicking their natural habitat (Fig. 3; Supp. Fig. 2). The well-illuminated microbialite surfaces demonstrated the highest gross O_2 production, though this varied markedly both within individual samples (e.g., Sample 2 ranged from 29.31 to 182.26 µmol O_2 L^{-1} min^{-1}) and among samples (Fig. 3; Supp. Fig. 2). Upon transition to darkness, O_2 was rapidly depleted, primarily through aerobic respiration and diffusion into the overlying water column (Fig. 3). The rapid O_2 consumption likely contributes to niches for the widespread facultative and obligate anaerobes at the millimetre-scale (Supp. Fig. 2). The O_2 generated via oxygenic photosynthesis also serves as an electron acceptor for several metabolic pathways, including organotrophy, hydrogenotrophy, carboxydotrophy, and nitrification. Correspondingly, all samples demonstrated oxidation of H_2, CO, and NH_4^+, albeit at varying rates (Fig. 3).

We observed consumption of H_2, CO, Fe^{2+}, and S^{2-} in nearly all anaerobic microcosms (Fig. 3), confirming the high metabolic flexibility of the anaerobic communities. This metabolic versatility aligns with genomic predictions made by previous studies in the literature[13,26,41] and reflects the capacity of microbialite communities to exploit alternative electron donors to sustain their nutrient web (Supp. Fig. 1). Such flexibility enables them to optimise energy generation across the dynamic physicochemical gradients of the microbialite environment. Consistent with these findings, physicochemical analyses revealed the presence of diverse electron acceptors in the microbialite samples. Specifically, SO_4^{2-} was present in high concentrations (av. 67.85 mg/L), whereas NO_3^- at moderate concentrations (av. 2 mg/L; Supp. Data 3).

Our genomic predictions further indicate that microbialites host microbial populations with high capacity for reductive metabolisms (Fig. 1). Microcosm experiments under prolonged anoxia confirmed the production of large amounts of reduced inorganic compounds (Fig. 3). Despite methanogens being low in abundance (av. *mcrA* 0.12% of the community), their elevated metabolic rates enable them to contribute notably to CH_4 production (av. 28.7 ppm/h; Fig. 3), a characteristic previously documented in other environments[45]. Similarly, sulfate reducers were scarce (av. *asrA, dsrA:* 2.44%) but released abundant S^{2-} (av. 7.4 µM/h; Fig. 3). On the other hand, fermenters (av. FeFe-hydrogenase: 34.37%; group 3 NiFe hydrogenase: 48.77%) and nitrogen fixer (av. *nifH:* 27.49%) abundance was mirrored in the high H_2

282 production (22.7 ppm/h; Fig. 3). These reduced compounds, diffusing through the
283 microbialite structure, fuel a diverse array of aerobic and anaerobic metabolic
284 pathways, thereby supporting microbial energy conservation and carbon fixation
285 within these dynamic ecosystems.

286
287
288

Chemosynthesis and photosynthesis contribute significantly to microbialites' carbon fixation

291
While there's long-standing evidence for photosynthesis in microbial mats[19,46,47],
recent genomic studies suggest chemosynthetic pathways also facilitate primary
production[25–27,48], though current evidence is fragmentary. To gain a comprehensive
understanding of the diversity of microbialite autotrophic populations, we performed
phylogenetic analyses of carbon fixation protein sequences. These phylogenetic
inferences were further supported by activity measurements using [14]C incorporation
assays, conducted with a range of supplemental electron donors (Fig. 4). We
analysed the phylogenies of protein sequences encoding the acetyl-CoA synthase
(AcsB), ribulose 1,5-bisphosphate (RbcL) and ATP-citrate lyase (AclB) (Fig. 4a-c).
This analysis revealed an extraordinary diversity of sequences affiliated with at least
16 phyla across the CBB, WLP and rTCA (Fig. 4a-c). These sequences
encompassed multiple novel autotrophic clades including two Nanoarchaeota MAGs
encoding the CBB, one JAHJDO01 MAG encoding the WLP, and one Polyangia
MAG encoding the rTCA (Fig. 4a-c).

306
Across the 140 RuBisCO (RbcL) sequences, we recovered four subtypes (Fig. 4a).
Notably, RuBisCO subtypes IA-D exhibit intermediate to high specificity for CO_2 and
are generally found across both aerobic and anaerobic habitats, whereas RuBisCO
subtype II has lower CO_2 specificity and is more commonly associated with
anaerobic or microaerophilic environments[49]. Consistent with the prevalent
anaerobic niches of most microbialites, we recovered 184 acetyl-CoA synthase
(AcsB) protein sequences for the WLP (Fig. 4b). However, a large proportion of
these sequences most likely were not encoded by autotrophs but rather by members
of the phyla Actinomycetota, Desulfobacterota and Chloroflexota that generally
utilise the reverse WLP for acetate oxidation coupled with dissimilatory sulfate
reduction. Conversely, sequences associated with the Firmicutes, Planctomycetota
and Spirochaetota likely represent carbon-fixing acetogens, consistent with prior
studies characterizing these phyla[50,51].

To validate the potential for carbon fixation highlighted through genomic and
phylogenetic analysis, we quantified relative chemosynthetic and photosynthetic

carbon fixation rates using radiolabeled carbon dioxide ($^{14}CO_2$). To capture both
aerobic and anaerobic carbon fixation processes, we conducted five-day
experiments, during which the microcosms transitioned to anaerobic conditions after
approximately 24 hours. This transition was inferred based on O_2 measurements
taken simultaneously on other microcosms. $^{14}CO_2$ incorporation was detected in all
microcosms, except for H_2-stimulated chemosynthesis in microbialite C (Fig. 4d,
Supp. Fig. 3), which most likely resulted from an experimental issue. As anticipated,
microcosms exposed to light supported higher $^{14}CO_2$ incorporation through
photosynthesis (av. 0.0122 nmol day^{-1} g wet^{-1}) compared to those incubated in the
dark (av. 0.00279 nmol day^{-1} g wet^{-1}; Fig. 4d). Dark incubations likely captured the
combined activity of anaplerotic processes and baseline chemosynthetic carbon
fixation. The addition of electron donors significantly boosted chemosynthetic carbon
fixation rates (Fig. 4d). Consistent with the high abundance of genes and MAGs
encoding *sqr* and *rdsrA* genes (Fig. 1), S^{2-}-supplemented microcosms exhibited the
highest chemosynthetic $^{14}CO_2$ incorporation rates (av. 0.0084 nmol day^{-1} g wet^{-1}).
Similarly, NH_4^+ and H_2 supplementation enhanced chemosynthetic carbon fixation
(av. 0.00741 and 0.0062 nmol day^{-1} g wet^{-1}, respectively; Fig. 4d). Contrary to the
prevailing view that microbial productivity is predominantly driven by
photoautotrophs, our findings reveal that inorganic chemical sources and gas
substrates are also critical for the primary productivity of these ecosystems,
highlighting the significant role of chemosynthesis in supporting stratified microbial
communities.

345
346

Conclusions

Through an integrative approach, our study sheds light on the intricate molecular
exchanges that underpin the carbon acquisition, energy conservation, and broader
nutrient cycling within a model microbialite community, offering a detailed view into
how taxonomic and metabolic diversity intersect in such ecosystems. Microbialite
communities in the hypersaline West Basin Lake exhibit a metabolic diversity
comparable to that observed in other microbialite ecosystems worldwide (Fig. 1).
This diversity likely arises from the presence of organisms with complementary traits,
driven by resource facilitation among community members[52,53]. In microbialite
communities, metabolic synergies occur between organisms inhabiting contrasting
physicochemical niches[26,54], which vary on adiel cycle The resultant mosaic of
ecological niches align with the physiological requirements of diverse microbial taxa.
Consistent with these observations, our analysis simultaneously reveals millimetre-
scale overlap of functional guilds throughout the microbialite structure (Supp. Fig. 2)
and cycling of key metabolites such as iron, nitrogen and sulfur compounds across

their major redox states through intricate molecular handoffs (Supp. Data 2). These
processes promote minimal energy loss and enhance ecosystem productivity[55]. In a
broader context, the exceptional efficiency of elemental cycling and carbon use
efficiency within microbialite communities, combined with the duality of light- and
chemically-driven metabolic pathways, suggests that these ecosystems have likely
served as hotspots of metabolic innovation throughout Earth's history.
368
369

Methods

Characterization of microbial mat organosedimentary structures

The eastern shoreline of West Basin Lake hosts abundant microbialites and living
microbial matgrounds[56]. Well-developed microbialite buildups and hardgrounds are
exposed on subaerially-exposed benches up to several metres above the current
lake level. Previous studies have documented extensive sub-aqueous, well-lithified
living microbialites with considerable relief above the lake floor, especially in the
adjacent East Basin Lake[56]. The living microbialites documented in this study occur
in shallow water of around 0.5 m depth, and are best-developed in benches
approximately 1-2 m from the current shoreline in West Basin Lake (Supp. Fig. 4).
These living microbial matgrounds appear dark orange-red and have an irregular,
undulating surface with a broad-scale relief of several centimetres. In cross section,
the mats have a dark orange-red film over a hard to friable, mineralised, cream-white
uppermost layer of up to several millimetres in thickness (Supp. Fig. 1). This
mineralised layer has an irregular, pustular appearance and is composed of
carbonate, likely hydromagnesite[56]. Underneath this, the mat consists of weakly
lithified to unlithified mud with several thin colour zones over a several millimetres
depth (dark green, purple and grey), followed by up to several centimetres of light
green mat (Supp. Fig. 1; Supp. Fig. 4). The sediment texture is commonly clotted
and unlaminated. The substrate for the mat is dark, organic-rich mud which may
have weak layering preserved through the presence of occasional coarse sediment
laminae. Here we use the term 'microbialite' to define these microbial matgrounds,
as they do form mineralised structures with cm-scale relief above the lake floor.
While these modern and recent microbialites are not as extensively developed as the
sub-aerially exposed microbialites, they do appear to form significant accretionary
structures over time (for example encrusting anthropogenic debris).

Sample collection, processing and physicochemical parameters

397 Forming microbialites were collected from West Basin Lake (38° 19' 24.6468'' S,
398 143° 26' 51.8928'' E), a hypersaline (salinity 6.5-9.5%) inland crater lake located in
399 Victoria, Australia. Microbialites samples were collected during three field trips in
400 November 2022, March 2024 and May 2024 using pre-sterilized spades and

transferred into pre-sterilized 5-gallon buckets containing lake water. At first collection in November 2022, the water temperature was 18.9 °C, dissolved oxygen 90%, pH 8.47 and oxidative reduction potential 2.72 mV. Irradiance at noon was measured using a Walz Universal Light Meter (ULM-500) equipped with a Mini Quantum Sensor (LS-C) at 30 cm and 50 cm below the water surface was ~650 µmol photons m^{-2} s^{-1} and ~450 µmol photons m^{-2} s^{-1}, respectively. Physicochemical parameters for lakewater and microbialite depth-profile subsamples were measured at the TrACEES Platform, University of Melbourne. The selected physicochemical parameters include total organic carbon, total carbon, inorganic carbon, total nitrogen, sodium, magnesium, potassium, calcium, sulfate and chlorum. Nitrate was measured at Water Studies, Monash University.

16S rRNA gene sequencing and community analysis

Total DNA was extracted from the layers of two different microbialite samples (three technical triplicate per layer) using the DNeasy PowerSoil Kit (Qiagen, Hilden, Germany) on 0.5g of material. Two sample-free negative control were also included. Extracted DNA samples were sent to AGRF (Melbourne, Victoria) for library preparation, PCR-amplification and sequencing of the 16S rRNA gene V1-V3 regions on an Illumina MiSeq platform, 2x300bp paired-end reads. Sequences were processed using the QIIME2 pipeline v 2022.2[57]. Primer sequences were trimmed using Cutadapt[58], while DADA2 was employed for merging forward and reverse reads, quality filtering, dereplication, and chimera removal[59]. Taxonomic classification was performed with QIIME2's feature-classifier plugin. The SILVA v132 QIIME release was used for 16S rRNA gene taxonomy[60].

Community DNA extraction and sequencing

Each microbialite sample was homogenised into a slurry. Total DNA was extracted from three different microbialite slurry samples (technical triplicate per sample for a total of nine samples) using the DNeasy PowerMax Soil Kit (Qiagen, Hilden, Germany) on 10 g of materials as per the manufacturer's protocol. A sample-free negative control was also included. Extracted DNA samples were sent to AGRF (Melbourne, Victoria) for library preparation and sequencing on two lanes using an Illumina NovaSeq SP Flow-cell, 2 x 150 bp for 500 cycles.

Reads quality control, assembly and binning

Across the three technical replicates of each microbialite sample, we obtained an average of over 16 million read pairs for sample 1, over 56 million read pairs for sample 2, and over 40 million read pairs for sample 3. Reads quality control, assembly and binning were implemented within the Metaphor pipeline[61]. Specifically,

raw reads derived from the nine metagenome libraries were quality-controlled by
trimming primers and adapters, followed by artefacts and low-quality read filtering
using fastp[62] with parameters *length_required 50*, *cut_mean_quality 30*, and *extra: --*
detect_adapter_for_pe. The nine quality-controlled metagenomes were coassembled
using MEGAHIT v1.2.9[63] with default parameters. Contigs shorter than 1,000 bp
were removed. Assembled contigs were binned using Vamb v4.1.3[64], MetaBAT
v2.12.1[65] and CONCOCT v1.1.0[66]. The three bin sets were then refined using DAS
Tool v1.1.6[67] and de-replicated using dRep v3.4.2[68] with 95% ANI integrated with
CheckM2[69]. Bins completeness and contamination were estimated using CheckM2[69].
After dereplication, we recovered 331 between medium (completeness >50%,
contamination <10%) and high-quality (completeness >90%, contamination <5%)
metagenome-assembled genomes (MAGs), according to the MIMAG standard[70].
MAG taxonomy was assigned according to Genome Taxonomy Database Release
R214[71] using GTDB-Tk v2.3.2[72]. CoverM v0.6.1 was used to calculate the relative
abundance of bins based on the metagenomic reads
(https://github.com/wwood/CoverM).
457
458

Functional annotation of binned contigs and unbinned contigs

The sequences of 51 marker genes representing energy conservation, carbon
fixation, trace gas metabolism, sulfur cycle, nitrogen cycle, arsenic cycle, iron cycle,
formate oxidation, phototrophy, and aerobic respiration were retrieved from the 331
MAGs and unbinned contigs. Open reading frames (ORFs) were predicted using
Prodigal v2.6.3[73], then annotated using DIAMOND blastp[74] homology-based
searches against a custom database[75] of 51 metabolic marker gene sets described
below. DIAMOND mapping was performed with a query coverage threshold of 80%
for all databases, and a percentage identity threshold of 80% (for *psaA*) 75% (for
mcr, hbsT), 70% (for *isoA, psbA, ygfK, aro, atpA*), 60% (*amoA, pmoA, mmoA, coxL,*
[FeFe], nxrA, rbc, nuoFL) or 50% (all other databases). MAGs were also annotated
using DRAM[76].
471
472

Metabolic annotation of metagenomic short reads

Paired-end reads from the nine samples were stripped of adapter and barcode
sequences, then contaminating PhiX and low-quality sequences were removed
(minimum quality score 20) using the BBDuk function of BBTools v.36.92
(https://sourceforge.net/projects/bbmap/). Resultant quality-filtered forward reads
with lengths of at least 100 bp were searched for the presence of the 51 marker
genes described above using the DIAMOND blastx algorithm[77]. Specifically, reads
were compared against the custom-made reference databases[75] of 51 metabolic
marker genes for energy conservation, carbon fixation, phototrophy, and hydrogen,

carbon monoxide, methane, sulfur, nitrogen, and iron cycling. A query coverage of
80% and an identity threshold of 80% (for *psaA*), 75% (for *hbsT*), 70% (for *atpA*,
psbA, isoA, ygfK, aro), 60% (for *amoA, mmoA, coxL, FeFe, nxrA, rbcL, nuoF*) and
50% (for all others) was used. The proportion of community members encoding each
gene was calculated by normalizing the gene's read count (measured in reads per
kilobase million [RPKM]) against the average RPKM of 14 universal single-copy
ribosomal marker genes.

489
490

Phylogenetic analysis

Maximum-likelihood phylogenetic trees for archaeal and bacterial MAGs were built
using GTDBtk[72] commands *identify* and *align* on high quality MAGs (completion
>90% and contamination <5%). The archaral and bacterial trees were built using IQ-
TREE v2.2.2.6[78,79] with 1,000 ultrafast bootstrap[80] using the LG+C10+F+G and
WAG+G20 models, respectively. MUSCLE[81] was used to align 36 AclB, 184 AcsB
and 140 RbcL proteins retrieved between the MAGs and unbinned contigs. Aclb,
AcsB and RbcL maximum-likelihood phylogenetic trees were built using IQ-TREE
v2.2.2.6[78,79] with 1,000 ultrafast bootstrap[80] and models LG+F+I+R6 for the AcsB
tree, LG+I+G4 for the AclB tree and LG+R5 for the RbcL tree. All trees were plotted
using iTOL v6[82].

502
503

Chemical imaging

The O_2 sensitive optode preparation included mixing 100 mg of polystyrene, 1.5 mg
of indicator (PT (II) meso-tetra(pentafluorophenyl)porphine), 1.5mg of reference
(Macrolex yellow®) and dissolved in 1 g of solvent (Tetrahydrofuran) to form a
cocktail. The O_2 cocktails were knife-coated on dust-free polyester foils
(goodfellow.com) and the final thickness of the coating was <2 μm. Once dry, the O_2
sensitive optode was coated with an antirefractory layer. The antirefractory cocktail
preparation included mixing 100 mg hydrogel D4, 100mg carbon black and 1 g of
100% ethanol. The antirefractory cocktail were knife-coated on top of the O_2
sensitive optode and the final thickness of the coating was <3 μm.

The experimental setup included a modified digital single-lens reflex camera (Canon
EOS 1000D) with its near-infrared (NIR) blocking filter removed and equipped with a
Sigma 50 mm F2.8 EX DG Macro lens. An emission filter (Schott 530 nm,
Uqgoptics.com) was fitted to the lens to detect oxygen fluorescence. Following the
protocol described by Larsen et al. (RED), O_2 sensitive optode were excited by four
high-power blue LEDs (I-peak = 445 nm, LXHL-LR3C, Luxeon, F = 340 mW at IF =
700 mA) combined with a 470 nm short-pass filter (blue dichroic color filter,
Uqgoptics.com). Microbialite sample cross-sections were illuminated using a Schott

Leica KL 2500 LCD Cold Light Source. All components were synchronized via a
trigger box (https://imaging.fish-n-chips.de) and controlled using the custom software
Look@RGB. Each planar optode was calibrated individually in an aquarium
maintained at a constant seawater temperature of 20 ± 1°C in a darkened room. The
calibration range for the O_2 sensitive optode was 0 – 360 µmol L^{-1}.

All experiments were conducted in a dark room at a constant 20 ± 1 °C to resemble
the lake temperature at the time of sampling. Three microbialite samples were cut
using a diamond saw exposing their cross-sections. Each of these samples was
placed in a 4 L glass aquarium pressing on the O_2 sensitive optode, which was
attached to the aquarium side. Following overnight acclimation in the aquaria,
microbialite sample cross-sections were illuminated from above with approximately
450 µmol m^{-2} s^{-1} of light. Irradiance levels in the experimental setup for defined lamp
settings were measured using a Walz Universal Light Meter (ULM-500) equipped
with a Mini Quantum Sensor (LS-C). Image sequences capturing O_2 dynamics
across the microbialite cross-sections were taken every 5 minutes. Lake water in the
aquarium was aerated with an aeration stone connected to an air pump.

Downstream data analysis was conducted using ImageJ v1.53K. Each image was
separated into Red, Green, Green2, and Blue RAW TIFF channels. The ImageJ
plugin Ratio Plus was used to calculate the ratio of the Red to Green channels (R/G).
The resulting ratio images were color-coded using the 'Fire' lookup table to visualise
O_2 dynamics. Calibration was performed using the Curve Fitting function, applying an
exponential fit with offset for O_2, based on planar optode calibration values.
Brightness and contrast settings were adjusted to display minimum and maximum
values of 0 – 360 µmol L^{-1} for O_2 images. To minimize influence of water infiltrating
in between the O_2 optode and microbialite cross-section and to quantify microbial O_2
production, the first image of each experiment was subtracted from subsequent
images using the Image Calculator function. To identify the photosynthetic regions
on the microbialite sample cross-sections, we overlaid images with the highest O_2
production onto microbialite cross-section images. The portions of each microbialite
sample that showed O_2 production were identified as regions on interes (ROI).
Subsequently values were extracted from the ROI. Control sample in Fig. 3 was an
image sequence of ROI measuring a deeper, anaerobic microbialite cross-section
portion simultaneously to O_2 measurements. Proxies for the net photosynthesis (PN)
and apparent dark respiration (RD) were calculated by subtracting images taken with
a 5 min interval when O_2 production and respiration were the highest, respectively.
Gross photosynthesis was estimated as PG = PN+|RD|.

Ex-situ biogeochemical measurements

We conducted microcosm experiments to evaluate the aerobic and anaerobic
metabolism of microbialite communities. Each experiment was performed in
triplicate, with three independent microbialite samples per replicate. Heat-killed
samples were prepared using gamma radiation followed by one autoclave cycle at
121 °C for 30 minutes and served as controls. These controls confirmed that the
observed element dynamics were attributable to biotic processes.

Aerobic and anaerobic microcosms, including trace gas and S^{2-} and Fe^{2+} additions,
were set up in 120 mL serum vials containing 50 mL of 0.22 µm-filtered lake water
and approximately 10 g of microbialite to create slurries. The vials were sealed with
butyl rubber septa. For aerobic microcosms, the headspace was left with ambient air,
while anaerobic microcosms were flushed with helium for 10 minutes to remove O_2
and then supplemented with 1.5 mM NO_3^- as an electron acceptor (except for S^{2-}
consumption). The weights of microbialite used in each microcosm were recorded
and used to normalise downstream calculations.

Trace gas microcosms were supplemented with 10 ppm H_2, CH_4, and CO in the
headspace. Sampling of the headspace began immediately after the addition of
electron donors and acceptors, with 2 mL of gas extracted at variable time intervals
as shown in Fig. 3. In anaerobic vials, the sampled gas volume was replaced with
He. Gas concentrations were analysed by gas chromatography using a pulsed
discharge helium ionization detector (model TGA-6791-W-4U-2, Valco Instruments
Company Inc.), with calibration based on certified standard mixtures of H_2, CH_4, and
CO (0, 10, 100 ppm in N_2, BOC Australia).

In anaerobic S^{2-} and Fe^{2+} microcosms, either 100 µM Na☐S·9H☐O (only for
consumption) or 6 mM FeCl☐ was added to helium-purged slurries. At each
timepoint, 3 mL of water was sampled and filtered through 0.45 µm pore-size filters.
For S^{2-} analysis, 2 mL of the filtered sample was preserved with ZnAc, while for Fe^{2+}
analysis, 1 mL was preserved with ferrozine. Both S^{2-} and Fe^{2+} concentrations were
measured using a GBC UV-Visible 918 spectrophotometer, following methods
described previously[83].

Microcosms for nitrification were prepared in uncapped 250 mL Schott bottles
containing 100 mL of 0.22 µm-filtered lake water, 100 µM NH_4^+, and approximately
10 g of microbialite sample. At each sampling timepoint, 10 mL of water was filtered
through 0.45 µm pore-size filters and stored frozen until further analysis. The filtered
samples were analysed for NOx (NO_2^- + NO_3^-) concentrations using a Lachat
QuikChem 8000 Flow Injection Analyzer (FIA) in accordance with APHA methods[84].
For oxygen measurements, microcosms were prepared in 120 mL vials containing
100 mL of 0.22 µm-filtered lake water and approximately 10 g of microbialite slurry.
Dissolved O_2 concentrations were monitored using a FireSting oxygen probe
(PyroScience) until the microcosms approached anoxic conditions.

0.25 g of homogenised microbialite sample with 1 mL of filtered lake water (0.22 µm)
were prepared in 7 mL scintillation vials with ambient air headspaces. Radiolabeled
sodium bicarbonate solution (NaH$_{14}$CO$_3$, Perkin Elmer, 53.1 mCi nmol^{-1}) was added
to a concentration ~0.1 µM. Triplicates of each sample were prepared and subjected
to five different conditions, namely light (40 µmol m^{-2} s^{-1}), dark, dark + H$_2$ (100 ppm),
dark + S^{2-} (800 µM) and dark + NH$_4^+$ (1 mM) and incubated for 5 days. These
experiments aimed at capturing both aerobic and anaerobic carbon incorporation
metabolisms and according to parallel O$_2$ measurements, microcosms likely
transitioned to anaerobic after approximately 24 h. After the incubation period,
concentrated HCl was added dropwise to each vial and left for 24h with intermittent
shaking to ensure excess unbound dissolved inorganic carbon (DIC) was acidified
and released as ^{14}CO$_2$. HCl was added equally to all vials until bubble production
ceased and then they were placed at 60 °C under a heat lamp to dry. When dry, 7
mL of scintillation liquid (EcoLume™, MP Biomedical) was added and ^{14}C measured
on an automated liquid scintillation counter (Tri-Carb 2810 TR, Perkin Elmer).
Photosynthetic ^{14}C incorporation values were adjusted to account for the
photosynthetic ROI areas present on the 0.25g portion of microbialite used in the
light treatment. Assuming that the photosynthetic ROI areas identified through
chemical imaging represent the microbialite sample regions performing oxygenic
photosynthesis, we normalised by applying the ratio of photosynthetic ROI area of
microbialite sample 3 which had the largest photosynthetic surface area
(photosynthetic ROI sample 1: 3.11%, sample 2: 2.73%, and sample 3: 6.35%) to
total microbialite area. We developed a hierarchical Bayesian model to analyze ^{14}C
incorporation rates under different conditions. The model is implemented in Stan and
R (see SI for code). The model consists of N total observations, with I samples, J
conditions, and K replicates for each sample condition combination. As the Dark+H$_2$
condition experiment failed for microbialite C, we ran two models: one excluding
microbialite C and included all conditions (where I=2, J=5, K=3 and N=30), and one
excluding the Dark+H$_2$ condition and included all samples (where I=3, J=4, K=3 and
N=36), so that in each case the model can be fitted to a complete dataset. The
likelihood function is given by:
632
633    $y_{ijk} \sim N(\alpha_i + \beta_j, \sigma^2)$,
634
where α_i describes the sample specific effect for the ith sample, β_j describes the
condition specific effect for the jth sample, and σ^2 is the unexplained variance in the
data. The Dark condition (j=1) was prescribed to be the control by setting $\beta_1 =$
$0.$ Weakly informative priors were used. Priors for α_i ($i \in \{1:I\}$) and β_j ($j \in \{2:J\}$)
were specified as normal distributions with variances an order of magnitude larger
than the variance of the total dataset. The prior for σ was a Cauchy distribution.
Posterior probability distributions for values of β are presented in figure 4D, with

Credible intervals defined between the 2.5th and 97.5th, and the 25th and 75th
percentiles.

Statistics and visualization

Downstream statistical analyses were performed in RStudio (version 1.2.5033) using
several R packages, including decontam[85], ggplot2[86], phyloseq[87], and vegan[88].
Illustrator v24.0.2 was utilised for figure editing.
649

Data availability

651 All data supporting the findings of the present study are available. We deposited all
652 the project's sequences at the Sequence Read Archive. Metagenomic sequences
653 have accession number PRJNA1194634, 16S rRNA gene sequences have
654 accession number PRJNA1194668, and MAGs have been submitted and awaiting
655 release.
656

Conflict of interest

658 The authors declare no conflict of interest.
659

Acknowledgments

We acknowledge Judy and Leon for kindly allowing us to conduct our study on their
property at West Basin Lake. F.R. was supported by the Early Career Postdoctoral
Fellowship (ECPF24-4273843556) awarded by the Faculty of Medicine, Nursing and
Health Science at Monash University, and internal funding awarded to H.M. from the
University of Melbourne. P.M.L. was supported by an ARC DECRA Fellowship
(DE250101210). C.G. was supported by an NHMRC EL2 Fellowship (APP1178715).
A.H. was supported by an ARC DECRA Fellowship (DE190100988). This study used
the MASSIVE M3 supercomputing infrastructure.
669

Author contribution

F.R., C.G., P.M.L., and H.M. conceptualised the study. Experimental planning and design were conducted by F.R., C.G., T.H., V.E., W.W.W., and P.C. Fieldwork was conducted by F.R., H.M. and A.H. A.H. provided microbialite description. Data analysis was led by F.R. Gas chromatography measurements were carried out by F.R., T.H., and T.N., while V.E. and W.W.W. oversaw nitrification measurements. Chemical imaging, oxygen, sulfide and ferrous oxide measurements were performed by F.R. Carbon fixation incubations were performed by T.H. Metagenome analysis, MAG construction, and annotation were completed by F.R., with extensive bioinformatics support provided by V.W.S. Resources, supervision, and funding were contributed by C.G., P.C., and H.M. The manuscript was written by F.R., with input from all authors.

References

1. Walter, M. R., Buick, R. & Dunlop, J. S. R. Stromatolites 3,400–3,500 Myr old from the North pole area, Western Australia. *Nature* **284**, 443–445 (1980).

2. Van Kranendonk, M. J., Philippot, P., Lepot, K., Bodorkos, S. & Pirajno, F. Geological setting of Earth's oldest fossils in the ca. 3.5 Ga Dresser formation, Pilbara Craton, Western Australia. *Precambrian Research* **167**, 93–124 (2008).

3. Dick, G. J., Grim, S. L. & Klatt, J. M. Controls on O_2 Production in Cyanobacterial Mats and Implications for Earth's Oxygenation. *Annu. Rev. Earth Planet. Sci.* **46**, 123–147 (2018).

4. Cavalier-Smith, T. Cell evolution and Earth history: stasis and revolution. *Phil. Trans. R. Soc. B* **361**, 969–1006 (2006).

5. Nisbet, E. G. & Fowler, C. M. R. Archaean metabolic evolution of microbial mats. *Proc. R. Soc. Lond. B* **266**, 2375–2382 (1999).

6. Hoehler, T. M., Bebout, B. M. & Des Marais, D. J. The role of microbial mats in the production of reduced gases on the early Earth. *Nature* **412**, 324–327 (2001).

7. Des Marais, D. J. Earth's early biosphere. *Gravitational and space biology bulletin: publication of the American Society for Gravitational and Space Biology* **11**, 23–30 (1998).

8. Hawes, I., Sumner, D. Y., Andersen, D. T., Jungblut, A. D. & Mackey, T. J. Timescales of growth response of microbial mats to environmental change in an ice-

covered Antarctic lake. *Biology* **2**, 151–176 (2013).

9.   Nold, S. C. *et al.* Underwater sinkhole sediments sequester Lake Huron's carbon.

Biogeochemistry **115**, 235–250 (2013).

10.   Boidi, F. J., Mlewski, E. C., Gomez, F. J. & Gérard, E. Characterization of

Microbialites and Microbial Mats of the Laguna Negra Hypersaline Lake (Puna of

Catamarca, Argentina). in *Microbial Ecosystems in Central Andes Extreme Environments*

(ed. Farías, M. E.) 183–203 (Springer International Publishing, Cham, 2020).

doi:10.1007/978-3-030-36192-1_13.

11.   Paul, V. G., Wronkiewicz, D. J., Mormile, M. R. & Foster, J. S. Mineralogy and

Microbial Diversity of the Microbialites in the Hypersaline Storr's Lake, the Bahamas.

Astrobiology **16**, 282–300 (2016).

12.   Bischoff, K. *et al.* Spherulitic microbialites from modern hypersaline lakes, Rottnest

Island, Western Australia. *Geobiology* **18**, 725–741 (2020).

13.   Osman, J. R. *et al.* Key energy metabolisms in modern living microbialites from

hypersaline Andean lagoons of the Salar de Atacama, Chile. *Science of The Total*

Environment **937**, 173469 (2024).

14.   Pace, A. *et al.* Microbial and diagenetic steps leading to the mineralisation of Great

Salt Lake microbialites. *Scientific Reports* **6**, 31495 (2016).

15.   Laval, B. *et al.* Modern freshwater microbialite analogues for ancient dendritic reef

structures. *Nature* **407**, 626–629 (2000).

16.   Valdespino-Castillo, P. M. *et al.* Exploring biogeochemistry and microbial diversity of

extant microbialites in Mexico and Cuba. *Frontiers in microbiology* **9**, 510 (2018).

17.   Centeno, C. M. *et al.* Microbialite genetic diversity and composition relate to

environmental variables. *FEMS microbiology ecology* **82**, 724–735 (2012).

18.   Myshrall, K. L. *et al.* Biogeochemical cycling and microbial diversity in the thrombolitic

microbialites of Highborne Cay, Bahamas. *Geobiology* **8**, 337–354 (2010).

19.   Dupraz, C. & Visscher, P. T. Microbial lithification in marine stromatolites and

hypersaline mats. *Trends in Microbiology* **13**, 429–438 (2005).

20.   Sabbe, K., Verleyen, E., Hodgson, D. A., Vanhoutte, K. & Vyverman, W. Benthic

diatom flora of freshwater and saline lakes in the Larsemann Hills and Rauer Islands,

East Antarctica. *Antarctic science* **15**, 227–248 (2003).

21.   Patterson, D. J. & Simpson, A. G. Heterotrophic flagellates from coastal marine and

hypersaline sediments in Western Australia. *European Journal of Protistology* **32**, 423–

448 (1996).

22.   Hauer, G. & Rogerson, A. Heterotrophic Protozoa from Hypersaline Environments. in

Adaptation to Life at High Salt Concentrations in Archaea, Bacteria, and Eukarya (eds.

Gunde-Cimerman, N., Oren, A. & Plemenitaš, A.) vol. 9 519–539 (Springer-Verlag,

Berlin/Heidelberg, 2005).

23.   Post, F. J., Borowitzka, L. J., Borowitzka, M. A., Mackay, B. & Moulton, T. The

protozoa of a Western Australian hypersaline lagoon. *Hydrobiologia* **105**, 95–113 (1983).

24.   Toneatti, D. M., Albarracín, V. H., Flores, M. R., Polerecky, L. & Farías, M. E.

Stratified bacterial diversity along physico-chemical gradients in high-altitude modern

stromatolites. *Frontiers in Microbiology* **8**, 1–12 (2017).

25.   Wong, H. L., Smith, D. L., Visscher, P. T. & Burns, B. P. Niche differentiation of

bacterial communities at a millimeter scale in Shark Bay microbial mats. *Scientific Reports*

5, 1–17 (2015).

26.   Shoemaker, A. *et al.* Wood–Ljungdahl pathway encoding anaerobes facilitate low-

cost primary production in hypersaline sediments at Great Salt Lake, Utah. *FEMS*

Microbiology Ecology **100**, fiae105 (2024).

27.   Wong, H. L., MacLeod, F. I., White, R. A., Visscher, P. T. & Burns, B. P. Microbial

dark matter filling the niche in hypersaline microbial mats. *Microbiome* **8**, 135 (2020).

28.   Saghaï, A. *et al.* Comparative metagenomics unveils functions and genome features

of microbialite☐associated communities along a depth gradient. *Environmental*

Microbiology **18**, 4990–5004 (2016).

29.   Gutiérrez-Preciado, A. *et al.* Functional shifts in microbial mats recapitulate early

Earth metabolic transitions. *Nat Ecol Evol* **2**, 1700–1708 (2018).

30.   Ragsdale, S. W. & Pierce, E. Acetogenesis and the Wood–Ljungdahl pathway of

CO_2 fixation. *Biochimica et Biophysica Acta (BBA)-Proteins and Proteomics* **1784**, 1873–

1898 (2008).

31.   Fuchs, G. Alternative Pathways of Carbon Dioxide Fixation: Insights into the Early

Evolution of Life? *Annu. Rev. Microbiol.* **65**, 631–658 (2011).

32.   Campbell, M. A. *et al.* Functional gene expression in Shark Bay hypersaline microbial

mats: adaptive responses. *Frontiers in Microbiology* **11**, 560336 (2020).

33.   Bernstein, H. C. *et al.* Trade-offs between microbiome diversity and productivity in a

stratified microbial mat. *The ISME journal* **11**, 405–414 (2017).

34.   Moorhead, D. L., Wolf, C. F. & Wharton, R. A. Impact of light regimes on productivity

patterns of benthic microbial mats in an antarctic lake: A modeling study. *Limnology &*

Oceanography **42**, 1561–1569 (1997).

35.   Williams, T. J., Allen, M. A., Berengut, J. F. & Cavicchioli, R. Shedding light on

microbial "dark matter": insights into novel Cloacimonadota and Omnitrophota from an

Antarctic Lake. *Frontiers in Microbiology* **12**, 741077 (2021).

36.   Li, Y., Liu, H., Xiao, Y. & Jing, H. Metagenome sequencing and 982 microbial

genomes from Kermadec and Diamantina Trenches sediments. *Scientific Data* **11**, 1067

(2024).

37.   Fang, Y. *et al.* Casting Light on the Adaptation Mechanisms and Evolutionary History

of the Widespread Sumerlaeota. *mBio* **12**, e00350-21 (2021).

38.   Edgcomb, V. P. *et al.* Active eukaryotes in microbialites from Highborne Cay,

Bahamas, and Hamelin Pool (Shark Bay), Australia. *The ISME journal* **8**, 418–429 (2014).

39.   Couradeau, E. *et al.* Prokaryotic and eukaryotic community structure in field and

cultured microbialites from the alkaline Lake Alchichica (Mexico). *PloS one* **6**, e28767

(2011).

40.   Bonacolta, A. M., Visscher, P. T., Del Campo, J. & White III, R. A. The eukaryome of

modern microbialites reveals distinct colonization across aquatic ecosystems. *npj Biofilms*

and Microbiomes **10**, 78 (2024).

41.   Breitbart, M. *et al.* Metagenomic and stable isotopic analyses of modern freshwater

microbialites in Cuatro Ciénegas, Mexico. *Environmental Microbiology* **11**, 16–34 (2009).

42.   Kanik, M. *et al.* Unexpected Abundance and Diversity of Phototrophs in Mats from

Morphologically Variable Microbialites in Great Salt Lake, Utah. *Appl Environ Microbiol*

86, e00165-20 (2020).

43.   Ricci, F. *et al.* Fine□scale mapping of physicochemical and microbial landscapes of

the coral skeleton. *Environmental Microbiology* 1462-2920.16369 (2023)

doi:10.1111/1462-2920.16369.

44.   Larsen, M., Borisov, S. M., Gunwald, B., Klimant, I. & Glud, R. N. A simple and

inexpensive, high-resolution color-ratiometric planar optode imaging approach:

Application to oxygen and pH sensing. *Limnology and Oceanography: Methods* **2**, 348–

360 (2011).

45.   Hall, N. *et al.* Aerotolerant methanogens use seaweed and seagrass metabolites to

drive marine methane emissions. *bioRxiv* 2024–10 (2024).

46.   Martínez-Alonso, M. *et al.* Distribution of phototrophic populations and primary

production in a microbial mat from the Ebro Delta, Spain. *International Microbiology* **7**,

19–25 (2004).

47.   Canfield, D. E. & Des Marais, D. J. Biogeochemical cycles of carbon, sulfur, and free

oxygen in a microbial mat. *Geochimica et Cosmochimica acta* **57**, 3971–3984 (1993).

48.   Kurth, D. *et al.* Arsenic metabolism in high altitude modern stromatolites revealed by

metagenomic analysis. *Scientific reports* **7**, 1024 (2017).

49.   Prywes, N., Phillips, N. R., Tuck, O. T., Valentin-Alvarado, L. E. & Savage, D. F.

Rubisco Function, Evolution, and Engineering. *Annu. Rev. Biochem.* **92**, 385–410 (2023).

50.   Garritano, A. N., Song, W. & Thomas, T. Carbon fixation pathways across the

bacterial and archaeal tree of life. *PNAS nexus* **1**, pgac226 (2022).

51.   Fuchs, G. CO2 fixation in acetogenic bacteria: variations on a theme. *FEMS*

Microbiology Reviews **2**, 181–213 (1986).

52.   Loreau, M. & Hector, A. Partitioning selection and complementarity in biodiversity

experiments. *Nature* **412**, 72–76 (2001).

53.   Cardinale, B. J. *et al.* Effects of biodiversity on the functioning of trophic groups and

ecosystems. *Nature* **443**, 989–992 (2006).

54.   Dunham, E. C. *et al.* An ecological perspective on dolomite formation in Great Salt

Lake, Utah. *Frontiers in Earth Science* **8**, 24 (2020).

55.   Seto, M. & Iwasa, Y. Microbial material cycling, energetic constraints and ecosystem

expansion in subsurface ecosystems. *Proc. R. Soc. B.* **287**, 20200610 (2020).

56.   Last, W. M. & De Deckker, P. Modern and Holocene carbonate sedimentology of two

saline volcanic maar lakes, southern Australia. *Sedimentology* **37**, 967–981 (1990).

57.   Bolyen, E. *et al.* Reproducible, interactive, scalable and extensible microbiome data

science using QIIME 2. *Nature biotechnology* **37**, 852–857 (2019).

58.   Martin, M. Cutadapt removes adapter sequences from high-throughput sequencing

reads. *EMBnet. journal* **17**, 10–12 (2011).

59.   Callahan, B. J. *et al.* DADA2: High-resolution sample inference from Illumina

amplicon data. *Nature Methods* **13**, 581–583 (2016).

60.   Quast, C. *et al.* The SILVA ribosomal RNA gene database project: improved data

processing and web-based tools. *Nucleic acids research* **41**, D590–D596 (2012).

61.   Salazar, V. W. *et al.* Metaphor—a workflow for streamlined assembly and binning of

metagenomes. *GigaScience* **12**, giad055 (2023).

62.   Chen, S. Ultrafast one□pass FASTQ data preprocessing, quality control, and

deduplication using fastp. *Imeta* **2**, e107 (2023).

63.   Li, D., Liu, C.-M., Luo, R., Sadakane, K. & Lam, T.-W. MEGAHIT: an ultra-fast single-

node solution for large and complex metagenomics assembly via succinct de Bruijn

graph. *Bioinformatics* **31**, 1674–1676 (2015).

64.   Nissen, J. N. *et al.* Improved metagenome binning and assembly using deep

variational autoencoders. *Nat Biotechnol* **39**, 555–560 (2021).

65.   Kang, D. D. *et al.* MetaBAT 2: an adaptive binning algorithm for robust and efficient

genome reconstruction from metagenome assemblies. *PeerJ* **7**, e7359 (2019).

66.   Alneberg, J. *et al.* Binning metagenomic contigs by coverage and composition.

Nature methods **11**, 1144–1146 (2014).

67.   Sieber, C. M. K. *et al.* Recovery of genomes from metagenomes via a dereplication,

aggregation and scoring strategy. *Nature Microbiology* **3**, 836–843 (2018).

68.   Olm, M. R., Brown, C. T., Brooks, B. & Banfield, J. F. dRep: a tool for fast and

accurate genomic comparisons that enables improved genome recovery from

metagenomes through de-replication. *The ISME Journal* **11**, 2864–2868 (2017).

69.   Chklovski, A., Parks, D. H., Woodcroft, B. J. & Tyson, G. W. CheckM2: a rapid,

scalable and accurate tool for assessing microbial genome quality using machine

learning. *Nature Methods* **20**, 1203–1212 (2023).

70.   Bowers, R. M. *et al.* Minimum information about a single amplified genome (MISAG)

and a metagenome-assembled genome (MIMAG) of bacteria and archaea. *Nature*

biotechnology **35**, 725–731 (2017).

71.   Parks, D. H. *et al.* GTDB: an ongoing census of bacterial and archaeal diversity

through a phylogenetically consistent, rank normalized and complete genome-based

taxonomy. *Nucleic acids research* **50**, D785–D794 (2022).

72.   Chaumeil, P.-A., Mussig, A. J., Hugenholtz, P. & Parks, D. H. GTDB-Tk v2: memory

friendly classification with the genome taxonomy database. *Bioinformatics* **38**, 5315–5316

(2022).

73.   Hyatt, D. *et al.* Prodigal: prokaryotic gene recognition and translation initiation site

identification. *BMC Bioinformatics* **11**, 119 (2010).

74.   Altschul, S. F. *et al.* Gapped BLAST and PSI-BLAST: a new generation of protein

database search programs. *Nucleic Acids Research* **25**, 3389–3402 (1997).

75.   Greening, C. Greening lab metabolic marker gene databases. (2021).

76.   Shaffer, M. *et al.* DRAM for distilling microbial metabolism to automate the curation of

microbiome function. *Nucleic acids research* **48**, 8883–8900 (2020).

77.   Buchfink, B., Xie, C. & Huson, D. H. Fast and sensitive protein alignment using

DIAMOND. *Nature methods* **12**, 59–60 (2015).

78.   Kalyaanamoorthy, S., Minh, B. Q., Wong, T. K., Von Haeseler, A. & Jermiin, L. S.

ModelFinder: fast model selection for accurate phylogenetic estimates. *Nature methods*

14, 587–589 (2017).

79.   Minh, B. Q. *et al.* IQ-TREE 2: new models and efficient methods for phylogenetic

inference in the genomic era. *Molecular biology and evolution* **37**, 1530–1534 (2020).

80.   Hoang, D. T., Chernomor, O., Von Haeseler, A., Minh, B. Q. & Vinh, L. S. UFBoot2:

improving the ultrafast bootstrap approximation. *Molecular biology and evolution* **35**, 518–

522 (2018).

81.   Edgar, R. C. Muscle5: High-accuracy alignment ensembles enable unbiased

assessments of sequence homology and phylogeny. *Nature Communications* **13**, 6968

(2022).

82.   Letunic, I. & Bork, P. Interactive Tree of Life (iTOL) v6: recent updates to the

phylogenetic tree display and annotation tool. *Nucleic Acids Research* gkae268 (2024).

83.   Islam, Z. F. *et al.* Two Chloroflexi classes independently evolved the ability to persist

on atmospheric hydrogen and carbon monoxide. *ISME J* **13**, 1801–1813 (2019).

84.   Rice, E. W., Bridgewater, L. & Association, A. P. H. *Standard Methods for the*

Examination of Water and Wastewater. vol. 10 (American public health association

Washington, DC, 2012).

85.   Davis, N. M., Proctor, D. M., Holmes, S. P., Relman, D. A. & Callahan, B. J. Simple

statistical identification and removal of contaminant sequences in marker-gene and

metagenomics data. *Microbiome* **6**, 1–14 (2018).

86.   Wickham, H. ggplot2. *WIREs Computational Stats* **3**, 180–185 (2011).

87.   McMurdie, P. J. & Holmes, S. phyloseq: an R package for reproducible interactive

analysis and graphics of microbiome census data. *PloS one* **8**, e61217 (2013).

88.   Dixon, P. VEGAN, a package of R functions for community ecology. *Journal of*

vegetation science **14**, 927–930 (2003).

Figures legend

Figure 1 | Maximum-likelihood genome trees depicting the taxonomic diversity of 331 archaeal and bacterial metagenome-assembled genomes (MAGs), built with 1000 ultrafast bootstrap replicates using the LG+C10+F+G and WAG+G20 models, respectively. The top heatmap highlights the metabolic potential of microbes co-encoding energy acquisition enzymes with carbon fixation pathways. The side bargraph shows the number of MAGs in each microbial family (GTDB taxonomy). The bottom heatmap shows the abundance of each gene in the metagenomic short reads in West Basin Lake samples (bold) and across 17 publicly available microbialite community metagenomes from five global sites. Homology-based searches were used to identify signature genes encoding enzymes associated with metabolic pathways. To infer abundance, read counts were normalised to gene length and the abundance of single-copy marker genes.

Figure 2 | Metabolic model predicted based on genome- and gene-resolved data of the dominant microbial guilds within the microbialite communities of West Basin Lake. This model represents the inferred metabolic pathways and interactions based on the genomic content of these guilds. Blue and brown lines indicate the direction of electron acceptors and donors, respectively. Note that the graphic is an artistic generalization of the data and should not be interpreted as an exact depiction of the microbial community structure. Asterisk (*) denotes that specific metabolic marker genes were exclusively recovered from metagenomic short-read data.

Figure 3 | Biogeochemical assays illustrating the metabolic activities of microbialite communities under aerobic (**a**) and anaerobic (**b**, **c**) conditions in microcosms. Oxygen dynamics were assessed using chemical imaging on independent microbialite samples incubated in 4 L glass aquaria, with data presented as the mean ± standard deviation for defined photosynthetic regions of interest. Trace gases, sulfide and ferrous ion measurements were taken in 120 ml sealed serum vial containing 10 g of microbialite slurry and 50 mL of 0.22 µm-filtered lake water. Trace gas microcosms were supplemented with 10 ppm H_2, CH_4, and CO in the headspace. S^{2-} and Fe^{2+} microcosms were supplied with either 100 µM $Na_2S \cdot 9H_2O$ (only for consumption) or 6 mM $FeCl_2$. All anaerobic microcosms except for S^{2-} production were supplemented with 1.5 mM NO_3^- as an electron acceptor. Nitrification (NOx = NO_2^- + NO_3^-) measurements were taken in uncapped 250 mL Schott bottles containing approximately 10 g of microbialite slurry, 100 mL of 0.22 µm-filtered lake water and 100 µM NH_4^+. In the oxygen plot, asterisks (*) indicate the time when light was switched off, mimicking the onset of darkness. All microcosm experiments were performed in triplicate, with results expressed as the mean ±

940   standard deviation across three replicates. All measurements except for oxygen
941   dynamics were normalised to microbialite slurry wet weight.
942

Figure 4 | Maximum-likelihood phylogenetic trees of 140 RbcL (**a**), 184 AcsB (**b**),
and 36 AclB (**c**) amino acid sequences obtained from the three microbialite samples,
constructed using 1000 ultrafast bootstrap replicates. The LG+R5 (**a**), LG+F+I+R6
(**b**), and LG+I+G4 (**c**) substitution models were applied. Sequences derived from
binned contigs are highlighted in bold, while those from unbinned contigs are shown
without emphasis. Bootstrap support values ≥ 90 are denoted by white circles (**a–c**).
Panel (**d**) Credible intervals of effect (electron donors) relative to dark condition
denoting 2.5th and 97.5th (thin bar) and 25th and 75th (thick bar) percentiles of
posterior probability of the effect of each assay condition relative to the dark
condition.
953

CO$_2$ + H$_2$O → NH$_3$

Cyano-bacteria

H$_2$ + organic carbon + O$_2$ N$_2$

anox. phototrophs → energy

H$_2$S or H$_2$ + CO$_2$

H$_2$ & CO oxidiser

H$_2$ or CO+ O$_2$ or NO$_3^-$/SO$_4^{-2}$ + CO$_2$ organic carbon + H$_2$O

SOB → SO$_4^{-2}$ organic carbon

O$_2$ or NO$_3^-$ + H$_2$S

denitrifier
N$_2$
N$_2$O
NO
NO$_2^-$ CO$_2$

organic carbon + NO$_3^-$

N$_2$ fixer
H$_2$
NH$_3$ N$_2$

DNRA
NO$_3^-$ + H$_2$ NH$_4^+$

fermenter
H$_2$, acetate, etc.   organic carbon

SRB
CO$_2$ H$_2$S   organic carbon or H$_2$ + SO$_4^{-2}$

acetogen
H$_2$ + CO$_2$ acetate

methanogen
*
H$_2$ + CO$_2$ CH$_4$

Oxygen concentration
+
−

Phyla

- Acidobacteriota
- Actinomycetota
- Actinobacteriota
- Armatimonadota
- Asgardarchaeota
- Bacteroidota
- Bdellovibrionota
- Cloacimonadota
- Campylobacterota
- Chlamydiota
- Chloroflexota
- Cyanobacteriota
- Deinococcota
- Desulfobacterota
- Fermentibacterota
- Firmicutes
- Gemmatimonadota
- Halobacteriota
- Hydrogenedentota
- Krumholzibacteriota
- Myxococcota
- Nanoarchaeota
- Nitrospinota
- Omnitrophota
- Patescibacteria
- Planctomycetota
- Proteobacteria
- Spirochaetota
- Sumerlaeota
- Thermoplasmatota
- Thermoproteota
- UBP6
- Verrucomicrobiota
- JAHJDO01