Neural bidomain model for multidimensional ephaptic coupling of neural spike propagation along myelinated fiber bundles with the nodes of Ranvier

Sehun Chun [1*], Linyu Peng [2], Hae-Jeong Park [3]

1 Underwood International College, Yonsei University, Seoul, South Korea
2 Department of Mechanical Engineering, Keio University, Yokohama, Japan
3 Graduate School of Medical Science, Brain Korea 21 Project, Department of Nuclear Medicine, Yonsei University College of Medicine, Seoul, South Korea

* sehun.chun@yonsei.ac.kr

Abstract

A novel mathematical model is proposed to investigate the effects of ephaptic coupling between general neural fiber bundles in a multidimensional space with anisotropic neural fiber bundles. Ephaptic coupling corresponds to the spatiotemporal interaction between propagating fiber bundles through the extracellular space. Adapted from a well-known model in cardiac electrophysiology, the bidomain model comprises of the nonoverlapping intracellular and extracellular space except the common nodes of Ranvier. The proposed two-variable model, a *neural bidomain model*, is derived from the classic Frankenhaeuser–Huxley model for neural spike propagation along the general neural fiber bundles. The governing equation is mathematically and computationally validated against existing one-dimensional models of neural fiber bundle propagations with aligned nodes of Ranvier. A high-order continuous Galerkin scheme is employed for efficient two-dimensional computational simulation with moving frames, or orthonormal basis vectors, representing the intracellular and extracellular conductivity and nonoverlapping domains. The proposed model is simulated in various two-dimensional configurations of neural fiber bundles that are little known to the community, such as fiber bundles with misaligned nodes of Ranvier, opposite-traveling fiber bundles, and curved fiber bundles.

Author summary

Ephaptic coupling is a communication method among fiber bundles, other than chemical interactions via synaptic junctions and electrical interactions via gap junctions. Ephaptic coupling is of a small magnitude compared to other communication methods. The effect is known to be on a large scale and is more affected by the intrinsic structure of neural fiber bundles. Consequently, the study of ephaptic coupling has attracted broad attention since the 1940s, when the term *ephaptic coupling* was first coined. However, the lack of a multidimensional model of neural spike propagation with various configuration of myelinated fiber bundles with the nodes of Ranvier prohibits further investigations of the effects of ephaptic coupling in various brain subregions. This paper proposes a multidimensional model of neural spike simulation in the context of partial differential equations to simulate neural spike propagations in various complex multidimensional fiber bundles and study the role of ephaptic coupling in such brain structures.

Introduction

Since Hodgkin and Huxley's pioneer work in the 1950s [1], neural spike propagation has been successfully modeled with ion channels or a smaller number of macroscopic dynamic variables [2]. A sharp increase in the neuron cell membrane potential is received at the dendrite and propagates along a neural axon to reach the axon terminal of a spiking neuron [3]. The neural fiber bundle, the multitude of neural axons linking distributed neural populations in parallel, is the central mediator of interactions within the brain via axon potentials. The signal transfer is often considered *digital* [4] because it does not depend on the pathways between the end points. The previous mathematical models of spike propagation have primarily been represented as ordinary differential equations; thus, the geometric and spatial configuration of the cable-like fiber bundle does not significantly affect signal propagation. This phenomenon contrasts the mathematical models of physical waves often represented in partial differential equations depending on the geometry of the spatiotemporal domain. This digital connectivity without space and time has inspired many virtual neural network algorithms but failed to provide insight into the unique morphology and configuration of brain fibers for complex and high-mode brain functions.

A complementary propagation of the neural spike propagation is achieved via the extracellular space of axons [3]. Closely packed neurons can laterally interact through the potential change in the extracellular space. This lateral interaction of the neural signal propagation through ion exchange in the extracellular medium is known as *ephaptic coupling*. Since Arvanitaki coined this term in the 1940s [5], it has been broadly used to include more general *electric field effects* [6–9] for the spatiotemporal electric potential distribution in the extracellular space for neural spike propagation and the consequent neural activities. This paper primarily focuses on ephaptic coupling among neighboring neural fibers by the extracellular potential created by the neural propagation of the fibers. The neural fibers are ephaptically coupled by the diffusion of ions occurring at the nodes of Ranvier, where the myelin partly unwraps from the axon. The myelin with high lipid composition insulates electricity along the axon; thus, the signal propagates along the myelinated axon with little electricity leakage. The interconnectivity of the nodes of Ranvier with the extracellular medium sends additional fluctuations of electric potential changes in the extracellular medium, as illustrated in Fig. 1. This complementary potential distribution appears to interfere with neural spike propagation along neural fibers, analogous to the complementary relationship between electric and magnetic fields for light propagation [10–12].

Fig 1. Illustration of ephaptic coupling via neural spike propagation along a single fiber.

The mathematical modeling of ephaptic coupling for myelinated fibers with nodes of Ranvier is often based on Frankenhaeuser and Huxley's 1960s model for the myelinated axon [13], a subsequent work based on the original model by Hodgkin and Huxley for an unmyelinated axon [1]. Goldman and Albus [14] investigated this model further to

demonstrate that the conduction velocity is proportional to the fiber diameter. In the 1970s, further developments of Frankenhaeuser and Huxley's model with updated parameter values were achieved by Waxman and Brill [15–17] and Markin [18–21] to study the conduction for various myelinated and unmyelinated configurations and dynamics. In 2003, Reutskiy et al. [22] adopted a similar model to investigate the dependency of the conduction velocity on various parameters including temperature and the conductivity ratio between intracellular and extracellular space. More recently, various mathematical and computational models were developed to study the interaction between fiber axon bundles the validate various experimental results, including spike propagation delay and sensory stimuli processing [8, 10, 11, 23]. All previous models are either derived in a one-dimensional (1D) domain of a neural fiber or unable to incorporate a complex configuration of myelin sheath and the nodes of Ranvier. This 1D model without complex fiber bundle configuration limits further understanding of multidimensional ephaptic coupling in the complex neural fiber bundles.

This paper proposes a multidimensional bidomain model for simulating the ephaptic coupling of neural spike propagation along general neural fibers. The concept of the bidomain model, first proposed by Schmitt in 1969 [24], was initially developed for neural propagation along nerve fibers [25, 26] but has become popular for propagating the cardiac electric signal in the heart [27]. This work derives a multidimensional Frankenhaeuser–Huxley model using the bidomain approach for non-overlapping intracellular and extracellular domains. In contrast to the bidomain model of cardiac electrophysiology, the two domains are non-overlapping, and the membrane potential and extracellular potential should be defined separately for each domain. Two different sets of orthogonal basis vectors, known as *moving frames* [28–33], are introduced for each variable in both domains to address this problem. The proposed multidimensional model is validated by demonstrating that it yields the previously known mathematical and computational results in 1D model.

In addition to proposing a multidimensional mathematical model, this paper aims to provide further goals in the context of computational neuroscience. This work explores multidimensional cases that are only possible with the proposed model. The first goal is to observe the synchronization of misaligned the nodes of Ranvier. The synchronization of myelinated fibers with aligned node of Ranvier is well-known. Nevertheless, the propagation behavior of misaligned nodes of Ranvier is largely unknown because of the required multidimensional model. The second goal is to observe the changes in conduction velocity in the traveling neural spikes that are not in the same direction. Many crossing fibers in the brain demand propagational velocity changes around that region, but no analysis or simulation has been conducted. A simple case is simulated in this paper:oppositely traveling neural spikes in parallel fibers. The third case is the conduction velocity changes in curved fibers. Brain fiber is primarily curved, and the conduction velocity is sufficiently rapid such that the curvedness of the neural fiber should affect its propagation and the corresponding synchronization. The proposed multidimensional model can simulate a neural spike propagation along curved fibers to offer insights into the function of the curvedness of the fibers.

Materials and methods

The previous mathematical and computational studies including the original model by Frakenhaeuser and Huxley [13] are primarily confined to 1D domain Ω^{1d} [10, 14, 22] with overlapped fibers. An axon bundle with a myelin sheath and node of Ranvier in Ω^{1d} is considered, where $\phi_i^k(x, t)$ and $\phi_e(x, t)$ denote the electric potential of the intracellular space of the kth fiber and the extracellular space in Ω^{1d}, respectively. The extracellular potential is *collective* at every point for all fibers. The membrane potential ϕ_m^k is

defined as $\phi_m^k = \phi_i^k - \phi_e$. The gradient of the membrane potential proportional to the electric current is expressed as follows:

$$\frac{\partial \phi_m^k(x,t)}{\partial x} = r_i I_i^k(x,t) - r_e I_e(x,t), \quad x \in \Omega^{1d}, \tag{1}$$

where r_i and r_e present the fiber resistance in the intercellular and extracellular media, respectively. If the values of r_i and r_e are the same for all fibers and I_i^k and I_e represent the electric current for the intercellular medium of the kth fiber and the extracellular medium, respectively. Substituting $I_e = (1/r_e)\partial \phi_e / \partial x$ into Eq. (1) yields the following intracellular electric flow for the kth fiber:

$$I_i^k = \frac{1}{r_i} \left(\frac{\partial \phi_m^k}{\partial x} + \frac{\partial \phi_e}{\partial x} \right). \tag{2}$$

The first component is the electric flow by the gradient of the membrane potential ϕ_m^k. The second component is the additional current caused by the gradient of the extracellular potential ϕ_e, known as the *ephaptic current* (point of contact) for the kth fiber. The electric flow of the membrane potential in the fiber is affected by the changes in the extracellular medium. In other words, two separate fibers are connected and communicated via the extracellular medium. The conservation of currents induces the dynamics of the extracellular potential ϕ_e contributed from all fibers as follows:

$$I_{eph} = \sum_{k=1}^{N} I_{eph}^k = \frac{\alpha_N}{r_i} \sum_{k=1}^{N} \frac{\partial \phi_m^k}{\partial x}. \tag{3}$$

The constant $\alpha_N := \rho/(1 + N\rho)$ is called the *coupling strength constant* [22], where ρ denotes the conductivity ratio between the intracellular and extracellular space, defined as $\rho = r_e/r_i$. Moreover, $\rho = 1$ corresponds to the case in which the conductivity of the intracellular space is the same as that of the extracellular space. The coupling strength constant α_N, which is usually less than 1.0, depends on the resistivity of the extracellular space and the surface ratio of the fiber. Eq. (3) is only possible by assuming a specific alignment between ϕ_m^k and ϕ_e to simplify the dynamics of ephaptic coupling in the 1D domain. Substituting Eq. (3) into Eq. (2) yields the following equation for the total current using the linearity of electric potentials:

$$I_i = \sum_k I_i^k = \frac{1}{r_i} \frac{\partial \phi_m}{\partial x} + I_{eph}, \tag{4}$$

where $\phi_m = \sum_k \phi_m^k$. Differentiating Eq. (4) with respect to x and incorporating the corresponding leakage or ion channel reaction yields the following equations for the membrane potential dynamics for the myelin region:

$$c_m \frac{\partial \phi_m(x,t)}{\partial t} = \frac{1}{r_i} \frac{\partial^2 \phi_m(x,t)}{\partial x^2} + \frac{\partial I_{eph}(x,t)}{\partial x} - g_m(x)\phi_m(x,t), \quad x \in \Omega^{1d}, \tag{5}$$

where c_m represents the myelin capacitance per unit length, 1.87×10^{-11} F/cm. In addition, g_m denotes the myelin conductance per unit length, 5.60×10^{-9} F/cm [16]. The membrane equation for the node of Ranvier is similarly derived as follows:

$$c_n \frac{\partial \phi_m(x,t)}{\partial t} = \frac{1}{r_i} \frac{\partial^2 \phi_m(x,t)}{\partial x^2} + \frac{\partial I_{eph}(x,t)}{\partial x} - I_{ion}(x,t) - I_{ext}(x,t), \quad x \in \Omega^{1d}, \tag{6}$$

where c_n indicates the nodal axolemma capacitance per unit length, 3.14×10^{-9} F/cm, and I_{ext} represents the external current. Moreover, I_{ion} denotes the ionic current per unit length adopted from the model by Frankenhaeuser–Huxley as follows [13, 22]:

$$I_{ion}(\phi_m, K, Na) = \pi d \left[I_K(\phi_m, K) + I_{Na}(\phi_m, Na) + I_p(\phi_m, Na) + I_L(\phi_m) \right], \quad (7)$$

where d denotes the fiber diameter, I_K represents the potassium current, I_{Na} indicates sodium current, I_p denotes the non-specific delayed current, and I_l indicates the leak current, defined as follows:

$$I_K(\phi_m, K) = P_K n^2 Z(\phi_m, K), \quad (8)$$
$$I_{Na}(\phi_m, Na) = P_{Na} m^2 h Z(\phi_m, Na), \quad (9)$$
$$I_p(\phi_m, Na) = P_p p^2 Z(\phi_m, Na), \quad (10)$$
$$I_L(\phi_m) = g_L(\phi_m - \phi_L), \quad (11)$$

where g_L indicates the leak conductance, $0.0303\ mho/cm^2$, and ϕ_L represents the leak current equilibrium potential, 0.026 mV. The ion variables m, h, n, and p correspond to the sodium permeability, inactivation of sodium permeability, potassium permeability, and non-specific permeability, respectively. The function Z in Eqs. (8) to (11) is defined as follows:

$$Z(\phi_m, Y) = 10^{-3} F \left[[Y]_i \mathcal{E} \left(-10^{-3} \frac{F\phi_m}{RT} \right) - [Y]_0 \mathcal{E} \left(10^{-3} \frac{F\phi_m}{RT} \right) \right],$$

where $[\cdot]_0$ and $[\cdot]_i$ are the corresponding ion density values in the extracellular and intracellular space, respectively. In addition, T denotes the absolute temperature, R presents the gas constant of 8.3145 (J/mol/K), and F indicates the Faraday constant of $96,484.6$ (C/mol). Moreover, the function $\mathcal{E}$ is defined as follows:

$$\mathcal{E}(z) = \begin{cases} \frac{z}{\exp(z)-1}, & \text{if } z \geq 10^{-4}, \\ 1 - \frac{z}{2}, & \text{if } z < 10^{-4}. \end{cases} \quad (12)$$

The corresponding constants and ion permeability dynamics for the variables (m, n, p, h) are the same as found in [13, 16, 22]. Additionally, Fig. (2) displays the time-dependent dynamics during excitation. The 1D single variable model in Eqs. (5) to (6) is compact and effective for predicting and analyzing ephaptic coupling effects. However, the restriction of this model are clear. For example, Eq. (3) assumes that all nodes of Ranvier should be allocated in the exact location with the same excitation initiation along the fibers. Consequently, this 1D model is only relevant to special case studies, such as the synchronization strength of neural spikes, failing to investigate the multidimensional ephaptic coupling effects with various fiber configurations.

Mathematical modeling of multidimensional neural spike propagation

This work considers a two-dimensional (2D) isolated domain Ω consisting of the intracellular and extracellular space. The intracellular domain corresponds to the region neurons occupy, denoted by Ω_i. The extracellular space corresponds to the space between neurons, denoted by Ω_e. The intracellular domain is considered independent of the extracellular space with a clear boundary between the domains because the membrane has high resistance with thousands of $\Omega \cdot cm^2$ compared to the resistance of the extracellular space with hundreds of $\Omega \cdot cm^2$ [34]. The intracellular space Ω_i consists of the nodes of Ranvier Ω_n and the internodal region Ω_m covered by a myelin sheath (Fig. 3), where

(a) Membrane potential   **(b)** Ion channels

Fig 2. Dynamics of membrane potential and ion channels during excitation using the Frankenhaeuser–Huxely model.

$$\Omega_i = \Omega_m \cup \Omega_n. \tag{13}$$

Ion channel reactions primarily occur in Ω_n. The extracellular space Ω_e corresponds to the area outside the intracellular domain but includes the nodes of Ranvier:

$$\Omega_e = \Omega_m^c = \Omega_i^c \cup \Omega_n, \tag{14}$$

where the superscript c represents the complement of the corresponding domain. The intracellular and extracellular space comprises the entire domain Ω to have the node of Ranvier in common:

$$\Omega_i \cup \Omega_e = \Omega, \quad \Omega_i \cap \Omega_e = \Omega_n. \tag{15}$$

Compared to the length of the myelinated node at 2 mm, the length of the node of Ranvier is approximately 1000 times smaller about 2.5 μ m [16, 22]. The width of an axon is approximately 2 μ m, the same for the myelinated node and node of Ranvier. The boundary between the myelinated cell and node of Ranvier is not specified but is distinguished as the boundary of elements with different conductivities.

Fig 3. Two-dimensional fiber bundle domain Ω with the node of Ranvier Ω_n and myelin sheath Ω_m. Ω_i: intercellular space, Ω_e: extracellular space.

The electric current along the neural fiber is denoted by $\mathbf{J}_i$ in the intracellular region Ω_i, and the conductivity tensor and electric potential are denoted by $\widehat{\sigma}_i$ and ϕ_i, respectively, in Ω_i. The hat sign implies that the corresponding quantity is a tensor with a magnitude and direction. Note that $\widehat{\sigma}_i$ is strongly anisotropic such that a strong conductivity exists along the axon and a weak conductivity exists orthogonal to the

axonal direction. Hence, the electric current along the fiber is expressed as the gradient of the electric potential ϕ_i along $\widehat{\sigma}_i$ as follows:

$$\mathbf{J}_i(\mathbf{x}, t) = \widehat{\sigma}_i(\mathbf{x}, t)\nabla\phi_i(\mathbf{x}, t), \quad \text{if } \mathbf{x} \in \Omega_i. \tag{16}$$

For the general 3D space, $\widehat{\sigma}_i$ can be represented as the following matrx.

$$\widehat{\sigma}_i(\mathbf{x}, t) = \begin{bmatrix} e_1^x & e_2^x & e_3^x \\ e_1^y & e_2^y & e_3^y \\ e_1^z & e_2^z & e_3^z \end{bmatrix} \begin{bmatrix} \sigma_1 & 0 & 0 \\ 0 & \sigma_2 & 0 \\ 0 & 0 & \sigma_3 \end{bmatrix} \begin{bmatrix} e_1^x & e_2^x & e_3^x \\ e_1^y & e_2^y & e_3^y \\ e_1^z & e_2^z & e_3^z \end{bmatrix}^T, \tag{17}$$

where $\mathbf{e}_1 = [e_1^x \ e_1^y \ e_1^z]$ indicates the unit direction vector of the neural fiber with a conductivity of σ_1. Alternatively, a curve $\gamma(s)$ is considered to be the parametrization of the neural fiber with the parameter s within a certain range, and $\mathbf{e}_1$ denotes the unit tangent vector of the curve $\gamma(s)$. Then, Eq. (16) is equivalently expressed as follows:

$$\mathbf{J}_i(\mathbf{x}, t) = \sigma_1\nabla\phi_i(\gamma(s), t), \quad \text{if } \gamma(s) \in \Omega_i. \tag{18}$$

Similarly, the extracellular current $\mathbf{J}_e$ is defined in the extracellular space Ω_e with diffusivity conductivity $\widehat{\sigma}_e$ and electric potential ϕ_e as

$$\mathbf{J}_e(\mathbf{x}, t) = \widehat{\sigma}_e(\mathbf{x}, t)\nabla\phi_e(\mathbf{x}, t), \quad \text{if } \mathbf{x} \in \Omega_e, \tag{19}$$

where $\widehat{\sigma}_e$ is assumed to have isotropic diffusivity with a magnitude of σ_e. Applying the divergence operator in Eqs. (16) and (19) yields the following conservation of current densities [27]:

$$\nabla \cdot (\widehat{\sigma}_i\nabla\phi_i) = \nabla \cdot \mathbf{J}_i = I_m, \tag{20}$$

$$\nabla \cdot (\widehat{\sigma}_e\nabla\phi_e) = \nabla \cdot \mathbf{J}_e = -I_m, \tag{21}$$

where I_m represents the membrane current density per unit area. The current flow across the membrane can be described by time-dependent and ionic currents as follows:

$$\nabla \cdot (\widehat{\sigma}_i\nabla\phi_i(\mathbf{x}, t)) = \begin{cases} c_m\frac{\partial\phi_m(\mathbf{x},t)}{\partial t} + g_m\phi_m(\mathbf{x}, t), & \mathbf{x} \in \Omega_m, \\ c_n\frac{\partial\phi_m(\mathbf{x},t)}{\partial t} + I_{ion}(\mathbf{x}, t), & \mathbf{x} \in \Omega_n, \end{cases} \tag{22}$$

where c_m and c_n denote the corresponding capacitances, the same as those in Eqs. (5) and (6). Moreover, g_m indicates the myelin conductance of the myelin sheath, and I_{ion} denotes the ionic current per unit length. Substituting the definition of the membrane potential $\phi_i = \phi_m + \phi_e$ into Eq. (22) yields

$$c_m\frac{\partial\phi_m(\mathbf{x}, t)}{\partial t} = \nabla \cdot (\widehat{\sigma}_i\nabla\phi_m(\mathbf{x}, t)) + \nabla \cdot (\widehat{\sigma}_i\nabla\phi_e(\mathbf{x}, t)) - g_m\phi_m(\mathbf{x}, t), \quad \mathbf{x} \in \Omega_m, \tag{23}$$

$$c_n\frac{\partial\phi_m(\mathbf{x}, t)}{\partial t} = \nabla \cdot (\widehat{\sigma}_i\nabla\phi_m(\mathbf{x}, t)) + \nabla \cdot (\widehat{\sigma}_i\nabla\phi_e(\mathbf{x}, t)) - I_{ion}(\mathbf{x}, t), \quad \mathbf{x} \in \Omega_n. \tag{24}$$

Each tessellated element is assumed to be either a myelin sheath or the node of Ranvier, not both regions in a single element. Table 1summarizes each region, defining each variable and tensor.

To derive the conservation of ions in Ω, each variable $\phi_i \in \Omega_i$ and $\phi_e \in \Omega_e$ defined in each non-overlapping domain is expanded into $\phi_i^\Omega \in \Omega$ and $\phi_e^\Omega \in \Omega$ in the entire domain Ω as follows:

$$\phi_i^\Omega(\mathbf{x}, t) = \begin{cases} \phi_i(\mathbf{x}, t), & \text{if } \mathbf{x} \in \Omega_i, \\ 0, & \text{otherwise}, \end{cases} \quad \text{and} \quad \phi_e^\Omega(\mathbf{x}, t) = \begin{cases} \phi_e(\mathbf{x}, t), & \text{if } \mathbf{x} \in \Omega_e, \\ 0, & \text{otherwise}. \end{cases} \tag{25}$$

	ϕ_i	ϕ_e	ϕ_m	$\widehat{\sigma}_i$	$\widehat{\sigma}_e$
Node of Ranvier	O	O	O	O	O
Myeline	O	O	O	O	X
Extracellular space	X	O	X	X	O

Table 1. Regions defining each variable and tensor (O: defined; X: not defined).

 Similarly, the conductivity tensor of $\widehat{\sigma}_i \in \Omega_i$ and $\widehat{\sigma}_e \in \Omega_e$ are expanded into $\sigma_i^\Omega \in \Omega$ and $\sigma_e^\Omega \in \Omega$ as

$$\widehat{\sigma}_i^\Omega(\mathbf{x},t) = \begin{cases} \widehat{\sigma}_i(\mathbf{x},t), & \text{if } \mathbf{x} \in \Omega_i, \\ 0, & \text{otherwise,} \end{cases} \quad \text{and} \quad \widehat{\sigma}_e^\Omega(\mathbf{x},t) = \begin{cases} \widehat{\sigma}_e(\mathbf{x},t), & \text{if } \mathbf{x} \in \Omega_e, \\ 0, & \text{otherwise.} \end{cases} \quad (26)$$

Adding Eqs. (16) and (19) with Eqs. (25) and (26) yields the following equation for the conservation of ions:

$$\nabla \cdot \left[\widehat{\sigma}_i^\Omega(\mathbf{x},t)\nabla\phi_i^\Omega(\mathbf{x},t)\right] = -\nabla \cdot \left[\widehat{\sigma}_e^\Omega(\mathbf{x},t)\nabla\phi_e^\Omega(\mathbf{x},t)\right], \quad \mathbf{x} \in \Omega. \quad (27)$$

For the electric potential ϕ, the corresponding quantity of $\nabla \cdot \hat{\sigma}\nabla\phi$ represents the current source density corresponding to the overall charged ion density [34–36]. The left- and right-hand sides of Eq. (27) corresponds to the current source density of the intracellular and extracellular space, respectively. Thus, Eq. (27) states that the sum of the current source density of the intracellular and extracellular space is zero, equivalent to the conservation of ions in a domain with two separate spaces. Substituting $\phi_i = \phi_m + \phi_e$ into Eq. (27) results in the following equations.

$$\nabla \cdot \left[(\widehat{\sigma}_e^\Omega(\mathbf{x},t) + \widehat{\sigma}_i^\Omega(\mathbf{x},t))\nabla\phi_e^\Omega(\mathbf{x},t)\right] = -\nabla \cdot \left[\widehat{\sigma}_i^\Omega(\mathbf{x},t)\nabla\phi_m^\Omega(\mathbf{x},t)\right], \quad \mathbf{x} \in \Omega. \quad (28)$$

Equations similar to Eqs. (23), (24) and (28) are known as *bidomain equations*, particularly in cardiac electric signal propagation [27, 37, 38]. The 2D decompositions of tensors $\widehat{\sigma}_i^\Omega$ and $\widehat{\sigma}_e^\Omega$ are obtained as follows, assuming a cylindrical shape of the fiber with a circular base of radius a:

$$\widehat{\sigma}_i^\Omega(\mathbf{x},t) = \begin{bmatrix} e_1^x & e_2^x & e_3^x \\ e_1^y & e_2^y & e_3^y \\ e_1^z & e_2^z & e_3^z \end{bmatrix} \begin{bmatrix} \pi a^2\sigma_i & 0 & 0 \\ 0 & \pi a^2\sigma_i & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} e_1^x & e_2^x & e_3^x \\ e_1^y & e_2^y & e_3^y \\ e_1^z & e_2^z & e_3^z \end{bmatrix}^T, \quad (29)$$

$$\widehat{\sigma}_e^\Omega(\mathbf{x},t) = \begin{bmatrix} e_1^x & e_2^x & e_3^x \\ e_1^y & e_2^y & e_3^y \\ e_1^z & e_2^z & e_3^z \end{bmatrix} \begin{bmatrix} \pi a^2\sigma_i & 0 & 0 \\ 0 & \pi a^2\sigma_i & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} e_1^x & e_2^x & e_3^x \\ e_1^y & e_2^y & e_3^y \\ e_1^z & e_2^z & e_3^z \end{bmatrix}^T, \quad (30)$$

where σ_i and σ_e denote the maximum conductivity in the intracellular and extracellular space, respectively. Substituting Eqs. (29) and (30) into Eq. (28) results in the following Poisson equation along the fiber direction $\mathbf{e}_1$ and its orthogonal direction $\mathbf{e}_2$ for $\mathbf{x} \in \Omega$,s

$$\nabla \cdot [(\sigma_i + \sigma_e)(\mathbf{e}_1 \cdot \nabla\phi_e^\Omega)\mathbf{e}_1] = -\nabla \cdot [\sigma_i(\mathbf{e}_1 \cdot \nabla\phi_m^\Omega)\mathbf{e}_1], \quad (31)$$

$$\nabla \cdot [(\pi a^2\sigma_i + \sigma_e)(\mathbf{e}_2 \cdot \nabla\phi_e^\Omega)\mathbf{e}_2] = -\nabla \cdot [\pi a^2\sigma_i(\mathbf{e}_2 \cdot \nabla\phi_m^\Omega)\mathbf{e}_2]. \quad (32)$$

Moreover, another variable is introduced to approximate the case with a significant extracellular potential, compared to the membrane potential, without including more fibers. If the strength of the extracellular potential is proportional to the square of the

fiber bundle radius, r_b, as modeled in [11], which is continuously filled with axons of radius a, then the cumulative extracellular potential at x outside the fiber can be derived as follows:

$$\phi_e(x) = \frac{2\pi}{\Omega(a)} \int_0^{r_b} r\phi_e^0(x,r)dr, \tag{33}$$

where $\Omega(a)$ is the cross-sectional area filled by the axons of radius a and ϕ_e^0 represent the magnitude of the extracellular potential of each fiber. If each induced extracellular potential by every axon is constant as ϕ_e^0 along the radial direction, then the following expression defines the extracellular potential inside the fiber bundle.

$$\phi_e(x) = \frac{\pi r_b^2}{\Omega(a)}\phi_e^0 = \frac{\rho_f g^2 r_b^2}{a^2}\phi_e^0, \quad 0 \le x \le r_b, \tag{34}$$

where $\Omega(a) = \pi a^2/(\rho g^2)$ is applied for the constant ρ_f as the relative size of the volume occupied by the fibers, and g is the ratio between the axon diameter and fiber diameter (the g-ratio). Substituting Eq. (34) in Eq. (32) results in the multidimensional Poisson equation:

$$\nabla \cdot [\pi a^2 \sigma_i + \sigma_e)(\mathbf{e}_2 \cdot \nabla\phi_e)\mathbf{e}_2] = -\alpha_f^2 \nabla \cdot [\pi a^2 \sigma_i(\mathbf{e}_2 \cdot \nabla\phi_m)\mathbf{e}_2], \tag{35}$$

where α_f denotes the amplified factor of the extracellular potential, defined as $\alpha_f^2 = 1/(\rho g^2 r_b^2)$ [11]. The increased fiber bundle radius (r_b) does not increase the conduction velocity by increasing the threshold of ion channels. The parameter d corresponding to the fiber diameter in Eq. (7) is constant, regardless of r_b, in the reaction functions I_{ion} at the nodes of Ranvier. However, the slight change in conduction velocity by various fiber radii is indirectly caused by the change in the extracellular potential and induced electric currents.

Validation of the governing equations

This work considers the dynamics of the membrane potential propagation, which is affected by the distribution of the extracellular potential of the node of Ranvier, to validate the bidomain equations (Eqs. (23) and (24)).

Proposition 1. *The bidomain equations (Eqs. (23) and (24)) along the fiber direction at the node of Ranvier in the 1D domain are equivalent to the 1D ephaptic coupling equations, as presented in Eqs. (5) and (6) with Eq. (3).*

Proof. The 1D projections of the multidimensional governing equation in Eqs. (23) and (24) are given for the 1D domain Ω^{1d} as follows:

$$c_m \frac{\partial \phi_m(x,t)}{\partial t} = \frac{1}{r_i}\frac{\partial^2 \phi_m(x,t)}{\partial x^2} + \frac{1}{r_i}\frac{\partial^2 \phi_e(x,t)}{\partial x^2} - g_m\phi_m(x,t), \quad x \in \Omega^{1d}, \tag{36}$$

$$c_n \frac{\partial \phi_m(x,t)}{\partial t} = \frac{1}{r_i}\frac{\partial^2 \phi_m(x,t)}{\partial x^2} + \frac{1}{r_i}\frac{\partial^2 \phi_e(x,t)}{\partial x^2} - I_{ion}(x,t), \quad x \in \Omega^{1d}, \tag{37}$$

where $\sigma_i = 1/r_i$. With the conductivity ratio ρ such that $\rho = \sigma_i/\sigma_e$, Eq.(28) along the fiber at the node of Ranvier is expressed as follows:

$$\left(\frac{1+\rho}{\rho}\right)\frac{\partial^2 \phi_e(x,t)}{\partial x^2} = -\frac{\partial^2 \phi_m(x,t)}{\partial x^2}, \quad x \in \Omega^{1d}. \tag{38}$$

Substituting Eq. (38) into Eqs. (36) and (37) results in

$$c_m \frac{\partial \phi_m}{\partial t} = \frac{1}{r_i(1+\rho)} \frac{\partial^2 \phi_m}{\partial x^2} - g_m \phi_m, \quad x \in \Omega^{1d}, \tag{39}$$

$$c_n \frac{\partial \phi_m}{\partial t} = \frac{1}{r_i(1+\rho)} \frac{\partial^2 \phi_m}{\partial x^2} - I_{ion}, \quad x \in \Omega^{1d}, \tag{40}$$

which are the same as Eqs. (5) and (6) using Eq. (3). □

Moreover, the dynamics orthogonal to the fiber direction at the node of Ranvier are considered, as indicated in Eq. (32).

Proposition 2. *The extracellular potential of Eq. (32) parallel to $\mathbf{e}_2$, or orthogonal to the fiber direction $\mathbf{e}_1$ at the node of Ranvier, is equivalent to the extracellular potential for a sufficiently small axon radius a as follows [11]:*

$$\phi_e(x, d) = \frac{\sigma_i a^2}{4\sigma_e} \int_{-\infty}^{\infty} \frac{\partial^2 \phi_m / \partial x^2}{\sqrt{(x - x')^2 + d^2}} dx', \tag{41}$$

where x denotesss the axis whose tangent vector is the same as $\mathbf{e}_2$. Moreover, d represents the distance from the axon, and a indicates the axon radius.

Proof. For a constant δ, the solution to the scalar potential of $\nabla^2 \phi_e = -\delta$ along an infinite line is derived as follows [11, 12]:

$$\phi_e(x, d) = \frac{1}{4\pi} \int_{-\infty}^{\infty} \frac{\delta}{\sqrt{(x - x')^2 + d^2}} dx'. \tag{42}$$

The electric potential is obtained by setting $d \to 0$. Using Eq. (42) for Eq. (32) yields the following expression for δ,

$$\delta = \left(\frac{\pi a^2 \sigma_i}{\pi a^2 \sigma_i + \sigma_e} \right) \frac{\partial^2 \phi_m}{\partial x^2}, \tag{43}$$

where $\mathbf{e}_2 \cdot \nabla \phi_m = \partial \phi_m / \partial x$. If σ_i and σ_e are approximately constant in each fiber, the solution of Eq. (32) is derived as follows:

$$\phi_e = \frac{1}{4\pi} \left(\frac{\pi a^2 \sigma_i}{\pi a^2 \sigma_i + \sigma_e} \right) \int_{-\infty}^{\infty} \frac{\partial^2 \phi_m / \partial x^2}{\sqrt{(x - x')^2 + d^2}} dx'. \tag{44}$$

For a significantly small fiber bundle radius (r_b), to obtain Eq. (41), $a \ll 1.0$ and $\pi a^2 \sigma_i + \sigma_e \approx \sigma_e$. □

Numerical schemes with moving frames

A set of orthonormal basis vectors $(\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3)$ in three-dimensional (3D) space, called a *moving frame* [28–33], is employed for the efficient numerical solution of the bidomain equations in Eqs. (23) to (28). A moving frame can be considered a hexahedron at point P with each edge aligned along the unit tangent vectors (Fig. 4). A moving frame is denoted by $\{\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3\}$ satisfying the following orthonormality.

$$\mathbf{e}_i \cdot \mathbf{e}_j = \delta_{ij}, \quad \|\mathbf{e}_i\| = 1, \quad 1 \le i, j \le 3, \tag{45}$$

where δ_{ij} denotes the Kronecker delta. Two tangent vectors $\mathbf{e}_1$ and $\mathbf{e}_2$ lie on the tangent plane of a 2D domain Ω, whereas $\mathbf{e}_3$ is aligned along its surface normal vector,.

Fig 4. Orthonormal (orthogonal with unit length) basis vectors, called moving frames.

By introducing the capacitance ratio $d = \sqrt{c_n/c_m}$, Eqs. (23) and (24) change to

$$\frac{\partial \phi_m(\mathbf{x}, t)}{\partial t} = \frac{1}{c_n} \left[\nabla \cdot \left(d^2 \widehat{\sigma}_i \nabla \phi_m(\mathbf{x}, t) \right) + \nabla \cdot \left(d^2 \widehat{\sigma}_i \nabla \phi_e(\mathbf{x}, t) \right) \right] - \frac{g_m}{c_m} \phi_m(\mathbf{x}, t), \quad \mathbf{x} \in \Omega_m, \tag{46}$$

$$\frac{\partial \phi_m(\mathbf{x}, t)}{\partial t} = \frac{1}{c_n} \left[\nabla \cdot \left(\widehat{\sigma}_i \nabla \phi_m(\mathbf{x}, t) \right) + \nabla \cdot \left(\widehat{\sigma}_i \nabla \phi_e(\mathbf{x}, t) \right) \right] - \frac{1}{c_n} I_{ion}(\mathbf{x}, t), \quad \mathbf{x} \in \Omega_n. \tag{47}$$

The following lemma is considered using the rescaled moving frames.

Lemma 1. *In a 2D domain Ω, the diffusion operator with a conductivity tensor $\widehat{\sigma}$ is equivalently computed by rescaling the moving frames as follows:*

$$\nabla \cdot \left(d^2 \widehat{\sigma}_i \nabla f \right) = \nabla^d \cdot \nabla^d f, \tag{48}$$

where ∇^d represents the differential operator with orthogonal basis vectors $\{\mathbf{d}_j\}$ of a magnitude of d, defined such that

$$\nabla^d \cdot \nabla^d f = \nabla \cdot \left[(\nabla f \cdot \mathbf{d_1})\mathbf{d_1} + (\nabla f \cdot \mathbf{d_2})\mathbf{d}_2 \right]. \tag{49}$$

Proof. If orthogonal basis vectors $\{\mathbf{e}_1, \mathbf{e}_2\}$ correspond to the diffusivity tensor $\widehat{\sigma}_i$, then $\widehat{\sigma}_i \nabla f$ can be computed using the orthogonal basis vectors $\{\mathbf{e}_1, \mathbf{e}_2\}$ as follows:

$$\widehat{\sigma}_i \nabla f = (\nabla f \cdot \mathbf{e}_1)\mathbf{e}_1 + (\nabla f \cdot \mathbf{e}_2)\mathbf{e}_2. \tag{50}$$

Similarly, the left-hand side of Eq. (48) can be computed as follows:

$$\nabla \cdot \left(d^2 \widehat{\sigma}_i \nabla f \right) = \nabla \cdot \left[(d^2 \nabla f \cdot \mathbf{e}_1)\mathbf{e}_1 + (d^2 \nabla f \cdot \mathbf{e}_2)\mathbf{e}_2 \right]. \tag{51}$$

A scaled basis vector $\mathbf{d}_j$ is introduced such that

$$\mathbf{d}_j = d\mathbf{e}_j, \quad \text{where} \ \ 1 \le j \le 2. \tag{52}$$

Then, Eq. (51) becomes

$$\nabla \cdot \left(d^2 \widehat{\sigma}_i \nabla f \right) = \nabla \cdot \left[(\nabla f \cdot \mathbf{d}_1)\mathbf{d}_1 + (\nabla f \cdot \mathbf{d}_2)\mathbf{d}_2 \right]. \tag{53}$$

The right-hand side of Eq. (53) is the expression of the diffusion operator with the orthogonal basis vectors $\{\mathbf{d}_j\}$, which is the same as the right-hand side of Eq. (48). $\square$

By Lemma 1, the diffusion-reaction equations in Eqs. (46) and (47) and the anisotropic Poisson equation in Eq. (28) yield the following equations, called *neural bidomain equation*:

$$\frac{\partial \phi_m(\mathbf{x}, t)}{\partial t} = \frac{1}{c_n} \left[\nabla^{d_i} \cdot \nabla^{d_i} \phi_m(\mathbf{x}, t) + \nabla^{d_i} \cdot \nabla^{d_i} \phi_e(\mathbf{x}, t) \right] - I(\mathbf{x}, t), \tag{54}$$

$$\nabla^{d_e} \cdot \nabla^{d_e} \phi_e(\mathbf{x}, t) = -\nabla^{d_i} \cdot \nabla^{d_i} \phi_m(\mathbf{x}, t), \tag{55}$$

where d_i and d_e represent the magnitude of the corresponding tangent vectors $\{\mathbf{e}_j\}$ for the intracellular and extracellular conductivity tensors $\widehat{\sigma}_i$ and $\widehat{\sigma}_e$, respectively, as presented in Table 2 and Fig. 5 for each domain.

	Myelin sheath (Ω_m)	Node of Ranvier (Ω_n)	Outside the fiber (Ω_i^c)
d_i	$\sqrt{c_n/c_m}$	1.0	0.0
d_e	$\sqrt{c_n/c_m}/\rho$	$1/\rho$	α_f
I	$g_m\phi_m(\mathbf{x},t)/c_m$	$I_{ion}(\mathbf{x},t)/c_n$	0.0

Table 2. Magnitude of the tangent vectors of the moving frames for intracellular and extracellular conductivity tensorss in each domain and the corresponding electric currents (I). In addition, d_i and d_e corresponds to the intracellular and extracellular conductivity tensors $\widehat{\sigma}_i$ and $\widehat{\sigma}_e$, respectively. Moreover, ρ represents the conductivity ratio, and α_f denotes the amplified factor of the extracellular potential.

(a) Distribution of d_i

(b) Distribution of d_e

Fig 5. Distribution of the magnitude of the tangent vectors of d_i and d_e corresponding to the intracellular and extracellular conductivity tensors $\widehat{\sigma}_i$ and $\widehat{\sigma}_e$, respectively, for a single fiber with $\rho = 1$.

Weak formulation

The following weak formulation is applied in the context of the Galerkin formulation to solve the partial differential equations of Eqs. (54) and (55). For a given test function φ of the same polynomial order as the solution, the following Galerkin formulation is computed for Eq. (54):

$$\int_{\Omega} c_n \frac{\partial \phi_m}{\partial t}\varphi d\mathbf{x} = \int_{\Omega} \nabla^d \cdot \widehat{\sigma}_i \nabla^d \phi_m \varphi d\mathbf{x} + \int_{\Omega} \nabla^d \cdot \widehat{\sigma}_i \nabla^d \phi_e \varphi d\mathbf{x} - \int_{\Omega} I\varphi d\mathbf{x}. \qquad (56)$$

The boundary conditions are considered as follows. The intercellular space is distinguished from the extracellular space by the membrane, represented by the boundary $\partial \Omega_i$. The electric insulation property of the membrane is expressed by the pure Neumann boundary condition as

$$\widehat{\sigma}_i \nabla \phi_i \cdot \mathbf{n} = 0 \qquad \text{on} \ \ \partial\Omega_i, \qquad (57)$$

where $\mathbf{n}$ indicates the edge normal vector of the elements orthogonal to the boundary. The following equality, $\phi_i = \phi_m + \phi_e$, obtains the boundary condition of Eq. (57) on $\partial\Omega_i$:

$$\widehat{\sigma}_i \nabla \phi_m \cdot \mathbf{n} = -\widehat{\sigma}_i \nabla \phi_e \cdot \mathbf{n} \qquad \text{on} \ \ \partial\Omega_i. \qquad (58)$$

If the conductivity tensor of the neural fiber is significantly larger in the orthogonal direction to $\mathbf{n}$ and the fiber is sufficiently thin compared to its length, then, the external potential change in the anisotropic fiber along the normal direction is almost negligible:

$$\widehat{\sigma}_i \nabla \phi_e \cdot \mathbf{n} \approx 0. \qquad (59)$$

Therefore, the boundary condition in Eq. (58) is approximated as follows:

$$\widehat{\sigma}_i \nabla \phi_m \cdot \mathbf{n} \approx 0 \qquad \text{on} \ \ \partial\Omega_i, \tag{60}$$

Moreover, this work considers the boundary conditions of the extracellular space Ω_e, or the space surrounding the cells. The outskirt boundary of the domain Ω_e is also insulated from any influx to have the following boundary condition:

$$\nabla \phi_e \cdot \mathbf{n} = 0 \qquad \text{on} \ \ \partial\Omega_e, \tag{61}$$

where the extracellular space is assumed to be isotropic. Furthermore, the following numerical schemes are introduced to address the numerical challenges, particularly regarding *numerical stiffness*. One difficulty in computationally solving Eq. (56) is the dramatic ratio between the node and myelin sheath length. The internodal length is approximately 2,000 μ m when the nodal length is $2.5 \sim 3.183$ μ m [16] [22]. Thus, the internodal region and node ratio is approximately between 1/200 and 1/1000. A minimal node size Δx is required, affecting the time-step size for the numerical solution of the corresponding partial differential equations. The time-step size iin the context of the finite difference scheme is approximately proportional to Δx^2 [39, 40]. A high-order implicit-explicit time marching scheme is adopted to alleviate the strong restriction of the time step [41]. Eq. (56) becomes the following by considering the equation at each time.

$$\int_\Omega \phi_m(\mathbf{x}, t^n)\varphi d\mathbf{x} = \int_\Omega \phi_m(\mathbf{x}, t^{n-1})\varphi d\mathbf{x} + \frac{\Delta t}{c_n} \int_\Omega \nabla^d \cdot \widehat{\sigma}_i \nabla^d \phi_m(\mathbf{x}, t^n)\varphi d\mathbf{x}$$
$$+ \frac{\Delta t}{c_n} \left[\int_\Omega \nabla^d \cdot \widehat{\sigma}_i \nabla^d \phi_e(\mathbf{x}, t^{n-1})\varphi d\mathbf{x} - \int_\Omega I(\mathbf{x}, t^{n-1})\varphi d\mathbf{x} \right]. \tag{62}$$

The Laplacian operator for ϕ_m should be solved *implicitly* to be computed at time t^n, leading to a large global matrix that must be solved iteratively for a given time t^n. This implicit scheme allows an approximately $10 \sim 100$ times larger time step than the explicit Laplacian operator for ϕ_m. The other components, such as the Laplacian operator for ϕ_e, the reaction function, and external flow $\mathbf{I}$, are time-marched *explicitly*. The explicit Laplacian operator for ϕ_e affects the time step less significantly than that for ϕ_m, possibly due to the smooth distribution of ϕ_e.

Another stiffness problem arises for the integration of the reaction function. For the four variables for permeabilities (m, n, p, h), we have the following:

$$\frac{dm}{dt} = \alpha_m(1-m) - \beta_m, \tag{63}$$

$$\frac{dn}{dt} = \alpha_n(1-n) - \beta_n, \tag{64}$$

$$\frac{dp}{dt} = \alpha_p(1-p) - \beta_p, \tag{65}$$

$$\frac{dh}{dt} = \alpha_h(1-m) - \beta_h. \tag{66}$$

As illustrated in Fig. 2, the dynamics of ion channels demonstrate sharp variations with respect to membrane potential changes. This stiffness in the reaction functions also significantly restricts the time-step size. The Rush–Larsen scheme [42] was adopted for the numerical scheme to alleviate the strong restriction by the stiff reaction function. Applying the Rush–Larsen scheme to Eqs. (63) - (66), the solution at the ith time stage for $A = (m, h, p, h)$ is obtained as follows:

$$A(t^i) = A_\infty - [A_\infty - A(0)]e^{-\Delta t/\tau_i}, \tag{67}$$

where $A_\infty = \alpha_A/(\alpha_A + \beta_A)$ and $\tau_i = 1/(\alpha_A + \alpha_B)$.

Anisotropic Helmholtz Solver

In the numerical solution of Eqs. (54) and (54), the following 2D Helmholtz equation is solved repeatedly, for example, in the implicit time marching of Eq. (54) and the anisotropic Poisson equation for the extracellular potential in Eqs. (31) and (32):

$$\nabla \cdot d^2\widehat{\sigma}\nabla u(\mathbf{x}) - \lambda u(\mathbf{x}) = f(\mathbf{x}), \qquad \mathbf{x} \in \mathbf{\Omega}, \;\; \lambda \in \mathbb{R}^+. \tag{68}$$

This work considers one tessellated element Ω^k of a 2D2 domain Ω such that $\cap_k \Omega^k = \Omega$ and $\cup_k \Omega^k = \emptyset$. For each Ω^k, Eq. (68) is integrated by part after multiplying a test function φ of the same polynomial degree as the solution function u, obtaining

$$\int_{\Omega^k} d^2\widehat{\sigma}\nabla u^k \cdot \nabla\varphi d\mathbf{x} + \int_{\Omega^k} \lambda u^k \varphi d\mathbf{x} - \int_{\partial\Omega^k} (d\widehat{\sigma}\nabla u^k \cdot \mathbf{n})\varphi ds = -\int_{\Omega^k} f^k \varphi d\mathbf{x}, \tag{69}$$

where u^k and f^k are the projections of u and k on the element Ω^k, respectively. An alternative expression is obtained by Lemma 1 as follows:

$$\int_{\Omega^k} \nabla^d u^k \cdot \nabla^d\varphi d\mathbf{x} + \int_{\Omega^k} \lambda u^k \varphi d\mathbf{x} - \int_{\partial\Omega^k} (\nabla^d u^k \cdot \mathbf{n})\varphi ds = -\int_{\Omega^k} f^k \varphi d\mathbf{x}. \tag{70}$$

An axis ξ_d^j is defined such that it satisfies the following expression of directional derivative along $\mathbf{d}_j$, that is,

$$\mathbf{d}_j \cdot \nabla u^k = \frac{\partial u^k}{\partial \xi_d^j}, \qquad \xi_d^j \in \Omega^k. \tag{71}$$

Using Lemma 1 again to incorporate the diffusivity tensor into moving frames yields the following discretization of the anisotropic Helmholtz solver for the solution u^k in an element Ω^k:

$$\mathbb{L}u^k + \lambda\mathbb{M}u^k - \mathbb{B}u^k = -\mathbb{M}f^k, \tag{72}$$

wher

$$\mathbb{L}[m][n] = \int_{\Omega^k} \sum_{i,j} \frac{\partial\phi_m}{\partial\xi_d^i}\frac{\partial\phi_n}{\partial\xi_d^j} d\Omega^k, \tag{73}$$

$$\mathbb{M}[m][n] = \int_{\Omega^k} \phi_m\phi_n d\Omega^k, \tag{74}$$

$$\mathbb{B}[m][n] = \int_{\Omega^k} \sum_j \frac{\partial\phi_m}{\partial\xi_d^j}(\mathbf{n} \cdot \nabla\phi_n)ds. \tag{75}$$

The following test problem is employed to demonstrate the accuracy of the proposed anisotropic Helmholtz solver in Eq. (72) in a 2D domain of $(x, y) \in [-1, 1] \times [-1, 1]$. The following Helmholtz equation with an tensor $\hat{\sigma}$ is considered:

$$\nabla \cdot \widehat{\sigma}\nabla u = -(\sigma_1 + \sigma_2)\omega^2 \sin(\omega x)\cos(\omega y), \tag{76}$$

(a) Solution **(b)** Error convergence

Fig 6. Anisotropic Helmholtz solver for a test problem with (a) the solution, and (b) error convergence for various σ_1 and σ_2.

where the conductivity tensor is given by

$$\widehat{\sigma} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} \sigma_1 & 0 \\ 0 & \sigma_2 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & 1 \\ 0 & 0 \end{bmatrix}^T. \tag{77}$$

The exact solution of this test problem is $u = \sin(\omega x)\cos(\omega y)$. In addition, Fig. 6 displays the exact solution with $\omega = \pi$ and the exponential error convergence for the isotropic case with unit conductivity ($\sigma_1 = \sigma_2 = 1.0$), the isotropic case with nonunit conductivity ($\sigma_1 = \sigma_2 = 4.0$), and the anisotropic case ($\sigma_1 = 0.5, \sigma_2 = 4.0$).

Results

The computational results were obtained by numerically solving the proposed multidimensional neural spike propagation model in Eqs. (54) to (55). The numerical scheme is given as a weak formulation with moving frames in the context of the continuous Galerkin formulation, as presented in Eqs. 56 with the boundary conditions in Eqs. (60) to (61). The scheme was implemented in the open-source spectral/hp element framework, called *Nektar++* [43]. 1D data are extracted along a line from 2D simulations to analyze the propagation along the fiber, particularly regarding arriving time and conduction velocity. Moreover, this work introduced the *arrival time map* (i.e., *time map*) denoted by $\mathcal{T}(\mathbf{x})$ for $\mathbf{x} \in \Omega$, for the comparison and analysis of conduction speed and synchronization. The time map, corresponding roughly to the time when the neural spike is excited, is computed as follows [44]:

$$\mathcal{T}(\mathbf{x}) = \left[\int t \left(\frac{d\phi_m}{dt} \right)^+ dt \bigg/ \int \left(\frac{d\phi_m}{dt} \right)^+ dt \right], \quad \mathbf{x} \in \Omega, \tag{78}$$

where the superscript $+$ denotes the maxmod function defined by

$$A^+ = maxmod\{A, 0\} = \begin{cases} A, & \text{if } A > 0, \\ 0, & \text{otherwise.} \end{cases} \tag{79}$$

In this paper, the width of the intracellular domain, regarded as a fiber bundle with identical fibers as a small fascicle, is primarily 0.01 cm in most cases. Considering that the fiber radius is approximately 2 μ m, the corresponding bundle with a width of

(a) Two-dimensional mesh of a single fiber

(b) Potential distribution at $t = 0.1$ ms (contour: ϕ_m, color: ϕ_e)

(c) Propagation along an unmyelinated axon **(d)** Propagation along a myelinated axon with nodes of Ranvier

Fig 7. Single-fiber propagation and comparison of neural spikes in the neural fiber in unmyelinated and myelinated axons with nodes of Ranvier. The unit of time is milliseconds and the length is centimeter. (a) The myelin sheath is marked gray, and the four nodes of Ranvier are slightly darker,
$(x, y) = [0.02, 0.03] \times [0.02 + k * 0.2, 0.03 + k * 0.2]$ for $0 \leq k \leq 3$. $x_0 = 0.01$, $y_0 = 0.01$, $x_L = 0.0$, $x_R = 0.05$, $y_L = 0.0$, $y_R = 0.65$, (b) Potential distribution at $t = 0.4$ ms initiated from the leftmost node. contour: ϕ_m, color: ϕ_e. (c) Simulated on a fiber of length 0.65 cm. (d) Simulated on the fiber with 20 nodes and a length of 4.1 cm, where $T = 24°C$, $\rho = 1.0$, and $r_b = 0.1$ cm.

0.01 cm contains around a thousand neural fibers for a packing ratio of 0.5. In the proposed mathematical model, ephaptic coupling between fiber bundles is only effective when ephaptic coupling within each fiber bundle is ignored. Thus, the conduction velocity of the neural spike along the fiber bundle is approximately the same as that of a single fiber bundle. This conduction velocity contrasts the fact that the conduction velocity in an ephaptically coupled fiber bundle slows [10, 22]. The mathematical models in Eqs. (54) to (55) can be further modified to include this slowing effect in a fiber bundle. However, this model was not considered in this paper because it is beyond the scope of work.

A. Single fiber bundle

This work considers a single fiber bundle embedded in a 2D plane, as presented in Fig. 7a. The domain of the fiber bundle is given as a rectangular region $(x, y) \in [x_0, x_0 + w] \times [y_0, y_0 + L]$, where w denotes the fiber width at 0.01 cm, and L represents the fiber length varying from 1 to 4 cm. Each node and myelin length is the same at 0.01 and 0.2 cm, respectively. Temperature is $24°C$, and conductivity ratio rho is 1. The neural spike is initiated at the left-most node for the duration of 0.01 ms with

a continuous external 0.4 mV stimulus. Without myelin, the shape of the neural spike is almost linear in the depolarization and repolarization phases, as depicted in Fig. 7c, coincident with the findings of previous studies [13, 22]. The conduction velocity in an unmyelinated axon is close to 2.5 m/s, consistent with the previously known value [16].

Fig 8. Distribution of the membrane potential (ϕ_m) and extracellular potential (ϕ_e) perpendicular to the propagation direction (along the x-axis). Time t is in milliseconds. Measured at a node of Ranvier, wherew $T = 24°C$, $\rho = 1.0$, and $r_b = 0.1$ cm.

The myelinated sheath increases the neural spike duration by more than 10 times larger than that of the unmyelinated axon. The wavelengths in the unmyelinated and myelinated axons with the nodes of Ranvier are approximately 0.2 cm and 2.2 cm, respectively. Moreover, the conduction velocity in the myelinated axon with nodes of Ranvier is also increases by more than 10 times that of the unmyelinated axon (i.e., approximately 25 m/s) in accordance with the measured quantity of a frog motor axon [45] and the corresponding computational simulations of previous studies [13, 22]. In Fig. 7d, the most striking feature of the neural spike for the myelinated axon is the dramatic potential change at the nodes of Ranvier, in contrast to slower changes in the myelin sheath. The reaction of the ion channels causes this staircase repolarization phase only at the nodes of Ranvier. The highest magnitude of the action potential for unmyelinated and myelinated axons remains approximately the same.

Furthermore, Fig. 8 displays the time sequential distribution of the extracellular potential (ϕ_e) distribution perpendicular to the propagation direction, i.e., along the x-axis. The membrane potential increases for the initial depolarization ($t = 0.05$ ms). Accordingly, the extracellular potential inside the axon increases ($t = 0.075$ ms). After the extracellular potential reaches a certain threshold, for example, approximately around 10 mV for a fiber radius of 0.1 cm ($t = 0.1$ ms), the extracellular potential becomes approximately zero when the membrane potential is approximately 60 mV ($t = 0.12$ ms). The extracellular potential becomes negative when the membrane potential is between 80 and 120 mV ($t = 0.15$ ms). Finally, it returns to zero when the membrane potential reaches the maximum ($t = 0.18$ ms). This extracellular potential fluctuation is consistent with the mathematical predictions for the oscillation between hyperpolarization and depolarization deduced by Markin in the 1970s [18–21].

Moreover, Fig. 9 displays the conduction velocity changes according to the

(a) Instantaneous velocity   **(b)** Average velocity

Fig 9. Conduction velocity along a single fiber bundle. The vertical dotted black line corresponds to the boundaries of the nodes of Ranvier, where ρ denotes the conductivity ratio between the intracellular and extracellular domains, $T = 24°C$, and $r_b = 0.1$ cm.

conductivity ratio ρ variation between the intracellular and extracellular space. In addition, Fig. 9a displays the variations of the instantaneous conduction velocity of a single neural spike initiated at the left-most node using the instantaneous velocity from the arrival time [44]. The contribution of the extracellular potential is proportional to the inverse of the square of the distance from the nodes of Ranvier, as deduced from Eq. (35), or deduced in [10,11]. Thus, the instantaneous velocity varies from $10 \sim 60$ (m/s) for the corresponding value of ρ as $\{1, 2, 4, 8, 12\}$.

The average velocity of the neural spike propagation along the myelinated axon with the nodes of Ranvier is obtained by averaging the instantaneous velocity, as illustrated in Fig. 9b. The average velocity depends on ρ, coincident with the result from [22]. Moreover, the average conduction velocity is approximately proportional to the square root of ρ. For example, the average velocity with $\rho = 1$ is between 36 and 52 m/s, approximately three times larger than the average velocity for $\rho = 12$ when the conduction velocity is between 13 and 20 m/s.

Fiber radius (cm)	0.01	0.1	0.2	0.4	0.8
Conduction velocity (m/s)	29.0	29.3	29.5	29.5	29.4
Temperature $(°C)$	24	28	32	36	40
Conduction velocity (m/s)	29.3	45.3	70.3	108.8	168.3

Table 3. Conduction velocity dependency along a single fiber. For, the conduction velocity versus the fiber radius, $T = 24°C$. For the conduction velocity versus temperature, r_b=0.1 cm. ρ=1.0.

Moreover, Table 3 demonstrates that the average conduction velocity depends on the fiber radius and temperature. In the mathematical model of Eq. (28) using the approximation of Eq. (35), the fiber radius does not significantly change the conduction velocity for a single fiber bundle. The induced extracellular potential primarily causes a slight change in the conduction velocity. The proposed mathematical model does not consider the direct conduction velocity change in I_{ion} by the fiber radius, as elaborated after Eq. (35), causing this non-dependency. In contrast, the increase in temperature directly yields a dramatic increase in the conduction velocity. Temperature is a critical component in the function $\mathcal{Z}$ in Eq. (12), affecting the threshold of ion channels. This computational simulation is consistent with the physiological observation in [46]. Table 3 reveals that the conduction velocity linearly increased up to $30°C$ but increased

more rapidly for higher temperatures when $\rho = 1$ [22]. 430

(a) Mesh of two parallel myelinated fiber bundles with five nodes of Ranvier.

(b) Potential distribution at $T = 0.4$ ms. contour: membrane potential, color: extracellular potential

(c) ϕ_e along the x-axis **(d)** ϕ_e along the y-axis

Fig 10. Extracellular potential distribution along each axis for two fiber bundles with aligned nodes of Ranvier. The first fiber bundle is in $x \in [0.01, 0.02]$ and the second is in $x \in [0.03, 0.04]$, where $T = 24°C$ and $\rho = 1.0$.

B. Fiber bundles with aligned nodes of Ranvier 431

This work considers two straight and parallel fiber bundles with aligned node of 432
Ranvier, as depicted in Fig. 10a. The two fiber bundles with the same width (0.01 cm) 433
are aligned along the y-axis with a gap of $w = 0.01$ cm between them along the x-axis. 434
The neural spike propagation is initiated at the nodes at $y \in [0.01, 0.02]$ for the first 435
fiber bundle in $x \in [0.01, 0.02]$ and the second in $x \in [0.03, 0.04]$. 436

First, the two fiber bundles are initiated simultaneously in the first node of each 437
fiber bundle. The propagation along the first and second fiber bundles is approximately 438
the same, with an arrival time error of 10^{-5} ms. Similar to the propagation in a single 439
fiber bundle, the strength of the extracellular potential ϕ_e also increases as the fiber 440
bundle radius increases, as illustrated in Fig. 10c. For example, the extracellular 441
potential between the fiber bundles with a radius 0.8 cm is approximately 3 mV when 442
the extracellular potential in each fiber bundle is approximately 9 mV. This potential is 443
significantly greater than the extracellular potential between the fiber bundles with a 444
radius of 0.1 cm, approximately 0.216 mV. Moreover, Fig. 10d demonstrates that the 445
extracellular potential changes drastically after depolarization where the magnitude is 446
approximately proportional to the fiber bundle radius. The increase in the propagation 447
velocity caused by the fiber bundle radius is relatively insignificant to other factors, 448
such as ρ and temperature. 449

In Fig. 11, the propagation in the myelin sheath of the two fiber bundles is faster 450
than that in the single fiber bundle, regardless of the conductivity ratio (ρ) and fiber 451
bundle radius (r_b). The difference in the conduction velocity between these two fiber 452

(a) Instantaneous velocity (m/s) versus conductivity ratio (ρ)

(b) Instantaneous velocity (m/s) versus fiber bundle radius (r_b))

Fig 11. Instant conduction velocity in two fiber bundles with simultaneous initiations. (a) Instant velocity along the first fiber bundle for various conductivity ratio ρ. $r_b = 0.1$ cm. *Double* and *single* correspond to the velocity for two and a single fiber, respectively. (b) Instant velocity along the first fiber bundle for various fiber bundle radius r_b ($\rho = 1.0$). The two fiber bundle' radii are 0.01 cm, and the gap between them is 0.01 cm ($T = 24°C$).

bundles is small for $r_b = 0.01$ cm, as illustrated in Fig. 11b; thus, this is likely caused by ephaptic coupling effects. Contrary to the conduction velocity in the myelin sheath, the conduction velocity in the nodes of Ranvier depends on the ephaptic coupling effects and the conductivity ratio ρ. Table 4 demonstrates that the conduction velocity at the nodes of the two fiber bundles is slower than that at the nodes of the single fiber bundle for a smaller ρ, less than about $\rho = 3.0$. The conduction velocity at the nodes of Ranvier of two fiber bundles with a ρ value larger than 3.0 is faster than that for a single fiber bundle. Hence, the collective average velocity in two fiber bundles is slower than that of a single fiber bundle for a ρ value less than 3.0. If ρ is higher than 3.0 with stronger ephaptic coupling effects, the propagation in two fiber bundles is faster than that of a single fiber bundle. The slowing of the propagation in the fiber bundles by ephaptic coupling for a low ρ value agrees with previous 1D studies [10, 14, 22]. However, the increase in speed of the conduction velocity for two fiber bundles for a large value of ρ was unknown.

This work considers the propagation of the two fiber bundles with a delayed initiation in one fiber bundle, where the initiation of the second fiber bundle is delayed without loss of generality. Both Fig. 12 and Table 5 demonstrate the effects of ephaptic coupling on unsynchronized propagation, agreeing with previous mathematical and computational studies [11, 22]. in Fig. 12a, synchronization continuously occurs between the two fiber bundles when the initiation delay is less than about 0.1 ms. This time is approximately when the neural spike in the first fiber leaves the first node ($y \in [2.2, 2.3]$) and moves into the second myelin ($y \in [2.3, 4.3]$).

Moreover, Fig. 12b confirms that synchronization is caused by the increased velocity of the delayed neural spike in the second fiber bundle while the velocity of the first fiber bundle remains approximately the same. The average velocity of the second fiber is at maximum when the initial delay is $0.04 \sim 0.08$ ms and is at minimum for the initial delay of more than 0.1 ms or less than 0.01 ms. The synchronization velocity and ratio are defined as follows to quantify the rate of each synchronization:

$$\text{Synchronization velocity} = \frac{d\mathcal{T}_{diff}}{ds}, \tag{80}$$

where $\mathcal{T}_{diff}$ denotes the arrival time difference between the fiber bundles, and s

ρ (conductivity ratio)		1	2	4	8	12
	No ephaptic			5.366		
Node velocity (m/s)	Two fiber bundles	3.661	2.804	2.096	1.543	1.572
	Single fiber bundle	3.723	2.795	2.068	1.542	1.481
	Difference (*)	-0.0618	0.0089	0.028	0.0007	0.0918
	No ephaptic			36.188		
Avg. velocity (m/s)	Two fiber bundles	26.005	22.246	18.637	15.341	13.482
	Single fiber bundle	26.245	22.352	18.614	15.213	13.272
	Difference (*)	-0.24	-0.11	0.023	0.13	0.21

Table 4. Conduction velocity at the fourth node and the total average velocity in the myelin and nodes for the propagation 1) without ephaptic coupling (no ephaptic), 2) with two fiber bundles, and 3) with a single fiber bundle. (*) Difference is measured as the conduction velocity of two fibers minus the conduction velocity of a single fiber bundle.

represents the parameter of the propagation distance. Similarly, the relative synchronization velocity is defined by dividing Eq. (80) by the maximum arrival time difference. Table 5 reveals that the synchronization velocity is linearly proportional to the magnitude of the initial delay, particularly up to a delay of 0.05 ms. This synchronization proportionality to the initial delay provides a constant synchronization ratio when the initial delay is less than 0.05 ms.

Delay (ms)	0.01	0.02	0.04	0.08	0.1	0.11
Sync. velocity (s/cm)	2.48	4.76	7.55	4.77	0.86	-0.93
Relative sync. velocity (%/cm)	25.8	24.5	19.8	6.3	0.9	-0.9

Table 5. Synchronization velocity and relative synchronization velocity defined in Eq. (80) for the two fiber bundles with aligned nodes of Ranvier. The initiation delay occurs in the second fiber bundle of $y = \in [0.03, 0.04]$. The two fiber bundles have a width of 0.01 cm and length of 2.2 cm with a gap of 0.01 cm between them, where $\rho = 1$, $r_b = 0.1\ cm$, and $T = 24°C$.

This work also considers three fiber bundles with aligned nodes of Ranvier, as illustrated in Fig. 13. The configuration of the fiber bundles is the same as the case for two fiber bundles. The three fiber bundles are aligned along the y-axis, and each domain is defined as follows: $y \in [0.01, 0.02]$, $y \in [0.03, 0.04]$, and $y \in [0.05, 0.06$ in that order. The neural spike initiate at the same location in the three fibers but with a slight delay in the second fiber bundle. The extracellular potential distribution for the initial delay of 0.04 ms in the second fiber bundle is displayed in Fig. 13c. The synchronization in the three fiber bundles is slightly faster than for the two for a shorter delay of approximately less than 0.05 ms, as presented in Table 6. However, synchronization is significantly slower or fails when the delay is longer than 0.08 ms, which is less than that for the two fiber bundles. In this paper, synchronization in similar conditions is reported for a delay of 0.5 ms in [22] for a packing ratio of 0.25, in contrast to a ratio of 0.5 for the configuration of the three fiber bundles. The packing ratio represents the ratio between the extracellular and intracellular space in a fiber bundle. This mechanism is likely due to the symmetric extracellular potential distribution. This

(a) Arrival time difference between two fiber bundles

(b) Arrival time of the second fiber bundle.

Fig 12. Propagation along two fiber bundles with an initiation delay. (a) Arrival time difference between the two fiber bundles when the second fiber bundle is initiated with various delays. (b) Conduction velocity of the second fiber bundle increases for synchronization, whereas the conduction velocity of the first fiber bundle remains about the same.

Delay (ms)	0.01	0.02	0.04	0.08	0.1	0.12
Sync. velocity (s/cm)	2.45	5.32	8.82	2.93	-0.23	-4.83
Relative sync. velocity (%/cm)	26.3	28.0	22.8	3.8	-0.2	-4.1

Table 6. Absolute and relative synchronization velocity defined in Eq. (80) for the three fiber bundles with aligned nodes of Ranvier. The initiation delay occurs in the second (middle) fiber bundle of $y = \in [0.03, 0.04]$. The threes fiber bundles have a width of 0.01 cm and length of 1.1 cm with a gap of 0.01 cm between them, where $\rho = 1$, $r_b = 0.1$ cm, and $T = 24°C$.

potential distribution increases the synchronization within a shorter delay but lowers   497
the delay threshold for synchronizing three fiber bundles.   498

C. Interference: Opposite-traveling neural spikes   499

The first simulation, which is only available in a multidimensional model, is the   500
interference between the traveling spikes in different directions. As an example of   501
interference, the neural spike propagations in two parallel but opposite-propagating   502
fiber bundles are simulated; one propagation in the normal (orthodromic) direction   503
along one fiber bundle and the other in the opposite (antidromic) direction along the   504
other fiber bundle. The configuration of the two fiber bundles is the same as that in the   505
previous section but with different initiation nodes. The first and second fiber bundles   506
initiate from the left-most ($y \in [0.01, 0.02]$) and right-most ($y \in [1.08, 1.09]$) nodes,   507
respectively. Further, Figs. 14b to 14d display the 1D distribution of the membrane and   508
extracellular potential at various time before and after the two spikes are closest.   509

The membrane potential in one fiber bundle does not directly affect the other   510
because each fiber domain is separated. However, the extracellular potential   511
distribution in the extracellular space is abruptly changed by interference. The   512
interference has two distinctive patterns, depending on when the two neural spikes are   513
closest; the first is when the two spikes are closest at a node and the second is when the   514
two spikes are closest at the myelin sheath.   515

The first case can be represented by two opposite-traveling fibers with an even   516

(a) Mesh of three parallel fiber bundles with aligned nodes of Ranvier

(b) Potential distribution at $T = 0.4$ ms. contour: membrane potential, color: extracellular potential

(c) ϕ_e distribution for an initiation delay of 0.04 ms in the second fiber bundle

(d) Synchronization for the initiation delay of the second fiber bundle

Fig 13. Propagation in the three neighboring fiber bundles. The first, middle, and third fiber bundles are located in $x \in [0.01, 0.02]$, $x \in [0.03, 0.04]$ and $x \in [0.05, 0.06]$, respectively. (a) Mesh of the three fiber bundles with the same thickness of 0.01 cm and a gap of 0.01 cm between them. (b) Distribution of the extracellular space for a delayed initiation of the middle fiber bundle. (c) Synchronization of neural spike propagation in the three-parallel fiber bundles for an initiation delay in the second fiber bundle.

number of nodes, as depicted in Fig. 10a. The two neural spikes are closest in the middle myelin when the spikes are initiated from opposite ends. Moreover, Fig. 14 displays the membrane and extracellular potential distribution of two opposite-traveling neural spikes. Without opposite-traveling spikes, the extracellular potential of the first fiber bundle is zero ahead of the depolarizing area. If the opposite-traveling spike occurs ahead of the depolarizing area, the extracellular potential distribution follows the same distribution as that of the second fiber bundle (Fig. 14b). This deformation of the extracellular potential ahead of the depolarization area generates an additional electric current to modify the electric current in the fiber. The extracellular potential gradients are presented in $y \in [0.6, 0.8]$, $[0.2, 0.6]$, and $[0.0, 0.2]$ for T=0.2, 0.4, and 0.6 ms, respectively. The extracellular potential gradients caused by the interference seem to propagate backward. This extracellular potential gradient induces the corresponding membrane potential changes, as depicted in Figs. 14b to 14d.

The second case is represented well by two opposite-traveling fibers with an odd number of nodes. When the spikes are initiated from opposite ends, the two neural spikes are closest to a node of Ranvier, as illustrated in Fig. 15b. Similar to fibers with an odd number of nodes, the extracellular potential gradient by interference emerges in $y \in [0.6, 0.8]$ and $[0.4, 0.8]$, and approximately 0.3 for $T = 0.3$, 0.5, and 0.7 ms, respectively. Compared to the fiber with an even number of the nodes of Ranvier, the

(a) Potential distribution at $T = 0.2$ ms. contour: membrane potential, color: extracellular potential

(b) $T = 0.2$ ms **(c)** $T = 0.4$ ms **(d)** $T = 0.6$ ms

Fig 14. Propagation along the two fiber bundles with six nodes of Ranvier in the normal and opposite directions. The mesh is the same as in Fig. 10a. The first fiber in $x \in [0.01, 0.02]$ is initiated at $y \in [0.01, 0.02]$ and the second fiber in $x \in [0.03, 0.04]$ is simultaneously initiated at $y \in [1.06, 1.07]$, where $\rho = 1$, $r_b = 0.1$ cm, and $T = 24°C$.

extracellular potential deformation is more substantial; thus, the membrane potential change in each fiber is more significant. Nevertheless, the deformation region seems smaller than that for fibers with an even number of nodes of Ranvier. The effect of this kind of interference in the fiber with an odd number of nodes of Ranvier appears reasonable because the two neural spikes remain closest at the neighboring nodes of Ranvier, with a stronger but narrower interference with the extracellular potential of other spikes.

Ephaptic coupling of traveling neural propagations depends on when the two neural spikes are closest. Thus, the configuration of the fiber bundle or the initiation delay can affect the interference magnitude. For example, a variation in the delay of the second fiber bundle can generate changes in the propagation velocity of the first fiber by interference, significantly modifying its arrival time. In addition, Fig. 16a confirms that the conduction velocity of the first fiber bundle is approximately proportional to the delay of the second fiber, up to 0.05 ms. Moreover, the fiber radius also affects the interference because the change in the fiber bundle radius yields a change in the magnitude of ephaptic coupling. Further, Fig. 16b confirms that the larger ephaptic coupling effect does not always yield a more significant change in the arrival time and conduction velocity. Significant slowing and greater interference occur at $r_b = 0.2$ cm.

D. Asynchronizing configuration: Misaligned nodes of Ranvier

The second available case using the multidimensional bidomain model is neural spike propagation in the fiber bundles with misaligned nodes of Ranvier. This work considers two parallel fibers with misaligned nodes of Ranvier with the first and second fiber bundle in $x \in [0.01, 0.02]$ and $x \in [0.03, 0.04]$, respectively, as illustrated in Fig. 17a and 17b. The nodes of Ranvier of the first fiber bundle are not aligned with those of the second fiber bundle but alternate every 0.1 cm, except the first and last nodes. For example, the third nodes of the first and second fiber bundles appear at $y \in [0.23, 0.24]$ and $[0.33, 0.34]$, respectively. The total length of the myelin and fiber is the same for the

(a) Mesh of two parallel fiber bundles with seven nodes of Ranvier

(b) $T = 0.3$ ms   **(c)** $T = 0.5$ ms   **(d)** $T = 0.7$ ms

Fig 15. Propagation along the two fiber bundles with seven nodes of Ranvier in the normal and opposite directions. The mesh is the same as in Fig. 10a. The first fiber bundle in $x \in [0.01, 0.02]$ is initiated at $y \in [0.01, 0.02]$, and the second fiber bundle in $x \in [0.03, 0.04]$ is simultaneously initiated at $y \in [1.27, 1.28]$, where $\rho = 1$, $r_b = 0.1$ cm, and $T = 24°C$.

two fibers.

If two fiber bundles are initiated simultaneously at the first nodes in each fiber bundle, without the ephaptic coupling effect, the neural spike reaches its final node simultaneously, or with a small delay as small as 7.63×10^{-4} ms for a fiber with a length of 2.22 cm with 10 misaligned nodes, represented as the dashed red line at 2.21 cm in Fig. 17c. This identical arrival time is due to the same length of the myelin sheath and nodes of Ranvier. With the ephaptic coupling effect, the neural spike in each fiber bundle reaches its final node at different arrival times. The arrival time difference between the two fibers is 0.0202 ms for the same fiber bundle, represented as the solid black line at 2.21 cm in Fig. 17c. This arrival time difference is significant, approximately 26.5 times larger than the propagation without ephaptic coupling.

The neural spike in the first fiber bundle initially propagates faster than that in the second fiber bundle, as presented in Fig. 17b. This distinctive propagation difference is generated by the second myelin in $y \in [0.13, 0.34]$ of the first fiber bundle, which is twice as lengthy as the second myelin in $y \in [0.13, 0.23]$ of the second fiber bundle. The advanced propagation in the first fiber bundle constantly slows the lagged propagation via ephaptic coupling in the second fiber bundle. Ephaptic coupling dramatically changes the instantaneous conduction velocity in the two fiber bundles with misaligned nodes, compared to a single fiber bundle propagation, as confirmed in Fig. 17d. The last myelin in the second fiber bundle is twice as long as the last myelin in the first fiber bundle, making the total length the same.

Nevertheless, the modified initial propagation velocity by the various myelin lengths produces a different arrival time for both fiber bundles at each final node. If the fiber is propagated reversely from the last node of Ranvier, the neural spike in the second fiber bundle would always be faster than that of the first. The direction of the propagation determines which neural spikes propagates faster in the same two fiber bundles. However, in both directions, the synchronization between these two misaligned fiber bundles is almost impossible unless ephaptic coupling is negligible.

Moreover, this work considered a small delayed initiation in either the first or second

(a) Arrival time difference for various delays of the second fiber bundle

(b) Arrival time difference for various radii of the fiber bundles

Fig 16. Arrival time difference of the first fiber bundle against (a) the delay time for the second fiber bundle and (b) the fiber bundle radius. The fiber bundles have an even number of nodes. The arrival time difference is computed as the arrival time for two opposite-traveling fiber minus the arrival time for a single fiber. The mesh is the same as in Fig. 10a. The first fiber bundle in $x \in [0.01, 0.02]$ is initiated at $y \in [0.01, 0.02]$, and the second fiber bundle in $x \in [0.03, 0.04]$ is initiated at $y \in [1.06, 1.07]$, where $\rho = 1$, $r_b = 0.1$ cm, and $T = 24°C$.

fiber bundle of 0.01 ms for each fiber bundle. When the first fiber bundle is delayed by 0.01 ms, Fig. 18a demonstrates that the arrival time difference between the fiber bundles is 0.0147 ms. The arrival time difference is 0.0055 ms shorter than that by the simultaneous initiation. Only a fraction of the delay of the first fiber bundle of 0.01 ms is reflected in the arrival time difference. If the neural spike initiation in the second fiber bundle is delayed by 0.01 ms, the arrival time difference between the fiber bundles is 0.0246 ms, 0.0043 ms longer than that via the simultaneous initiation. Similar to the delay of the first fiber bundle, only a fraction of that of the second fiber bundle of 0.01 ms is reflected in the arrival time difference. Thus, the magnitude of the initiation delay is not linear to the arrival time difference in a misaligned fiber. This nonlinear phenomenon seems to be due to the dramatically changed conduction velocity owing to the complexity of ephaptic coupling between the nodes of Ranvier, as illustrated in Figs. 18b and 18c.

E. Curvature effect: Two curved fiber bundles

The final case in the multidimensional bidomain model is the neural spike propagation in curved fiber bundles. This work considers two curved fiber bundles embedded in a rectangular 2D domain of $(x, y) \in [0, 1] \times [0, 1]$. The curved fiber bundle is a circular shell with a constant radius r, centered at $[0, 0]$, and the fiber width is significantly small. Thus, the fiber curvature is approximately constant. For convenience, the polar coordinate system (r, θ) is introduced to represent the fiber bundle geometry, where θ advances counter-clockwise. The first fiber bundle lies in $r \in [0.81, 0.82]$ and $\theta \in [0.075, 1.388]$, whereas the second fiber bundle lies in $r \in [0.83, 0.84]$ and $\theta \in [0.075, 1.388]$. The node and myelin lengths are similar to the previous linear fiber bundles, approximately 0.2 ms and 0.01 ms, respectively. The initial excitation is assumed to start at the nodes of $\theta = [0.0561, 0.0686]$ in both fiber bundles.

The extracellular potential distribution of the curved fiber bundles is not linear, as presented in Fig. 19. A Poisson distribution with a few sources at the nodes along the curved fiber bundle generates a unique extracellular potential distribution. The distribution at T=0.6 ms is striking because the extracellular potential distribution

(a) Mesh of two misaligned fibers.

(b) Potential distribution at $T = 0.4$ ms. contour = membrane potential, color = extracellular potential

(c) Arrival time difference between two fiber bundles

(d) Instant velocity compared to a single fiber bundle

Fig 17. Arrival time and instant velocity differences in fiber bundles with misaligned nodes of Ranvier. The fiber bundle configuration is the same as before, except for the misaligned and alternating nodes of Ranvier. (a) Mesh of the misaligned nodes of Ranvier (dark square). Fiber bundles 1 and 2 belong to the domain of $x \in [0.010.02]$ and $x \in [0.03, 0.04]$, respectively. (b) Membrane and extracellular potential distribution at $T = 0.4$ ms. (c) Significant arrival time difference (0.0202 ms) via ephaptic coupling at the final node of Ranvier, compared to the negligible time difference (7.63×10^{-4} ms) at the final node of Ranvier via nonephaptic propagation. (d) Instant velocity difference for each fiber bundle displays complex velocity changes via ephaptic coupling in fiber bundles with misaligned nodes, where $\rho = 1$, $r_b = 0.1$ cm, and $T = 24°C$.

affects the membrane potential in the backward part of the propagation in the fiber bundle. The nonlinear distribution of the extracellular potential deforms the propagation features of its own and neighboring fiber bundles's. This distribution contrasts the extracellular potential of the linear fiber bundles, which is approximately linear and orthogonal to the membrane and boundary (Fig. 20).

Next, Fig. 21 demonstrates the arrival time and conduction velocity difference between the two fiber bundles, compared to the propagations in a single fiber bundle and the propagation without ephaptic coupling. In Fig. 21a, the single curved fiber bundle with ephaptic coupling (blue line) displays a similar conduction velocity along each fiber bundle. The slight difference in the arrival time for the same θ value is caused by the slight radius difference. The two fiber bundles with ephaptic coupling (black line) illustrate the decreased arrival time difference, indicating that the second fiber with a slightly larger radius has a higher conduction velocity. This phenomenon is coincident with ephaptic coupling for linear fiber bundles. In addition, Fig. 21b displays the instantaneous conduction velocity difference between the two fiber bundles. This figure confirms that the second fiber bundle speeds up during propagation along curved fiber

(a) Arrival time difference by initiation delays

(b) Instant velocity difference by the delay of the first fiber bundle

(c) Instant velocity difference by the delay of the second fiber bundle

Fig 18. Arrival time and instant velocity of the misaligned fibers with a delay of 0.01 ms for either fiber bundle. (a) Arrival time difference by the delay of the first and second fiber bundle 0.01 ms. (b) Instant velocity changes of the two fiber bundles by the delay of the first fiber bundle 0.01 ms, compared to a single fiber bundle propagation. (c) Instant velocity changes in the two fiber bundles by the delay of the second fiber bundle 0.01 ms, compared to a single fiber bundle propagation where $\rho = 1$, $r_b = 0.1$ cm, and $T = 24°C$.

bundles. Therefore, the synchronized neural spikes along curved fiber bundles maintain synchronization at the first nodes, although the total length of the fiber bundles differs.

Discussion

This paper proposes a multidimensional bidomain model for neural spike propagation to study the ephaptic coupling effects in a multidimensional complex domain. In contrast to the 1D differential equations for neural spike propagation studies, the proposed model is expressed as partial differential equations for the general domain of multi-dimensional space. Orthonormal basis vectors (moving frames) are employed in the numerical scheme to tackle the two separate domains for the membrane and extracellular potential. Computational simulations in 2D domains for the known results are provided to validate the proposed model and confirm the known results in a 1D domain. Moreover, additional simulations are provided on three new cases which are only possible in a multidimensional model; interference by opposite-traveling neural spikes, propagation in misaligned neural fiber bundles, and curved fiber bundles.

The proposed model and corresponding numerical scheme have the following limitations. First, the ratio between the width and length of a single neural fiber is significantly large, such that the simulation of a single fiber is extremely expensive or impossible, due to numerous elements representing the geometry. As a solution, the proposed model instead fits a fiber bundle with a width of more than about 0.01 cm, which contains tens of thousands of identical neural fibers, although the strength of the extracellular potential should be adjusted accordingly. Second, the computational cost of solving the proposed model is relatively high, even in the domain of a fiber bundle with a reasonable number of elements. This high computational cost is caused by the numerical stiffness in (a) the ion channel reaction function, (b) the Laplacian operator with strong anisotropy, and (c) the repetitive anisotropic Poisson solver in the bidomain equation. The two fiber bundles with a length of 1.1 cm with approximately 500 triangular elements take about 50 hours for up to 1 ms at a workstation with a single

(a) $T = 0.2$ ms **(b)** $T = 0.4$ ms **(c)** $T = 0.6$ ms

Fig 19. Extracellular potential distribution for the propagation in the domain embedding for two curved fiber bundles. (a) Extracellular potential distribution at (a) $T = 0.2$ ms, (b) $T = 0.4$ ms, (c) $T = 0.6$ ms. The neural spike initiates at $\theta \in [0.0561, 0.0686]$, where $\rho = 1$, $r_b = 0.1$ cm, and $T = 24°C$.

(a) $T = 0.2$ ms **(b)** $T = 0.4$ ms **(c)** $T = 0.6$ ms

Fig 20. Extracellular potential distribution for the propagation in the domain embedding for two fiber bundles. (a) Extracellular potential distribution at (a) $T = 0.2$ ms, (b) $T = 0.4$ ms, (c) $T = 0.6$ ms. The neural spike initiates at $y \in [0.01, 0.02]$, where $\rho = 1$, $r_b = 0.1$ cm, and $T = 24°C$.

central processing unit (CPU). A parallel CPU computation reduces the computational time inversely proportional to the number of CPUs. Further model reduction or 3D–1D modeling seem necessary to enhance computational efficiency, possibly for full 3D simulations of real fiber bundles of the human brain.

Acknowledgments

The research of the first author is supported by the National Research Foundation of Korea (NRF-2021R1A2C109297811). The research of the second author is supported by JSPS Grant-in-Aid for Scientific Research (24K06852), JST CREST (JPMJCR1914), and Keio University (Fukuzawa Memorial Fund).

References

1. Hodgkin AL, Huxley AF. Currents carried by sodium and potassium ions through the membrane of the giant axon of Loligo. J Physio. 1952;116(4):449–72.

(a) Arrival time difference between two curved fiber bundles

(b) Conduction velocity difference between two curved fiber bundles

Fig 21. Arrival time and conduction velocity difference between the two curved fibers. Two fiber bundles with ephaptic coupling (black), a single fiber bundle with ephaptic coupling (blue), and propagation without ephaptic coupling (red). (a) Arrival time difference in milliseconds, and (b) conduction velocity difference in meters per second, where $\rho = 1$, $r_b = 0.1$ cm, and $T = 24°C$.

2. Izhikevich EM. Dynamical Systems in Neuroscience. The MIT Press; 2007.

3. Hall JE, Hall ME. Guyton and Hall Textbook of Medical Physiology. 11th ed. Elsevier; 2021.

4. Sapolsky RM. Determined: A Science of Life without Free Will. Penguin Press; 2023.

5. Arvanitaki A. Effects evoked in an axon by the activity of a contiguous one. J Neurophysiol. 1942;5:89–108.

6. Bizsáki G, Anastassiou CA, Koch C. The origin of extracellular fields and currents - EEG, ECOG, LFP and spikes. Nat Rev Neurosci. 2012;13:407–420.

7. Anastassiou C, Koch C. Ephaptic coupling to endogenous electric field activity: why bother? Curr Opin Neurobiol. 2015;31:95–103.

8. Schmidt H, Knösche TR. Modelling the effect of ephaptic coupling on spike propagation in peripheral nerve fibres. Biol Cyber. 2022;116:461–473.

9. Capllonch-Juan M, SepulvedaI F. Modelling the effects of ephaptic coupling on selectivity and response patterns during artificial stimulation of peripheral nerves. PLOS Comput Biol. 2020;16(6):1007826.

10. Schmidt H, Knösche TR. Action potential propagation and synchronisation in myelinated axons. PLOS Comput Biol. 2019;15(10):e1007004.

11. Schmidt H, Hahn G, Deco G, Knösche TR. Ephaptic coupling in whiter matter fibre bundles modulates axonal transmission delays. PLOS Comput Biol. 2021;17(2):e1007004.

12. Jackson JD. Classical Electrodynamics. 3rd ed. Wiley; 2021.

13. Frankenhaeuser B, Huxley AF. The action potential in the myelinated nerve fibre of xenopus laevis as computed on the basis of voltage clamp data. J Physiol. 1964;171:302–315.

14. Goldman L, Albus JS. Computation of Impulse Conduction in Myelinated Fibers; Theoretical Basis of the Velocity-Diameter Relation. Biophys J. 1968;8(5):596–607.

15. Waxman SG. Conduction in myelinated, unmyelinated, and demyelinated fibers. Arch Neurol. 1977;34(10):585–9.

16. Waxman SG, Brill MH. Conduction through demyelinated plaques in multiple sclerosis: computer simulations of facilitation by short internodes. J Neurol. 1978;41:408–416.

17. Brill MH, Waxman SG, Moore JW, Joyner RW. Conduction velocity and spike configuration in myelinated fibres: computed dependence on internode distance. J Neurol Neurosurg Psychiatry. 1977;40(8):769–774.

18. Markin VS. Electric interaction of parallel non-myelinated nerve fibers. 1. Change in excitability of the adjacent fibre. Biofizika. 1970;15(1):122–133.

19. Markin VS. Electric interaction of parallel non-myelinated nerve fibers. 2. Shared conduction of impulses. Biofizika. 1970;15(4):681–9.

20. Markin VS. Electric interaction of parallel non-myelinated nerve fibers. 3. Interaction in nerve trunk. Biofizika. 1973;18(2):314–321.

21. Markin VS. Electric interaction of parallel non-myelinated nerve fibers. 4. The role of anatomic non uniformities of the nerve trunk. Biofizika. 1973;18(3):512–518.

22. Reutskiy S, Rossoni E, Tirozzi B. Conduction in bundles of demyelinated nerve fibers: computer simulation. Biol Cybern. 2003;89:439–448.

23. Sheheitli H, Jirsa VK. A mathematical model of ephaptic interactions in neuronal fiber pathways: Could there be more than transmission along the tracts? Netw Neurosci. 2020;4(3):595–610.

24. Schmitt OH. Information processing in the nervous system; 1968. Proceedings of a symposium held at the State University of New York at Buffalo, 21st-24th October.

25. Tung L. A Bi-Domain Model for Describing Ischemic Myocardial D-C Potentials [PhD thesis]. MIT; 1978.

26. Roth BJ. Electrical conductivity values used with the bidomain model of cardiac tissue. IEEE Trans Bio Eng. 1997;44(4):326–328.

27. Pullan AJ, Buist ML, Cheng LK. Mathematically modelling the electrical activity of the heart. World Scientific; 2005.

28. Cartan E. Riemannian Geometry in an Orthogonal Frame. World Scientific Pub. Co. Inc.; 2002.

29. Cartan E. Geometry of Riemannian Spaces. Math. Sci. Press; 2001.

30. Chun S. Method of moving frames to solve conservation laws on curved surfaces. J Sci Comput. 2012;53(2):268–294.

31. Chun S. Method of moving frames to solve (an)isotropic diffusion equations on curved surfaces. J Sci Comput. 2013;59(3):626–666.

32. Chun S, Eskilsson C. Method of moving frames to solve the shallow water equations on arbitrary rotating curved surfaces. J Compt Phys. 2017;333:1–23.

33. Chun S. Method of moving frames to solve the time-dependent Maxwell's equations on anisotropic curved surfaces: Applications to invisible cloak and ELF propagation. J Compt Phys. 2017;340:85–104.

34. Mitzdorf U. Current source-density method and application in cat cerebral cortex: investigation of evoked potentials and EEG phenomena. Physiol Rev. 1985;65(1):37–100.

35. Nicholson C, Freeman JA. Theory of current source-density analysis and determination of conductivity tensor for anuran cerebellum. J Neurophys. 1975;38:356–368.

36. Pitts W. Investigations on synaptic transmission. Cyber Trans 9th Conf. 1952; p. 159–162.

37. Roth BJ, Beaudoin DL. Approximate analytical solutions of the Bidomain equations for electrical stimulation of cardiac tissue with curving fibers. Phys Rev E. 2003;67:051925.

38. Kandel SM. The Electrical Bidomain Model: A Review. Sch Acad J Biosci. 2015;3(7):633–639.

39. Versteeg H, Malalasekra W. An Introduction to Computational Fluid Dynamics. Pearson; 2007.

40. Lewis RW, Nithiarasu P, Seetharamu KN. Fundamentals for the Finite Element Method for Heat and Fluid Flow. Wiley; 2004.

41. Ascher UM, Ruuth SJ, Spiteri RJ. Implicit-Explicit Runge-Kutta Methods for Time-Dependent Partial Differential Equations. Appl Numer Math. 1997;25:151–167.

42. Rush S, Larsen H. A Practical Algorithm for Solving Dynamic Membrane Equations. IEEE Trans Bio Eng. 1978;25(4):389–392.

43. Moxey D, Cantwell CD, Bao Y, Cassinelli A, Castiglioni G, Chun S, et al. Nektar++: enhancing the capability and application of high-fidelity spectral/hp element methods. Compt Phys Comm. 2020;249:107110.

44. Chun S, Jung J. Method of time map spatializing the dynamics of biological electric signal propagation and its applications to quick multidimensional simulation and deformed conductivity reconstruction. submitted. 2024;.

45. Tasaki I. Nervous Trasmission. Thomas; 1953.

46. Rasminsky M. The Effects of Temperature on Conduction in Demyelinated Single Nerve Fibers. Arch Neurol. 1973;28:287–292.