The effect of an increasing elephant population on the allometry of trees

Lucie Thel[1,*], Dietre Stols[1,*], Sarah Orth[2,3,4], D.D. Georgette Lagendijk[5,6,7], Rob Slotow[8], Jan A. Venter[1,3,#], Herve Fritz[3,4,9,#]

Authors details[a]

[1] Department of Conservation Management, Faculty of Science, Nelson Mandela University, George Campus, George, South Africa.

[2] Master de Biologie, École Normale Supérieure de Lyon, Université Claude Bernard Lyon 1, Université de Lyon, 69342 Lyon Cedex 07, France.

[3] Sustainability Research Unit, George Campus, Nelson Mandela University, George, South Africa.

[4] REHABS International Research Laboratory, CNRS-Université Lyon 1-Nelson Mandela University, George Campus, Nelson Mandela University, George, South Africa.

[5] School of Life Sciences, University of KwaZulu-Natal, Private Bag X01, Scottville, 3209 South Africa.

---

[a] **Lucie Thel**: Conceptualization - Methodology - Formal analysis - Investigation - Data Curation - Writing: Original Draft - Writing: Review & Editing – Visualisation; **Dietre Stols**: Conceptualization - Methodology - Formal analysis - Investigation - Resources - Data Curation - Writing: Original Draft - Writing: Review & Editing; **Sarah Orth:** Data Curation - Writing: Original Draft - Writing: Review & Editing; **Georgette Lagendijk**: Methodology - Data Curation - Validation - Writing: Review & Editing; **Rob Slotow**: Methodology - Data Curation - Validation - Writing: Review & Editing - Funding acquisition; **Jan Venter**: Conceptualization - Validation - Writing: Review & Editing - Supervision - Project administration - Funding acquisition; **Hervé Fritz**: Conceptualization - Validation - Methodology - Writing: Review & Editing - Supervision - Project administration - Funding acquisition.

[6] Conservation Ecology Group, Groningen Institute for Evolutionary Life Sciences

(GELIFES), University of Groningen, PO Box 11103, 9700 CC Groningen, The Netherlands.

[7] BirdEyes, Centre for Global Ecological Change at the Faculty of Science & Engineering

and Campus Fryslân, University of Groningen, Wirdumerdijk 34, 8911 CE Leeuwarden, The

Netherlands.

[8] Oppenheimer Fellow in Functional Biodiversity, Centre for Functional Biodiversity, School

of Life Science, University of Kwazulu-Natal, Pietermaritzburg 3209, South Africa.

[9] Université Claude Bernard Lyon 1, 43 Bd du 11 Novembre 1918, 69100 Villeurbanne,

France.

*These authors share first authorship

[#]These authors co-supervised the project

Corresponding authors: Herve Fritz, herve.fritz@cnrs.fr; Lucie Thel,

lucie.mc.thel@gmail.com; Dietre Stols, dietrestols@yahoo.com

Increasing elephant densities in fenced reserves reshape landscapes and modify vegetation.

By comparing with an elephant-free zone, we uncover how these giants leave their mark in

the trees structure.

Abstract

African savannas, which support a diverse range of plant species, from evergreen trees to deciduous shrubs, are regulated by complex interactions between climatic, fire, soil and herbivory processes. African elephants (*Loxodonta africana*) exert substantial top-down control over vegetation dynamics by utilising large amounts of plant biomass, as well as through various forms of non-consumptive effects. However, how much elephants affect the pattern of proportional growth between different tree components such as roots, stems or branches (known as allometry), remains understudied. Using vegetation data collected in 2000-2001 and 2019 for more than 3500 individual trees inside and outside Madikwe Game Reserve, South Africa, we determined the long-term effects of an increasing elephant population on allometric relationships in 13 tree species. The presence of elephants reduced logged canopy volume, and increasing elephant population in Madikwe reduced logged tree height and logged initial growth rate of the allometric relationship between tree height and tree diameter, without modifying asymptotic change in growth rate. Although not significantly, elephants seemed to increase asymptotic change in growth rate of canopy volume according to tree diameter, suggesting that compensatory mechanisms for the loss of height might happen in the longer-term. Although the species-specific elephant use index increased over the 20 years of the study, no consistent effect on allometric coefficients was detectable. Our study emphasises the necessity of long-term, species-specific studies to improve our understanding of how elephants affect allometric relationships in trees, which will ultimately guide management of savanna systems with increasing elephant numbers.

Keywords: large herbivore ecology, *Loxodonta africana*, canopy volume, tree height, tree diameter, long-term monitoring, savanna ecosystem.

Introduction

African savannas, which cover approximately half of sub-Saharan Africa (Du Toit and Cumming, 1999), are characterised by a "continuous grass understory interspersed by a discontinuous woody overstory" (Scholes and Archer, 1997). The productivity of these savannas is shaped by a complex interplay of micro- and macro-herbivory, soil dynamics, as well as fire, rainfall and frost regimes (Van Langevelde *et al*., 2003; Sankaran *et al*., 2005; Hoffmann *et al*., 2019), which contribute to the diversity of plant species and the structure of the vegetation (Sankaran *et al*., 2005). Although fire is commonly considered the dominant factor shaping savanna vegetation by reducing canopy and maintaining a continuous grass understory competing with trees (Sankaran *et al*., 2008), the herbivore community also strongly influences both vegetation structure and plant community composition through the consumption of biomass (Danell *et al*., 2006; O'Connor *et al.*, 2007; Lagendijk *et al*., 2015). This effect parallels the impacts of fire at a local scale and contributes to the stabilisation of the "tree-grass" coexistence in this ecosystem (Bond *et al*., 2005). Herbivory can lead to a decrease of up to 30.6 % in tree abundance and 57 % in grass abundance (Staver *et al*., 2021). In African savannas, the woody vegetation and biomass are influenced by mega-herbivores, particularly African elephants (Cumming *et al*., 1997; Kerley and Landman, 2006; Guldemond and Van Aarde, 2008; Valeix *et al*., 2011; Morrison *et al*., 2016). As ecosystem engineers (Jones *et al*., 1994), elephants directly affect tree height (O'Connor *et al*., 2024), canopy volumes (Guldemond and Van Aarde, 2010; O'Connor *et al*., 2024; Sorokina *et al*., 2024) and understorey composition (Ferry *et al*., 2021). Elephants affect species diversity by reducing the density of more palatable tree species, such as *Senegalia mellifera* and *Sclerocarya birrea caffra* (Skarpe *et al.*, 2004; Guldemond and Van Aarde, 2008; Sankaran *et al.*, 2013; Wigley *et al*., 2014; Bakker *et al.*, 2016; Cook *et al.*, 2017; Guldemond *et al.*, 2017; Sorokina *et al.*, 2024). Indeed, by utilising certain species, elephants can cause rapid

declines and even local extirpation in species that are unable to cope with their feeding

behaviour (*e.g.*, *Adansonia digitata*, *Aloe marlothii*, *Commiphora ugogensis*, *Sclerocarya*

birrea, and *Vachellia* species; Barnes, 1983; Kerley and Landman, 2006; O'Connor *et al.*,

2007; Guldemond and Van Aarde, 2008; Helm *et al.*, 2011).

Elephants can also significantly influence the allometry (*i.e.*, the proportional relationships

between different components of an individual through growth; Huxley and Teissier, 1936)

of the woody plant species (Makhabu *et al.*, 2006; Shannon *et al.*, 2008; Moncrieff, *et al.*,

2011; Lagendijk *et al.*, 2015). Some species utilised by elephants change growth form,

allowing them to persist (*e.g.*, *Colophospermum mopane*; Lewis, 1991; Kerley and Landman,

2006). They may experience reductions in height and total basal area (*e.g.*, *Senegalia*

nigrescens; Moncrieff *et al.*, 2011), or to the contrary thicker diameters and wider canopies

(*e.g.*, *Vachellia drepanolobium* and *Vachellia karroo*; Archibald and Bond 2003; Moncrieff

et al., 2011).

However, while earlier studies suggested that elephant activity could transform woodlands

and forests into grasslands or open savannas across Africa, a concern known as "the elephant

problem" (Laws, 1970; Caughley, 1976; Owen-Smith, 1988; Trollope *et al.*, 1998), more

recent research indicates that elephant effect on savanna vegetation is highly variable

(Guldemond and Van Aarde, 2008). A meta-analysis highlighted that these effects depend

significantly on environmental conditions, such as rainfall and geographical constraints,

leading to outcomes that range from negative to neutral, and even positive effects

(Guldemond and Van Aarde, 2008). While numerous studies have suggested that elephants

may have a negative impact on their habitats especially under sustained high population

densities, the expected changes might be less striking in open systems (Skarpe *et al.*, 2004;

Ferry *et al.*, 2021), or when measured over longer periods (Guldemond and Van Aarde,

2008).

In this study, we determined the effect of elephants on allometric relationships in 13 tree species. We used data from woody vegetation monitoring conducted in 2000-2001 and repeated in 2019 in Madikwe Game Reserve (hereafter MGR), which is exposed to an increasing elephant population, and in the adjacent Barokologadi Communal Property (hereafter BCP) in 2019, which does not contain elephants. We examined how elephants affected the allometric relationship between tree diameter and tree height, and the allometric relationship between tree diameter and canopy volume. We expected that repeated utilisation by elephants would lead to mature individual trees with large diameter stems being shorter and showing a reduced canopy volume, translating into the reduction of the allometric coefficients in the elephant zone (MGR) compared to the elephant-free zone (BCP, Stoinski et al., 2000; Holdo, 2007). Conversely, through growth stimulation, we could alternatively expect that elephants may have a positive effect on allometric relationships, as mature trees subject to long-term elephant exposure could exhibit greater height and larger canopy volumes if they grow taller to escape browsing (Canham et al., 1994). Eventually, we expected the possibility for compensatory growth which would translate into the reduction of the tree height parameters and the increase of the canopy volume parameters in their respective allometric relationships with tree diameter (Makhabu et al., 2006).

Material and methods

Study site and species

MGR (24°49′S - 26°13′E, South Africa), is a fenced reserve currently covering approximately 750 km² (Supporting information 1). MGR is situated within the central bushveld bioregion, as part of the broader savanna biome, and experiences a dry season from mid-April to mid-November (Rutherford et al., 2011). The average annual rainfall is approximately 475-520 mm (Trinkel et al., 2010).

In 2001, the elephant population was estimated at 389 individuals (*i.e.*, 0.60 elephants/km^2;

North West Parks and Tourism Board, 2024). By 2019, the elephant population almost

increased threefold, with an estimated 1318 individuals (*i.e.*, 1.76 elephants/km^2; North West

Parks and Tourism Board, 2024). Elephants constituted approximately 70 % of the herbivore

biomass in MGR at the time of the study (North West Parks and Tourism Board, 2024). Other

browsing herbivores in MGR include giraffe (*Giraffa camelopardalis*), greater kudu

(*Tragelaphus strepsiceros*), springbuck (*Antidorcas marsupialis*) and impala (*Aepyceros*

melampus). The density of the latter increased substantially between 2011 and 2019 from

3.84 individuals/km^2 to 6.30 individuals/km^2 (Supporting information 2).

BCP, covering 160 km^2 (Supporting information 1), is separated from MGR by an elephant-

proof fence, rendering it an elephant-free zone. BCP has been used for cattle, goat, and sheep

farming for meat production since 2009, with occasional firewood collection (Barokologadi

Communal Property Association, unpublished report). Bushbuck (*Tragelaphus scriptus*),

greater kudu and impala roam freely in BCP in unknown numbers. Although unfortunately

there is no written report available, BCP experienced more frequent fires than MGR over the

study period, where fire frequency is controlled and maintained at a very low level (park

management services, personal communication).

Sampling design

Tree morphological measurements were collected at 35 sites in MGR during three two-weeks

periods in January 2000, July 2000 and January 2001, following a stratified random sampling

protocol (*i.e.*, stratified across vegetation types and randomised within vegetation types, Page

and Slotow, 2001). The minimal distance between sites was approximately 100 m. In each

site, a 50-metre transect of variable width (from 2 m to 50 m depending on the woody

vegetation density of the site) was used to record all living woody plants. A nested sampling

approach was used to identify species not found within the core transect by extending

sampling to surrounding areas with a maximum transect size of 100 x 50 m. The 35 original

sites from 2000-2001 were re-sampled between February and June 2019 (except for 2 sites

sampled in October), as well as 27 new sites in BCP, following the same methodology. In

BCP, 23 sites were sampled during a two-week period in October, and four sites were

sampled between April and June. The minimum distance between sites in BCP was

approximately 240 m.

We defined three spatio-temporal treatments: i) "Old", using MGR data from 2000-2001 and

exposed to a relatively low elephant density (n = 2331 trees, 0.60 elephants/km^2), ii) "New",

using MGR data from 2019 and exposed to a higher elephant density (n = 1583 trees, 1.76

elephants/km^2) and iii) "Out", using BCP data from 2019 and excluded from elephants (n =

1026 trees, 0 elephants/km^2). Although not completely untouched (*e.g.*, occasional livestock

browsing and firewood collection), the "Out" treatment provides a good comparison point for

the allometric relationships of trees in the absence of any elephant pressure.

For each living tree, we recorded the number of live stems and measured stem diameters. For

the analyses, we only used the stems within the six largest stem diameters for a given

individual (*i.e.*, all the stems of a similar diameter up to six classes, hereafter called "stem

classes", range = 0.1 to 75 cm). We measured the total tree height (**Height**, in metres) as the

vertical distance from the ground to the tree's highest point, and the height of the bottom of

the canopy (*HBC*, in metres) as the vertical distance from the ground to the height where the

main canopy starts. For trees larger than 2 m in height, measurements were estimated. We

calculated the total tree diameter (**Diameter**, in centimetres) as the diameter of a disk of an

area equal to the sum of all the stem diameters, each stem being modelled as a perfect

cylinder, as explained in equation (1):

$$\boldsymbol{Diameter} \ = \ \sqrt{\textstyle\sum_{i=1}^{n} \boldsymbol{nb\ stem_i \times d_i}} \qquad \textit{equation (1)},$$

where d_i is the diameter of the stem class i and *nb stem$_i$* is the number of stems belonging to

the stem class i. We calculated the canopy volume (**Canopy Volume**, in cubic metres) using

the formula of the volume of an ellipsoid (equation (2)):

$$\boldsymbol{Canopy\ Volume\ =\ \frac{4}{3} \times \Pi \times \left(\frac{Diameter\ max}{2}\right)^2 \times \left(\frac{Canopy\ height}{2}\right)}$$ *equation (2)*,

where *Diameter max* (in metres) corresponds to the widest horizontal diameter of the

ellipsoid formed by the canopy, and *Canopy height* corresponds to the difference between

Height and *HBC* and represents the vertical diameter of the ellipsoid formed by the canopy.

For each individual tree, we estimated elephant utilisation as the percentage of biomass (*i.e.*,

branches, leaves, stems, bark and/or roots) removed using eight categories: 0 (0 %), 1 (1 to 10

%), 2 (11 to 25 %), 3 (26 to 50 %), 4 (51 to 75 %), 5 (76 to 90 %), 6 (91 to 99 %), and 7 (100

%). We then calculated the species-specific mean elephant utilisation per treatment (hereafter

called **Elephant Use**).

Statistical analyses

We identified the species that were present in all three treatments with sufficient sample size

recorded (*i.e.*, $\geq$ 15 trees in each treatment) and included only these in our analyses to ensure

between-treatments comparability and statistical robustness ($n = 13$ species; $n_{Old} = 1693$ trees,

$n_{New} = 1392$ trees, $n_{Out} = 713$ trees; Supporting information 3). In other words, this study

focussed on the allometric responses of common tree species in this landscape. We first

investigated the changes of tree **Height** and **Canopy Volume** between treatments. We

conducted a one-way ANOVA on the log-transformed (natural log to meet assumptions of

normality and heteroscedasticity) tree **Height** and **Canopy Volume** of all species pooled

together. We subsequently conducted a post-hoc pairwise comparison test using Tukey's test

to identify significant differences between treatments. We conducted the same analysis for

each species individually (see details in Supporting information 4).

We then determined the species-specific coefficients of the allometric relationships between

tree **Height** and tree **Diameter**, and between **Canopy Volume** and tree **Diameter**. One

commonly expected relationship in tree allometry (Kearsley *et al.*, 2013) can be written as

(equation (3)):

$$Y \;=\; A \times X^B + C \quad \text{equation (3),}$$

where X and Y are two tree measurements (X is tree diameter and Y is either tree height or

canopy volume), A corresponds to the Initial Growth Rate, B corresponds to the **Asymptotic**

Change in Growth Rate, and C is a constant close to 0 (C is 0 in nature but to fit the model

we allow for some parameter adjustment in the fit; if C is significantly different from 0 it

suggests poor fit of the model). Mathematically, the **Initial Growth Rate** illustrates the

intrinsic growth linking the two parameters X and Y, namely the growth potential that can be

reached. The **Asymptotic Change in Growth Rate** corresponds to the speed of the increase

in growth, namely how fast the curve increases: if $B < 1$ the growth rate in height, or canopy

volume, decreases as the tree diameter increases; if $B > 1$ the growth rate increases with

increasing tree diameter. However, in our study, visual inspection revealed non-normality and

heteroscedasticity associated with an important variance in the tree **Height** and **Canopy**

Volume and creating a poor fit of the data with equation (3). Therefore, we used the log-

transformed formula, and we fitted a linear regression instead (equation (4)):

$$log(Y) \;=\; I + B \times log(X) \quad \text{equation (4),}$$

where X and Y are the same two tree measurements, B remains the **Asymptotic Change in**

Growth Rate (equation (3)) and I corresponds to the Logged Initial Growth Rate coupled

with the constant term (A and C, respectively in equation (3)), hereafter named **Logged**

Initial Growth Rate for simplicity. We fitted such a relationship for each species in each

treatment separately, and extracted the coefficients B and I of the allometric relationship

linking the dependent (Y, *i.e.*, tree **Height** and **Canopy Volume**, respectively) and the

explanatory (X, *i.e.*, tree **Diameter**) variables. Eventually, we tested the correlation between

tree **Height** and **Canopy Volume** using the Spearman's test of correlation, as these two

variables are known to be correlated (*e.g.*, Arzai and Aliyu, 2010).

We then determined how the treatment and the **Elephant Use** affected the coefficients B and

I of the allometric relationships (between **Height** and **Diameter**, and between **Canopy**

Volume and **Diameter**, respectively). We first conducted a one-way ANOVA to explore the

effect of treatment on these coefficients, followed by a post-hoc pairwise comparison test

using Tukey's test to identify significant differences between pairs of treatments. We then

used linear regressions with an interaction between treatment and **Elephant Use** to determine

the effect of **Elephant Use** on the coefficients B and I in the "Old" versus "New" treatment.

All statistical analyses were conducted using R (version 4.4.0, R Core Development Team

2024).

Results

We identified 13 species present in all three treatments, which were represented by more than

15 individuals in each treatment: *Combretum hereroense*, *Dichrostachys cinerea*, *Euclea*

undulata, *Flueggea virosa*, *Grewia flava*, *Grewia monticola*, *Gymnosporia buxifolia*,

Senegalia erubescens, *Senegalia mellifera*, *Vachellia karroo*, *Vachellia tortilis*, *Ximenia*

americana and *Ziziphus mucronata* (Supporting information 3). The most abundant species

within all three treatments was *Dichrostachys cinerea* (n_{old} = 327, n_{new} = 482, n_{out} = 194), and

the least abundant species was *Senegalia mellifera* in the "Old" and "New" treatment (n_{old} =

23, n_{new} = 15, n_{out} = 67), as well as *Vachellia karroo* in the "Out" treatment (n_{old} = 108, n_{new} =

54, n_{out} = 16). Ten species experienced a marked decrease in occurrence (between -10 and -50

%) between the "Old" and "New" treatments, while *Combretum hereroense*, *Grewia*

monticola and *Dichrostachys cinerea* increased, the latter by almost 50 %. The number of

individuals of *Euclea undulata*, *Gymnosporia buxifolia*, *Vachellia karroo*, *Senegalia*

erubescens, *Ximenia caffra* and *Ziziphus mucronata* decreased by almost 50 % between the

"Old" and "New" treatments (Supporting information 3).

Overall, the logged tree **Height** was significantly lower in the "New" than in the "Old" ($P <$

0.01) and the "Out" ($P < 0.01$) treatments, and significantly higher in the "Out" than in the

"Old" treatment ($P < 0.01$, Fig. 1a). The logged **Canopy Volume** was similar in the "New"

and "Old" treatments ($P = 0.39$), but it was significantly larger in the "Out" than in both other

elephant treatments (both $P < 0.01$, Fig. 1b). Although these results were mostly consistent

among five species (*Dichrostachys cinerea*, *Grewia monticola*, *Vachellia karroo*, *Ximenia*

americana and *Ziziphus mucronata*), there were species-specific patterns (Supporting

information 4). Notably, *Flueggea virosa* increased in logged tree **Height** and logged

Canopy Volume between the "Old" and "New" treatments, and the differences in logged tree

Height and logged **Canopy Volume** were less pronounced between the two elephant

treatments ("Old" and "New") and the elephant free treatment ("Out") for *Gymnosporia*

buxifolia. No significant differences were detected for *Senegalia mellifera*.

We found a significant positive relationship between logged tree **Diameter** and logged tree

Height for all 13 species in each treatment, except for *Senegalia mellifera* in the "New"

treatment (Supporting information 5). Similarly, we found a significant positive relationship

between logged tree **Diameter** and logged **Canopy Volume** for all 13 species in each

treatment (Supporting information 6). Additionally, we found a significant and very high

correlation between logged tree **Height** and logged **Canopy Volume** in all species in all three

treatments (all $\rho > 0.6$, all $P \leq 0.01$, Table 1).

The **Logged Initial Growth Rate** (*I*) was significantly lower in the "New" than in the "Old"

($P = 0.01$) and the "Out" ($P = 0.04$) treatments for the tree **Height** - tree **Diameter**

relationship (Fig. 2b). Although not significant, we found a similar trend for the **Logged**

Initial Growth Rate (*I*) of the **Canopy Volume** - tree **Diameter** relationship ($P = 0.06$, Fig.

2d). The **Asymptotic Change in Growth Rate** (*B*) was not significantly different in any

treatment for the tree **Height** - tree **Diameter** relationship ($P = 0.10$, Fig. 2a). However, it

was significantly larger in the "Out" than in the "Old" treatment for the **Canopy Volume** -

tree **Diameter** relationship ($P = 0.01$, Fig. 2c).

On average, **Elephant Use** varied between 0.0 (for *Ximenia americana* in the "Old"

treatment) and 2.1 (for *Senegalia erubescens* in the "New" treatment). The least used species

were *Ximenia americana* and *Euclea undulata* (**Elephant Use** ranging between 0.0 and 0.2 in

the "Old" and "New" treatments). Overall, there was an increase in **Elephant Use** between

the "Old" and "New" treatments, except for *Euclea undulata* for which the index remained

stable, and *Gymnosporia buxifolia* which was the only species to experience a decrease (from

0.8 to 0.6 in the "Old" and "New" treatments respectively, Fig. 3). *Senegalia erubescens*,

Grewia monticola and *Grewia flava* experienced the largest increase in **Elephant Use** ($\geq$ 1.3

increase between the "Old" and "New" treatments). **Elephant Use** did not affect variation of

the **Logged Initial Growth Rate** (*I*), or variation of the **Asymptotic Change in Growth**

Rate (*B*) for either allometric relationship (Table 2, Fig. 3).

Discussion

In this study, we highlighted the effect of an increasing elephant population on the structure

of the woody vegetation at the community scale. Elephants reduced the **Height** and **Canopy**

Volume of trees, and reduced the **Logged Initial Growth Rate** of the allometric relationship

between tree **Height** and tree **Diameter**, without modifying the **Asymptotic Change in**

Growth Rate. Our results suggest that a compensatory mechanism for the loss of tree **Height**

could start to take place after 20 years of continuously increasing exposure to elephants, with

a detectable but not significant increase in the **Asymptotic Change in Growth Rate** of the

Canopy Volume. Although the species-specific **Elephant Use** index increased over the

period of the study, no general trend of its effect on allometric coefficients was detectable,

suggesting species-specific patterns.

Trees were shorter and had a smaller **Canopy Volume** (Fig. 1) in MGR ("Old" and "New"

treatments) than in BCP ("Out" treatment). Even at low density, elephants might reduce the

abundance of very large trees, consistent with similar studies conducted in African savannas

(*e.g.*, Shannon *et al*., 2008; Mapaure and Moe, 2009; Gandiwa *et al*., 2011; Cook *et al*.,

2017). The smaller **Logged Initial Growth Rate** (*I*) of tree **Height** in the "New" compared

to the "Out" treatments, coupled with the absence of modification of the **Asymptotic Change**

in Growth Rate (*B*), denotes a reduction of the overall growth potential of trees in the former

treatment, but not of the speed at which the final growth potential is reached. This suggests

that elephants might participate in the maintenance of small trees in the landscape (Kalwij *et*

al., 2010). However, the allometric coefficients (tree **Height** - tree **Diameter** relationship) in

the "Out" and "Old" treatments were not significantly different, suggesting that elephants

affect the allometric relationship between tree **Height** and tree **Diameter**, but only

subsequent to a long-term population increase.

Although there was a high correlation between tree **Height** and **Canopy Volume** (Table 1;

Blanchard *et al.*, 2016; Davies *et al.*, 2018), there was a trend towards an increase of the

Asymptotic Change in Growth Rate (*B*) only in the allometric relationship between

Canopy Volume and tree **Diameter** in MGR (*mean*$_{Old}$ = 2.02 ± 0.27, *mean*$_{New}$ = 2.20 ± 0.53,

Fig. 2c) with an increasing elephant population. Such a phenomenon could denote the

development of a long-term compensation mechanism, trees being shortened by elephants,

they would increase in canopy volume. When exposed to herbivory, trees can develop

compensatory mechanisms to balance the loss of biomass (McNaughton, 1983; Paige 1992;

Zhang *et al*., 2018), although species-specific patterns are common (Focardi and Tinelli,

2005). Alternatively, elephant presence is usually associated with a decreased density in trees

(O'Connor *et al.*, 2007; Thompson *et al.*, 2022), which might reduce light competition,

leading to increased tree growth (Sevillano *et al.*, 2016). This increase of the **Asymptotic**

Change in Growth Rate of **Canopy Volume** was however not significant ($P = 0.54$), and

further exploration is needed to confirm if this trend becomes significant in the longer-term.

In MGR and BCP, besides elephants, other browsing herbivores also occur, which can affect

tree structure. In MGR, impala, a mixed feeder (Owen-Smith, 2008), is known for its

negative impact on the recruitment of trees in both the seedling and sapling classes, but also

reducing recruitment into taller height classes as they restrain tree growth within the

browsing trap (O'Kane *et al.*, 2012; Lagendijk *et al.*, 2015). Similarly, the BCP area was

affected by fuelwood collection and livestock (e.g. goats), which can both affect the structure

of the woody vegetation (Stuart-Hill, 1992; Scogings and Macanda, 2005; Tripathi et al.,

2019). During the study period, MGR has experienced an increase in its impala population

(from 2879 individuals in 2011 to 6110 individuals in 2021, Supporting information 2),

which has likely contributed to the modification of the allometric relationships between the

"Old" and "New" treatments (Staver and Bond 2014). Unfortunately, there was no data

available regarding fire regimes in the MGR and BCP, although several studies showed that

fire interacts with herbivory and contributes to modifications in the woody vegetation

dynamics (Holdo *et al.*, 2009; Shannon *et al.*, 2011; Vanak *et al.*, 2012).

Although there was a general increase of **Elephant Use** in MGR, we did not find evidence

for a modification of the **Logged Initial Growth Rate** (I) and the **Asymptotic Change in**

Growth Rate (B) in response to this increase (Fig. 3, Table 2). The wide dispersion of these

species-specific coefficients prevents detecting any significant trend, and rather denotes

species-specific patterns as found by Levick and Rogers (2008). Some species characterised

by higher abundance or palatability can be preferentially selected by herbivores (e.g. kudu

and elephant select preferentially *Dichrostachys cinerea*, Owen-Smith and Cooper 1987; de Garine-Wichatitsky *et al.*, 2004; Makhabu, 2005). Depending on the tree and herbivore species, different parts of the tree can be preferentially selected and exposed to various types of modifications. Elephants can topple, strip and break stems (e.g. in *Vachellia tortilis*, Boundja and Midgley, 2010) and/or consume the bark (e.g. in *Senegalia nigrescens*; Watson *et al.*, 2020), while giraffes browse exclusively on leaves (e.g. in *Acacia spp.*; Mahenya *et al.*, 2016). Such preferences generate the potential for species-specific allometric responses in trees (MacGregor and O'Connor, 2004).

Our study provides an overview of the influence of elephants on tree allometry (reduced **Height** and **Canopy Volume**, reduced **Logged Initial Growth Rate** of the allometric relationship between tree **Height** and tree **Diameter**). However, allometric relationships in trees are the result of a complex interplay between tree species traits (e.g. regeneration by regrowth in *Acacia nigrescens*, Fornara and du Toit, 2007), herbivore decisions (e.g. different herbivores browse different plant parts; Cameron and du Toit, 2007) and abiotic factors (e.g. fire regimes; Schafer and Mack, 2014). Further exploration is thus needed to better understand how elephants interact with these coexisting factors and modify the allometric relationships at the species-specific level in African savannas.

Acknowledgments

We acknowledge and thank Bruce Page, who designed the sampling protocol, and supervised the sampling in 2000-2001. We thank all University of Kwazulu-Natal students and volunteers involved in the 2000-2001 sampling. We thank the fieldworkers who collected the data in 2019 and managed the data for this study: Yentl Swartz, Chris Brooke, Adriaan Venter, Michelle Marais, Jandre Vermaak, Terry-Lee Honiball. We also thank the North-West Parks and Tourism Board for their logistical support for the study.

Declarations

Funding: The 2000-2001 sampling was funded by the Amarula Elephant Research Programme through a donation by Distell (PTY) Ltd, and the University of Kwazulu-Natal, with in-kind support from North West Parks and Tourism Board. Postgraduate Research Scholarship funding from Nelson Mandela University and REHABS funding to DS.

Conflict of interest: The authors declare that they have no conflict of interest.

Ethical approval: Not applicable.

Consent to participate: Not applicable.

Consent for publication: Not applicable.

Availability of data and material: The data was deposited in the Zenodo Repository under the reference number https://doi.org/10.5281/zenodo.14268022.

Code availability: Not applicable.

Authors' contributions: Lucie Thel: Conceptualization - Methodology - Formal analysis - Investigation - Data Curation - Writing: Original Draft - Writing: Review & Editing - Visualisation; **Dietre Stols**: Conceptualization - Methodology - Formal analysis - Investigation - Resources - Data Curation - Writing: Original Draft - Writing: Review & Editing; **Sarah Orth:** Data Curation - Writing: Original Draft - Writing: Review & Editing

Georgette Lagendijk: Methodology - Data Curation - Validation - Writing: Review &

Editing; **Rob Slotow**: Methodology - Data Curation - Validation - Writing: Review & Editing

- Funding acquisition; **Jan Venter**: Conceptualization - Validation - Writing: Review &

Editing - Supervision - Project administration - Funding acquisition; **Hervé Fritz**:

Conceptualization - Validation - Methodology - Writing: Review & Editing - Supervision -

Project administration - Funding acquisition.

References

Archibald, S. and Bond, W.J., 2003. Growing tall vs growing wide: tree architecture and allometry of *Acacia karroo* in forest, savanna, and arid environments. *Oikos*, 102(1), pp.3-14.

Arzai, A., and Aliyu B., 2010. The relationship between canopy width, height and trunk size in some tree species growing in the Savanna zone of Nigeria. *Bayero Journal of Pure and Applied Sciences*, 3(1).

Bakker, E.S., Gill, J.L., Johnson, C.N., Vera, F.W., Sandom, C.J., Asner, G.P. and Svenning, J.C., 2016. Combining paleo-data and modern exclosure experiments to assess the impact of megafauna extinctions on woody vegetation. *Proceedings of the National Academy of Sciences*, 113(4), pp.847-855.

Barnes, R.F.W., 1983. Effects of elephant browsing on woodlands in a Tanzanian National Park: measurements, models and management. *Journal of Applied Ecology*, pp.521-539.

Barokologadi Communal Property Association. The Path Followed by Barokologadi CPA (BCPA) Since the Claim of Melorane, Barokologadi's Ancestral Land. *Unpublished report*.

Blanchard, E., Birnbaum, P., Ibanez, T., Boutreux, T., Antin, C., Ploton, P., Vincent, G., Pouteau, R., Vandrot, H., Hequet, V. and Barbier, N., 2016. Contrasted allometries between stem diameter, crown area, and tree height in five tropical biogeographic areas. *Trees*, 30, pp.1953-1968.

Bond, W.J., Woodward, F.I., Midgley, G.F., 2005. The global distribution of ecosystems in a world without fire. *New Phytologist* 165, pp. 525–538.

Boundja, R.P. and Midgley, J.J., 2010. Patterns of elephant impact on woody plants in the Hluhluwe-Imfolozi park, Kwazulu-Natal, South Africa. *African Journal of Ecology*, 48(1), pp.206-214.

Cameron, E. Z. and du Toit, J. T., 2007. Winning by a neck: tall giraffes avoid competing

with shorter browsers. *The American Naturalist*, 169(1), pp.130-135.

Canham, C. D., McAninch, J. B. and Wood, D. M., 1994. Effects of the frequency, timing,

and intensity of simulated browsing on growth and mortality of tree seedlings. *Canadian*

Journal of Forest Research, 24(4), pp.817-825.

Caughley, G., 1976. The elephant problem–an alternative hypothesis. African Journal of

Ecology, 14(4), pp.265-283.

Cook, R.M., Witkowski, E.T.F., Helm, C.V., Henley, M.D. and Parrini, F., 2017. Recent

exposure to African elephants after a century of exclusion: Rapid accumulation of marula

tree impact and mortality, and poor regeneration. *Forest Ecology and Management*, 401,

pp.107-116.

Cumming, D.H., Fenton, M.B., Rautenbach, I.L., Taylor, R.D., Cumming, G.S., Cumming,

M.S., Dunlop, J.M., Ford, A.G., Hovorka, M.D., Johnston, D.S. and Kalcounis, M., 1997.

Elephants, woodlands and biodiversity in southern Africa. *South African Journal of*

Science, 93(5), pp.231-236.

Danell, K., Bergström, R., Duncan, P. and Pastor, J. eds., 2006. Large herbivore ecology,

ecosystem dynamics and conservation (Vol. 11). Cambridge University Press.

Davies, A.B., Gaylard, A. and Asner, G.P., 2018. Megafaunal effects on vegetation structure

throughout a densely wooded African landscape. *Ecological applications*, 28(2), pp.398-

408.

de Garine-Wichatitsky, M., Fritz, H., Gordon, I.J. and Illius, A.W., 2004. Bush selection

along foraging pathways by sympatric impala and greater kudu. *Oecologia*, 141, pp.66-75.

Du Toit, J.T., Cumming, D.H.M., 1999. Functional significance of ungulate diversity in

African savannas and the ecological implications of the spread of pastoralism, *Biodiversity*

and Conservation 8, pp.1643-1661.

459 Ferry, N., Dray, S., Fritz, H., Ipavec, A., Wigley, B.J., Charles-Dominique, T., Bourgarel, M.,

460   Sebele, L. and Valeix, M., 2021. Long-term high densities of African elephants clear the

461   understorey and promote a new stable savanna woodland community. *Journal of*

462   *Vegetation Science*, 32(6), p.e13101.

463 Focardi, S. and Tinelli, A., 2005. Herbivory in a Mediterranean forest: browsing impact and

464   plant compensation. *Acta Oecologica*, 28(3), pp. 239-247.

465 Fornara, D. A. and Toit, J. D., 2007. Browsing lawns? Responses of *Acacia nigrescens* to

466   ungulate browsing in an African savanna. *Ecology*, 88(1),pp. 200-209.

467 Gandiwa, E., Magwati, T., Zisadza, P., Chinuwo, T. and Tafangenyasha, C., 2011. The

468   impact of African elephants on *Acacia tortilis* woodland in northern Gonarezhou National

469   Park, Zimbabwe. *Journal of Arid Environments*, 75(9), pp 809-814.

470 Guldemond, R. and Van Aarde, R., 2008. A meta-analysis of the impact of African elephants

471   on savanna vegetation. *The Journal of Wildlife Management*, 72(4), pp.892-899.

472 Guldemond, R. and Van Aarde, R., 2010. The influence of tree canopies and elephants on

473   sub-canopy vegetation in a savannah. *African Journal of Ecology*, 48(1), pp.180-189.

474 Guldemond, R.A., Purdon, A. and Van Aarde, R.J., 2017. A systematic review of elephant

475   impact across Africa. *PloS one*, 12(6), p.e0178935.

476 Helm, C.V., Scott, S.L. and Witkowski, E.T.F., 2011. Reproductive potential and seed fate of

477   *Sclerocarya birrea* subsp. *caffra* (marula) in the low altitude savannas of South Africa.

478   *South African Journal of Botany*, 77(3), pp.650-664.

479 Hoffmann, W.A., Flake, S.W., Abreu, R.C.R., Pilon, N.A.L., Rossatto, D.R., Durigan, G.,

480   2019. Rare frost events reinforce tropical savanna–forest boundaries. *Journal of Ecology*

481   107, pp. 468–477.

Holdo, R. M., Holt, R. D. and Fryxell, J. M., 2009. Grazers, browsers, and fire influence the

extent and spatial pattern of tree cover in the Serengeti. *Ecological Applications*, 19(1),

pp.95-109.

Holdo, R.M., 2007. Elephants, fire, and frost can determine community structure and

composition in Kalahari woodlands. *Ecological applications*, 17(2), pp.558-568.

Huxley, J. S. and G. Teissier., 1936. Terminology of relative growth. *Nature*, 137, pp. 780–

781.

Jones, C. G., Lawton, J. H. and Shachak, M., 1994. Organisms as ecosystem engineers.

Oikos, 69(3): pp.373-386.

Kalwij, J. M., De Boer, W. F., Mucina, L., Prins, H. H. T., Skarpe, C. and Winterbach, C.,

2010. Tree cover and biomass increase in a southern African savanna despite growing

elephant population. *Ecological Applications*, 20(1), pp.222-233.

Kearsley, E., De Haulleville, T., Hufkens, K., Kidimbu, A., Toirambe, B., Baert, G.,

Huygens, D., Kebede, Y., Defourny, P., Bogaert, J. and Beeckman, H., 2013.

Conventional tree height–diameter relationships significantly overestimate aboveground

carbon stocks in the Central Congo Basin. *Nature communications*, 4(1), p.2269.

Kerley, G.I. and Landman, M., 2006. The impacts of elephants on biodiversity in the Eastern

Cape Subtropical Thickets: elephant conservation. *South African Journal of Science*,

102(9), pp.395-402.

Lagendijk, D.G., Thaker, M., De Boer, W.F., Page, B.R., Prins, H.H. and Slotow, R., 2015.

Change in mesoherbivore browsing is mediated by elephant and hillslope position. *PloS*

one, 10(6), p.e0128340.

Laws, R.M., 1970. Elephants as agents of habitat and landscape change in East Africa. *Oikos*,

pp.1-15.

Levick, S. and Rogers, K., 2008. Patch and species specific responses of savanna woody

vegetation to browser exclusion. *Biological Conservation*, 141(2), pp.489-498.

Lewis, D., 1991. Observations of tree growth, woodland structure and elephant damage on

Colophospermum mopane in Luangwa Valley, Zambia. *African Journal of Ecology*, 29(3),

pp.207-221.

MacGregor, S.D. and O'Connor, T.G., 2004. Response of *Acacia tortilis* to utilization by

elephants in a semi-arid African savanna. *South African Journal of Wildlife Research*,

34(1), pp.55-66.

Mahenya, O., Ndjamba, J.K., Mathisen, K.M. and Skarpe, C., 2016. Giraffe browsing in

response to plant traits. *Acta Oecologica*, 75, pp.54-62.

Makhabu, S. W., 2005. Resource partitioning within a browsing guild in a key habitat, the

Chobe Riverfront, Botswana. *Journal of Tropical Ecology*, 21(6), pp.641-649.

Makhabu, S.W., Skarpe, C. and Hytteborn, H., 2006. Elephant impact on shoot distribution

on trees and on rebrowsing by smaller browsers. *Acta oecologica*, 30(2), pp.136-146.

Mapaure, I. and Moe, S.R., 2009. Changes in the structure and composition of miombo

woodlands mediated by elephants (*Loxodonta africana*) and fire over a 26-year period in

north-western Zimbabwe. *African journal of ecology*, 47(2), pp.175-183.

McNaughton, S. J., 1983. Compensatory plant growth as a response to herbivory. *Oikos*,

pp.329-336.

Moncrieff, G.R., Chamaillé-Jammes, S., Higgins, S.I., O'Hara, R.B., Bond, W.J., 2011. Tree

allometries reflect a lifetime of herbivory in an African savanna. *Ecology*, 92, pp. 2310–

2315.

Morrison, T.A., Holdo, R.M., Anderson, T.M., 2016. Elephant damage, not fire or rainfall,

explains mortality of overstorey trees in Serengeti. *Journal of Ecology* 104,pp. 409–418.

North West Parks and Tourism Board, 2024, Madikwe Game Reserve Annual Game Census

Report. *Unpublished report.*

O'Connor, T.G., Goodman, P.S. and Clegg, B., 2007. A functional hypothesis of the threat of

local extirpation of woody plant species by elephant in Africa. *Biological Conservation*,

136(3), pp.329-345.

O'Connor, T., Ferguson, A., Clegg, B.W., Pallett, N., Midgley, J.J. and Shimbani, J., 2024.

Emergent trees in *Colophospermum mopane* woodland: influence of elephant density on

persistence versus attrition. *PeerJ*, *12*, p.e16961.

O'Kane, C.A., Duffy, K.J., Page, B.R. and Macdonald, D.W., 2012. Heavy impact on

seedlings by the impala suggests a central role in woodland dynamics. *Journal of Tropical*

Ecology, 28(3), pp.291-297.

Owen-Smith, N. and Cooper, S. M., 1987. Palatability of woody plants to browsing

ruminants in a South African savanna. *Ecology*, 68(2), pp.319-331.

Owen-Smith, N., 2008. The Comparative Population Dynamics of Browsing and Grazing

Ungulates. In: Gordon, I.J., Prins, H.H.T. (eds) *The Ecology of Browsing and Grazing.*

Ecological Studies, 195. Springer, Berlin, Heidelberg. pp.149-177.

Owen-Smith, R.N., 1988. Megaherbivores: the influence of very large body size on ecology.

Cambridge university press.

Page, B., Slotow, R., 2001. The Influence of Browsers, Particularly Elephants, on Vegetation

(Diversity) at Madikwe Game Reserve. *Preliminary Project Report*, April 2001. Amarula

Elephant Research Programme, School of Life and Environmental Science, University of

Natal, Durban.

Paige, K. N., 1992. Overcompensation in response to mammalian herbivory: from mutualistic

to antagonistic interactions. *Ecology*, 73(6), pp.2076-2085.

R Core Development Team., 2024. R: A language and environment for statistical computing,

version 4.4.0. R Foundation for Statistical Computing. Available at: https://www.R-

project.org/

Rutherford, M.C., Mucina, L., Lötter, M.C., Bredenkamp, G.J., Smit, J.H.L., Scott-Shaw,

C.R., Hoare, D.B., *et al.*, 2011. *Savanna Biome.* In: L. Mucina & M.C. Rutherford, eds.

The Vegetation of South Africa, Lesotho and Swaziland. Strelitzia 19. Pretoria: South

African National Biodiversity Institute (SANBI), pp.439–538.

Sankaran, M., Augustine, D.J. and Ratnam, J., 2013. Native ungulates of diverse body sizes

collectively regulate long-term woody plant demography and structure of a semi-arid

savanna. *Journal of Ecology*, 101(6), pp.1389-1399.

Sankaran, M., Hanan, N.P., Scholes, R.J., Ratnam, J., Augustine, D.J., Cade, B.S., Gignoux,

J., Higgins, S.I., Le Roux, X., Ludwig, F. and Ardo, J., 2005. Determinants of woody

cover in African savannas. *Nature*, 438(7069), pp.846-849.

Sankaran, M., Ratnam, J. and Hanan, N., 2008. Woody cover in African savannas: the role of

resources, fire and herbivory. *Global ecology and biogeography*, 17(2), pp.236-245.

Schafer, J. L. and Mack, M. C., 2014. Growth, biomass, and allometry of resprouting shrubs

after fire in scrubby flatwoods. *The American Midland Naturalist*, 172(2), pp.266-284.

Scholes, R.J. and Archer, S.R., 1997. Tree-grass interactions in savannas. *Annual review of

Ecology and Systematics*, 28(1), pp.517-544.

Scogings, P., and Macanda, M. 2005. *Acacia karroo* responses to early dormant season

defoliation and debarking by goats in a semi-arid subtropical savanna. *Plant Ecology*, 179,

pp.193-206.

Sevillano, I., Short, I., Grant, J. and O'Reilly, C., 2016. Effects of light availability on

morphology, growth and biomass allocation of *Fagus sylvatica* and *Quercus robur*

seedlings. *Forest Ecology and Management*, 374, pp. 11-19.

Shannon, G., Druce, D.J., Page, B.R., Eckhardt, H.C., Grant, R. and Slotow, R., 2008. The

utilization of large savanna trees by elephant in southern Kruger National Park. *Journal of*

tropical ecology, 24(3), pp.281-289.

Shannon, G., Thaker, M., Vanak, A.T., Page, B.R., Grant, R. and Slotow, R., 2011. Relative

impacts of elephant and fire on large trees in a savanna ecosystem. *Ecosystems*, 14,

pp.1372-1381.

Skarpe, C., Aarrestad, P.A., Andreassen, H.P., Dhillion, S.S., Dimakatso, T., du Toit, J.T.,

Duncan, J.H., Hytteborn, H., Makhabu, S., Mari, M. and Marokane, W., 2004. The return

of the giants: ecological effects of an increasing elephant population. *AMBIO: A Journal*

of the Human Environment, 33(6), pp.276-282.

Sorokina, H.E., Nunes, M.H., Heiskanen, J., Munyao, M., Mwang'ombe, J., Pellikka, P.,

Raumonen, P. and Maeda, E.E., 2024. East African megafauna influence on vegetation

structure permeates from landscape to tree level scales. *Ecological Informatics*, 79,

p.102435.

Staver, A.C., Abraham, J.O., Hempson, G.P., Karp, A.T. and Faith, J.T., 2021. The past,

present, and future of herbivore impacts on savanna vegetation. *Journal of Ecology*,

109(8), pp.2804-2822.

Staver, A. C. and Bond, W. J., 2014. Is there a 'browse trap'? Dynamics of herbivore impacts

on trees and grasses in an African savanna. *Journal of Ecology*, 102(3), pp.595-602.

Stoinski, T.S., Daniel, E. and Maple, T.L., 2000. A preliminary study of the behavioral

effects of feeding enrichment on African elephants. *Zoo Biology: Published in affiliation*

with the American Zoo and Aquarium Association, 19(6), pp.485-493.

Stuart-Hill, G. C., 1992. Effects of elephants and goats on the Kaffrarian succulent thicket of

the eastern Cape, South Africa. *Journal of Applied Ecology*, pp. 699-710.

Thompson, K.E., Ford, A., Esteban, G., Poupard, A., Zoon, K. and Pettorelli, N., 2022.

Impacts of African savannah elephants (*Loxodonta africana*) on tall trees and their

recovery within a small, fenced reserve in South Africa. *African Journal of Ecology*,

60(3), pp.357-366.

Trinkel, M., Funston, P., Hofmeyr, M., Hofmeyr, D., Dell, S., Packer, C. and Slotow, R.,

2010. Inbreeding and density-dependent population growth in a small, isolated lion

population. *Animal Conservation*, 13(4), pp.374-382.

Tripathi, H. G., Mzumara, T. I., Martin, R. O., Parr, C. L., Phiri, C. and Ryan, C. M., 2019.

Dissimilar effects of human and elephant disturbance on woodland structure and

functional bird diversity in the mopane woodlands of Zambia. *Landscape Ecology*, 34,

357-371.

Trollope, W.S.W., Trollope, L.A., Biggs, H.C., Pienaar, D. and Potgieter, A.L.F., 1998.

Long-term changes in the woody vegetation of the Kruger National Park, with special

reference to the effects of elephants and fire. *Koedoe*, 41(2), pp.103-112.

Valeix, M., Fritz, H., Sabatier, R., Murindagomo, F., Cumming, D. and Duncan, P., 2011.

Elephant-induced structural changes in the vegetation and habitat selection by large

herbivores in an African savanna. *Biological Conservation*, 144(2), pp.902-912.

Vanak, A. T., Shannon, G., Thaker, M., Page, B., Grant, R. and Slotow, R., 2012.

Biocomplexity in large tree mortality: interactions between elephant, fire and landscape in

an African savanna. *Ecography*, 35(4),pp.315-321.

Van Langevelde, F., Van De Vijver, C.A., Kumar, L., Van De Koppel, J., De Ridder, N., Van

Andel, J., Skidmore, A.K., Hearne, J.W., Stroosnijder, L., Bond, W.J. and Prins, H.H.,

2003. Effects of fire and herbivory on the stability of savanna ecosystems. *Ecology*, 84(2),

pp.337-350.

Watson, L.H., Cameron, M.J. and Iifo, F., 2020. Elephant herbivory of knob-thorn *(Senegalia*

nigrescens) and ivory palm (*Hyphaene petersiana*) in Bwabwata National Park, Caprivi,

Namibia: The role of ivory palm as a biotic refuge. *African Journal of Ecology*, 58(1),

pp.14-22.

Wigley, B.J., Coetsee, C., Fritz, H. and Bond, W.J., 2014. Herbivores shape woody plant

communities in the Kruger National Park: lessons from three long-term exclosures.

Koedoe: African protected area conservation and science, 56(1), pp.1-12.

Zhang, B., Liu, X., DeAngelis, D. L., Zhai, L., Rayamajhi, M. B. and Ju, S., 2018. Modeling

the compensatory response of an invasive tree to specialist insect herbivory. *Biological*

Control, 117, 128-136.

Tables

Table 1: Species-specific Spearman rank correlation (ρ) between tree height and canopy volume and their significance (P) in Madikwe Game Reserve (MGR) and Barokologadi Communal Property (BCP), South Africa. "Old": data collected in MGR in 2000 and 2001 at relatively low elephant density; "New", data collected in MGR in 2019 at relatively high elephant density; "Out", data collected in BCP in 2019 in the absence of elephants.

Species	ρ_{old}	P_{old}	ρ_{new}	P_{new}	ρ_{out}	P_{out}
Combretum hereroense	0.95	< 0.01	0.9	< 0.01	0.94	< 0.01
Dichrostachys cinerea	0.91	< 0.01	0.95	< 0.01	0.86	< 0.01
Euclea undulata	0.82	< 0.01	0.89	< 0.01	0.97	< 0.01
Flueggea virosa	0.85	< 0.01	0.65	< 0.01	0.78	< 0.01
Grewia flava	0.82	< 0.01	0.87	< 0.01	0.72	< 0.01
Grewia monticola	0.89	< 0.01	0.9	< 0.01	0.84	< 0.01
Gymnosporia buxifolia	0.91	< 0.01	0.8	< 0.01	0.62	< 0.01
Senegalia erubescens	0.93	< 0.01	0.94	< 0.01	0.94	< 0.01
Senegalia mellifera	0.97	< 0.01	0.68	0.01	0.92	< 0.01
Vachellia karroo	0.76	< 0.01	0.86	< 0.01	0.92	< 0.01
Vachellia tortilis	0.94	< 0.01	0.9	< 0.01	0.93	< 0.01
Ximenia americana	0.77	< 0.01	0.93	< 0.01	0.81	< 0.01
Ziziphus mucronata	0.93	< 0.01	0.95	< 0.01	0.92	< 0.01

Table 2: Effect of the Elephant Use on the species-specific Asymptotic Change in Growth

Rate (*B*) and Logged Initial Growth Rate (*I*) for the tree Height - tree Diameter and the

Canopy Volume - tree Diameter relationships according to the treatment, in Madikwe Game

Reserve (MGR) and Barokologadi Communal Property (BCP), South Africa. "Old": data

collected in MGR in 2000 and 2001 at relatively low elephant density; "New", data collected

in MGR in 2019 at relatively high elephant density.

Treatment	Allometric relationship	Allometric coefficient	Estimate ± SE	P
Old	Height - Diameter	B	0.18 ± 0.21	0.38
New	Height - Diameter	B	0.00 ± 0.10	0.99
Old	Height - Diameter	I	-0.14 ± 0.35	0.68
New	Height - Diameter	I	-0.08 ± 0.17	0.64
Old	Canopy Volume - Diameter	B	-0.14 ± 0.49	0.77
New	Canopy Volume - Diameter	B	-0.04 ± 0.24	0.85
Old	Canopy Volume - Diameter	I	-0.22 ± 0.83	0.79
New	Canopy Volume - Diameter	I	-0.01 ± 0.40	0.99

Figure legends

Figure 1: Variation of a) the logged tree Height and b) the logged Canopy Volume, between the three treatments in Madikwe Game Reserve (MGR) and Barokologadi Communal Property (BCP), South Africa. "Old", data collected in MGR in 2000 and 2001 at relatively low elephant use; "New", data collected in MGR in 2019 at relatively high elephant use; "Out", data collected in BCP in 2019 in the absence of elephants. All 13 tree species were pooled.

Figure 2: Asymptotic Change in Growth Rate (B) and Logged Initial Growth Rate (I) according to the treatment in Madikwe Game Reserve (MGR) and Barokologadi Communal Property (BCP), South Africa for the tree Height - tree Diameter and the Canopy Volume - tree Diameter relationships: a) Asymptotic Change in Growth Rate (B) for the tree Height - tree Diameter relationship; b) Logged Initial Growth Rate (I) for the tree Height - tree Diameter relationship; c) Asymptotic Change in Growth Rate (B) for the Canopy Volume - tree Diameter relationship; d) Logged Initial Growth Rate (I) for the Canopy Volume - tree Diameter relationship. "Old": data collected in MGR in 2000 and 2001 at relatively low elephant use; "New", data collected in MGR in 2019 at relatively high elephant use; "Out", data collected in BCP in 2019 in the absence of elephants. Letters denote significant differences between treatments, the absence of letters denoting the absence of significant difference.

Figure 3: Species-specific Asymptotic Change in Growth Rate (B) and Logged Initial Growth Rate (I) according to the Elephant Use in each treatment in Madikwe Game Reserve (MGR) and Barokologadi Communal Property (BCP), South Africa, for the tree Height - tree

Diameter and the Canopy Volume - tree Diameter relationships: a) Asymptotic Change in

Growth Rate (B) for the tree Height - tree Diameter relationship; b) Logged Initial Growth

Rate (I) for the tree Height - tree Diameter relationship; c) Asymptotic Change in Growth

Rate (B) for the Canopy Volume - tree Diameter relationship; d) Logged Initial Growth Rate

(I) for the Canopy Volume - tree Diameter relationship. "Old": data collected in MGR in

2000 and 2001 at relatively low elephant use, in green; "New", data collected in MGR in

2019 at relatively high elephant use, in red. Dots correspond to the observed values, solid

lines correspond to the predicted values from the model, shaded areas correspond to the 95 %

confidence intervals of the model predictions.

Figures

Figure 1

Figure 2

Figure 3

698