Starvation of the bacterium *Vibrio atlanticus* induces simultaneous attacks

on the dinoflagellate *Alexandrium pacificum*

Jean Luc Rolland[1, 2†*], Estelle Masseret[3], Mohamed Laabir[3], Guillaume Tetreau[4], Benjamin Gourbal[4],

Anne Thebault[5], Eric Abadie[6], Alice Rodrigues-Stien[7,8], Carole Veckerlé[7,8], Elodie Servanne-Meunier[1],

Delphine Destoumieux-Garzón[1], Arnaud Lagorce[4,9] , Raphaël Lami[7,9].

Affiliations

[1]IHPE, Univ. Montpellier, CNRS, IFREMER, Université de Perpignan Via Domitia, Montpellier, France.

[2]MARBEC, Univ. Montpellier, IRD, IFREMER, CNRS, Sète, France.

[3]MARBEC, Univ. Montpellier, IRD, IFREMER, CNRS, Montpellier, France.

[4]IHPE, Univ Montpellier, CNRS, IFREMER, Université de Perpignan Via Domitia, Perpignan, France

[5]Agence nationale de sécurité sanitaire de l'alimentation, de l'environnement et du travail (ANSES),

ANSES-DER 27-31, avenue du Général Leclerc, 94701 Maisons-Alfort, France

[6]IFREMER, 9 rue Jean Bertho, 97822, Le Port, la Réunion, France

[7]Sorbonne Université, CNRS, LBBM USR3579, Observatoire Océanologique de Banyuls sur Mer,

Avenue Pierre Fabre, 66650 Banyuls-sur-Mer, France.

[8]Sorbonne Université, CNRS, Bio2MAR, Observatoire Océanologique de Banyuls sur Mer, Avenue Pierre

Fabre, 66650 Banyuls-sur-Mer, France.

[9]These authors contributed equally

†Current address

*Corresponding author: Jean Luc Rolland

Email: Jean.Luc.Rolland@ifremer.fr

ABSTRACT

Phytoplankton serve as a source of nutrients for bacteria in the marine environment. The interactions between algae and bacteria are known to include mutualism, commensalism, competition or antagonism. This occurs in the microenvironment surrounding phytoplankton cells, the phycosphere, an interface rich in nutrients and organic molecules exuded by the cell. Here, based on *in situ* observations and on an *in vitro* interaction study, we report on a novel form of starvation-induced hunting that the cells of selected Vibrio species exert on dinoflagellates. The results showed that *Vibrio atlanticus* was capable of attacking and killing the dinoflagellate *Alexandrium pacificum* ACT03. Briefly, the observed mechanism of algal-killing consists of first, the 'immobilization stage' involving the secretion of algicidal metabolites that disrupt the flagella of the algae. In the 'attack stage', Vibrios simultaneously surround algal cells at high density for a brief period without invading them. Finally, the 'killing stage' in which the lysis and consumption of the dinoflagellates occur. By using a combination of biochemical, proteomic, molecular and fluorescence microscopy approaches, we showed that this relationship is not related to the decomposition of algal organic matter, Vibrio quorum sensing pathways, toxicity of the algae or pathogenicity of the bacterium but is conditioned by nutrient stress, iron availability and linked to the iron-vibrioferrin transport system of *Vibrio atlanticus*. This is the first evidence of a new mechanism that could be involved in regulating Alexandrium spp. blooms and giving Vibrio a competitive advantage in obtaining nutrients from the environment. The interaction model we propose here suggests that Vibrio could play a role in regulating the proliferation of Alexandrium spp., giving it a competitive advantage in obtaining nutrients from the environment.

KEYWORDS

Interaction; Bacterium; Alga; Predation; Environment; Iron-related predation

INTRODUCTION

Harmful algal blooms (HABs) have experienced an increase in their occurrence, intensity, and geographical distribution on a global scale, resulting in adverse environmental, health, and socioeconomic impacts (Marampouti et al., 2021). HABs have a considerable impact on human health as a result of direct exposure to volatile toxins or by toxic seafood consumption (Burkholder et al., 2018). From an ecological point of view, the expansion of HABs can result in the erosion of biodiversity, because they cause massive mortality of marine species and they are generally monospecific in nature (Chai et al., 2020). In coastal areas, understanding the biological interactions that control toxic algal blooms is therefore a major ecological challenge. Among HAB-causing organisms, a number of *Alexandrium* species have been placed on the list of invasive Mediterranean species. Among them, *Alexandrium pacificum* is a flagellated eukaryotic unicellular organism that together with *Alexandrium tamarense* and *Alexandrium fundyense* forms the "Alexandrium tamarense" complex responsible for paralytic shellfish poisoning worldwide (Hadjadji et al., 2020). Since 1998, *A. pacificum* (former *A. catenella*) was monitored by the French phytoplankton observation and monitoring network (Rephy) in the Thau lagoon (French Mediterranean) because it produces paralytic shellfish toxins (PSTs) resulting in paralytic shellfish poisoning (PSP) syndrome.

Laanaia (Laanaia et al., 2013) showed that in Thau lagoon, a water temperature around 20°C for several days and organic and inorganic nutrients in sufficient concentrations are parameters favoring the development of *A. pacificum*, whose massive blooms occur in autumn. Algal blooms are seasonal events resulting in a rapid increase in the concentration of a species of algae in an aquatic environment. Depending on the species of phytoplankton, tolerance to physicochemical parameters varies, which influences when blooms occur (Leblad et al., 2020). Interestingly, although the collapse of phytoplankton blooms has been previously attributed to viruses (Pal et al., 2020), some ecological studies have suggested an important role of algicidal bacteria (Su et al., 2007; Wang et al., 2010). Among them several are belonging to the Vibrio genus (Li et al., 2014; Wang et al., 2020).

Vibrio (class γ-proteobacteria) are common microorganisms in marine systems worldwide (Baker-Austin

et al., 2017; Mavian et al., 2020), where they are important components of the food chain, particularly in

biodegradation, nutrient regeneration and biogeochemical cycles (Oberbeckmann et al., 2012). Vibrio is

one of the most studied bacterial taxa due to their ubiquity in coastal marine systems and their capacity to

cause infections in humans and animals, leading sometimes to epizootic or zoonotic epidemics (LeRoux et

al., 2015; Mavian et al., 2020). Vibrio are extremely adaptable to their environment (Johnson, 2013). The

main factors influencing their occurrence and distribution in water are temperature, salinity, nutrient

availability (Wang et al., 2020), multiple strategies such as biofilm formation on biotic and abiotic

surfaces (Espinoza-Vergara et al., 2020), or interactions with a multitude of other organisms such as

eukaryotic predators (Drebes Dörr and Blokesch, 2020) or plankton (Lopez-Joven et al., 2018) are used by

Vibrio in the environment. There is also evidence that global climate change has increased Vibrio-

associated illnesses affecting humans and animals (Brumfield et al., 2021; Muhling et al., 2017).

However, the drivers and dynamics of Vibrio survival and propagation in the marine environment are not

yet fully understood.

A substantial number of research articles have highlighted the potential of γ-proteobacteria to exert

algicidal activity against dinoflagellates, supporting the hypothesis that γ-proteobacteria such as Vibrio

play a role in the control of algal blooms *in situ* (Coyne et al., 2022). However, the mechanisms behind

Vibrio-driven algal lysis in the environment remain to be elucidated. Particularly, it is unclear how in the

water column, algicidal compounds secreted by bacteria can concentrate around the algae to exert their

lytic effect.

This study aims to describe observations made in the natural environment between Vibrio bacteria and

Alexandrium algal blooms, and to determine *in vitro* the main factors involved in this relationship. Using

a combination of biochemical, proteomic, molecular, and fluorescence microscopy approaches, we

explored the role in algal toxicity, bacterial pathogenicity and the quorum sensing pathway on this

relationship and showed the important role of nutrient stress and the iron uptake pathway in this unique

Vibrio/Alexandrium interaction.

METHODS

Quantification of *Alexandrium* algae and Vibrio bacteria in the environment by qPCR.

Seawater samples were collected in the Thau Lagoon (southern France, a shallow Mediterranean

ecosystem open to the sea (Abadie et al., 1999), during spring and autumn 2015. Briefly, samples were

collected from the subsurface (-50 cm) near an oyster table at a phytoplankton surveillance site (part of the

REPHY network, N 43°26.058' and E 003°39.878'). Once a week during spring and autumn 2015, during

field sampling campaigns, 20 L of water was filtered on board through a 180 µm pore-size nylon

membrane. At the laboratory, according to Lopez-Joven et al. (Lopez-Joven et al., 2018) seawater was

fractionated into two size classes as follows: 2 L of the above filtrate was filtered through a 0.8 µm pore-

size polycarbonate Whatman Nuclepore membrane to obtain organisms in the 0.8–180 µm range

corresponding to plankton-associated Vibrio and living *Alexandrium* forms. Then, the filtrate from the 0.8

µm membrane was filtered again through a 0.2 µm pore-size polycarbonate Whatman Nuclepore

membrane until the membrane was saturated. *Alexandrium* cells, ranging from 25 to 40 µm, belong to the

microphytoplankton and are therefore retained in the 0.8–180 µm fraction. Any Vibrio cells potentially

associated with or attached to *Alexandrium* cells will also be retained in this fraction. Vibrio cells are

approximately 0.5-0.8 µm thick. The fraction between 0.2 and 0.8 µm therefore includes the free-living

Vibrio. The bacterial population collected on 0.8-µm-pore-size filters was designated the particle-

associated community, and the population on 0.2-µm-pore-size filters was designated as the free-living

community. Membranes (in triplicate) were then conserved in 500 µL of 100% EtOH at -20°C.

Environmental DNA (eDNA) was extracted from the MF Millipore membrane using the Macherey-Nagel

NucleoSpin Tissue Kit and resuspended in 100 µL of water. The samples were then stored at -20°C after

eDNA quantity and purity were assessed using a NanoDrop system (NanoDrop Technologies,

Wilmington, DE, USA). PCR amplification reactions were done on a Roche LightCycler 480 Real-Time

thermocycler (qPHD platform, University of Montpellier, France) using specific primer pairs (Table 3).

Typically, the reactions contained 1 µL of template DNA (the DNA concentration for all samples varied

from 1 to 40 µg mL^{-1}), 0.5 µL of each primer (3.33 µM) and 4 µL of reaction mixture (SYBR Green

Master Mix) in a total volume of 6 µL. The reaction parameters were as follows: 5 minutes at 95°C (initial

denaturation) and 40 cycles of 10 s at 95°C (denaturation), 10 s at the corresponding hybridization

temperature (Table 3) and 10 s at 72°C (elongation). Melting curve profiles were generated by increasing

the temperature from 65°C to 95°C at 0.5°C s^{-1}. Amplification products were analysed using LightCycler

software (Roche Diagnostics). *Vibrio spp*. and *A. pacificum* and *A. tamarense* were quantified by

constructing calibration curves based on DNA from the *V. atlanticus* LGP32 reference strain (former *V.*

tasmaniensis LGP32) and from the *A. pacificum* reference strain (ACT03: *A. catenella* strain isolated from

Thau in 2003) and the *A. tamarense* reference strain (ATT07: *A. tamarense* isolated from Thau in 2007)

(not shown).

Strains and growth conditions.

Vibrio strains. Wild-type and isogenic mutants of *V. atlanticus* LGP32 (Table 1) were used in this study.

Deletion-mutants included Δ*luxR*, Δ*luxS*, Δ*luxM* and Δ*pvuB* isogenic strains. The Δ*pvuB* mutant was

constructed here by allelic exchange as described previously by Le Roux (Le Roux et al., 2007). We also

used *V. atlanticus* LGP32 carrying the pSW3654T-GFP plasmid (Le Roux et al., 2007), hereafter referred

to as *V. atlanticus* LGP32-GFP. Bacterial strains were grown at 22 ± 1°C in Zobell medium (0.38 µM iron

(III)). When needed, 25 µg mL^{-1} chloramphenicol (Cm) was added to cultures of *V. atlanticus* LGP32-

GFP (Le Roux et al., 2007).

Phytoplankton strains. Non-axenic phytoplankton species (Table 2) were grown in batch culture in

enriched natural sea water (ENSW, 6.55 µM iron (III)) with a salinity of 36 practical salinity units (PSU)

at $22 \pm 1°C$ under cool white fluorescent illumination (100 µmol photons m^{-2} s^{-1}) and a 12 h:12 h

light:dark cycle (Harrison et al., 1980). The algae were used for experiments in their exponential growth

phase.

Co-culture assay

For each tested phytoplankton species (Table 2), 2 x 10^4 cells harvested in their exponential growth phase

(doubling time between 5 and 7 days) were placed in 20 mL of ENSW medium in a 50 mL suspension

culture flask (Cellstar® PS, Greiner bio-one). After incubation for 24 h at $22 \pm 1°C$ under cool white

fluorescent illumination, 40 µL of Vibrio strains (Table 1) grown for 12, 36, 60 and 156 h in Zobell

medium or in Zobell medium supplemented with $FeCl_3$ (6 µM iron (III)) or with boron (H_3BO_3, 0.47 mM)

or the corresponding culture supernatant were added to the phytoplankton cells (Table 2). After incubation

at $22 \pm 1°C$ for 0, 15, 30, 45 and 60 min under cool white fluorescent lights, living, non-swimming,

attacked and lysed phytoplankton cells were counted in a sedimentation chamber under an inverted

microscope. The number of lysed cells corresponded to phytoplankton cells showing disrupted

membranes. Non-swimming algae were not counted as lysed cells. For Vibrio analysis, 100 µL of a 1:10

serial dilution mixture in ENSW (from 10^{-2} to 10^{-10}) was plated on Vibrio Selective TCBS (thiosulfate-

citrate-bile salts-sucrose) agar (in triplicate). After incubation for 24 h at $22 \pm 1°C$, the number of living

Vibrio cells was determined by counting colony-forming units (CFUs). The data come from three

independent experiments using independent phytoplankton cultures and independent bacterial cultures.

Microscope observations

The dynamic of the interaction between *A. pacificum* ACT03 isolated from the French Thau Lagoon,

south of France (Laabir et al., 2011) and *V. atlanticus* LGP32, which is an oyster pathogen isolated from

the French Atlantic coast (Gay et al., 2004) and present in the Thau Lagoon (Lopez-Joven et al., 2018)

was surveyed. As the *A. pacificum* ACT03 strain (Table 2) used in the study is not axenic, there is

potential for bacteria other than *V. atlanticus* LGP32 to be present in the experiments. To elucidate the

interaction without thoroughly accounting for the non-axenic cells, interaction was observed under

a Zeiss Axio upright fluorescence microscope equipped with an AxioCamMRm 2 digital microscope

camera using *V. atlanticus* LGP32 tagged with green fluorescent protein (GFP). Lasers were used at

excitation wavelength (λex) 488 nm for GFP (emission wavelengths (λem): 505–530 nm) and λex 532 nm

for plankton chloroplasts (λem 560–630 nm). Images were taken sequentially to avoid cross-

contamination between fluorochromes. Sequences of images were merged during the Vibrio-*Alexandrium*

interaction using ZEN 2012 (blue edition) software. Interaction events between Vibrio strains and

phytoplankton strains were also observed under a Leica TCS SPE confocal laser scanning system

connected to a Leica DM 2500 upright microscope camera (Montpellier RIO Imaging Platform,

University of Montpellier, France).

Comparative proteomic analysis

Vibrio sampling and protein extraction. V. atlanticus LGP32 was grown for 60 h at 22°C in artificial

seawater (high nutrient stress, cond. 1) or 12 h at 22°C in Zobell media (no nutrient stress, cond. 2). After

10 min centrifugation at 8000 rpm, crude protein extracts of *V. atlanticus* LGP32 in each culture condition

(triplicates) were obtained by sonication on ice at 20% amplitude for 20 s in 200 µL of ice-cold denaturing

buffer (7 M urea, 2 M thiourea, 4% CHAPS in 30 mM Tris-HCl, pH 8.5) and clarified by centrifugation at

2000 x *g*, 15 min, 4°C. The protein concentration of the supernatant was estimated using the 2D Quant Kit

(Cytiva™, MERCK) and samples were stored at -80°C until use.

Two-dimensional gel electrophoresis (2D gel). Protein extracts were individually analysed on 2D gel

electrophoresis (6 gels per condition each corresponding to different biological replicates). To do so, 100

µg of proteins from each extract was added to rehydration buffer (7 M urea, 2 M thiourea, 4% CHAPS, 65

mM DTT) for a total volume of 350 µL. They were then individually loaded onto 17 cm isoelectric

focusing strips (Bio-Rad) with a stabilized non-linear pH ranging from 3 to 10. Due to the high

complexity of the protein profile in the acidic part (left) of the gel pH 3–10 (Fig. S3), we conducted

additional 'close-up' analyses in gels using 17 cm isoelectric focusing strips (Bio-Rad) with a narrower,

stabilized pH gradient ranging from 4 to 7. Strips were rehydrated passively for 5 h at 22°C, followed by

active rehydration for 14 h under a 50 V current at 22°C (to help large proteins enter the strips).

Thereafter, isoelectrofocusing was carried out using the following programme: 50 V for 1 h, 250 V for 1

h, 8000 V for 1 h and a final step at 8000 V for a total of 140 000 V.h with a slow ramping voltage

(quadratic increasing voltage) at each step. Focused proteins were reduced by incubating the strip twice in

equilibration buffer (1.5 M Tris, 6 M urea, 2% SDS, 30% glycerol; bromophenol blue, pH 8.8) containing

DTT (130 mM) at 55°C. Then, they were alkylated by incubation with equilibration buffer containing

iodoacetamide (135 mM) on a rocking agitator (400 rpm) at room temperature protected from light.

Proteins were also separated according to their molecular weight (second dimension) on 12%

acrylamide/0.32% piperazine diacrylamide gels run at 25 mA per gel for 30 min followed by 75 mA per

gel for 8 h using a Protean II XL system (Bio-Rad). Gels were stained using an MS-compatible silver

staining protocol and scanned using a ChemiDoc MP Imaging System (Bio-Rad) associated with Image

Lab software version 4.0.1 (Bio-Rad).

Comparative bioinformatics analysis of 2D gels. Twelve gels (six per condition) were selected for

comparative analysis on PD-Quest v. 7.4.0 (Bio-Rad) to identify changes in protein abundance between

the proteomic profiles of *V. atlanticus* LGP32 cultured in contrasting nutrient conditions (ENSW/Zobell).

Spots whose mean intensity across six replicates per strain was two times higher or lower than those from

the other strain, with a *P* <0.01 (Mann-Whitney U-test), were considered significantly different in terms of

abundance between the two conditions (quantitative difference). Differentially represented spots were then

excised from the gels, destained, trypsin-digested and the obtained peptides were identified by tandem

mass spectrometry (MS-MS) using the PISSARO platform facility (University of Rouen, France). To

identify protein(s) present in each spot, the obtained peptides were compared with *V. atlanticus* LGP32

reference genome *(https://vibrio.biocyc.org/)*. The genes whose peptides matched strongly were retrieved

and used for a BLASTx query against non-redundant databases to determine the protein identity of the

best match. A gene was considered as strongly matched when at least two peptides matched the sequence

with a coverage of > 6%. Their theoretical isoelectric point (pI) and molecular weights were also

calculated using the Expasy server (https://www.expasy.org/) to compare them with the location of the

spot on the gel. Altogether, these complementary analyses made it possible to characterize the protein

identity of each spot with confidence.

Gene expression analysis

Vibrio sampling and RNA extraction.   *V. atlanticus* LGP32 was grown in Zobell media for 36 h and 60 h

at 22°C (decline phase of growth, nutrient stress) or for 12 h at 22°C (exponential growth phase, no

nutrient stress). Total RNA was isolated from *V. atlanticus* LGP32 using the standard TRIzol method

(Invitrogen Life Technologies SAS, Saint-Aubin, France) and then treated with DNase (Invitrogen) to

eliminate genomic DNA contamination. After sodium acetate precipitation, the quantity and quality of the

total RNA were determined using a NanoDrop spectrophotometer and agarose gel electrophoresis.

Following heat denaturing (70°C for 5 minutes), reverse transcription was performed using 1 μg of RNA

prepared with 50 ng μL^{-1} oligo-(dT) 12–18 mer in a 20-μL reaction containing 1 mM dNTPs, 1 unit μL^{-1}

RNAseOUT and 200 units μL^{-1} Moloney murine leukaemia virus reverse transcriptase (M-MLV RT) in

reverse transcriptase buffer, according to the manufacturer's instructions (Invitrogen Life Technologies

SAS, Saint-Aubin, France).

PCR amplification. Amplification reactions were analysed using a Roche LightCycler 480 Real-Time

thermocycler (Bio-Environnement platform, University of Perpignan, France). In this study, several PCR

primer pairs were designed using Primer3 software (optimal primer size: 20 bases; Tm: 60°C; primer

GC%: 50; 2GC clamp and product size range: 150–200 bp) and calibrated with *V. atlanticus* LGP32

genomic DNA (Table 1). To determine the qPCR efficiency of each primer pair used, standard curves

were generated using seven serial dilutions of genomic DNA (10^{-2}, 10^{-3}, 10^{-4}, 10^{-5}, 10^{-7} and 10^{-8}) (not

shown); the qPCR efficiencies of the tested genes varied between 1.85 and 2.08 (Table 3). For gene

expression, reverse transcription was performed with 1 µg of total RNA using random hexamers and

SuperScript IV reverse transcriptase (Invitrogen). The total qPCR reaction volume was 10 µL and

consisted of 5 µL of cDNA (diluted 1:5), 2.5 µL of SensiFAST SYBR No-ROX Mix (Bioline) and 100

nM or 300 nM PCR primer pair (Table 3). The reaction parameters were as follows: 2 min at 95°C (initial

denaturation) and 40 cycles of 5 s at 95°C (denaturation), 10 s at 59°C (annealing) and 20 s at 72°C

(elongation). The specificity of each PCR was checked by measuring fluorescent signals during melting

curve analysis (PCR product heated from 65°C to 95°C continuously and slowly at $0.1°C\ s^{-1}$). Relative

expression was calculated by normalization to the expression of two constitutively expressed

housekeeping genes, namely, 6PKF (VS_2913) and CcmC (VS_0852), using the delta-delta threshold

cycle (ΔΔCt) method (Pfaffl, 2001).

Detection of quorum-sensing signalling molecules.

Vibrio culture. To detect the QS molecules (AI-2, AI-1 and CAI-1) *V. atlanticus* LGP32 was grown

in Zobell media for 12 h (exponential growth phase, control) and 60 h (decline phase of growth, nutrient

stress).

AI-2 analysis. Bioluminescence assay using the QS bioluminescent of *Vibrio campbellii* MM32

(*luxN*::Cm, *luxS*::Tn5Kan) was used to detect AI-2 molecules in culture supernatants. Briefly, *Vibrio*

cultures were centrifuged at 17,000 x *g* for 10 min, and the resulting supernatants were filtered on 0.22

µm. Then 20 µL of the filtrates were mixed with 180 µL of *V. campbellii* MM32 diluted 1:5000 then

incubated at 30°C and 100 rpm. Luminescence and cell density (OD620) were collected in triplicate and

analysed according to Tourneroche et al.(Tourneroche et al., 2019).

AI-1 and CAI-1 extraction and LC-MS analysis. Chemical analyses were conducted with a Q Exactive

Focus Orbitrap System coupled to an Ultimate 3000™ ultrahigh-performance liquid chromatography

(UHPLC) system (Thermo Fisher Scientific) according to Rodrigues et al. (Rodrigues et al., 2022).

Briefly, ethyl acetate (2 mL) was added into each culture (2 mL). This mixture was shaken overnight at

room temperature (150 rpm). The two phases were then separated and the aqueous phase was extracted

once again. The two obtained organic phases were pooled and the solvent was evaporated under vacuum.

The crude extracts were dissolved in 500 µL LC-MS grade methanol for analysis. The experiments were

performed using biological triplicates, each of which was analyzed in triplicate. Analyses of extracts and

standards (3 μL injected) were performed in electrospray positive ionization mode in the 50–750 m/z

range in centroid mode. The parameters were as follows: spray voltage: 3 kV; sheath flow rate: 75; aux

gas pressure: 20; capillary temperature: 350°C; heater temperature: 430°C. The analysis was conducted in

Full MS data-dependent MS2 mode (Discovery mode). Resolution was set to 70,000 in Full MS mode,

and the AGC (automatic gain control) target was set to 1×10^6. In MS2, resolution was 17,500, AGC target

was set to 2×10^5, isolation window was 0.4 m/z, and normalized collision energy was stepped to 15, 30

and 40 eV. The UHPLC column was a Phenomenex Luna Omega Polar C18 1.6 μm, 150 x 2.1 mm. The

column temperature was set to 42°C, and the flow rate was 0.4 mL min^{-1}. The solvent system was a

mixture of water (A) with increasing proportions of acetonitrile (B), with both solvents modified with

0.1% formic acid. The gradient was as follows: 1% B 3 min before injection, then from 1 to 15 min, a

gradient increase of B up to 100% (curve 5), followed by 100% B for 5 min. The flow was injected into

the mass spectrometer starting immediately after injection. All data were acquired and processed using

FreeStyle 1.5 software (Thermo Fisher Scientific).

Chemicals and solvents. N-acyl-homoserine lactones (AHL) were obtained from Cayman Chemical (Ann

Arbor, MI, USA). Stock solutions were obtained by dissolving standards in methanol or dichloromethane

(C18-AHL) at a concentration of 1 mg mL^{-1} and stored at -80°C. Standard solutions for UHPLC- high-

resolution tandem mass spectrometry (HRMS) analyses were prepared by diluting each individual

standard solution with methanol in order to obtain a concentration range from 2000 to 20 ng mL^{-1}. LC-MS

grade methanol, acetonitrile and formic acid were purchased from Biosolve (Biosolve Chimie, Dieuze,

France), analytical-grade ethyl acetate was obtained from Sigma-Aldrich. Pure water was obtained from

Elga Purelab Flex System (Veolia LabWater STI, Antony, France).

Nature of lytic compounds secreted by *V. atlanticus* LGP32.

To determine the temperature-sensitivity of the lytic compounds secreted, *V. atlanticus* LGP32 grown for

60 h in Zobell media at 22 °C was filtered through a 10 kDa membrane (Amicon$^{®}$ Ultra-4 filter unit).

The eluate containing molecules with MW below 10 kDa was then incubated in a water bath at 100°C for

30 min. Boiled filtrates (0.1 mL) were subsequently used to inoculate *A. pacificum* (ACT03 strain)

cultures, then lytic activity was observed under the Leica TCS SPE confocal laser scanning system. Zobell

media with the same treatment was used as control.

Statistical analysis. *Environmental data*. Statistical analyses were performed using R 3.6.3 software (R

Core Team, 2024) using Rstudio. The relationship between *Alexandrium* and Vibrio was explored

separately in spring and autumn. We used a generalized linear model specifying a Gaussian family. For

spring, the dataset for salinity and temperature was complete (14 periods of observation). An influential

period was detected and removed from the dataset because no dead *Alexandrium* cells were observed. The

effects of explanatory variables such as log10 (Vibrio+1), salinity and temperature were centred, reduced

and tested as fixed effects with a linear relationship. Model selection was performed using the Akaike

information criterion corrected for small sample size (AIC_c). Models were considered different whenever

the difference between their AIC_c value and the lowest AIC_c value (ΔAIC_c) was lower than 2 (Burnham

and Anderson, 2002). *Alexandrium* distribution and model residuals were checked for normal distribution

assumptions (QQ plot and Shapiro-Wilk test). For autumn, the dataset was complete for 10 periods of

observation. Salinity and temperature were missing for three periods. We explored the relationship

between *Alexandrium* and Vibrio alone using the method detailed above for spring.

In vitro data. Statistical analyses were performed using one-way ANOVA (analysed by pair) followed by

Tukey's test (Statistica 10.0 software, StatSoft, Maison-Alfort, France). *P <0.05, **P <0.01, ***P

<0.001.

RESULTS

Concomitant occurrence of *A. pacificum* ACT03, *A. tamarense* ATT07 and free-living *Vibrio spp.* in

the Thau lagoon. In the spring and autumn of 2015, in the Thau Lagoon (Fig. 1A), we detected

Alexandrium algae (*A. pacificum* ACT03 *and A. tamarense* ATT07, both alive and in degraded cell forms)

and free-living Vibrio, but no plankton-associated Vibrio were observed (Fig. 1B, Data S1). Using model

selection based on AICc, we found no significant relationship between *Alexandrium* (*A. pacificum*

ACT03, *A. tamarense*) and *Vibrio spp.* abundances in autumn. This result is consistent with Vibrio's

difficulty in growing at temperatures below 20°C, as well as with the many environmental factors that can

influence the dynamics of algae proliferation (Laanaia et al., 2013). Interestingly, in spring 2015, the mean

densities of all *Alexandrium* cells (degraded and alive) and of free-living Vibrio were positively

correlated. The lowest AICc was obtained with the model explaining degraded form of *Alexandrium*

density based on the free Vibrio density (Fig. 1C). Given that, this model is not so different from the

model with only the intercept, but better than any other linear combination with other potentially

interfering drivers, such as temperature and salinity (Fig. 1D), we searched for evidence of a relationship

between Vibrio and *Alexandrium* by studying their interaction *in vitro*.

V. atlanticus **LGP32 feeds on** *Alexandrium pacificum* **ACT03**.

To investigate whether *Alexandrium* interacts with Vibrio, we incubated in Enriched Natural SeaWater

(ENSW) *A. pacificum* ACT03 (2.0 x 10^4 cells) with *V. atlanticus* LGP32 previously grown for 12 hours in

Zobell media (initial concentration of 8.8 x 10^7 cells mL^{-1}). In interaction with *V. atlanticus* LGP32, *A.*

pacificum ACT03 cell abundance decreased significantly from 2.1 x 10^4 cells mL^{-1} after 1h of exposure to

1.1 x 10^4 cells mL^{-1} after 48 h of exposure (Fig. 2A), while the *V. atlanticus* LGP32 concentration grew

significantly after 26 h of interaction, reaching a maximum peak density of 7.6 x 10^7 CFU mL^{-1} at 34 h

(Fig. 2B). In the control experiment where *A. pacificum* ACT03 was cultured alone in ENSW, the algal

concentration remained stable over time (Fig. 2A) and no bacteria were on the corresponding TCBS plates

(Vibrio selective medium). In the control where *V. atlanticus* LGP32 was grown alone in ENSW, the

bacterial concentration decreased from $7.0 \times 10\square$ CFU/mL after 1 hour of incubation to $1.1 \times 10\square$

CFU/mL after 48 hours of incubation (Fig. 2B). These results show that the interaction between *V.*

atlanticus LGP32 and *A. pacificum* ACT03 leads to a decline in the algal population and promotes the

growth of *V. atlanticus* LGP32. This suggests that *V. atlanticus* LGP32 is able to feed on *A. pacificum*

ACT03.

***V. atlanticus* LGP32 performs attacks on *A. pacificum* ACT03**.

Epifluorescence microscopy observation of GFP-labelled *V. atlanticus* LGP32 (previously grown in

Zobell medium) in interaction showed that *V. atlanticus* LGP32 cells are capable of simultaneously

attacking *A. pacificum* ACT03 cells (Fig. 2C and Video 1). The attacks were extremely rapid, with empty

thecae (algal envelopes) observed in the medium after less than 60 s (Fig. 2D and Video 2). During the

attack, *V. atlanticus* LGP32 did not invade the algal cell but remained clustered on the cell surface (Fig.

2C).

Attack of *A. pacificum* ACT03 is activated by *V. atlanticus* LGP32 starvation.

In 2002, Martin hypothesized that nutritional stress induces bacteria to lyse algae (Martin, 2002). To test

this hypothesis, we monitored *V. atlanticus* LGP32 behaviour in response to starvation (Fig. 3). We

observed that *V. atlanticus* LGP32 in exponential growth phase (12 h of culture in Zobell medium, 8.8 x

10^7 cells mL^{-1}) did not interact with *A. pacificum* ACT03 cells for the first hour of contact (Fig. 3A, B, C).

In contrast, *V. atlanticus* LGP32 in the decline phase (36 h of culture in Zobell medium, 1.3 x 10^7 cells

mL^{-1}) induced a significant decrease in the number of motile algae cells by 8.9% after 15 min and by

43.3% after 60 min (Fig. 3A). This phenomenon corresponded to the degradation and/or disruption of

algal flagella (Video 3). The flagella no longer functioned correctly, which caused irregular swimming of

the algae (Video 3, left cell). This was followed by a complete cessation of swimming. When the

flagellum detached from the algae (Video 3, right cell), the attack occurred. With starved *V. atlanticus*

LGP32 (60 h of culture in Zobell medium, 0.6 x 10^7 cells mL^{-1}), algae immobilization was fast and

significant (91.4% in 15 min, Fig. 3A), and algae were attacked individually being targeted by *V.*

atlanticus LGP32 cells (Video 4). The percentage of cells attacked and killed peaked at 30% after 15–30

min of contact (Fig. 3B and 3B1) and then decreased. After 1 h, attacks had stopped with approximately

40% of the algal cells still alive (Fig. 3B1 and 3C). Although it remains unclear whether the attacks occur

during a specific phase of growth, it is evident that the cells are already weakened before attack as they

have all lost their flagella. An old-starved culture of *V. atlanticus* LGP32 (126 h of culture in Zobell

medium, < 0.1 x 10^7 cells mL^{-1}) significantly immobilized *A. pacificum* ACT03 cells within a few

minutes, with lysis occurring immediately (Fig. 3A and 3C), making it impossible to detect attacks by *V.*

atlanticus LGP32 (Fig. 3B). The lysis phase corresponded to initial vesicle formation followed by the

bursting of *A. pacificum* ACT03 cells (Fig. 3C and 3C1). Importantly, Vibrio densities decreased with

culture age, ruling out the possibility that the stronger predation observed in older cultures was driven by

higher bacterial densities.

We next tested whether this lytic effect was linked to Vibrio culture supernatant and mediated by

thermostable molecule(s) secreted by Vibrio. The culture supernatant of starved culture of *V. atlanticus*

LGP32 (36 h) filtered through a 10 kDa membrane and then incubated at 100°C for 30 min still possessed

its lytic properties, indicating that the algicidal compounds produced by *V. atlanticus* LGP32 are small

thermostable molecules unlikely to be lytic enzymes, or lysins able to digest the algae cell. However, these

experimental observations clearly show the key role of nutrient limitation in triggering the attack

behaviour and the secretion of lytic compounds of *V. atlanticus* LGP32.

Attack occurs on *A. pacificum* ACT03 in exponential phase of growth.

Here, we wondered whether the live/dead status of algae is important for *V. atlanticus* LGP32-mediated

attacks targeting. To this end, *A. pacificum* ACT03 in exponential growth phase was first exposed for 30

minutes to the supernatant of a 126-hour culture of *V. atlanticus* LGP32, which induced lysis of 70% of

the *A. pacificum* ACT03 cells (Figures 3C and 3C1 (arrow 2) and Video 4). Next, cells of *V. atlanticus*

LGP32 from a 60-hour culture, capable of attacking *A. pacificum* ACT03 cells (Fig. 3B), were added. For

1 hour of exposure, no attack was observed on the previously lysed algae. This result is similar to what is

observed on the Video 1 with flash attacks only on immobilized, but not degraded *A. pacificum* ACT03

cells (red), and not on lysed cells (green). In addition, no attacks occurred on cells from an old *A.*

pacificum ACT03 culture (1-month culture). Together, with the very short duration of attacks (Video 1),

these results indicate that *V. atlanticus* LGP32 attacked exponentially growing cells of *A. pacificum*

ACT03, but not decomposing algae, suggesting that this behaviour is not just an opportunistic response of

heterotrophic bacteria to organic substrates.

407   Attack is independent of quorum sensing.

Considering the simultaneous action of *V. atlanticus* LGP32 in attacks on *A. pacificum* ACT03, we tested

whether the attack process depended on the key physiological mechanism that regulates many functions in

marine microbial cells, quorum sensing (QS) (Lami, 2019; Papenfort and Bassler, 2016), a type of cell-

cell communication. Although QS is a cell-density-dependent mechanism, our results showed no attack

from a 12 h culture of *V. atlanticus* LGP32 up to a concentration of 4×10^6 Vibrio mL^{-1} (Fig 4A). Attacks

were only observed with *V. atlanticus* LGP32 from a 60 h culture at low concentration of 5×10^3 Vibrio

mL^{-1} to the highest concentration tested of 5×10^5 Vibrio mL^{-1} (Fig 4A), consistent with the hypothesis

that the attacks were independent of Vibrio density.

The analysis of the expression of genes involved in the known QS pathways in *Vibrio* cell (Fig. S3A),

highlighted that only the AI-2 pathway was induced during nutrient stress of *V. atlanticus* LGP32, because

only the expression of the AI synthase (*LuxS*) and its receptor (*LuxP*) increased significantly (Fig. 4B;

ANOVA $p < 0.05$). This was confirmed by a QS bioluminescence assay, which showed AI-2 molecules

(unquantified) in the Zobell culture supernatant of *V. atlanticus* LGP32 after 60 h of culture but not after

12 h of culture and not in the ENSW supernatant of *V. atlanticus* LGP32 after 12 or 60 h of culture.

UHPLC-HRMS/MS provided no evidence of detectable HAI-1 and CAI-1 in any experiments.

Targeted mutagenesis of key genes involved in two of the three known QS pathways in vibrios (Fig. S3),

ΔluxM (HAI-1 production), ΔluxS (AI-2 production), and ΔluxR (main high-density QS regulator), did

not result in any changes in the attack behavior of *V. atlanticus* LGP32 (Fig. 4C). Combined with the

absence of overexpression of the CqsS gene (inducible by CAI-1) involved in the last known QS pathway

in Vibrio (Fig. S3), these results indicated that the attack by *V. atlanticus* LGP32 is most likely unrelated

to QS.

Attack related to the availability of iron.

The comparative analysis of the proteome of *V. atlanticus* LGP32 incubated 60 h in artificial seawater

(ENSW) versus *V. atlanticus* LGP32 grown 12 h in Zobell nutrient-rich medium revealed 10 proteins that

were differentially abundant under these two contrasting conditions (Fig. S2). The two most down-

regulated proteins correspond to be ß-ketoacyl-(acyl-carrier-protein) synthase II (-22-fold in ENSW

compared to Zobell), a key regulator of bacterial fatty acid synthesis, and the dihydroorotase (-6.6-fold in

ENSW compared to Zobell), an enzyme essential for pyrimidine biosynthesis and thus bacterial

proliferation and growth. The low expression of these proteins in ENSW is consistent with *V. atlanticus*

LGP32 nutritional starvation. The most up-regulated protein in starved *V. atlanticus* LGP32, with an

increase of more than 6-fold, was glucosamine-6-phosphate deaminase, an enzyme involved in bacterial

energy metabolism probably necessary for its survival. Among the other up-regulated proteins, one was an

iron siderophore-binding protein (Spot 4413, Fig. S2A) corresponding to the vibrioferrin outer membrane

receptor PvuB, whose gene is part of the *pvu* operons involved in iron transport (Fig. S3B). Interestingly,

the corresponding gene *pvuB* as well as the vibrioferrin membrane receptor gene *pvuA2* (Fig. S3B) were

both significantly induced in Vibrio under nutrient stress (Fig. 4B; ANOVA p <0.01) but not the one

involved in the vibrioferrin biosynthesis, *pvsA* (Fig. 4B). Remarkably, of the 10 proteins identified by

proteomic analysis and eliminated by mutation, only elimination of PvuB prevented *V. atlanticus* from

attacking *A. pacificum* ACT03 (Fig. 4C; ANOVA p <0.001). In the absence of the *pvuB* gene, *V.*

atlanticus LGP32 was unable to simultaneously attack *A. pacificum* ACT03. In addition, *V. atlanticus*

LGP32 cells that had been washed with ENSW to remove their culture supernatant metabolites also failed

to attack *A. pacificum* ACT03 (Fig. 4C; ANOVA p <0.001), which is congruent with the hypothesis that

attacks depend on the *V. atlanticus* LGP32 vibrioferrin transport system. Finally, attacks increased

significantly when $FeCl_3$ was added to the Vibrio culture medium (Fig. 4D) but not with boron known to

be capable of being transported by vibrioferrin (see Fig. 4D). Taken together, those results are consistent

with the hypothesis that attacks are regulated by iron.

Attack is a *Vibrio spp.* behaviour specific to *Alexandrium spp*.

To evaluate the dinoflagellates specificity of the attack behaviour, a selection of *Vibrio spp.* was co-

cultured with a selection of dinoflagellate strains commonly found in the Mediterranean Sea. The results

showed that, among the *Vibrio spp.* tested (pathogenic or not) all, when under nutrient stress, were able to

secrete algicidal compounds, immobilize, attack and lyse *A. pacificum* ACT03 cells (Table 1) and no link

to their pathogenicity for fish of invertebrates was observed (Table 1). Among the sixteen dinoflagellates

species tested, only *Alexandrium spp*. (non-toxic and paralytic shellfish toxin (PST) producers) and

Gymnodinium catenatum (PST producer) were immobilized, attacked and lysed by *V. atlanticus* LGP32

(Table 2), but no link to PSTs was revealed (Table 2).

DISCUSSION

Predation is a widespread mode of interaction for survival in the natural world (Finke and Denno, 2004;

Sinclair et al., 2003). Predatory bacteria are found in a wide variety of environments and are commonly

described as feeding on other bacteria, although some cases of predation on microbial eukaryotes have

also been hypothesized (Johnke et al., 2014; Perez et al., 2016). Conceiving predators as free-living

organisms that kill other organisms and feed on them, this study suggests that Vibrio engage in a novel

form of predation in which they kill and feed on algae.

In fact, the strategy developed by Vibrio to kill algae may be reminiscent of strategies previously

described in the prokaryotes (Johnke et al., 2014). As shown in Video 1, the interaction between *V.*

atlanticus LGP32 and *A. pacificum* ACT03 proceeds in three stages (Fig. 5). The first stage, the

'**immobilization stage**', recalls the strategy used by *Streptomyces* to immobilize its prey (Kumbhar et al.,

2014) based on the secretion of algicidal metabolites that disrupt the flagella. The second stage, the 'attack

stage' corresponding to the physical contact between Vibrios and Alexandrium, is similar to the strategy

used by *Myxococcus xanthus* and Lysobacter. These bacteria may require close proximity to their prey to

cause lysis and utilize their biomass, although some can also kill prey at a distance through diffusible

secretions (Martin, 2002; Genovesi et al., 2013; Perez et al., 2016; Zhang et al., 2020). *V. atlanticus*

LGP32 also surrounds *A. pacificum* ACT03 cells at high density for a very short time, but does not invade

the algal cell. Visually, this phenomenon resembles bacteria clustering around lysed ciliate cells

(Blackburn et al., 1998). The third stage, the '**killing stage**', is similar to that of epibiotic bacterial

predators, which induce the lysis of bacteria from the outside (Rashidan and Bird, 2001). Overall, these

observations suggest that *V. atlanticus* LGP32 can exhibit a predatory behavior.

The attack behaviour of *V. atlanticus* LGP32 was linked to iron absorption, as mutants with impaired iron

absorption completely lost its ability to attack the algae. Iron is an essential element for growth in most

organisms, including phytoplankton (Martin and Fitzwater, 1988) and bacteria (Neilands, 1981), and its

concentration in seawater is known to be very low with measurements in the open ocean surface at 0.2 nM

and in deep waters at 0.6 nM (Millero, 1998).

Moreover, its low solubility in seawater limits its availability (Bruland et al., 1994; Wu and Luther, 1994).

To acquire iron, bacteria have developed systems based on the secretion (and subsequent uptake) of iron-

chelating siderophores to obtain this element from the environment (Amin et al., 2009a). Therefore, many

Vibrio spp. produce a siderophore known as vibrioferrin, which is synthesized and secreted by proteins

encoded by the *pvsABCDE* gene cluster (Fig. S3). Given that boron is known for its role in regulating a

global bacterial cellular response to phytoplankton and to bind to vibrioferrin (Romano et al., 2013;

Weerasinghe et al., 2013), we tested its potential involvement in simultaneous Vibrio attacks. Compared

to the Zobell control, no effect on the number of attacks was observed. For iron-vibrioferrin uptake,

Vibrio parahaemolyticus uses a membrane siderophore receptor, called PvuA, which is coupled to an

inner membrane ATP-binding cassette (ABC). This ABC transporter system, which is comprised of

proteins encoded by the pvuABCDE gene cluster (Fig. S3), is required for the transport of the siderophore

across the inner membrane (Tanabe et al., 2003). Siderophores are not only iron carriers but also important

regulators of virulence (Miethke and Marahiel, 2007) and mediators of bacterial interaction with

phytoplankton (Amin et al., 2009b; Kramer et al., 2020). We showed here a pivotal role of iron in the

interaction between *V. atlanticus* LGP32 and *A. pacificum* ACT03. This mirrors the mutualistic interaction

observed between *Gymnodinium catenatum* and *Marinobacter* (Amin et al., 2009b). In fact, in natural

settings, the co-occurrence of *Marinobacter* and *G. catenatum* is suggested to depend on a mutually

beneficial utilization of iron and carbon resources (Bolch et al., 2011). As in the present study, iron seems

to play a key role in the interaction. Indeed, the labile iron released through the photolysis of ferric

chelates with vibrioferrin providing a crucial iron source for phytoplankton, which need substantial

amounts of iron to support carbon fixation through photosynthesis (Amin et al., 2009a; Yang et al., 2021).

This fixed carbon, in turn, sustains the growth of both the phytoplankton and their associated bacterial

counterparts (Amin et al., 2009b; Kramer et al., 2020). Interestingly, if a general nutrient deficiency

causes attacks, iron supplementation increases this number of attacks (Figure 4D), suggesting the

importance of iron absorption in the attack behavior. Future studies should determine whether a nutrient

deficiency increases the iron absorption capacity of Vibrio bacteria and whether this could play a major

role in the attack mechanism.

This study showed that quorum sensing is not involved in microbial attacks. Thus, *V. atlanticus* LGP32

mutants lacking the genes involved in known quorum sensing pathways exhibit the same phenotype as

wild-type *V. atlanticus* LGP32, and Vibrio density does not induce attacks. However, V. atlanticus LGP32

produces AI-2 during attacks. Quorum sensing and iron acquisition are sometimes interconnected in

Vibrio (McRose et al., 2018). For example, in *Vibrio vulnificus*, the production of vulnibactin (a

siderophore) is known to be controlled by AI-2 (Kim and Shin, 2011). Similarly, AI-2 could be involved

in the production of vibrioferrin in *V. atlanticus* LGP32.

In the natural environment, associations between bacteria and algae have already been observed (Lopez-

Joven et al., 2018; Miller et al., 2005; Rosales et al., 2022; Xu et al., 2022). We have shown here that algal

attacks by Vibrio can be carried out *in vitro*. Environmental data collected in the Thau lagoon showed a

correlation between the presence of Alexandrium and that of Vibrio, suggesting that such interactions

could also occur in the marine environment. If that were the case, this behaviour would provide an

important ecological advantage to Vibrio to obtain nutrients in the environment, where *Alexandrium spp*.

and *Gymnodinium catenatum* form blooms.

With more than 30 species distributed all over the world (Anderson et al., 2012; Hallegraeff et al., 2021),

Alexandrium spp. and *Gymnodinium spp*. considered as invasive species by the Delivering Alien Invasive

Species Inventories for Europe (http://www.europe-aliens.org)), could play an unexpected and important

role in maintaining, structuring and regulating Vibrio populations in the ecosystem. In turn, Vibrio could

contribute to the regulation and control of their blooms.

To conclude, this study reveals the capacity of some *Vibrio spp*. to act as facultative predator-like bacteria

that hunt specific algae. In the current context of climate change, which is favourable to their

development, monitoring the invasive algae *Alexandrium spp*. and *Gymnodinium catenatum* should be

considered not only for their potent harmful effect on humans and animals, but also because they may

represent a potential source of nutrients for the expansion of Vibrio, particularly pathogenic species

(Lemire et al., 2015).

DATA AVAILABILITY

All data on environmental study are included in the article and/or supporting information files. All other

study data are available on request from the corresponding author.

REFERENCES

Abadie, E., Amzil, Z., Belin, C., Comps, M.A., Elzière-Papayanni, P., Lassus, P., le Bec, C., Baut, M.L., Nezan,
E., Poggi, R., 1999. Contamination de l'étang de Thau par Alexandrium tamarense (No. Rapport
884). Ifremer.
Abadie, E., Kaci, L., Berteaux, T., Hess, P., Sechet, V., Masseret, E., Rolland, J.L., Laabir, M., 2015. Effect of
Nitrate, Ammonium and Urea on Growth and Pinnatoxin G Production of Vulcanodinium
rugosum. Mar. Drugs 13, 5642–5656. https://doi.org/10.3390/md13095642
Amin, S.A., Green, D.H., Hart, M.C., Kuepper, F.C., Sunda, W.G., Carrano, C.J., 2009a. Photolysis of iron-
siderophore chelates promotes bacterial-algal mutualism. Proc. Natl. Acad. Sci. U. S. A. 106,
17071–17076. https://doi.org/10.1073/pnas.0905512106
Amin, S.A., Green, D.H., Kuepper, F.C., Carrano, C.J., 2009b. Vibrioferrin, an Unusual Marine Siderophore:
Iron Binding, Photochemistry, and Biological Implications. Inorg. Chem. 48, 11451–11458.
https://doi.org/10.1021/ic9016883
Anderson, D.M., Alpermann, T.J., Cembella, A.D., Collos, Y., Masseret, E., Montresor, M., 2012. The
globally distributed genus Alexandrium: Multifaceted roles in marine ecosystems and impacts on
human health. Harmful Algae 14, 10–35. https://doi.org/10.1016/j.hal.2011.10.012
Baker-Austin, C., Trinanes, J., Gonzalez-Escalona, N., Martinez-Urtaza, J., 2017. Non-Cholera Vibrios: The
Microbial Barometer of Climate Change. Trends Microbiol. 25, 76–84.
https://doi.org/10.1016/j.tim.2016.09.008
Ben-Gharbia, H., Yahia, O.K.-D., Amzil, Z., Chomerat, N., Abadie, E., Masseret, E., Sibat, M., Triki, H.Z.,
Nouri, H., Laabir, M., 2016. Toxicity and Growth Assessments of Three Thermophilic Benthic
Dinoflagellates (Ostreopsis cf. ovata, Prorocentrum lima and Coolia monotis) Developing in the
Southern Mediterranean Basin. Toxins 8, 297. https://doi.org/10.3390/toxins8100297
Blackburn, N., Fenchel, T., Mitchell, J., 1998. Microscale nutrient patches in planktonic habitats shown by
chemotactic bacteria. Science 282, 2254–2256. https://doi.org/10.1126/science.282.5397.2254
Bolch, C.J.S., Subramanian, T.A., Green, D.H., 2011. THE TOXIC DINOFLAGELLATE *GYMNODINIUM*
CATENATUM (DINOPHYCEAE) REQUIRES MARINE BACTERIA FOR GROWTH[1]. J. Phycol. 47, 1009–
1022. https://doi.org/10.1111/j.1529-8817.2011.01043.x
Bruland, K., Orians, K., Cowen, J., 1994. Reactive Trace-Metals in the Stratified Central North Pacific.
Geochim. Cosmochim. Acta 58, 3171–3182. https://doi.org/10.1016/0016-7037(94)90044-2
Brumfield, K.D., Usmani, M., Chen, K.M., Gangwar, M., Jutla, A.S., Huq, A., Colwell, R.R., 2021.
Environmental parameters associated with incidence and transmission of pathogenic *Vibrio spp* .
Environ. Microbiol. 23, 7314–7340. https://doi.org/10.1111/1462-2920.15716
Burkholder, J.M., Shumway, S.E., Glibert, P.M., 2018. Food Web and Ecosystem Impacts of Harmful
Algae, in: Harmful Algal Blooms. John Wiley & Sons, Ltd, pp. 243–336.
https://doi.org/10.1002/9781118994672.ch7

Burnham, K.P., Anderson, D.R. (Eds.), 2002. Summary, in: Model Selection and Multimodel Inference: A
588   Practical Information-Theoretic Approach. Springer, New York, NY, pp. 437–454.
589   https://doi.org/10.1007/978-0-387-22456-5_8
590   Chai, Z.Y., Wang, H., Deng, Y., Hu, Z., Tang, Y.Z., 2020. Harmful algal blooms significantly reduce the
591   resource use efficiency in a coastal plankton community. Sci. Total Environ. 704, 135381.
592   https://doi.org/10.1016/j.scitotenv.2019.135381
593   Coyne, K.J., Wang, Y., Johnson, G., 2022. Algicidal Bacteria: A Review of Current Knowledge and
594   Applications to Control Harmful Algal Blooms. Front. Microbiol. 13, 871177.
595   https://doi.org/10.3389/fmicb.2022.871177
596   Drebes Dörr, N.C., Blokesch, M., 2020. Interbacterial competition and anti-predatory behaviour of
597   environmental *Vibrio cholerae* strains. Environ. Microbiol. 22, 4485–4504.
598   https://doi.org/10.1111/1462-2920.15224
599   Espinoza-Vergara, G., Hoque, M.M., McDougald, D., Noorian, P., 2020. The Impact of Protozoan
600   Predation on the Pathogenicity of Vibrio cholerae. Front. Microbiol. 11, 17.
601   https://doi.org/10.3389/fmicb.2020.00017
602   Fertouna-Bellakhal, M., Dhib, A., Fathalli, A., Bellakhal, M., Chomérat, N., Masseret, E., Laabir, M., Turki,
603   S., Aleya, L., 2015. Alexandrium pacificum Litaker sp. nov (Group IV): Resting cyst distribution and
604   toxin profile of vegetative cells in Bizerte Lagoon (Tunisia, Southern Mediterranean Sea). Harmful
605   Algae 48, 69–82. https://doi.org/10.1016/j.hal.2015.07.007
606   Finke, D.L., Denno, R.F., 2004. Predator diversity dampens trophic cascades. Nature 429, 407–410.
607   https://doi.org/10.1038/nature02554
608   Gay, M., Renault, T., Pons, A.M., Le Roux, F., 2004. Two Vibrio splendidus related strains collaborate to
609   kill Crassostrea gigas: taxonomy and host alterations. Dis. Aquat. Organ. 62, 65–74.
610   https://doi.org/10.3354/dao062065
611   Genovesi, B., Mouillot, D., Laugier, T., Fiandrino, A., Laabir, M., Vaquer, A., Grzebyk, D., 2013. Influences
612   of sedimentation and hydrodynamics on the spatial distribution of Alexandrium
613   catenella/tamarense resting cysts in a shellfish farming lagoon impacted by toxic blooms.
614   Harmful Algae 25, 15–25. https://doi.org/10.1016/j.hal.2013.02.002
615   Genovesi, B., Shin-Grzebyk, M.-S., Grzebyk, D., Laabir, M., Gagnaire, P.-A., Vaquer, A., Pastoureaud, A.,
616   Lasserre, B., Collos, Y., Berrebi, P., Masseret, E., 2011. Assessment of cryptic species diversity
617   within blooms and cyst bank of the Alexandrium tamarense complex (Dinophyceae) in a
618   Mediterranean lagoon facilitated by semi-multiplex PCR. J. Plankton Res. 33, 405–414.
619   https://doi.org/10.1093/plankt/fbq127
620   Hadjadji, I., Laabir, M., Frihi, H., Collos, Y., Shao, Z.J., Berrebi, P., Abadie, E., Amzil, Z., Chomérat, N.,
621   Rolland, J.L., Rieuvilleneuve, F., Masseret, E., 2020. Unsuspected intraspecific variability in the
622   toxin production, growth and morphology of the dinoflagellate Alexandrium pacificum R.W.
623   Litaker (Group IV) blooming in a South Western Mediterranean marine ecosystem, Annaba Bay
624   (Algeria). Toxicon 180, 79–88. https://doi.org/10.1016/j.toxicon.2020.04.005
625   Hallegraeff, G.M., Anderson, D.M., Belin, C., Bottein, M.-Y.D., Bresnan, E., Chinain, M., Enevoldsen, H.,
626   Iwataki, M., Karlson, B., McKenzie, C.H., Sunesen, I., Pitcher, G.C., Provoost, P., Richardson, A.,
627   Schweibold, L., Tester, P.A., Trainer, V.L., Yñiguez, A.T., Zingone, A., 2021. Perceived global
628   increase in algal blooms is attributable to intensified monitoring and emerging bloom impacts.
629   Commun. Earth Environ. 2, 117. https://doi.org/10.1038/s43247-021-00178-8
630   Harrison, P., Waters, R., Taylor, F., 1980. A Broad-Spectrum Artificial Seawater Medium for Coastal and
631   Open Ocean Phytoplankton. J. Phycol. 16, 28–35. https://doi.org/10.1111/j.1529-
632   8817.1980.tb00724.x

Johnke, J., Cohen, Y., de Leeuw, M., Kushmaro, A., Jurkevitch, E., Chatzinotas, A., 2014. Multiple micro-
predators controlling bacterial communities in the environment. Curr. Opin. Biotechnol. 27, 185–
190. https://doi.org/10.1016/j.copbio.2014.02.003
Johnson, C.N., 2013. Fitness Factors in Vibrios: a Mini-review. Microb. Ecol. 65, 826–851.
https://doi.org/10.1007/s00248-012-0168-x
Kim, C.M., Shin, S.H., 2011. Modulation of Iron-Uptake Systems by a Mutation of luxS Encoding an
Autoinducer-2 Synthase in Vibrio vulnificus. Biol. Pharm. Bull. 34, 632–637.
https://doi.org/10.1248/bpb.34.632
Kitatsukamoto, K., Oyaizu, H., Nanba, K., Simidu, U., 1993. Phylogenetic-Relationships of Marine-
Bacteria, Mainly Members of the Family Vibrionaceae, Determined on the Basis of 16s
Ribosomal-Rna Sequences. Int. J. Syst. Bacteriol. 43, 8–19. https://doi.org/10.1099/00207713-
43-1-8
Kramer, J., Oezkaya, O., Kuemmerli, R., 2020. Bacterial siderophores in community and host interactions.
Nat. Rev. Microbiol. 18, 152–163. https://doi.org/10.1038/s41579-019-0284-4
Kumbhar, C., Mudliar, P., Bhatia, L., Kshirsagar, A., Watve, M., 2014. Widespread predatory abilities in
the genus Streptomyces. Arch. Microbiol. 196, 235–248. https://doi.org/10.1007/s00203-014-
0961-7
Laabir, M., Jauzein, C., Genovesi, B., Masseret, E., Grzebyk, D., Cecchi, P., Vaquer, A., Perrin, Y., Collos, Y.,
2011. Influence of temperature, salinity and irradiance on the growth and cell yield of the
harmful red tide dinoflagellate Alexandrium catenella colonizing Mediterranean waters. J.
Plankton Res. 33, 1550–1563. https://doi.org/10.1093/plankt/fbr050
Laanaia, N., Vaquer, A., Fiandrino, A., Genovesi, B., Pastoureaud, A., Cecchi, P., Collos, Y., 2013. Wind and
temperature controls on Alexandrium blooms (2000-2007) in Thau lagoon (Western
Mediterranean). Harmful Algae 28, 31–36. https://doi.org/10.1016/j.hal.2013.05.016
Labreuche, Y., Le Roux, F., Henry, J., Zatylny, C., Huvet, A., Lambert, C., Soudant, P., Mazel, D., Nicolas, J.-
L., 2010. Vibrio aestuarianus zinc metalloprotease causes lethality in the Pacific oyster
Crassostrea gigas and impairs the host cellular immune defenses. Fish Shellfish Immunol. 29,
753–758. https://doi.org/10.1016/j.fsi.2010.07.007
Lami, R., 2019. Quorum Sensing in Marine Biofilms and Environments, in: Quorum Sensing, Academic
Press. Elsevier, pp. 55–96. https://doi.org/10.1016/B978-0-12-814905-8.00003-4
Le Roux, F., Binesse, J., Saulnier, D., Mazel, D., 2007. Construction of a *Vibrio splendidus* Mutant Lacking
the Metalloprotease Gene *vsm* by Use of a Novel Counterselectable Suicide Vector. Appl.
Environ. Microbiol. 73, 777–784. https://doi.org/10.1128/AEM.02147-06
Leblad, B.R., Amnhir, R., Reqia, S., Sitel, F., Daoudi, M., Marhraoui, M., Abdellah, M.K.O., Veron, B., Er-
Raioui, H., Laabir, M., 2020. Seasonal variations of phytoplankton assemblages in relation to
environmental factors in Mediterranean coastal waters of Morocco, a focus on HABs species.
Harmful Algae 96, 101819. https://doi.org/10.1016/j.hal.2020.101819
Lemire, A., Goudenege, D., Versigny, T., Petton, B., Calteau, A., Labreuche, Y., Le Roux, F., 2015.
Populations, not clones, are the unit of vibrio pathogenesis in naturally infected oysters. Isme J.
9, 1523–1531. https://doi.org/10.1038/ismej.2014.233
LeRoux, F., Wegner, K.M., Baker-Austin, C., Vezzulli, L., Osorio, C.R., Amaro, C., Ritchie, J.M., Defoirdt, T.,
Destoumieux-Garzon, D., Blokesch, M., Mazel, D., Jacq, A., Cava, F., Gram, L., Wendling, C.C.,
Strauch, E., Kirschner, A., Huehn, S., 2015. The emergence of Vibrio pathogens in Europe:
ecology, evolution, and pathogenesis (Paris, 11-12th March 2015). Front. Microbiol. 6, 830.
https://doi.org/10.3389/fmicb.2015.00830

Li, D., Zhang, H., Fu, L., An, X., Zhang, B., Li, Y., Chen, Z., Zheng, W., Yi, L., Zheng, T., 2014. A Novel
Algicide: Evidence of the Effect of a Fatty Acid Compound from the Marine Bacterium, Vibrio sp
BS02 on the Harmful Dinoflagellate, Alexandrium tamarense. Plos One 9, e91201.
https://doi.org/10.1371/journal.pone.0091201
Liu, P.-C., Lee, K.-K., Yii, K.-C., Kou, G.-H., Chen, S.-N., 1996. News & Notes: Isolation of Vibrio harveyi
from Diseased Kuruma Prawns Penaeus japonicus. Curr. Microbiol. 33, 129–132.
https://doi.org/10.1007/s002849900087
Lopez-Joven, C., Rolland, J.-L., Haffner, P., Caro, A., Roques, C., Carre, C., Travers, M.-A., Abadie, E.,
Laabir, M., Bonnet, D., Destoumieux-Garzon, D., 2018. Oyster Farming, Temperature, and
Plankton Influence the Dynamics of Pathogenic Vibrios in the Thau Lagoon. Front. Microbiol. 9,
2530. https://doi.org/10.3389/fmicb.2018.02530
Mandel, M.J., Stabb, E.V., Ruby, E.G., 2008. Comparative genomics-based investigation of resequencing
targets in Vibrio fischeri: Focus on point miscalls and artefactual expansions. Bmc Genomics 9,
138. https://doi.org/10.1186/1471-2164-9-138
Marampouti, C., Buma, A.G.J., De Boer, M.K., 2021. Mediterranean alien harmful algal blooms: origins
and impacts. Environ. Sci. Pollut. Res. 28, 3837–3851. https://doi.org/10.1007/s11356-020-
10383-1
Martin, J., Fitzwater, S., 1988. Iron-Deficiency Limits Phytoplankton Growth in the Northeast Pacific
Subarctic. Nature 331, 341–343. https://doi.org/10.1038/331341a0
Martin, M.O., 2002. Predatory prokaryotes: An emerging research opportunity. J. Mol. Microbiol.
Biotechnol. 4, 467–477.
Mavian, C., Paisie, T.K., Alam, M.T., Browne, C., Beau De Rochars, V.M., Nembrini, S., Cash, M.N., Nelson,
E.J., Azarian, T., Ali, A., Morris, J.G., Salemi, M., 2020. Toxigenic Vibrio cholerae evolution and
establishment of reservoirs in aquatic ecosystems. Proc. Natl. Acad. Sci. 117, 7897–7904.
https://doi.org/10.1073/pnas.1918763117
McRose, D.L., Baars, O., Seyedsayamdost, M.R., Morel, F.M.M., 2018. Quorum sensing and iron regulate
a two-for-one siderophore gene cluster in *Vibrio harveyi*. Proc. Natl. Acad. Sci. 115, 7581–7586.
https://doi.org/10.1073/pnas.1805791115
Miethke, M., Marahiel, M.A., 2007. Siderophore-Based Iron Acquisition and Pathogen Control. Microbiol.
Mol. Biol. Rev. 71, 413–451. https://doi.org/10.1128/MMBR.00012-07
Miller, S.D., Haddock, S.H.D., Elvidge, C.D., Lee, T.F., 2005. Detection of a bioluminescent milky sea from
space. Proc. Natl. Acad. Sci. U. S. A. 102, 14181–14184.
https://doi.org/10.1073/pnas.0507253102
Millero, F.J., 1998. Solubility of Fe(III) in seawater. Earth Planet. Sci. Lett. 154, 323–329.
https://doi.org/10.1016/S0012-821X(97)00179-9
Muhling, B.A., Jacobs, J., Stock, C.A., Gaitan, C.F., Saba, V.S., 2017. Projections of the future occurrence,
distribution, and seasonality of three Vibrio species in the Chesapeake Bay under a high-emission
climate change scenario. Geohealth 1, 278–296. https://doi.org/10.1002/2017GH000089
Neilands, J., 1981. Iron-Absorption and Transport in Microorganisms. Annu. Rev. Nutr. 1, 27–46.
https://doi.org/10.1146/annurev.nu.01.070181.000331
Oberbeckmann, S., Fuchs, B.M., Meiners, M., Wichels, A., Wiltshire, K.H., Gerdts, G., 2012. Seasonal
Dynamics and Modeling of a Vibrio Community in Coastal Waters of the North Sea. Microb. Ecol.
63, 543–551. https://doi.org/10.1007/s00248-011-9990-9
Pal, M., Yesankar, P.J., Dwivedi, A., Qureshi, A., 2020. Biotic control of harmful algal blooms (HABs): A
brief review. J. Environ. Manage. 268, 110687. https://doi.org/10.1016/j.jenvman.2020.110687

Papenfort, K., Bassler, B.L., 2016. Quorum sensing signal-response systems in Gram-negative bacteria.
Nat. Rev. Microbiol. 14, 576–588. https://doi.org/10.1038/nrmicro.2016.89

Perez, J., Moraleda-Munoz, A., Javier Marcos-Torres, F., Munoz-Dorado, J., 2016. Bacterial predation: 75
years and counting! Environ. Microbiol. 18, 766–779. https://doi.org/10.1111/1462-2920.13171

Pfaffl, M.W., 2001. A new mathematical model for relative quantification in real-time RT-PCR. Nucleic
Acids Res. 29, e45. https://doi.org/10.1093/nar/29.9.e45

R Core Team, 2024. R: A Language and Environment for Statistical Computing. https://www.R-
project.org/

Rashidan, K.K., Bird, D.F., 2001. Role of predatory bacteria in the termination of a cyanobacterial bloom.
Microb. Ecol. 41, 97–105.

Rodrigues, A.M.S., Lami, R., Escoubeyrou, K., Intertaglia, L., Mazurek, C., Doberva, M., Pérez-Ferrer, P.,
Stien, D., 2022. Straightforward *N* -Acyl Homoserine Lactone Discovery and Annotation by LC–
MS/MS-based Molecular Networking. J. Proteome Res. 21, 635–642.
https://doi.org/10.1021/acs.jproteome.1c00849

Rolland, J.-L., Pelletier, K., Masseret, E., Rieuvilleneuve, F., Savar, V., Santini, A., Amzil, Z., Laabir, M.,
2012. Paralytic Toxins Accumulation and Tissue Expression of alpha-Amylase and Lipase Genes in
the Pacific Oyster Crassostrea gigas Fed with the Neurotoxic Dinoflagellate Alexandrium
catenella. Mar. Drugs 10, 2519–2534. https://doi.org/10.3390/md10112519

Romano, A., Trimble, L., Hobusch, A.R., Schroeder, K.J., Amin, S.A., Hartnett, A.D., Barker, R.A., Crumbliss,
A.L., Carrano, C.J., 2013. Regulation of iron transport related genes by boron in the marine
bacterium Marinobacter algicola DG893. Metallomics 5, 1025.
https://doi.org/10.1039/c3mt00068k

Rosales, D., Ellett, A., Jacobs, J., Ozbay, G., Parveen, S., Pitula, J., 2022. Investigating the Relationship
between Nitrate, Total Dissolved Nitrogen, and Phosphate with Abundance of Pathogenic Vibrios
and Harmful Algal Blooms in Rehoboth Bay, Delaware. Appl. Environ. Microbiol. 88, e00356-22.
https://doi.org/10.1128/aem.00356-22

Sinclair, A.R.E., Mduma, S., Brashares, J.S., 2003. Patterns of predation in a diverse predator-prey system.
Nature 425, 288–290. https://doi.org/10.1038/nature01934

Su, R.Q., Yang, X.R., Zheng, T.L., Tian, Y., Jiao, N.Z., Cai, L.Z., Hong, H.S., 2007. Isolation and
characterization of a marine algicidal bacterium against the toxic dinoflagellate Alexandrium
tamarense. Harmful Algae 6, 799–810. https://doi.org/10.1016/j.hal.2007.04.004

Tanabe, T., Funahashi, T., Nakao, H., Miyoshi, S.-I., Shinoda, S., Yamamoto, S., 2003. Identification and
Characterization of Genes Required for Biosynthesis and Transport of the Siderophore
Vibrioferrin in *Vibrio parahaemolyticus*. J. Bacteriol. 185, 6938–6949.
https://doi.org/10.1128/JB.185.23.6938-6949.2003

Thompson, F.L., Thompson, C.C., Swings, J., 2003. Vibrio tasmaniensis sp. nov., isolated from Atlantic
Salmon (Salmo salar L.). Syst. Appl. Microbiol. 26, 65–69.
https://doi.org/10.1078/072320203322337326

Tourneroche, A., Lami, R., Hubas, C., Blanchet, E., Vallet, M., Escoubeyrou, K., Paris, A., Prado, S., 2019.
Bacterial–Fungal Interactions in the Kelp Endomicrobiota Drive Autoinducer-2 Quorum Sensing.
Front. Microbiol. 10, 1693. https://doi.org/10.3389/fmicb.2019.01693

Vanhove, A.S., Rubio, T.P., Nguyen, A.N., Lemire, A., Roche, D., Nicod, J., Vergnes, A., Poirier, A.C.,
Disconzi, E., Bachere, E., Le Roux, F., Jacq, A., Charriere, G.M., Destoumieux-Garzon, D., 2016.
Copper homeostasis at the host vibrio interface: lessons from intracellular vibrio transcriptomics.
Environ. Microbiol. 18, 875–888. https://doi.org/10.1111/1462-2920.13083

Wang, X., Li, Z., Su, J., Tian, Y., Ning, X., Hong, H., Zheng, T., 2010. Lysis of a red-tide causing alga,
769   Alexandrium tamarense, caused by bacteria from its phycosphere. Biol. Control 52, 123–130.
770   https://doi.org/10.1016/j.biocontrol.2009.10.004

771 Wang, Y., Li, S., Liu, G., Li, X., Yang, Q., Xu, Y., Hu, Z., Chen, C.-Y., Chang, J.-S., 2020. Continuous
772   production of algicidal compounds against Akashiwo sanguinea via a Vibrio sp. co-culture.
773   Bioresour. Technol. 295, 122246. https://doi.org/10.1016/j.biortech.2019.122246

774 Weerasinghe, A.J., Amin, S.A., Barker, R.A., Othman, T., Romano, A.N., Parker Siburt, C.J., Tisnado, J.,
775   Lambert, L.A., Huxford, T., Carrano, C.J., Crumbliss, A.L., 2013. Borate as a Synergistic Anion for
776   *Marinobacter algicola* Ferric Binding Protein, FbpA: A Role for Boron in Iron Transport in Marine
777   Life. J. Am. Chem. Soc. 135, 14504–14507. https://doi.org/10.1021/ja406609s

778 Wu, J., Luther, G., 1994. Size-Fractionated Iron Concentrations in the Water Column of the Western
779   North-Atlantic Ocean. Limnol. Oceanogr. 39, 1119–1129.
780   https://doi.org/10.4319/lo.1994.39.5.1119

781 Xu, Q., Wang, P., Huangleng, J., Su, H., Chen, P., Chen, X., Zhao, H., Kang, Z., Tang, J., Jiang, G., Li, Z., Zou,
782   S., Dong, K., Huang, Y., Li, N., 2022. Co-occurrence of chromophytic phytoplankton and the Vibrio
783   community during Phaeocystis globosa blooms in the Beibu Gulf. Sci. Total Environ. 805, 150303.
784   https://doi.org/10.1016/j.scitotenv.2021.150303

785 Yang, Q., Feng, Q., Zhang, B., Gao, J., Sheng, Z., Xue, Q., Zhang, X., 2021. Marinobacter alexandrii sp. nov.,
786   a novel yellow-pigmented and algae growth-promoting bacterium isolated from marine
787   phycosphere microbiota. Antonie Van Leeuwenhoek Int. J. Gen. Mol. Microbiol. 114, 709–718.
788   https://doi.org/10.1007/s10482-021-01551-5

789 Zhang, W., Wang, Y., Lu, H., Liu, Q., Wang, C., Hu, W., Zhao, K., 2020. Dynamics of Solitary Predation by
790   Myxococcus xanthus on Escherichia coli Observed at the Single-Cell Level. Appl. Environ.
791   Microbiol. 86, e02286-19. https://doi.org/10.1128/AEM.02286-19

ACKNOWLEDGMENT

The authors thank F. Le Roux and Y. Labreuche for their gift of *Vibrio* mutants; E. Garcés, S. Turki, H.

Frehi, A. Bouquet and W. Medhioub for their contribution in obtaining Mediterranean phytoplankton

strains; K. Escoubeyrou and D. Stien for advice on HRMS/MS data; V. Diakou-Verdin and E. Jublanc

from the Montpellier RIO Imaging Platform for access to microscope facilities (www.mri.cnrs.fr); P. Clair

from the Montpellier qPHD platform and the Technoviv platform of the UPVD for access to PCR

facilities; Luc Markiw, N. Brunet, A. Lang, A. Payelleville, T. Milhau and M. Leroy for their technical

assistance; The IFREMER LER-LR laboratory for access to the Thau Lagoon; the BIO2MAR platform at

the Observatoire Océanologique de Banyuls for providing access to UHPLC-HRMS/MS facilities.

FUNDING

European VIVALDI grant 678589, French EC2CO ROSEOCOM grant and IFREMER ALGOVIR grant.

     FIGURES, TABLES and VIDEOS

Figure 1. Dynamics of *Alexandrium* and Vibrio in the environment. (A) Location of the monitoring
station in the Thau Lagoon (southern France). **(B)** Mean abundance (DNA equiv.) of *Vibrio spp*. (16S) and
Alexandrium *spp. (A. pacificum* ACT03 + *A. tamarense* ATT07). Vibrio cells (black line with diamond
dot) and degraded *Alexandrium* cells (grey line with round dot) were detected in the 0.2–0.8 μm fraction
(free Vibrio fraction) in spring and autumn 2015. Living *Alexandrium* cells (grey line with round dot) but
no plankton-associated *Vibrio spp*. (black line with diamond dot) were evidence in the 0.8–180 μm in
spring and autumn. **(C)** Results of the Akaike Information Criterion (AICc) test conducted to select a
model for explaining the mean value of dead Alexandrium (degraded cells) in spring. **(D)** Wald test of the
AICc model explaining the mean value of dead Alexandrium in spring by free Vibrio.

Figure 2. Incubation of *V. atlanticus* LGP32 and *Alexandrium pacificum* ACT03 in enriched natural seawater (ENSW). **(A)** *A. pacificum* ACT03 cultured alone (grey bar) and incubated with *V. atlanticus* LGP32 (black bar) in ENSW. **(B)** *V. atlanticus* LGP32 cultured alone (grey bar) and incubated with *A. pacificum* ACT03 (black bar) in ENSW. **(C)** Snapshot of the interaction between *V. atlanticus* LGP32-GFP cells (60-hour culture) and one cell of *A. pacificum* ACT03 taken at 8h00 of co-culture. **(D)** Chronological snapshots of the interaction (70 pictures, one per second). *V. atlanticus* LGP32 (small green cells) and *A. pacificum* ACT03 cell (large red cell). All experiments were done in triplicate. Asterisks indicate significant differences in a multiple comparison test (One-way ANOVA with post hoc Tukey test), *P ≤ 0.05.

827
Figure 3. Role of *V. atlanticus* LGP32 starvation in the interspecific interaction process. Experiments
were conducted by incubating *A. pacificum* ACT03 with *V. atlanticus* LGP32 previously grown for 12, 36,
60 and 126 h in Zobell medium. (**A**) Cumulative percentage of motile *A. pacificum* ACT03 cells. (**B**)
Cumulative number of cells attacked by *V. atlanticus* LGP32 and (**C**) Cumulative cell lysis after 0, 15, 30,
45 and 60 minutes of interaction. Corresponding pictures showing (**A1**) Black arrows indicate unhooked
and degraded flagellum from *A. pacificum* ACT03 flagellum, (**B1**) Chronological sequence of five
snapshots showing *V. atlanticus* LGP32-GFP cells (60-hour culture) and *A. pacificum* ACT03 cells,
during the first hour of their interaction. *V. atlanticus* LGP32 (small green cells), living *A. pacificum*
ACT03 (large red cells) and dead *A. pacificum* ACT03 (large green cell). (**C1**) Black arrow 1 indicates
vesicle formation on *A. pacificum* ACT03 cell and black arrow 2 indicates lysed *A. pacificum* ACT03 cell.
All percentages were determined based on a minimum of 2,000 cells of *A. pacificum* ACT03. All
experiments were done in triplicate. Asterisks indicate significant differences in a multiple comparison
test (One-way ANOVA with post hoc Tukey test), ***P ≤ 0.001

Figure 4. Role of quorum sensing and the vibrioferrin iron uptake pathway in the interaction process. **(A)** Effect of *V. atlanticus* LGP32 cell density on the attack process. *A. pacificum* ACT03 cells (1 x 10^3 cells mL^{-1}) were incubated with *V. atlanticus* LGP32 grown for 60 hours in Zobell medium at concentrations ranging from 5.10^3 to 5×10^5 cells mL^{-1} (black bars). For comparison, *A. pacificum* ACT03 incubated with *V. atlanticus* LGP32 grown for 12 hours in Zobell medium at concentrations ranging from 5×10^3 to 4×10^6 cells mL–1 (grey bars). The image on the bars indicates either unaffected algae (live red algae) or algae attacked by Vibrio (algae covered with green vibrio cell) during the interaction **(B)** CqsS, luxM, luxN, luxS, and luxP quorum sensing and PvsA, PvuB and PvuA2 vibrioferrin pathway genes expression in *V. atlanticus* LGP32 grown for 12, 36 and 60 h in Zobell medium. **(C)** Effect of *V. atlanticus* LGP32 mutants on the attack process. Experiments were conducted by incubating *A. pacificum* ACT03 with *V. atlanticus* LGP32, *V. atlanticus* LGP32 tagged with GFP, *V. atlanticus* LGP32 washed with ENSW or *V. atlanticus* LGP32 mutant ΔPvuB, ΔluxM, ΔluxR and ΔluxS previously grown 60 h in Zobell media (control), The percentage of *A. pacificum* ACT03 attacked was determined during the first 30 min of exposure. **(D)** Effect of *V. atlanticus* LGP32 culture media composition on the attack process. Experiments were conducted by incubating *A. pacificum* ACT03 with *V. atlanticus* LGP32 grown 60 h in Zobell media supplemented with boron (H_3BO_3) or $FeCl_3$. The results were compared with an exposure to *V. atlanticus* LGP32 grown 60 h in Zobell media. All percentages were determined based on a minimum of 2,000 cells of *A. pacificum* ACT03. All experiments were done in triplicate. Asterisks indicate significant differences in a multiple comparison test (One-way ANOVA with post hoc Tukey test), **P $\leq$ 0.01, ***P $\leq$ 0.001.

864
Figure 5. Schematic representation of a putative strategy developed by *Vibrio spp*. to feed on
Alexandrium *spp*. and *G. catenatum* in the environment. (A) Vibrio in the environment when subjected
to starvation secrete non-protein lytic compounds. **(B)** Some of these lytic compounds degrade the
flagella, immobilizing the alga (immobilization stage). **(C)** Then Vibrio swims and clusters around its prey
(attack stage). **(D)** Lytic compounds released by Vibrio where able to concentrate around the algal cells,
thereby lysing the algae (killing stage). **(E)** Feeding on the released nutrients, Vibrio multiply and then
spread in the environment. Yellow clouds: Lytic compound release by Vibrio, Grey clouds: Algal
nutrients released upon lysis.

874
Figure S1. Time and dose-dependent effects of the *V. atlanticus* LGP32 culture supernatant on *A. pacificum* ACT03 motility. (**A**) A time dependence experiment was conducted by incubating *A. pacificum* ACT03 for 1, 5 or 20 h with 1/1000 v/v (1 µL/mL) of culture supernatant from *V. atlanticus* LGP32 previously grown for 60 h in Zobell culture media. (**B**) A dose-dependence experiment was conducted by incubating *A. pacificum* ACT03 for 1 h with 1/1000 to 1/20 v/v (1-50 µL/mL) of culture supernatant from *V. atlanticus* previously grown for 60 h in Zobell media. The percentage of motile *A. pacificum* ACT03 was determined after 1 hour of exposure. All percentages were determined based on a minimum of 2,000 cells of *A. pacificum* ACT03. Error bars represent the standard deviation of the mean of three independent experiments. Asterisks indicate significant differences in a multiple comparison test (One-way ANOVA with post hoc Tukey test), *P ≤ 0.05, ***P ≤ 0.001.

Figure S2. *V. atlanticus* **LGP32 proteome analysis following nutrient stress. (A)** Example of 2D gel, the numbers in white on the gel 4-7 correspond to the number and position of the protein spots analyzed. **(B)** Proteins identified by LC-MS/MS as differentially represented in the 2D gel comparative approach following nutrient stress. ND: Not determined; ENSW (artificial seawater)**.**

Figure S3. Quorum sensing and the vibrioferrin iron uptake pathway in Vibrio. (A) Putative quorum sensing (QS) pathways at low and high cell density in Vibrio according to Lami et al.(Lami, 2019). **(B)** Genetic organization of the vibrioferrin utilization gene cluster on *V. atlanticus* LGP32 chromosome 2. The Pvu and Pvs operons are involved in the secretion and the transport of ferric vibrioferrin and biosynthesis of vibrioferrin, respectively. Arrows indicate the transcriptional directions of the genes. VSII1126, VSII1137 and VSII1129 corresponding to PvuB, PvuA2 and PvsA genes respectively.

899 **Table 1.** Ability of Vibrio strains to attack and to lyse *Alexandrium pacificum.* ND: not determined.

Vibrio species	Strains	Virulence for fish or invertebrates	References	Attack *A. pacificum*	Lyse *A. pacificum*
Vibrio atlanticus LGP32	WT	Yes	(Gay et al., 2004)	+	+
//	WT + pSW3654T-GFP	Yes	(Le Roux et al., 2007)	+	+
//	ΔLuxM	ND	Ifremer Institute, France	+	+
//	ΔLuxS	ND	//	+	+
//	ΔLuxR	ND	//	+	+
//	ΔPvuB	ND	This work	-	+
Vibrio tasmaniensis	J5-9	Yes	(Lemire et al., 2015)	+	+
//	LMG20012^T	No	(Thompson et al., 2003)	+	+
Vibrio crassostreae	J2.9	Yes	(Lemire et al., 2015)	+	+
//	J2-8	No	//	+	+
Vibrio fischeri	ES114	ND	(Mandel et al., 2008)	+	+
Vibrio harveyi	ATCC14126	Yes	(Liu et al., 1996)	+	+
Vibrio aestuarianus	janv-32	Yes	(Labreuche et al., 2010)	+	+

Table 2. Ability of *V. atlanticus* LGP32 to degrade flagella, attack and lyse the targeted dinoflagellates *spp*. commonly found in the Mediterranean Sea. ND not determined.

Dinoflagellates species	Strains	Toxicity for human	References	Flagella degraded	Cells Attacked	Cells Lysed
Alexandrium pacificum	ACT03, Thau, France	Yes	(Laabir et al., 2011)	+	+	+
Alexandrium catenella	Bizerte, Tunisia	Yes	(Fertouna-Bellakhal et al., 2015)	+	+	+
//	F3-9F, Tarragona, Spain	ND	//	+	+	+
//	C10-5, Annaba, Algeria	Yes	(Hadjadji et al., 2020)	+	+	+
Alexandrium tamarense	ATT07, Thau, France	No	(Rolland et al., 2012)	+	+	+
Alexandrium spp.	Golf of Tunis, Tunisia	ND	Algal collection University of Montpellier, France	+	+	+
//	Bizerte, Tunisia	ND	//	+	+	+
//	Mediterranean coast, Morocco	ND	//	+	+	+
Prorocentrum lima	PLBZT14, Bizerte, Tunisia	Yes	(Ben-Gharbia et al., 2016)	-	-	-
Coolia monotis	CMBZT14, Bizerte, Tunisia	ND	//	-	-	-
Vulcanodinium rugosum	IFR-VRU-01, Ingril, France	Yes	(Abadie et al., 2015)	-	-	-
Karenia selliformis	Golf of Gabes, Tunisia	ND	Algal collection University of Montpellier, France	-	-	-
Scripsiella trochoidae	Mellah Lagoon, Algeria	-	//	-	-	-
Gyrodinium impudicum	Golf of Tunis, Tunisia	-	//	-	-	-
Amphidium carterae	SAMS, Scotland	ND	SAMS laboratory, Scotland	-	-	-
Gymnodinium catenatum	M'diq Bay, Morocco	+	(Leblad et al., 2020)	+	+	+

904 **Table 3.** Oligonucleotide sequences of primers used for RNA expression analysis.

Species	genes	Primer Sequences	Tm (°C)	Efficiency	References
Alexandrium pacificum (ACT03)	18S – 28S rRNA ITS region	TGATATTGTGGGCAACTGTAA AACATCTGTTAGCTCACGGAA	54		(Genovesi et al., 2011)
Alexandrium tamarense (ATT07)	18S – 28S rRNA ITS region	TGGTAATTCTTCATTGATTACAATG AACATCTGTTAGCTCACGGAA	54		//
Vibrios spp.	16S	CGGTGAAATGCGTAGAGAT TTACTAGCGATTCCGAGTTC	62		(Kitatsukamoto et al., 1993)
Vibrio atlanticus LGP32	LuxN (VS_II0260)	CACTTGCTAGTATCATCGC ATCGAGTTAGCAAGAGCAC	60	1,92	This work
//	LuxM (VS_II0261)	TCCACTTATCACAAACAGG ACTGTACTTCCATTTGTCG	60	1,91	//
//	LuxP (VS_II0355)	AAGTTCAGGATGAACCTATC CAAAGAGATACTTTGCTGAG	60	1,89	//
//	LuxS (VS_2562)	ACTCTCGAGCACCTATACG GAAGGCGTACCAATCAAGC	60	1,85	//
//	CqsS (VS_1725)	GACATCTATTGATGTTATGC TCACCCACTTCACGTAACTG	60	1,91	//
//	PvsA (VS_II0355),Vibrioferrin biosynthesis protein	CAGAGCAAGAGCTAGAACC TCGTTGAGAACCTGACGAG	59	1,91	//
//	PvuB (VS_II1126), ABC transporter vibrioferrin uptake FecB	TAGTGCAACCATGGGAATCG TAAACCGTACGTAGACGCTC	57	2,01	//
//	PvuA2 (VS_II1127), Vibrioferrin receptor FecA	GGAGCTACAAGCATTCGTTC TTCGTCATATGGTCGCTTCG	57	2,08	//
//	Housekeeping gene 1, CcmC (VS_0852)	ATTGCCGCCTTTATCGGTTT CAAGCACCCCACATTGGTTT	60		(Vanhove et al., 2016)
//	Housekeeping gene 2, 6-phosphofructokinase (VS_2913)	GCCGTCACTGTGGTGACCTT TGCTTCTTGCCTTTCGCAAT	60		//

VIDEOS

Video 1. Dynamics of *V. atlanticus* LGP32-*Alexandrium pacificum* ACT03 interaction. GFP-tagged *V. atlanticus* (small green cells); living *A. pacificum* (large red cells); lysed *A. pacificum* (large green cells) filmed under an epifluorescence microscope.

Video 2. Second-by-second timing of *V. atlanticus* LGP32 attacking *Alexandrium pacificum* ACT03. GFP-tagged *V. atlanticus* (small green cells); *A. pacificum* living cell (large red cells) filmed under an epifluorescence microscope**.**

Video 3. Degradation and disruption of *Alexandrium pacificum* ACT03 flagella. Effect of Vibrio supernatant on the first stage of the interaction filmed under a confocal microscope.

Video 4. Attacks of *V. atlanticus* LGP32 on target *Alexandrium pacificum* ACT03. This video, recorded under a confocal microscope, shows Vibrios simultaneously attacking a first immobilized Alexandrium cell, then moving on to attack a second cell without ever targeting the other cells present, suggesting active communication between the Vibrio bacteria. *V. atlanticus* LGP32 (small cells); *A. pacificum* ACT03 (large cells).

Video 5. Vesicle formation and bursting of an *Alexandrium pacificum* ACT03. Direct effect of Vibrio supernatant on *Alexandrium* after 126 h of culture filmed under a confocal microscope.

Model degraded Alexandrium cells	AICc
Intercept+ free Vibrio	171.77
intercept	172.57
Intercept + salinity	173.41
Intercept + temperature	173.94
Intercept+ free Vibrio + salinity	174.99
Intercept+ free Vibrio + temperature	175.37
Intercept + temperature + salinity	177.72
Intercept+ free Vibrio + temperature + salinity	180.56

	Estimate	Std error	T value	Pr(>|t|)
Intercept	235.41	38.69	6.08	7.91×10^{-05}
Vibrio	83.19	40.27	2.07	0.06

A

Alexandrium mL^{-1}

B

Vibrio mL^{-1}

Incubation time (Hours)

C

Alexandrium   Merge   Vibrio GFP

D

Chronological snapshots of the event
(1 snapshot per second)

Vibrio starved

A

Lytic compound release

Vibrio migration

E

Vibrio grow then spread in the environment

Lytic compound accumulation

nutrient release

Fe

Flagellum degradation

Alexandrium

B

Immobilization stage

C

Attack stage

D

Killing stage

A

B

Spot #	Value ENSW	Value Zobell	Ratio (ENSW/Zobell)	Best ID Accession	#Peptides	-10logP Prot	-10logP Pept. Min	-10logP Pept. Max	Seq coverage	CDD accession	Best ID Description	Expected mass	Expected pI
7304	3620,4	570,7	6,34	WP_012601215.1	3	172,83	67,81	92,01	14%	PRK00443	glucosamine-6-phosphate deaminase	29610	5,53
4115	1203,8	191,9	6,27	ND									
0403	3697,4	750,2	4,93	WP_004733797.1	16	253,95	23,25	124,97	33%	cd13641	ABC transporter substrate-binding protein	36195	4,52
4413	**640,2**	**267,4**	**2,39**	**WP_012600975.1**	**21**	**241,28**	**25,81**	**101,45**	**51%**	**PRK11411**	**iron siderophore-binding protein, PvuB**	**36111**	**5,73**
7009	616,6	316,3	1,95	ND									
3704	683,2	1303	-1,91	WP_009848030.1	30	360,8	29,21	121,35	54%	PRK09344	phosphoenolpyruvate carboxykinase (ATP)	59755	4,85
4114	315,2	812,6	-2,58	WP_009847387.1	2	58,71	58,71	58,71	8%	PRK00129	uracil phosphoribosyltransferase	22658	5,21
3412	57,7	238,2	-4,13	WP_012604509.1	10	244,21	29,11	105,11	38%	cd13674	C4-dicarboxylate ABC transporter	36946	5,26
5612	211,4	1395,9	-6,60	WP_012600249.1	2	79,11	40,49	40,83	6%	PRK05451	dihydroorotase	37599	5,36
9601	76,9	1717,1	-22,33	WP_004736554.1	12	208,83	33,74	100,38	29%	PRK08722	beta-ketoacyl-(acyl-carrier-protein)synthase II	43007	5,95

A

Low cell density

High cell density

B

pvuE VSII_1123
pvuD VSII_1124
pvuC VSII_1125
pvuB VSII_1126
pvuA2 VSII_1127
pvuA1 VSII_1128
pvsA VSII_1129
pvsB VSII_1130
pvsC VSII_1131
pvsD VSII_1132
pvsE VSII_1133

Vibrioferrin transport operon

Vibrioferrin biosynthesis and efflux operon