Starvation of the bacteria *Vibrio atlanticus* promotes lightning group-attacks on the

dinoflagellate *Alexandrium pacificum*

Jean Luc Rolland[1, 2†*], Estelle Masseret[3], Mohamed Laabir[3], Guillaume Tetreau[4], Benjamin Gourbal[4],

Anne Thebault[5], Eric Abadie[6], Alice Rodrigues-Stien[7,8], Carole Veckerlé[7,8], Elodie Servanne-Meunier[1],

Delphine Destoumieux-Garzón[1], Arnaud Lagorce[4,9], Raphaël Lami[7,9].

Affiliations

[1]IHPE, Univ. Montpellier, CNRS, IFREMER, Université de Perpignan Via Domitia, Montpellier, France.

[2]MARBEC, Univ. Montpellier, IRD, IFREMER, CNRS, Sète, France.

[3]MARBEC, Univ. Montpellier, IRD, IFREMER, CNRS, Montpellier, France.

[4]IHPE, Univ Montpellier, CNRS, IFREMER, Université de Perpignan Via Domitia, Perpignan, France

[5]Agence nationale de sécurité sanitaire de l'alimentation, de l'environnement et du travail (ANSES),

ANSES-DER 27-31, avenue du Général Leclerc, 94701 Maison-Alfort, France

[6]IFREMER, 9 rue Jean Bertho, 97822, Le port, la Reunion, France

[7]Sorbonne Université, CNRS, LBBM USR3579, Observatoire Océanologique de Banyuls sur Mer,

Avenue Pierre Fabre, 66650 Banyuls-sur-Mer, France.

[8]Sorbonne Université, CNRS, Bio2MAR, Observatoire Océanologique de Banyuls sur Mer, Avenue Pierre

Fabre, 66650 Banyuls-sur-Mer, France.

[9]These authors contributed equally

[†]Current address

*Corresponding author: Jean Luc.Rolland

Email: Jean.Luc.Rolland@ifremer.fr

ABSTRACT

Algae serve as a source of nutrients for bacteria in the marine environment. The interactions between algae and bacteria at the phycosphere interface are known to include mutualism, commensalism, competition or antagonism. Here, based on *in situ* observation and on an *in vitro* interaction study, we report on a novel form of starvation-induced hunting that the cells of selected *Vibrio* species exert on the dinoflagellate species. Results showed that *Vibrio atlanticus* was able to coordinate lightning group attacks then kill the dinoflagellate *Alexandrium pacificum* ACT03. Briefly, the observed coordinated mechanism of algal-killing consists of first, the 'immobilization stage' involving the secretion of algicidal metabolites that disrupt the flagella of the prey. The 'attack stage' resembles the 'wolf-pack attack' strategy, during which *Vibrios* surrounds algal cells at high density for a brief period without invading them. Finally, the 'killing stage' induces the lysis and degradation of the dinoflagellates. By using a combination of biochemical, proteomic, genetic and fluorescence microscopy approaches, we showed that this relationship is not related to the decomposition of algal organic matter, *Vibrio* quorum sensing pathways, to the toxicity of the algae or to the pathogenicity of the bacterium but is conditioned by nutrient stress, iron availability and link to the iron-vibrioferrin transport system of *V. atlanticus*. This is the first evidence of a new mechanism that could be involved in regulating Alexandrium spp. blooms and giving Vibrio a competitive advantage in obtaining nutrients from the environment.

KEYWORDS

Interaction; Bacterium; Alga; Predation; Environment; Iron related predation

INTRODUCTION

Harmful algal blooms (HABs) have experienced an increase in their occurrence, intensity, and geographical distribution on a global scale, resulting in adverse environmental, health, and socioeconomic impacts (Marampouti et al., 2021). HABs have a considerable impact on human health as a result of direct exposure to volatile toxins or by toxic seafood consumption (Burkholder et al., 2018). From an ecological point of view, the expansion of HABs can result in the erosion of biodiversity, because they cause massive mortality of marine species and they are generally monospecific in nature (Chai et al., 2020). In coastal areas, understanding the biological interactions that control toxic algal blooms is therefore a major ecological challenge. Among HABs, a number of *Alexandrium* species have been placed on the list of invasive Mediterranean species. Among them, *Alexandrium pacificum* is a flagellated eukaryotic unicellular organism that form with *Alexandrium tamarense* and *Alexandrium fundyense* the "Alexandrium tamarense" complex (Hadjadji et al., 2020). Since 1998, *A. pacificum* (former *A. catenella*) was monitored by Rephy survey in the Thau lagoon (French Mediterranean) because it produces paralytic shellfish toxins (PSTs) resulting of paralytic shellfish poisoning (PSP) syndrome.

Laania (Laanaia et al., 2013) showed that in Thau lagoon, a water temperature around 20°C for several days and organic and inorganic nutrients in sufficient concentrations are parameters favoring the developemnt of *A. pacificum*, whose massive blooms occur in autumn.

Algal blooms are seasonal events resulting in a rapid increase in the concentration of a species of algae in an aquatic environment. Depending on the phytoplankton species, the tolerance range of physicochemical parameters is different and influences the time of appearance of blooms (Leblad et al., 2020).

Interestingly, although the collapse of phytoplankton blooms has been previously attributed to viruses (Pal et al., 2020), some ecological studies have suggested an important role of algicidal bacteria (Su et al., 2007; Wang et al., 2010), such as some Vibrio *spp*. (Li et al., 2014; Wang et al., 2020).

Vibrio (class γ-proteobacteria) are common microorganisms in marine systems worldwide (Baker-Austin et al., 2017; Mavian et al., 2020), where they are important components of the food chain, particularly in

biodegradation, nutrient regeneration and biogeochemical cycles (Oberbeckmann et al., 2012). Vibrio is

one of the most studied bacterial taxa due to their ubiquity in coastal marine systems and their capacity to

cause infections in humans and animals, leading sometimes to epizootic or zoonotic epidemics (LeRoux et

al., 2015; Mavian et al., 2020). *Vibrio* are extremely adaptable to their environment (Johnson, 2013). The

main factors influencing their occurrence and distribution in water are temperature, salinity, nutrient

availability (Wang et al., 2020), multiple strategies such as biofilm formation on biotic and abiotic

surfaces (Espinoza-Vergara et al., 2020), or interactions with a multitude of other organisms such as

eukaryotic predators (Drebes Dörr and Blokesch, 2020) or plankton (Lopez-Joven et al., 2018) are used by

Vibrio in the environment. There is also evidence that global climate change has increased *Vibrio*-

associated illnesses affecting humans and animals (Brumfield et al., 2021; Muhling et al., 2017).

However, the drivers and dynamics of *Vibrio* survival and propagation in the marine environment are not

yet fully understood.

A substantial number of research articles have highlighted the potential of γ-proteobacteria to exert

algicidal activity against dinoflagellates, supporting the hypothesis that γ-proteobacteria such as *Vibrio*

play a role in the control of algae blooms *in situ* (Coyne et al., 2022). However, the mechanisms behind

Vibrio-driven algal lysis in the environment remain to be elucidated. Particularly, it is unclear how in the

water column, algicidal compounds secreted by bacteria can concentrate around the algae to exert their

lytic effect.

Based on observations from the natural environment showing a potent relationship between *Vibrio* and

Alexandrium algae bloom events, this study aim to determine *in vitro,* the main factors implicated in this

relationship. Using a combination of biochemical, proteomic, genetic, and fluorescence

microscopy approaches, we explored the role of algal toxicity, bacterial pathogenicity and the

quorum sensing pathway on this relationship and showed the important role of nutrient stress and

the iron uptake pathway in this unique Vibrio/Alexandrium predator/prey interaction.

METHODS

Quantification of *Alexandrium* algae and *Vibrio* bacteria in the environment.

Seawater samples were collected in the Thau Lagoon (southern France, a shallow Mediterranean

ecosystem open to the sea (Abadie et al., n.d.), during spring and autumn 2015. Briefly, samples were

collected from the subsurface (-50 cm) near an oyster table at a phytoplankton surveillance site (part of the

REPHY network, N 43°26.058' and E 003°39.878'). Once a week during spring and autumn 2015, during

field sampling campaigns, 20 L of water was filtered on board through a 180 µm pore-size nylon

membrane. At the laboratory, according to Lopez-Joven et al. (Lopez-Joven et al., 2018) seawater was

fractionated into two size classes as follows: 2 L of the above filtrate was filtered through a 0.8 µm pore-

size polycarbonate Whatman Nuclepore membrane to obtain organisms in the 0.8–180 µm range

corresponding to plankton-associated *Vibrio* and living *Alexandrium* forms. Then, the filtrate from the 0.8

µm membrane was filtered again through a 0.2 µm pore-size polycarbonate Whatman Nuclepore

membrane until the membrane was saturated. *Alexandrium* cells, ranging from 25 to 40 µm, belong to the

microphytoplankton and are therefore retained in the 0.8–180 µm fraction. Any *Vibrio* cells potentially

associated with or attached to *Alexandrium* cells will also be retained in this fraction. *Vibrio* cells are

approximately 0.5-0.8 µm-thick. The fraction between 0.2 and 0.8 µm therefore includes the free-living

Vibrio. The bacterial population collected on 0.8-µm-pore-size filters was designated the particle-

associated community, and the population on 0.2-µm-pore-size filters was designated as the free-living

community. Membranes (in triplicate) were then conserved in 500 µL of 100% EtOH at -20°C.

Environmental DNA (eDNA) was extracted from the MF Millipore membrane using the Macherey-Nagel

NucleoSpin Tissue Kit and resuspended in 100 µL of water. The samples were then stored at -20°C after

eDNA quantity and purity were assessed using a NanoDrop system (NanoDrop Technologies,

Wilmington, DE, USA). PCR amplification reactions were done on a Roche LightCycler 480 Real-Time

thermocycler (qPHD platform, University of Montpellier, France).using specific primer pairs (Table 3).

Typically, the reactions contained 1 μL of template DNA (the DNA concentration for all samples varied

from 1 to 40 μg mL^{-1}), 0.5 μL of each primer (3.33 μM) and 4 μL of reaction mixture (SYBR Green

Master Mix) in a total volume of 6 μL. The reaction parameters were as follows: 5 minutes at 95°C (initial

denaturation) and 40 cycles of 10 s at 95°C (denaturation), 10 s at the corresponding hybridization

temperature (Table 3) and 10 s at 72°C (elongation). Melting curve profiles were generated by increasing

the temperature from 65°C to 95°C at 0.5°C s^{-1}. Amplification products were analysed using LightCycler

software (Roche Diagnostics). *Vibrio spp.* and *A. pacificum* and *A. tamarense* were quantified by

constructing calibration curves based on DNA from the *V. atlanticus* LGP32 reference strain (former *V.*

tasmaniensis LGP32) and from the *A. pacificum* reference strain (ACT03: *A. catenella* strain isolated form

Thau in 2003) and the *A. tamarense* reference strain (ATT07: *A. tamarense* isolated from Thau in 2007)

(not shown).

Strains and growth conditions.

Vibrio strains. Wild-type and isogenic mutants of *Vibrio atlanticus* LGP32 (Table 1) were used in this

study. Deletion-mutants included Δ*luxR*, Δ*luxP*, Δ*luxM* and Δ*pvuB* isogenic strains. The Δ*pvuB* mutant

was constructed here by allelic exchange as described previously by Le Roux (Le Roux et al., 2007). We

also used *V. atlanticus* LGP32 carrying the pSW3654T-GFP plasmid (Le Roux et al., 2007), hereafter

referred to as *V. atlanticus* LGP32-GFP. Bacterial strains were grown at 22 ± 1°C in Zobell medium (0, 38

μM iron (III)). When needed, 25 μg mL^{-1} chloramphenicol (Cm) was added to cultures of *V. atlanticus*

LGP32-GFP (Le Roux et al., 2007). Oligonucleotides used for RNA expression analysis are specified in

table 3.

Phytoplancton strains. Non-axenic phytoplankton species (Table 2) were grown in batch culture in

enriched natural sea water (ENSW, 6,55 μM iron (III)) with a salinity of 36 practical salinity units (PSU)

at 22 ± 1°C under cool white fluorescent illumination (100 μmol photons m^{-2} s^{-1}) and a 12 h:12 h

light:dark cycle (Harrison et al., 1980). The algae were used for experiments in their exponential growth

phase.

Co-culture assay

For each tested phytoplankton species, 2.10^4 cells harvested in their exponential growth phase were placed

in 20 mL of ENSW medium in a 50 mL suspension culture flask (Cellstar® PS, Greiner bio-one). After

incubation for 24 h at $22 \pm 1°C$ under cool white fluorescent illumination, 40 µL of *Vibrio* strains grown

for 12, 36, 60 and 156 h in Zobell medium or in Zobell medium supplemented with $FeCl_3$ (6 µM iron (III))

or with H_3BO_3 (0.47 mM) or the corresponding culture supernatant were added to the algal culture. After

incubation at $22 \pm 1°C$ under cool white fluorescent lights, living, non-swimming, group-attacked and

lysed phytoplankton cells were counted in a sedimentation chamber under an inverted microscope. The

number of lysed cells corresponded to algae showing disrupted membranes. Non-swimming algae were

not counted as lysed cells. For *Vibrio* analysis, 100 µL of a 1:10 serial dilution mixture in ENSW (from

10^{-2} to 10^{-10}) was plated on *Vibrio* Selective TCBS (thiosulfate-citrate-bile salts-sucrose) agar (in

triplicate). After incubation for 24 h at $22 \pm 1°C$, the number of living *Vibrio* cells was determined by

counting colony-forming units (CFUs). Data are the means of three independent experiments.

Microscope observations

The dynamic of the interaction between *A. pacificum* ACT03 and *V. atlanticus* LGP32, which is an oyster

pathogen isolated from the French Atlantic coast (Gay et al., 2004) and present in the Thau Lagoon, south

of France (Lopez-Joven et al., 2018)) was surveyed. As the algal strain used in the study is not axenic,

means that additional bacteria, other than the *V. atlanticus* LGP32, are present in the experiments.

To elucidate the interaction without thoroughly accounting for the non-axenic cells, interaction was

observed under a Zeiss Axio upright fluorescence microscope equipped with an AxioCamMRm 2 digital

microscope camera using *V. atlanticus* LGP32 tagged with green fluorescent protein (GFP). Lasers were

used at excitation wavelength (λex) 488 nm for GFP (emission wavelengths (λem): 505–530 nm) and λex

532 nm for plankton chloroplasts (λem 560–630 nm). Images were taken sequentially to avoid cross-

contamination between fluorochromes. Sequences of images were merged during the *Vibrio-Alexandrium*

interaction using ZEN 2012 (blue edition) software. Interaction events between *Vibrio* strains and

phytoplankton strains were also observed under a Leica TCS SPE confocal laser scanning system connected

to a Leica DM 2500 upright microscope camera (Montpellier RIO Imaging Platform, University of

Montpellier, France).

Comparative proteomic analysis

Vibrio sampling and protein extraction. V. atlanticus LGP32 was grown for 60 h at 22°C in artificial

seawater (high nutrient stress, cond. 1) or 12 h at 22°C in Zobell media (low nutrient stress, cond. 2). After

10 min centrifugation at 8000 rpm, crude protein extracts of *V. atlanticus* LGP32 in each culture condition

(triplicates) were obtained by sonication on ice at 20% amplitude for 20 s in 200 µL of ice-cold denaturing

buffer (7 M urea, 2 M thiourea, 4% CHAPS in 30 mM Tris-HCl, pH 8.5) and clarified by centrifugation at

2000 x *g*, 15 min, 4°C. The protein concentration of the supernatant was estimated using the 2D Quant Kit

(Cytiva™, MERCK) and samples were stored at -80°C until use.

Bi-dimensional gel electrophoresis (2D gel). Proteins extracts were individually analysed on 2D gel

electrophoresis (6 gels per condition each corresponding to different biological replicates). To do so, 100

µg of proteins from each extract was added to rehydration buffer (7 M urea, 2 M thiourea, 4% CHAPS, 65

mM DTT) for a total volume of 350 µL. They were then individually loaded onto 17 cm isoelectric

focusing strips (Bio-Rad) with a stabilized non-linear pH ranging from 3 to 10. Due to the high

complexity of the protein profile in the acidic part (left) of the gel pH 3–10 (Fig. S3), we conducted

additional 'close-up' analyses in gels using 17 cm isoelectric focusing strips (Bio-Rad) with a narrower,

stabilized pH gradient ranging from 4 to 7. Strips were rehydrated passively for 5 h at 22°C, followed by

active rehydration for 14 h under a 50 V current at 22°C (to help large proteins enter the strips).

Thereafter, isoelectrofocusing was carried out using the following programme: 50 V for 1 h, 250 V for 1

h, 8000 V for 1 h and a final step at 8000 V for a total of 140 000 V.h with a slow ramping voltage

(quadratic increasing voltage) at each step. Focused proteins were reduced by incubating the strip twice in

equilibration buffer (1.5 M Tris, 6 M urea, 2% SDS, 30% glycerol; bromophenol blue, pH 8.8) containing

DTT (130 mM) at 55°C. Then, they were alkylated by incubation with equilibration buffer containing

iodoacetamide (135 mM) on a rocking agitator (400 rpm) at room temperature protected from light.

Proteins were also separated according to their molecular weight (second dimension) on 12%

acrylamide/0.32% piperazine diacrylamide gels run at 25 mA per gel for 30 min followed by 75 mA per

gel for 8 h using a Protean II XL system (Bio-Rad). Gels were stained using an MS-compatible silver

staining protocol and scanned using a ChemiDoc MP Imaging System (Bio-Rad) associated with Image

Lab software version 4.0.1 (Bio-Rad).

Comparative bioinformatics analysis of 2D gels. Twelve gels (six per condition) were selected for

comparative analysis on PD-Quest v. 7.4.0 (Bio-Rad) to identify changes in protein abundance between

the proteomic profiles of *V. atlanticus* LGP32 cultured in contrasting nutrient conditions (ENSW/Zobell).

Spots whose mean intensity across six replicates per strain was two times higher or lower than those from

the other strain, with a $P <0.01$ (Mann-Whitney U-test), were considered significantly different in terms of

abundance between the two conditions (quantitative difference). Differentially represented spots were then

excised from the gels, destained, trypsin-digested and the obtained peptides were identified by tandem

mass spectrometry (MS-MS) using the PISSARO platform facility (University of Rouen, France). To

identify protein(s) present in each spot, the obtained peptides were compared with *V. atlanticus* LGP32

reference genome *(https://vibrio.biocyc.org/)*. The genes whose peptides matched strongly were retrieved

and used for an BLASTx query against non-redundant databases to determine the protein identity of the

best match. A gene was considered as strongly matched when at least two peptides matched the sequence

with a coverage of> 6%. Their theoretical isoelectric point (pI) and molecular weights were also

calculated using the Expasy server (https://www.expasy.org/) to compare them with the location of the

spot on the gel. Altogether, these complementary analyses made it possible to characterize the protein

identity of each spot with confidence.

Gene expression analysis

Vibrio sampling and RNA extraction. V. atlanticus LGP32 was grown in Zobell media for 36 h and 60 h at

22°C (decline phase of growth, nutrient stress) or for 12 h at 22°C (exponential grow phase, poor nutrient

stress). Total RNA was isolated from *V. atlanticus* LGP32 using the standard TRIzol method (Invitrogen

Life Technologies SAS, Saint-Aubin, France) and then treated with DNase (Invitrogen) to eliminate

genomic DNA contamination. After sodium acetate precipitation, the quantity and quality of the total

RNA were determined using a NanoDrop spectrophotometer and agarose gel electrophoresis. Following

heat denaturing (70°C for 5 minutes), reverse transcription was performed using 1 μg of RNA prepared

with 50 ng μL^{-1} oligo-(dT) 12–18 mer in a 20-μL reaction containing 1 mM dNTPs, 1 unit μL^{-1}

RNAseOUT and 200 units μL^{-1} Moloney murine leukaemia virus reverse transcriptase (M-MLV RT) in

reverse transcriptase buffer, according to the manufacturer's instructions (Invitrogen Life Technologies

SAS, Saint-Aubin, France).

PCR amplification. Amplification reactions were analysed using a Roche LightCycler 480 Real-Time

thermocycler (Bio-Environnement platform, University of Perpignan, France). In this study, several PCR

primer pairs were designed using Primer3 software (optimal primer size: 20 bases; Tm: 60°C; primer

GC%: 50; 2GC clamp and product size range: 150–200 bp) and calibrated with *V. atlanticus* LGP32

genomic DNA (Table 1). To determine the qPCR efficiency of each primer pair used, standard curves

were generated using seven serial dilutions of genomic DNA (10^{-2}, 10^{-3}, 10^{-4}, 10^{-5}, 10^{-5}, 10^{-7} and 10^{-8}) (not

shown); the qPCR efficiencies of the tested genes varied between 1.85 and 2.08 (Table 3). For gene

expression, reverse transcription was performed with 1 μg of total RNA using random hexamers and

SuperScript IV reverse transcriptase (Invitrogen). The total qPCR reaction volume was 10 μL and

consisted of 5 μL of cDNA (diluted 1:5), 2.5 μL of SensiFAST SYBR No-ROX Mix (Bioline) and 100

nM or 300 nM PCR primer pair (Table 3). The reaction parameters were as follows: 2 min at 95°C (initial

denaturation) and 40 cycles of 5 s at 95°C (denaturation), 10 s at 59°C (annealing) and 20 s at 72°C

(elongation). The specificity of each PCR was checked by measuring fluorescent signals during melting

curve analysis (PCR product heated from 65°C to 95°C continuously and slowly at $0.1°C \ s^{-1}$). Relative

expression was calculated by normalization to the expression of two constitutively expressed

housekeeping genes, namely, 6PKF (VS_2913) and CcmC (VS_0852), using the delta-delta threshold

cycle (ΔΔCt) method(Pfaffl, 2001).

Detection of quorum-sensing signalling molecules.

Vibrio culture. To detect the QS molecules (AI-2, AI-1 and CAI-1) *V. atlanticus* LGP32 was grown

in Zobell media for 12 h (exponential growth phase, control) and 60 h (decline phase of growth, nutrient

stress).

AI-2 analysis. Bioluminescence assay using the QS bioluminescent of *Vibrio campbellii* MM32

(*luxN*::Cm, *luxS*::Tn5Kan) was used to detect AI-2 molecules in culture supernatants. Briefly, *Vibrio*

cultures were centrifuged at 17,000 x *g* for 10 min, and the resulting supernatants were filtered on 0.22

μm. Then 20 μL of the filtrates were mixed with 180 μL of *V. campbellii* MM32 diluted 1:5000 then

incubated at 30°C and 100 rpm. Luminescence and cell density (OD620) were collected in triplicate and

analysed according to Tourneroche et al.(Tourneroche et al., 2019).

AI-1 and CAI-1 extraction and LC-MS analysis. Chemical analyses were conducted with a Q Exactive

Focus Orbitrap System coupled to an Ultimate 3000™ ultrahigh-performance liquid chromatography

(UHPLC) system (Thermo Fisher Scientific) according to Rodrigues et al. (Rodrigues et al., 2022).

Briefly, ethyl acetate (2 mL) was added into each culture (2 mL). This mixture was shaken overnight at

room temperature (150 rpm). The two phases were then separated and the aqueous phase was extracted

once again. The two obtained organic phases were pooled and the solvent was evaporated under vacuum.

The crude extracts were dissolved in 500 μL LC-MS grade methanol for analysis. All experiments were

conducted in triplicate. Analyses of extracts and standards (3 μL injected) were performed in electrospray

positive ionization mode in the 50–750 m/z range in centroid mode. The parameters were as follows:

spray voltage: 3 kV; sheath flow rate: 75; aux gas pressure: 20; capillary temperature: 350°C; heater

temperature: 430°C. The analysis was conducted in Full MS data-dependent MS2 mode (Discovery

mode). Resolution was set to 70,000 in Full MS mode, and the AGC (automatic gain control) target was

set to 1×10^6. In MS2, resolution was 17,500, AGC target was set to 2×10^5, isolation window was 0.4 m/z,

and normalized collision energy was stepped to 15, 30 and 40 eV. The UHPLC column was a

Phenomenex Luna Omega Polar C18 1.6 μm, 150 x 2.1 mm. The column temperature was set to 42°C,

and the flow rate was 0.4 mL min^{-1}. The solvent system was a mixture of water (A) with increasing

proportions of acetonitrile (B), with both solvents modified with 0.1% formic acid. The gradient was as

follows: 1% B 3 min before injection, then from 1 to 15 min, a gradient increase of B up to 100% (curve

5), followed by 100% B for 5 min. The flow was injected into the mass spectrometer starting immediately

after injection. All data were acquired and processed using FreeStyle 1.5 software (Thermo Fisher

Scientific).

Chemicals and solvents. N-acyl-homoserine lactones (AHL) were obtained from Cayman Chemical (Ann

Arbor, MI, USA). Stock solutions were obtained by dissolving standards in methanol or dichloromethane

(C18-AHL) at a concentration of 1 mg mL^{-1} and stored at -80°C. Standard solutions for UHPLC- high-

resolution tandem mass spectrometry (HRMS) analyses were prepared by diluting each individual

standard solution with methanol in order to obtain a concentration range from 2000 to 20 ng mL^{-1}. LC-MS

grade methanol, acetonitrile and formic acid were purchased from Biosolve (Biosolve Chimie, Dieuze,

France), analytical-grade ethyl acetate was obtained from Sigma-Aldrich. Pure water was obtained from

Elga Purelab Flex System (Veolia LabWater STI, Antony, France).

Nature of lytic compounds secreted by *V. atlanticus* LGP32.

To determine the nature of the lytic compounds secreted, *V. atlanticus* LGP32 grown for 60 h in Zobell

media at 22 °C was filtered through a 10 kDa membrane (Amicon® Ultra-4 filter unit). The eluate

containing molecules with MW below 10 KDa was then incubated in a water bath at 100°C for 30 min.

Boiled filtrates (0.1 mL) were subsequently used to inoculate *A. pacificum* (ACT03 strain) cultures, then

lytic activity was observed under the Leica TCS SPE confocal laser scanning system. Zobell media with

the same treatment was used as control.

Statistical analysis.*Environmental data*. Statistical analyses were performed using R 3.6.3 software. The

relationship between *Alexandrium* and *Vibrio* was explored separately in spring and autumn. We used a

generalized linear model specifying a Gaussian family. For spring, the dataset for salinity and temperature

was complete (14 periods of observation). One influential period was detected and removed from the

dataset whenever no dead *Alexandrium* cells were observed. The effects of explanatory variables such as

log10 (Vibrio+1), salinity and temperature were centred, reduced and tested as fixed effects with a linear

relationship. Model selection was performed using the Akaike information criterion corrected for small

sample size (AIC$_c$). Models were considered different whenever the difference between their AIC$_c$ value

and the lowest AIC$_c$ value (ΔAIC$_c$) was lower than 2 (Burnham and Anderson, 2002). *Alexandrium*

distribution and model residuals were checked for normal distribution assumptions (QQ plot and Shapiro-

Wilk test). For autumn, the dataset was complete for 10 periods of observation. Salinity and temperature

were missing for three periods. We explored the relationship between *Alexandrium* and *Vibrio* alone using

the method detailed above for spring.

In vitro data. Statistical analyses were performed using one-way ANOVA (analysed by pair) followed by

Tukey's test (Statistica 10.0 software, StatSoft, Maison-Alfort, France). $*P < 0.05$, $**P < 0.01$, $***P$

< 0.001

RESULTS

Concomitant occurrence of *A. pacificum* ACT03, *A. tamarense* ATT07 and free-living *Vibrio spp.* in

the Thau lagoon. In the spring and autumn of 2015, in the Thau Lagoon (Fig. 1A), we detected

Alexandrium algae (*A. pacificum* ACT03 *and A. tamarense* ATT07, both alive and in degraded cell forms)

and free-living *Vibrio*, but no plankton-associated *Vibrio* were observed (Fig. 1B, Data S1). Using model

selection based on AICc, we found no significant relationship between *Alexandrium* (*A. pacificum*

ACT03*, A. tamarense*) and *Vibrio Spp.* abundances in autumn. This result is consistent with the difficulty

that *Vibrio* has in growing at temperatures below 20°C and with the complex interacting factors driving

bloom dynamics (Laanaia et al., 2013). Interestingly, in spring 2015, the mean densities of *Alexandrium*

and of free-living *Vibrio* were positively correlated. The lowest AICc was obtained with the model

explaining degraded form of *Alexandrium* density based on the free *Vibrio* density (Fig. 1C). Given that,

this model is not so different from the model with only the intercept, but better than any other linear

combination with other potentially interfering drivers, such as temperature and salinity (Fig. 1D), we

searched for evidence of a relationship between *Vibrio* and *Alexandrium* by studying their interaction *in*

vitro.

335 *Vibrio atlanticus* **LGP32 feeds on** *Alexandrium pacificum* **ACT03**.

336 To investigate whether *Alexandrium* interacts with *Vibrio*, we incubated in Enriched Natural SeaWater

337 (ENSW) *A. pacificum* ACT03 with *V. atlanticus* LGP32 previously grown for 12 hours in Zobell media.

338 In interaction with *V. atlanticus* LGP32, *A. pacificum* ACT03 cell abundance decreased significantly from

339 2.10×10^4 cells mL^{-1} to 1.07×10^4 cells mL^{-1} after 48 h (Fig. 2A), while the *V. atlanticus* LGP32

340 concentration grew significantly after 26 h of interaction, reaching a maximum peak density of 7.67×10^7

341 CFU mL^{-1} at 34 h (Fig. 2B). In the control experiment where *A. pacificum* ACT03 was cultured alone in

342 ENSW, the algal concentration remained stable over time (Fig. 2A) and no bacteria were on the

343 corresponding TCBS plates (*Vibrio* selective medium). In the control where *V. atlanticus* LGP32 was

344 cultured alone in ENSW, the concentration of the bacteria tended to increase during the first 24 h (from

345 2.0×10^5 to 7.0×10^5 CFU mL^{-1}) and then decreased at 48h, below the initial starting concentration (Fig.

346 2B). These results show that the interaction between *V. atlanticus* LGP32 and *A. pacificum* ACT03 leads

347 to a decline in the algal population and promotes the growth of *V. atlanticus* LGP32. This suggests that *V.*

348 *atlanticus* LGP32 is able to feed on *A. pacificum* ACT03.

349 *V. atlanticus* **LGP32 performs group-attacks on** *A. pacificum* **ACT03**.

350 Epifluorescence microscopy observation of GFP-labelled *V. atlanticus* LGP32 (previously grow in Zobell

351 medium) in interaction showed that *A. pacificum* ACT03 cells that had lost their motility were attacked

352 individually by groups of *V. atlanticus* LGP32 before being lysed (Movie 1). Attacks were extremely

353 rapid, with empty thecae (algal envelopes) observed in the medium after less than 60 s (Movie 2). During

354 the attack, *V. atlanticus* LGP32 did not invade the algal cell but remained clustered on the cell surface

355 (Fig. 3A2).

356 **Group attack of** *A. pacificum* **ACT03 is activated by** *V. atlanticus* **LGP32 starvation.**

357 In 2002, Martin hypothesized that nutritional stress induces bacteria to lyse algae (Martin, 2002). To test

358 this hypothesis, we monitored *V. atlanticus* LGP32 behaviour in response to starvation (Fig. 3). We

observed that *V. atlanticus* LGP32 in exponentially growth phase (12 h of culture in Zobell medium) did

not interact with *A. pacificum* ACT03 cells for the first hour of contact (Fig. 3A, B, C). In contrast, *V.*

atlanticus LGP32 in the decline phase (36 h of culture in Zobell medium) induced a significant decrease in

the number of motile algae cells by 8.9% after 15 min and by 43.3% after 60 min (Fig. 3A). This

phenomenon corresponded to the degradation and/or disruption of algal flagella (Movie 3). The flagella no

longer functioned correctly, which caused erratic swimming of the algae (Video 3, left cell). This was

followed by a complete cessation of swimming. When the flagellum detached from the algae (Video 3,

right cell), the attack occurred. With starved *V. atlanticus* LGP32 (60 h of culture in Zobell medium),

algae immobilization was fast and significant (91.4% in 15 min, Fig. 3A), and algae were attacked

individually being targeted by groups of *V. atlanticus* LGP32 (Movie 4). The percentage of cells attacked

and killed peaked at 30% after 15–30 min of contact (Fig. 3B) and then decreased. After 1 h, attacks had

stopped with approximately 40% of the algal cells still alive (Fig. 3C). Although it remains unclear

whether the attacks occur during a specific phase of growth, it is evident that the cells are already

weakened before attack as they have all lost their flagella. An old-starved culture of *V. atlanticus* LGP32

(126 h) significantly immobilized *A. pacificum* ACT03 cells within a few minutes, with lysis occurring

immediately (Fig. 3A, C), making it impossible to detect attacks by *V. atlanticus* LGP32 (Fig. 3B). The

lysis phase corresponded to initial vesicle formation followed by the bursting of *A. pacificum* ACT03 cells

(Movie 5) and was induced by the old-starved culture supernatant of *V. atlanticus* LGP32 (Fig. S1).

We next tested whether this lytic effect was mediated by thermostable molecule (s) secreted by *Vibrio*.

The culture supernatant of starved culture of *V. atlanticus* LGP32 (36 h) filtered through a 10 kDa

membrane and then incubated at 100°C for 30 min still possessed its lytic properties, indicating that the

algicidal compounds produced by *V. atlanticus* LGP32 are small thermostable molecules unlikely to be

lytic enzymes, or lysins able to digest the algae cell. However, these experimental observations clearly

show the key role of nutrient limitation in triggering the 'group-attack' behaviour and the secretion of lytic

compounds of *V. atlanticus* LGP32.

Group attack occurs on *A. pacificum* ACT03 in exponential phase of growth.

Here, we wondered whether the live/dead status of algae is important for *V. atlanticus* LGP32-mediated group attacks targeting. To this aim, *A. pacificum* ACT03 in exponential growth phase was first exposed for 30 min to supernatant from 60 hours starved *V. atlanticus* LGP32 Zobell media that induced 25% lysis of *A. pacificum* ACT03 cells and next to the corresponding *V. atlanticus* LGP32 cells. Group attacks were observed on non-degraded *A. pacificum* ACT03 cells, but not on lysed cells. This result is similar to what is observed on the Movie 1 with flash attacks only on immobilized, but not degraded *A. pacificum* ACT03 cells (red), and not on lysed cells (green). In addition, no group attacks occurred on cells from an old *A. pacificum* ACT03 culture (1-month culture). Together, with the very short duration of group attacks (Movie 1), these results indicate that *V. atlanticus* LGP32 group-attacked exponentially growing cells of *A. pacificum* ACT03, but not decomposing algae, suggesting that this behaviour is not just an opportunistic response of heterotrophic bacteria to organic substrates.

Group attack is independent of quorum sensing.

Considering the coordinated action of *V. atlanticus* LGP32 in attacks on *A. pacificum* ACT03, we tested whether the group-attack process depended on the key physiological mechanism that regulates many functions in marine microbial cells, quorum sensing (QS) (Lami, 2019; Papenfort and Bassler, 2016), a type of cell-cell communication. Although QS is a cell-density-dependent mechanism, our results showed no attack from a 12 h culture of *V. atlanticus* LGP32 up to a concentration of 4.10^6 *Vibrio* mL^{-1} (Fig 4A). Attacks were only observed with *V. atlanticus* LGP32 from a 60 h culture at low concentration of 5.10^3 *Vibrio* mL^{-1} to the highest concentration tested of 5.10^5 *Vibrio* mL^{-1} (Fig 4A), consistent with the hypothesis that the attacks were independent of *Vibrio* density.

The analysis of the expression of genes involved in the known QS pathways in *Vibrio* cell (Fig. S3A),

highlighted that only the AI-2 pathway was induced during nutrient stress of *V. atlanticus* LGP32, because

only the expression of the AI synthase (*LuxS*) and its receptor (*LuxP*) increased significantly (Fig. 4B;

ANOVA p <0.05). This was confirmed by a QS bioluminescence assay, which showed a AI-2 molecules

(unquantified) in the Zobell culture supernatant of *V. atlanticus* LGP32 after 60 h of culture but not after

12 h of culture and not in the ENSW supernatant of *V. atlanticus* LGP32 after 12 or 60 h of culture.

UHPLC-HRMS/MS provided no evidence of detectable HAI-1) and CAI-1 in any experiments.

By targeted mutagenesis of key genes involved in QS pathways Δ*luxM* (HAI-1 production), Δ*luxS* (AI-2

production) and Δ*luxR* (high-density QS master regulator) did not lead to any change in the group-attack

behaviour of *V. atlanticus* LGP32 (Fig. 4C). Taken together these results showed that group-attack by *V.*

atlanticus LGP32 is not coordinated by QS.

Group attack related to the availability of iron.

The comparative analysis of the proteome of *V. atlanticus* LGP32 incubated 60 h in artificial seawater

(ENSW) versus *V. atlanticus* LGP32 grown 12 h in Zobell nutrient-rich medium revealed 10 proteins

modulated by nutrient stress (Fig. S2). The two most down-regulated proteins correspond to be ß-

ketoacyl-(acyl-carrier-protein) synthase II (-22-fold in ENSW compared to Zobell), a key regulator of

bacterial fatty acid synthesis, and the dihydroorotase (-6.6-fold in ENSW compared to Zobell), an enzyme

essential for pyrimidine biosynthesis and thus bacterial proliferation and growth. The low expression of

these proteins in ENSW is consistent with *V. atlanticus* LGP32 nutritional starvation. The most up-

regulated protein in starved *V. atlanticus* LGP32, with an increase of more than 6-fold, was glucosamine-

6-phophate deaminase, an enzyme involved in bacterial energy metabolism probably necessary for its

survival. Among the other up -regulated proteins, one was an iron siderophore-binding protein (Spot 4413,

Fig. S2A) corresponding to the vibrioferrin outer membrane receptor PvuB, whose gene is part of the *pvu*

operons involved in iron transport (Fig. S3B). Interestingly, the corresponding gene *pvuB* as well as the

vibrioferrin membrane receptor gene *pvuA2* (Fig. S3B) were both significantly induced in *Vibrio* under

nutrient stress (Fig. 4B; ANOVA p <0.01) but not the one involved in the vibrioferrin biosynthesis, *pvsA*

(Fig. 4B). Remarkably, among the 10 proteins identified by proteomic analysis only *V. atlanticus* LGP32

mutant lacking *pvuB* failed to attack *A. pacificum* ACT03 (Fig. 4C; ANOVA p <0.001). In the absence of

the *pvuB* gene, *V. atlanticus* LGP32 was unable to attack in group *A. pacificum* ACT03. In addition, *V.*

atlanticus LGP32 cells that had been washed with ENSW to remove their culture supernatant metabolites

also failed to attack *A. pacificum* ACT03 (Fig. 4C; ANOVA p <0.001), which is congruent with the

hypothesis that attacks depend on the *V. atlanticus* LGP32 vibrioferrin transport system. Finally, group-

attacks increased significantly, when $FeCl_3$ was added to the *Vibrio* culture medium (Fig. 4D) but not with

H_3BO_4 (Fig. 4D). Taken together, those results are consistent with the hypothesis that attacks are regulated

by iron.

Group attack is a *Vibrio spp.* behavior specific to *Alexandrium spp.*

To evaluate the dinoflagellates specificity of the group-attack behaviour, a selection of *Vibrio spp.* was co-

cultured with a selection of dinoflagellate strains commonly found in the Mediterranean Sea. The results

showed that, among the *Vibrio spp.* tested (pathogenic or not) all, under nutrient stress, were able to

secrete algicidal compounds, immobilize, 'group-attack' and lyse *A. pacificum* ACT03 cells (Table 1) and

no tropism linked to their pathogenesis for fish of invertebrates was observed (Table 1). Among the

sixteen dinoflagellates species tested, all eight *Alexandrium spp.* (non-toxic and paralytic shellfish toxin

(PST) producers) and *Gymnodinium catenatum* (PST producer) were immobilized, 'group-attacked' and

lysed by *V. atlanticus* LGP32 (Table 2), but no tropism linked to PSTs was revealed (Table 2).

DISCUSSION

Predation is a widespread mode of interaction for survival in the natural world (Finke and Denno, 2004;

Sinclair et al., 2003). Among predators, predatory bacteria are found in a wide variety of environments,

and like bacteriophages and predatory protists, they have been reported to prey exclusively on other

bacteria (Johnke et al., 2014; Perez et al., 2016). Considering predator as a free organism that feeds at the

expense of another, this study is the first evidence of the capacity of some *Vibrio* to develop a predatory

strategy against an alga. This behaviour differs from parasitism, because the survival of *Vibrio* is not

exclusively dependent on the algae.

The strategy developed by *Vibrio* to kill algae may be reminiscent of strategies previously described in the

prokaryotes (Johnke et al., 2014). As shown in Movie 1, the interaction between *V. atlanticus* LGP32 and

A. pacificum ACT03 proceeds in three stages (Fig. 5). The first stage, the '**immobilization stage**', recalls

the strategy used by *Streptomyces* to immobilize its prey (Kumbhar et al., 2014) based on the secretion of

algicidal metabolites that disrupt the flagella. The second stage, the '**attack stage**', resembles the 'wolf-

pack attack' strategy described for *Myxococcus xanthus* and *Lysobacter* (Martin, 2002; Perez et al.,

2014). *V. atlanticus* LGP32 also surrounds *A. pacificum* ACT03 cells at high density for a very short time,

but does not invade the algal cell. This phenomenon is comparable to that of bacteria clustering around

lysed ciliate cells (Blackburn et al., 1998). The third stage, the '**killing stage**', is similar to that of epibiotic

bacterial predators, which induces the lysis of bacteria from the outside (Rashidan and Bird, 2001).

Overall, these observations show that *V. atlanticus* LGP32 is able of wolf-pack hunting behaviour.

Results showed that quorum sensing does not control algal predation by *Vibrio*, in contrast to iron

acquisition systems, which play a major role in regulating this phenotype, as we demonstrated genetically.

Indeed, our quorum sensing mutants exhibited the same predatory phenotype as the wild-type *Vibrio*

atlanticus LGP32, whereas mutants impaired in iron uptake completely lost this phenotype. However,

quorum sensing and iron acquisition are sometimes interconnected in *Vibrio* (McRose et al., 2018). For

instance, in *Vibrio vulnificus*, the production of vulnibactin (a siderophore) is controlled by AI-2 (Kim and

Shin, 2011). Whether the production of AI-2 by *V. atlanticus* LGP32, induced during group attacks of *A.*

pacificum ACT03, promotes the production of vibrioferrin (Lami, 2019) remains to be investigated.

We further show that the group-attack behaviour of *V. atlanticus* LGP32 depends on iron availability. Iron

is an essential element for growth in most organisms, including phytoplankton (Martin and Fitzwater,

1988) and bacteria (Neilands, 1981), and its concentration in seawater is known to be very low with

measurements in the open ocean surface at 0.2 nM and in deep waters at 0.6 nM (Millero, 1998).

Moreover, Its low solubility in seawater limits its availability (Bruland et al., 1994; Wu and Luther, 1994).

To acquire iron, bacteria have developed systems based on the secretion (and subsequent uptake) of iron-

chelating siderophores to obtain this element from the environment (Amin et al., 2009a). Therefore, many

Vibrio spp. produce a siderophore known as vibrioferrin, which is synthesized and secreted by proteins

encoded by the *pvsABCDE* gene cluster (Fig. S3). Boron being known to be a regulator or capable of

being transported by vibrioferrin (Romano et al., 2013; Weerasinghe et al., 2013) we tested its potential

involvement in the interaction but no effect was evidenced here. For iron-vibrioferrin uptake, *Vibrio*

parahaemolyticus uses a membrane siderophore receptor, called PvuA, which is coupled to an inner

membrane ATP-binding cassette (ABC). This ABC transporter system comprised of proteins encoded by

the pvuABCDE gene cluster (Fig. S3) is required for transporting the siderophore across the inner

membrane (Tanabe et al., 2003). Siderophores are not only iron carriers but also important regulators of

virulence (Miethke and Marahiel, 2007) and mediators of bacterial interaction with phytoplankton (Amin

et al., 2009b; Kramer et al., 2020). We showed here a pivotal role of iron in the interaction between *V.*

altlanticus LGP32 and *A. pacificum* ACT03. This mirrors the mutualistic interaction observed between

Gymnodinium catenatum and *Marinobacter* (Amin et al., 2009b). In fact, in natural settings, the co-

occurence of *Marinobacter* and *G. catenatum* is suggested to depend on a mutually beneficial utilization

of iron and carbon resources (Bolch et al., 2011). As in the present study, iron seems to play a key role in

the interaction. Indeed, the labile iron released through the photolysis of ferric chelates with vibrioferrin

providing a crucial iron source for phytoplankton, which need substantial amounts of iron to support

carbon fixation through photosynthesis (Amin et al., 2009a; Yang et al., 2021). This fixed carbon, in turn,

sustains the growth of both the phytoplankton and their bacterial counterparts (Amin et al., 2009b; Kramer

et al., 2020). Future study could demonstrate further the role of vibrioferrin in group attack, by adding

iron-saturated vibrioferrin to algae-*Vibrio* co-cultures.

In the natural environment, associations between bacteria and algae have already been observed (Lopez-

Joven et al., 2018; Miller et al., 2005; Rosales et al., 2022; Xu et al., 2022). Although we do not have

definitive proof of a predator/prey interaction *in situ* due to confounding environmental variables,

environmental data acquired in the Thau Lagoon showed a correlation between *Alexandrium* and *Vibrio*

occurrence, suggesting that such interaction could occur in nature. Evolution of such a predatory behavior

would provide an important ecological advantage to *Vibrio* to obtain nutrients in environment, where

Alexandrium spp. and *Gymnodinium catenatum* form blooms.

With more than 30 species distributed all over the world (Anderson et al., 2012; Hallegraeff et al., 2021),

Alexandrium spp. and *Gymnodinium spp.* considered as invasive species by the Delivering Alien Invasive

Species Inventories for Europe (http://www.europe-aliens.org)), could play an unexpected and important

role in maintaining, structuring and regulating *Vibrio* populations in the ecosystem. In turn, *Vibrio* could

contribute to the regulation and control of their blooms.

To conclude, this study reveals the capacity of some *Vibrio spp.* to be facultative predators that hunt

specific algae. In the current context of climate change favourable to their development monitoring the

invasive algae, *Alexandrium spp.* and *Gymnodinium catenatum* should be considered not only for their

potent harmful effect on humans and animals, but also as a potential source of nutrients for the expansion

of *Vibrio,* particularly pathogenic species (Lemire et al., 2015).

DATA AVAILABILITY

All data on environmental study are included in the article and/or supporting information files. All other

study data are available on request from the corresponding author.

REFERENCES

Abadie, E., Amzil, Z., Belin, C., Comps, M.A., Elzière-Papayanni, P., Lassus, P., le Bec, C., Baut, M.L., Nezan, E., Poggi, R., n.d. Contamination de l'étang de Thau par Alexandrium tamarense (No. Rapport 884). Ifremer.

Abadie, E., Kaci, L., Berteaux, T., Hess, P., Sechet, V., Masseret, E., Rolland, J.L., Laabir, M., 2015. Effect of Nitrate, Ammonium and Urea on Growth and Pinnatoxin G Production of Vulcanodinium rugosum. Mar. Drugs 13, 5642–5656. https://doi.org/10.3390/md13095642

Amin, S.A., Green, D.H., Hart, M.C., Kuepper, F.C., Sunda, W.G., Carrano, C.J., 2009a. Photolysis of iron-siderophore chelates promotes bacterial-algal mutualism. Proc. Natl. Acad. Sci. U. S. A. 106, 17071–17076. https://doi.org/10.1073/pnas.0905512106

Amin, S.A., Green, D.H., Kuepper, F.C., Carrano, C.J., 2009b. Vibrioferrin, an Unusual Marine Siderophore: Iron Binding, Photochemistry, and Biological Implications. Inorg. Chem. 48, 11451–11458. https://doi.org/10.1021/ic9016883

Anderson, D.M., Alpermann, T.J., Cembella, A.D., Collos, Y., Masseret, E., Montresor, M., 2012. The globally distributed genus Alexandrium: Multifaceted roles in marine ecosystems and impacts on human health. Harmful Algae 14, 10–35. https://doi.org/10.1016/j.hal.2011.10.012

Baker-Austin, C., Trinanes, J., Gonzalez-Escalona, N., Martinez-Urtaza, J., 2017. Non-Cholera Vibrios: The Microbial Barometer of Climate Change. Trends Microbiol. 25, 76–84. https://doi.org/10.1016/j.tim.2016.09.008

Ben-Gharbia, H., Yahia, O.K.-D., Amzil, Z., Chomerat, N., Abadie, E., Masseret, E., Sibat, M., Triki, H.Z., Nouri, H., Laabir, M., 2016. Toxicity and Growth Assessments of Three Thermophilic Benthic Dinoflagellates (Ostreopsis cf. ovata, Prorocentrum lima and Coolia monotis) Developing in the Southern Mediterranean Basin. Toxins 8, 297. https://doi.org/10.3390/toxins8100297

Blackburn, N., Fenchel, T., Mitchell, J., 1998. Microscale nutrient patches in planktonic habitats shown by chemotactic bacteria. Science 282, 2254–2256. https://doi.org/10.1126/science.282.5397.2254

Bolch, C.J.S., Subramanian, T.A., Green, D.H., 2011. THE TOXIC DINOFLAGELLATE *GYMNODINIUM CATENATUM* (DINOPHYCEAE) REQUIRES MARINE BACTERIA FOR GROWTH[1]. J. Phycol. 47, 1009–1022. https://doi.org/10.1111/j.1529-8817.2011.01043.x

Bruland, K., Orians, K., Cowen, J., 1994. Reactive Trace-Metals in the Stratified Central North Pacific. Geochim. Cosmochim. Acta 58, 3171–3182. https://doi.org/10.1016/0016-7037(94)90044-2

Brumfield, K.D., Usmani, M., Chen, K.M., Gangwar, M., Jutla, A.S., Huq, A., Colwell, R.R., 2021. Environmental parameters associated with incidence and transmission of pathogenic *Vibrio spp* . Environ. Microbiol. 23, 7314–7340. https://doi.org/10.1111/1462-2920.15716

Burkholder, J.M., Shumway, S.E., Glibert, P.M., 2018. Food Web and Ecosystem Impacts of Harmful Algae, in: Harmful Algal Blooms. John Wiley & Sons, Ltd, pp. 243–336. https://doi.org/10.1002/9781118994672.ch7

Burnham, K.P., Anderson, D.R. (Eds.), 2002. Summary, in: Model Selection and Multimodel Inference: A Practical Information-Theoretic Approach. Springer, New York, NY, pp. 437–454. https://doi.org/10.1007/978-0-387-22456-5_8

Chai, Z.Y., Wang, H., Deng, Y., Hu, Z., Tang, Y.Z., 2020. Harmful algal blooms significantly reduce the resource use efficiency in a coastal plankton community. Sci. Total Environ. 704, 135381. https://doi.org/10.1016/j.scitotenv.2019.135381

Coyne, K.J., Wang, Y., Johnson, G., 2022. Algicidal Bacteria: A Review of Current Knowledge and Applications to Control Harmful Algal Blooms. Front. Microbiol. 13, 871177. https://doi.org/10.3389/fmicb.2022.871177

572 Drebes Dörr, N.C., Blokesch, M., 2020. Interbacterial competition and anti-predatory behaviour of
573    environmental *Vibrio cholerae* strains. Environ. Microbiol. 22, 4485–4504.
574    https://doi.org/10.1111/1462-2920.15224
575 Espinoza-Vergara, G., Hoque, M.M., McDougald, D., Noorian, P., 2020. The Impact of Protozoan
576    Predation on the Pathogenicity of Vibrio cholerae. Front. Microbiol. 11, 17.
577    https://doi.org/10.3389/fmicb.2020.00017
578 Fertouna-Bellakhal, M., Dhib, A., Fathalli, A., Bellakhal, M., Chomérat, N., Masseret, E., Laabir, M., Turki,
579    S., Aleya, L., 2015. Alexandrium pacificum Litaker sp. nov (Group IV): Resting cyst distribution and
580    toxin profile of vegetative cells in Bizerte Lagoon (Tunisia, Southern Mediterranean Sea). Harmful
581    Algae 48, 69–82. https://doi.org/10.1016/j.hal.2015.07.007
582 Finke, D.L., Denno, R.F., 2004. Predator diversity dampens trophic cascades. Nature 429, 407–410.
583    https://doi.org/10.1038/nature02554
584 Gay, M., Renault, T., Pons, A.M., Le Roux, F., 2004. Two Vibrio splendidus related strains collaborate to
585    kill Crassostrea gigas: taxonomy and host alterations. Dis. Aquat. Organ. 62, 65–74.
586    https://doi.org/10.3354/dao062065
587 Genovesi, B., Shin-Grzebyk, M.-S., Grzebyk, D., Laabir, M., Gagnaire, P.-A., Vaquer, A., Pastoureaud, A.,
588    Lasserre, B., Collos, Y., Berrebi, P., Masseret, E., 2011. Assessment of cryptic species diversity
589    within blooms and cyst bank of the Alexandrium tamarense complex (Dinophyceae) in a
590    Mediterranean lagoon facilitated by semi-multiplex PCR. J. Plankton Res. 33, 405–414.
591    https://doi.org/10.1093/plankt/fbq127
592 Hadjadji, I., Laabir, M., Frihi, H., Collos, Y., Shao, Z.J., Berrebi, P., Abadie, E., Amzil, Z., Chomérat, N.,
593    Rolland, J.L., Rieuvilleneuve, F., Masseret, E., 2020. Unsuspected intraspecific variability in the
594    toxin production, growth and morphology of the dinoflagellate Alexandrium pacificum R.W.
595    Litaker (Group IV) blooming in a South Western Mediterranean marine ecosystem, Annaba Bay
596    (Algeria). Toxicon 180, 79–88. https://doi.org/10.1016/j.toxicon.2020.04.005
597 Hallegraeff, G.M., Anderson, D.M., Belin, C., Bottein, M.-Y.D., Bresnan, E., Chinain, M., Enevoldsen, H.,
598    Iwataki, M., Karlson, B., McKenzie, C.H., Sunesen, I., Pitcher, G.C., Provoost, P., Richardson, A.,
599    Schweibold, L., Tester, P.A., Trainer, V.L., Yñiguez, A.T., Zingone, A., 2021. Perceived global
600    increase in algal blooms is attributable to intensified monitoring and emerging bloom impacts.
601    Commun. Earth Environ. 2, 117. https://doi.org/10.1038/s43247-021-00178-8
602 Harrison, P., Waters, R., Taylor, F., 1980. A Broad-Spectrum Artificial Seawater Medium for Coastal and
603    Open Ocean Phytoplankton. J. Phycol. 16, 28–35. https://doi.org/10.1111/j.1529-
604    8817.1980.tb00724.x
605 Johnke, J., Cohen, Y., de Leeuw, M., Kushmaro, A., Jurkevitch, E., Chatzinotas, A., 2014. Multiple micro-
606    predators controlling bacterial communities in the environment. Curr. Opin. Biotechnol. 27, 185–
607    190. https://doi.org/10.1016/j.copbio.2014.02.003
608 Johnson, C.N., 2013. Fitness Factors in Vibrios: a Mini-review. Microb. Ecol. 65, 826–851.
609    https://doi.org/10.1007/s00248-012-0168-x
610 Kim, C.M., Shin, S.H., 2011. Modulation of Iron-Uptake Systems by a Mutation of luxS Encoding an
611    Autoinducer-2 Synthase in Vibrio vulnificus. Biol. Pharm. Bull. 34, 632–637.
612    https://doi.org/10.1248/bpb.34.632
613 Kitatsukamoto, K., Oyaizu, H., Nanba, K., Simidu, U., 1993. Phylogenetic-Relationships of Marine-
614    Bacteria, Mainly Members of the Family Vibrionaceae, Determined on the Basis of 16s
615    Ribosomal-Rna Sequences. Int. J. Syst. Bacteriol. 43, 8–19. https://doi.org/10.1099/00207713-
616    43-1-8

Kramer, J., Oezkaya, O., Kuemmerli, R., 2020. Bacterial siderophores in community and host interactions.
Nat. Rev. Microbiol. 18, 152–163. https://doi.org/10.1038/s41579-019-0284-4
Kumbhar, C., Mudliar, P., Bhatia, L., Kshirsagar, A., Watve, M., 2014. Widespread predatory abilities in
the genus Streptomyces. Arch. Microbiol. 196, 235–248. https://doi.org/10.1007/s00203-014-
0961-7
Laabir, M., Jauzein, C., Genovesi, B., Masseret, E., Grzebyk, D., Cecchi, P., Vaquer, A., Perrin, Y., Collos, Y.,
2011. Influence of temperature, salinity and irradiance on the growth and cell yield of the
harmful red tide dinoflagellate Alexandrium catenella colonizing Mediterranean waters. J.
Plankton Res. 33, 1550–1563. https://doi.org/10.1093/plankt/fbr050
Laanaia, N., Vaquer, A., Fiandrino, A., Genovesi, B., Pastoureaud, A., Cecchi, P., Collos, Y., 2013. Wind and
temperature controls on Alexandrium blooms (2000-2007) in Thau lagoon (Western
Mediterranean). Harmful Algae 28, 31–36. https://doi.org/10.1016/j.hal.2013.05.016
Labreuche, Y., Le Roux, F., Henry, J., Zatylny, C., Huvet, A., Lambert, C., Soudant, P., Mazel, D., Nicolas, J.-
L., 2010. Vibrio aestuarianus zinc metalloprotease causes lethality in the Pacific oyster
Crassostrea gigas and impairs the host cellular immune defenses. Fish Shellfish Immunol. 29,
753–758. https://doi.org/10.1016/j.fsi.2010.07.007
Lami, R., 2019. Quorum Sensing in Marine Biofilms and Environments, in: Quorum Sensing, Academic
Press. Elsevier, pp. 55–96. https://doi.org/10.1016/B978-0-12-814905-8.00003-4
Le Roux, F., Binesse, J., Saulnier, D., Mazel, D., 2007. Construction of a *Vibrio splendidus* Mutant Lacking
the Metalloprotease Gene *vsm* by Use of a Novel Counterselectable Suicide Vector. Appl.
Environ. Microbiol. 73, 777–784. https://doi.org/10.1128/AEM.02147-06
Leblad, B.R., Amnhir, R., Reqia, S., Sitel, F., Daoudi, M., Marhraoui, M., Abdellah, M.K.O., Veron, B., Er-
Raioui, H., Laabir, M., 2020. Seasonal variations of phytoplankton assemblages in relation to
environmental factors in Mediterranean coastal waters of Morocco, a focus on HABs species.
Harmful Algae 96, 101819. https://doi.org/10.1016/j.hal.2020.101819
Lemire, A., Goudenege, D., Versigny, T., Petton, B., Calteau, A., Labreuche, Y., Le Roux, F., 2015.
Populations, not clones, are the unit of vibrio pathogenesis in naturally infected oysters. Isme J.
9, 1523–1531. https://doi.org/10.1038/ismej.2014.233
LeRoux, F., Wegner, K.M., Baker-Austin, C., Vezzulli, L., Osorio, C.R., Amaro, C., Ritchie, J.M., Defoirdt, T.,
Destoumieux-Garzon, D., Blokesch, M., Mazel, D., Jacq, A., Cava, F., Gram, L., Wendling, C.C.,
Strauch, E., Kirschner, A., Huehn, S., 2015. The emergence of Vibrio pathogens in Europe:
ecology, evolution, and pathogenesis (Paris, 11-12th March 2015). Front. Microbiol. 6, 830.
https://doi.org/10.3389/fmicb.2015.00830
Li, D., Zhang, H., Fu, L., An, X., Zhang, B., Li, Y., Chen, Z., Zheng, W., Yi, L., Zheng, T., 2014. A Novel
Algicide: Evidence of the Effect of a Fatty Acid Compound from the Marine Bacterium, Vibrio sp
BS02 on the Harmful Dinoflagellate, Alexandrium tamarense. Plos One 9, e91201.
https://doi.org/10.1371/journal.pone.0091201
Liu, P.-C., Lee, K.-K., Yii, K.-C., Kou, G.-H., Chen, S.-N., 1996. News & Notes: Isolation of Vibrio harveyi
from Diseased Kuruma Prawns Penaeus japonicus. Curr. Microbiol. 33, 129–132.
https://doi.org/10.1007/s002849900087
Lopez-Joven, C., Rolland, J.-L., Haffner, P., Caro, A., Roques, C., Carre, C., Travers, M.-A., Abadie, E.,
Laabir, M., Bonnet, D., Destoumieux-Garzon, D., 2018. Oyster Farming, Temperature, and
Plankton Influence the Dynamics of Pathogenic Vibrios in the Thau Lagoon. Front. Microbiol. 9,
2530. https://doi.org/10.3389/fmicb.2018.02530

Mandel, M.J., Stabb, E.V., Ruby, E.G., 2008. Comparative genomics-based investigation of resequencing targets in Vibrio fischeri: Focus on point miscalls and artefactual expansions. Bmc Genomics 9, 138. https://doi.org/10.1186/1471-2164-9-138

Marampouti, C., Buma, A.G.J., De Boer, M.K., 2021. Mediterranean alien harmful algal blooms: origins and impacts. Environ. Sci. Pollut. Res. 28, 3837–3851. https://doi.org/10.1007/s11356-020-10383-1

Martin, J., Fitzwater, S., 1988. Iron-Deficiency Limits Phytoplankton Growth in the Northeast Pacific Subarctic. Nature 331, 341–343. https://doi.org/10.1038/331341a0

Martin, M.O., 2002. Predatory prokaryotes: An emerging research opportunity. J. Mol. Microbiol. Biotechnol. 4, 467–477.

Mavian, C., Paisie, T.K., Alam, M.T., Browne, C., Beau De Rochars, V.M., Nembrini, S., Cash, M.N., Nelson, E.J., Azarian, T., Ali, A., Morris, J.G., Salemi, M., 2020. Toxigenic Vibrio cholerae evolution and establishment of reservoirs in aquatic ecosystems. Proc. Natl. Acad. Sci. 117, 7897–7904. https://doi.org/10.1073/pnas.1918763117

McRose, D.L., Baars, O., Seyedsayamdost, M.R., Morel, F.M.M., 2018. Quorum sensing and iron regulate a two-for-one siderophore gene cluster in *Vibrio harveyi*. Proc. Natl. Acad. Sci. 115, 7581–7586. https://doi.org/10.1073/pnas.1805791115

Miethke, M., Marahiel, M.A., 2007. Siderophore-Based Iron Acquisition and Pathogen Control. Microbiol. Mol. Biol. Rev. 71, 413–451. https://doi.org/10.1128/MMBR.00012-07

Miller, S.D., Haddock, S.H.D., Elvidge, C.D., Lee, T.F., 2005. Detection of a bioluminescent milky sea from space. Proc. Natl. Acad. Sci. U. S. A. 102, 14181–14184. https://doi.org/10.1073/pnas.0507253102

Millero, F.J., 1998. Solubility of Fe(III) in seawater. Earth Planet. Sci. Lett. 154, 323–329. https://doi.org/10.1016/S0012-821X(97)00179-9

Muhling, B.A., Jacobs, J., Stock, C.A., Gaitan, C.F., Saba, V.S., 2017. Projections of the future occurrence, distribution, and seasonality of three Vibrio species in the Chesapeake Bay under a high-emission climate change scenario. Geohealth 1, 278–296. https://doi.org/10.1002/2017GH000089

Neilands, J., 1981. Iron-Absorption and Transport in Microorganisms. Annu. Rev. Nutr. 1, 27–46. https://doi.org/10.1146/annurev.nu.01.070181.000331

Oberbeckmann, S., Fuchs, B.M., Meiners, M., Wichels, A., Wiltshire, K.H., Gerdts, G., 2012. Seasonal Dynamics and Modeling of a Vibrio Community in Coastal Waters of the North Sea. Microb. Ecol. 63, 543–551. https://doi.org/10.1007/s00248-011-9990-9

Pal, M., Yesankar, P.J., Dwivedi, A., Qureshi, A., 2020. Biotic control of harmful algal blooms (HABs): A brief review. J. Environ. Manage. 268, 110687. https://doi.org/10.1016/j.jenvman.2020.110687

Papenfort, K., Bassler, B.L., 2016. Quorum sensing signal-response systems in Gram-negative bacteria. Nat. Rev. Microbiol. 14, 576–588. https://doi.org/10.1038/nrmicro.2016.89

Perez, J., Jimenez-Zurdo, J.I., Martinez-Abarca, F., Millan, V., Shimkets, L.J., Munoz-Dorado, J., 2014. Rhizobial galactoglucan determines the predatory pattern of Myxococcus xanthus and protects Sinorhizobium meliloti from predation. Environ. Microbiol. 16, 2341–2350. https://doi.org/10.1111/1462-2920.12477

Perez, J., Moraleda-Munoz, A., Javier Marcos-Torres, F., Munoz-Dorado, J., 2016. Bacterial predation: 75 years and counting! Environ. Microbiol. 18, 766–779. https://doi.org/10.1111/1462-2920.13171

Pfaffl, M.W., 2001. A new mathematical model for relative quantification in real-time RT-PCR. Nucleic Acids Res. 29, e45. https://doi.org/10.1093/nar/29.9.e45

Rashidan, K.K., Bird, D.F., 2001. Role of predatory bacteria in the termination of a cyanobacterial bloom. Microb. Ecol. 41, 97–105.

Rodrigues, A.M.S., Lami, R., Escoubeyrou, K., Intertaglia, L., Mazurek, C., Doberva, M., Pérez-Ferrer, P.,
Stien, D., 2022. Straightforward *N* -Acyl Homoserine Lactone Discovery and Annotation by LC–
MS/MS-based Molecular Networking. J. Proteome Res. 21, 635–642.
https://doi.org/10.1021/acs.jproteome.1c00849

Rolland, J.-L., Pelletier, K., Masseret, E., Rieuvilleneuve, F., Savar, V., Santini, A., Amzil, Z., Laabir, M.,
2012. Paralytic Toxins Accumulation and Tissue Expression of alpha-Amylase and Lipase Genes in
the Pacific Oyster Crassostrea gigas Fed with the Neurotoxic Dinoflagellate Alexandrium
catenella. Mar. Drugs 10, 2519–2534. https://doi.org/10.3390/md10112519

Romano, A., Trimble, L., Hobusch, A.R., Schroeder, K.J., Amin, S.A., Hartnett, A.D., Barker, R.A., Crumbliss,
A.L., Carrano, C.J., 2013. Regulation of iron transport related genes by boron in the marine
bacterium Marinobacter algicola DG893. Metallomics 5, 1025.
https://doi.org/10.1039/c3mt00068k

Rosales, D., Ellett, A., Jacobs, J., Ozbay, G., Parveen, S., Pitula, J., 2022. Investigating the Relationship
between Nitrate, Total Dissolved Nitrogen, and Phosphate with Abundance of Pathogenic Vibrios
and Harmful Algal Blooms in Rehoboth Bay, Delaware. Appl. Environ. Microbiol. 88, e00356-22.
https://doi.org/10.1128/aem.00356-22

Sinclair, A.R.E., Mduma, S., Brashares, J.S., 2003. Patterns of predation in a diverse predator-prey system.
Nature 425, 288–290. https://doi.org/10.1038/nature01934

Su, R.Q., Yang, X.R., Zheng, T.L., Tian, Y., Jiao, N.Z., Cai, L.Z., Hong, H.S., 2007. Isolation and
characterization of a marine algicidal bacterium against the toxic dinoflagellate Alexandrium
tamarense. Harmful Algae 6, 799–810. https://doi.org/10.1016/j.hal.2007.04.004

Tanabe, T., Funahashi, T., Nakao, H., Miyoshi, S.-I., Shinoda, S., Yamamoto, S., 2003. Identification and
Characterization of Genes Required for Biosynthesis and Transport of the Siderophore
Vibrioferrin in *Vibrio parahaemolyticus*. J. Bacteriol. 185, 6938–6949.
https://doi.org/10.1128/JB.185.23.6938-6949.2003

Thompson, F.L., Thompson, C.C., Swings, J., 2003. Vibrio tasmaniensis sp. nov., isolated from Atlantic
Salmon (Salmo salar L.). Syst. Appl. Microbiol. 26, 65–69.
https://doi.org/10.1078/072320203322337326

Tourneroche, A., Lami, R., Hubas, C., Blanchet, E., Vallet, M., Escoubeyrou, K., Paris, A., Prado, S., 2019.
Bacterial–Fungal Interactions in the Kelp Endomicrobiota Drive Autoinducer-2 Quorum Sensing.
Front. Microbiol. 10, 1693. https://doi.org/10.3389/fmicb.2019.01693

Vanhove, A.S., Rubio, T.P., Nguyen, A.N., Lemire, A., Roche, D., Nicod, J., Vergnes, A., Poirier, A.C.,
Disconzi, E., Bachere, E., Le Roux, F., Jacq, A., Charriere, G.M., Destoumieux-Garzon, D., 2016.
Copper homeostasis at the host vibrio interface: lessons from intracellular vibrio transcriptomics.
Environ. Microbiol. 18, 875–888. https://doi.org/10.1111/1462-2920.13083

Wang, X., Li, Z., Su, J., Tian, Y., Ning, X., Hong, H., Zheng, T., 2010. Lysis of a red-tide causing alga,
Alexandrium tamarense, caused by bacteria from its phycosphere. Biol. Control 52, 123–130.
https://doi.org/10.1016/j.biocontrol.2009.10.004

Wang, Y., Li, S., Liu, G., Li, X., Yang, Q., Xu, Y., Hu, Z., Chen, C.-Y., Chang, J.-S., 2020. Continuous
production of algicidal compounds against Akashiwo sanguinea via a Vibrio sp. co-culture.
Bioresour. Technol. 295, 122246. https://doi.org/10.1016/j.biortech.2019.122246

Weerasinghe, A.J., Amin, S.A., Barker, R.A., Othman, T., Romano, A.N., Parker Siburt, C.J., Tisnado, J.,
Lambert, L.A., Huxford, T., Carrano, C.J., Crumbliss, A.L., 2013. Borate as a Synergistic Anion for
Marinobacter algicola Ferric Binding Protein, FbpA: A Role for Boron in Iron Transport in Marine
Life. J. Am. Chem. Soc. 135, 14504–14507. https://doi.org/10.1021/ja406609s

Wu, J., Luther, G., 1994. Size-Fractionated Iron Concentrations in the Water Column of the Western
North-Atlantic Ocean. Limnol. Oceanogr. 39, 1119–1129.
https://doi.org/10.4319/lo.1994.39.5.1119
Xu, Q., Wang, P., Huangleng, J., Su, H., Chen, P., Chen, X., Zhao, H., Kang, Z., Tang, J., Jiang, G., Li, Z., Zou,
S., Dong, K., Huang, Y., Li, N., 2022. Co-occurrence of chromophytic phytoplankton and the Vibrio
community during Phaeocystis globosa blooms in the Beibu Gulf. Sci. Total Environ. 805, 150303.
https://doi.org/10.1016/j.scitotenv.2021.150303
Yang, Q., Feng, Q., Zhang, B., Gao, J., Sheng, Z., Xue, Q., Zhang, X., 2021. Marinobacter alexandrii sp. nov.,
a novel yellow-pigmented and algae growth-promoting bacterium isolated from marine
phycosphere microbiota. Antonie Van Leeuwenhoek Int. J. Gen. Mol. Microbiol. 114, 709–718.
https://doi.org/10.1007/s10482-021-01551-5
763

ACKNOWLEDGMENT

The authors thank F. Le Roux and Y. Labruche for their gift of *Vibrio* mutants; E. Garcés, S. Turki, H. Frehi, A. Bouquet and W. Medhioub for their contribution in obtaining Mediterranean phytoplankton strains; K. Escoubeyrou and D. Stien for advice on HRMS/MS data; V. Diakou-Verdin and E. Jublanc from the Montpellier RIO Imaging Platform for access to microscope facilities (www.mri.cnrs.fr); P. Clair from the Montpellier qPHD platform and the Technoviv platform of the UPVD for access to PCR facilities; N. Brunet, A. Lang, A. Payelleville, T. Milhau and M. Leroy for their technical assistance; The IFREMER LER-LR laboratory for access to the Thau Lagoon; the BIO2MAR platform at the Observatoire Océanologique de Banyuls for providing access to UHPLC-HRMS/MS facilities.

FUNDING

European VIVALDI grant 678589, French EC2CO ROSEOCOM grant and IFREMER ALGOVIR grant.

Figure 1. Dynamics of *Alexandrium* and *Vibrio* in the environment. (A) Location of the monitoring
station in the Thau Lagoon (southern France). **(B)** Mean abundance (DNA equiv.) of *Vibrio spp.* (16S) and
Alexandrium *spp. (A. pacificum* ACT03 *+ A. tamarense* ATT07). *Vibrio* cells (black line with diamond
dot) and degraded *Alexandrium* cells (grey line with round dot) were evidence in the 0.2–0.8 µm fraction
(free *Vibrio* fraction) in spring and autumn 2015. Living *Alexandrium* cells (grey line with roundot) but no
plankton-associated *Vibrio spp.* (black line with diamond dot) were evidence in the 0.8–180 µm in spring
and autumn. **(C)** Result of Akaike information criterion (AICc) models tested to explain the mean value of
degraded Alexandrium cells (dead cells) in spring. **(D)** Wald test of the AICc model attributing the mean
value of degraded cells of Alexandrium in spring to free *Vibrio*.

Figure 2. Incubation of *Vibrio atlanticus* LGP32 and *Alexandrium pacificum* ACT03 in enriched natural seawater (ENSW). (A) *A. pacificum* ACT03 cultured alone (grey bar) and incubated with *V. atlanticus* LGP32 (black bar) in ENSW. **(B)** *V. atlanticus* LGP32 cultured alone (grey bar) and incubated with *A. pacificum* ACT03 (black bar) in ENSW. **(C)** Images of the interaction of *V. atlanticus* LGP32-GFP with *A. pacificum* ACT03 taken at 8h00 and 28h00 of co-culture. *V. atlanticus* LGP32 (small green cells), living *A. pacificum* ACT03 cells (large red cells) and degraded *A. pacificum* ACT03 cells (large green cells). $*P < 0.01$ (analysis by pairs).

A₁ **B₁** **C₁**

Figure 3. Role of *Vibrio atlanticus* LGP32 starvation in the interspecific interaction process.
Experiments were conducted by incubating *A. pacificum* ACT03 with *V. atlanticus* LGP32 previously
grown for 12, 36, 60 and 126 h in Zobell medium. **(A)** Percentage of motile *A. pacificum* ACT03. **(B)** *A.
pacificum* ACT03 group-attacked by *V. atlanticus* LGP32 and **(C)** *A. pacificum* ACT03 lysis after 0, 15,
30, 45 and 60 min of interaction. Corresponding pictures showing **(A1)** Black arrows indicate unhooked
and degrade flagellum from *A. pacificum* ACT03 flagellum, **(B1)** Image of the interaction of *V. atlanticus*
LGP32-GFP with *A. pacificum* ACT03 taken at 8h00 of co-culture. *V. atlanticus* LGP32 (small green
cells), living *A. pacificum* ACT03 (large red cells) and dead *A. pacificum* ACT03 (large green cell). **(C1)**
Black arrow 1 indicate vesicle formation on *A. pacificum* ACT03 cell and black arrow 2 indicate exploded
A. pacificum ACT03 cell. ***P <0.001.

Figure 4. Role of quorum sensing and the vibrioferrin iron uptake pathway in the interaction process. **(A)** Effect of *V. atlanticus* LGP32 cell density on the predation process. Experiments examined the impact of *V. atlanticus* LGP32 cell density on its ability to attack *A. pacificum* ACT03. *A. pacificum* ACT03 cells (2.10^4 cells) were incubated with *V. atlanticus* LGP32 grown for 60 hours in Zobell medium at concentrations ranging from 5×10^3 to 5×10^5 cells mL^{-1} (represented by black bars). For comparison, *A. pacificum* ACT03 was also incubated with *V. atlanticus* LGP32 grown for only 12 hours in Zobell medium at concentrations ranging from 5×10^3 to 4×10^6 cells mL-1 (represented by grey bars). The term "Att" indicates cells of *A. pacificum* ACT03 that were subjected to group attacks by *V. atlanticus* LGP32. **(B)** CqsS, luxM, luxN, luxS, and luxP quorum sensing and PvsA, PvuB and PvuA2 vibrioferrin pathway genes expression in *V. atlanticus* LGP32 grown for 12, 36 and 60 h in Zobell medium. **(C)** Effect of *V. atlanticus* LGP32 mutants on the group-attacked process. Experiments were conducted by incubating *A. pacificum* ACT03 with *V. atlanticus* LGP32, *V. atlanticus* LGP32 tagged with GFP, *V. atlanticus* LGP32 washed with ENSW or *V. atlanticus* LGP32 mutant ΔPvuB, ΔluxM, ΔluxR and ΔluxS previously grown 60 h in Zobell media (control), The percentage of *A. pacificum* ACT03 group-attacked was determined during the first 30 min of exposure. **(D)** Effect of *V. atlanticus* LGP32 cultures media composition on the group-attacked process. Experiments were conducted by incubating *A. pacificum* ACT03 with *V. atlanticus* LGP32 grown 60 h in Zobell media supplemented with H_3BO_4 or $FeCl_3$. The results were compared with an exposure to *V. atlanticus* LGP32 grown 60 h in Zobell media. **P <0.01, ***P <0.001

Figure 5. Schematic representation of a putative strategy developed by *Vibrio spp*. to feed on Alexandrium *spp*. and *G. catenatum* in the environment. (A) *Vibrio* in the environment when subjected to starvation secrete non-protein lytic compounds. **(B)** Some of these lytic compounds degrade the flagella, immobilizing the alga (immobilization stage). **(C)** Then *Vibrio* swims and clusters around its prey coordinating group attacks (group-attack stage). **(D)** Lytic compounds released by *Vibrio* where able to concentrate around the algae cells, thereby lysing the algae (killing stage). **(E)** Feeding on the released nutrients, *Vibrio* multiply and then spread in the environment. Yellow clouds: Lytic compound release by *Vibrio,* Grey clouds: Algal nutrients released upon lysis.

Figure S1.Time and dose-dependent effects of the *V. atlanticus* LGP32 culture supernatant on *A.*

***pacificum* ACT03 motility. (A)** A time dependence experiment was conducted by incubating *A.*

pacificum ACT03 for 1, 5 or 20 h with 1/1000 v/v (1 µL/mL) of culture supernatant from *V. atlanticus*

LGP32 previously grown for 60 h in Zobell culture media. **(B)** A dose dependence experiment was

conducted by incubating *A. pacificum* ACT03 for 1 h with 1/1000 to 1/20 v/v (1-50 µL/mL) of culture

supernatant from *V. atlanticus* previously grown for 60 h in Zobell media. The percentage of motile *A.*

pacificum ACT03 was determined after 1 hours of exposure.*P < 0.05, ***P < 0.001

Figure S2. *Vibrio atlanticus* **LGP32 proteome analysis following nutrient stress. (A)** Example of 2D
gel, the numbers in white on the gel 4-7 correspond to the number and position of the protein spots
analyzed. **(B)** Proteins identified by LC-MS/MS as differentially represented in the 2D gel comparative
approach following nutrient stress. ND: Not determined; ENSW (artificial seawater)**.**
854

Figure S3. Quorum sensing and the vibrioferrin iron uptake pathway in *Vibrio*. (A) Putative quorum sensing (QS) pathways at low and high cell density in *Vibrio* according to Lami et al.(Lami, 2019). **(B)** Genetic organization of the vibrioferrin utilization gene cluster on *V. atlanticus* LGP32 chromosome 2. The Pvu and Pvs operons are involved in the secretion and the transport of ferric vibrioferrin and biosynthesis of vibrioferrin, respectively. Arrows indicate the transcriptional directions of the genes. VSII1126, VSII1137 and VSII1129 corresponding to PvuB, PvuA2 and PvsA genes respectively.

866 **Table 1.** Ability of Vibrio *spp*. and strains to secrete algicidal compounds and attack *Alexandrium*
867 *pacificum* LGP32 in groups. NT: not determined.

Vibrio species	Strains	Virulence for fish or invertebrates	References	Secretion of algicidal compounds (This study)	Group attack (This study)
Vibrio atlanticus LGP32	WT	+	(Gay et al., 2004)	+	+
//	WT + pSW3654T-GFP	+	(Le Roux et al., 2007)	+	+
//	ΔLuxM	ND	Ifremer Institute, France	+	+
//	ΔLuxS	ND	//	+	+
//	ΔLuxR	ND	//	+	+
//	ΔPvuB	ND	This work	+	-
Vibrio tasmaniensis	J5-9	+	(Lemire et al., 2015)	+	+
//	LMG20012$^{\text{T}}$	-	(Thompson et al., 2003)	+	+
Vibrio crassostreae	J2.9	+	(Lemire et al., 2015)	+	+
//	J2-8	-	//	+	+
Vibrio fischeri	ES114	ND	(Mandel et al., 2008)	+	+
Vibrio harveyi	ATCC14126	+	(Liu et al., 1996)	+	+
Vibrio aestuarianus	janv-32	+	(Labreuche et al., 2010)	+	+

869 **Table 2.** Ability of *Vibrio Atlanticus* LGP32 to degrade flagella, attack in group and lysed the targeted
870 dinoflagellates *spp*. commonly found in the Mediterranean Sea. ND not determined.

Dinoflagellates species	Strains	Toxicity for human	References	Degraded flagella (This study)	Attacked in groups (This study)	Lysed cells (This study)
Alexandrium pacificum	ACT03, Thau, France	+	(Laabir et al., 2011)	+	+	+
Alexandrium catenella	Bizerte, Tunisia	+	(Fertouna-Bellakhal et al., 2015)	+	+	+
//	F3-9F, Tarragona, Spain	ND	//	+	+	+
//	C10-5, Annaba, Algeria	+	(Hadjadji et al., 2020)	+	+	+
Alexandrium tamarense	ATT07, Thau, France	-	(Rolland et al., 2012)	+	+	+
Alexandrium spp.	Golf of Tunis, Tunisia	ND	Algal collection university of Montpellier, France	+	+	+
//	Bizerte, Tunisia	ND	//	+	+	+
//	Mediterranean coast, Morocco	ND	//	+	+	+
Prorocentrum lima	PLBZT14, Bizerte, Tunisia	+	(Ben-Gharbia et al., 2016)	-	-	-
Coolia monotis	CMBZT14, Bizerte, Tunisia	ND	//	-	-	-
Vulcanodinium rugosum	IFR-VRU-01, Ingril, France	+	(Abadie et al., 2015)	-	-	-
Karenia selliformis	Golf of Gabes, Tunisia	NT	Algal collection university of Montpellier, France	-	-	-
Scripsiella trochoidae	Mellah Lagoon, Algeria	-	//	-	-	-
Gyrodinium impudicum	Golf of Tunis, Tunisia	-	//	-	-	-
Amphidium carterae	SAMS, Scotland	NT	SAMS laboratory, Scotland	-	-	-
Gymnodinium catenatum	M'diq Bay, Morocco	+	(Leblad et al., 2020)	+	+	+

Table 3. Oligonucleotide sequences of primers used for RNA expression analysis.

Species	genes	Primers Sequences	Tm (°C)	Efficiency	References
Alexandrium pacificum (ACT03)	18S – 28S rRNA ITS region	TGATATTGTGGGCAACTGTAA AACATCTGTTAGCTCACGGAA	54		(Genovesi et al., 2011)
Alexandrium tamarense (ATT07)	18S – 28S rRNA ITS region	TGGTAATTCTTCATTGATTACAATG AACATCTGTTAGCTCACGGAA	54		//
Vibrios spp.	16S	CGGTGAAATGCGTAGAGAT TTACTAGCGATTCCGAGTTC	62		(Kitatsukamoto et al., 1993)
Vibrio atlanticus LGP32	LuxN (VS_II0260)	CACTTGCTAGTATCATCGC ATCGAGTTAGCAAGAGCAC	60	1,92	This work
//	LuxM (VS_II0261)	TCCACTTATCACAAACAGG ACTGTACTTCCATTTGTCG	60	1,91	//
//	LuxP (VS_II0355)	AAGTTCAGGATGAACCTATC CAAAGAGATACTTTGCTGAG	60	1,89	//
//	LuxS (VS_2562)	ACTCTCGAGCACCTATACG GAAGGCGTACCAATCAAGC	60	1,85	//
//	CqsS (VS_1725)	GACATCTATTGATGTTATGC TCACCCACTTCACGTAACTG	60	1,91	//
//	PvsA (VS_II0355),Vibrioferrin biosynthesis protein	CAGAGCAAGAGCTAGAACC TCGTTGAGAACCTGACGAG	59	1,91	//
//	PvuB (VS_II1126), ABC transporter vibrioferrin uptake FecB	TAGTGCAACCATGGGAATCG TAAACCGTACGTAGACGCTC	57	2,01	//
//	PvuA2 (VS_II1127), Vibrioferrin receptor FecA	GGAGCTACAAGCATTCGTTC TTCGTCATATGGTCGCTTCG	57	2,08	//
//	Housekiping gene 1, CcmC (VS_0852)	ATTGCCGCCTTTATCGGTTT CAAGCACCCCACATTGGTTT	60		(Vanhove et al., 2016)
//	Housekiping gene 2, 6-phosphofructokinase (VS_2913)	GCCGTCACTGTGGTGACCTT TGCTTCTTGCCTTTCGCAAT	60		//

VIDEOS

Video 1. Dynamics of *Vibrio atlanticus* LGP32-*Alexandrium pacificum* ACT03 interaction. GFP-tagged *V. atlanticus* (small green cells); living *A. pacificum* (large red cells); degraded *A. pacificum* (large green cells) filmed under an epifluorescence microscope.

Video 2. Second-by-second timing of *Vibrio Atlanticus* LGP32 attacking *Alexandrium pacificum* ACT03. GFP-tagged *V. atlanticus* (small green cells); *A. pacificum* living cell (large red cells) filmed under an epifluorescence microscope**.**

Video 3. Degradation and disruption of *Alexandrium pacificum* ACT03 flagella. Effect of *Vibrio* supernatant on the first stage of the interaction filmed under a confocal microscope.

Video 4. Group attacks of *Vibrio atlanticus* LGP32 on target *Alexandrium pacificum* ACT03. *Vibrio*, filmed under a confocal microscope, attacks in groups one immobilized *Alexandrium* cell then moves on to attack — still as a group — another cell without touching the other whole cells, suggesting active communication between *Vibrio* cells. *V. atlanticus* LGP32 (small cells); *A. pacificum* ACT03 (large cells).

Video 5. Vesicle formation and bursting of an *Alexandrium pacificum* ACT03. Direct effect of *Vibrio* supernatant on *Alexandrium* after 126 h of culture filmed under a confocal microscope