Single molecule imaging of RecB *in vivo* reveals dynamics of DNA Double Strand Break repair in *Escherichia coli*

Daniel Thédié[1], **Alessia Lepore**[1,2], **Lorna McLaren**[1], and **Meriem El Karoui**[1,3]

[1]Institute of Cell Biology, University of Edinburgh, Edinburgh, UK
[2]Laboratory for Optics and Biosciences, École Polytechnique, Institut Polytechnique de Paris, Palaiseau, France
[3]Ecole Normale Supérieure, Paris-Saclay, France

Repairing DNA damage is of primary importance for all living organisms. DNA double-strand breaks (DSBs) are one of the most serious types of DNA damage, as they lead to loss of genetic information and death when not repaired. In *E. coli*, they are recognised and processed by the RecBCD complex, which initiates repair by homologous recombination. Although the repair dynamics downstream of RecBCD has been well characterised, it is still unclear how long this complex stays attached to DNA and what triggers its dissociation *in vivo*. To answer these questions, we imaged RecB at the single-molecule level, and quantified its dynamic behaviour in bacterial cells exposed to ciprofloxacin, an antibiotic that induces DSBs. Our results show that RecB forms long-lived complexes with DSBs (10 seconds), and that its dissociation from DNA is an intrinsic property of the complex, that does not depend on the amount of DNA damage, nor the following steps in the repair pathway. Moreover, we show that we can use RecB binding to DSBs as a marker to estimate the rate of damage formation. This study provides a detailed quantitative insight into the interaction of RecBCD with DNA double-strand ends in *E. coli in vivo*, and into the bacterial response to DSBs induced by ciprofloxacin.

DNA repair | RecBCD | Imaging | *in vivo*
Correspondence: *meriem.elkaroui@ed.ac.uk*

Introduction

Repairing DNA damage and ensuring integrity of the genome is essential to all living organisms. Among the different types of damage, DNA double-strand breaks (DSBs) pose a serious threat to bacterial cells: when left unrepaired, they can lead to cell death, genome rearrangements or mutagenesis (1). In *E. coli*, DSBs are repaired through the RecBCD pathway. RecBCD is a heterotrimeric protein complex, which combines several enzymatic activities (2). Upon recognition of a DSB, RecBCD translocates along the DNA at a speed of ∼1.6 kb/s (3), and digests both DNA strands until it meets a specific 8-base pair sequence (5'-GCTGGTGG-3') called a Chi-site. Upon recognition of the Chi-site, RecBCD stops degrading the 3' end, leading to a 3' DNA overhang. Of note, Chi recognition by RecBCD is not systematic, and was previously reported to occur with ∼40% probability (4, 5). As RecBCD creates the 3' overhang, the RecB subunit promotes loading of the RecA protein onto single-stranded DNA (ss-DNA) (6, 7). The resulting RecA-DNA nucleoprotein filament is used as a template to search for an intact homologous

sequence and perform homologous recombination (8, 9). It has been suggested that RecBCD dissociates from DNA by disassembly of the three subunits following Chi recognition (10); however, it is still unclear what specifically triggers the dissociation, and how long RecBCD stays bound to DNA before dissociating.

A common source of DSBs in *E. coli* is the collision of the replication fork with DNA-bound proteins, which leads to the disassembly of the replisome (11). It has been reported that per cell cycle, ∼18% of cells showed signs of endogenous DNA damage, most likely from replication fork breakage (12). Replisome disassembly results in a Y-shaped DNA structure that is bound by the RuvAB complex, which pulls the DNA strands (in a process called "branch migration") to create a "chicken-foot" four-way DNA structure (13). The free end of this structure is a DNA double strand end, and is bound by RecBCD, which either (i) digests the whole DNA tail and displaces RuvAB, allowing the replisome to be reloaded in a PriA-dependent manner (13), or (ii) recognises a Chi-site and loads the RecA protein onto the 3' end of the DNA tail, leading to SOS induction (14). Since this part of the DNA was recently replicated, a homologous sequence is likely to be present in close proximity, and can be used for homologous recombination.

E. coli cells might also experience DSBs from exogenous sources, such as antibiotics. Ciprofloxacin is an antibiotic of the fluoroquinolones family, which causes DSBs. In *E. coli*, it does so by binding covalently to the topoisomerase II DNA gyrase (and with lower affinity to topoisomerase IV), and trapping it in a DNA-bound conformation (15). The exact mechanism through which DSBs are created is still unclear, but ciprofloxacin has been suggested to cause replication-dependent and independent DSBs (16). Since poisoned gyrase is closely bound to DNA, it is likely to cause replication fork collisions (17, 18), and the subsequent repair of a chicken foot structure. Furthermore, the ciprofloxacin-poisoned gyrase is trapped in the "open" stage of its catalytic cycle, where it holds together two disjointed DNA ends. Therefore, after gyrase removal [through a process that involves ExoVII nuclease (19)], it can be reasonably expected that a DSB is formed independently of DNA replication (20). DSB repair is the first step of a general response to DNA damage known as the SOS response (21). The formation of the RecA nucleoprotein filament triggers the auto-proteolysis

of the LexA transcriptional repressor, which results in the activation of the genes of the SOS regulon. Among those are RecA itself, inhibitors of cell division such as SulA, and the SMC (Structural Maintenance of Chromosomes)-like protein RecN. As a result, the cells elongate (22), and their DNA is compacted at the cell centre (23).

Mutations in proteins of the repair pathway have varying effects on the cell's ability to repair DSBs. Cells that are recA-deficient ($\Delta recA$) are entirely unable to repair DSBs through homologous recombination, and the prolonged action of RecBCD on the DNA results in the digestion of large parts of the chromosome (24, 25). recA-deficient cells are, however, capable of processing DSBs arising from fork reversal, thanks to RecBCD's exonuclease activity (13, 14). The $D_{1080} \to A$ point mutation in the RecB subunit of the RecBCD complex ($RecB_{1080}$) abolishes RecB's nuclease activity, as well as its ability to load the RecA protein on DNA (26, 27). The other activities of the complex (DSB recognition, DNA unwinding, Chi-recognition) are unaffected (28). Despite the loss of its RecA loading activity, the $recB_{1080}$ mutant is still able to repair DSBs. To do so, the $RecB_{1080}CD$ complex unwinds DNA at the break without degrading it, and the RecJ exonuclease degrades the 5' ssDNA. Finally, the RecFOR complex loads RecA, allowing for homologous recombination to occur (29).

Previous microscopy studies have shed light on different aspects of DSB repair by RecBCD. Several in vitro studies have quantified the rate of DNA unwinding by RecBCD (30, 31), as well as the growth rate of RecA filaments (32–34) and the mechanism of homology search (35, 36). Complimentary to the in vitro work, in vivo studies have provided insight into DNA end resection by RecBCD (3), confirming its translocation speed and high processivity. More recently, we investigated RecB mobility in vivo at the single-molecule level (37), finding that RecB's engagement in repair initiation is proportional to the level of DNA damage and that its interactions with DNA are consistent with its role in initiating the repair process. The spatial organisation of RecA during DSB repair has been extensively characterised. Studies reported the presence of RecA foci (38–41), filaments (42), or bundles of filaments (43, 44). The homology search and pairing processes were also observed in vivo, showing that DSBs are brought to the centre of the cell during the repair process (9, 45). Our previous observations (37) highlighted how the coordination of all RecBCD activities facilitated rapid and efficient repair. Nevertheless, several important aspects of RecBCD's processing of DSBs in vivo remain to be clarified: how long does RecBCD stay bound to DNA after recognising a DSB? Does the binding time change under different levels of DNA damage? Does RecA play a role in triggering RecBCD dissociation from DNA? These open questions are fundamental to the complete understanding of DSB repair in E. coli.

To address these questions, we used single-molecule imaging to quantify RecB binding to DSBs in living E. coli cells exposed to various concentrations of ciprofloxacin. We measured the duration of the RecB-DSB interaction and found that it was not affected by the ciprofloxacin concentration within the tested range (0 to 30 ng/ml). Concurrent imaging of RecA and the bacterial nucleoid showed RecA structures similar to previously observed bundles and compaction of the bacterial nucleoid, both of which scaled with ciprofloxacin concentration and duration of exposure. Furthermore, we imaged RecB in the $\Delta recA$ and $recB_{1080}$ mutants, and revealed that RecB dissociation depends only on the intrinsic biochemical activities of the complex.

Materials and Methods

Strain construction. E. coli MG1655 and derivatives were used in this study. A list of all strains and plasmids used are presented in Supp. Tables 1 and 2. MEK2623 (RecB-Halo RecA-SYFP2) was constructed by transferring the RecA-SYFP2 tandem fusion by P1 transduction, using EL2514 (9) as a donor and MEK65 (46) as a receiver. The clones were selected on Kanamycin plates, and checked by Sanger sequencing. MEK2622 ($\Delta recA$) was constructed by transferring the recA deletion by P1 transduction, using DL654 as a donor, and MEK65 as a receiver. MEK2629 was obtained by transforming the pDT6 plasmid (gift from Mark Dillingham) into the MEK65 strain by heat-shock.

Microscopy samples preparation. For all experiments, the cells were grown in M9 supplemented with 0.2% (w/v) glucose, 2 mM $MgSO_4$, 0.1 mM $CaCl_2$, 1X MEM Essential and MEM non-essential amino acids (Gibco).

Cell culture. Cells were grown overnight from a -80°C glycerol stock in 5 mL medium at 37°C, shaking at 150 rpm. In the morning, the cells were diluted 1:300 into 15 mL of medium and grown at 37°C, shaking at 150 rpm until they reached an OD_{600} of 0.2-0.3 (mid-exponential phase). Alternatively, a 15 mL culture was inoculated using a -80°C glycerol stock, and placed at 37°C with stirring in a turbidostat device (OGIBio), which measured OD_{600} and diluted cells at 3 min intervals, in order to keep them at a constant OD_{600} of 0.2.

Halo labelling. The same labelling protocol was used from a previous study (37). A volume of cells equivalent to 1 mL at OD_{600}=0.2 was centrifuged (8000 rpm) and resuspended in 1 mL of fresh medium. 5 μL of JF549 dye (Janelia Fluor Halo-tag Ligand, Promega) was added (final concentration 1 μM) and the cells were further incubated for 1h at 37°C in the dark, shaking at 150 rpm.

DNA labelling. When relevant, cellular DNA was labelled by adding Sytox Green (Invitrogen) at a concentration of 500 nM, at the same time as the JF549 dye. It was incubated as described above.

Gam overexpression. For experiments where Gam was overexpressed, arabinose was added to the cells during the labelling step, at 1% final concentration. Arabinose (at the same concentration) was also added to the medium for subsequent washes, and in the agarose-pad to maintain exposure during imaging.

Thédié *et al.* | Single molecule imaging of RecB *in vivo* reveals dynamics of DSB repair in *E. coli*

Dye removal. The cells were centrifuged for 3 min at 8000 rpm, the supernatant discarded, and the pellet resuspended in 1 mL of fresh medium and transferred to a new tube to avoid the dyes sticking to the plastic of the tube. This procedure was repeated 3 times to fully remove unbound dye. The cells were finally resuspended in 200 μL medium.

Sample preparation. Agarose-pads were prepared by dissolving 2% agarose in culture medium. In experiments with ciprofloxacin, the antibiotic was added to the agarose-pad at the desired final concentration. After the dye removal step, 5 μL of cells were added on the agarose-pad, and left to settle for 10 min at 37°C in the dark before imaging.

Microscopy.

Microscope set-up. Imaging was performed using an inverted microscope (Nikon Ti-E) equipped with a 100X TIRF Nikon objective (NA 1.49, oil immersion) and a 1.5X Nikon magnification lens (pixel size = 107 nm). Fluorescence excitation was achieved using 488- and 561-nm lasers (Coherent OBIS) in HILO (Highly Inclined Laminar Optical sheet) configuration. Excitation light and fluorescence emission were separated using a dual-wavelength dichroic filter (TRF59904, Chroma), and the fluorescence signal was detected on an EM-CCD (Electron Multiplying Charge-Coupled Device) camera (iXion Ultra 897, Andor). The hardware was controlled and images were saved using MetaMorph (Molecular Devices; v7.8.13.0). The HILO configuration was established using the iLas variable angle TIRF control window. All experiments were performed at 37°C, using an Okolab microscope cage incubator equipped with dark panels.

Data acquisition. For each sample, a Metamorph journal was used to acquire images on 40 different positions, each separated by 200 μm. The full camera sensor (512x512 pixels) was used. Acquisition parameters are described in Supp. Table 3. The prolonged imaging of the RecB-Halo fusion was only possible thanks to the exceptional photostability of the JF549 dye, which displayed slow photobleaching and no blinking (Supp. Note 1 and Supp. Fig. 1).

Data analysis.

The general data analysis workflow is described in Supp. Figure 2. In brief, after acquisition, the raw microscopy images were stored in .tif format on a dedicated Omero server (OME). They were directly accessed by the BACMMAN ImageJ plugin (47), which performed image processing tasks such as denoising of fluorescence images, cell segmentation (including manual curation of incorrectly segmented cells), nucleoid segmentation, and fluorescent spot detection. Measurement tables in CSV format were exported from BACMMAN and loaded in Jupyter Notebooks using the PyBerries package (version 0.2.25) in Python (version 3.11). The data tables were manipulated using PyBerries and the pandas library (version 2.0.2). Figures were generated in Jupyter Notebooks using the Seaborn library (version 0.13.2). Figures that contained several panels were assembled in Inkscape (version 1.3.2).

BACMMAN pipeline.

Deep-learning denoising of fluorescence images We improved a self-supervised denoising neural network by making it use the previous and next frames in the timelapse and performing deconvolution simultaneously. Our method is based on a previously developed method (48) which uses an encoder-decoder convolutional neural network that performs denoising (DNet) as well as a fully connected neural network that recovers the noise distribution used to compute the self-supervised loss. In order to include previous and next frames, we modified the architecture of DNet so that the encoder is shared between frames (t-1, t and t+1), i.e. it processes each frame independently. At each contraction level, the residual tensors corresponding to each frame are combined using a 1x1 convolution and fed to the decoder. The same applies for the feature tensors and the last level of the decoder. The decoder only predicts the central frame (t) and the same self-supervised loss is used. We observe a dramatic improvement of denoising performance, which implies that the proposed architecture benefits from using information from adjacent frames. It has been shown that performing denoising and deconvolution simultaneously dramatically improves deconvolution performance (49). We computed the experimental PSF of our system by averaging hundreds of bead images cropped in a 33x33 pixels area around each bead, and normalised so that values sum to one. The obtained averaged images were used during training as kernel for a 2D convolution at the end of the neural network. At prediction time, convolution was removed.

Cell segmentation The 16-image brightfield Z-stack was first cropped to 5 images on one side of the focus, as required by our segmentation algorithm. The 5-image stack was used as input to Talissman, a U-net-based segmentation algorithm. In brief, the U-net model predicts a Euclidean distance map, where the value of each pixel is its predicted distance to the nearest background pixel. A watershed algorithm is then applied to retrieve cell contours. This approach allowed us to accurately segment cells from bright-field images, including when they formed tight clusters. Any overlapping cells were manually removed from the analysis. Following segmentation, post-filters were applied to dilate the segmented regions slightly and to remove any cells that were in contact with the edge of the image and might, therefore, be cropped. Finally, all segmentation masks were visually inspected and curated to remove cells that were incorrectly segmented (∼1% of total cells).

Detection of RecB spots Fluorescent spots were segmented using a seeded watershed algorithm on the Laplacian transform of the denoised RecB image. The quality of the segmentation was visually assessed by overlaying the segmentation mask on the raw fluorescence images. The segmentation parameters were optimised, but no manual curation was applied to avoid introducing user bias in the results.

Classification of RecA structures We designed a U-net-based deep-learning model capable of classifying objects (in

this case, cells) based on the type of RecA structure they contained (Supp. Figure 3). In brief, the model was given as input the cell segmentation mask and the RecA-SYFP2 fluorescence image, and it provided as output a single class for each object: diffuse fluorescence; RecA focus; or RecA filament. The model's predictions were 84% accurate, as evaluated against a test set of manually labelled data that was not used in training.

Nucleoid segmentation   Sytox Green fluorescence images were processed with a rolling-ball noise subtraction (radius 6 pixels), and nucleoids segmented by a seeded watershed algorithm. The seed threshold was determined by Otsu's method (50). Any segmented regions that were in contact with each other within the same cell were merged, and segmented regions smaller than 5 pixels were excluded. Regions with weak signal-to-noise ratio (SNR < 2) were also excluded. For each segmented region, the SNR was computed as follows:

$$SNR = \frac{\overline{F} - \overline{B}}{std(B)} \tag{1}$$

with $\overline{F}$ the mean fluorescence intensity in the object, $\overline{B}$ the mean background intensity, and $std(B)$ the standard deviation of background intensities.

Measurements   As the final step of the pipeline, BACMMAN performed several measurements on the objects created (cells, RecB spots and nucleoids). These measurements included cell length, cell area, raw fluorescence intensity, background-subtracted fluorescence intensity, number of RecB spots in the cell, and position of the RecB spots and nucleoids along the long- and short-axis of the cell. For each dataset, BACMMAN produced one csv table per object.

PyBerries.

Data import and format   The PyBerries package, combined with the pandas library, allowed to easily import the multiple CSV tables produced by BACMMAN, and perform operations such as grouping, aggregations, normalisation and curve fitting.

Fitting RecB spot lifetimes   The lifetime of individual RecB spots was computed in BACMMAN. The resulting lifetime histogram was fitted with a bi-exponential decay function of the type $y = a_1.e^{-k_1.t} + a_2.e^{-k_2.t}$. To improve the reliability of the fits, a bootstrapping procedure was used. For each group, 100 resamplings with replacement were made, and fitted individually. Each fit parameter was estimated as the median of all fitted parameters.

Because we were unable to differentiate whether the disappearance of a fluorescent spot was a result of photobleaching or of unbinding and returning to the pool of diffusing molecules, we considered that the fitted "spot disappearance rates" k_1 and k_2 were a sum of RecB's dissociation rate from DNA and the dye's bleaching rate ($k_1 = k_{d1} + k_b$ and $k_2 = k_{d2} + k_b$, with k_{d1} and k_{d2} the two RecB dissociation

rates, and k_b the bleaching rate). The bleaching rate was calculated for each dataset, as described in Supp. Note 1 and Supp. Figure 1 and subtracted to retrieve the true RecB dissociation rates from DNA.

Estimation of the rate of RecB recruitment to DSBs   For each timelapse, we quantified the total number of RecB spots that were bound to DSBs. This could not be achieved by thresholding on the lifetime histograms, as a significant number of short-lived spots correspond to DSB-bound RecB molecules. To estimate the total number of DSB-bound RecB, we considered that all RecB spots that disappeared with a slower rate (as determined by the bi-exponential decay fit) were DSB-bound. We multiplied the total number of spots that appeared during the timelapse by the proportion of slow-dissociating spots retrieved from the lifetime histogram fits, which gave us an estimate of the number of DSB-bound RecB per timelapse, from which a number of recruitment events per cell per hour was calculated. This rate was corrected for photobleaching of the fluorescent dye, to account for non-fluorescent RecBCD complexes binding to DSBs.

Code availability. Instructions to install the BACMMAN ImageJ plugin can be found on the BACMMAN wiki. The PyBerries package can be found on the Python Package Index, and its source code is available on Gitlab. The deep-learning model used to classify cells according to their RecA-associated fluorescence is available on Gitlab. The Jupyter Notebooks used to make the article figures are available on Github.

Results

RecB forms long-lived spots when recruited to DSBs caused by ciprofloxacin. To quantify RecB binding time to DSBs in live *E. coli*, we used a Halo-tag fusion to the RecB subunit, conjugated to the JF549 fluorescent dye (Figure 1A). The fusion was previously fully characterised, ensuring specific one-to-one labelling of RecB molecules without adverse effects on the DNA repair process (37, 46). As a result of the large size and topological constraints of the bacterial chromosome, DSB-bound RecB diffuses very slowly (37). Therefore, we applied the previously developed technique of localisation enhancement (51, 52): since RecB is present at low copy numbers in *E. coli* [∼5 molecules per cell on average (46, 53)], imaging live cells with a long exposure time (1 second) made fast-diffusing RecB molecules appear as weak homogeneous signal in the cell, while very slow-diffusing RecB molecules formed diffraction-limited spots (referred to throughout this article as RecB spots, Figures 1B and 1C). The complete absence of similar spots in cells expressing the free Halo-tag from a plasmid confirmed that these spots were specific to RecB (Supp. Figure 4). However, their exact nature needed to be confirmed. Although DSB-bound molecules are expected to diffuse very slowly, the RecBCD-Halo complex diffuses in the crowded and constrained environment of the cytoplasm, and might undergo transient interactions with DNA (37), which could lead to

RecB spots forming independently of DSB binding. To gain more insight into the nature of RecB spots, we increased the number of DSBs by adding ciprofloxacin to agar pads before imaging, and measured the lifetime of the spots by recording 100-second timelapse videos with 2-sec interval between frames at different positions in the sample, for a total time of 60 min following exposure to ciprofloxacin (Figure 2A). The 1-second exposure time was optimised to detect RecB spots with good signal-to-noise while limiting dye photobleaching and keeping a sufficient time resolution to determine spot lifetimes. Visual inspection of the timelapse data showed that RecB spot lifetime was heterogenous, with a majority of spots being visible only for a few frames, and a small number persisting for longer times. These longer-lived spots became more frequent as the ciprofloxacin concentration increased. Inspecting background-subtracted intensity time-traces for single RecB spots showed that spots lost their intensity in a single step (Supp. Figure 5), consistent with single-molecule photobleaching or transitioning from a very slowly diffusing state to a rapid one. Therefore, our imaging setup allowed us to visualise single, very slow diffusing RecB molecules, and to measure the time they spent in this diffusive state.

Fig. 1. Detecting RecB binding to DSBs. **(A)** Scheme of our experimental protocol. The RecB subunit of the RecBCD complex is fused to a Halo-tag, bound by the JF549 fluorescent dye (37, 46). **(B)** A long exposure time (1 sec) makes diffusing molecules appear as a diffuse signal in the cell, while DSB-bound molecules are visible as bright, diffraction-limited spots. **(C)** Example image of a RecB spot (white arrow).

RecB forms long-lived complexes with DSBs. Using the time-lapse data, we built histograms of RecB spot lifetimes at the different ciprofloxacin concentrations (Figure 2B). In the absence of ciprofloxacin, the majority of spots lasted less than 2 seconds. Under increasing ciprofloxacin concentrations, the distribution showed a clear shift towards longer-lived spots, with the histogram at 20 and 30 ng/ml ciprofloxacin displaying a clear "tail" of long-lived events (> 10 sec). Fitting these histograms with a mono-exponential decay model ($y = a.e^{-k.t}$) did not match the experimental data accurately (Supp. Figure 6). In particular, it did not account for the tail of longer-lived spots that formed at high

ciprofloxacin concentrations. Fitting with a bi-exponential decay model ($y = a_1.e^{-k_1.t} + a_2.e^{-k_2.t}$) accounted better for the longer-lived RecB spots (Figure 2B). The bi-exponential fit outlined two populations of spots with different lifetimes: a short-lived one, with average lifetimes ranging from 1.4 to 1.7 sec; and a longer-lived one, with average lifetimes ranging from 10 to 14 sec (Table 1). Even though the short-lived spots always represented a majority of events (over 90% of the spots), the proportion of long-lived spots increased under higher ciprofloxacin exposure (from 1.4 ± 0.3% at 3 ng/ml ciprofloxacin to 5.5 ± 1.4% at 30 ng/ml ciprofloxacin).

Table 1. Parameters derived from the spot lifetime histogram fits (Figures 2B and 2C). The lifetime was calculated as the inverse of the fitted dissociation rate. Values are given as the median ± standard deviation over at least 3 independent datasets. N_{cells} = 66,764. N_{spots} = 170,138

Strain	Cipro.	Type	Lifetime (sec)	Population (%)
WT	0 ng/ml	Short	1.4 ± 0.2	98.2 ± 1.0
		Long	9.5 ± 9.7	1.8 ± 1.0
	3 ng/ml	Short	1.5 ± 0.1	98.6 ± 0.3
		Long	10.5 ± 1.3	1.4 ± 0.3
	10 ng/ml	Short	1.7 ± 0.2	97.6 ± 0.6
		Long	13.7 ± 2.3	2.4 ± 0.6
	20 ng/ml	Short	1.6 ± 0.2	96.6 ± 0.8
		Long	11.8 ± 2.1	3.4 ± 0.8
	30 ng/ml	Short	1.7 ± 0.2	94.5 ± 1.4
		Long	13.3 ± 2.2	5.5 ± 1.4
Gam	0 ng/ml	Short	1.7 ± 0.2	99.2 ± 0.4
		Long	15.5 ± 9.1	0.8 ± 0.4
	30 ng/ml	Short	1.5 ± 0.1	98.2 ± 0.9
		Long	8.2 ± 1.8	1.8 ± 0.9

To determine whether short- or long-lived spots resulted from RecB binding to DSBs, we measured RecB spot lifetime in the presence and absence of ciprofloxacin while over-expressing the Gam protein of phage λ from a plasmid. The Gam protein was previously shown to bind to the RecBCD complex instead of DNA double-strand ends (54), and its overexpression is therefore expected to abolish RecBCD binding to DSBs. Accordingly, cells that overexpressed Gam and were exposed to high ciprofloxacin (30 ng/ml) showed little elongation compared to cells that did not overexpress Gam (Supp. Figure 7), indicating that most cells did not induce the SOS response, as a result of the inability of the RecBCD-Gam complex to bind to DSBs and load RecA. The resulting RecB spot lifetimes showed a similar distribution to the one obtained in the absence of ciprofloxacin (Supp. Figure 8). This was confirmed by fitting the histogram with our bi-exponential decay model (Figure 2C), which found a proportion of long-lived spots in the presence of 30 ng/ml ciprofloxacin (1.8 ± 0.9%, Table 1) equivalent to wild-type cells that were not exposed to ciprofloxacin (1.8 ± 1%). The residual amount of long-lived RecB spots in the presence of Gam could be a result of residual DSB binding despite the

Fig. 2. RecB spot lifetimes under ciprofloxacin exposure. **(A)** Example kymographs (2-sec interval, 100 sec total) of cells containing short- and long-lived RecB spots. **(B)** RecB spot lifetime histograms at 0, 3, 10, 20 and 30 ng/ml ciprofloxacin (bars), fitted with a bi-exponential decay model (black line, fit components shown as dashed lines). N_{cells} = 66,764. N_{spots} = 170,138. **(C)** RecB spot lifetime histograms (bars) for cells over-expressing the Gam protein, fitted with a bi-exponential decay model (black line, fit components shown as dashed lines). N_{cells} = 8,812. N_{spots} = 18,698. **(D)** Probability of a spot corresponding to a DSB-bound RecB molecule. Black dots show averages for individual datasets; the black line is the average between them, and the red dashed line shows the smallest lifetime at which RecB spots have a 95% probability of being DSB-bound. N_{cells} = 8,812. N_{spots} = 18,698.

presence of Gam. This residual binding could also explain the small increase in cell length observed in Gam-expressing cells under 30 ng/ml ciprofloxacin (Supp. Figure 7). Since Gam overexpression prevents RecB binding to DSBs, and specifically caused long-lived spots to disappear, we conclude that long-lived spots correspond to RecB molecules bound to a DSB (and we will refer to them as DSB-bound RecB throughout this article). Given that under Gam overexpression, 95% of RecB spots had a lifetime shorter than 10 sec (Figure 2D), we considered any spot with a lifetime over 10 sec to be a DSB-bound RecB molecule. The appearance of short-lived spots in the presence of Gam is likely a result of a combination of short-lived DNA interactions and slow, confined diffusion of the RecBCD-Gam-Halo complex (Supp. Note 2 and Supp. Figure 9). Therefore, spots that are visible for less than 10 seconds cannot be reliably assigned as either DNA-interacting or not.

Using the Gam protein allowed us to identify DSB-bound RecB molecules as the longer-lived population in our bi-exponential fits of the RecB spot lifetime histograms. We determined that when it processes a DSB, RecB stays bound to it for 10 to 14 seconds on average, regardless of the ciprofloxacin concentration (Table 1). This is consistent with

processing of DSBs by RecBCD taking place independently of the number of DSBs in the cell. In addition, to test whether the duration of ciprofloxacin exposure affected spot lifetimes, we performed separate fits at 15 min intervals for cells exposed to 30 ng/ml ciprofloxacin (Supp. Figure 10). The RecB spot lifetimes obtained were consistent with those reported in Table 1 (10-15 sec for long-lived spots, and 1.5-2 sec for short-lived spots), indicating that RecB binding was not affected by the duration of ciprofloxacin exposure.

RecB dissociation time is independent of RecA loading. Since the duration of the RecB-DSB interaction was unaffected by the amount of DNA damage, we wondered if it was an intrinsic property of RecBCD, or if dissociation of the complex was triggered by the following step in the repair process, the loading of RecA on DNA. To assess this, we used the Δ*recA* mutant, which lacks the RecA protein, and the *recB*₁₀₈₀ mutant, where RecB has lost its RecA loading activity. We imaged both strains in the presence and absence of ciprofloxacin (30 ng/ml), computed histograms of RecB spot lifetimes and fitted them with a bi-exponential decay model (Supp. Figure 11), from which we extracted the binding times of RecB on DSBs (Table 2). In the Δ*recA* mu-

tant, the DSB-binding lifetime of RecB was similar to the wild-type (~12 sec), both in the presence and absence of ciprofloxacin. This suggests that RecA loading by RecBCD does not play a role in triggering the dissociation of wild-type RecBCD from DNA. In contrast, $RecB_{1080}$ stayed bound to DSBs for longer on average than wild-type RecB (~16 to 19 sec). This could be a result of the inability of $RecB_{1080}$ to digest the DNA it unwinds, making dissociation more challenging because of the DNA strands threading through the RecBCD complex.

Table 2. Parameters derived from the spot lifetime histogram fits for the $\Delta recA$ and $recB_{1080}$ strains (Supp. Figure 11). The lifetime was calculated as the inverse of the fitted dissociation rate. Values are given as the median $\pm$ standard deviation over at least 3 independent datasets. N_{cells} = 56,131. N_{spots} = 177,646.

Strain	Cipro.	Type	Lifetime (sec)	Population (%)
$\Delta recA$	0 ng/ml	Short	1.8 ± 0.2	96.6 ± 1.0
		Long	13.0 ± 1.8	3.5 ± 1.0
	30 ng/ml	Short	2.1 ± 0.2	90.5 ± 3.4
		Long	12.4 ± 3.5	9.5 ± 3.4
$recB_{1080}$	0 ng/ml	Short	1.8 ± 0.2	97.4 ± 0.9
		Long	16.7 ± 5.5	2.6 ± 0.9
	30 ng/ml	Short	2.1 ± 0.1	96.5 ± 0.7
		Long	19.4 ± 7.1	3.5 ± 0.7

RecB binding can be used to estimate the rate of DSB formation. Whereas the lifetime of RecB spots informs us on how long RecB stays bound to DSBs, the rate of appearance of spots can be used to estimate the rate of recruitment of RecB to DSBs under each DNA damage condition. To do so, we used the RecB spot lifetime histogram fits to estimate the total number of slow-dissociating spots per time-lapse. Since we determined that the appearance of a slow-dissociating spot corresponds to the binding of RecB to a DSB, we could calculate the number of RecB recruitments to DSBs per cell per hour (Figure 3A). We estimated that 1.3 $\pm$ 1.1 RecB molecules are recruited to DSBs per hour in the absence of ciprofloxacin. As expected, the recruitment rate of RecB scaled with ciprofloxacin concentration, up to 8.2 $\pm$ 3.6 RecB recruited to DSBs per hour at 30 ng/ml. It has been reported that DSB repair from the point of RecBCD binding takes ~15 min in *E. coli* (9); therefore, it appears consistent that undergoing 8 DSBs per hour at 30 ng/ml ciprofloxacin (a concentration higher than the Minimum Inhibitory Concentration, MIC) will cause multiple DSBs to be repaired simultaneously, and be lethal to most cells.

Surprisingly, the rates of recruitment of RecB to DSBs in the $\Delta recA$ and $recB_{1080}$ mutants were higher than in the wild-type, both in the absence and in the presence of ciprofloxacin (Figure 3B). In both cases, the rate of DSB formation is expected to be the same between the wild-type and the mutants. Therefore, we hypothesise that this higher rate of RecB recruitment to DSBs is a result of multiple recruitment events on the same original DSB that cannot be repaired, as was previously reported in the case of the $\Delta recA$ mutant (55, 56). In the $recB_{1080}$ mutant, RecB recruitment is higher than in WT cells, but equivalent (in the absence of ciprofloxacin) or lower (in the presence of 30 ng/ml ciprofloxacin) to the level of RecB recruitment in the $\Delta recA$ mutant (Figure 3B). This might reflect the ability of the $recB_{1080}$ mutant to repair DSBs in a RecA-dependent manner, albeit with lower efficiency than wild-type cells (see Discussion).

Ciprofloxacin exposure induces formation of RecA filaments. During DSB processing, RecBCD facilitates the loading of the RecA protein on ssDNA. To broaden our view of the repair process downstream of RecBCD, we imaged a tandem fusion of RecA with the fluorescent protein SYFP2, which was previously shown to preserve RecA functionality (9). Based on the fluorescence distribution in the cells (Supp. Figure 12A), we identified three states of RecA: diffuse; forming a bright focus; or forming an elongated structure (called "filament" thereafter), which matches previous *in vivo* observations of RecA (9). Because of the large diversity of shapes observed, particularly for RecA filaments, detecting RecA structures using rule-based algorithms was challenging. Therefore, we designed and trained a deep-learning algorithm capable of classifying individual cells based on the three types of spatial distributions cited above (Supp. Figures 3 and 12B). In cells not exposed to ciprofloxacin, the RecA-associated fluorescence was mostly diffuse, and occasionally formed a bright focus or filament. This is consistent with the expectation that RecA diffuses freely in the

Fig. 3. Recruitment of RecB to DSBs. **(A)** Recruitment rate of RecB to DSBs per cell at different ciprofloxacin concentrations. Black points show averages for individual datasets, and bars the median value between them. N_{cells} = 66,764. N_{spots} = 170,138. **(B)** Recruitment rate of RecB to DSBs per cell for the $\Delta recA$ and $recB_{1080}$ mutants. Black points show averages for individual datasets, and bars the median value between them. N_{cells} = 23,540. N_{spots} = 99,891.

cell without DNA damage, and polymerises on DNA in the event of an endogenous DSB. After one hour of exposure, the proportion of cells that contained a RecA filament had increased to ~40% at 20 ng/ml ciprofloxacin, and ~60% at 30 ng/ml. RecA foci, on the other hand, formed quickly following ciprofloxacin exposure (present in ~40% of the cells after 15 min of exposure to 30 ng/ml ciprofloxacin) and came back to endogenous damage level after ~1 hour. Taken together, these results show that RecA foci are transient structures in the repair process that do not accumulate under high DNA damage, whereas RecA filaments accumulate under constant exposure to ciprofloxacin (see Discussion).

DSB-bound RecB colocalises with the bacterial nucleoid. RecA loading on ssDNA triggers the SOS response, which inhibits cell division, leading to cell filamentation. Previous studies have reported that the SOS response also triggers compaction of the bacterial nucleoid (23). To see if this was the case for breaks induced by ciprofloxacin in our experimental conditions, we stained DNA using the Sytox Green dye. The use of a green dye allowed us to image RecB concomitantly, and to correlate the position of DSB-bound RecB molecules (defined as RecB spots with a lifetime >10 sec, see Figure 2D) with that of the nucleoid. In cells that were not exposed to ciprofloxacin, the nucleoid was often observed to be bi-lobed (Figure 4A), consistently with our previous observations (37). After 60 min of exposure to ciprofloxacin, the cells appeared elongated, and the nucleoid was often compacted in the centre of the cell. As a result of nucleoid compaction, the total fraction of the cell occupied by the nucleoid decreased, from 40% on average in untreated cells to 31% in cells exposed to 30 ng/ml of ciprofloxacin for an hour (Supp. Figure 13). In addition to the compaction, we observed that nucleoids were increasingly centred, and the cells elongated upon increasing exposure to ciprofloxacin, both in time and concentration (Supp. Figure 14).

Long-lived RecB spots were found close to the nucleoid (Figures 4A and 4B). This proximity was also observed after different durations of ciprofloxacin exposure (Supp. Figure 15). This observation strengthens our interpretation that long-lived spots are DSB-bound RecB molecules. Upon exposure to increasing ciprofloxacin concentrations, induction of the SOS response caused the cells to elongate significantly (from 3.1 ± 0.19 µm on average in the absence of ciprofloxacin to 5.5 ± 0.71 µm after 75 min of exposure to 30 ng/ml ciprofloxacin, Supp. Figure 7). Despite this, and because of compaction and centring, the nucleoid remained within ~2 µm either side of the cell centre. The distribution of DSB-bound RecB molecules overlapped strongly with nucleoid density (Figure 4B), indicating that as the nucleoid was compacted, newly formed DSBs triggered RecB recruitment at the centre of the cell, where the nucleoid was located.

Nucleoid compaction requires RecA loading. The spatial distribution of nucleoid density and DSB-bound RecB was affected in the $\Delta recA$ and $recB_{1080}$ mutants (Figures 4C and 4D). In the $\Delta recA$ mutant, exposure to ciprofloxacin did not trigger nucleoid compaction and centring as in wild-type cells. This confirmed that nucleoid compaction under ciprofloxacin exposure requires RecA loading, and most likely the induction of the SOS response. Despite the absence of compaction, the nucleoid occupied a smaller fraction of the cell in the $\Delta recA$ mutant after exposure to ciprofloxacin (Supp. Figure 16). This can be attributed to the progressive degradation of the bacterial chromosome by the repeated cycles of RecBCD binding (Figure 3B).

In the $recB_{1080}$ mutant, the nucleoid often formed irregular shapes and small segregated regions in the cell (Figure 4C). This disorganisation was especially visible in the presence of 30 ng/ml ciprofloxacin. Furthermore, in the $recB_{1080}$ mutant the nucleoid did not undergo compaction within the 75 min of our experiment (Supp. Figure 16). This might reflect the less efficient RecA loading in this mutant, leading to different time dynamics of the SOS induction and either impaired or significantly delayed nucleoid compaction.

In both mutant strains, the localisation of DSB-bound RecB overlapped with the bacterial nucleoid (Figure 4D), similarly to the wild-type. In the $\Delta recA$ mutant, addition of 30 ng/ml ciprofloxacin did not change the spatial distribution of nucleoid density or DSB-bound RecB. This result was expected, as $\Delta recA$ cells were unable to induce the SOS response, and, therefore, did not undergo nucleoid compaction. In the $recB_{1080}$ mutant in the absence of ciprofloxacin, both nucleoid density and DSB-bound RecB distributions were more centred than in wild-type cells, despite the nucleoid occupying the same fraction of the cell area (Supp. Figure 16). This might be because of the slower growth rate of the $RecB_{1080}$ mutant resulting in a larger proportion of cells having a single centred nucleoid, rather than a bi-lobed nucleoid as seen in the wild-type and the $\Delta recA$ mutant. Upon exposure of the $recB_{1080}$ mutant to ciprofloxacin, the absence of change in the distribution of nucleoid density and DSB-bound RecB position reinforced the hypothesis that nucleoid compaction is impaired or delayed in this mutant.

Discussion

RecB dissociation does not depend on subsequent repair steps. In this work, we used imaging of RecB bound to DSBs to elucidate key aspects of RecBCD's processing of DSBs in vivo, focusing on how long RecBCD remains bound to DNA after DSB recognition and whether this binding time depends on the level of DNA damage. We found that in wild-type *E. coli*, RecB stays bound to DSBs for 10 to 15 seconds on average (Table 1), independently of the concentration of ciprofloxacin. Additionally, we investigated whether RecA might play a role in promoting RecBCD dissociation from DNA. Imaging RecB in a $\Delta recA$ mutant indicated that this is not the case, and that RecB dissociation occurs independently of RecA loading (Table 2). Interestingly, the $recB_{1080}$ mutant had a reduced dissociation rate compared to wild-type cells, which highlights the importance of RecB's exonuclease activity in the process and the possibility that digestion of the unwound DNA strands is a pre-requisite for RecBCD dissociation.

Nucleoid - RecB

Fig. 4. Colocalisation of RecB spots with the bacterial nucleoid. **(A)** Representative images of cells (segmented outline in grey) showing the nucleoid (green) and RecB-associated fluorescence (magenta). RecB spots (indicated by arrows) are located in close proximity to the nucleoid. **(B)** Overlay of nucleoid density (green area) and position of DSB-bound RecB molecules (magenta bars) along the cell's long axis, for different ciprofloxacin concentrations (0 to 30 ng/ml). N_{cells} = 24,014. N_{spots} = 79,969. $N_{nucleoids}$ = 31,441. **(C)** Representative overlay images of RecB-associated fluorescence (magenta) and the bacterial nucleoid (green) in the $\Delta recA$ and $recB_{1080}$ mutants. Segmented cell outline shown in grey. **(D)** Overlay of nucleoid density (green area) and position of DSB-bound RecB molecules (magenta bars) along the cell's long axis for the $\Delta recA$ and $recB_{1080}$ mutants at 0 and 30 ng/ml ciprofloxacin. N_{cells} = 7,804. N_{spots} = 36,595. $N_{nucleoids}$ = 11,215.

RecB can be used as a marker to quantify DSB formation *in vivo*. Given RecBCD's high affinity for DSBs *in vivo* and our ability to detect its binding to DSBs, the system is a useful tool to detect the presence of DSBs in *E. coli*. Our experiments allowed us to estimate the number of recruitments of RecB to DSBs per hour, under different ciprofloxacin concentrations (Figure 3A), and in the $\Delta recA$ and $recB_{1080}$ mutants (Figure 3B). In the absence of ciprofloxacin, we estimated that 1.3 ± 1.1 RecB molecules are recruited to DSBs per hour. A previous study had estimated that 18% of cells generated an endogenous DSB per cell cycle (12), with a division time of 20 min. This would correspond to a rate of ~0.6 DSBs per hour, a lower value than our own estimate. This discrepancy could be due to the significant differences in the methods used to produce these estimates.

In the wild-type, the general pathway for DSB repair suggests that RecBCD is recruited once per DSB (Figure 5). We can, therefore, reasonably expect that the observed number of RecB recruitments matches the number of DSBs formed

(1 to 8 per hour, depending on ciprofloxacin concentration). In the $\Delta recA$ and $recB_{1080}$ mutants, however, the DSB repair pathway disruptions lead to multiple RecB recruitments per DSB.

It should be noted that in some cases, ciprofloxacin is expected to produce double-sided DSBs, hence, leading to two RecB recruitments per break. We did not take this into account in our estimation of the DSB formation rate, as it has been previously suggested that the two sides of a DSB are kept in close proximity during DNA repair (57, 58), and our imaging setup would not allow us to separate two RecB spots located close together. Indeed, our timelapse images of RecB binding seldom show two spots next to each other, further supporting the idea that we cannot resolve RecBCD binding to both sides of a DSB.

Imaging DSBs in live cells is challenging and was previously achieved using the Gam protein of bacteriophage Mu, which binds to free double-stranded DNA ends (59). However, Gam binding to DSBs prevents RecBCD action and, therefore, the

repair of the breaks. Using RecBCD as a probe allows us to monitor DSB formation in live *E. coli* in real-time, with single-cell sensitivity and without perturbing the repair process. This tool could be further improved by substituting the Halo tag with a bright and photostable fluorescent protein [as was used in our previous work (46)], enabling long-term imaging, for example, in a mother-machine microfluidic device.

RecB is recruited to DSBs multiple times in the Δ*recA* and *recB*$_{1080}$ mutants. Our observations of RecB recruitment to DSBs (Figure 3) and nucleoid position (Figure 4) in the different mutant strains have led us to the model of RecB recruitment described in Figure 5, which confirms and extends the model we previously proposed in (37). In wild-type cells, a DSB is recognised by RecBCD, which promotes RecA loading. The RecA filament triggers the SOS response, and is used for homology search and repair. In the Δ*recA* mutant, the 3' ssDNA generated by RecBCD is first coated by SSB, and then degraded by the SbcCD and ExoI nucleases (60). This leads to blunting of the DNA end, creating a new substrate on which RecBCD can bind. This circle leads to multiple RecBCD recruitments per DSB, and eventually to full chromosome degradation. In the *recB*$_{1080}$ mutant, RecBCD unwinds DNA without degrading it. Our data suggests that two competing pathways take place: either DNA degradation by SbcCD and ExoI leading to DNA-end blunting and re-recruitment of RecBCD; or RecA loading by RecFOR, leading to SOS induction and homologous repair.

RecA filaments accumulate under high DNA damage. Under exposure to high ciprofloxacin concentrations (20-30 ng/ml), we observed an accumulation of cells that contained a RecA filament (Supp. Figures 12A and 12B). At these ciprofloxacin concentrations, we have determined that cells undergo frequent DSBs, leading to multiple recruitments of RecB to DSBs over the course of the experiment (up to 8 per hour, Figure 3A). Given RecBCD's high processivity (3), such a high number of RecB recruitments would likely result in several kilobases of DNA being digested at different chromosomal locations, which could result in the complete absence of a homologous copy of the DSB site. In this case, we expect that the repair process will stall at the homology search stage, after formation of the RecA filament, which is consistent with our observation that a large proportion of cells contain RecA filaments following exposure to over-MIC ciprofloxacin concentrations.

RecB colocalises with the bacterial nucleoid. Imaging RecB simulteanously to the bacterial nucleoid reinforced our interpretation that long-lived RecB spots (>10 sec) are DSB-bound RecB molecules, and helped us understand their spatial distribution in the cell. Nucleoid compaction in *E. coli* had previously been reported after UV irradiation (23), and we were able to confirm that a similar compaction occurs under ciprofloxacin treatment (Supp. Figure 13). As expected, DSB-bound RecB molecules colocalised with the nucleoid. In the Δ*recA* and *recB*$_{1080}$ mutants, nucleoid imag-

ing pointed out the involvement of RecA loading (and probably SOS induction) in nucleoid compaction, again confirming previous observations (23). The absence of any significant nucleoid compaction in the *recB*$_{1080}$ mutant despite its ability to induce the SOS response was surprising, and could result from a significant delay in SOS induction as a result of the loading of RecA by the RecFOR alternative pathway.

Taken together, this work provides a detailed picture of RecBCD's interaction with DNA following DSBs. We found that RecB dissociation is independent of RecA and that binding time is unaffected by DNA damage levels. Additionally, we observed changes in nucleoid conformation in response to DNA damage and examined the effects of high antibiotic levels on the formation of RecA filaments. The model we developed offers an overview of RecBCD recruitment to DSBs and dissociation from DNA, depending on the strain's ability to effectively carry out or complete repair. These findings contribute valuable insights into the dynamics of DNA repair mechanisms in *E. coli*, suggesting that RecBCD dissociation from DNA is an intrinsic property of the complex, largely unaffected by external factors, such as DNA damage levels or RecA loading.

FUNDING
This work has been supported by Wellcome Trust Investigator Awards (Grant No. 205008/Z/16/Z awarded to M.E.K.), a BBSRC BB/S008012/1 responsive mode award (to M.E.K.), and a Marie Skłodowska-Curie Personal Fellowship (Grant No. 101063725-BARTAS) awarded to A.L.

ACKNOWLEDGEMENTS
We would like to thank Prof. David Leach for insightful discussions on DNA repair in *E. coli*. We thank Mark Dillingham for sharing his knowledge on the biochemistry of RecBCD. We thank Johan Elf for kindly sharing the RecA-SYFP2 tandem fusion strain. We thank Jean Ollion for the development of BACMMAN and its associated tools, as well as the image analysis support he provided. We thank Elise Darmon for discussing results, and reviewing the manuscript.

AUTHOR CONTRIBUTIONS
M.E.K. and D.T. conceived the experiments and designed the data analysis. D.T., A.L., and L.McL. built the strains. D.T. collected and analysed the data. M.E.K., D.T. and A.L. discussed the data. D.T. wrote the manuscript's first draft and created the figures. M.E.K., D.T. and A.L. revised the manuscript. L.McL., as the lab manager, oversaw order placements and maintained the lab organisation. M.E.K. provided funding. All authors read, edited and approved the final manuscript.

COMPETING FINANCIAL INTERESTS
The authors declare no competing financial interests.

Bibliography

1. Claire Wyman and Roland Kanaar. Dna double-strand break repair: All's well that ends well. *Annual Review of Genetics*, 40:363–383, 12 2006. ISSN 00664197. doi: 10.1146/annurev.genet.40.110405.090451.

2. Mark S. Dillingham and Stephen C. Kowalczykowski. Recbcd enzyme and the repair of double-stranded dna breaks. *Microbiology and Molecular Biology Reviews*, 72:642–671, 12 2008. ISSN 1092-2172.

3. Jakub Wiktor, Marit Van Der Does, Lisa Büller, David J. Sherratt, and Cees Dekker. Direct observation of end resection by recbcd during double-stranded dna break repair in vivo. *Nucleic Acids Research*, 46:1821–1833, 2 2018. ISSN 0305-1048. doi: 10.1093/NAR/GKX1290.

4. Andrew F. Taylor and Gerald R. Smith. Recbcd enzyme is altered upon cutting dna at a chi recombination hotspot. *Proceedings of the National Academy of Sciences*, 89:5226–5230, 6 1992. ISSN 00278424. doi: 10.1073/PNAS.89.12.5226.

5. Charlotte A. Cockram, Milana Filatenkova, Vincent Danos, Meriem El Karoui, and David R.F. Leach. Quantitative genomic analysis of reca protein binding during dna double-strand break repair reveals recbcd action in vivo. *Proceedings of the National Academy of Sciences of the United States of America*, 112:E4735–E4742, 8 2015. ISSN 10916490. doi: 10.1073/pnas.1424269112.

6. Jason J. Churchill and Stephen C. Kowalczykowski. Identification of the reca protein-loading domain of recbcd enzyme. *Journal of Molecular Biology*, 297:537–542, 3 2000. ISSN 0022-2836. doi: 10.1006/JMBI.2000.3590.

7. Maria Spies and Stephen C. Kowalczykowski. The reca binding locus of recbcd is a general domain for recruitment of dna strand exchange proteins. *Molecular Cell*, 21:573–580, 2 2006. ISSN 10972765. doi: 10.1016/j.molcel.2006.01.007.

Thédié *et al.* | Single molecule imaging of RecB *in vivo* reveals dynamics of DSB repair in *E. coli*

Fig. 5. RecBCD recruitment pathways in wild-type *E. coli*, Δ*recA* and *recB*$_{1080}$ mutants. **(Wild-type)** After DSB recognition, RecBCD degrades DNA until it recognises a Chi-site. It switches activity to create a 3' ssDNA overhang, and promotes RecA loading. The RecA-coated ssDNA can then be used for DNA repair by homologous recombination. **(Δreca)** In the absence of RecA, the 3' ssDNA is coated by SSB, and eventually digested by cellular nucleases. Blunting of the DNA end by digestion of the ssDNA creates a new substrate for binding of RecBCD. **(RecB$_{1080}$)** Following DSB recognition, RecB$_{1080}$ unwinds DNA without digesting it. The unwound ssDNA can either be digested by nucleases, leading to a new blunt dsDNA end and RecBCD re-recruitment; or the RecFOR complex displaces SSB to load RecA, allowing DNA repair by homologous recombination to proceed.

8. Claire Wyman and Roland Kanaar. Homologous recombination: Down to the wire. *Current Biology*, 14:R629–R631, 8 2004. ISSN 09609822. doi: 10.1016/j.cub.2004.07.049.

9. Jakub Wiktor, Arvid H. Gynnå, Prune Leroy, Jimmy Larsson, Giovanna Coceano, Ilaria Testa, and Johan Elf. Reca finds homologous dna by reduced dimensionality search. *Nature 2021 597:7876*, 597:426–429, 9 2021. ISSN 1476-4687. doi: 10.1038/s41586-021-03877-6.

10. Taylor A and Smith G. Regulation of homologous recombination: Chi inactivates recbcd enzyme by disassembly of the three subunits. *Genes and development*, 1999.

11. Bénédicte Michel, S. Dusko Ehrlich, and Marilyne Uzest. Dna double-strand breaks caused by replication arrest. *The EMBO Journal*, 16:430–438, 1 1997. ISSN 02614189. doi: 10.1093/EMBOJ/16.2.430.

12. Anurag Kumar Sinha, Christophe Possoz, Adeline Durand, Jean Michel Desfontaines, François Xavier Barre, David R.F. Leach, and Bénédicte Michel. Broken replication forks trigger heritable dna breaks in the terminus of a circular chromosome. *PLOS Genetics*, 14: e1007256, 3 2018. ISSN 1553-7404. doi: 10.1371/JOURNAL.PGEN.1007256.

13. Marie Seigneur, Vladimir Bidnenko, S. Dusko Ehrlich, and Bénédicte Michel. Ruvab acts at arrested replication forks. *Cell*, 95:419–430, 10 1998. ISSN 00928674. doi: 10.1016/S0092-8674(00)81772-9.

14. B. Michel, M. J. Flores, E. Viguera, G. Grompone, M. Seigneur, and V. Bidnenko. Rescue of arrested replication forks by homologous recombination. *Proceedings of the National Academy of Sciences*, 98:8181–8188, 7 2001. ISSN 00278424. doi: 10.1073/PNAS.111008798.

15. Michael A. Kohanski, Daniel J. Dwyer, and James J. Collins. How antibiotics kill bacteria: from targets to networks. *Nature Reviews Microbiology 2010 8:6*, 8:423–435, 5 2010. ISSN 1740-1534. doi: 10.1038/nrmicro2333.

16. Nikola Ojkic, Elin Lilja, Susana Direito, Angela Dawson, Rosalind J. Allen, and Bartlomiej Waclaw. A roadblock-and-kill mechanism of action model for the dna-targeting antibiotic ciprofloxacin. *Antimicrobial Agents and Chemotherapy*, 64, 9 2020. ISSN 10986596. doi: 10.1128/AAC.02487-19/SUPPL_FILE/AAC.02487-19-S0001.PDF.

17. L. M. Wentzell and A. Maxwell. The complex of dna gyrase and quinolone drugs on dna forms a barrier to the t7 dna polymerase replication complex. *Journal of Molecular Biology*, 304:779–791, 12 2000. ISSN 0022-2836. doi: 10.1006/JMBI.2000.4266.

18. Karl Drlica, Muhammad Malik, Robert J. Kerns, and Xilin Zhao. Quinolone-mediated bacterial death. *Antimicrobial Agents and Chemotherapy*, 52:385–392, 2 2008. ISSN 00664804.

19. Shar Yin N. Huang, Stephanie A. Michaels, Brianna B. Mitchell, Nadim Majdalani, Arnaud Vanden Broeck, Andres Canela, Yuk Ching Tse-Dinh, Valerie Lamour, and Yves Pommier. Exonuclease vii repairs quinolone-induced damage by resolving dna gyrase cleavage complexes. *Science Advances*, 7, 3 2021. ISSN 23752548.

20. Xilin Zhao, Muhammad Malik, Nymph Chan, Alex Drlica-Wagner, Jian Ying Wang, Xinying Li, and Karl Drlica. Lethal action of quinolones against a temperature-sensitive dnab replication mutant of escherichia coli. *Antimicrobial Agents and Chemotherapy*, 50:362–364, 1 2006. ISSN 00664804.

21. Zeynep Baharoglu and Didier Mazel. Sos, the formidable strategy of bacteria against aggressions. *FEMS Microbiology Reviews*, 38:1126–1145, 11 2014. ISSN 0168-6445. doi: 10.1111/1574-6976.12077.

22. Julia Bos, Qiucen Zhang, Saurabh Vyawahare, Elizabeth Rogers, Susan M. Rosenberg, and Robert H. Austin. Emergence of antibiotic resistance from multinucleated bacterial filaments. *Proceedings of the National Academy of Sciences of the United States of America*, 112:178–183, 1 2015. ISSN 10916490.

23. Ingvild Odsbu and Kirsten Skarstad. Dna compaction in the early part of the sos response is dependent on recn and reca. *Microbiology (United Kingdom)*, 160:872–882, 5 2014. ISSN 14652080.

24. Z. I. Horii and K. Suzuki. Degradation of the dna of escherichia coli k12 rec- (jc1569b) after irradiation with ultraviolet light. *Photochemistry and Photobiology*, 8:93–105, 8 1968. ISSN 1751-1097. doi: 10.1111/J.1751-1097.1968.TB05850.X.

25. Kin-Hoe Chow and Justin Courcelle. Recbcd and recj/recq initiate dna degradation on distinct substrates in uv-irradiated escherichia coli. *Radiation Research*, 168:499–506, 10 2007. ISSN 0033-7587. doi: 10.1667/RR1033.1.

26. Misook Yu, Jehanne Souaya, and Douglas A. Julin. Identification of the nuclease active site in the multifunctional recbcd enzyme by creation of a chimeric enzyme. *Journal of Molecular Biology*, 283:797–808, 11 1998. ISSN 0022-2836. doi: 10.1006/JMBI.1998.2127.

27. Jingdi Wang, Ruiwu Chen, and Douglas A. Julin. A single nuclease active site of the escherichia coli recbcd enzyme catalyzes single-stranded dna degradation in both directions. *Journal of Biological Chemistry*, 275:507–513, 1 2000. ISSN 00219258. doi: 10.1074/jbc.275.1.507.

28. Daniel G. Anderson, Jason J. Churchill, and Stephen C. Kowalczykowski. A single mutation, recb(d1080a), eliminates reca protein loading but not chi recognition by recbcd enzyme. *Journal of Biological Chemistry*, 274:27139–27144, 9 1999. ISSN 00219258. doi: 10.1074/jbc.274.38.27139.

29. Ivana Ivančić-Baće, Petra Peharec, Sunčana Moslavac, Nikolina Škrobot, Erika Salaj-Šmic, and Krunoslav Brčić-Kostić. Recfor function is required for dna repair and recombination in a reca loading-deficient recb mutant of escherichia coli. *Genetics*, 163:485–494, 2 2003. ISSN 00166731. doi: 10.1093/GENETICS/163.2.485.

30. Maria Spies, Piero R. Bianco, Mark S. Dillingham, Naofumi Handa, Ronald J. Baskin, and Stephen C. Kowalczykowski. A molecular throttle: The recombination hotspot χ controls dna translocation by the recbcd helicase. *Cell*, 114:647–654, 9 2003. ISSN 0092-8674. doi: 10.1016/S0092-8674(03)00681-0.

31. Bian Liu, Ronald J. Baskin, and Stephen C. Kowalczykowski. Dna unwinding heterogeneity by recbcd results from static molecules able to equilibrate. *Nature 2013 500:7463*, 500: 482–485, 7 2013. ISSN 1476-4687. doi: 10.1038/nature12333.

32. Chirlmin Joo, Sean A. McKinney, Muneaki Nakamura, Ivan Rasnik, Sua Myong, and Taekjip Ha. Real-time observation of reca filament dynamics with single monomer resolution. *Cell*, 126:515–527, 8 2006. ISSN 00928674. doi: 10.1016/j.cell.2006.06.042.

33. Roberto Galletto, Ichiro Amitani, Ronald J. Baskin, and Stephen C. Kowalczykowski. Direct observation of individual reca filaments assembling on single dna molecules. *Nature 2006 443:7113*, 443:875–878, 9 2006. ISSN 1476-4687. doi: 10.1038/nature05197.

34. Naofumi Handa, Ichiro Amitani, Nathan Gumlaw, Steven J. Sandler, and Stephen C. Kowalczykowski. Single molecule analysis of a red fluorescent reca protein reveals a defect in nucleoprotein filament nucleation that relates to its reduced biological functions. *Journal of Biological Chemistry*, 284:18664–18673, 7 2009. ISSN 00219258. doi: 10.1074/jbc.M109.004895.

35. Anthony L. Forget and Stephen C. Kowalczykowski. Single-molecule imaging of dna pairing by reca reveals a three-dimensional homology search. *Nature 2012 482:7385*, 482:423–427, 2 2012. ISSN 1476-4687. doi: 10.1038/nature10782.

36. Kaushik Ragunathan, Cheng Liu, and Taekjip Ha. Reca filament sliding on dna facilitates homology search. *eLife*, 2012, 12 2012. ISSN 2050084X. doi: 10.7554/ELIFE.00067.

37. Alessia Lepore, Daniel Thédié, Lorna McLaren, Benura Azeroglu, Oliver J. Pambos, Achillefs N. Kapanidis, and Meriem El Karoui. In vivo single-molecule imaging of recb reveals efficient repair of dna damage in escherichia coli. *bioRxiv*, page 2023.12.22.573010, 12 2023. doi: 10.1101/2023.12.22.573010.

38. Nicholas Renzette, Nathan Gumlaw, Jared T. Nordman, Marlee Krieger, Su Ping Yeh, Edward Long, Richard Centore, Ruethairat Boonsombat, and Steven J. Sandler. Localization of reca in escherichia coli k-12 using reca–gfp. *Molecular Microbiology*, 57:1074–1085, 8 2005. ISSN 1365-2958. doi: 10.1111/J.1365-2958.2005.04755.X.

39. Nicholas Renzette, Nathan Gumlaw, and Steven J. Sandler. Dini and recx modulate reca–dna structures in escherichia coli k-12. *Molecular Microbiology*, 63:103–115, 1 2007. ISSN 1365-2958. doi: 10.1111/J.1365-2958.2006.05496.X.

40. Richard C. Centore and Steven J. Sandler. Uvrd limits the number and intensities of reca-green fluorescent protein structures in escherichia coli k-12. *Journal of Bacteriology*, 189: 2915–2920, 4 2007. ISSN 00219193.

41. Vincent Amarh, Martin A. White, and David R.F. Leach. Dynamics of reca-mediated repair of replication-dependent dna breaks. *Journal of Cell Biology*, 217:2299–2307, 7 2018. ISSN 0021-9525. doi: 10.1083/JCB.201803020.

42. Dawit Kidane and Peter L. Graumann. Dynamic formation of reca filaments at dna double strand break repair centers in live cells. *Journal of Cell Biology*, 170:357–366, 9 2005. ISSN 00219525.

43. Christian Lesterlin, Graeme Ball, Lothar Schermelleh, and David J. Sherratt. Reca bundles mediate homology pairing between distant sisters during dna break repair. *Nature 2013 506:7487*, 506:249–253, 12 2013. ISSN 1476-4687. doi: 10.1038/nature12868.

44. Harshad Ghodke, Bishnu P. Paudel, Jacob S. Lewis, Slobodan Jergic, Kamya Gopal, Zachary J. Romero, Elizabeth A. Wood, Roger Woodgate, Michael M. Cox, and Antoine M.Van Oijen. Spatial and temporal organization of reca in the escherichia coli dna-damage response. *eLife*, 8, 2 2019. ISSN 2050084X. doi: 10.7554/ELIFE.42761.

45. Anjana Badrinarayanan, Tung B.K. Le, and Michael T. Laub. Rapid pairing and resegregation of distant homologous loci enables double-strand break repair in bacteria. *Journal of Cell Biology*, 210:385–400, 8 2015. ISSN 0021-9525. doi: 10.1083/JCB.201505019.

46. Alessia Lepore, Hannah Taylor, Dirk Landgraf, Burak Okumus, Sebastián Jaramillo-Riveri, Lorna McLaren, Somenath Bakshi, Johan Paulsson, and M. El Karoui. Quantification of very low-abundant proteins in bacteria using the halotag and epi-fluorescence microscopy. *Scientific Reports*, 9:1–9, 12 2019. ISSN 20452322. doi: 10.1038/s41598-019-44278-0.

47. Jean Ollion, Marina Elez, and Lydia Robert. High-throughput detection and tracking of cells and intracellular spots in mother machine experiments. *Nature Protocols 2019 14:11*, 14: 3144–3161, 9 2019. ISSN 1750-2799. doi: 10.1038/s41596-019-0216-9.

48. Jean Ollion, Charles Ollion, Elisabeth Gassiat, Luc Lehéricy, Sylvain Le Corff, and Laboratoire J A Dieudonné. Joint self-supervised blind denoising and noise estimation. *arxiv*, 2 2021.

49. Hirofumi Kobayashi, Ahmet Can Solak, Joshua Batson, and Loic A Royer. Image deconvolution via noise-tolerant self-supervised inversion. *arxiv*, 2020.

50. N Otsu. A threshold selection method from gray-level histograms. *IEEE Trans SMC*, 9:62–, 1979. doi: 10.1371/JOURNAL.PONE.0035550.

51. Ji Yu, Jie Xiao, Xiaojia Ren, Kaiqin Lao, and X. Sunney Xie. Probing gene expression in live cells, one protein molecule at a time. *Science*, 311:1600–1603, 3 2006. ISSN 00368075.

52. Johan Elf, Gene Wei Li, and X. Sunney Xie. Probing transcription factor dynamics at the single-molecule level in a living cell. *Science*, 316:1191–1194, 5 2007. ISSN 00368075.

53. Irina Kalita, Ira Alexandra Iosub, Lorna McLaren, Louise Goossens, Sander Granneman, and Meriem El Karoui. An hfq-dependent post-transcriptional mechanism fine tunes recb expression in escherichia coli. *eLife*, 13, 5 2024. doi: 10.7554/ELIFE.94918.1.

54. Martin Wilkinson, Lucy Troman, Wan AK Wan Nur Ismah, Yuriy Chaban, Matthew B Avison, Mark S Dillingham, and Dale B Wigley. Structural basis for the inhibition of recbcd by gam and its synergistic antibacterial effect with quinolones. *eLife*, 5, 12 2016. doi: 10.7554/ELIFE.22963.

55. Florence N. Capaldo and Stephen D. Barbour. Dna content, synthesis and integrity in dividing and non-dividing cells of rec- strains of escherichia coli k12. *Journal of Molecular Biology*, 91:53–66, 1 1975. ISSN 0022-2836. doi: 10.1016/0022-2836(75)90371-X.

56. K. Skarstad and E. Boye. Degradation of individual chromosomes in reca mutants of escherichia coli. *Journal of Bacteriology*, 175:5505–5509, 1993. ISSN 00219193. doi: 10.1128/JB.175.17.5505-5509.1993.

57. Elise Vickridge, Charlene Planchenault, Charlotte Cockram, Isabel Garcia Junceda, and Olivier Espéli. Management of e. coli sister chromatid cohesion in response to genotoxic stress. *Nature Communications 2017 8:1*, 8:1–12, 3 2017. ISSN 2041-1723. doi: 10.1038/ncomms14618.

58. Kenji Keyamura and Takashi Hishida. Topological dna-binding of structural maintenance of chromosomes-like recn promotes dna double-strand break repair in escherichia coli. *Communications Biology 2019 2:1*, 2:1–12, 11 2019. ISSN 2399-3642. doi: 10.1038/s42003-019-0655-4.

59. Chandan Shee, Ben D. Cox, Franklin Gu, Elizabeth M. Luengas, Mohan C. Joshi, Li Ya Chiu, David Magnan, Jennifer A. Halliday, Ryan L. Frisch, Janet L. Gibson, Ralf Bernd Nehring, Huong G. Do, Marcos Hernandez, Lei Li, Christophe Herman, P. J. Hastings, David Bates, Reuben S. Harris, Kyle M. Miller, and Susan M. Rosenberg. Engineered proteins detect spontaneous dna breakage in human and bacterial cells. *eLife*, 2013, 10 2013. ISSN 2050084X. doi: 10.7554/ELIFE.01222.001.

60. Ksenija Zahradka, Maja Buljubašić, Mirjana Petranović, and Davor Zahradka. Roles of exol and sbccd nucleases in "reckless" dna degradation in reca mutants of escherichia coli. *Journal of Bacteriology*, 191:1677–1687, 3 2009. ISSN 00219193. doi: 10.1128/jb.01877-07.

61. Daniel Thédié, Romain Berardozzi, Virgile Adam, and Dominique Bourgeois. Photoswitching of green meos2 by intense 561 nm light perturbs efficient green-to-red photoconversion in localization microscopy. *Journal of Physical Chemistry Letters*, 8:4424–4430, 9 2017. ISSN 19487185. doi: 10.1021/ACS.JPCLETT.7B01701/ASSET/IMAGES/LARGE/JZ-2017-01701K_0004.JPEG.

62. Jonathan B. Grimm, Brian P. English, Jiji Chen, Joel P. Slaughter, Zhengjian Zhang, Andrey Revyakin, Ronak Patel, John J. Macklin, Davide Normanno, Robert H. Singer, Timothée Lionnet, and Luke D. Lavis. A general method to improve fluorophores for live-cell and single-molecule microscopy. *Nature Methods 2015 12:3*, 12:244–250, 1 2015. ISSN 1548-7105. doi: 10.1038/nmeth.3256.

63. Kenneth F. Wertman, Arlene R. Wyman, and David Botstein. Host/vector interactions which affect the viability of recombinant phage lambda clones. *Gene*, 49:253–262, 1 1986. ISSN 0378-1119. doi: 10.1016/0378-1119(86)90286-6.