Barley disease screening: a multiplex digital droplet PCR approach for the detection of *Ramularia collo-cygni, Rhynchosporium graminicola* and *Pyrenophora teres*

Rabisa Zia[1,2], Zoë A. Popper[2], Steven Kildea[1*]

[1] Department of Crops Science, Teagasc, Crops, Environment and Land Use Programme, Oak Park, Carlow, Ireland

[2] Botany and Plant Science and Martin Ryan Institute, School of Natural Sciences, College of Science and Engineering, University of Galway, Ireland

[*]Corresponding author: S. Kildea; E-mail: stephen.kildea@teagasc.ie

Keywords: barley pathogens, Ramularia leaf spot, Barley leaf scald, Net blotch, endpoint quantification, TaqMan assay, seed screening

ABSTRACT

Barley is the most widely grown cereal crop in Ireland and faces substantial yield loss annually due to airborne, seed-borne, and soil-borne pathogens. The most prevalent and destructive of these include Ramularia leaf spot, Barley leaf scald and Net blotch, caused by *Ramularia collo-cygni, Rhynchosporium graminicola* and *Pyrenophora teres* respectively. Infected seeds are considered as the major source of inoculum for these diseases. The absence of varietal resistance for these fungal pathogens suggests the need for better disease management strategies. Accurate pathogen detection is one of the most widely accepted disease management strategies. Here we demonstrate two digital droplet PCR (ddPCR) assays, based on TaqMan probe chemistry, for the detection of the above pathogens. ddPCR is a third generation of PCR, relying on sample partition into thousands of droplets in which each droplet acts as a reaction unit, enabling pathogen detection at extremely low levels. We developed a multiplex assay capable of detecting *R. graminicola*, *R.collo-cygni* and *P. teres*. For *P. teres* samples, a further assay was designed to distinguish between the two economically significant forms of *P. teres* i.e. *P. teres* f. maculata (Ptm; spot form of Net blotch) and *P. teres* f. teres (Ptt; net form of Net blotch). The assay offers sensitive and selective pathogen detection from barley seeds to pico moles/µl level. Finally, the efficacy of the assay was demonstrated by screening a collection of historical barley seeds.

1 | INTRODUCTION

After wheat and corn, barley is the 3rd most widely produced cereal crop in Europe, making up almost 20% of the total cereals produced (European Union Crop Production, 2024). Due to the combination of agronomic and climatic factors, the potential for high yields and favourable end-markets, barley accounts for approximately 70% of all cereals produced in Ireland (CSO, 2023). Unfortunately, as with all crops, barley can succumb to a spectrum of diseases that if left unchecked can seriously impact potential yields, both quantitatively and qualitatively. Most notable amongst these are the fungal foliar diseases ramularia leaf spot (RLS) caused by *Ramularia collo-cygni,* barley scald caused by *Rhynchosporium graminicola,* and the spot and net forms of net blotch (SFNB and NFNB), caused by *Pyrenophora teres* f. *maculata* (Ptm) and *P. teres* f. *teres* (Ptt) respectively (Huss et al. 1992; Jalli et al. 2020; Zhang et al. 2020).

All three diseases impact yield by reducing the healthy leaf area of the crop available for photosynthesis, and as barley is typically more sink limited than source the earlier disease develops in the crop the greater the potential impact on yield (Bingham et al. 2007). As is the case with all plant diseases and pests, limiting initial inoculum/pest sources is critical to achieving sustainable control and is the principal tactic of integrated pest management (Barzman et al. 2015). For *R. collo-cygni*, *R. graminicola* and *P. teres* trash and stubble from the previous season's crops, volunteers and infected seed, all act as potential sources of initial inoculum. For *R. graminciola* and *P. teres* it has been previously demonstrated that seed-borne infection represents a significant means of disease transmission and can directly impact disease levels and subsequent impacts on barley yields (Fountaine et al. 2010; Hysing & Wiik, 2013). For *R. collo-cygni*, whilst the pathogen can be readily detected in seed, and it has been

demonstrated that it can grow from the seed into the developing seedling, the importance of

seed-borne infection in the aetiology of RLS remains to be determined (Havis et al. 2006).

Traditionally seed has been tested for the presence of pathogens using culture dependent

techniques, whereby the test seeds are cultured on a selective growth media and subsequent

microbial growth is examined morphologically and/or microscopically to confirm the

contaminating species (Hiddink et al. 2023). However, there is a major disadvantage of

culture-dependent assays which is that they are restricted to those pathogens that can be

easily grown in culture and have a relatively fast growth rate. Of the three major barley

pathogens identified above, it is only possible to routinely identify *P. teres* through culture-

dependent assays as both *R. graminicola* and *R. collo-cygni* are relatively slow growing and

are outcompeted by other fungi (pathogenic or non-pathogenic) present in barley seed.

Identification of *P. teres* in culture can also be difficult as it can be mistaken for *P. graminea*,

which exhibits a similar phenology in culture. Similarly, it is not possible to distinguish

between Ptt and Ptm when grown in culture or by using osmotic or blotter tests (Brodal,

1997). More recently developed detection methods that rely on detection of nucleic acids

(either DNA or RNA) have improved specificity compared to culture-based detection

methods. They also allow a wider range of pathogens to be detected, whilst simultaneously

reducing costs and run time (Elnifro Elfath et al. 2000).

Seed samples are unfortunately often complex matrixes, which can inhibit or impair DNA

based reactions and assay sensitivities (Lu et al. 2020), in particular those such as qPCR on

which quantification is based on comparable efficiencies of reactions required between the

test sample and samples/standards of a known DNA concentration (Hindson et al. 2013).

Digital droplet PCR (ddPCR, also referred to as digital PCR (dCPR) or crystal digital PCR (cdPCR)

depending on the platform used) is a third generation of PCR in which the starting DNA sample

is partitioned through the use of microfluidics into thousands of samples which subsequently

undergo PCR reactions. By coupling partitioning and PCR with fluorescent dye based DNA

probes it is possible to use endpoint analysis to confirm the presence or absence of the

targeted DNA partitioned samples. Through the use of Poisson's Law it is then possible to

determine the total copies of the target DNA in the starting sample. As quantification is based

on an end point reaction it is not as sensitive to PCR inhibitors etc which can impact methods

such as qPCR which can be strongly influenced by the efficiencies of the PCR reaction (Basu,

2017). Therefore, ddPCR allows for the accurate and sensitive quantification of specific DNA

targets, even in the most complex of matrices, making it an ideal platform for the detection

of pathogens in cereal seeds.

Different PCR based methods have been developed for each of the commonly occurring

barley pathogens, with qPCR assays available for both *P. teres* and *R. graminicola* (Bates et al.

2001; Fountaine et al. 2010), and a dPCR based method for detection of *R. collo-cygni* recently

published by Knight et al. (2022). To exploit the multi-spectral capacity of the Stilla 3-colour

Nacia® ddPCR system here we combine the *R. graminicola* and *R. collo-cygni* assays of

Fountaine et al. (2010) and Knight et al. (2022), with a *P. teres* probe based assay to provide

a multiplex ddPCR assay to simultaneously detect and quantify the presence of the three

pathogens. A second multiplex ddPCR is also described to further differentiate those samples

which were positive for *P. teres* in order to determine whether form i.e. Ptm or Ptt.

2 | MATERIALS AND METHODS

A multistage process of assay design, optimisation and application was developed and is

outlined in Figure 1.

2.1 | Fungal strains and DNA extraction

All fungal cultures used in the study (Supplementary Table 1) were acquired from the phytopathological collection in Teagasc, Oak Park, Co. Carlow. DNA was extracted from these cultures following lyophilisation and homogenisation using a Qiagen DNeasy Plant Mini kit (Qiagen, Germany), following the manufacturer's instructions. Extracted DNA was quantified using Invitrogen™ Qubit™ Fluorometer with a Qubit™ 1X dsDNA BR Assay (Invitrogen, US).

2.1 | Assay designs and initial optimization

For the initial ddPCR multiplex assay to detect and distinguish between the three target fungal pathogens, the published fluorescence probe based assays for the detection of *R. collo-cygni* (Knight et al. 2022) and *R. secalis* (Fountaine et al. 2010) were selected. A separate set of primers and probe to detect *P. teres* was designed using Primer 3Plus (https://www.primer3plus.com/index.html) based on the Internal Transcribed Spacer (ITS) region of *P. teres* (OM471876.1). To further distinguish between Ptt and Ptm a set of primers and probe was designed using PrimerQuest™ (Integrated DNA Technologies) based on differences in their mRNA region as described by Poudel et al. (2017) (accession numbers Ptt: KX909556.1 and Ptm: KX909562.1). Details of all primers and probes are summarized in Table 1. Initial primer and probe optimisation, including identifying annealing temperatures, was conducted using routine PCR and visualised using agarose gel electrophoresis (see Supplementary Figure S1 for further details).

2.2 | ddPCR optimisation

Following these initial validation steps, the assays were transferred to ddPCR system (naica® Crystal Digital PCR 3 colour system, Stilla Technologies, France). ddPCR using the naica®

Crystal Digital PCR 3 colour is separated into four stages: 1) the reaction sample (primers,

probes, DNA, etc) is partitioned into nano-sized droplets on a chip (up to 17k on a Ruby chip

or 30k on a Sapphire chip), 2) partitioned samples undergo up to 50 PCR cycles, 3) once the

PCR is complete a scan of the chip is taken under different wavelengths (corresponding to the

excitation of the FAM, HEX and Cy5 dyes) is undertaken, and 4) scanned chips are analysed to

identify droplets with the target DNA amplified based on presence or absence of the specified

molecular probes and associated dyes. In the present study all reactions were conducted

using the Ruby Chip (up-to 17k droplets per chamber) and 5X naica® multiplex PCR MIX (Stilla

technologies, France) and only reactions with >10k analysable droplets were included.

An initial series of reactions were performed to optimize concentrations of primers and

probes required to differentiate between positive and negative droplets. Reaction mixtures

were prepared in accordance with the guidelines provided by Stilla Technologies, consisting

of 1 µl of Buffer A, 0.2 µl of Buffer B, 0.25 µl (both supplied as part of the 5X naica® multiplex

PCR MIX each of the primers (0.05 µM final concentration), 0.15 µl of each probe (0.06 µM

final concentration), 0.85 µl of Barley DNA, 1 µl of template DNA (0.1 ng/µl) and nuclease free

water (Fisher Scientific, UK) was added to reach a final volume of 5 µl. All PCR reactions were

performed in the presence of pathogen-free barley DNA as a matrix, extracted from two-week

barley seedlings (variety: RGT Planet, Goldcrop Ltd, Ireland) grown under controlled

conditions (16 h of light, 8 h of dark, at 20 and 16 °C) Thermal cycling conditions for both

assays are as follows; partition at 25°C followed by 95°C for 3 minutes, then 5 cycles of 95°C

for 10 seconds followed by 66°C for 30 seconds, and then 35 cycles at 95°C for 10 second and

64°C for 15 second. A compensation matrix was designed according to the guidelines provided

by Stilla Technologies, to normalize the fluorescence overlap and spill over.

2.2 | Determining Limits of Blank (LoB) and Limits of Detection (LoD)

Limit of blank (LoB) and Limit of detection (LoD) for the assays were calculated in multiplex

fashion and in accordance with the guidelines provided by Stilla Technologies. LoB can be

defined as the maximum target concentration expected in the blank sample with a probability

PLoB = 1 − α (typically 1 − α = 95% where α = 5% of false positives), where α is the probability

of making a false positive decision in a real blank sample (no DNA of any target except the

background DNA). Whereas, LoD is defined as the minimum concentration, a non-zero,

statistically significant value, that can be detected with a probability of 1 − β (typically 1 − β =

95% where β = 5% of false negatives), β is the probability making a false negative decision in

a real positive sample.

To determine the LoB, 30 blank (negative for the target for which LoB is being performed)

reaction replications were performed for each target. To calculate LoD, ten-fold serial

dilutions of a known target DNA concentration was performed and a non-zero, statistically

significant concentration detected by the assay, designated as low-level sample was

calculated. Afterwards, five independent low-level samples were prepared and six reaction

replications of each were performed. Based on the copy per microliter (cp/μl) measured by

the fluorescence detector, LoD and LoB was calculated by using the online tool provided by

Stilla Technologies (https://www.stillatechnologies.com/digital-pcr/statistical-tools/limit-

detection/).

2.3 | Linearity of the assay

To calculate the relationship between the expected and measured quantities of each of the

target pathogen DNA, linearity of the assay was calculated. To achieve this, a 10-fold dilution

series (0.01-0.00001 ng/μl) of each pathogen DNA (*R. collo-cygni*, *R. graminicola* and *P. teres*)

was prepared and tested individually whilst keeping a constant DNA concentration of the

other two pathogens (0.1 ng/µl) in the reaction mixture. The log10 values of concentration

against copy number/µl (cp/ µl) were plotted and correlation was calculated.

2.4 | Specificity of the assays

Specificity of both multiplex assays was performed by challenging the system with a number

of barley pathogens including *Botrytis cinerea (Bc), Parastagonospora nodorum (Pn),*

(Alternaria alternate (Aa), Fusarium graminearum (Fg), Microdochium nivale (Mn) and

Fusarium culmorum (Fc) accessions that are stored as part of the Phytopathological collection

in Teagasc, Oak Park, Co. Carlow. Additionally, specificity of the Ptt and Ptm assay was also

verified against *R. collo-cygni* and *R. graminicola* in a singleplex reaction. Each of the

pathogens was tested in an individual ddPCR reaction.

2.5 | Detection of pathogens in leaf and seed samples

To validate the reliability of the assay for naturally infected barley samples, historical winter

and spring barley seed DNA from 2016–2018, with variable susceptibility to leaf blotches,

were evaluated using the assays. Any droplet population above LoD of the respective assay

was considered positive. Samples positive for *P. teres* were further tested by the form specific

assay to confirm *P. teres* form. Volunteers are one of the sources of inoculum for RLS, barley

leaf scald and net blotch, efficacy of the assay was also assessed in wheat volunteers from

barley field with visual symptoms of RLS (Teagasc wheat trials in Knockbeg, Ireland (2023).

Leaves were freeze dried and DNA was extracted using kit (Qiagen, Germany). Extracted DNA

(1 µl) was used as a template in the reaction mixture described in section 2.2.

3 | Results

3.1 | Assay design and optimization

For the detection of *R. collo-cygni* and *R. graminicola*, primers and probes were deployed

from published assays (Fountaine et al. 2010; Knight et al. 2022). However, for *P. teres*

detection and to distinguish between both Ptt and Ptm it was necessary to design novel

fluorescent probes. The initial multiplex detection assay was specific for the three pathogens,

with no fluorescence overlap observed between the different probes-pathogen combinations

(Figure 2A). Similarly, for the *P. teres* form specific assay there was no fluorescence overlap

between the probe-pathogen combination, demonstrating specificity of the assay (Figure 2B).

For *R. graminicola, R. collo-cygni and P. teres* the LoB were calculated to be 0.68, 0 and 1.34

cp/µl respectively. For the *P. teres* form specific assay and LoB of 0 cp/µl was calculated for

both Ptt and Ptt. Limits of detection (LoD) for *R. graminicola, R. collo-cygni and P. teres* in the

multiplex assay were calculated as 1.67 cp/µl, 1.06 cp/µl and 4.21 cp/µl respectively.

Additionally, LoD was calculated as 1.44 for Ptt assay and 0.82 for Ptm assay. No positive

droplets, irrespective of probe, were detected in negative control (barley DNA only) (Figure

2A & B).

3.2 | Linearity of the assay

To determine the accuracy of the assay, the linearity of the assay was ascertained by

correlating the projected DNA copies of each target in a sample and copy numbers detected.

For each probe-pathogen combination R^2 values of >0.99 were calculated. Linearity of the

assay also indicated the lower limit of detection of the assay, which is 0.00001 ng/µl (Figure

3). The linearity of the *P. teres* form specific assay was not determined as it was specifically

designed to distinguish between forms following initial detection of *P. teres* using the

multiplex assay.

3.3 | Selectivity of the assay

Considering the potential application of our assays, selectivity of both the assays was

investigated against range of fungi that may be found on or in barley seeds and leaves. No

positive droplets were detected in any of the fluorescent channels for any of the fungi tested,

indicating that the assays are highly specific (Figure 4A & B).

3.4 | Detection of barley pathogens in wheat and rye

The sequence similarity between the wheat pathogen *R. graminicola* and the rye pathogen *R.*

secalis, suggested that the developed assay could be used for detection of the pathogen from

rye seed. Rye seed and leaf DNA was tested against the assay. Rye seed samples that did not

receive the standard fungicide regime were found to be positive for the presence of the

pathogen *R. secalis* (Figure 5). Wheat leaf DNA was also tested for the presence of *R. collo-*

cygni. The assay showed a considerable droplet population for *R. collo-cygni*. However, wheat

samples displayed positive droplets for the presence of *R. graminicola and P. teres* DNA in the

samples (Figure 6).

3.5 | Detection of *R. collo-cygni*, *R. graminicola* and *P. teres* in barley seed samples

To determine the application of assays to test the presence of pathogens in naturally infected

barley samples, DNA previously extracted from seeds of winter and spring barley from 2016

(Kildea et al. 2024) were tested. Only samples exhibiting cp/µl equal to or above the limits of

detection, as identified above, were deemed positive for the respective pathogens. Of the 35

winter barley samples tested, 16 samples were positive for *R. collo-cygni*, two samples were

positive for *R. graminicola,* while 19 were positive for *P. teres.* For the 32 spring barley

samples, 15 were positive for *R. graminicola*, 25 samples were positive for *R. collo-cygni,* and

25 positive for *P. teres*. Nine of the spring and winter barley samples that were found to be

positive for *P. teres* were tested to determine the form determination. None of the samples

tested positive for the presence of *P. teres* f. maculata. In the case of winter barley, eight out

of nine samples were positive for *P. teres* f. maculata and two samples were positive for *P.*

teres f. teres as well as *P. teres* f. maculata (Supplementary information).

4 | Discussion

Clean seeds are undeniably important for healthy crops and this is reliant on early detection

of specific disease agents, which can further facilitate the execution of IPM based disease

control strategies (Dussart et al. 2020; Tsedaley, 2015). The ddPCR assays described, offer a

selective, reliable and highly sensitive means of detecting and quantifying three of the most

economically important barley pathogens in a single reaction. As the assay is based on the

digital PCR concept and is therefore end-point in nature which makes it ideal for detecting

the different pathogens in complex plant materials, such as seeds/grains (Liu et al. 2011).

To validate the efficacy of our assays and investigate the prevalence of *R. collo-cygni, R.*

graminicola and *P. teres* in real samples we screened a selection of historical barley seed

samples from 2016 to 2018, rye grains harvested in 2023 and wheat leaves sampled from

volunteer wheat plants in a winter barley field in 2023. Unsurprisingly, the barley seed

samples were contaminated with variable levels of the three pathogens, with a proportion

coinfected with all three pathogens. Only *R. graminicola* was detected in the rye grains, whilst

all three pathogens were detected in the winter wheat leaves, the latter result is unsurprising

given the fact that the leaves tested were collected from wheat volunteers collected from a

winter barley field. Whilst *R. collo-cygni* has previously been reported in wheat in Ireland (Zia

et al. unpublished), the detection of *R. gramincola* and *P. teres* in the sample may simply

reflect the presence of both pathogens on the wheat leaf surface rather than colonisation of

the leaf tissue. This highlights a potential limitation of the assays. As they are based on the

detection of DNA of the respective pathogens, they cannot determine if the detected

pathogen was actively colonising the plant tissue, or if the pathogen detected was even viable.

However, it is important to note that these limitations are not restricted to the assays

developed and described in this paper, but are a drawback of all DNA-based detection

methods (Ficetola et al. 2015). It is therefore critical that any positive detections are assessed

in the context of detection (host, etc.) and the potential risk(s) posed by the pathogen. As

mentioned previously it is more likely that the positive detections for *R. graminciola* and *P.*

teres from the wheat volunteers are consequential to levels of both diseases in the

surrounding barley crop. However, it is worth highlighting that in addition to *R. collo-cygni*

being noted as able to colonise wheat, Ptt has been reported as causing disease on wheat in

Hungary (Tóth et al. 2008), Russia (Mikhailova et al. 2010) and most recently in Brazil (Garozi

et al. 2020). Further analysis of wheat grains and/or a wider wheat disease survey is

warranted to determine if either pathogen is more widespread in wheat crops.

Those barley seed samples that presented positive for *P. teres* were subsequently tested to

determine what form of *P. teres* was present. Amongst the spring barley samples tested only

Ptt was present. For the winter barley seed samples both Ptt and Ptm was detected, with a

number of samples positive for both. All samples with detectable levels of Ptm, irrespective

of the presence of Ptt, were from the cultivar KWS Tower, which had by the late 2010s

become highly susceptible to *P. teres*, and in particular Ptm (Recommendation list in

supplementary information). Positive detection of both Ptt and Ptm could indicate of a

potential Ptt/Ptm hybrid, however based upon the number of droplets positive for both in

the test samples this appears unlikely. However, such detections do suggest such hybrids may

arise, with recent reductions in fungicide sensitivity and efficacy in Western Australia

associated with such a hybrid (Mair et al. 2020; Turo et al. 2021).

It is worth noting that within the multi-pathogen assay the region targeted for *P. teres* was

ITS region, whilst in the form specific assay we targeted the Ptt form-specific region PttQ5

mRNA sequence and the Ptm form-specific region PtmQ11 mRNA sequence. The copy

number for ITS in different fungi can vary considerably (Lofgren et al. 2019) and the copy

number of PtmQ11 and PttQ5 are unknown in *P. teres.* As a number of samples that tested

positive for *P. teres* in the multi-pathogen assay, were neither positive for either Ptt or Ptm

suggests PtmQ11 and PttQ5 have a lower copy number compared to ITS region in *P. teres.*

Therefore, ~ 15 cp/µl of *P. teres* sequence are required to confirm the presence of Ptt or Ptm

from the samples.

Infected seed is the most likely source of introduction of new pathogens and resistance

mechanisms across the globe for all of the three pathogens discussed (Stam et al. 2019).

Therefore, seed screening for pathogen levels and presence of resistance alleles should be

regular practice prior to seed transportation. Furthermore, establishment of pathogen

threshold levels and rejection of seeds with high pathogen levels can minimize the risk of

future outbreaks. Seed is only one of the sources of these diseases however prevention is the

key to suppression.

Acknowledgments

This study is part of BioCrop, a project funded by The Irish Department of , Agriculture, Food

and The Marine under their Research Stimulus Fund (Ref: 2019PROG705). We would like to

thank Dr. Richie Hackett (Teagasc, Oak Park) for providing rye seed samples and Dr. Diana

Bucur (Teagasc, Oak Park) for fungal isolates for assay selectivity testing.

Conflict of interest declaration

None declared.

Data availability statement

278   The data that support the findings of this study are available on reasonable request from the

279   corresponding author.

References

Barzman, M., Bàrberi, P., Birch, A. N. E., Boonekamp, P., Dachbrodt-Saaydeh, S., Graf, B., Hommel,
B., Jensen, J. E., Kiss, J., Kudsk, P., Lamichhane, J. R., Messéan, A., Moonen, A.-C., Ratnadass,
A., Ricci, P., Sarah, J.-L., & Sattin, M. (2015) Eight principles of integrated pest management.
Agronomy for Sustainable Development, 35(4), 1199-1215. https://doi.org/10.1007/s13593-
015-0327-9

Basu, A. S. (2017) Digital assays part I: Partitioning statistics and digital PCR. *SLAS Technology, 22*(4),
369-386. https://doi.org/https://doi.org/10.1177/2472630317705680

Bates, J. A., Taylor, E. J., Kenyon, D. M., & Thomas, J. E. (2001) The application of real-time PCR to the
identification, detection and quantification of *Pyrenophora* species in barley seed. *Mol Plant
Pathol, 2*(1), 49-57. https://doi.org/10.1046/j.1364-3703.2001.00049.x

Bingham, I. J., Blake, J., Foulkes, M. J., & Spink, J. (2007) Is barley yield in the UK sink limited?: II.
Factors affecting potential grain size. *Field Crops Research, 101*(2), 212-220.
https://doi.org/https://doi.org/10.1016/j.fcr.2006.11.004

Brodal, G. (1997) Comparative Tests with the osmotic blotter method for detection of *Drechslera*
spp. in barley seeds. In: Seed Health Testing – Progress Towards the 21st Century. (Eds.) *J. D.
Hutchins & J. C. Reeves*, pp. 211–218.

Dussart, F., Creissen, H. E., & Havis, N. D. (2020) *Ramularia collo-cygni*-an enemy in waiting. In: *eLS*
(pp. 1-8).

Elnifro Elfath, M., Ashshi Ahmed, M., Cooper Robert, J., & Klapper Paul, E. (2000) Multiplex PCR:
optimization and application in diagnostic virology. *Clinical Microbiology Reviews, 13*(4), 559-
570. https://doi.org/10.1128/cmr.13.4.559

Ficetola, G. F., Pansu, J., Bonin, A., Coissac, E., Giguet-Covex, C., De Barba, M., Gielly, L., Lopes, C. M.,
Boyer, F., & Pompanon, F. (2015) Replication levels, false presences and the estimation of
the presence/absence from eDNA metabarcoding data. *Molecular Ecology Resources, 15*(3),
543-556.

Fountaine, J. M., Shaw, M. W., Ward, E., & Fraaije, B. A. (2010) The role of seeds and airborne
inoculum in the initiation of leaf blotch (*Rhynchosporium secalis*) epidemics in winter barley.
Plant Pathology, 59(2), 330-337. https://doi.org/https://doi.org/10.1111/j.1365-
3059.2009.02213.x

318
319 Garozi, R., Angelotti, P., Feksa, H. R., & Tessmann, D. J. (2020) First report of *Pyrenophora teres* f.
320   teres causing leaf spot on wheat in Brazil. *Plant Disease, 104*(12), 3254-3254.
321   https://doi.org/10.1094/pdis-03-20-0570-pdn

322
323 Havis, N., Pastok, M., Pyzalski, S., Oxley, S., & Heilbronn, T. (2006) Investigating the life cycle of
324   *Ramularia collo-cygni*. Proceedings crop protection in Northern Britain, Dundee,UK

325
326 Hiddink, G. A., Willmann, R., Woudenberg, J. H. C., & Souza-Richards, R. (2023) Seed health testing:
327   doing things right. *PhytoFrontiers™, 3*(1), 71-74. https://doi.org/10.1094/phytofr-03-22-
328   0029-fi

329
330 Hindson, C. M., Chevillet, J. R., Briggs, H. A., Gallichotte, E. N., Ruf, I. K., Hindson, B. J., Vessella, R. L.,
331   & Tewari, M. (2013) Absolute quantification by droplet digital PCR versus analog real-time
332   PCR. *Nature Methods, 10*(10), 1003-1005. https://doi.org/10.1038/nmeth.2633

333
334 Huss, H., Mayrhofer, H., & Iugolič, E. (1992) *Ramularia collo-cygni* Sutton & Waller fungi imperfecti,
335   ein wirtschaftlich bedeutender parasit der gerste in der Steiermark. *Mitteilungen des
336   Naturwissenschaftlichen Vereines für Steiermark*, 122, 87-95.

337
338 Hysing, S.-C., & Wiik, L. (2013) The role of seed infection level and fungicide seed treatments in
339   control of net blotch in barley. *European Journal of Plant Pathology, 137*(1), 169-180.
340   https://doi.org/10.1007/s10658-013-0230-7

341
342 Jalli, M., Kaseva, J., Andersson, B., Ficke, A., Nistrup-Jørgensen, L., Ronis, A., Kaukoranta, T., Ørum, J.-
343   E., & Djurle, A. (2020) Yield increases due to fungicide control of leaf blotch diseases in
344   wheat and barley as a basis for IPM decision-making in the Nordic-Baltic region. *European
345   Journal of Plant Pathology, 158*(2), 315-333. https://doi.org/10.1007/s10658-020-02075-w

346
347 Knight, N. L., Moslemi, A., Begum, F., Dodhia, K. N., Covarelli, L., Hills, A. L., & Lopez-Ruiz, F. J. (2022)
348   Detection of *Ramularia collo-cygni* from barley in Australia using triplex quantitative and
349   droplet digital PCR. *Pest Management Science, 78*(4), 1367-1376.
350   https://doi.org/https://doi.org/10.1002/ps.6753

351
352 Liu, Z., Ellwood, S. R., Oliver, R. P., & Friesen, T. L. (2011) *Pyrenophora teres*: profile of an increasingly
353   damaging barley pathogen. *Molecular plant pathology, 12*(1), 1-19.

354
Lofgren, L. A., Uehling, J. K., Branco, S., Bruns, T. D., Martin, F., & Kennedy, P. G. (2019) Genome-
based estimates of fungal rDNA copy number variation across phylogenetic scales and
ecological lifestyles. *Molecular ecology, 28*(4), 721-730.

Lu, Y., Zhang, H.-j., Zhao, Z.-j., Wen, C.-l., Wu, P., Song, S.-h., Yu, S.-c., Luo, L.-x., & Xu, X.-l. (2020)
Application of droplet digital PCR in detection of seed-transmitted pathogen *Acidovorax*
citrulli. *Journal of Integrative Agriculture, 19*(2), 561-569.
https://doi.org/https://doi.org/10.1016/S2095-3119(19)62673-0

Mair, W. J., Thomas, G. J., Dodhia, K., Hills, A. L., Jayasena, K. W., Ellwood, S. R., Oliver, R. P., &
Lopez-Ruiz, F. J. (2020) Parallel evolution of multiple mechanisms for demethylase inhibitor
fungicide resistance in the barley pathogen *Pyrenophora teres* f. sp. maculata. *Fungal*
Genetics and Biology, 145, 103475.
https://doi.org/https://doi.org/10.1016/j.fgb.2020.103475

Mikhailova, L. A., Ternyuk, I. G., & Mironenko, N. V. (2010) *Pyrenophora teres*, an agent causing
wheat leaf spot. *Microbiology, 79*(4), 561-565. https://doi.org/10.1134/S0026261710040223

Poudel, B., Ellwood, S. R., Testa, A. C., McLean, M., Sutherland, M. W., & Martin, A. (2017) Rare
Pyrenophora teres hybridization events revealed by development of sequence-Specific PCR
markers. *Phytopathology, 107*(7), 878-884. https://doi.org/10.1094/phyto-11-16-0396-r

European Union Crop Production. (2024)
https://ipad.fas.usda.gov/countrysummary/default.aspx?id=E4.

Stam, R., Sghyer, H., Tellier, A., Hess, M., & Hückelhoven, R. (2019) The current epidemic of the
barley pathogen *Ramularia collo-cygni* derives from a population expansion and shows
Global Admixture. *Phytopathology®, 109*(12), 2161-2168. https://doi.org/10.1094/phyto-04-
19-0117-r

Tóth, B., Csősz, M., Kopahnke, D., & Varga, J. (2008) First report on *Pyrenophora teres* causing lesions
of wheat leaves in Hungary. *Plant Pathology, 57*(2), 385-385.
https://doi.org/https://doi.org/10.1111/j.1365-3059.2007.01746.x

Tsedaley, B. (2015) Review on seed health tests and detection methods of seedborne diseases.
Journal of Biology, Agriculture and Healthcare, 5(5).

Turo, C., Mair, W., Martin, A., Ellwood, S., Oliver, R., & Lopez-Ruiz, F. (2021) Species hybridisation
and clonal expansion as a new fungicide resistance evolutionary mechanism *in Pyrenophora*
teres spp. *bioRxiv*, 2021.2007. 2030.454422.

Zhang, X., Ovenden, B., & Milgate, A. (2020) Recent insights into barley and *Rhynchosporium*
commune interactions. *Molecular Plant Pathology, 21*(8), 1111-1128.
https://doi.org/10.1111/mpp.12945

List of Figures

Figure 1

Summarized workflow of the digital droplet PCR assay

Figure 2

Specificity of the multiplex ddPCR assays, pathogen detected are indicated on x-axis (A) Multiplex assay for the detection of *R. graminicola* , *R. collo-cygni* and *P. teres* (B) Multiplex assay for the detection of forms of *P. teres*, FAM channel detects *P. teres* f. teres while HEX channel is to detect *P. teres* f. maculata. Cy5 was used as an internal control.

Figure 3

Linearity of the multiplex assay for the detection of (A) *P. teres,* (B) *R. graminicola* and (C) *R. collo-cygni*.

Figure 4

Selectivity of the multiplex assays (A) Multiplex assay for the detection of *R. collo-cygni*, R. *graminicola* and *P. teres* challenged with various barley pathogens (B) *P. teres* form specific assay exposed to various barley pathogens.

Figure 5

Detection of *R. secalis* (FAM channel) and *R. collo-cygni* (HEX channel) in Rye seed DNA (3 and 5); sample 15 and 17 indicate positive and negative controls for the multiplex assay respectively.

Figure 6

Detection of *R. collo-cygni* from wheat leaf DNA (sample 9, 11 and 13) collected from wheat volunteers in the barley field, sample 16 and 32 indicate positive and negative controls for

434   the multiplex assay. Droplet population in FAM and Cy5 channel indicate the possible cross

435   contamination of the wheat leaves with *R. graminicola* and *P. teres* spores from the barley.

436