Impact of early life antibiotic and probiotic treatment on gut microbiome and resistome of very-low-birth-weight preterm infants

Raymond Kiu[1,2,3], Elizabeth M Darby[1,2#], Cristina Alcon-Giner[3#], Antia Acuna-Gonzalez[3#], Anny Camargo[3,4,5], Lisa E Lamberte[2], Kathleen Sim[6], Alexander G Shaw[6], Paul Clarke[7,8], Willem van Schaik[1,2], J Simon Kroll[6], Lindsay J Hall[1,2,3,8*]

[1] Department of Microbes, Infection and Microbiomes, School of Infection, Inflammation and Immunology, College of Medicine and Health, University of Birmingham, Birmingham, UK
[2] Institute of Microbiology and Infection, University of Birmingham, Birmingham, UK
[3] Food, Microbiome and Health, Quadram Institute Bioscience, Norwich Research Park, Norwich, UK
[4] Centro de Investigaciones en Microbiología y Biotecnología-UR (CIMBIUR), Facultad de Ciencias Naturales, Universidad del Rosario, Bogotá, Colombia
[5] Health Sciences Faculty, Universidad de Boyacá, Tunja, Colombia
[6] Faculty of Medicine, Imperial College London, London, UK
[7] Norfolk and Norwich University Hospital, Norwich, UK
[8] Norwich Medical School, University of East Anglia, Norwich, UK
[#] These authors have contributed equally to this work
* Corresponding authors:
Prof Lindsay J Hall l.hall.3@bham.ac.uk

Abstract
Preterm infants (<37 weeks' gestation) are often administered broad-spectrum antibiotics in hospitals due to their vulnerability to severe morbidity, including necrotising enterocolitis and sepsis. However, antibiotics can disrupt the development of early-life microbiota, potentially impairing gut immunity and colonisation resistance. Evidence shows that probiotics (e.g., certain *Bifidobacterium* strains) may help restore healthy gut microbiota. In this study, we examined the effects of probiotics and antibiotics on the preterm gut microbiome and resistome in two unique cohorts of 34 very-low-birth-weight, human-milk-fed preterm infants (moderate to very preterm), with one cohort receiving probiotics. Within each group, some infants were treated with antibiotics (benzylpenicillin and/or gentamicin) while others served as non-antibiotic treated controls. We performed shotgun metagenomic sequencing on 93 longitudinal faecal samples from 34 infants, generated >300 metagenome-assembled genomes, and obtained ~90 isolate genomes through targeted culturomics, enabling analysis of the microbiome/resistome at species and strain levels. Additionally, we investigated *in vitro* horizontal gene transfer (HGT) capacity of preterm infant-derived multidrug-resistant (MDR) pathogen *Enterococcus* via neonatal gut models. Overall, probiotic supplementation significantly reduced antibiotic resistance gene prevalence, MDR pathogen load, and helped restore a typical early-life microbiota. However, the persistence of MDR pathogens like *Enterococcus*, with high HGT potential, highlights the need for ongoing surveillance in neonatal care. Our findings underscore the complex interactions between antibiotics, probiotics, and HGT in shaping the neonatal microbiome and support further research into probiotics for antimicrobial stewardship in preterm populations.

Introduction

The World Health Organisation (WHO) estimates that over 10% of infants are born prematurely each year worldwide, defined as gestation of <37 weeks[1]. Among newborns, Very Low Birth Weight (VLBW) infants, those born weighing below 1500g, represent around 1.1-1.4%[2,3]. VLBW preterm infants have underdeveloped immune systems, making them particularly susceptible to morbidities such as Necrotising Enterocolitis (NEC)[4,5] and sepsis[6,7], often involving antibiotic-resistant bacteria. Due to these risks, preterm infants frequently reside in Neonatal Intensive Care Unit (NICUs) and are routinely administered broad-spectrum antibiotics, typically benzylpenicillin and gentamicin or derivatives, during their first days and weeks of life[8,9]. However, this early-life antibiotic exposure can disrupt the normal development of the gut microbiota[8,10,11].

The use of antibiotics in preterm infants can also lead to an enrichment of antibiotic resistance genes (ARGs), collectively known as the gut resistome[12]. ARGs within gut bacterial communities can spread rapidly, primarily through horizontal gene transfer (HGT), occurring both within and between species, including from commensals to pathogens or vice versa, and this transfer often occurs via mobile genetic elements such as plasmids[13]. Multidrug-resistant bacteria, including *Staphylococcus*, *Klebsiella*, *Enterococcus* and *Escherichia* are common in the preterm infant gut[5,8,14], with their presence frequently linked to prolonged hospitalisation[15], late-onset bloodstream infections[16], and nosocomial infections in hospital settings[17-19].

In response to these challenges, WHO has recommended probiotic supplementation for very preterm (<32 weeks' gestation), human-milk-fed infants[20]. Probiotics, particularly *Bifidobacterium* spp. and *Lactobacillus* spp., are now increasingly used in NICUs[21], with approximately 40% of NICUs in the UK adopting this practice[8,9]. Probiotic supplementation has been associated with reduced NEC incidence, lower mortality, fewer gut pathogens, and enhanced immune maturation[8,14,21-28]. Importantly, probiotics have also been observed to reduce the abundance of ARGs in the gut microbiota of preterm infants, bringing it closer to levels seen in full-term infants[29-32].

To investigate the impact of probiotics on the preterm gut microbiome and resistome, we studied two cohorts of VLBW preterm infants, all exclusively fed human milk. Using shotgun metagenomics and genome-resolved approaches, we assessed microbiome species and strain dynamics and the effects of probiotics and antibiotics during the first three weeks of life. Additionally, a plasmid transfer experiment using *Enterococcus* in an infant gut model explored *ex vivo* ARG transfer.

Results

We analysed the gut microbiome from 34 VLBW preterm infants (moderate to very preterm), divided into two main cohorts: the Probiotic-Supplemented (PS) cohort and the Non-Probiotic-Supplemented (NPS) cohort. Infants in the PS cohort received probiotics containing *B. bifidum* and *L. acidophilus*, while those in the NPS cohort did not receive any probiotics. These infants were selected as a subset from a larger study[8]. Within each cohort, some infants received empirical antibiotic treatment with benzylpenicillin and gentamicin, while others served as controls without antibiotic exposure (Fig. 1a). Faecal samples were collected weekly from preterm infants during the first three weeks of life, when possible (Table 1). These samples underwent processing, sequencing, and computational analysis to characterise the gut microbiome.

Table 1. Summary of cohort characteristics: NPS cohort vs PS cohort. Cohort characteristics were compared using either Wilcoxon test, Student's t-test or Fisher's exact test where appropriate.

Characteristic	NPS (n=19)	PS (n=15)
Demographics		
Male	6 (32)	10 (67)
Female	13 (68)	5 (33)
Median birth weight (IQR), g	980 (1000-1325)	1382* (1243-1451)
Median gestation at birth (IQR), weeks + days	28+6 (28+2 - 30+1)	30+0 (29+5 – 30+1)
Mode of delivery		
Caesarean section	17 (79)	7 (47)
Normal vaginal delivery	4 (21)	8 (53)
Antibiotic use in the first 3 weeks of life		
Median No. of days of treatment in treated infants (IQR)	3.0 (3.0-5.0)	3.0 (3.0-4.0)
No. of infants without treatment	10 (53)	3 (20)
No. of infants on short-course (1-3 day) treatment	6 (32)	8 (53)
No. of infants on long-course (>3 days) treatment	3 (16)	4 (27)
Diet		
Breastmilk/donor breastmilk (week 1)	19 (100)	15 (100)
Breastmilk/donor breastmilk (week 2)	19 (100)	15 (100)
Breastmilk/donor breastmilk (week 3)	18 (95)	12 (80)
Sample collection		
Median No. of samples collected (per infant)	3.0	3.0
Week 1 median time point, day (IQR), No. of infants (%)	7.0 (7.0-7.8), 18 (95)	6.0* (5.0-7.0), 11 (73)
Week 2 median time point, day (IQR), No. of infants (%)	14.0 (14.0-14.0), 19 (100)	13.0 (11.5-14.5), 15 (100.0)
Week 3 median time point, day (IQR), No. of infants (%)	21.0 (20.8-21.3), 16 (84)	20.0 (20.0-22.0), 13 (87)
Sequencing depth		
Mean No. of sequencing reads	10,578,191	10,869,232
Mean metagenome size (bp)	1,322,088,528	1,358,155,375

Data are presented as No. (%) unless otherwise specified. IQR, interquartile range. * P<0.05.

Probiotics support restoration of the early life gut microbiota
Gut microbiome diversity differed significantly between the NPS and PS cohorts, with the NPS cohort showing an increase in microbiome diversity during weeks 2 and 3, while the PS cohort maintained similar levels of diversity throughout (Fig. 1b). The overall gut microbiome profiles were markedly distinct between the NPS and PS groups (Fig. 1c). The NPS infant gut microbiomes were characterised at the genus level by early-life pathobionts (Fig. 1d), including *Klebsiella*, *Enterobacter*, *Escherichia*, *Enterococcus*, and *Staphylococcus*, which on the species level were identified as *Klebsiella pneumoniae, Klebsiella grimontii*, *Enterobacter hormaechei*, *Escherichia coli*, *Staphylococcus epidermidis*, and *Staphylococcus haemolyticus* (Fig. 1e). In contrast, the gut microbiomes of PS infants were dominated by the genus *Bifidobacterium*, particularly *Bifidobacterium bifidum*, a major component of the Infloran probiotic provided to the infants, followed by *Bifidobacterium breve* and *Bifidobacterium longum* in weeks 2 and 3, reflecting the impact of probiotic supplementation (Fig. 1e). Notably, *B. breve* and *B. longum*, both associated with breastfeeding and recognised for promoting a healthy infant gut by breaking down complex carbohydrates, including human milk oligosaccharides (HMOs), were more abundant in PS infants compared to NPS infants (Fig. 1f).

Comparative analysis showed that *Bifidobacterium* was significantly more abundant in the PS cohort microbiota, while *Enterobacter*, *Escherichia*, and *Klebsiella* were more abundant in NPS infants (Fig. 1g). *Staphylococcus* was notably dominant only during the first week of life and declined in both cohorts by weeks 2 and 3, indicating its transient presence in the

126   infant gut. Importantly, *Enterococcus faecalis*, a frequently multidrug-resistant bacterium,
127   was also a prominent member of the gut microbiota in both cohorts.
128
Next, we examined the functional pathways within the gut microbiomes. Clustering of these
pathways showed increasingly distinct patterns between the PS and NPS cohorts starting
from week 2 (Fig. 1h). Across all samples, a total of 407 functional pathways were identified,
with 84 pathways showing significantly different abundances between the two cohorts (Fig.
1i). Among these, 14 pathways (3.4%) were unique to the PS cohort, including sucrose
degradation pathways, while 370 pathways (90.9%) were shared between both cohorts. In
addition, 23 pathways (5.7%) were exclusively associated with NPS infants.
136
Probiotic intervention is linked to suppression of ARGs in the gut resistome
We investigated the gut resistome in both NPS and PS cohorts, comprising infants treated
empirically with benzylpenicillin and/or gentamicin, to assess the impact of these antibiotics
on early gut microbiomes and resistomes over the first 3 weeks of life (Fig. 2a). Notably, the
abundance of ARGs was significantly higher in NPS infants compared to PS infants across
the first 3 weeks (Fig. 2b). Analysis of ARG diversity, defined by the number of
antibiotic/drug classes represented, showed that NPS infant guts contained significantly more
ARG types than those in the PS cohort, particularly in weeks 2 and 3 (Fig. 2c). Key ARG
types in both cohorts included genes conferring resistance to aminoglycosides, macrolides-
lincosamides-streptogramines (MLS), beta-lactamases, trimethoprim, and tetracycline, while
ARGs conferring resistance to fluoroquinolones and colistin were exclusively found in NPS
infants (Fig. 2d). Clustering analysis of resistome profiles revealed shared ARGs in both
cohorts (48 ARGs), with 46 unique ARGs in NPS infants and 11 in PS infants, though the
separation between groups was not clearly distinct (Fig. 1e, f).
151
To examine the impact of empirical antibiotic therapy on the gut microbiome, we compared
Antibiotic-treated vs Control (non-antibiotic-treated) groups within each cohort. Overall, no
significant difference in microbiome diversity was observed between Antibiotic and Control
groups (Fig. 2g). In the NPS cohort, the Antibiotic group showed higher gut microbiome
diversity from the first week, which continued through weeks 2 and 3, while the Control
group's microbiome diversity increased in later weeks; both groups were dominated by key
pathobionts (Fig. 2h). In the PS cohort, apart from an initial increase in *Staphylococcus* in
Antibiotic-treated infants and *Enterococcus* in Controls, *Bifidobacterium* dominated in weeks
2 and 3. Taxonomic abundance analysis revealed no significant overall differences in
abundance for Antibiotic vs Control in the NPS cohort, except for *Bifidobacterium* and
Lactobacillus (Fig. 2i, top). In the PS cohort, Antibiotic-treated infants had significantly
higher levels of *Cutibacterium*, *Escherichia*, and *Lactobacillus*, with reduced *Enterococcus*
and *Streptococcus* over 3 weeks (Fig. 2i, bottom).
165
To further understand antibiotic impacts on the resistome, we compared ARG profiles in
Antibiotic vs Control groups within each cohort. No significant differences were observed in
ARG counts between these groups in either cohort (Fig. 2j). However, both Antibiotic-treated
and Control infants in the NPS cohort had higher ARG counts than their counterparts in the
PS cohort, suggesting a potential ARG-suppressive effect of *Bifidobacterium bifidum* in the
preterm gut (Fig. 2k).
172
We also assessed whether microbiome diversity correlated with ARG abundance in preterm
infants. Statistical testing indicated a weak, but statistically significant, positive correlation
between increased microbiome diversity and ARG levels (correlation coefficient R=0.25;

P=0.00074; Fig. 2l). Notably, a colistin resistance gene, *mcr-9.1*, was detected with 100%
nucleotide identity in the gut microbiome of a preterm infant around 2011-2012 (Fig. 2m),
predating its identification as a resistance gene in 2019, though it could not be
computationally mapped to a specific bacterial strain in this study.
180
Strain-level gut-derived pathobiont resistome analysis
To examine gut resistomes at the strain level, we applied genome-resolved metagenomics,
recovering 322 high-quality metagenome-assembled genomes (MAGs) from shotgun
metagenome sequences and incorporating 89 novel isolate genomes cultured from preterm
infant faecal samples. Together, these genomes represent 27 genera and 47 species.
Resistance-gene searches and statistical analyses were conducted on these bacterial genomes
to characterise the early-life gut resistomes.
188
Following dereplication of all genomes, including MAGs, at a strain-level cut-off of 99.9%
ANI, we obtained 195 representative genomes for further resistome analysis (Fig. 3a). In
total, 10 ARG types were identified across bacterial strains (Fig. 3a), with *Enterococcus*,
Escherichia, *Klebsiella*, and *Staphylococcus* identified as the top four resistant genera, each
exhibiting a significantly higher ARG count per strain vs the rest (Fig. 3b). Further
examination of these genera showed a trend towards lower ARG counts in *Enterococcus*,
Escherichia, and *Klebsiella* strains within the PS cohort compared to NPS, though
differences were not statistically significant (Fig. 3c).
197
A deeper analysis of multidrug resistance (MDR) efflux pumps across these resistant genera
revealed multiple efflux pumps, with *Escherichia* and *Klebsiella* strains encoding the highest
number (n=13), followed by *Staphylococcus* (n=7) and *Enterococcus* (n=6) (Fig. 3d). ARG
type analysis further demonstrated the widespread presence of beta-lactamase resistance
genes in *Escherichia*, *Klebsiella*, and *Staphylococcus* strains, with aminoglycoside resistance
genes prevalent in NPS *Enterococcus* strains (Fig. 3e, top). Analysis of MDR capacity
showed that *Enterococcus* ranked highest among these resistant genera, with over half of its
strains harbouring ARGs conferring resistance to more than three antibiotic classes, followed
by *Staphylococcus*. Notably, none of the PS-associated *Klebsiella* (n=5) or *Escherichia*
(n=12) strains were MDR, while 47.6% of NPS-associated *Escherichia* (n=31) strains
exhibited MDR characteristics (Fig. 3e, bottom).
To assess clinical relevance, we sequence-typed a total of 193 key resistant pathobiont
genomes from *Escherichia*, *Klebsiella*, *Enterococcus*, and *Staphylococcus* (Fig. 3f) using the
multi-locus sequence typing (MLST)[33] scheme. Dominant sequence types (STs) included
ST1193, ST127, ST394, and ST681 for *E. coli*, while ST432 was prominent for *K.*
pneumoniae. In *Enterococcus*, 12 STs were identified, including six potential novel STs. *S.*
epidermidis displayed 14 STs, with ST32 in 10 genomes and 12 potential novel STs. Of
particular note, key *S. haemolyticus* STs included methicillin-resistant ST1, linked to NICUs
previously[34] and ST49, associated with children and often resistant to methicillin[35,36]. We
also performed surface antigen typing for *Klebsiella*, identifying KL10 (n=10) as the most
common capsule (K) type and O1/O2v1 (n=15) as the most prevalent LPS (O) type in the
Klebsiella genomes (Fig. 3f, right).
Strain-level mobilome analysis and circulation of strains within NICUs
To investigate whether plasmid carriage is linked to the resistome, we performed a plasmid
replicon search across 195 strain-level bacterial genomes (Fig. 4a). Comparison between PS
and NPS cohorts revealed significantly higher plasmid counts in the NPS cohort (Fig. 4b). To

further assess the role of antibiotics in the mobilome, we compared plasmid counts between
Antibiotic-treated and Control groups within both cohorts. Although Antibiotic-treated
strains showed a higher plasmid count, the difference was not statistically significant in either
cohort (Fig. 4c) or in the overall comparison (Fig. 4d).

Next, we examined the correlation between plasmid replicon count and ARG count, finding a
weak positive correlation (R=0.33; P=0.00001; Fig. 4e). Notably, the Antibiotic-treated
group exhibited a significantly higher frequency of potential horizontal gene transfer events
compared to the Control group (Fig. 4f). Among key pathobiont genera, *Enterococcus*,
Escherichia, and *Staphylococcus* were identified as top plasmid carriers, with *Enterococcus*
ranking highest in both ARG and plasmid carriage (Fig. 4g). Given its prominence as an
MDR pathogen and its relative understudy in preterm infants, we focused on tracking
Enterococcus transmission among infants in both cohorts.

Average nucleotide identity (ANI) analysis at a strain-level cut-off of 99.9% indicated 6
Enterococcus strains were circulating among infants, with four pairs of infants (unrelated) in
Hospital A (one of which was a twin) and one pair in Hospital B carrying identical strains
(Fig. 4h). Additionally, in Hospital B, three other unrelated infants were colonised with the
same *Enterococcus* strain, underscoring the potential for nosocomial transmission of this
MDR pathogen.

AMR plasmid transfer between Enterococcus strains occurs within an infant gut model
Given the clinical significance of *Enterococcus faecium* as a multidrug-resistant (MDR)
pathogen and its relatively limited study in infant microbiomes, we hypothesised that ARG-
encoding plasmids could be transferred between *E. faecium* strains within the infant gut. To
test this, we used two donor strains, both originated from preterm infants, and a plasmid-free,
gentamicin-sensitive recipient strain (64/3) in a plasmid transfer experiment (Fig. 5a). We
utilised a gut model using *Enterococcus*-free infant faecal slurry as the culture medium to
simulate the neonatal gut environment (Fig. 4b). Following 24 hours incubation, we isolated
Enterococcus colonies on selective media and performed WGS. Genome analysis revealed
the successful transfer of a ~137kb mega-plasmid encoding the aminoglycoside resistance
gene *aac6-aph2* to plasmid-free recipient strain 64/3 (transformants D1 and D3), thereby
conferring gentamicin resistance to these strains (Fig. 5b, 5c and 5d).

Discussion
In this study, we used genome-resolved metagenomics to analyse the resistome and
microbiome of preterm infants during their first 3 weeks of life in neonatal units, highlighting
the complex effects of antibiotics, probiotics, and HGT on the gut microbiome. Our findings
suggest that probiotic supplementation with *B. bifidum* and *L. acidophilus* not only supports
beneficial microbial communities but also plays a role in reducing MDR bacteria and overall
ARG carriage.
261
One of the strengths of this study is the inclusion of both PS and NPS infants who did not
receive antibiotics, which is uncommon considering the high antibiotic prescription rates
among these at-risk patients (55-87% for very to extremely preterm infants in the UK)[37],
allowing us to begin disentangling the specific effects of probiotics and antibiotics on the
preterm gut microbiome and resistome. This carefully selected sub-cohort (n=34) from a
larger observational study (n=234) consists exclusively of preterm infants (<33 weeks'
gestation) who were matched for diet and birth mode, providing a relatively uniform
baseline[8]. Moreover, in cases where empirical antibiotic therapy was administered, the

duration and types of antibiotics (benzylpenicillin and/or gentamicin, the most frequently
prescribed antibiotics in UK NICUs[37]) were strictly limited and standardised to less than 1
week. Despite this exposure, we observed minimal impact on overall microbiome diversity
within both PS and NPS cohorts, suggesting that early-life antibiotic exposure over a short
period may not have an immediate effect on preterm microbiome diversity. However, it is
possible that the limited 3-week sampling window did not capture potential longer-term shifts
induced by antibiotic exposure. This observation aligns with previous studies that reported
minimal immediate impact of antibiotics on early microbiome composition, while indicating
that more pronounced changes may emerge over extended periods[38].
279
However, the administration of antibiotics, though limited to an average duration of median 3
days, did specifically impact microbiome composition by promoting the proliferation of
certain genera with pathogenic potential, including *Klebsiella* and *Enterococcus*, in the first
week, with some of these strains persisting into the second week. *Enterobacter*, a genus less
associated with multiple ARGs in this study, was observed to decline in the NPS cohort after
antibiotic exposure, indicating potential susceptibility to the antibiotic regimen. In contrast,
Bifidobacterium - particularly in the PS cohort - appeared to exert a protective effect, as
evidenced by a marked reduction in *Enterococcus* abundance. This decrease could potentially
be attributed to the carbon source depletion promoted by *Bifidobacterium* specialised HMO
utilisation in human milk-fed gut environment, and reduced niche pH, which may inhibit
pathobiont colonisation and growth[39,40].
291
Overall, we identified a diverse range of ARGs in the gut microbiomes of preterm infants,
with key ARG classes including genes encoding resistance to aminoglycosides, macrolides-
lincosamides-streptogramins (MLS), beta-lactamases, trimethoprim, and tetracyclines. These
ARG classes are clinically relevant in NICUs, where aminoglycosides and beta-lactams, e.g.
gentamicin and benzylpenicillin, are frequently empirically administered from birth to cover
possible infection[8,37]. The presence of these ARGs, underscores the potential for reduced
antibiotic efficacy in this vulnerable population and highlights the need for careful and
improved antibiotic stewardship in NICU settings. Efflux pumps, which facilitate multidrug
resistance by expelling a broad range of antibiotics[41], were also prevalent across multiple
strains, particularly within *Escherichia* and *Klebsiella*, further complicating treatment options
for infections by these pathogens in preterm infants. Notably, we detected an *mcr-9.1* gene,
conferring resistance to colistin - a last-resort antibiotic -within the gut microbiome of a NPS
infant. This gene was identical to a homolog found in *E. coli* strain 68A though it could not
be definitively linked to a specific bacterial species in our cohort due to limitations in short-
read sequencing data, as it appeared only as a single contig (Fig. 2m). The *mcr-9.1* gene was
first identified in *Salmonella enterica* in 2019[42], where it conferred phenotypic resistance to
colistin, and has since been associated with *Enterobacter* and *Klebsiella* species, often on
mobile IncHI2 plasmids[43]. The presence of *mcr-9.1* in our cohort raises concerns about the
silent circulation of colistin resistance genes within the gut microbiome of preterm infants,
which may complicate future treatment options, particularly in cases where conventional
antibiotics fail.
313
We also observed an increase in ARG abundance over time in NPS cohort, indicating that
ARG acquisition may be driven by the introduction of new bacteria, likely hospital-acquired,
rather than ARG evolution within the closed ecological system of the infant gut (at least
across the short sampling widow). This is further supported by our strain-level analyses,
where we observed the persistence of certain pathobionts, such as *Enterococcus*, across
multiple time points. This persistence, along with the detection of shared strains among

infants within the same hospital, suggests a high likelihood of nosocomial transmission of
MDR bacteria. These findings align with reports from other NICU studies, where
Enterococcus, *Escherichia*, and *Klebsiella* are common nosocomial pathogens and are often
implicated in late-onset infections in preterm infants[32,44,45]. Indeed, we identified specific STs
linked to neonatal infections, such as *E. coli* ST1193[46], ST73 and ST95[46,47], and *K.*
pneumoniae ST432[48] strains, which are typically drug-resistant strains and of clinical
significance. *Enterococcus*, frequently identified as a prominent ARG carrier in our cohort,
exhibited multiple STs, including novel ones not previously reported, and certain strains were
shared across different individual infants within the same hospital. This sharing among
unrelated infants further supports the notion of nosocomial transmission of *Enterococcus*, a
common cause of late-onset infections in NICUs. Thus, the frequent colonisation with
hospital-acquired, ARG-bearing bacteria in the absence of probiotics underscores the
importance of continuous surveillance and infection control in NICUs.

Our findings on the high prevalence of strains carrying plasmids, and in particular MDR
plasmids, is particularly concerning, as it indicates that these strains could serve as reservoirs
for resistance genes that may spread to other non-MDR strains within the same species as
well as other genera via HGT. Notably, our findings indicate that antibiotic treatment
promotes a higher prevalence of LGT events within the neonatal gut, likely due to the
selective pressure antibiotics place on microbial populations. This pressure appears to favour
the retention and spread of mobile genetic elements carrying ARGs, increasing HGT
dynamics in antibiotic-treated infants. Indeed, our plasmid transfer experiment within *E.*
faecium further emphasises the significance of plasmid-mediated ARG transfer in the
neonatal gut as we demonstrated the successful transfer of a ~137kb mega-plasmid carrying
an aminoglycoside resistance gene (*aac6-aph2*) to a plasmid-free, gentamicin-sensitive
recipient strain, conferring phenotypic gentamicin resistance. It is important to note that our
genomic analysis indicated only a weak correlation between overall ARG abundance and
plasmid replicon count, suggesting that plasmids are not the sole contributors to ARG
dissemination, and alternative routes of HGT are likely driving ARG spread[49].

The high-resolution sequencing data in this study underscored the beneficial impact of
probiotic supplementation on microbiome development in preterm infants. PS infants showed
an earlier and more robust colonisation of infant-associated *Bifidobacterium* species,
particularly *B. breve* and *B. longum* subsp. *infantis*, compared to NPS infants[50-52]. The
probiotic *B. bifidum* likely facilitated the establishment of these beneficial species through
cross-feeding on HMOs[53,54], promoting a microbiome composition more similar to that of
term infants. Indeed, this early and abundant presence of multiple *Bifidobacterium* species
appeared to enhance pathogen colonisation resistance significantly, as evidenced by the
limited presence of pathobionts, such as *Klebsiella* and *E. coli*, within the PS cohort.

Consistent with previous studies, our results also showed that PS infants harboured
significantly fewer ARGs than NPS infants[29]. *Bifidobacterium* supplementation has been
associated with lower ARG abundance in the infant gut microbiome, likely due to the
competitive exclusion of ARG-rich pathobionts[55,56]. In our study, the higher abundance of
Bifidobacterium in the PS cohort correlated with a reduced ARG load, particularly in genera
commonly associated with MDR pathogens, such as *Enterococcus*, *Escherichia*, and
Klebsiella. Notably, *Escherichia* and *Klebsiella* genomes in PS infants did not exhibit MDR
characteristics, highlighting the potential for probiotic supplementation to mitigate MDR
pathogen prevalence in the preterm gut. However, while plasmid analysis revealed a greater
number of plasmids in the NPS cohort compared to PS infants, suggesting that probiotics

may help reduce the plasmid pool carrying ARGs and, by extension, limit the ARG reservoir,
we still observed plasmid transfer *in vitro* even in the presence of high levels of
Bifidobacterium, suggesting that presence of beneficial bacteria may not fully inhibit HGT.
This observation underscores a critical point: while probiotics such as *Bifidobacterium*
support beneficial microbiome composition and may suppress ARG-rich pathobionts, they do
not necessarily prevent HGT or plasmid transfer, particularly in the antibiotic-exposed gut
where HGT may be more common. This emphasises the need for further studies to evaluate
the role of probiotics not only in microbial colonisation but also in their potential impact on
HGT dynamics, particularly in environments where antibiotics are heavily used.

Finally, one limitation of our study was the relatively short sampling period, which cannot
capture longer-term impacts of antibiotics and probiotics on microbiome development.
Additionally, the study involved samples from two separate hospital sites, which allowed us
to examine baseline and probiotic-supplemented conditions but inevitably introduces
potential confounders from local variability in practices. Future studies involving larger
sample sizes, extended sampling windows, and multi-site designs could provide more
comprehensive insights into the dynamic interactions between antibiotics, probiotics, and the
neonatal microbiome.

In conclusion, our investigation provides a comprehensive overview of the preterm gut
resistome and microbiome, demonstrating the role of probiotic supplementation in reducing
ARG prevalence and pathogen load. However, the persistence of MDR pathogens such as
Enterococcus, with its high plasmid carriage and demonstrated capacity for ARG transfer,
even in a *Bifidobacterium*-dominant ecosystem, underscores the need for continued
surveillance and targeted intervention strategies in NICUs to minimise the risk of
colonisation and subsequent infections by MDR bacteria. Our findings highlight the complex
interplay between antibiotics, probiotics, and HGT in shaping the neonatal microbiome, and
provide a platform for further research into the role of probiotics in antimicrobial stewardship
and infection control in vulnerable preterm populations.
394

Methods

Cohort and sample selection

395
396
A sub-set of 93 samples in total were selected from a previously published observational
cohort study[8]. All NICUs in this study applied similar protocols for feeding, prescription of
antibiotics and the use of prophylactic antifungal drugs. Hospital for PS cohort routinely
prescribed all VLBW infants with oral probiotic supplementation (Infloran®, Desma
Healthcare, Switzerland) twice daily (from birth until ~34 weeks post-menstrual age), while
infants in NPS cohort were not supplemented with probiotics. Probiotic supplements
contained *Bifidobacterium bifidum* (1×10^9 cfu) and *Lactobacillus acidophilus* (1×10^9 cfu).
Samples from both cohorts (NPS and PS) were selected for this study based on the following
criteria: VLBW preterm infants with gestational age < 34 weeks, infants were solely given
antibiotics benzylpenicillin and/or gentamicin, and age-matched infants without antibiotics,
infants were all fed with breastmilk and/or donor breastmilk in both cohorts. A total of 34
VLBW preterm infants were selected for this study. Longitudinally collected faecal samples
at week 1, week 2 and week 3 (first 3 weeks of their NICU stay) were recruited into this
study (Supplementary Table 1).
411

412 *Cohort characteristics*
413 This study is a 'controlled' sub-study from the original observational longitudinal study[8] as
414 published previously. Summary of the cohort characteristics are presented in **Table 1**. Faecal
415 slurry for the plasmid transfer experiment was originally from five infants recruited in
416 PEARL study[57].
417
Ethical approval
Faecal sample collection from Norfolk and Norwich University Hospital (BAMBI study) was
approved by the Faculty of Medical and Health Sciences Ethics Committee at the University
of East Anglia (UEA), and followed protocols laid out by the UEA Biorepository (License
no: 11208). Faecal sample collection Imperial Healthcare NICUs (NeoM study) was
approved by West London Research Ethics Committee (REC) under the REC approval
reference number 10/H0711/39. In all cases, medical doctors and nurses recruited infants
after parents gave written consent. The PEARL study has been reviewed and agreed by the
Human Research Governance Committee at the Quadram Institute Bioscience and the
London-Dulwich Research Ethics Committee (reference 18/LO/1703) and received written
ethical approval by the Human Research Authority. IRAS project ID number 241880[57].
429
Genomic DNA extraction and Shotgun Metagenomic Sequencing
FastDNA Spin Kit for Soil (MP Biomedicals) was utilised to extract genomic DNA from
infant faeces following manufacturer instructions, with an extended 3 min bead-beating on
FastPrep tissue homogeniser (MP Biomedicals), additionally DNA was eluted with 55°C
sterile pure water. Genomic DNA concentration quantification followed using a Qubit 2.0
fluorometer (Invitrogen). DNA samples were then subject to established Illumina paired-end
library preparation prior to Shotgun Metagenomic Sequencing on Illumina HiSeq 2500 to
generate 125-bp paired-end reads (FASTQ), which was performed at the Wellcome Trust
Sanger Institute (Supplementary Table 2).
439
Bacterial isolation work, DNA extraction
A targeted bacterial isolation work was carried out to isolate the most abundant taxa present
in the preterm infant faecal samples – *Bifidobacterium*, *Enterococcus*, *Staphylococcus*,
Klebsiella and *Escherichia* genera. Briefly, 25-50 mg of faecal samples were homogenised in
5 ml of phosphate buffer saline (PBS) by vortexing. Homogenates were then serially diluted
to 10^{-4} in PBS and aliquots of 100 ml each were spread plated on selective agar respectively:
MacConkey (Oxoid; targeting Gram-negative bacteria including *Escherichia* and *Klebsiella*),
De Man-Rogosa-Sharpe (MRS; Difco) with 50 mg/L mupirocin (Oxoid; targeting
Bifidobacterium), Baird-Parker (Oxoid; targeting *Staphylococcus*) and Slanetz and Bartley
(Oxoid; targeting *Enterococcus*). Agar plates were then incubated both aerobically
(MacConkey, Baird-Parke, and Slanetz and Bartley) and anaerobically (MRS agar only) at
37°C for 3 days. Next, 5 colonies from each agar plate were picked and were re-streaked for
3 consecutive times onto newly prepared agar plates to obtain pure isolates.
453
Bacterial isolates were cultured in appropriate media for overnight prior to genomic DNA
extraction. DNA extraction was performed using the phenol-chloroform extraction method as
described previously[58]. Briefly, PBS-washed bacterial cell pellets were resuspended in 2 ml
of 25% sucrose in 10 mM Tris and 1 mM EDTA at pH 8.0, followed by enzymatic lysing
step using 50 µl of lysozyme at 100 mg ml^{-1} (Roche). Next, 100 µl of Proteinase K at 20 mg
ml^{-1} (Roche), 30 µl of RNase A at 10 mg ml^{-1} (Roche), 400 µl of 0.5 M EDTA at pH 8.0 and
250 µl of 10% Sarkosyl NL30 (Thermo Fisher Scientific) were added sequentially into the
lysed suspension. The suspension was then subjected to an 1 h incubation on ice and then a

50°C water bath for overnight. On the next day, 3 rounds of phenol-chloroform-isoamyl
alcohol (Merck) extraction were performed using 15 ml gel-lock tubes (QIAGEN), followed
by chloroform-isoamyl alcohol (Merck) extraction prior to ethanol precipitation step and 70%
ethanol cell wash for 10 min (twice). The DNA pellets were then air dried overnight, re-
suspended in 300 μl of sterile pure water and stored in -80°C freezer prior to further analyses.
467
Bacterial pure isolate Whole Genome Sequencing (WGS) and genome assembly
WGS of each pure bacterial isolate was performed on Illumina NextSeq 500 at the Quadram
Institute (Norwich, UK) to generate 151-bp paired-end reads. Raw sequence reads (FASTQ)
were firstly quality filtered (-q 20) with fastp v0.20.0[59] before *de novo* genome assembly via
genome assembler optimiser Unicycler v0.4.9b[60] at default parameters that involved *de novo*
genome assembler SPAdes v3.11.1[61], also Bowtie2 v2.3.4.1[62], SAMtools v1.7[63] and sequence
polishing software Pilon v1.22[64]. Contigs with lengths <500 bp were discarded from each
draft genome assembly before subsequent analysis. All draft genome assemblies underwent
sequence contamination check via CheckM v1.1.3[65], and contaminated (sequence
contamination >5%) and/or incomplete (genome completeness <90%) genome assemblies
were excluded from further analysis (n=1) resulting in a total of 89 high quality pure isolate
genomes subject to subsequent genome analyses. Taxonomic assignment (species level
identification) was performed using gtdb-tk v1.5.1[66]. Genome assembly statistics were
generated via sequence-stats v1.0[67] and genome performed using Prokka v1.14.6[68]
(Supplementary Table 3).
483
Recovery of Metagenome-Assembled Genomes
Shotgun metagenome raw reads (FASTQ) were firstly trimmed, adaptor-removed and
quality-filtered using fastp v0.20.0[59] (-q 20). Subsequently, host-associated sequences were
removed via KneadData v0.10.0 with human genome (GRCh38.p13 - no ALT) bowtie2 index
file retrieved from https://benlangmead.github.io/aws-indexes/bowtie (with options --bypass-
trim and –reorder) to generate clean FASTQ reads that consist of purely bacterial sequences.
These metagenome reads were then co-assembled with MEGAHIT v1.2.9[69] prior to the
reconstruction of Metagenome-Assembled Genomes (MAGs). Next, MetaWRAP v1.3.2[70]
was utilised to extract MAGs based on metagenome co-assemblies generated and
metagenome clean reads via three binning software included in the pipeline MetaBAT
v2.12.1[71], MaxBin v2.2.6[72] and CONCOCT v1.1.0[73] using sub-module *metawrap binning*.
MAGs were then refined using sub-module *metawrap bin_refinement* to select the high-
quality bins (MAGs) from each sample with completeness >90% and contamination <5% via
checkm v1.1.3[65]. All MAGs were taxonomically ranked using gtdb-tk v1.5.1[66] via module
gtdbtk classify_wf. A total of 411 high-quality (completeness >90%, contamination <5%)
MAGs (n=322) and isolate genomes (n=89) were recovered for subsequent analyses. Next,
all MAGs and isolate genomes were dereplicated using dRep v3.2.2[74] at ANI 99.9% as the
strain-level inference cut-off (n=195).
502
Taxonomic profiling for faecal metagenomes
After quality filtering steps as described above, briefly, shotgun metagenome raw reads
(FASTQ) were quality-filtered using fastp v0.20.0[59] (-q 20) and host-associated sequences
were removed using KneadData; Kraken v2.1.2[75] was utilised for taxonomic assignment for
purified metagenome reads (Kraken2 standard Refseq indexes retrieved from
https://benlangmead.github.io/aws-indexes/k2, May 2021), with confidence level set at 0.1.
Bracken v2.6.2[76] was then deployed to re-estimate the relative abundance of taxa at both
genus and species level (-t set at 10 as recommended to reduce false positive) from Kraken2
outputs as recommended. To simplify data for visualisation purpose, minimal genera with

<1000 reads across all samples were removed prior to further analysis (with average 520,000
resultant genera reads across all samples). Minor genera with <2% (relative abundance)
across all samples were classified as 'Others' to simplify representations prior to data
visualisation in R[77] using library ggplot2[78].
516

517   *Phylogenetic relatedness estimation*
518   The Mash-distance sequence tree consisted of 411 genomes (including MAGs) was generated
519   using Mashtree v1.2.0[79] with 100 bootstrap replicates and option --mindepth 0, as in other
520   cross-genus/species trees in this study. Distance tree was then mid-point rooted and
521   visualised in iTOL v6[80].
522

Metagenomic functional profiling
Filtered shotgun metagenomic raw reads as described in previous method section (using fastp
v0.20.0) were concatenated into a single FASTQ file per sample prior to running Humann
v3.0.0 and Metaphlan v3.0.13 based on CHOCOPhlAn_201901 metaphlan database (built
using bowtie2) retrieved from https://zenodo.org/record/3957592#.YSZDVS1Q3_o. Path
abundance output files were then normalised and used for visualisation of microbiome
functionalities.
530

Antibiotic resistance genes, Multi-Locus Sequencing Typing (MLST) and plasmid replicons
Antibiotic resistance genes (ARGs) sequence search on all sequence data was performed via
ABRicate v1.0.1 with options --minid=95 and --mincov=90 based on nucleotide sequence
database ARG-ANNOT[81] NT v6 (retrieved from https://www.mediterranee-
infection.com/acces-ressources/base-de-donnees/arg-annot-2/), individually validated against
ResFinder v4.0[82] (only acquired ARGs) databases. The resistome analysis of infant gut
metagenome was carried out via sequence search on metagenome co-assemblies of each
sample, whilst strain-level resistome analysis was performed on individual genome
assemblies. Antibiotic classes were sorted using outputs from ABRicate v1.0.1 based on
antibiotic classes specified in ARG-ANNOT. Multidrug resistance efflux pumps were
predicted using METABOLIC v4.0[83].
542

543   MLST was predicted via mlst v2.19.0[84] at default parameters through scanning contig files
544   against the PubMLST[85] typing schemes (https://pubmlst.org/) sited at the University of
545   Oxford.
546

547   Plasmid replicons were predicted using ABRicate v1.0.1[86] via PlasmidFinder sequence
548   database v2[87] with options --minid=90 and --mincov=90 on individual strain-level genome
549   assemblies. Plasmid replicons were reclassified in two key groups - *inc* families and *rep*
550   groups, respectively prior to visualisation using iTOL v6[80].
551

552   *Klebsiella surface antigen typing*
553   Kleborate v2.3.2[88] was invoked with the –all flag to type the capsule (K) and LPS (O) antigen
554   loci, using default parameters.
555

Horizontal gene transfer analysis
Waafle v1.0[89] was used with default parameters to find incidences of horizontal gene transfer
in the dereplicated MAGs and whole genome sequences. To determine statistical significance
between the antibiotic positive (n=237) and antibiotic negative (n=174) cohorts, a random
permutation test was conducted to normalise the sample sizes. Here, the number of LGT
events predicted by Waafle from either population were randomly sub-sampled 100 times,

and these values were then subject to a Mann Whitney statistical test, and averaged, using
Graphpad Prism v10.
564
Enterococcus-associated plasmid transfer study via ex vivo colon model
Enterococcus faecium ARMA59 and ARMA73 were isolated from preterm infant stool
samples that were part of the Baby-Associated Microbiota of the Intestine (BAMBI) study.
Isolates were selectively cultured from stool samples using Slanetz and Bartley Medium
(Oxoid), on which *Enterococcus* forms dark red colonies. Colonies were then streaked on
Brain Heart Infusion agar (Merck) before being sent for whole genome sequencing. *E.*
faecium 64/3, a plasmid-free derivative exhibiting high-level resistance to both rifampicin
and fusidic acid, was utilized as the recipient strain for conjugation assays. This strain was
originally isolated from a stool sample of a hospital patient[90].
574
575   *Enterococcus faecium* ARMA59 and ARMA73 were used as donor strains in the *ex vivo* gut
576   model experiment, while *Enterococcus* 64/3 was used as a recipient strain. Both ARMA59
577   and ARMA73 were confirmed for their gentamicin resistance (>130µg/ml) prior to the key
578   experiment, while recipient strains 64/3 was phenotypically resistant to rifampicin (>25
579   µg/ml) and fusidic acid (>25 µg/ml).
580
Next, faecal slurry was prepared using *Enterococcus*-negative infant stool samples
determined by selective agar plates (5 infants from PEARL study). Stool samples (1,965 mg)
were mixed with 7 ml of reduced PBS to constitute the faecal slurry for the colon model
experiment. Colon media were formulated based on our previous study with added vitamin
solutions[91]. Briefly, 6 different media, M1 (500 ml), M2 (100 ml), M3 (100 ml), M4 (100
ml), M5 (200 ml), M6 (200 ml) were mixed aseptically in a sterile container. In addition, 200
ml of milk and vitamin mix (50 ml; pantothenate 10 mg/L, nicotinamide 5mg/L, thiamine
4mg/L, biotin 2mg/L, vitamin B12 0.5mg/L, menadione, 1mg/L and p-aminobenzoic acid
5mg/L) were also added, making the total volume 2000ml, and this was adjusted to a
maximum 700 ml final total volume (for each of the media or additives). These colon media
were then pH-adjusted to pH6.8.
592
For bacterial strain preparation, frozen bacterial stocks were plated on Brain Heart Infusion
(BHI) agar plates for 24 h anaerobically at 37°C, followed by inoculating single colonies in
5ml BHI broth for overnight (16h) anaerobically at 37°C. Next, the cultures were
homogenised and mixed (200 µl) with 10 ml of colon media (equivalent to 1:50 dilution),
incubated at 37°C anaerobically overnight. The cultures were next washed (PBS + 3%
cysteine) twice and pelleted.
599
Prior to experiments, faecal slurry was determined to be *Enterococcus*-free by both culturing
(on Slanetz and Bartley Agar) and Whole Genome Sequencing approaches. DNA extraction
of faecal slurry was performed via FastDNA Spin Kit for Soil (MP Biomedicals). Genomic
DNA was then subjected to Whole Genome Sequencing via Illumina NovaSeq X to generate
151-bp paired-end reads. Raw sequencing reads (FASTQ) was processed as mentioned in
previous section on metagenomic sequence analysis. Briefly, raw sequences were firstly
quality-filtered with fastp v0.20.0 prior to taxonomic classification of sequence reads using
Kraken v2.1.2[75] (Kraken2 standardDB; --confidence 0.1) and further classified by Bracken
v2.8[76] (-t 10).
609
To set up the *ex vivo* colon model, the micro-Matrix fermentation system (Applikon
Biotechnology) was used to model the human distal colon[92]. This system utilised a 24-well

plate for batch fermentation. Four conditions were set up in the 24-well plate: 1) no donor, no
recipient (control), 2) donor, no recipient, 3) no donor, recipient only, 4) donor and recipient
strains. We have used a total volume of 5 ml in each well (slurry 26.6μl, donor strain
inoculum 100μl, recipient strain inoculum 200μl, colon media negative control 100/200μl
depending, colon media 4.67ml). The basic parameters of the colon model was set at pH6.8,
temperature 37°C, and duration of experiment at 24 h. Samples were collected at 0h and 24h.
618
After the experiments, samples from 4 conditions then were re-streaked on Slanetz-Bartley
agar plates supplemented or not with appropriate antibiotics: 1) no antibiotics, 2) 130 µg/ml
gentamicin, 3) 25 µg/ml rifampicin + 25 µg/ml fusidic acid, 4) 130 µg/ml gentamicin + 25
µg/ml rifampicin + 25 µg/ml fusidic acid, to allow the selection of donor and recipient
strains, and transformants. After 48h anaerobic incubation at 37°C, pure colonies were
selected and cultured in media broth to obtain bacterial pellets sufficient for DNA extraction.
Genomic DNA extraction of each isolate was performed via FastDNA Spin Kit for Soil (MP
Biomedicals) as per manufacturer's instructions. Well contents from 0h were also sampled
and DNA extracted with FastDNA Spin Kit for Soil (MP Biomedicals) for shotgun
metagenome sequencing to confirm the absence of *Enterococcus*.
629
Genomic DNA was whole-genome sequenced on the Nanopore MinION platform (R10.4.1
flowcell) to generate long-read raw sequences with basecalling via Dorado[93]. Raw sequences
were filtered using filtlong v0.2.1[94] to remove reads with <1000 bp. Filtered reads were then
assembled via Flye v2.9[95] (--nano-hq –scaffold -g 3.6m). Genome assemblies were
subsequently polished using Medaka v1.11.3[96] (*medaka_consensus* -m
r1041_e82_400bps_sup_v4.3.0) to generate high-quality genome assemblies for subsequent
analyses.
637
638   Isolate DNA was also whole-genome sequenced on Illumina NextSeq 2000 to generate 151-
639   bp paired-end short reads (FASTQ) in parallel with long-read sequencing as described in
640   previous paragraph. Raw reads were firstly trimmed with fastp v0.20.0[59] (-q 20) prior to de
641   novo genome assembly via genome assembler optimiser Unicycler v0.5.0[60] at default
642   parameters to generate draft genome assemblies for further analyses.
643
Average Nucleotide Identity (ANI) estimation and bacterial strain-level identification
ANI was computed using fastANI v1.34[97] at default parameters to compare genomes. Strain-
level ANI cut-off was set at 99.9%, species-level cut-off was 95% (species boundary). To
classify bacterial genomes at strain-level, dRep v3.2.2[74] (*dRep dereplicate --*
ignoreGenomeQuality) was used at 99.9% ANI (-sa 0.999) to separate highly similar
genomes.
650
Data visualisation
Various bar plots, box plots, dot plots, pie charts and scatter plots were graphed via R
v4.1.2[77], using R libraries tidyverse v1.3.1[98], ggplot2 v3.3.5[78] and ggpubr v0.6.0[99]. R library
vegan v2.6.2[100] was used to construct non-metric multidimensional scaling (NMDS) plot for
gut microbiome (taxonomic/functional profiling) data. Genomic region of contigs/gene
comparison was performed via R library genoplotr v0.8.11[101] or GenoFig v1.1.1[102].
657
Statistics
Statistical tests were performed via R base packages stats v4.1.2[77], including Wilcoxon's test
and Shapiro–Wilk normality test which was used to formally test for data normality where
appropriate. Spearman correlation tests were carried out via R library ggpubr v0.6.0[99]

function ggscatterhist. LEfSe[103] was used to perform linear discriminant analysis (LDA) on
functional profiling output data from Humann3[104]. Shannon diversity index of metagenome
data was estimated via R library vegan v2.6.2[100]. Genome statistics were generated via
sequence-stats v1.0[67]. R library dplyr v1.0.2[105] was used frequently for data handling. Cohort
characteristics were compared using either Wilcoxon test, Student's t-test or Fisher's exact
test where appropriate, via R packages rstatix v0.6.0[106].
668
Data and code availability
Infant faecal sample metagenome sequencing raw reads are publicly available in the NCBI
Sequence Read Archive (SRA) under accession no. PRJNA1191223. Sequencing raw reads
and draft genome assemblies for 89 isolates generated in the present study are made available
in the NCBI SRA and GenBank respectively, under accession no. PRJNA119225.
Sequencing raw reads and draft genome assemblies from long-read WGS on *Enterococcus*
plasmid transfer study are publicly available in SRA (for raw sequencing reads) and
GenBank (genome assemblies) respectively under accession no. PRJNA119226. All 322
metagenome-assembled genomes recovered from gut metagenome sample in this study and
associated R scripts (data visualisation and statistical analysis) are available and shared via
GitHub repository: https://github.com/raymondkiu/Infant-Resistome-Study
680

Acknowledgements
This research was supported in part by the NBI Research Computing through the provision of
a high-performance computing cluster. We would also like to thank the DNA sequencing
team at Wellcome Trust Sanger Institute for performing genomic DNA sequencing for this
study. This work was funded via a Wellcome Trust Investigator Award to L.J.H.
(100974/C/13/Z) and support of the BBSRC Norwich Research Park Bioscience Doctoral
Training Grant (BB/M011216/1; supervisor, L.J.H.; student, C.A.G.), Institute Strategic
Programme (ISP) grant for Gut Health and Food Safety, BB/J004529/1 (L.J.H.), and ISP
grant for Gut Microbes and Health BB/R012490/1 and its constituent project(s),
BBS/E/F/000PR10353 and BBS/E/F/000PR10355 to L.J.H.; L.E.L., A.A.G., W.v.S and
L.J.H were also supported by BBSRC grant BB/S017941/1. Work at Imperial Healthcare
NICUs was supported by a programme grant from the Winnicott Foundation to J.S.K. and the
National Institute for Health Research (NIHR) Biomedical Research Centre based at Imperial
Healthcare NHS Trust and Imperial College London. K.S. was funded by an NIHR Doctoral
Research Fellowship (NIHR-DRF-2011-04-128). We sincerely thank all clinical nurses at
NNUH and Imperial Healthcare NICUs for collecting stool samples. We would like to give a
special mention to research nurses Karen Few, Hayley Aylmer, and Zoe McClure for
obtaining consent from parents and collecting samples.
699

Author contributions
R.K., C.A. and L.J.H. conceived the study. R.K., A.A.G., L.E.L. and C.A. provided the
methodology. R.K. and E.M.D. provided the software. L.J.H. and E.M.D. validated the study.
R.K., E.M.D., A.A.G., L.E.L., A.C. and C.A. did the formal analysis. R.K., E.M.D., A.A.G.,
C.A. and A.C. carried out the investigations. K.S., L.E.L., A.G.S., P.C. and J.S.K. provided
the resources. R.K., C.A. and K.S. curated the data. R.K. and L.J.H. wrote the original draft
of the manuscript. W.vS., J.S.K. and L.J.H. supervised the study. R.K., A.A.G., W.vS.,
J.S.K., P.C. and L.J.H. reviewed and edited the manuscript. R.K. and C.A. administered the
project. J.S.K. and L.J.H. acquired funds.
709

Competing interests

The authors declare no competing interests.

References

Ohuma, E. O. *et al.* National, regional, and global estimates of preterm birth in 2020, with trends from 2010: a systematic analysis. *Lancet* **402**, 1261-1271 (2023). https://doi.org/10.1016/S0140-6736(23)00878-4

Victora, J. D. *et al.* Prevalence, mortality and risk factors associated with very low birth weight preterm infants: an analysis of 33 years. *J Pediatr (Rio J)* **96**, 327-332 (2020). https://doi.org/10.1016/j.jped.2018.10.011

Martin, J. A., Hamilton, B. E. & Osterman, M. J. K. Births in the United States, 2023. *NCHS Data Brief* (2024). https://doi.org/10.15620/cdc/158789

Neu, J. & Walker, A. W. Necrotizing enterocolitis. *The New England Journal of Medicine* **364**, 255-264 (2011).

Sim, K. *et al.* Dysbiosis anticipating necrotizing enterocolitis in very premature infants. *Clinical infectious diseases : an official publication of the Infectious Diseases Society of America* **60**, 389-397 (2015). https://doi.org/10.1093/cid/ciu822

Stoll, B. J. *et al.* Changes in pathogens causing early-onset sepsis in very-low-birth-weight infants. *N Engl J Med* **347**, 240-247 (2002). https://doi.org/10.1056/NEJMoa012657

Hornik, C. P. *et al.* Early and late onset sepsis in very-low-birth-weight infants from a large group of neonatal intensive care units. *Early human development* **88 Suppl 2**, S69-74 (2012). https://doi.org/10.1016/S0378-3782(12)70019-1

Alcon-Giner, C. *et al.* Microbiota Supplementation with Bifidobacterium and Lactobacillus Modifies the Preterm Infant Gut Microbiota and Metabolome: An Observational Study. *Cell Rep Med* **1**, 100077 (2020). https://doi.org/10.1016/j.xcrm.2020.100077

Patel, N. *et al.* How frequent is routine use of probiotics in UK neonatal units? *BMJ Paediatr Open* **7** (2023). https://doi.org/10.1136/bmjpo-2023-002012

Gasparrini, A. J. *et al.* Persistent metagenomic signatures of early-life hospitalization and antibiotic treatment in the infant gut microbiota and resistome. *Nat Microbiol* **4**, 2285-2297 (2019). https://doi.org/10.1038/s41564-019-0550-2

Van Daele, E. *et al.* Effect of antibiotics in the first week of life on faecal microbiota development. *Arch Dis Child Fetal Neonatal Ed* **107**, 603-610 (2022). https://doi.org/10.1136/archdischild-2021-322861

Fredriksen, S., de Warle, S., van Baarlen, P., Boekhorst, J. & Wells, J. M. Resistome expansion in disease-associated human gut microbiomes. *Microbiome* **11**, 166 (2023). https://doi.org/10.1186/s40168-023-01610-1

Redondo-Salvo, S. *et al.* Pathways for horizontal gene transfer in bacteria revealed by a global map of their plasmids. *Nat Commun* **11**, 3602 (2020). https://doi.org/10.1038/s41467-020-17278-2

Shao, Y. *et al.* Primary succession of Bifidobacteria drives pathogen resistance in neonatal microbiota assembly. *Nat Microbiol* **9**, 2570-2582 (2024). https://doi.org/10.1038/s41564-024-01804-9

Brooks, B. *et al.* Strain-resolved analysis of hospital rooms and infants reveals overlap between the human and room microbiome. *Nat Commun* **8**, 1814 (2017). https://doi.org/10.1038/s41467-017-02018-w

Carl, M. A. *et al.* Sepsis from the gut: the enteric habitat of bacteria that cause late-onset neonatal bloodstream infections. *Clinical infectious diseases : an official publication of the Infectious Diseases Society of America* **58**, 1211-1218 (2014). https://doi.org/10.1093/cid/ciu084

Wandro, S. *et al.* The Microbiome and Metabolome of Preterm Infant Stool Are Personalized and Not Driven by Health Outcomes, Including Necrotizing Enterocolitis and Late-Onset Sepsis. *mSphere* **3** (2018). https://doi.org/10.1128/mSphere.00104-18

18   Jo, K. J. *et al.* Enterococcal bacteremia in children: Clinical Significance of vancomycin
resistance. *Pediatr Neonatol* (2024). https://doi.org/10.1016/j.pedneo.2024.01.007

19   Köser, C. U. *et al.* Rapid whole-genome sequencing for investigation of a neonatal MRSA
outbreak. *N Engl J Med* **366**, 2267-2275 (2012). https://doi.org/10.1056/NEJMoa1109910

20   Group, C. o. P. o. L. B. I. New World Health Organization recommendations for care of
preterm or low birth weight infants: health policy. *EClinicalMedicine* **63**, 102155 (2023).
https://doi.org/10.1016/j.eclinm.2023.102155

21   Neumann, C. J. *et al.* Clinical NEC prevention practices drive different microbiome profiles
and functional responses in the preterm intestine. *Nat Commun* **14**, 1349 (2023).
https://doi.org/10.1038/s41467-023-36825-1

22   Aceti, A. *et al.* Probiotics for prevention of necrotizing enterocolitis in preterm infants:
systematic review and meta-analysis. *Ital J Pediatr* **41**, 89 (2015).
https://doi.org/10.1186/s13052-015-0199-2

23   Robertson, C. *et al.* Incidence of necrotising enterocolitis before and after introducing routine
prophylactic Lactobacillus and Bifidobacterium probiotics. *Arch Dis Child Fetal Neonatal Ed*
(2019). https://doi.org/10.1136/archdischild-2019-317346

24   Batta, V. K., Rao, S. C. & Patole, S. K. Bifidobacterium infantis as a probiotic in preterm
infants: a systematic review and meta-analysis. *Pediatr Res* **94**, 1887-1905 (2023).
https://doi.org/10.1038/s41390-023-02716-w

25   Plummer, E. L. *et al.* Gut microbiota of preterm infants supplemented with probiotics: sub-
study of the ProPrems trial. *BMC Microbiol* **18**, 184 (2018). https://doi.org/10.1186/s12866-
018-1326-1

26   Hui, Y. *et al.* The effect of early probiotic exposure on the preterm infant gut microbiome
development. *Gut microbes* **13**, 1951113 (2021).
https://doi.org/10.1080/19490976.2021.1951113

27   Mohan, R. *et al.* Effects of Bifidobacterium lactis Bb12 supplementation on body weight,
fecal pH, acetate, lactate, calprotectin, and IgA in preterm infants. *Pediatr Res* **64**, 418-422
(2008). https://doi.org/10.1203/PDR.0b013e318181b7fa

28   O'Neill, I., Schofield, Z. & Hall, L. J. Exploring the role of the microbiota member
Bifidobacterium in modulating immune-linked diseases. *Emerging Topics in Life Sciences* **1**,
333-349 (2017). https://doi.org/10.1042/ETLS20170058

29   Guitor, A. K. *et al.* Capturing the antibiotic resistome of preterm infants reveals new benefits
of probiotic supplementation. *Microbiome* **10**, 136 (2022). https://doi.org/10.1186/s40168-
022-01327-7

30   Taft, D. H. *et al.* Bifidobacterial Dominance of the Gut in Early Life and Acquisition of
Antimicrobial Resistance. *mSphere* **3** (2018). https://doi.org/10.1128/mSphere.00441-18

31   van Best, N. *et al.* Influence of probiotic supplementation on the developing microbiota in
human preterm neonates. *Gut microbes* **12**, 1-16 (2020).
https://doi.org/10.1080/19490976.2020.1826747

32   Esaiassen, E. *et al.* Effects of Probiotic Supplementation on the Gut Microbiota and
Antibiotic Resistome Development in Preterm Infants. *Front Pediatr* **6**, 347 (2018).
https://doi.org/10.3389/fped.2018.00347

33   Jolley, K. A., Bray, J. E. & Maiden, M. C. J. Open-access bacterial population genomics:
BIGSdb software, the PubMLST.org website and their applications. *Wellcome Open Res* **3**,
124 (2018). https://doi.org/10.12688/wellcomeopenres.14826.1

34   Magnan, C. *et al.* Emergence of multidrug-resistant *Staphylococcus haemolyticus* in neonatal
intensive care unit in Southern France, a genomic study. *Emerg Microbes Infect* **13**, 2353291
(2024). https://doi.org/10.1080/22221751.2024.2353291

35   Ocloo, R., Newton-Foot, M., Ziebuhr, W. & Whitelaw, A. C. Molecular epidemiology and
antibiotic resistance of staphylococci other than *Staphylococcus aureus* in children in Cape
Town, South Africa. *Front Microbiol* **14**, 1239666 (2023).
https://doi.org/10.3389/fmicb.2023.1239666

36   Dewi, D. A. P. R. *et al.* Detection and genetic characterization of multidrug-resistant
staphylococci isolated from public areas in an international airport. *Sci Rep* **14**, 27738 (2024).
https://doi.org/10.1038/s41598-024-79447-3

37   Al-Turkait, A., Szatkowski, L., Choonara, I. & Ojha, S. Drug utilisation in neonatal units in
England and Wales: a national cohort study. *Eur J Clin Pharmacol* **78**, 669-677 (2022).
https://doi.org/10.1007/s00228-021-03267-x
38   Patangia, D. V. *et al.* Early life exposure of infants to benzylpenicillin and gentamicin is
associated with a persistent amplification of the gut resistome. *Microbiome* **12**, 19 (2024).
https://doi.org/10.1186/s40168-023-01732-6
39   Sprockett, D., Fukami, T. & Relman, D. A. Role of priority effects in the early-life assembly
of the gut microbiota. *Nat Rev Gastroenterol Hepatol* **15**, 197-205 (2018).
https://doi.org/10.1038/nrgastro.2017.173
40   Horrocks, V., King, O. G., Yip, A. Y. G., Marques, I. M. & McDonald, J. A. K. Role of the
gut microbiota in nutrient competition and protection against intestinal pathogen colonization.
Microbiology (Reading) **169** (2023). https://doi.org/10.1099/mic.0.001377
41   Darby, E. M. *et al.* Molecular mechanisms of antibiotic resistance revisited. *Nature reviews.*
Microbiology **21**, 280-295 (2023). https://doi.org/10.1038/s41579-022-00820-y
42   Carroll, L. M. *et al.* Identification of Novel Mobilized Colistin Resistance Gene. *mBio* **10**
(2019). https://doi.org/10.1128/mBio.00853-19
43   Macesic, N. *et al.* Silent spread of mobile colistin resistance gene mcr-9.1 on IncHI2
'superplasmids' in clinical carbapenem-resistant Enterobacterales. *Clin Microbiol Infect* **27**,
1856.e1857-1856.e1813 (2021). https://doi.org/10.1016/j.cmi.2021.04.020
44   Gibson, M. K. *et al.* Developmental dynamics of the preterm infant gut microbiota and
antibiotic resistome. *Nat Microbiol* **1**, 16024 (2016).
https://doi.org/10.1038/nmicrobiol.2016.24
45   Bargheet, A. *et al.* Development of early life gut resistome and mobilome across gestational
ages and microbiota-modifying treatments. *EBioMedicine* **92**, 104613 (2023).
https://doi.org/10.1016/j.ebiom.2023.104613
46   Nhu, N. T. K. *et al.* High-risk *Escherichia coli* clones that cause neonatal meningitis and
association with recrudescent infection. *Elife* **12** (2024). https://doi.org/10.7554/eLife.91853
47   Riley, L. W. Pandemic lineages of extraintestinal pathogenic Escherichia coli. *Clin Microbiol*
Infect **20**, 380-390 (2014). https://doi.org/10.1111/1469-0691.12646
48   Weng, Y. *et al.* Multicenter Study of Colistin Heteroresistance in Carbapenem-Resistant
Klebsiella pneumoniae Strains in China. *Microbiol Spectr* **11**, e0221822 (2023).
https://doi.org/10.1128/spectrum.02218-22
49   McInnes, R. S., McCallum, G. E., Lamberte, L. E. & van Schaik, W. Horizontal transfer of
antibiotic resistance genes in the human gut microbiome. *Curr Opin Microbiol* **53**, 35-43
(2020). https://doi.org/10.1016/j.mib.2020.02.002
50   Underwood, M. A., German, J. B., Lebrilla, C. B. & Mills, D. A. Bifidobacterium longum
subspecies infantis: champion colonizer of the infant gut. *Pediatr Res* **77**, 229-235 (2015).
https://doi.org/10.1038/pr.2014.156
51   Turroni, F. *et al.* Diversity of bifidobacteria within the infant gut microbiota. *PloS one* **7**,
e36957 (2012). https://doi.org/10.1371/journal.pone.0036957
52   Samara, J. *et al.* Supplementation with a probiotic mixture accelerates gut microbiome
maturation and reduces intestinal inflammation in extremely preterm infants. *Cell Host*
Microbe **30**, 696-711.e695 (2022). https://doi.org/10.1016/j.chom.2022.04.005
53   O'Connell Motherway, M. *et al.* Carbohydrate Syntrophy enhances the establishment of
Bifidobacterium breve UCC2003 in the neonatal gut. *Sci Rep* **8**, 10627 (2018).
https://doi.org/10.1038/s41598-018-29034-0
54   Xiao, M. *et al.* Cross-feeding of bifidobacteria promotes intestinal homeostasis: a lifelong
perspective on the host health. *NPJ Biofilms Microbiomes* **10**, 47 (2024).
https://doi.org/10.1038/s41522-024-00524-6
55   Casaburi, G. *et al.* Early-life gut microbiome modulation reduces the abundance of antibiotic-
resistant bacteria. *Antimicrob Resist Infect Control* **8**, 131 (2019).
https://doi.org/10.1186/s13756-019-0583-6
56   Samarra, A., Cabrera-Rubio, R., Martínez-Costa, C. & Collado, M. C. The role
of *Bifidobacterium* genus in modulating the neonate microbiota: implications for antibiotic

871		resistance acquisition in early lifeThe role of. *Gut microbes* **16**, 2357176 (2024).
872		https://doi.org/10.1080/19490976.2024.2357176
873	57	Phillips, S. *et al.* The Pregnancy and EARly Life study (PEARL) - a longitudinal study to
874		understand how gut microbes contribute to maintaining health during pregnancy and early life.
875		*BMC Pediatr* **21**, 357 (2021). https://doi.org/10.1186/s12887-021-02835-5
876	58	Kiu, R. *et al.* Particular genomic and virulence traits associated with preterm infant-derived
877		toxigenic Clostridium perfringens strains. *Nat Microbiol* **8**, 1160-1175 (2023).
878		https://doi.org/10.1038/s41564-023-01385-z
879	59	Chen, S., Zhou, Y., Chen, Y. & Gu, J. fastp: an ultra-fast all-in-one FASTQ preprocessor.
880		*Bioinformatics* **34**, i884-i890 (2018). https://doi.org/10.1093/bioinformatics/bty560
881	60	Wick, R. R., Judd, L. M., Gorrie, C. L. & Holt, K. E. Unicycler: Resolving bacterial genome
882		assemblies from short and long sequencing reads. *PLoS Comput Biol* **13**, e1005595 (2017).
883		https://doi.org/10.1371/journal.pcbi.1005595
884	61	Bankevich, A. *et al.* SPAdes: a new genome assembly algorithm and its applications to
885		single-cell sequencing. *J Comput Biol* **19**, 455-477 (2012).
886		https://doi.org/10.1089/cmb.2012.0021
887	62	Langmead, B. & Salzberg, S. L. Fast gapped-read alignment with Bowtie 2. *Nat Methods* **9**,
888		357-359 (2012). https://doi.org/10.1038/nmeth.1923
889	63	Danecek, P. *et al.* Twelve years of SAMtools and BCFtools. *Gigascience* **10** (2021).
890		https://doi.org/10.1093/gigascience/giab008
891	64	Walker, B. J. *et al.* Pilon: an integrated tool for comprehensive microbial variant detection
892		and genome assembly improvement. *PLoS one* **9**, e112963 (2014).
893		https://doi.org/10.1371/journal.pone.0112963
894	65	Parks, D. H., Imelfort, M., Skennerton, C. T., Hugenholtz, P. & Tyson, G. W. CheckM:
895		assessing the quality of microbial genomes recovered from isolates, single cells, and
896		metagenomes. *Genome Res* **25**, 1043-1055 (2015). https://doi.org/10.1101/gr.186072.114
897	66	Chaumeil, P. A., Mussig, A. J., Hugenholtz, P. & Parks, D. H. GTDB-Tk: a toolkit to classify
898		genomes with the Genome Taxonomy Database. *Bioinformatics* (2019).
899		https://doi.org/10.1093/bioinformatics/btz848
900	67	Kiu, R. *sequence-stats: generate sequence statistics from FASTA and FASTQ files*,
901		<https://github.com/raymondkiu/sequence-stats> (2020).
902	68	Seemann, T. Prokka: rapid prokaryotic genome annotation. *Bioinformatics* **30**, 2068-2069
903		(2014). https://doi.org/10.1093/bioinformatics/btu153
904	69	Li, D., Liu, C. M., Luo, R., Sadakane, K. & Lam, T. W. MEGAHIT: an ultra-fast single-node
905		solution for large and complex metagenomics assembly via succinct de Bruijn graph.
906		*Bioinformatics* **31**, 1674-1676 (2015). https://doi.org/10.1093/bioinformatics/btv033
907	70	Uritskiy, G. V., DiRuggiero, J. & Taylor, J. MetaWRAP-a flexible pipeline for genome-
908		resolved metagenomic data analysis. *Microbiome* **6**, 158 (2018).
909		https://doi.org/10.1186/s40168-018-0541-1
910	71	Kang, D. D. *et al.* MetaBAT 2: an adaptive binning algorithm for robust and efficient genome
911		reconstruction from metagenome assemblies. *PeerJ* **7**, e7359 (2019).
912		https://doi.org/10.7717/peerj.7359
913	72	Wu, Y. W., Simmons, B. A. & Singer, S. W. MaxBin 2.0: an automated binning algorithm to
914		recover genomes from multiple metagenomic datasets. *Bioinformatics* **32**, 605-607 (2016).
915		https://doi.org/10.1093/bioinformatics/btv638
916	73	Alneberg, J. *et al.* Binning metagenomic contigs by coverage and composition. *Nat Methods*
917		**11**, 1144-1146 (2014). https://doi.org/10.1038/nmeth.3103
918	74	Olm, M. R., Brown, C. T., Brooks, B. & Banfield, J. F. dRep: a tool for fast and accurate
919		genomic comparisons that enables improved genome recovery from metagenomes through
920		de-replication. *The ISME journal* **11**, 2864-2868 (2017).
921		https://doi.org/10.1038/ismej.2017.126
922	75	Wood, D. E., Lu, J. & Langmead, B. Improved metagenomic analysis with Kraken 2.
923		*Genome Biol* **20**, 257 (2019). https://doi.org/10.1186/s13059-019-1891-0
924	76	Lu, J., Breitwieser, F., Thielen, P. & Salzberg, S. Bracken: estimating species abundance in
925		metagenomics data. *PeerJ Comput Sci* **3:e104** (2017). https://doi.org/10.7717/peerj-cs.104

77 R Development Core Team. in *R Foundation for Statistical Computing*   (Vienna, Austria,
2010).
78 Wickham, H. *ggplot2: Elegant Graphics for Data Analysis.* (Springer-Verlag, 2016).
79 Katz, L. S. *et al.* Mashtree: a rapid comparison of whole genome sequence files. *J Open*
Source Softw **4** (2019). https://doi.org/10.21105/joss.01762
80 Letunic, I. & Bork, P. Interactive Tree of Life (iTOL) v6: recent updates to the phylogenetic
tree display and annotation tool. *Nucleic Acids Res* **52**, W78-W82 (2024).
https://doi.org/10.1093/nar/gkae268
81 Gupta, S. K. *et al.* ARG-ANNOT, a new bioinformatic tool to discover antibiotic resistance
genes in bacterial genomes. *Antimicrob Agents Chemother* **58**, 212-220 (2014).
https://doi.org/10.1128/AAC.01310-13
82 Bortolaia, V. *et al.* ResFinder 4.0 for predictions of phenotypes from genotypes. *J Antimicrob*
Chemother **75**, 3491-3500 (2020). https://doi.org/10.1093/jac/dkaa345
83 Zhou, Z. *et al.* METABOLIC: high-throughput profiling of microbial genomes for functional
traits, metabolism, biogeochemistry, and community-scale functional networks. *Microbiome*
10, 33 (2022). https://doi.org/10.1186/s40168-021-01213-8
84 mlst (GitHub, 2022).
85 Jolley, K. A. & Maiden, M. C. BIGSdb: Scalable analysis of bacterial genome variation at the
population level. *BMC Bioinformatics* **11**, 595 (2010). https://doi.org/10.1186/1471-2105-11-
595
86 Seemann, T. *ABRicate: Mass screening of contigs for antimicrobial and virulence genes*,
<https://github.com/tseemann/abricate> (2018).
87 Carattoli, A. & Hasman, H. PlasmidFinder and In Silico pMLST: Identification and Typing of
Plasmid Replicons in Whole-Genome Sequencing (WGS). *Methods Mol Biol* **2075**, 285-294
(2020). https://doi.org/10.1007/978-1-4939-9877-7_20
88 Lam, M. M. C. *et al.* A genomic surveillance framework and genotyping tool for Klebsiella
pneumoniae and its related species complex. *Nat Commun* **12**, 4188 (2021).
https://doi.org/10.1038/s41467-021-24448-3
89 Hsu, T. Y. *et al.* Profiling novel lateral gene transfer events in the human microbiome.
bioRxiv (2023). https://doi.org/10.1101/2023.08.08.552500
90 Bender, J. K., Fiedler, S., Klare, I. & Werner, G. Complete Genome Sequence of the Gut
Commensal and Laboratory Strain Enterococcus faecium 64/3. *Genome Announc* **3** (2015).
https://doi.org/10.1128/genomeA.01275-15
91 Pungel, D. *et al.* Bifidobacterium breve UCC2003 Exopolysaccharide Modulates the Early
Life Microbiota by Acting as a Potential Dietary Substrate. *Nutrients* **12** (2020).
https://doi.org/10.3390/nu12040948
92 O'Donnell, M. M., Rea, M. C., Shanahan, F. & Ross, R. P. The Use of a Mini-Bioreactor
Fermentation System as a Reproducible, High-Throughput ex vivo Batch Model of the Distal
Colon. *Frontiers in microbiology* **9**, 1844 (2018). https://doi.org/10.3389/fmicb.2018.01844
93 Hariharan, J., Wright, K. & Passalacqua, P. dorado: A Python package for simulating passive
particle transport in shallow-water flows. *Journal of Open Source Software* **5**, 2585   (2020).
https://doi.org/https://doi.org/10.21105/joss.02585
94 Wick, R. *Filtlong*, <github.com/rrwick/Filtlong> (2024).
95 Kolmogorov, M., Yuan, J., Lin, Y. & Pevzner, P. A. Assembly of long, error-prone reads
using repeat graphs. *Nat Biotechnol* **37**, 540-546 (2019). https://doi.org/10.1038/s41587-019-
0072-8
96 Wright, C. & Wykes, M. *Medaka*, <https://github.com/nanoporetech/medaka> (2024).
97 Jain, C., Rodriguez, R. L., Phillippy, A. M., Konstantinidis, K. T. & Aluru, S. High
throughput ANI analysis of 90K prokaryotic genomes reveals clear species boundaries. *Nat*
Commun **9**, 5114 (2018). https://doi.org/10.1038/s41467-018-07641-9
98 Wickham, H. *et al.* Welcome to the Tidyverse. *Journal of open source software* **4**, 1686
(2019).
99 Kassambara, A. ggpubr:'ggplot2'based publication ready plots. *R package version*, 2 (2018).
100 Dixon, P. VEGAN, a package of R functions for community ecology. *J Veg Sci* **14**, 927-930
(2003). https://doi.org/https://doi.org/10.1111/j.1654-1103.2003.tb02228.x

101   Guy, L., Kultima, J. R. & Andersson, S. G. genoPlotR: comparative gene and genome
982   visualization in R. *Bioinformatics* **26**, 2334-2335 (2010).
983   https://doi.org/10.1093/bioinformatics/btq413
984   102   Branger, M. & Leclercq, S. O. GenoFig: a user-friendly application for the visualization and
985   comparison of genomic regions. *Bioinformatics* **40** (2024).
986   https://doi.org/10.1093/bioinformatics/btae372
987   103   Segata, N. *et al.* Metagenomic biomarker discovery and explanation. *Genome Biol* **12**, R60
988   (2011). https://doi.org/10.1186/gb-2011-12-6-r60
989   104   Beghini, F. *et al.* Integrating taxonomic, functional, and strain-level profiling of diverse
990   microbial communities with bioBakery 3. *Elife* **10** (2021).
991   https://doi.org/10.7554/eLife.65088
992   105   dplyr: A Grammar of Data Manipulation (2022).
993   106   rstatix: Pipe-Friendly Framework for Basic Statistical Tests (2023).
994

Fig. 1. Neonatal-period preterm infant gut microbiome analysis and functional profiling. a, Schematic of this study with key details and relevant analyses. **b,** NMDS plots of 34 preterm infant gut metagenomes samples within the first 21 days of life (stratified by weeks and coloured by cohorts) and clustered based on genus-level Bray-Curtis dissimilarity. Significant taxa were labelled in the respective plots. **c,** Gut microbiome profiles of 34 infants in relative abundance (%) based on genus level data. Only top 10 most abundant genera across all samples were shown here. **d,** Gut microbiome profiles (as in **c**) of 34 infants based on species-level taxonomic assignment. Only top 10 most abundant bacterial species were shown here. **e,** Gut microbiome diversity indices of 34 infants in 2 cohorts, stratified by weeks (two-sided Wilcoxon test). **f,** Comparison of taxonomic abundance (%) of three key *Bifidobacterium* species across 2 cohorts (NPS vs PS) stratified by weeks. Percentages indicated are mean relative abundances in each species. Statistical significance was performed using two-sided Wilcoxon test (NPS vs PS with respect to weeks). * *P*<0.05, **P*<0.01, ***P*<0.001, ****P*<0.0001. **g,** Mean taxa proportion of top 10 most abundant bacterial genera in both cohorts as shown in the bubble plots. The significance (two-sided Wilcoxon test) was compared NPS vs PS with respect to the time points (week). * *P*<0.05, **P*<0.01, ***P*<0.001, ****P*<0.0001. **h,** NMDS plots of 34 preterm infant gut microbiome functional profiles stratified by weeks, NPS vs PS. **i,** Functional profiling heatmap showing 41 significant pathways profiled (alpha=0.05, LDA scores>2.0). The box-within-figure shows a Venn diagram of overlapping/unique functional pathways NPS vs PS based on a total of 407 pathways identified computationally.

Fig. 2. **An in-depth profiling of neonatal preterm infant gut resistome. a,** Schematic of antibiotics regime timeline in the preterm infants. **b,** Comparison of ARGs between NPS and PS cohorts. The significance was compared using two-sided Wilcoxon test. * $P<0.05$, **$P<0.01$, ***$P<0.001$. **c,** Number of ARG types (antibiotic/drug classes against which encoding ARGs are resistant) comparison between NPS and PS cohorts stratified by time points. Statistical analysis was performed using two-sided Wilcoxon test. * $P<0.05$, **$P<0.01$. **d,** Proportions of ARG types in both NPS and PS cohorts stratified by weeks. The mini bar chart within sub-figure represents the mean number of ARGs in both NPS and PS cohorts coloured by ARG types. Statistical analysis was performed using two-sided Wilcoxon test. ***$P<0.001$. **e,** NMDS plot of ARG profiles of individuals in bot NPS and PS cohorts. **f,** Venn diagram showing number/percentages of unique and sharing ARGs between NPS and PS cohorts. **g,** Shannon index (diversity index) comparison between Antibiotic vs Control (untreated) in NPS and PS cohorts. **h,** Preterm gut microbiome dynamics (Top 10 genera were shown) in the first three weeks of life, stratified by antibiotic treated individuals and control group (without antibiotic treatment). **i,** Increase/reduction of taxonomic abundance (Top 10 genera) in Antibiotic group (vs Control) in both NPS and PS cohorts. Shaded area represents antibiotic treatment period, infants in Antibiotic group were treated with antibiotic only in their first week of life (3 median days). Statistical analysis was performed using two-sided Wilcoxon test. . * $P<0.05$, **$P<0.01$. **j,** No. of ARGs comparison between Antibiotic vs Control (untreated) in both NPS and PS cohorts. **k,** Number of ARGs was compared NPS vs PS, stratified by antibiotic usage. Statistical analysis was performed using two-sided Wilcoxon test. * $P<0.05$, **$P<0.01$. **l,** Correlation plot of (Shannon) diversity index vs ARGs using all available data points. Scatter plot was coloured by cohorts accordingly. R represents Kendall rank correlation coefficient. **m,** Genomic mapping of a computationally extracted sequence contig with *mcr-9.1* ARG gene against a *mcr-9.1* ARG from *E. coli* (NCBI nucleotide database) with 100% matching nucleotide identity.

Fig. 3. Strain-level gut-derived pathobiont resistome analysis. a, Strain-level neighbor-joining tree of bacterial strains (n=195) aligned with clinical metadata, ARG profiles and gene counts, including both isolate genomes and metagenome-assembled genomes (MAGs) derived from a total of 411 individual bacterial genomes. **b,** ARG count per strain comparison across Top 10 genera. Statistical analysis were performed using Kruskal Wallis test and post-hoc Dunn's test (FDR adjusted). Significance was compared against *Bifidobacterium*. **P<0.01, **** P<0.0001. **c,** ARG count per strain comparison between Top 4 drug-resistant pathobiont genera including *Enterococcus*, *Escherichia*, *Klebsiella* and *Staphylococcus*. Statistical analysis between cohorts were performed using two-sided Wilcoxon test. ns: non-significant. **d,** Multidrug resistance efflux pump profiles (n=20) of Top 4 drug-resistant pathobiont genera *Enterococcus*, *Escherichia*, *Klebsiella* and *Staphylococcus* genomes (n=205). **e,** (Top) Proportion of ARG type per genome (n=205) in both NPS and PS cohorts. (Bottom) Proportion of MDR/non-MDR genomes in both NPS and PS cohorts. MDR is defined as having three or more predicted ARG types (resistant against drugs of three or more antibiotic classes). Statistical test was performed using Fisher Exact test. ****P<0.0001. **f,** Multi-locus Sequence Typing (MLST) analysis of key pathobiont genomes (n=193) including Top 4 pathobiont genera *Enterococcus*, *Staphylococcus*, *Escherichia* and *Klebsiella*. Additional surface antigen typing (Kaptive) for *Klebsiella* spp as shown in the box-within-figure.

Fig. 4. Strain-level mobilome analysis and persistence of strains in preterm infant gut microbiome. a, Plasmid replicon profiles of preterm infant gut-associated bacterial strains (n=195) aligned with a strain-level neighbor-joining tree. Plasmid and ARG counts were also included for comparison. **b,** Gut microbiome plasmid count comparison NPS vs PS. Statistical test was performed using Wilcoxon test. *P<0.05. **c,** Gut microbiome plasmid count comparison Antibiotic vs Control, stratified by cohort NPS vs PS. Statistical test was performed using Wilcoxon test. ns: non-significant. **d,** Gut microbiome plasmid count comparison between Antibiotic vs Control in all preterm infant metagenome samples. **e,** Correlation plot of ARGs vs plasmid count (in all preterm infant metagenome samples) using all available data points. Scatter plot was coloured by Antibiotic usage accordingly, where purple represents infants administered with antibiotic while white indicates untreated infants. *R* represents Kendall rank correlation coefficient. **f,** HGT events per genome (n=411) between Antibiotic groups and Control. Statistical analysis was performed using Wilcoxon test. ****P<0.0001. **g,** Plasmid replicon count per strain comparison across the Top 10 genera. Statistical analysis were performed using Kruskal Wallis test and post-hoc Dunn's test (FDR adjusted). Significance was compared against *Bifidobacterium*. *P<0.05, **** P<0.0001. **h,** Potential dissemination of *Enterococcus* strains among certain preterm infants in the NICUs. Strain-level cut-off ANI>99.9%.

Fig. 5. Gentamicin-resistant plasmid transfer study in *Enterococcus*. a, Schematic of *in vitro* study workflow and relevant analyses with the study design in 24-well plate template. **b,** Antibiotic resistance gene profiling of donor and recipient strains, colonies singly picked from selective agars supplemented with appropriate antibiotics and whole-genome sequenced via long-read/short-read sequencing platform. A neighbour-joining tree was constructed to reflect the relatedness between isolate genomes, aligned with plasmid replicon predicted in the genomes in tandem with calculated genome sizes. The box-within-box (top left) was shotgun metagenome contents (microbial taxa) of infant faecal slurry that was used in the gut model experiment. **aac6-Ii* is known as a chromosomal-encoded aminoglycoside acetyltransferase in *Enterococcus* spp. **c,** Comparison of computationally extracted antibiotic resistance gene *aac-aph2* from representative donor strain B1, recipient (transformant) strains D1-1 and D3-1. **d,** Comparison of complete contig sequences that encoded gentamicin resistance gene *aac6-aph2,* predicted to be either whole or part of potential mobile genetic elements including plasmids.