FabF and FadM Cooperate to Recycle Fatty Acids and Rescue Δ*plsX* Lethality in *Staphylococcus aureus*

Paprapach Wongdontree[1], Milya Palmier[1], Clara Louche[1], Vincent Leguillier[1], Carine Machado Rodrigues[2], Karine Gloux[1], David Halpern[1], Céline Henry[2], Jamila Anba-Mondoloni[1], Alexandra Gruss[1#]

[1] Université Paris-Saclay, INRAE, AgroParisTech, Micalis Institute, 78350, Jouy-en-Josas, France.

[#] For correspondence: alexandra.gruss@inrae.fr

Keywords: *Staphylococcus aureus,* MRSA, fatty acid, FASII, phospholipid synthesis, *plsX* deletion suppressors, FabF, platensimycin, FadM, acyl-CoA thioesterase, acyl-ACP thioesterase, ACP binding, membrane, β-lactam sensitivity.

Short title: FabF-FadM role in fatty acid recycling

Abstract

Phospholipids are essential components of most cell membranes. In *Staphylococcus aureus*, PlsX acyltransferase is considered indispensable for initiating phospholipid synthesis, unless exogenous fatty acids (FAs) are available to bypass this requirement. We report that *S. aureus* can capture internal FA sources to overcome PlsX essentiality in a Δ*plsX* mutant *via* point mutations in either of two genes: *fabF*, which encodes the FA synthesis enzyme 3-oxoacyl-(acyl-carrier-protein) synthase II, or *fadM*, which encodes an understudied bifunctional acyl-CoA thioesterase and ACP binding protein. Despite growth rescue, both Δ*plsX* suppressors differ from the parental strain by producing phospholipids with shortened FA lengths suggesting that both suppressors lead to premature FA release during synthesis. Additionally, both suppressors display increased sensitivity to β-lactam antibiotics. The similar behavior of both suppressors led us to show that *fabF* suppressors require the presence of *fadM*, indicative of FabF-FadM cooperation. We propose that reduced processivity of FabF suppressor variants, or greater availability of FadM for ACP binding in FadM variants, facilitates FA release from FabF-acyl-ACP intermediates. A FabF-FadM relay leading to FA release may contribute to homeostasis between FASII and phospholipid synthesis pathways.

Significance

Phospholipids are vital cell membrane components. The essential *Staphylococcus aureus* phospholipid synthesis enzyme PlsX uses acyl-ACP, the end-product of fatty acid (FA) synthesis (FASII), to initiate phospholipid production. Despite its central role, PlsX can be substituted by exogenous FAs whose phosphorylation yields the same product. We discovered that without FA supplementation, mutants arise that rescue growth, indicating that internal FAs are released. Mutations occurred in either FabF, a FASII enzyme, or in FadM, an incompletely characterized protein. Our analyses give evidence that FabF and FadM proteins cooperate, and facilitate FA availability when either protein is mutated. We propose that in normal conditions, FadM might act as an "overflow valve" by releasing FAs from the FabF intermediate, which prevents buildup of FASII intermediates, and ensures FA-phospholipid balance. Remarkably, while this pathway rescues *S. aureus* growth, it sensitizes the MRSA strain to β-lactam antibiotics.

Introduction

Lipids provide the essential scaffolding for cell membranes and life. Bacterial phospholipid membranes comprise fatty acids (FA) that are attached to a glycerophosphate backbone [1]. The phospholipid FAs are either produced by the FA synthesis (FASII) pathway or captured from the environment [2]. The FASII product, acyl-Acyl Carrier Protein (acyl-ACP or FA-ACP) is a substrate for three enzymes: FabF for FA elongation *via* FASII, and PlsX and PlsC for respectively the first and second steps of phospholipid synthesis (**Fig. 1**).

The membrane-associated acyltransferase PlsX catalyzes the first dedicated step of phospholipid synthesis in Bacillota bacteria, by reversibly catalyzing conversion of acyl-ACP, the FASII product, to acyl-phosphate (acyl-PO$_4$) [3]. An alternative phospholipid initiation mechanism, not requiring PlsX, involves assimilation of exogenous FAs *via* phosphorylation by the FA kinase Fak [4]. The membrane glycerol-3-phosphate acyltransferase PlsY then transfers the FA moiety of acyl-PO$_4$ to the 1-position of the glycerophosphate backbone, forming lysophosphatidic acid. However, reverse PlsX activity may also reconvert acyl-PO$_4$ to acyl-ACP when exogenous FAs are available, which then re-enter the FASII system or are used by PlsC for phospholipid synthesis [4, 5] (**Fig. 1**). Providing acyl-ACP by PlsX reverse activity on exogenous FAs, once phosphorylated, allows bacteria to overcome the need for FASII, making them insensitive to FASII inhibitors [6-8]. In *Bacillus subtilis*, PlsX is membrane-associated for phospholipid synthesis, which could facilitate acyl-PO$_4$ transfer to PlsY, but likely not for its reverse activity [9]. PlsX is thus a sorting station that initiates phospholipid synthesis, or returns acyl groups for FASII elongation and PlsC processing.

When exogenous FAs are absent, PlsX may still be dispensable in some bacteria. Numerous Pseudomonadota bacteria encode PlsB, which has functional redundancy to PlsX-PlsY [3]. In *E. coli*, suppressors of the synthetic lethal $\Delta plsX$ $\Delta plsY$ mutation (despite the existence of the functionally redundant enzyme PlsB [3]) [10] displayed increased pools of glycerol-3-phosphate (Gly-3-P), the backbone for FA attachment to produce phospholipids [10]. Increased Gly-3-P was suggested to compensate the low efficiency acyl-transferase activity of PlsB compared to PlsY. Among Bacillota pathogens, some deploy an acyl-ACP thioesterase (acyl-ACP TE) that liberates endogenous FAs from acyl-ACP**,** which are then phosphorylated by Fak to initiate phospholipid synthesis [11-14]. Acyl-ACP TE fully restores growth of *Streptococcus pneumoniae* $\Delta plsX$ [12], and conditionally rescues *Enterococcus faecalis* $\Delta plsX$ growth; however, the acyl-ACP TE preferentially cleaves acyl-ACP comprising

unsaturated FAs, while growth is stimulated by saturated FAs [13]. These alternative systems in Bacillota all generate an internal FA supply to initiate phospholipid synthesis when PlsX function is impaired.

The major human pathogen *Staphylococcus aureus* lacks PlsB, and an acyl-ACP TE to liberate FAs, and only exogenous FAs or a cloned gene encoding heterologous acyl-ACP TE reportedly complemented growth of a *plsX* deletion strain [5]. Our previous work on FASII bypass revealed that exogenous FAs rescue *S. aureus* growth by one of two processes: point mutations that disable *fabD* [7], or by non-mutational adaptation in which FAs are efficiently incorporated after a 6-8 h time lag [6-8]. These studies led us to now investigate the step following FASII, mediated by PlsX. For this we constructed Δ*plsX* mutants, and unexpectedly identified suppressors conferring growth. In this report, we characterized activity-altering mutations mapping to two distinct loci that rescue Δ*plsX* viability without exogenous FAs. Mutations in FASII enzyme FabF, a 3-oxoacyl-ACP synthase II, or in an incompletely characterized acyl-CoA thioesterase and ACP binding protein designated as FadM, rescue Δ*plsX* growth. Our findings reveal that *plsX* essentiality is abolished by point mutations in *fabF* or *fadM*. We demonstrate interdependency of FabF and FadM in rescuing the absence of PlsX. These findings may indicate a general role for FadM as a newly found actor shuttling FAs from FASII towards phospholipid synthesis. Remarkably, the *fabF* and *fadM* suppressors rescue Δ*plsX* growth, but sensitize an MRSA *S. aureus* strain to the β-lactam antibiotic amoxicillin.

Results

Construction and confirmation of a Δ*plsX* in-frame mutant. *plsX* is the second gene in a multicistronic operon also comprising FASII genes (**Supplementary Fig. S1**). We constructed in-frame Δ*plsX* mutants of RN-R (RN4220 repaired for a defective *fakB1* gene in the 8325 lineage [15]) and of USA300_FPR3757 JE2 strains. Both Δ*plsX* mutants failed to grow on BHI solid medium in the absence of FAs. Deletions were confirmed by PCR and whole genome sequencing (**Table S1**) and by RN-R Δ*plsX* growth complementation in the absence of FAs using a plasmid-carried copy of the intact *plsX* gene (**Supplementary Fig. S2**).

FA auxotrophy of *S. aureus* Δ*plsX* mutants is suppressed by a point mutation in *fabF*. During phenotypic characterization of the JE2 Δ*plsX* mutant, we noted the appearance of colonies

on BHI medium without added FA (**Fig. 2A**, left plate). Note that BHI reportedly contains traces of FA [16], but which are insufficient to confer $\Delta plsX$ growth. Two colonies subjected to whole genome sequencing confirmed the in-frame *plsX* deletion and also revealed point mutations mapping to FabF (SAUSA300_0886), the 3-oxoacyl-ACP synthase II of the FASII elongation cycle (see **Fig. 1**), producing FabF^{A119E} (alanine to glutamic acid; Sup1) and FabF^{D266A} (aspartic acid to alanine; Sup2). The suppressor carrying FabF^{D266A} was not recoverable from the primary stocked strain, possibly because the mutation was deleterious for growth. Sup1 and Sup2 also carried the same point mutation in *csa1A* (SAUSA300_0100), encoding a tandem lipoprotein.

We expanded our analyses by isolating 31 other $\Delta plsX$ suppressors from RN-R or JE2 backgrounds (16 and 15 suppressors, respectively), and screened for those carrying *fabF* mutations by PCR followed by DNA sequence analyses (**Table S2**). Among them, 14 carried *fabF* mutations, which mapped to the same *fabF* nucleotide position in both strain backgrounds, resulting in a FabF^{A119E} alteration as in Sup1. This allele is referred to as *fabF1*. *csa1A*, which carried a mutation in Sup1 and Sup2 mutants, was not mutated in the 14 *fabF* mutants, indicating that *fabF* was the suppressor allele. Emergence of the same *fabF* mutation in both suppressor strain backgrounds could suggest that few FabF point modifications would allow both FASII activity and $\Delta plsX$ complementation.

We attempted to prove that the *fabF* mutation was directly responsible for the $\Delta plsX$ suppressor phenotype by PCR-amplifying the WT *fabF* and mutant *fabF1* alleles, cloning on a medium copy-number vector and constitutive promoter, and establishing plasmids in the $\Delta plsX$ and/or *fabF1$\Delta plsX$* strains, with the goal of reversing the suppressor phenotype. A clone expressing *fabF1*, but not WT *fabF* could be established in the WT RN-R strain by direct transformation.; this is consistent with the putative reduced activity of this allele. We further failed to establish the *fabF1* clone in the $\Delta plsX$ background, even in FA-supplemented medium. This suggested that *fabF* overexpression was toxic in *S. aureus*, as reported in *Escherichia coli* [17], and that our strategy was not feasible.

As an alternate approach, we used a FabF inhibitor, platensimycin, to determine whether disabling FabF was responsible for $\Delta plsX$ suppression. This antibiotic binds to the acyl-ACP-FabF intermediate, and competes with malonyl-ACP entry, thus preventing FA elongation [18]. However, it also leads to increased levels of FabF and other FASII enzymes in *Bacillus subtilis* [19]. We tested platensimysin, like the FabF mutant, would overcome the $\Delta plsX$ growth defect, presumably by slowing FabF activity. RN-R $\Delta plsX$ cultures were plated on BHI solid medium, and platensimycin (1.5 µg) was deposited on plates, which were then

incubated 96 h at 37°C (**Fig. 2A**, right plate). In addition to the emergence of ΔplsX suppressors, a distinct homogeneous growth ring appeared around the platensimycin spots. Similar experiments were performed in liquid medium using 2-fold dilutions of platensimycin concentrations ranging from 1 ng to 500 ng/ml (**Supplementary Fig. S3**). Compared to BHI medium, growth between 7 and 12 h was significantly stimulated by the presence of 125 ng/ml platensimycin (P=≤0.05). However, experiments in liquid medium are potentially flawed as mutant emergence and platensimycin stimulation cannot be distinguished. Ring formation on solid medium appears to be a more accurate means of assessing growth stimulation. Since the *fabF* mutants and a FabF inhibitor both restore ΔplsX growth, we consider it likely that the FabF mutation is directly responsible for ΔplsX suppression. Reducing *fabF* efficacy by mutation or subinhibitory platensimycin treatments could destabilize the FabF-acyl-ACP intermediate, to liberate free FAs and complement the ΔplsX growth defect (**Fig. 2B**).

FA auxotrophy of *S. aureus* ΔplsX mutants is suppressed by mutations in *fadM* that map to the acyl-CoA binding cavity. We searched for mutations that conferred ΔplsX growth on BHI among the 17 remaining suppressors carrying a wild type *fabF* sequence. Genomic DNA sequencing was performed on three such isolates (one derived from RN-R ΔplsX and two from JE2 ΔplsX backgrounds). A single common gene target, SAOUHSC_01348 in RN-R or SAUSA300_1247 in JE2, was mutated (**Table S1**, **Table S2**). While annotated as a 1,4-dihydroxy-2-naphthoyl-coenzyme A (DHNA-CoA) thioesterase implicated in menaquinone synthesis, the enzyme was shown to have acyl-CoA thioesterase activity [20]. Moreover, the 155-amino-acid ORF shares ~40% similarity and a conserved serine, histidine and aspartic acid catalytic triad with FadM, an *E. coli* acyl-CoA thioesterase and ACP binding protein (PBD database protein 1NJK), [20-24]. The *S. aureus* gene, referred to as *fadM*, was PCR-amplified and sequenced in the 17 other suppressors lacking mutations in *fabF*. All carried mutations in *fadM*. A single variant (FadM[Y90F], [encoded by *fadM2*]) was identified in the four isolates selected from RN-R ΔplsX; three distinct FadM variants were identified among the JE2 ΔplsX suppressors (FadM[I38T] [*fadM1*], FadM[Y90F] [*fadM2*], and FadM[Y133F] [*fadM3*]) (**Table S1**, **Table S2**). These results point to FabF and FadM as the main suppressors rescuing ΔplsX growth.

The *S. aureus* FadM crystal structure (PDB: 6FDG) and Alphafold predictions (https://alphafold.ebi.ac.uk/entry/A0A0H2XFE3) provide detailed information for mapping the identified FadM variants conferring ΔplsX suppression. The variants coincided with or

197	mapped adjacent to amino acids reportedly involved in forming an FA binding cavity that
198	favors long FA substrates [20] (**Fig. 3A**). The grouped location of these variants leads us to
199	speculate that acyl-CoA thioesterase activity may be defective in the *fadM* suppressors.
200	If FadM acyl-CoA thioesterase activity were needed for Δ*plsX* suppression, then its
201	substrate, acyl-CoA, would need to be produced. In *S. aureus*, acyl-CoA is synthesized by acyl-
202	CoA synthetase (encoded by *fadE*), which is part of the recently elucidated FA degradation
203	(*fad*) system (**Fig. 1**; [25, 26]). The *fad* genes are subject to CcpA-mediated glucose repression
204	[25-27]. However, glucose did not affect the *fadM* mutant Δ*plsX* suppressor phenotypes
205	(**Supplementary Fig. S4**). Based on above results, we consider it reasonable to rule out a role
206	for FadM acyl-CoA thioesterase activity in Δ*plsX* growth rescue.
207	
208	**A *fadM* null mutant does not confer Δ*plsX* growth.** Suppression by *fadM* point mutants could
209	be due to FadM-mediated ACP binding, or to the loss of all FadM activity. We asked whether
210	inactivation of *fadM*, the downstream gene in a 2-gene operon comprising *acnA* ([27];
211	**Supplementary Fig. S1**), suppresses the Δ*plsX* phenotype. For this, a *fadM* transposon
212	insertion from the Nebraska Transposon Mutant Library library, interrupting FadM at amino
213	acid position 5, was transferred by φ80-mediated transduction into the JE2 Δ*plsX fadM2*
214	suppressor (strain 7.1; **Table S2**) [28]. The Δ*plsX fadM::*Tn strain failed to grow on FA-free
215	medium (**Fig. 3B**). Since total loss of FadM does not promote Δ*plsX* suppression, we conclude
216	that some FadM activity is required for *S. aureus* Δ*plsX* suppression. As FadM I38T, Y90F, and
217	Y133F mutations each map to the acyl-CoA thioesterase binding tunnel, we propose that the
218	FadM ACP binding activity is involved in Δ*plsX* suppression (**Fig. 3C**).
219	
220	**Impact of Δ*plsX* suppressors on bacterial growth properties.** Growth of Δ*plsX* suppressors
221	was assessed in RN-R and JE2 backgrounds in non-FA-supplemented BHI medium (**Fig. 4A**).
222	The initial Δ*plsX* deletion strains fail to grow unless supplemented with FAs. Both *fabF* and
223	*fadM* suppressor mutants grew similarly, but their growth phenotypes differed according to
224	the parental strains: in the RN-R background, growth of both tested suppressors was
225	intermediate between the WT strain and the Δ*plsX* mutant. In contrast, JE2 derivative
226	suppressors grew like the parental strain. A distinguishing feature between the two *S. aureus*
227	strains is the production of staphyloxanthin pigment by JE2 but not by RN-R (an RN4220
228	derivative), whose hydrophobic long-chain isoprenyl moiety of staphyloxanthin [29] could
229	conceivably stabilize the membrane in suppressor strains. However, reduced pigmentation

of Δ*plsX*, and both *fabF* and *fadM* suppressors compared to WT and *fadM::*Tn controls (**Fig.**
4B), indicate that other factors are involved in improved rescue in the JE2 background.

Membrane FAs are shortened in *fabF* and *fadM* suppressors of Δ*plsX*. We considered that
the *fabF1* point mutation in the Δ*plsX fabF1* suppressor strain might alter FabF processivity,
suggesting a shift in FASII products. We also asked whether the mutations in *fadM,* encoding
a possible FabF partner, might have similar effects. We therefore assessed membrane FA
composition of both Δ*plsX* suppressors in RN-R and JE2 backgrounds in medium devoid of
FAs (**Fig. 4C**, **Table S3**). The *fabF* suppressors produced membrane FAs of markedly shorter
overall length in both strain backgrounds. Notably, straight-chain FAs C18:0 and C20:0,
represent 23-29% of total FAs in parental strains but are essentially absent in *fabF*
suppressors. This suggests that processivity of FabF^{A119E} expressed from the *fabF1* suppressor
allele is reduced compared to WT FabF. In *S. aureus*, branched chain precursors are preferred
substrates for FabH (see **Fig. 1**; [30]), which might lead to higher proportions of branched
chain FAs (*ai*15 and *ai*17) in *fabF1* suppressors. FA profiles in the *fadM2* suppressor showed
a 2-fold decrease in proportions of C18:0 and C20:0 compared to those in the respective
parental strains (**Fig. 4C**, **Table S3**). These results confirm that both *fabF* and *fadM* mutant
suppressors of Δ*plsX* lead to decreased proportions of long chain saturated FAs in
phospholipids.

Δ*plsX* suppressors display accrued β-lactam sensitivity. β-lactam resistance of methicillin
resistant *S. aureus* (MRSA) is associated with the presence of penicillin binding protein
Pbp2A' (encoded by *mecA*), and is a main cause of treatment failure [31, 32]. Interestingly,
disturbance of membrane lipid microdomains affects Pbp2A' oligomerization and
compromises resistance [33]. We asked whether the membrane changes induced by Δ*plsX*
suppressors, notably their shorter FAs (**Fig. 4C**) might also impact β-lactam antibiotic
resistance profiles. For this, amoxicillin minimal inhibitory concentrations (MICs) were
measured by Etests. Tests were performed on BHI solid medium without and with 2.5 μM
C18:1; the FA addition allowed us to include Δ*plsX* in these comparisons (**Supplementary Fig.**
S5). Control strains (WT, *fadM::*Tn, and Δ*plsX*) showed comparable resistance to amoxicillin
in both media (0.25-0.38 μg/ml by eTest). Remarkably, all Δ*plsX* suppressors, carrying *fabF*
or *fadM* mutant alleles, showed accrued sensitivity, with similar antibiotic resistance
phenotypes (0.064-0.094 μg/ml).

Acyl-ACP species vary in JE2 WT, Δ*plsX*, and Δ*plsX* suppressors. Since PlsX uses acyl-ACP, the FASII end product, to initiate phospholipid production, its deletion provokes acyl-ACP accumulation (**Fig. 1**; [5, 34]). Exogenous FA addition resolves accumulated acyl-ACP in a Δ*plsX* mutant [12]. We assessed acyl-ACP intermediate accumulation in the JE2 Δ*plsX* mutant and derivative suppressors by non-denaturing polyacrylamide gel electrophoresis, and probing with anti-ACP antibodies [7, 35] (**Fig. 4D**). Unlike the WT strain, the Δ*plsX* mutant was distinguished by the presence of a fast-migrating band corresponding to ACP bound to long-chain FA generated by FASII, likely the dominant FA *ai*15. The Δ*plsX* strain assimilates acyl-ACP to form phosphatidic acid (PA) only if an exogenous FA (e.g. C18:1) is added to complete PA synthesis (**Fig. 1**). The long acyl-ACP species was absent in Δ*plsX fabF1* and Δ*plsX fadM1, M2*, or *M3* suppressors. This observation, and good growth of suppressor strains (**Fig. 4A**) indicate that acyl-ACP is processed to initiate phospholipid synthesis and bypass the phospholipid synthesis block in Δ*plsX*.

Δ*plsX* suppressors do not compensate reverse PlsX activity, which is required to bypass the FASII pathway. The above results show that *fabF* and/or *fadM* suppressors compensate the absence of PlsX in normal growth conditions, i.e., in converting acyl-ACP into acyl-PO$_4$. We asked whether *fabF* and/or *fadM* suppressor strains could also complement the reverse PlsX activity, i.e., conversion of acyl-PO$_4$ to acyl-ACP (**Fig. 1, Fig. 5A**). If this were the case, FASII inhibition in Δ*plsX* suppressors would be bypassed by FA supplementation to growth medium. To test this, WT strain JE2, and isogenic Δ*plsX*, Δ*plsX fabF1*, and Δ*plsX fadM2* strains were grown in FA-supplemented medium (SerFA, containing 250μM of an FA mixture and 10% fetal calf serum; see Materials and Methods), without and with the FASII inhibitor AFN-1252 [36] (**Fig. 5B**). All strains grew to equivalent densities in the absence of inhibitor. The JE2 WT strain overcame the presence of AFN-1252 in SerFA [6, 8]. In contrast, neither the Δ*plsX* mutant nor the Δ*plsX* suppressors could bypass the FASII inhibitor. These findings show that *fabF* or *fadM* suppressor mutations do not compensate the PlsX reverse reaction, and that FASII must be active for Δ*plsX* suppressor growth.

As the Δ*plsX fabF* and *fadM* suppressors behaved similarly in all the above phenotypic tests (Figs. 4,5, S4, and S5), we considered that these alleles might cooperate in a single pathway.

FadM is required for emergence of Δ*plsX* suppressors mapping to FabF. As *fadM* interruption abolished Δ*plsX* suppression (**Fig. 3**), we reasoned that suppressors arising in the

Δ*plsX fadM::*Tn mutant would map to *fabF*. This hypothesis was tested by comparing

suppressor emergence in JE2 Δ*plsX* and Δ*plsX fadM::*Tn backgrounds on solid medium

without FA supplementation. We also tested for a stimulatory effect of the FabF inhibitor

platensimycin as seen for the Δ*plsX* mutant (**Fig. 2**). Unexpectedly, no suppressors arose after

extended (> 1 week) plate incubation at 37°C, compared to frequent mutant emergence in

the Δ*plsX* background after 1-2 days (**Fig. 6A, upper**). Moreover, no stimulatory growth ring

was detected surrounding the platensimycin spot (**Fig. 6A, lower**). These results affirm that

FadM is needed to obtain FabF suppressors of Δ*plsX*. Finally, we introduced the *fadM::*Tn

allele into the Δ*plsX fabF1* suppressor mutant. Compared to good growth of Δ*plsX fabF1*,

introduction of the *fadM::*Tn allele arrested grow in the absence of exogenous FA, proving

that *fadM* is required for *fabF1* rescue (**Fig. 6B**).

The potential relevance of FabF-FadM cooperation was examined in a WT background by

comparing platensimycin inhibition zones in the WT and *fadM::*Tn derivative on BHI medium

(**Supplementary Fig. S6**). While inhibitory zones were equivalent, the WT strain displayed a

distinct growth ring (24 h incubation) that is not formed in the *fadM::*Tn mutant. Growth ring

rescue from the anti-FabF platensimycin inhibitor also requires FadM in the WT background.

Based on all the above findings, we conclude that FadM and FabF have cooperative roles.

Discussion

S. aureus requires the phospholipid synthesis enzyme PlsX for growth when FAs are not

available in its environment. Here, we show that mutations affecting either of two proteins,

FabF (a FASII core enzyme), or FadM (a bifunctional protein with acyl-CoA thioesterase and

ACP binding activities), allow *S. aureus* to grow even without this essential enzyme [20, 23,

27]. To explain growth rescue, we propose that the FabF^{A119E} variant Is responsible for

premature release of the FA-ACP intermediate. This explanation is consistent with the

shortened FAs in membrane phospholipids and with Δ*plsX* growth stimulation by low anti-

FabF concentrations. The FadM I38T, Y90F, Y133F variants all map to the acyl binding tunnel

as based on crystallographic and biochemical evidence, and likely impair acyl-CoA

thioesterase activity [20, 37]. A functional connection is established between FabF and FadM,

as both alleles must be present in order to obtain Δ*plsX* suppressors. Interestingly, all tested

phenotypes of the *fabF* and *fadM* suppressors, e.g., growth kinetics, membrane FA
shortening, and antibiotic sensitivities, were similar. While the nature of FabF-FadM
cooperation, whether direct or indirect, remains to be elucidated, these results identify FadM
as a key modulator linking FASII and phospholipid homeostasis in *S. aureus*.

We propose a working model to explain $\Delta plsX$ growth rescue by FabF and FadM variants
[38] (**Fig. 7**). In this model, the FabF^{A119E} variant (or FabF in the presence of subinhibitory
amounts of platensimycin) may stall and release its FA-ACP cargo before malonyl-ACP entry,
which is needed for FA elongation. ACP binding by FadM would facilitate ACP removal from
the FabF cavity, thus favoring FA release. The FA made available would then be
phosphorylated by Fak for entry into the phospholipid synthesis pathway.

FadM suppressor variants with modifications in their FA-binding cavity are predicted to
exhibit reduced acyl-CoA thioesterase activity, as previously observed [20] and proposed
here. This would increase the availability of FadM to facilitate ACP removal and FA release
from the FabF intermediate. In contrast, a FadM null mutation would exclude this process,
consistent with our findings that *fadM* is required for $\Delta plsX$ suppression.

The newly uncovered contribution of FadM to FabF activity may have relevance to WT *S.*
aureus FASII-phospholipid homeostasis by acting as an FA release valve that decreases FabF
processivity, leading to shortened phospholipid FAs, e.g., under stress conditions. This
hypothesis is currently being investigated.

Remarkably, the $\Delta plsX$ suppressor strains are β-lactam-sensitive as compared to the MRSA
JE2 parent or its $\Delta plsX$, and *fadM::*Tn derivatives (**Supplementary Fig. S5**). Both *fabF* and
fadM suppressors display similar greater amoxicillin sensitivity, as expected if FabF and FadM
act on the same pathway. The mechanisms underlying β-lactam-sensitivity and the potential
role of membrane synthesis or composition remain unclear, but may involve destabilization
of PBP2', as reported [33]. Thus, rerouting phospholipid synthesis rescues *S. aureus* $\Delta plsX$
growth but reduces the bacterial response to β-lactam antibiotics. These observations open
perspectives for understanding how membrane perturbations, as occur in $\Delta plsX$ suppressor
strains, might resensitize bacteria to this important class of antibiotics.

Materials and methods

Strains, growth conditions, and plasmids. Strains and plasmids are listed in **Table S4**. *S. aureus* strains are derived from USA300_FPR3757 strain JE2 [28] and RN4220-R (called here RN-R; [15]). RN-R is a readily transformable strain derived from RN4220, of the NCTC 8325 lineage [39]. It was repaired to restore a functional *fakB1* gene, which is defective in the entire NCTC 8325 lineage [15]). *S. aureus* parental strains and $\Delta plsX$-suppressor mutants were grown in BHI medium (Brain Heart Infusion; Gibco, France) or in LB (Lysogeny broth; [40]) as specified, at 37°C with shaking. Medium was supplemented with C18:1 (oleic acid) at a concentration of 250 µM (Larodan, Sweden) when indicated. The FabI inhibitor AFN-1252 (Med ChemExpress, France) and FabF inhibitor platensimycin (Bioaustralis-Tebubio France) were used as specified [18, 41].

$\Delta plsX$ **mutant construction.** The $\Delta plsX$ mutants were constructed in RN-R (RN4220 repaired for a defective *fakB1* gene in the NCTC8325 lineage; [15]) and JE2 backgrounds. The preparatory plasmid for *plsX* deletion was constructed using the Gibson assembly method to insert DNA segments flanking the *plsX orf* into Sma1-linearized pMAD, a thermosensitive plasmid expressing erythromycin resistance [42, 43]. Oligonucleotides for $\Delta plsX$ construction are in **Table S5**. Constructions were verified by DNA sequencing of PCR-amplified fragments. The double cross-over gene replacements were obtained as described [15, 43] in RN-R and JE2 strains. Deletions removed nucleotides 1,228,616-1,229,594 in JE2, and 1,148,670-1,149,648 in RN-R, as per respective annotated genomes. The resultant strain genotypes were confirmed by whole genome sequencing (Eurofins, Germany).

plsX **cloning and complementation of the** $\Delta plsX$ **mutant in RN-R and JE2 strains.** To confirm that strain characteristics were due to the *plsX* deletion, the intact *plsX* gene was cloned with its natural promoter (upstream of *fapR*) on multicopy plasmid pIM-locus1 (a pIMAY derivative [44, 45]). Cloning was performed by Gibson assembly with oligonucleotide primers (**Table S5**).

Isolation of suppressor mutants. RN-R $\Delta plsX$ and JE2 $\Delta plsX$ strains were first streaked on solid BHI medium containing 250 µM C18:1, an FA that promotes growth. For each strain, a single colony was used to inoculate a liquid culture (in BHI+C18:1). When the culture reached exponential phase, bacteria were centrifuged, resuspended in BHI, and adjusted to OD_{600} = 0.1. Dilutions were plated on solid medium to obtain ~10^5 to ~10^7 colony forming units per plate on solid BHI medium without FAs. Colonies were scored after two or more days of

incubation at 37°C, and selected at random for restreaking on solid BHI medium. Liquid cultures were prepared in BHI, and stocked in 20% glycerol for storage at -70°C. Selected mutant genotypes were confirmed by whole genome sequencing or by PCR fragment sequencing (Eurofins, Germany) to identify mutated loci.

Genomic DNA preparation for whole genome sequencing. Bacterial cell pellets, prepared from 3 mL liquid exponential phase cultures, were centrifuged at 8,000 rpm for 5 min. Pellets were re-suspended in 800 μL Tris-EDTA buffer (10 mM Tris- HC1, pH 8.0/1 mM EDTA) plus 20 μL lysostaphin 5 mg/mL, and incubated for 45 min at 37°C. DNA was prepared by DNeasy Blood & Tissue Kit (Qiagen, France) and samples were outsourced for whole genome DNA sequencing using 2x150 bp paired end chemistry and bioinformatics analyses (Eurofins Genomics, France) [6]. SNPs that differed in non-antibiotic-treated JE2 and RN-R strains from those in the reference sequence (GenBank Nucleotide accession codes NC_007793.1 and NC_007795.1 respectively) were subtracted prior to variant identification. Variants were identified as representing ≥80% of reads in sequences for which there were at least 100 reads.

PCR amplification and confirmation by DNA sequencing. Selected candidate colonies were suspended in 100 μL of 0.9% NaCl with 1 μL of lysostaphin from a 5 mg/mL stock. Tubes containing suspensions were shaken for 45 min at 37°C. Contents were then transferred to tubes containing glass beads and samples were vortexed for 1 min. After a 30 second centrifugation at 8000 rpm, PCR amplification was performed on 1 μl of supernatant, using primers (**Table S5**) and Phusion DNA polymerase (ThermoFisher Scientific, France) according to manufacturer protocol, using a SimpliAmp Thermal Cycler (ThermoFisher Scientific, France). PCR products were analyzed on a 0.8% agarose gel prepared in 40 mM Tris acetate 1 mM EDTA buffer containing 0.5 μg/mL ethidium bromide. Gels were run at 100 V for 30 min in an electrophoresis chamber (Mupid ONE; Dutscher, France), and photographed under UV light using the EBOX VX5 imaging system (Vilber Lourmat, France). PCR samples were purified using the NucleoSpin® Gel and PCR Clean-up kit (Macherey-Nagel, France), and sent for DNA sequencing (Eurofins, France).

Effects of the FabF inhibitor platensimycin on Δ*plsX* and Δ*plsX fadM::*Tn strains. Strains were grown in BHI containing C18:1 (250 μM) overnight, washed once in BHI medium, and resuspended in BHI to a final density of OD_{600} = 0.1. From this, 100 μl of the bacterial suspension was plated on solid BHI medium. Platensimycin (1.5 μg in 3 μl) was spotted in the center of plates, which were then incubated at 37°C for 4-7 days and photographed.

Effect of glucose on growth of Δ*plsX* suppressors. Strains were precultured in liquid LB (n.b., LB lacks glucose) or LB with C18:1 for Δ*plsX* at 37°C. After overnight growth, cultures were diluted to OD_{600} = 0.05 and grown for 3-4 hours. Δ*plsX* cultures were first centrifuged and resuspended in fresh LB medium to remove free FAs prior to regrowth. Bacterial densities were then adjusted to OD_{600} = 1. Cultures were streaked on solid medium, corresponding to LB or LB containing 0.5% glucose, which represses *fad* operon expression [25, 27]. Plates were incubated at 37°C for 24 hours and photographed.

Growth kinetics. Three mL precultures were prepared in BHI for parental and Δ*plsX* suppressor strains, or in BHI containing 250 μM C18:1 for Δ*plsX* mutants. After overnight growth, culture OD_{600} were adjusted to OD_{600} = 0.05 in 200 μL BHI medium without or with C18:1 (250 μM), in 96-well microtiter plates. Growth was followed on a Spark M10 plate reader (Tecan, France). For experiments in which samples were taken for FA determinations or for Western blot experiments, cultures were prepared in 3 to 10 ml liquid cultures, and were monitored manually by OD_{600} measurements every two hours. Amoxicillin sensitivity was determined using AMOXICILLIN AC 256 eTests (BioMérieux, France).

Determination of membrane FA profiles. Bacterial cultures (equivalent of OD_{600} = 2) were centrifuged, and pellets were washed once in 1 ml of 0.02% Triton in 0.9% NaCl, then twice with 1 ml of 0.9% NaCl. FA extractions were performed as described [46] [6]. Briefly, 0.5 mL of 1N sodium methoxide in methanol was added to bacterial pellets in 2 ml microfuge tubes, and then subjected to a 5 min ultrasound bath. Then 0.2 mL heptane spiked with methyl-10-undecenoate (Sigma-Aldrich) as internal standard was added, followed by 1 min vortexing, and short centrifugation for phase separation. FA methyl esters were recovered in the heptane phase. Analyses were performed in a split-splitless injection mode on an AutoSystem XL Gas Chromatograph (Perkin-Elmer) equipped with a ZB-Wax capillary column (30 m x 0.25 mm x 0.25 μm; Phenomenex, France). Data were recorded and analyzed by TotalChrom Workstation (Perkin-Elmer). *S. aureus* FA peaks were detected between 12 and 32 min of elution, and identified with retention times of purified esterified FA standards.

Detection of ACP products. Pellets from OD_{600}=10 equivalent of exponential phase cultures were collected by centrifugation (8000 rpm for 5 min), and then washed twice with PBS (Phosphate Buffer Saline)-Triton X100 0.1%, and once with PBS. Pellets were re-suspended in 600 μL PBS supplemented with 6 μL antiprotease (100X), 50 μg/mL lysostaphin and 50 μg/mL DNAse and incubated with shaking for 25 min at 37°C. Protein concentrations were determined [47], and extracts (10 μg) were loaded onto non-denaturing gels containing 1M urea and 14% acrylamide gel as described [5, 7]. After electrophoresis, gels were subject to

electroblotting for transfer to a PVDF membrane (Biorad, France) using a current of 25 V and

1.3 A on a power blotter semi-dry transfer system (Invitrogen, France) for 7 min. Primary

anti-ACP antibody (Covalabs, France; [7]) was used at 1:500 dilution and secondary anti-

rabbit IgG conjugated with peroxidase (Invitrogen, France) at a 1:5000 dilution. ACP-reactive

bands were identified using the ECL kit (Perkin Elmer, France) and the fluorescent signal was

recorded using ChemiDoc (BioRad, France).

Phage transductions. Transductions of *fadM::*Tn transposon insertion from JE2 into Δ*plsX*

suppressors *fadM2* and *fabF1* were performed as described using ɸ80 phage stock [48] [49].

The *fadM* transposon insertion [28] was transduced into JE2 Δ*plsX fadM2* suppressor strain

7.1 and into JE2 Δ*plsX fabF1* suppressor strain 7.8, giving rise respectively to Δ*plsX fadM::*Tn

and Δ*plsX fabF1 fadM::*Tn, using erythromycin (5 µg/ml) as selection (**Table S4**). Insertions

were confirmed by PCR.

FASII bypass of WT and Δ*plsX* derivative strains. Anti-FASII bypass was evaluated as

described [6]. Strains were grown overnight in liquid SerFA medium (SerFA, BHI containing

10 % newborn calf serum and an equimolar mixture of 83 µM each C14:0, C16:0, and C18:1

[Larodan, Sweden]). Cultures were then diluted to OD_{600}=0.05 in SerFA containing or not 0.5

µg/ml of the anti-FASII AFN-1252 (here called AFN; [36]; MedChemExpress, France). Growth

was monitored using the Spark plate reader (Tecan, France).

Whole-genome sequencing data of S. aureus Δ*plsX* mutants and suppressors. Data is

available in the Mendeley database: DOI: 10.17632/mzw2429w3m.1 .

Acknowledgements and funding

We are grateful to Kam Pou Han and Philippe Bouloc (Bouloc lab, I2BC, Orsay, France) for

providing transducing phage and plasmid; Florence Dubois-Brissonet and Katia de Oliveira

(Micalis, INRAE) for expert help and advice with CpG analyses; Candice Rigoulay for a

simplified phage transduction protocol. We thank Micalis Institute colleagues Vincent Juillard

for discussion of PlsX functions, and Philippe Gaudu, Jasmina Vidic, and team members for

stimulating discussion and pertinent comments. We thank the PAPPSO proteomics facility

(https://doi.org/10.15454/1.5572393176364355E12 ; supported by Paris-Saclay University,

INRAE, AgroParisTech, the Ile-de-France Regional Council, and the IBiSA network) for an

exploratory study that inspired β-lactam sensitivity testing.

This work was supported by the French Agence Nationale de la Recherche ANR (ANR-16-

CE15-0013; AG), the Joint Programming Initiative on Antimicrobial Resistance (JPIAMR)

Data availability

All relevant data are within the manuscript, supporting information files, and repositories.

Any biological materials are available upon request.

Author Contributions

All authors performed, analyzed, and interpreted experiments, and approved the

manuscript. PW, JAM, and AG conceived the project, designed research, and wrote the

paper.

Competing interests

The authors declare no competing interests.
509
510
511

References

1.   Zhang YM, Rock CO. Membrane lipid homeostasis in bacteria. Nat Rev Microbiol. 2008;6(3):222-33. Epub 2008/02/12. doi: 10.1038/nrmicro1839. PubMed PMID: 18264115.

2.   Silbert DF, Ruch F, Vagelos PR. Fatty acid replacements in a fatty acid auxotroph of *Escherichia coli*. J Bacteriol. 1968;95(5):1658-65. Epub 1968/05/01. doi: 10.1128/jb.95.5.1658-1665.1968. PubMed PMID: 4870280; PubMed Central PMCID: PMCPMC252192.

3.   Lu YJ, Zhang YM, Grimes KD, Qi J, Lee RE, Rock CO. Acyl-phosphates initiate membrane phospholipid synthesis in Gram-positive pathogens. Mol Cell. 2006;23(5):765-72. Epub 2006/09/05. doi: 10.1016/j.molcel.2006.06.030. PubMed PMID: 16949372.

4.   Parsons JB, Broussard TC, Bose JL, Rosch JW, Jackson P, Subramanian C, et al. Identification of a two-component fatty acid kinase responsible for host fatty acid incorporation by *Staphylococcus aureus*. Proc Natl Acad Sci U S A. 2014;111(29):10532-7. Epub 2014/07/09. doi: 10.1073/pnas.1408797111. PubMed PMID: 25002480; PubMed Central PMCID: PMCPMC4115530.

5.   Parsons JB, Frank MW, Jackson P, Subramanian C, Rock CO. Incorporation of extracellular fatty acids by a fatty acid kinase-dependent pathway in *Staphylococcus aureus*. Mol Microbiol. 2014;92(2):234-45. doi: 10.1111/mmi.12556. PubMed PMID: 24673884; PubMed Central PMCID: PMC4007170.

6.   Kenanian G, Morvan C, Weckel A, Pathania A, Anba-Mondoloni J, Halpern D, et al. Permissive Fatty Acid Incorporation Promotes Staphylococcal Adaptation to FASII Antibiotics in Host Environments. Cell Rep. 2019;29(12):3974-82 e4. Epub 2019/12/19. doi: 10.1016/j.celrep.2019.11.071. PubMed PMID: 31851927.

7.   Morvan C, Halpern D, Kenanian G, Hays C, Anba-Mondoloni J, Brinster S, et al. Environmental fatty acids enable emergence of infectious *Staphylococcus aureus* resistant to FASII-targeted antimicrobials. Nat Commun. 2016;7:12944. doi: 10.1038/ncomms12944. PubMed PMID: 27703138; PubMed Central PMCID: PMC5059476.

8.   Wongdontree P, Millan-Oropeza A, Upfold J, Lavergne JP, Halpern D, Lambert C, et al. Oxidative stress is intrinsic to staphylococcal adaptation to fatty acid synthesis antibiotics. iScience. 2024;27(4):109505. Epub 2024/04/05. doi: 10.1016/j.isci.2024.109505. PubMed PMID: 38577105; PubMed Central PMCID: PMCPMC10993138.

9.   Sastre DE, Pulschen AA, Basso LGM, Benites Pariente JS, Marques Netto CGC, Machinandiarena F, et al. The phosphatidic acid pathway enzyme PlsX plays both catalytic and channeling roles in bacterial phospholipid synthesis. J Biol Chem. 2020;295(7):2148-59. Epub 2020/01/11. doi: 10.1074/jbc.RA119.011147. PubMed PMID: 31919098; PubMed Central PMCID: PMCPMC7029103.

10.   Rex AN, Simpson BW, Bokinsky G, Trent MS. PlsX and PlsY: Additional roles beyond glycerophospholipid synthesis in Gram-negative bacteria. mBio. 2024:e0296924. Epub 2024/10/30. doi: 10.1128/mbio.02969-24. PubMed PMID: 39475235.

11.   Jing F, Cantu DC, Tvaruzkova J, Chipman JP, Nikolau BJ, Yandeau-Nelson MD, et al. Phylogenetic and experimental characterization of an acyl-ACP thioesterase family reveals significant diversity in enzymatic specificity and activity. BMC Biochem. 2011;12:44. Epub 2011/08/13. doi: 10.1186/1471-2091-12-44. PubMed PMID: 21831316; PubMed Central PMCID: PMCPMC3176148.

12.   Parsons JB, Frank MW, Eleveld MJ, Schalkwijk J, Broussard TC, de Jonge MI, et al. A thioesterase bypasses the requirement for exogenous fatty acids in the *plsX* deletion of

Streptococcus pneumoniae. Mol Microbiol. 2015;96(1):28-41. Epub 2014/12/24. doi: 10.1111/mmi.12916. PubMed PMID: 25534847; PubMed Central PMCID: PMCPMC4415690.

13.   Zou Q, Dong H, Cronan JE. Growth of *Enterococcus faecalis* Δ*plsX* strains is restored by increased saturated fatty acid synthesis. mSphere. 2023;8(4):e0012023. Epub 2023/06/08. doi: 10.1128/msphere.00120-23. PubMed PMID: 37289195; PubMed Central PMCID: PMCPMC10449490.

14.   Cross B, Garcia A, Faustoferri R, Quivey RG. PlsX deletion impacts fatty acid synthesis and acid adaptation in *Streptococcus mutans*. Microbiology (Reading). 2016;162(4):662-71. Epub 2016/02/07. doi: 10.1099/mic.0.000252. PubMed PMID: 26850107; PubMed Central PMCID: PMCPMC4892098.

15.   Pathania A, Anba-Mondoloni J, Gominet M, Halpern D, Dairou J, Dupont L, et al. (p)ppGpp/GTP and Malonyl-CoA Modulate *Staphylococcus aureus* Adaptation to FASII Antibiotics and Provide a Basis for Synergistic Bi-Therapy. mBio. 2021;12(1). Epub 2021/02/04. doi: 10.1128/mBio.03193-20. PubMed PMID: 33531402; PubMed Central PMCID: PMCPMC7858065.

16.   Rashid R, Nair ZJ, Chia DMH, Chong KKL, Cazenave Gassiot A, Morley SA, et al. Depleting Cationic Lipids Involved in Antimicrobial Resistance Drives Adaptive Lipid Remodeling in *Enterococcus faecalis*. mBio. 2023;14(1):e0307322. Epub 2023/01/12. doi: 10.1128/mbio.03073-22. PubMed PMID: 36629455; PubMed Central PMCID: PMCPMC9973042.

17.   Subrahmanyam S, Cronan JE, Jr. Overproduction of a functional fatty acid biosynthetic enzyme blocks fatty acid synthesis in *Escherichia coli*. J Bacteriol. 1998;180(17):4596-602. Epub 1998/08/29. doi: 10.1128/JB.180.17.4596-4602.1998. PubMed PMID: 9721301; PubMed Central PMCID: PMCPMC107473.

18.   Wang J, Soisson SM, Young K, Shoop W, Kodali S, Galgoci A, et al. Platensimycin is a selective FabF inhibitor with potent antibiotic properties. Nature. 2006;441(7091):358-61. doi: 10.1038/nature04784. PubMed PMID: 16710421.

19.   Wenzel M, Patra M, Albrecht D, Chen DY, Nicolaou KC, Metzler-Nolte N, et al. Proteomic signature of fatty acid biosynthesis inhibition available for in vivo mechanism-of-action studies. Antimicrob Agents Chemother. 2011;55(6):2590-6. Epub 2011/03/09. doi: 10.1128/AAC.00078-11. PubMed PMID: 21383089; PubMed Central PMCID: PMCPMC3101464.

20.   Murad AM, Brognaro H, Falke S, Lindner J, Perbandt M, Mudogo C, et al. Structure and activity of the DHNA Coenzyme-A Thioesterase from *Staphylococcus aureus* providing insights for innovative drug development. Sci Rep. 2022;12(1):4313. Epub 2022/03/14. doi: 10.1038/s41598-022-08281-2. PubMed PMID: 35279696; PubMed Central PMCID: PMCPMC8918352.

21.   Swarbrick CMD, Nanson JD, Patterson EI, Forwood JK. Structure, function, and regulation of thioesterases. Prog Lipid Res. 2020;79:101036. Epub 2020/05/18. doi: 10.1016/j.plipres.2020.101036. PubMed PMID: 32416211.

22.   Butland G, Peregrin-Alvarez JM, Li J, Yang W, Yang X, Canadien V, et al. Interaction network containing conserved and essential protein complexes in *Escherichia coli*. Nature. 2005;433(7025):531-7. Epub 2005/02/04. doi: 10.1038/nature03239. PubMed PMID: 15690043.

23.   Cherkasov A, Hsing M, Zoraghi R, Foster LJ, See RH, Stoynov N, et al. Mapping the protein interaction network in methicillin-resistant *Staphylococcus aureus*. J Proteome Res. 2011;10(3):1139-50. Epub 2010/12/21. doi: 10.1021/pr100918u. PubMed PMID: 21166474.

24.   Gully D, Bouveret E. A protein network for phospholipid synthesis uncovered by a variant of the tandem affinity purification method in *Escherichia coli*. Proteomics. 2006;6(1):282-93. Epub 2005/11/19. doi: 10.1002/pmic.200500115. PubMed PMID: 16294310.

25.   Kuiack RC, Tuffs SW, Dufresne K, Flick R, McCormick JK, McGavin MJ. The *fadXDEBA* locus
of *Staphylococcus aureus* is required for metabolism of exogenous palmitic acid and in vivo growth.
Mol Microbiol. 2023;120(3):425-38. Epub 2023/07/28. doi: 10.1111/mmi.15131. PubMed PMID:
37501506.
26.   Menjivar C, DeMars ZR, Wiemels RE, Carroll RK, Bose JL. Characterizing the *Staphylococcus*
aureus fatty acid degradation operon. J Bacteriol. 2025;207(8):e0008925. Epub 2025/07/17. doi:
10.1128/jb.00089-25. PubMed PMID: 40673679; PubMed Central PMCID: PMCPMC12369327.
27.   Mader U, Nicolas P, Depke M, Pane-Farre J, Debarbouille M, van der Kooi-Pol MM, et al.
Staphylococcus aureus Transcriptome Architecture: From Laboratory to Infection-Mimicking
Conditions. PLoS Genet. 2016;12(4):e1005962. doi: 10.1371/journal.pgen.1005962. PubMed PMID:
27035918; PubMed Central PMCID: PMC4818034.
28.   Fey PD, Endres JL, Yajjala VK, Widhelm TJ, Boissy RJ, Bose JL, et al. A genetic resource for
rapid and comprehensive phenotype screening of nonessential *Staphylococcus aureus* genes. MBio.
2013;4(1):e00537-12. doi: 10.1128/mBio.00537-12. PubMed PMID: 23404398; PubMed Central
PMCID: PMC3573662.
29.   Pelz A, Wieland KP, Putzbach K, Hentschel P, Albert K, Gotz F. Structure and biosynthesis of
staphyloxanthin from *Staphylococcus aureus*. J Biol Chem. 2005;280(37):32493-8. Epub
2005/07/16. doi: 10.1074/jbc.M505070200. PubMed PMID: 16020541.
30.   Qiu X, Choudhry AE, Janson CA, Grooms M, Daines RA, Lonsdale JT, et al. Crystal structure
and substrate specificity of the beta-ketoacyl-acyl carrier protein synthase III (FabH) from
Staphylococcus aureus. Protein Sci. 2005;14(8):2087-94. Epub 2005/07/01. doi:
10.1110/ps.051501605. PubMed PMID: 15987898; PubMed Central PMCID: PMCPMC2279320.
31.   Song MD, Wachi M, Doi M, Ishino F, Matsuhashi M. Evolution of an inducible penicillin-
target protein in methicillin-resistant *Staphylococcus aureus* by gene fusion. FEBS Lett.
1987;221(1):167-71. Epub 1987/08/31. doi: 10.1016/0014-5793(87)80373-3. PubMed PMID:
3305073.
32.   Antimicrobial Resistance C. Global burden of bacterial antimicrobial resistance in 2019: a
systematic analysis. Lancet. 2022;399(10325):629-55. Epub 2022/01/24. doi: 10.1016/S0140-
6736(21)02724-0. PubMed PMID: 35065702.
33.   Garcia-Fernandez E, Koch G, Wagner RM, Fekete A, Stengel ST, Schneider J, et al. Membrane
Microdomain Disassembly Inhibits MRSA Antibiotic Resistance. Cell. 2017;171(6):1354-67 e20.
Epub 2017/11/07. doi: 10.1016/j.cell.2017.10.012. PubMed PMID: 29103614; PubMed Central
PMCID: PMCPMC5720476.
34.   Machinandiarena F, Nakamatsu L, Schujman GE, de Mendoza D, Albanesi D. Revisiting the
coupling of fatty acid to phospholipid synthesis in bacteria with FapR regulation. Mol Microbiol.
2020;114(4):653-63. Epub 2020/07/17. doi: 10.1111/mmi.14574. PubMed PMID: 32671874.
35.   Cronan JE, Thomas J. Bacterial fatty acid synthesis and its relationships with polyketide
synthetic pathways. Methods Enzymol. 2009;459:395-433. Epub 2009/04/14. doi: 10.1016/S0076-
6879(09)04617-5. PubMed PMID: 19362649; PubMed Central PMCID: PMCPMC4095770.
36.   Karlowsky JA, Kaplan N, Hafkin B, Hoban DJ, Zhanel GG. AFN-1252, a FabI inhibitor,
demonstrates a Staphylococcus-specific spectrum of activity. Antimicrob Agents Chemother.
2009;53(8):3544-8. doi: 10.1128/AAC.00400-09. PubMed PMID: 19487444; PubMed Central
PMCID: PMC2715641.
37.   Jumper J, Evans R, Pritzel A, Green T, Figurnov M, Ronneberger O, et al. Highly accurate
protein structure prediction with AlphaFold. Nature. 2021;596(7873):583-9. Epub 2021/07/16. doi:
10.1038/s41586-021-03819-2. PubMed PMID: 34265844; PubMed Central PMCID:

PMCPMC8371605 have filed non-provisional patent applications 16/701,070 and
PCT/EP2020/084238, and provisional patent applications 63/107,362, 63/118,917, 63/118,918,
63/118,921 and 63/118,919, each in the name of DeepMind Technologies Limited, each pending,
relating to machine learning for predicting protein structures. The other authors declare no
competing interests.

38. Mindrebo JT, Patel A, Kim WE, Davis TD, Chen A, Bartholow TG, et al. Gating mechanism of
elongating beta-ketoacyl-ACP synthases. Nat Commun. 2020;11(1):1727. Epub 2020/04/09. doi:
10.1038/s41467-020-15455-x. PubMed PMID: 32265440; PubMed Central PMCID:
PMCPMC7138838.

39. Herbert S, Ziebandt AK, Ohlsen K, Schafer T, Hecker M, Albrecht D, et al. Repair of global
regulators in *Staphylococcus aureus* 8325 and comparative analysis with other clinical isolates.
Infect Immun. 2010;78(6):2877-89. Epub 2010/03/10. doi: 10.1128/IAI.00088-10. PubMed PMID:
20212089; PubMed Central PMCID: PMCPMC2876537.

40. Bertani G. Studies on lysogenesis. I. The mode of phage liberation by lysogenic *Escherichia
coli*. J Bacteriol. 1951;62(3):293-300. Epub 1951/09/01. doi: 10.1128/jb.62.3.293-300.1951.
PubMed PMID: 14888646; PubMed Central PMCID: PMCPMC386127.

41. Hunt T, Kaplan N, Hafkin B. Safety, tolerability and pharmacokinetics of multiple oral doses
of AFN-1252 administered as immediate release (IR) tablets in healthy subjects. Journal of
chemotherapy. 2016;28(3):164-71. doi: 10.1179/1973947815Y.0000000075. PubMed PMID:
26431470.

42. Arnaud M, Chastanet A, Debarbouille M. New vector for efficient allelic replacement in
naturally nontransformable, low-GC-content, gram-positive bacteria. Appl Environ Microbiol.
2004;70(11):6887-91. Epub 2004/11/06. doi: 10.1128/AEM.70.11.6887-6891.2004. PubMed PMID:
15528558; PubMed Central PMCID: PMCPMC525206.

43. Biswas I, Gruss A, Ehrlich SD, Maguin E. High-efficiency gene inactivation and replacement
system for gram-positive bacteria. J Bacteriol. 1993;175(11):3628-35. Epub 1993/06/01. doi:
10.1128/jb.175.11.3628-3635.1993. PubMed PMID: 8501066; PubMed Central PMCID:
PMCPMC204764.

44. Coronel-Tellez RH, Pospiech M, Barrault M, Liu W, Bordeau V, Vasnier C, et al. sRNA-
controlled iron sparing response in Staphylococci. Nucleic Acids Res. 2022;50(15):8529-46. Epub
2022/07/30. doi: 10.1093/nar/gkac648. PubMed PMID: 35904807; PubMed Central PMCID:
PMCPMC9410917.

45. Monk IR, Shah IM, Xu M, Tan MW, Foster TJ. Transforming the untransformable: application
of direct transformation to manipulate genetically *Staphylococcus aureus* and *Staphylococcus
epidermidis*. mBio. 2012;3(2). Epub 2012/03/22. doi: 10.1128/mBio.00277-11. PubMed PMID:
22434850; PubMed Central PMCID: PMCPMC3312211.

46. Rozes N, Garbay S, Denayrolles M. A rapid method for the determination of bacterial fatty
acid composition. Lett Appl Microbiol. 1993;17(3):126-31. doi: https://doi.org/10.1111/j.1472-
765X.1993.tb01440.x.

47. Bradford MM. A rapid and sensitive method for the quantitation of microgram quantities
of protein utilizing the principle of protein-dye binding. Anal Biochem. 1976;72:248-54. Epub
1976/05/07. doi: 10.1006/abio.1976.9999. PubMed PMID: 942051.

48. Olson ME. Bacteriophage Transduction in *Staphylococcus aureus*. Methods Mol Biol.
2016;1373:69-74. Epub 2015/02/04. doi: 10.1007/7651_2014_186. PubMed PMID: 25646608.

49. Bowring JZ, Su Y, Alsaadi A, Svenningsen SL, Parkhill J, Ingmer H. Screening for Highly
Transduced Genes in *Staphylococcus aureus* Revealed Both Lateral and Specialized Transduction.

Microbiol Spectr. 2022;10(1):e0242321. Epub 2022/02/10. doi: 10.1128/spectrum.02423-21. PubMed PMID: 35138167; PubMed Central PMCID: PMCPMC8826898.

50.   Fujita Y, Matsuoka H, Hirooka K. Regulation of fatty acid metabolism in bacteria. Mol Microbiol. 2007;66(4):829-39. Epub 2007/10/09. doi: 10.1111/j.1365-2958.2007.05947.x. PubMed PMID: 17919287.

51.   Brown ED. Microbiology: antibiotic stops 'ping-pong' match. Nature. 2006;441(7091):293-4. Epub 2006/05/20. doi: 10.1038/441293a. PubMed PMID: 16710404.

52.   Pettersen EF, Goddard TD, Huang CC, Couch GS, Greenblatt DM, Meng EC, et al. UCSF Chimera--a visualization system for exploratory research and analysis. J Comput Chem. 2004;25(13):1605-12. Epub 2004/07/21. doi: 10.1002/jcc.20084. PubMed PMID: 15264254.

53.   Rock CO, Cronan JE, Jr., Armitage IM. Molecular properties of acyl carrier protein derivatives. J Biol Chem. 1981;256(6):2669-74. Epub 1981/03/25. PubMed PMID: 7009596.

54.   Cronan JE, Jr. Molecular properties of short chain acyl thioesters of acyl carrier protein. J Biol Chem. 1982;257(9):5013-7. Epub 1982/05/10. PubMed PMID: 7040391.

55.   Brinster S, Lamberet G, Staels B, Trieu-Cuot P, Gruss A, Poyart C. Type II fatty acid synthesis is not a suitable antibiotic target for Gram-positive pathogens. Nature. 2009;458(7234):83-6. doi: 10.1038/nature07772. PubMed PMID: 19262672.

56.   Kuhn S, Slavetinsky CJ, Peschel A. Synthesis and function of phospholipids in *Staphylococcus aureus*. Int J Med Microbiol. 2015;305(2):196-202. Epub 2015/01/18. doi: 10.1016/j.ijmm.2014.12.016. PubMed PMID: 25595024.

57.   Monk IR, Stinear TP. From cloning to mutant in 5 days: rapid allelic exchange in *Staphylococcus aureus*. Access Microbiol. 2021;3(2):000193. Epub 2021/06/22. doi: 10.1099/acmi.0.000193. PubMed PMID: 34151146; PubMed Central PMCID: PMCPMC8209637.

58.   Christie GE, Matthews AM, King DG, Lane KD, Olivarez NP, Tallent SM, et al. The complete genomes of *Staphylococcus aureus* bacteriophages 80 and 80alpha--implications for the specificity of SaPI mobilization. Virology. 2010;407(2):381-90. Epub 2010/09/28. doi: 10.1016/j.virol.2010.08.036. PubMed PMID: 20869739; PubMed Central PMCID: PMCPMC2952651.

Fig. 1: Roles of the reversible PlsX enzyme in phospholipid synthesis. The FASII pathway schematized here produces acyl-ACP (FA-ACP), which may either be elongated by FASII using FabF (blue), or used by PlsX (purple) to initiate phospholipid (PL) synthesis. The PlsX product FA-PO$_4$ is a substrate for PlsY, a membrane protein that joins the acyl group to the glycerophosphate backbone in position 1 (①, bottom). PlsC then uses this intermediate product (lysophosphatidic acid), to add the second acyl moiety from FA-ACP to position 2 (②), generating phosphatidic acid (PA, left). Reverse PlsX activity uses phosphorylated exogenous FAs (green) produced by the FA kinase Fak [4], to produce FA-ACP. FASII inhibition (dashed red T) is bypassed by exogenous FAs [6], which are converted to FA-PO$_4$. The latter is used by PlsY to fill position 1, and by reverse PlsX, followed by PlsC, to fill position 2, thus producing PA. In a Δ*plsX* mutant, FAs charging position 1 are exogenous, while those charging position 2 are FASII-synthesized; in this case, the two FA sources are disconnected. See [1, 6, 9, 12] for references. Exogenous FA are subject to β-oxidation by FA degradation enzymes particularly in glucose-starved conditions (Fad, lower right; [25, 26, 50]). Enzymes circled in grey are not discussed in this study.

Fig. 2. *fabF* mutants or the FabF inhibitor platensimycin rescues Δ*plsX* growth in the absence of exogenous FAs. A. Lawns of RN-R Δ*plsX* were spread on BHI medium without FA addition, without (left) or with (right) 1.5 µg platensimycin (P) deposited in the center. Plates were photographed after 4 days at 37°C. Black arrowheads, examples of suppressors, some of which map to *fabF*. Dashed arrow (left), a homogenous growth ring surrounds the platensimycin spot. N=4. **B.** Working model for Δ*plsX* growth rescue by a FabF mutation (FabF mut) or by partial FabF inhibition by platensimycin (anti-FabF). FabF perturbations would reduce enzyme processivity, such that the unstable FabF-FA intermediate releases FAs prior to second substrate malonyl-ACP loading, as needed to bypass the Δ*plsX* defect. FabF in blue; FAs in green. Symbols for FabF are based on [51].

763
764
Fig. 3. FadM variants mapping to the predicted FA binding tunnel, but not FadM
inactivation, confer the *fadM* suppressor phenotype. A. Schematized *S. aureus* FadM
monomer binding to FA, based on crystal structure and Alphafold predictions ([20, 37],
designed with https://alphafoldserver.com and http://www.cgl.ucsf.edu/chimera [52]). FA
moiety (black line) binding in the FadM cavity involves amino acids Ile38 (pink), Tyr45; Met48,
Leu122, Tyr125, and Phe126 (purple) [20]. The Δ*plsX* suppressors mapping to FadM affect
amino acids Ile38Thr, Tyr90Phe, or Tyr133Phe (pink, and 2 red, respectively), which cluster
around the FA-binding cavity; Ile38 (pink) is common to structural predictions and the
FadM[I38T] Δ*plsX* suppressor. **B.** FadM inactivation abolishes Δ*plsX* suppression: Growth of
Δ*plsX fadM2* and the Δ*plsX fadM::*Tn derivative was compared on solid BHI medium without
or with C18:1. Plates were photographed after 48 h growth at 37°C. Background growth on
BHI (left) may be due to FA carryover from pre-cultures or to FA traces in medium. Plates are
representative of biological triplicates. **C.** Working model for FadM *versus* FadM mutant
function. Binding of the acyl-CoA FA moiety (zigzag line) in the FadM tunnel is needed for
acyl-CoA thioesterase activity (scissor). FadM mutations in the tunnel (red dot) prevent this
activity (red X) but conserve ACP binding (thick arrow, lower right).

781
782
Fig. 4. Behavior of *fabF* and *fadM* suppressors of Δ*plsX* without exogenous FAs. A. Growth
of WT *S. aureus* (RN-R or JE2), Δ*plsX*, and Δ*plsX fabF* or Δ*plsX fadM* suppressors in BHI
medium**.** Left: RN-R WT, and derivatives: Δ*plsX*, and Δ*plsX* suppressors *fabF1* (encoding
FabF^{A119E}) and *fadM2* (encoding FadM^{Y90F}). Right: JE2 WT, and derivatives: Δ*plsX*, and Δ*plsX*
suppressors *fabF1*, *fadM1*, *fadM2*, and *fadM3* (encoding FabF^{A119E}, FadM^{I38T}, FadM^{Y90F}, and
FadM^{Y133F} respectively). Mean and standard deviation of independent triplicate cultures are
shown. **B.** BHI + C18:1-grown overnight cultures of the indicated strains were pelleted, and
photographed to compare pigmentation. C18:1 was added to include Δ*plsX* in comparisons.
C. FA profiles of RN-R (left)- and JE2 (right)- derived WT, Δ*plsX*, and *fabF* or *fadM* suppressors
determined from BHI-grown cultures. FA assignments are as indicated. *ai*, anteiso; even
numbered FAs are saturated (e.g., C16 is C16:0). The main FA component of WT *S. aureus* is
*ai*15 [7]. Peak heights correspond to relative responses (mV) of each FA in a sample. The
proportions (in %) of C18:0 plus C20:0 of total FAs are indicated over the brackets. **Table S3**
provides complete FA profile information. **D**. Acyl-ACP species in *S. aureus* JE2-derived Δ*plsX*,
and Δ*plsX* suppressor strains. JE2 WT, Δ*plsX fadM1*, Δ*plsX fadM3*, and Δ*plsX fabF1* (isolates
7.5, 8.7, and 7.7, respectively; **Table S2**) were grown in BHI. The Δ*plsX* strain was pre-grown
in BHI+C18:1 medium, and FA-starved for 2h in BHI prior to harvesting. Extracts were loaded
on non-denaturing acrylamide gels, and immunoblotted using anti-ACP [7, 35, 53]. Longer FA
moieties on acyl-ACP migrate faster on gels (wedge; [35, 54]). White arrow, accumulated
acyl-ACP produced by FASII is not processed to phospholipids [5] (**Fig. 1**). Results are
representative of biological duplicates.

Fig. 5. Synthetic lethal effect of a combined FASII and PlsX block in Δ*plsX* and suppressor
strains. A. Expected effects of FASII inhibition on an *S. aureus* Δ*plsX* mutant. In a WT strain,
FASII inhibition by mutation or by antibiotic action is overcome by exogenous FAs, which are
phosphorylated by Fak [4]. FA-PO$_4$ is the substrate for PlsY and for reverse PlsX activities, and
is thus used for phospholipid (PL) synthesis at both the 1 and 2 positions. If both FASII and
PlsX are blocked as shown, exogenous FAs can only fill position 1. The FASII defect or PlsX
absence can be rescued separately by FA supplementation, but not the combination. Green,
exogenous FAs and FA kinase; purple, *plsX* deletion; grey, pathways and products affected by
combined FASII and PlsX block. **B.** Effects of anti-FASII treatment on JE2 Δ*plsX*, and Δ*plsX*
suppressor strains in FA-supplemented medium. Strains were precultured in SerFA, and
grown in SerFA without or with addition of 0.5 μg/ml anti-FASII AFN-1252. When the anti-
FASII is added, WT *S. aureus* strains incorporate FAs to constitute membrane phospholipids,
thus bypassing inhibition [6, 8, 55]. As Δ*plsX* suppressors cannot generate FA-ACP needed for
FASII bypass [6, 9, 12], they remain sensitive to anti-FASII. Mean and standard deviation of
independent triplicate cultures are shown. Black lines, no anti-FASII, red lines with anti-FASII.
fadM2 encodes FadM^{Y90F}; *fabF1* encodes FabF^{A119E}.

Fig. 6. *fabF* suppressors of Δ*plsX* require the presence of a *fadM* allele. A. FadM is present
in JE2 Δ*plsX* (left) and absent in Δ*plsX fadM::*Tn (right, constructed from strain 7.1: **Table S2**).
Strains were plated on BHI solid medium (100 µl of OD_{600} = 0.005), and in the lower set,
spotted with platensimycin (P; 1.5 µg). Plates were photographed after 6 days at 37°C. Black
arrowheads point to Δ*plsX* suppressor clones. The dashed curved line indicates the growth
ring surrounding platensimycin spotting. The small colonies surrounding platensimycin on
Δ*plsX fadM::*Tn lawns (right) failed to regrow on BHI medium. **B.** FadM is present in the JE2
Δ*plsX fabF1* suppressor strain (left) and absent in Δ*plsX fabF1 fadM::*Tn (right, constructed
from strain 7.8: **Table S2**). Strains were streaked on BHI solid medium (upper), or BHI medium
containing 250 µM C18:1 and 5 µg/ml erythromycin (Ery; lower). Plates were photographed
after 24 h incubation at 37°C. Results are representative of three independent cultures and
platings.

Fig. 7. Proposed role of FabF - FadM interactions in controlling FA release and recycling from FabF intermediates. Our study showed epistasis between FabF and FadM, and leads us to a working model of how these proteins interact (depicted in the inset). Left, FASII and phospholipid synthesis (PL) coupled pathways [1, 56]. The *plsX* deletion dissociates the pathways, such that exogenous FA is required for survival. Symbols are as in **Fig. 1**. **Inset:** FabF - FadM interactions coordinately overcome Δ*plsX* growth arrest when either function is mutated. In the first FASII step, FabF (blue packman and blue arrow) cleaves acyl-ACP (FA-ACP) to release ACP, which then allows malonyl-ACP entry into the FabF cavity to extend the acyl intermediate by 2 carbons [18, 38]. The FabF[A119E] variant (FabFmut), or addition of anti-FabF inhibitor platensimycin, impedes malonyl-CoA entry, leading to FA release (FA symbols in green). FadM (gold) is bifunctional, with acyl-CoA thioesterase and ACP binding activities [20, 22-24]. Suppressor mutations in FadM (FadMmut red dot) alter the FadM acyl binding cavity, which we propose decreases acyl-CoA thioesterase activity as reported for FadM mutants also mapping to the binding cavity [20]. More FadMmut is thus available for ACP binding (boxed FadMmut). FadM-ACP or FadMmut-ACP binding could favor ACP release from the FabF pocket and leakage of FAs, which are then processed for PL synthesis *via* FA kinase (Fak) [4]. Symbols for FabF are based on [51].

898 **Fig. S1. FASII and phospholipid gene organization in RN-R and JE2 strains.** Locus tags of
899 genes related to this study are presented. Dark green, FASII initiation; blue, FASII elongation;
900 black, acyl carrier protein (*acpP*); purple, phospholipid synthesis; olive, FASII bypass fatty acid
901 kinase; pink, regulatory; orange, FadM; grey, not directly related. Locus tags are given for
902 USA300 FPR_3797 8325-4 as references.

Fig. S2. Complementation of Δ*plsX* by a plasmid-carried *plsX* gene. RN-R Δ*plsX* strains containing an empty plasmid carrying a chloramphenicol resistance cassette (pIMAY; left side of plates), or a plasmid carrying the *plsX* gene expressed from its native promoter on pIMAY (pPlsX; right side of plates), were streaked on BHI-chloramphenicol (Cm) 10 µg/ml solid medium, without or with 250 µM C18:1 to bypass the Δ*plsX* defect. Background growth on BHI (left) may be due to FA carryover from pre-cultures or to FA traces in medium. Plates were photographed after 48h growth at 30°C. The Δ*plsX* strain carrying the empty plasmid failed to grow in the absence of C18:1 supplementation, while the pPlsX-complemented strain grew in both media.

Fig. S3. JE2ΔplsX growth stimulation by platensimycin in liquid culture. JE2 ΔplsX was grown
overnight in BHI medium containing C18:1. After washing, the strain was resuspended to OD_{600} = 0.05
and used to inoculate BHI medium to which platensimycin was added at the concentrations indicated
below graphs. Left, growth curves with the different platensimycin concentrations added to medium.
Right, the optimal concentration leading to improved growth with platensimycin (red line) compared
to growth in BHI medium (black line). The averages of three independent triplicates are shown.
Differences are statistically significant (P=≤0.05) by 2-tailed bilateral T-test for all points comparing
ΔplsX growth in BHI and BHI plus platensimycin 0.125 μg/ml) between 7 h and 12 h growth.

Fig. S4. FadM^m acyl-CoA thioesterase activity is not needed for Δ*plsX* suppression. A. The FadM^m suppressor phenotype is not subject to glucose repression, suggesting that the Fad-mediated acyl-CoA synthesis degradation pathway (Fad) is not involved. Fad enzyme FadE converts FA to acyl-CoA, a FadM substrate [20]. The *fad* operon is subject to CcpA-glucose repression [25-28], suggesting that acyl-CoA pools would drop in glucose-containing medium. Strains precultured in LB and LB with C18:1 for Δ*plsX* (Materials and Methods) were streaked directly on solid LB medium without or with 0.5% glucose. Background growth on LB (left) may be due to FA carryover from pre-cultures or FA traces in medium. Upper and lower rows, strains are derived from JE2 and from RN-R, respectively. Results represent biological triplicates.

Fig. S5. MRSA JE2 ΔplsX suppressor mutants are sensitized to the β-lactam antibiotic amoxicillin. The indicated JE2 and ΔplsX derivative strains were grown overnight in BHI medium without (**A**) or with 250 μM C18:1 (**B**). Cultures were resuspended to OD_{600} = 0.1 and 100 μl were spread on the cognate solid media. Amoxicillin Etest strips were placed on plates, followed by 48 h incubation at 37°C. The amoxicillin MICs are indicated in micrograms per milliliter below each photo.

JE2   *fadM*::Tn

Fig. S6. FadM is required for platensimycin-stimulated ring formation in the WT background.
Overnight BHI cultures of JE2 and *fadM::*Tn (SAUSA300_1247) were washed in 0.9% NaCl, pellets were
resuspended to OD_{600} = 0.1, and 250 µl was spread on 120 x 120 mM square petri plates containing
solid BHI medium. Three µg platensimycin were spotted on plates ('P' in white circle; N=2). Plates were
photographed at 48 h. Vertical arrow, inhibition zone; left arrow, growth ring surrounds the
platensimycin deposit site in the WT, but not the *fadM::*Tn strain.

Table S2. Δ*plsX* suppressors [a,b].

	Gene : protein mutations in Δ*plsX* suppressor mutants				
	SAUSA300_0886 FabF (3-oxoacyl-ACP synthase II)		SAUSA300_1247 FadM (acyl-CoA thioesterase/ACP binding [20, 22-24])		
Strain name	*fabF1*: FabF^{A119E}	*fabF2*: FabF^{D266A}	*fadM1*: FadM^{I38T}	*fadM2*: FadM^{Y90F}	*fadM3*: FadM^{Y133F}
RN4220 5.1	X		NT	NT	NT
RN4220 5.2	X		NT	NT	NT
RN4220 5.3	X		NT	NT	NT
RN4220 5.4	X		NT	NT	NT
RN4220 5.5	X		NT	NT	NT
RN4220 5.6				X	
RN4220 5.7				**X**	
RN4220 5.8				X	
RN4220 6.1	X		NT	NT	NT
RN4220 6.2	X		NT	NT	NT
RN4220 6.3	X		NT	NT	NT
RN4220 6.4	X		NT	NT	NT
RN4220 6.5	X		NT	NT	NT
RN4220 6.6	X		NT	NT	NT
RN4220 6.7				X	
RN4220 6.8	X		NT	NT	NT
JE2 7.1				**X**	
JE2 7.2			X		
JE2 7.3				X	
JE2 7.4			X		
JE2 7.5			X		
JE2 7.7	X		NT	NT	NT
JE2 7.8	X		NT	NT	NT
JE2 8.1			X		
JE2 8.2				X	
JE2 8.3			X		
JE2 8.4			X		
JE2 8.5			X		
JE2 8.6			X		
JE2 8.7					X
JE2 8.8					X
JE2 Sup1 [b]	X				
JE2 Sup2 [b,c]		X			

[a] 'X' indicates that the mutation is present in the tested isolate. NT, not tested; without indication, no mutation detected by DNA sequencing. Suppressors identified from whole genome sequencing are in bold. [b] Sup1 and Sup2 suppressors of Δ*plsX* had the same secondary mutation in SAUSA300_100 (generating Csa1A^{K185Q}), which was absent in subsequently isolated suppressors mapping to *fabF*. [c] Sup2, encoding FabF^{D266A}, could not be propagated after storage.

[b] Accompanying Excel Table S1. Sequence comparisons of Δ*plsX* suppressors and reference strains.

 Table S4. Genetic materials.

Strains			
Host strain	**Strain**	**Description**	**Reference**
JE2 USA300_FPR3757	JE2 (WT)	USA300_FPR3757 cured of its three native plasmids	[28]
	Δ*plsX*	In-frame *plsX* deletion leaving operon intact	This study
	Δ*plsX* suppressor in *fadM*; Δ*plsX fadM1*	*fadM1* encodes FadM[I38T]	This study
	Δ*plsX* suppressor in *fadM*; Δ*plsX fadM2*	*fadM2* encodes FadM[Y90F]	This study
	Δ*plsX* suppressor in *fadM*; Δ*plsX fadM3*	*fadM3* encodes FadM[Y133F]	This study
	Δ*plsX* suppressor in *fabF*; Δ*plsX fabF1*	*fabF1* encodes FabF[A119E]	This study
	fadM Tn insertion	JE2 derivative containing an erythromycin-resistance-marked transposon (Tn) insertion in SAUSA300_1247 at position 1384915 (based on JE2 genome entry NZ_CP020619)	[28]
	Δ*plsX fadM::*Tn	Φ80 phage transduction of the SAUSA300_1247 Tn insertional mutant into the Δ*plsX fadM2* (7.1, **Table S2**) suppressor	This study
RN-R 4220	RN-R 4220 (WT)	RN4220 derivative repaired for *fakB1*	[15]
	Δ*plsX*	In-frame deletion of *plsX* leaving operon intact	This study
	Δ*plsX* suppressor in *fadM*; Δ*plsX fadM2*	*fadM2* encodes FadM[Y90F]	This study
	Δ*plsX* suppressor in *fabF*; Δ*plsX fabF1*	*fabF1* encodes FabF[A119E]	This study
Plasmids			
	Plasmid	**Description**	**Reference**
	pMAD	Thermosensitive plasmid used to generate double-crossover deletion events	[42, 43]
	pMAD-*fapR*-*fabD* fragments	Used to generate *plsX* deletion	This study
	pIMAY	Used for *plsX* cloning	[44, 57]
	pIMAY-*plsX*	Used for *plsX* complementation in Δ*plsX*	This study
Transducing phage			
	Phage	**Description**	**Reference**
	φ80	Used to transduce erythromycin-resistance-marked Tn insertion of SAUSA300_1247 into Δ*plsX fadM2* and Δ*plsX fabF1* backgrounds	[58]

1010   **Table S5. Oligonucleotides (5' to 3') used for genetic constructions and mutant detection**.

Construction type and primers used for detection		Description
Plasmid for Δ*plsX* construction on pMAD linearized by SmaI		
pMAD_FapR_F	TCTATCGATGCATGCCATGGTACCCCTTCATCACAGACCATGAACTAAGC	*fapR* fragment
FapR_FabD_R	CGGAAAAATAATTGCTGTTTTACTCATAATTTTATCCTCGCTTATCATAAAAC	
FapR_FabD_F	GTTTTATGATAAGCGAGGATAAAATTATGAGTAAAACAGCAATTATTTTTCCG	*fabD* fragment
FabD_pMAD_R	GCAGAAGCTTCTAGAATTCGAGCTCCCTGTGACCTGAAACAACAATTTGACC	
Verifications for pMAD constructions		
pMAD_F	GTTACGTTACACATTAACTAGACAG	
pMAD_R	TGGACAGCATGGCCTGCAACGC	
Cloning *plsX* on pIMAY-derived locus1 plasmid		
pIMAY_FapR_F	AGTTTGTGGTTTAATGGGGGTTGACACTTGAACTATTTAAA	*fapR* fragment
FapR_PlsX_R	TCAATTGCTAATTTAACCATTTTTTAGTACCTAGTCTTAA	
FapR_PlsX_F	TTAAGACTAGGTACTAAAAAATGGTTAAATTAGCAATTGA	*plsX* fragment
plsX_pIMAY_R	TTGTTGCTATAATGAAATGCGTTCTTACTCATTTGATTCACCTA	
pIMAY_FapR_R	TTTAAATAGTTCAAGTGTCAACCCCATTAAACCACAAACT	pIMAY linearized by reverse PCR
plsX_pIMAY_F	TAGGTGAATCAAATGAGTAAGAACGCATTTCATTATAGCAACAA	
Δ*plsX* suppressor verifications		
FabF_F	GGAATCAGCTAGAGAACGC	SAUSA300_0886
FabF_R	CATTTTTCAACCCACAAGAGG	
1247_F	AATCGGTATTTGAGGCAC	SAUSA300_1247
1247_R	ATCTTCACGCTAATTACATT	
0100_F	CGTGGCCTGGATATGATGAT	SAUSA300_0100 *csa1*
0100_R	GATTATCCTGAGCTACTTCG	
Verifications of Tn insertion in SAUSA300_1247		
1247_F2	GGATTACTATCGTCACGGTGG	Transposon insertion
UpBursa_F	CTCGATTCTATTAACAAGGG	
1247_R2	GTGTCCTGAACCTCGGAAGTTG	

1011