Title: Known unknowns and model selection in ecological evidence synthesis

Author: Shane A. Blowes, German Centre for Integrative Biodiversity Research (iDiv) Halle-

Jena-Leipzig, Germany; Department of Computer Science, Martin Luther University Halle-

Wittenberg, Halle (Salle), Germany. shane.blowes@idiv.de

Corresponding author: Shane Blowes, German Centre for Integrative Biodiversity Research

(iDiv) Halle-Jena-Leipzig, Deutscher Platz 5e, Leipzig 04103, Germany. shane.blowes@idiv.de.

Abstract (200 word limit)

Quantitative evidence synthesis is a prominent path towards generality in ecology. Generality is typically discussed in terms of central tendencies, such as an average effect across a compilation of studies, and the role of heterogeneity for advancing understanding is not as well developed. Heterogeneity quantifies the consistency of ecological effects across studies, and between-study heterogeneity is typically assumed as constant. Here, I relax the assumption of constant heterogeneity, and show how cross validation can further the generality goals of quantitative evidence synthesis. First, I examine scale-dependent heterogeneity for a meta-analysis of plant native-exotic species richness relationships, quantifying the relationships among unexplained effect size variation, spatial grain and extent. Second, I examine relationships among patch size, study-level covariates and unexplained variation in species richness using a database of fragmentation studies. Both case studies show that relaxing the assumption of constant unexplained variation can provide a more detailed description of available evidence - for example, where effects can be transferred with more or less certainty. Moreover, cross validation shows for both case studies that assuming constant heterogeneity can constrain model predictive performance, and in particular, the ability of model predictions to generalise to new studies.

Introduction

Quantitative evidence synthesis aims for general insights into the direction, magnitude and variability of ecological effects (Gurevitch et al. 2018, Fox 2019, Spake et al. 2022). Two common forms of quantitative synthesis are meta-analysis (i.e., analyses of effect sizes collated or calculated from existing studies), and analyses of raw data compilations (referred to as full data analysis; Spake et al. 2022). For both approaches, generalities typically come in the form of central tendencies, such as an average effect size across studies. While both approaches

frequently quantify the heterogeneity of effect sizes across studies to examine how consistent

ecological effects are (Senior et al. 2016, Gurevitch et al. 2018, Spake et al. 2022, Nakagawa et

al. 2023a), any role for heterogeneity in assessing generality is less well developed. Critically,

quantitative syntheses typically assume either constant between-study heterogeneity (meta-

analysis) or residual variation (i.e., homoscedasticity for full data analyses). This assumption is

equivalent to assuming that included studies sample effect sizes from a single population

(Williams et al. 2021). Despite the rapid rise of meta-analysis in ecology and conservation

biology (Gurevitch et al. 2018, Anderson et al. 2021), neither the consequences of this

assumption, nor the opportunities available when it is relaxed, have received much attention

from quantitative ecologists doing evidence synthesis.

Ecological meta-analyses often fit "random effects" or multilevel meta-analytic models

(Gurevitch et al. 2018, Nakagawa et al. 2023a). These models incorporate two forms of

uncertainty. First, sampling error or variance is represented by uncertainty in the effect size (i.e.,

response variable), and assumed known (with methods available for inferring missing values for

some commonly used effect sizes, e.g., Nakagawa et al. 2023b). The second form of uncertainty,

between-study heterogeneity or variance, is typically modelled as a varying (or random)

intercept, and assumed to be constant across studies; i.e., estimated with a single standard

deviation across all studies. Multilevel meta-analytic models estimate additional within study

heterogeneity (via an additional varying intercept), and are recommended when included studies

provide multiple (non-independent) effect sizes (Nakagawa et al. 2023a). Recently introduced

(location-scale) statistical models for meta-analysis relax the assumption of constant between-

study uncertainty with an extension to allow predictors of heterogeneity (i.e., of the standard

deviation of the varying intercepts; Williams et al. 2021, Viechtbauer & Lopez-Lopez 2022,

Rodriguez et al. 2023). Simulations show little is lost in terms of power and type-I error rates

(false rejection of the null hypothesis) when the assumption of constant between-study variances

is relaxed (Rodriguez et al. 2023). Evidence syntheses using full data compilations typically

make qualitatively similar assumptions: effects are allowed to vary between studies (e.g., varying

slopes and intercepts), and most often estimated assuming constant residual variation across

studies (i.e., homoscedasticity). Importantly, for both meta-analyses (Williams et al. 2021) and

full data analyses (Pinherio & Bates 2000, Zuur et al. 2009), relaxing the assumption of constant

heterogeneity or heteroscedasticity can impact estimates of other parameters, such as the overall

mean or study-level effects.

Examining unexplained variation can open up new research questions (Williams et al.

2021, Viechtbauer & Lopez-Lopez 2022). Quantifying known unknowns - for example, by using

covariates to describe patterns in unexplained variation – represents an extension to the majority

of existing ecological evidence syntheses, and can help identify promising directions for both

empirical and theoretical research. For example, covariates strongly associated with unexplained

variation could also help to more clearly define limits to the generality of quantitative evidence

syntheses (Spake et al. 2022). If different groups, such as taxon groups or geographic regions are

associated with more or less unexplained variation, researchers could better communicate where

effect sizes can be transferred with more or less uncertainty.

To compare models assuming homogeneous versus heterogeneous unexplained variation

I use cross validation. To date, model selection in evidence synthesis has largely focused on the

amount of variance explained (e.g., R^2, Nakagawa et al. 2023a), or on predictive performance

assessed with information criteria (Cinar et al. 2021). R^2 describes how much variation is

accounted for by different model components (e.g., meta-regression, or hierarchical partitions of

heterogeneity), and allow for comparisons of the total amount of variation explained by different

covariates or model components. In contrast, cross validation and information criteria assess

model predictive performance (Hastie et al. 2009, Cinar et al. 2021). However, explicit

consideration of the data used to train versus test models (Hastie et al. 2009), and the need to

define a function to be optimized (Hastie et al. 2009, Yates et al. 2022) make cross validation

more flexible than information criteria. For example, data splitting can be designed to compare

the ability of models to make different types of predictions (e.g., Merkle et al. 2019, Yates et al.

2022). In the context of quantitative evidence synthesis, I consider two types of predictions: (1)

to a new datum or data within the existing studies; or (2), for a new study, i.e., data outside the

sample of studies used for model fitting.

I show how location-scale models and cross-validation for model selection can be used in

quantitative evidence synthesis for ecology using two biodiversity synthesis case studies. First, I

extend a recent meta-analysis of spatial scale-dependence in plant native-exotic species richness

relationships (Peng et al. 2019) to quantify relationships among unexplained variation, grain size

and spatial extent. The second case study uses a data compilation of habitat fragment diversity

studies (Chase et al. 2019). I focus on the relationship between fragment size and local (i.e.,

patch-scale) species richness, examine whether residual variation is related to fragment size and

other study-level predictors. For both case studies, the assumption that unexplained variation

comes from a single population limits model predictive performance, particularly when making

predictions to new studies.

Cross validation for model comparison in evidence synthesis

Cross validation uses data splitting techniques to test model predictive performance.

Models are fit to a 'training' data set and assessed on their ability to predict the 'test' data set

(Hastie et al 2009, Yates et al. 2022). To quantify predictive performance, I used pointwise

expected log predictive density (*elpd*; see Yates et al. 2022 for further discussion of other

common loss functions). To minimize an inherent bias to support overfit models, I used the

modified one-standard-error rule (Yates et al. 2021) that aims to select the least complex model

that is comparable to the best scoring model. I evaluate model performance for making two

different types of predictions (Merkle et al. 2019, Yates et al. 2022): (i) conditional predictions

that examine model performance when making predictions to existing studies, and (ii) marginal

predictions that assess predictive performance to new studies. Conditional predictions are

qualitatively similar (and quantitively the same for some loss functions, Yates et al. 2022) to

those made by other information criteria methods (e.g., AIC). However, if generality is

considered as the focus for a quantitative evidence synthesis, it might be more valuable for

models to make good predictions to new studies (i.e., outside the sample of studies used to

parameterize the model). To illustrate this, I use model selection based on comparisons of

marginal predictions for a left-out study as an example in the two case studies.

I assess conditional and marginal predictions using two different data splitting

techniques. For conditional predictions, approximate or exact leave-one-out (loo) cross

validation is considered the gold standard (Yates et al. 2022). However, approximate loo

diagnostics (Vehtari et al. 2017) showed many influential observations for the models I fit in the

case studies (which reduces the reliability of performance estimates, Vehtari et al. 2017), and

because exact loo can be computationally expensive for large data compilations (due to the need

to refit the model as many times as there are data points), I approximated loo using *k*-fold cross

validation. *k*-fold cross validation converges on loo when the number of blocks (*k*) into which the

data are split equals the number of data points. The majority of quantitative evidence syntheses,

including the case studies presented here, use mixed or multilevel models, and so I use a

stratified *k*-fold approximation of loo. Stratified *k*-fold cross validation balances data splitting

among subgroups when creating the *k* blocks of data to which models are refit, ensuring that the

relative group frequencies (here, the data coming from individual studies included in my

analyses) are maintained. Similar to loo (and other information criteria such as AIC), *k*-fold cross

validation is focused on making predictions to new data points within existing studies

conditional on the model parameters. However, the most common goals of evidence synthesis –

intervention evaluation and generality (Gurevitch et al. 2018) - might also be assessed by

examining how well models make predictions to new studies. To assess the predictive

performance of models to new studies, I use leave-one-group-out cross validation, where

individual studies are removed one at a time, and models are assessed on their ability to predict

the held-out data.

Models and model fitting

Location-scale statistical models allowing predictor variables for both the mean (location)

and residual variation (scale) are well studied (Pinheiro & Bates 2000 and references therein). To

relax the assumption of constant heterogeneity, recent meta-analytic models fit predictors for the

standard deviation of varying (random) intercepts (e.g., Williams et al. 2021, Rodriguez et al.

2022, Viechtbauer & Lopez-Lopez 2022), covariates that appear as predictors for the location

parameter. These models admit only the known sampling error associated with individual effect

sizes to contribute to the scale component of the likelihood. Here, I describe an alternative

location-scale model for meta-analysis, where unexplained variation is modelled by covariates

for the scale parameter (that are added to the known error variances; see model 1.2 - 1.4 below; a

brief comparison of the two approaches shows that they have very similar parameter estimates,

but differ slightly in the model comparisons for the meta-analytic case study presented here:
Appendix S1). Questions being addressed with analyses of full data compilations can also model
heteroscedasticity in multiple ways. Models can be fit with predictors for the scale parameter
(e.g., when using Gaussian or log-normal likelihoods), or for parameters related to the scale (e.g.,
negative binomial or beta distributions). Many R packages commonly used by ecologists allow
for mixed (or multilevel) location-scale models to be fit using either frequentist (e.g., *nlme*,
Pinheiro & Bates 2024; *glmmTMB*, Brooks et al. 2017) or Bayesian methods (e.g., *brms*, Bürkner
2017), which I adopt for the case-studies presented here.

To robustly build increasingly complex models of heterogeneity and heteroscedasticity I
used simulation-based calibration (Gelman et al. 2018, Talts et al. 2020, Modrák et al. 2023).
This process uses simulated data to check model calibration, and I focus on whether models can
recover known parameter values using simulated data sets of the same size, shape and structure
as that found in each of the case studies. Simulation-based calibration requires a data generating
model, and I followed to same procedure for each model in both case studies. First, the model is
fit to the empirical (i.e., real) data. Then, the same model with priors informed by the fit to the
empirical data is used to simulate many new data sets (with the same size, shape and structure as
the empirical data). Then, based on the results of the calibration (i.e., are known parameters of
interest recovered with reasonable coverage?), I proceed with model building towards the goal of
models that characterize heterogeneity and heteroscedasticity. All models were fit using the
Hamiltonian Monte Carlo (HMC) sampler Stan (Carpenter et al. 2017) due to the availability of
tools for diagnosing model pathologies (Betancourt 2016, Monnahan et al. 2017), as well as for
cross validation (Vehtari et al. 2017). Models fit to empirical data used weakly regularizing
priors, four chains each with 4000 iterations, with 2000 used for warmup (details for case study

one in Appendix S2, and case study two Appendix S3). Models were coded using the 'brms'

package (Bürkner 2017). Code for all analyses is available at

https://github.com/sablowes/heterogeneity-evidence-synthesis (that will be archived following

acceptance).

Case study one: Scale-dependent uncertainty in plant native-exotic richness relationships

Communities with more species are often thought to be more resistant to invasion by

exotic (or non-native) species than communities with fewer species (Elton 1958), an idea

referred to as the 'biotic resistance hypothesis'. However, the spatial scale-dependence of

biodiversity complicates any straightforward interpretation of this hypothesis. Negative

relationships between the numbers of native and non-native species are often thought more likely

at small scales, with a switch to positive relationships expected at larger spatial scales (Levine

2000). Such scale-dependence has been linked to niche opportunities (Shea & Chesson 2002),

whereby at relatively small scales, more species rich native communities leave fewer niche

opportunities for exotics to invade. As spatial scale increases, so too does heterogeneity in

resources, natural enemies and the physical environment, which creates greater niche availability

for exotic species to become established (Shea & Chesson 2002).

Recently, Peng et al. (2019) used multilevel meta-regressions to synthesize evidence for

relationships between spatial scale (both grain and extent) and the correlation of native and

exotic species richness. Using 101 observational (i.e., unmanipulated, non-experimental) studies

and 204 effect sizes, Peng et al. (2019) found that the average effect size was positive, and that

the positive correlations between native and exotic species richness got stronger with increasing

(log) grain size. No relationship was found between spatial extent and the native-exotic richness

relationship. All of the models examined by Peng et al. (2019) assumed between-study variation

in the effect size associated with the native-exotic richness relationship, and adjusted for the non-

independence of multiple effect sizes coming from some studies with (nested) multilevel varying

(random) intercepts. Here I relax the assumption that between study heterogeneity is constant,

and examine whether unexplained variation in the effect size is related to either grain size or

extent. I compare how all models perform when making predictions to existing and new studies

using cross validation.

203   Peng et al. (2019) transformed all correlations to Pearson product moment correlation

coefficient (r), and calculated effect sizes using the Fisher z-transformation: $z = \frac{1}{2} log \left(\frac{1+r}{1-r} \right)$,

with variance estimated as $Var(z) = \frac{1}{n-3}$, log refers to the natural logarithm (as it does

hereafter), and n is the sample size (i.e., number of native-exotic species pairs; effect size data

sourced from Peng et al. 2019). I start by reproducing the main result from Peng et al. (2019)

using a multilevel linear meta-regression model for grain size that assumes the effect sizes, z_{ij},

are normally distributed with known within-case variance (s_{ij}^2) and constant between-study

variance (τ^2), which can be expressed as following:

211                             $$z_{ij} \sim N(\mu_{ij}, s_{ij}),$$

212                             $$\mu_{ij} = \beta_0 + \beta_{0i} + \beta_{0ij} + \beta_1 X_{ij},$$

213                             $$\beta_{0i} \sim N(0, \tau),$$

214                             $$\beta_{0ij} \sim N(0, \omega), \hspace{3cm} \text{(Model 1.1)}$$

where cases (j) are nested within studies (i), and have among-case (within study) variance, ω^2;

study-level heterogeneity, β_{0i}, has variance τ^2, and varies around the overall linear relationship

with intercept, β_0, slope β_1, and predictor X_i (here the natural logarithm of grain size in study i).

In this model, both τ and ω quantify variation not accounted for by the population (fixed) linear

predictor ($\beta_0 + B_1 X_{ij}$), and are sometimes referred to residual heterogeneity and interpreted as

residuals (e.g., Viechtbauer & Lopez-Lopez 2022, Nakagawa et al. 2023a). Given the strong

relationship between effect size and grain size (Peng et al. 2019), I retain (log) grain size as a

predictor in all subsequent models.

Peng et al. (2019) observed that variation around the average (overall positive)

relationship between grain size and effect sizes decreased with increasing grain size. That is

heterogeneity decreased as a function of grain size, though they did not explicitly model this

relationship. Indeed, the most widely used statistical computing software for meta-analysis has

only recently been extended to allow predictors for residual heteroscedasticity (Viechtbauer &

Lopez-Lopez 2022). Here I start by fitting a model that estimates additional parameters for

residual variation (i.e., the scale parameter of the likelihood) from the data. Recall that s_{ij} is the

known within-study standard deviation of the effect size estimate, and unexplained variation in

existing meta-analytic location-scale models include predictors for a varying intercept (e.g., τ or

ω in model 1.1). An alternative is to include another parameter to be estimated from the data for

residual variation, and I fit a model where an additional parameter (σ_{ij}) for residual variation is a

function of grain size:

235                             $$z_{ij} \sim N(\mu_{ij}, s_{ij} + \sigma_{ij}),$$

236                             $$\mu_{ij} = \beta_0 + \beta_{0i} + \beta_1 X_i,$$

237                             $$\beta_{0i} \sim N(0, \tau),$$

238                             $$log(\sigma_{ij}) = \beta_0^\sigma + \beta_1^\sigma X_{ij}, \qquad \text{(Model 1.2)}$$

where X_{ij} is grain size on a log-scale for the jth case in study i. I note that this model did not

produce reliable estimates when I included the case-level random intercept (ω), likely due to

identification problems associated with multiple parameters (i.e., σ and ω) describing the same

level of variation in the data, and the case-level random intercept was dropped from the model.

To model extent, Peng et al. (2019) discretized spatial extent into bins with a range of one

order of magnitude in km^2, i.e., (0, 10), [10, 100), [10^2, 10^3), …, [10^6,). Peng et al. (2019) fit

multiple meta-regression models to examine for an effect of spatial extent on effect sizes,

including models with an interaction between grain and extent. While Peng et al. (2019) did not

report a strong influence of spatial extent on effect sizes, they did describe a complex interaction

between extent and grain. Here, I examine whether unexplained variation (i.e., after adjusting for

the relationship between effect size and grain size) was related to extent using a model where

residual variation was a function of spatial extent. I fit extent as a categorical predictor of the

residual variation (σ_{ij}, i.e., same as previous model, but without the intercept (β_0^ω), and extent

categories coded as indicator variables for the predictor, X_{ij} [Model 1.3]).

Finally, I fit a model that allowed study-level residual variation. I note that where data are

limited to one effect size per study this model will have identification problems. Here, the data

have some replication within studies (mean = 2; median = 1; range = 1-17 effect sizes per study),

and simulation-based calibration suggested reasonable coverage for parameters in the order of

those found for the empirical data with the observed sample sizes (Appendix S2: Fig. Sx). Again,

I fit a model fit with (log) grain size as a predictor for the mean effect size, and estimated

residual variation with studies varying around an overall average:

$$z_{ij} \sim N(\mu_{ij}, s_{ij} + \sigma_{ij}),$$

$$\mu_{ij} = \beta_0 + \beta_{0i} + \beta_1 X_i,$$

$$\beta_{0i} \sim N(0, \tau),$$

$$log(\sigma_{ij}) = \beta_0^\sigma + \beta_{0i}^\sigma,$$

264
$$\beta_{0i}^{\sigma} \sim N(0, \zeta), \qquad\qquad \text{(Model 1.4)}$$

265   where β_0^{σ} is the average residual variation (on a log-scale), and β_{0i}^{σ} is a normally distributed

266   study-level varying intercept with zero mean and ζ standard deviation.

267   *Case study one: results and discussion*

268   All models were able to reproduce the observed data well, and simulation-based

calibration showed that all models had reasonable coverage for parameters of interest (Appendix

S2: Figs. S1-S16). Model selection identified the same model (1.4) as best for making

predictions within existing studies (Fig. 1a) and to new studies (Fig. 1b), though subsequent

ordering among the other models depended on predictive task (Fig. 1). All models produced

similar estimates for the parameters they shared (i.e., β_0, β_1, τ; Appendix S1: Fig. S3), and

support for model 1.4 is notable because this model is atypical of meta-analyses. The more

widely used constant heterogeneity meta-regression (model 1.1) performed worst for making

predictions within the observed data (Fig. 1a), and was tied for second (with all other models)

when making predictions for a new study (Fig. 1b). On average, unexplained variation decreased

with increasing grain size (as was observed qualitatively by Peng et al. 2019), though there is

considerable uncertainty surrounding the relationship for the smallest and largest grains (Fig. 1c),

and the 90% credible interval overlapped zero (model 1.2: β_1^{σ} = -0.08 – 0.015). Finally, the model

that included spatial extent (i.e., the total geographic area covered by the samples) showed that

after adjusting for the positive relationship between effect sizes and grain size, unexplained

variation was on average the greatest at the smallest extents (Fig. 1d).

Case study two: Variation in patch-scale effects of ecosystem decay is largest for the

smallest patches

Habitat loss is a major driver of biodiversity loss (Diaz et al. 2019). As habitat is removed

from landscapes, biodiversity reductions in remaining habitat fragments can be thought of as

arising from one of two processes (Chase et al. 2020). First, because fewer individuals and

species can exist in smaller habitat fragments, the species found small habitat fragments might

simply be a random (or passive) sample of those found in large habitat fragments. Alternatively,

if smaller habitat fragments negatively impact demographic rates relative to larger habitat

fragments, then ecosystem decay might result in greater diversity declines compared to those

expected from a solely passive sampling process (Chase et al. 2020).

To test these competing hypotheses for (patch-scale) diversity fragment size

relationships, Chase et al. (2019) compiled assemblage data (counts of individuals of each

species) in effort-standardized (or standardizable) samples across habitat fragments of different

sizes within landscapes. Across 123 studies comprising 1509 habitat fragments, neither the

number of individuals nor the species found in small fragments were simply a random sample of

those in large habitat fragments (Chase et al. 2020). Instead, altered demographic rates (e.g., due

to edge effects, reduced dispersal or demographic stochasticity, and collectively referred to as

ecosystem decay) in smaller habitat fragments, resulted on average in fewer individuals, fewer

species, and less even communities than expected from a passive sample of larger fragments

(Chase et al. 2020). Here, I revisit this analysis to examine for heteroscedasticity, and to see

whether modelling heteroscedasticity impacts model predictive performance. For simplicity, I

focus on species richness, and start by refitting the same multilevel model used by Chase et al.

(2020) that assumed residual variation was constant across all of the studies. I then fit a series of

models of increasing complexity that relax the assumption of homoscedasticity.

The model fit by Chase et al. (2020) to effort-standardized for species richness (S_{ij}) in

fragment j from study i took the form:

$$S_{ij} \sim lognormal(\mu_{ij}, \sigma)$$

$$\mu_{ij} = \beta_0 + \beta_{0i} + (\beta_1 + \beta_{1i})X_{ij},$$

$$[\beta_{0i}, \beta_{1i}]' \sim MVN(0, \boldsymbol{SRS}),$$

$$\boldsymbol{S} = \begin{bmatrix} \sigma_{0i} & 0 \\ 0 & \sigma_{1i} \end{bmatrix},$$

$$\boldsymbol{R} = \begin{bmatrix} 1 & \rho_{\sigma_{0i}\sigma_{1i}} \\ \rho_{\sigma_{0i}\sigma_{1i}} & 1 \end{bmatrix}, \qquad \text{(Model 2.1)}$$

where X_{ij} is the fragment size on a (natural) log-scale, which was centered by subtracting the

overall mean from each observation before modelling; β_{0i} and β_{1i} are study-level departures from

the overall intercept and slope, respectively, drawn from a multivariate normal (*MVN*)

distribution with standard deviations, σ_{0i} and σ_{1i}, that allowed for correlations (*R* matrix)

between the varying intercepts and slopes.

The first heteroscedastic model is motivated similarly to the varying study-level

intercepts and slopes. Just as average richness and its relationship with fragment size might be

expected to varying among studies around an overall average, so to might residual variation.

Therefore, I allowed varying study-level residuals around an overall average residual standard

deviation. First, I estimated study-level residual variation independently of other varying study-

level parameters for the mean:

$$S_{ij} \sim lognormal(\mu_{ij}, \sigma_i)$$

$$\mu_{ij} = \beta_0 + \beta_{0i} + (\beta_1 + \beta_{1i})X_{ij},$$

$$[\beta_{0i}, \beta_{1i}]' \sim MVN(0, \boldsymbol{SRS}),$$

329
$$S = \begin{bmatrix} \sigma_{0i} & 0 \\ 0 & \sigma_{1i} \end{bmatrix},$$

330
$$R = \begin{bmatrix} 1 & \rho_{\sigma_{0i}\sigma_{1i}} \\ \rho_{\sigma_{0i}\sigma_{1i}} & 1 \end{bmatrix},$$

331
$$log(\sigma_i) = \beta_0^\sigma + \beta_{0i}^\sigma,$$

332
$$\beta_{0i}^\sigma \sim N(0, \sigma_{0i}^\sigma), \qquad\qquad \text{(Model 2.2)}$$

where β_0^σ is the overall average of residual variation (on a log-scale), and β_{0i}^σ are the varying

study-level departures (for the residual variation) drawn from a normal distribution with zero

mean and standard deviation σ_{0i}^σ.

Next, I extend this model to include (log) fragment size as a predictor of residual

variation, and again allow it to varying between studies independently of varying study-level

parameters for the mean:

339
$$S_{ij} \sim lognormal(\mu_{ij}, \sigma_i)$$

340
$$\mu_{ij} = \beta_0 + \beta_{0i} + (\beta_1 + \beta_{1i})X_{ij},$$

341
$$[\beta_{0i}, \beta_{1i}]' \sim MVN(0, \boldsymbol{SRS}),$$

342
$$S = \begin{bmatrix} \sigma_{0i} & 0 \\ 0 & \sigma_{1i} \end{bmatrix},$$

343
$$R = \begin{bmatrix} 1 & \rho_{\sigma_{0i}\sigma_{1i}} \\ \rho_{\sigma_{0i}\sigma_{1i}} & 1 \end{bmatrix},$$

344
$$log(\sigma_i) = \beta_0^\sigma + \beta_{0i}^\sigma + (\beta_1^\sigma + \beta_{1i}^\sigma)X_i,$$

345
$$[\beta_{0i}^\sigma, \beta_{1i}^\sigma]' \sim MVN(0, \boldsymbol{S^\sigma R^\sigma S^\sigma}),$$

346
$$S^\sigma = \begin{bmatrix} \sigma_{0i}^\sigma & 0 \\ 0 & \sigma_{1i}^\sigma \end{bmatrix},$$

347
$$R^\sigma = \begin{bmatrix} 1 & \rho_{\sigma_{0i}^\sigma\sigma_{1i}^\sigma} \\ \rho_{\sigma_{0i}^\sigma\sigma_{1i}^\sigma} & 1 \end{bmatrix}, \qquad\qquad \text{(Model 2.3)}$$

348   where β_0^σ is the overall average residual variation, and β_1^σ is the overall average slope of residual

349   variation with fragment size; β_{0i}^σ and β_{1i}^σ are the varying study-level departures from the intercept

350   and slope, respectively, and were drawn from a multivariate normal distribution with zero mean

351   and standard deviation σ_{0i}^σ and σ_{1i}^σ, with correlations estimated in matrix $\boldsymbol{R}^\sigma$.

352   Finally, I estimate models with study-level residual variation with and without fragment

353   size as a predictor that allow for correlations between varying (study-level) parameters for the

354   mean (μ) and residual variation (σ). The model without fragment size is:

355
$$S_{ij} \sim lognormal(\mu_{ij}, \sigma_i)$$

356
$$\mu_{ij} = \beta_0 + \beta_{0i} + (\beta_1 + \beta_{1i})X_{ij},$$

357
$$log(\sigma_i) = \beta_0^\sigma + \beta_{0i}^\sigma,$$

358
$$[\beta_{0i}, \beta_{1i}, \beta_{0i}^\sigma]' \sim MVN(0, \boldsymbol{SRS}),$$

359
$$\boldsymbol{S} = \begin{bmatrix} \sigma_{0i} & 0 & 0 \\ 0 & \sigma_{1i} & 0 \\ 0 & 0 & \sigma_{0i}^\sigma \end{bmatrix},$$

360
$$\boldsymbol{R} = \begin{bmatrix} 1 & \rho_{\sigma_{0i}\sigma_{1i}} & \rho_{\sigma_{0i}\sigma_{0i}^\sigma} \\ \rho_{\sigma_{0i}\sigma_{1i}} & 1 & \rho_{\sigma_{1i}\sigma_{0i}^\sigma} \\ \rho_{\sigma_{0i}\sigma_{0i}^\sigma} & \rho_{\sigma_{1i}\sigma_{0i}^\sigma} & 1 \end{bmatrix}.$$
    (Model 2.4)

361   The model with fragment size as a predictor of both the mean and residuals that allows for

362   correlations among the varying study-level parameters is:

363
$$S_{ij} \sim lognormal(\mu_{ij}, \sigma_i)$$

364
$$\mu_{ij} = \beta_0 + \beta_{0i} + (\beta_1 + \beta_{1i})X_{ij},$$

365
$$log(\sigma_i) = \beta_0^\sigma + \beta_{0i}^\sigma + (\beta_1^\sigma + \beta_{1i}^\sigma)X_i,$$

366
$$[\beta_{0i}, \beta_{1i}, \beta_{0i}^\sigma, \beta_{1i}^\sigma]' \sim MVN(0, \boldsymbol{SRS}),$$

367
$$S = \begin{bmatrix} \sigma_{0i} & 0 & 0 & 0 \\ 0 & \sigma_{1i} & 0 & 0 \\ 0 & 0 & \sigma_{0i}^{\sigma} & 0 \\ 0 & 0 & 0 & \sigma_{1i}^{\sigma} \end{bmatrix},$$

368
$$R = \begin{bmatrix} 1 & \rho_{\sigma_{0i}\sigma_{1i}} & \rho_{\sigma_{0i}\sigma_{0i}^{\sigma}} & \rho_{\sigma_{0i}\sigma_{1i}^{\sigma}} \\ \rho_{\sigma_{0i}\sigma_{1i}} & 1 & \rho_{\sigma_{1i}\sigma_{0i}^{\sigma}} & \rho_{\sigma_{1i}\sigma_{1i}^{\sigma}} \\ \rho_{\sigma_{0i}\sigma_{0i}^{\sigma}} & \rho_{\sigma_{1i}\sigma_{0i}^{\sigma}} & 1 & \rho_{\sigma_{0i}^{\sigma}\sigma_{1i}^{\sigma}} \\ \rho_{\sigma_{0i}\sigma_{1i}^{\sigma}} & \rho_{\sigma_{1i}\sigma_{1i}^{\sigma}} & \rho_{\sigma_{0i}^{\sigma}\sigma_{1i}^{\sigma}} & 1 \end{bmatrix}.$$ (Model 2.5)

Case study two: results and discussion

All models were able to reproduce the observed data well, and simulation-based

calibration showed that all models had reasonable coverage for parameters of interest (Appendix

S3: Fig. S1-S20). Models that allowed for heteroscedasticity outperformed the model with

constant residual variation for making predictions of species richness in fragments for existing

studies (Fig. 2a), and to new studies (Fig. 2b). Modelling heteroscedasticity, however, did not

qualitatively impact the support for the ecosystem decay hypothesis (Chase et al. 2020) for

species richness (Appendix S3: Fig. S21). Yet, the best model for predictions to existing studies

differed from that favored for new studies. For making predictions to new data in existing

studies, cross validation supported the most complex model (model 2.5; Fig. 2a); this model

shows that residual variation was a decreasing function of fragment size (Fig. 2c), meaning

unexplained variation in the fragment size-species richness relationship is greatest for small

patches, and suggesting considerable potential for context-dependency in the response of

taxonomic diversity in small habitat fragments. Moreover, this model suggests that there were

some relationships between varying study-level effects and residual variation (Appendix S3: Fig.

S22). For example, the fragment size study-level slopes for the mean effect and residual variation

were negatively correlated (Appendix S3: Fig. S22a), meaning that the strongest ecosystem

decay effects were associated with the strongest decline in residual variation with increasing

fragment size (Appendix S3: Fig. S23).

For predictions to new studies leave-one-group-out cross validation supported a simpler,

multilevel model for residual variation (Fig. 2b), where study-level residuals varied around an

overall mean (model 2.2), meaning each study gets its own estimate of residual (unexplained)

variation. Because study encodes many other covariates (e.g., taxon group, matrix quality, time

since fragmentation), this model can be used to further examine for systematic variation in the

unexplained variation among studies by plotting posterior samples of study-level residual

variation (σ_i) against study-level covariates (see Chase et al. 2020 for a similar approach to

examining predictors of ecosystem decay effect size estimates). For example, studies of

amphibians and reptiles had more unexplained variation than average, while studies focused on

birds had less unexplained variation than average (Fig. 2d). The relatively small sample sizes (n

= 14 and n = 15 studies, respectively) precludes strong inference, and I highlight this result to

show how examining residual variation can provide a more complete description in a

quantitative evidence synthesis. For example, I might hypothesize less variation in connectivity

between patches for birds compared to amphibians and reptiles could drive this difference in

unexplained patch-scale diversity variation, and design either an empirical or theoretical study to

further examine the role of varying connectivity for fragmentation effects on patch-scale

diversity. More generally, this shows how explicit models of unexplained variation can help

communicate where the evidence for patch-scale fragmentation effects is more or less uncertain.

Discussion

When will it be reasonable to assume that studies compiled for evidence synthesis come

from single (statistical) population? And on what criteria should ecologists compare models for

evidence synthesis? Here, I relaxed the assumption that data compilations sampled a single

statistical population for two case studies, and compared model performance for two different

prediction goals. Statistical models that relaxed the assumption of homoscedastic residuals

performed best for making both types of predictions - to new data within the included studies,

and to unobserved data from a new study. Moreover, I found that model performance (i.e., the

comparative rank of candidate models) can depend on whether predictions are being made to

existing studies or to new studies, highlighting the flexibility of cross-validation for assessing

models based on different goals in an evidence synthesis context.

Meta-analysis and quantitative syntheses of compiled data are a prominent path to

generality in ecology and evolution (Gurevitch et al. 2018, Fox 2019, Anderson et al. 2021).

While heterogeneity of effect sizes in meta-analyses are frequently quantified, they are often not

reported (Senior et al. 2016, Nakagawa et al. 2023a), and the role of heterogeneity in assessing

claims of generality is not well developed (Spake et al. 2022, Yang et al. 2023). Relaxing the

assumption of constant heterogeneity in meta-analytical statistical models is relatively new

(Williams et al. 2021, Rodriguez et al. 2022, Viechtbauer & Lopez-Lopez 2022), and here I

examined an alternative to existing models with predictors for the scale parameter of the

likelihood. For the meta-analysis re-examined here, this new model produced qualitatively

similar estimates of the shared parameters with the other models examined (Appendix S1: Fig.

S3). Moreover, the estimated relationship between unexplained variation and grain size was

essentially indistinguishable between the model with the additional scale parameter (i.e., model

1.2 above) and a model comparable to existing models (Williams et al. 2021, Rodriguez et al.

2022, Viechtbauer & Lopez-Lopez 2022; Appendix S1: Fig. S2). The two models performed

similarly for predictions to new data within studies (Appendix S1: Fig. S4a), but the new model

was favored for making predictions to a new study (Appendix S1: Fig. S4b). Indeed, among the

models examined for each case study, the best predictions to a new study were made by models

with very similar structures.

Quantitative evidence syntheses in ecology and evolution routinely estimate variation in

effect sizes among studies (Gurevitch et al. 2018, Spake et al. 2024). However, unexplained

variation is most often assumed as constant, and therefore, as coming from a single (statistical)

population. In the frequently heterogeneous data that is compiled for evidence synthesis, this

assumption warrants careful examination. Here, for predictions to new data from an unobserved

study, both case studies favored a model where the scale (or residual variation) was modelled

similarly to the varying effects for the location (mean). In effect, these models admit greater

uncertainty, and this conservatism is likely what led to them being favored when making

predictions to unobserved studies.

Models that relaxed the assumption of constant heterogeneity (or homoscedasticity)

yielded new quantitative insights for both case studies presented here. First, I quantified the

pattern of decreasing unexplained variation in the number native and exotic species with

increasing grain size observed by Peng et al. (2019). I found support for the negative relationship

on average, though the model revealed considerable uncertainty in the relationship (especially

for large spatial grains where there is little data, Fig. 1c), and the 90% credible interval of the

effect overlapped zero. Similar to Peng et al. (2019), I also found some evidence for an

association between unexplained variation (i.e., after adjusting for grain size) and spatial extent,

indicating that there is considerable context dependency in the relationship between numbers of

native and exotic species, particularly at small spatial scales (i.e., small grains and small extents).

The second case study relaxed the assumption of homoscedasticity for the relationship

between patch-scale (standardized) species richness and fragment size. I found a negative

relationship between fragment size and residual variation, and that this negative relationship was

strongest for landscapes that had the steepest (positive) relationship between standardized

species richness and fragment size. Taken together, this means that variation in the patch-scale

richness is greatest for the smallest patches in fragmented landscapes, and that unexplained

variation decreases most strongly in landscapes with the strongest (patch-scale) ecosystem decay

effects. In effect, relaxing the assumption of homoscedasticity has identified another empirical

pattern that theory describing processes associated with fragmentation (e.g., Zhang et al. 2024)

might explore. I also used a simpler heteroscedastic model with varying study-level residuals to

show how quantitative descriptions of unexplained variation can reveal limitations to the

generality of an evidence synthesis (e.g., do some groups have more unexplained variation than

others), as well as providing promising new new paths for empirical and theoretical research

(e.g., *why* do some groups have more or less unexplained variation than others?). Importantly,

model selection in both case studies showed that relaxing the assumption of constant

heterogeneity or homoscedasticity can result in models with greater predictive ability.

Model selection in evidence synthesis has typically used either variance explained (R^2,

Nakagawa et al. 2023a) or information criterion methods (Cinar et al. 2021). Here, I showed how

cross-validation can be used to compare model performance for different predictive tasks.

Specifically, I compared models based on their ability to predict new data within existing studies

(conditional predictions), and on their ability to predict data for a new (unobserved) study (refs).

Cross-validation for conditional prediction (e.g., approximate or exact leave-one-out, and

different k-fold approximations to leave-one-out) are similar to information criteria in their goal

of making predictions to new data within the existing studies. However, it is in the assessment of

predictions to new studies (marginal predictions) that cross-validation can provide new insights

into model predictive performance for quantitative evidence synthesis. Here I broadly focused on

a new study in general, and defined individual studies as the group in the leave-one-group-out

cross validation, but other choices could be made depending on a particular research question or

goal for evidence synthesis. For example, limits beyond which effect size estimates or

relationships become unreliable (sometime called the transfer domain, Spake et al. 2022) of an

evidence synthesis could be assessed using cross-validation to partition studies into different

groups, e.g., based on taxonomy, geography, biogeography, phylogeny, etc. Moreover, the

ranking of models was prediction task-dependent for both case studies presented here (though

the same model was best for both predictions in case study one), which suggests that using cross-

validation to assess different prediction goals will be a useful addition to the evidence synthesis

toolbox.

Author contributions

SAB conceived and did the study.

Acknowlegements

I thank the Biodiversity Synthesis lab for constructive discussions, and for providing much

fodder for thinking about models for evidence synthesis. SAB was supported by the German

Centre for Integrative Biodiversity Research (iDiv) Halle-Jena-Leipzig, funded by the German

Research Foundation (FZT 118), and by the European Union. Views and opinions expressed are

however those of the author only, and do not necessarily reflect those of the European Union or

the European Research Council. Neither the European Union nor the granting

authority can be held responsible for them.

Conflict of interest

I declare no conflicts of interest.

References

Anderson SC, Elsen PR, Hughes BB, Tonietto RK, Bletz MC, Gill DA, et al. Trends in ecology
and conservation over eight decades. Frontiers in Ecol & Environ. 2021 Jun;19(5):274–82.

Betancourt, M. 2016, April 3. Diagnosing Suboptimal Cotangent Disintegrations in Hamiltonian
Monte Carlo. arXiv.

Brooks ME, Kristensen K, Van Benthem KJ, Magnusson A, Berg CW, Nielsen A, et al.
glmmTMB balances speed and flexibility among packages for zero-inflated generalized linear
mixed modeling. The R journal. 2017;9(2):378–400.

Bürkner PC. brms: An R package for Bayesian multilevel models using Stan. Journal of
Statistical Software. 2017;80(1):1–28.

Carpenter B, Gelman A, Hoffman MD, Lee D, Goodrich B, Betancourt M, et al. Stan: A
probabilistic programming language. Journal of statistical software. 2017;76(1).

Chase JM, Blowes SA, Knight TM, Gerstner K, May F. Ecosystem decay exacerbates
biodiversity loss with habitat loss. Nature. 2020;584(7820):238–43.

Chase JM, Liebergesell M, Sagouis A, May F, Blowes SA, Berg \AAke, et al. FragSAD: A
database of diversity and species abundance distributions from habitat fragments. Ecology.
2019;100(12).

Cinar, O., J. Umbanhowar, J. D. Hoeksema, and W. Viechtbauer. 2021. Using information-
theoretic approaches for model selection in meta-analysis. Research Synthesis Methods
12:537–556.

Díaz S, Settele J, Brondízio ES, Ngo HT, Agard J, Arneth A, et al. Pervasive human-driven

decline of life on Earth points to the need for transformative change. Science.

2019;366(6471).

Elton CS. The ecology of invasions by animals and plants. London, UK.: Methuen and

Company; 1958.

Fox JW. The many roads to generality in ecology. Philosophical Topics. 2019;47(1):83–104.

Gurevitch J, Koricheva J, Nakagawa S, Stewart G. Meta-analysis and the science of research

synthesis. Nature. 2018;555(7695):175–82.

Hastie, T., R. Tibshirani, and J. Friedman. (2009). The Elements of Statistical Learning. Second

edition. Springer New York, NY.

Levine JM. Species Diversity and Biological Invasions: Relating Local Process to Community

Pattern. Science. 2000 May 5;288(5467):852–4.

Merkle EC, Furr D, Rabe-Hesketh S. Bayesian Comparison of Latent Variable Models:

Conditional Versus Marginal Likelihoods. Psychometrika. 2019 Sep;84(3):802–29.

Modrák, M., A. H. Moon, S. Kim, P. Bürkner, N. Huurre, K. Faltejsková, A. Gelman, and A.

Vehtari. 2023. Simulation-based calibration checking for Bayesian computation: The choice

of test quantities shapes sensitivity. Bayesian Analysis 1:1–28.

Monnahan, C. C., J. T. Thorson, and T. A. Branch. 2017. Faster estimation of Bayesian models

in ecology using Hamiltonian Monte Carlo. Methods in Ecology and Evolution 8:339–348.

Nakagawa, S., Y. Yang, E. L. Macartney, R. Spake, and M. Lagisz. 2023a. Quantitative evidence

synthesis: a practical guide on meta-analysis, meta-regression, and publication bias tests for

environmental sciences. Environmental Evidence 12:8.

Nakagawa, S., D. W. A. Noble, M. Lagisz, R. Spake, W. Viechtbauer, and A. M. Senior. 2023b.

A robust and readily implementable method for the meta-analysis of response ratios with and

without missing standard deviations. Ecology Letters 26:232–244.

Peng S, Kinlock NL, Gurevitch J, Peng S. Correlation of native and exotic species richness: a

global meta-analysis finds no invasion paradox across scales. Ecology. 2019

Jan;100(1):e02552.

Pinheiro J, Bates D. Mixed-effects models in S and S-PLUS [Internet]. Springer science &

business media; 2006.

Rodriguez JE, Williams DR, Bürkner P. Heterogeneous heterogeneity by default: Testing

categorical moderators in mixed-effects meta-analysis. Brit J Math & Statis. 2023

May;76(2):402–33.

Schmid C H, Mengersen K. Bayesian Meta-analysis. In: Koricheva J, Gurevitch J, Mengersen K,

editors. Handbook of Meta-analysis in Ecology and Evolution [Internet]. Princeton University

Press; 2013 p. 145–73.

Senior AM, Grueber CE, Kamiya T, Lagisz M, O'Dwyer K, Santos ESA, et al. Heterogeneity in

ecological and evolutionary meta-analyses: its magnitude and implications. Ecology. 2016

Dec;97(12):3293–9.

Spake R, O'dea RE, Nakagawa S, Doncaster CP, Ryo M, Callaghan CT, et al. Improving

quantitative synthesis to achieve generality in ecology. Nature Ecology & Evolution.

2022;6(12):1818–28.

Talts, S., M. Betancourt, D. Simpson, A. Vehtari, and A. Gelman. 2020, October 21. Validating

Bayesian Inference Algorithms with Simulation-Based Calibration. arXiv.

Vehtari A, Gelman A, Gabry J. Practical Bayesian model evaluation using leave-one-out cross-

validation and WAIC. Stat Comput. 2017 Sep;27(5):1413–32.

Viechtbauer, W., and J. A. López-López. 2022. Location-scale models for meta-analysis.

Research Synthesis Methods 13:697–715.

Williams DR, Rodriguez JE, Bürkner PC. Putting variation into variance: modeling between-

study heterogeneity in meta-analysis. 2021.

Yang, Y., D. W. Noble, R. Spake, A. M. Senior, M. Lagisz, and S. Nakagawa. 2023. A

pluralistic framework for measuring and stratifying heterogeneity in meta-analyses.

ecoevoRxiv. https://doi.org/10.32942/X2RG7X

Yates LA, Aandahl Z, Richards SA, Brook BW. Cross validation for model selection: A review

with examples from ecology. Ecological Monographs. 2023 Feb;93(1):e1557.

Zhang, H., J. M. Chase, and J. Liao. 2024. Habitat amount modulates biodiversity responses to

fragmentation. Nature Ecology & Evolution 8:1437–1447.

Zuur AF, Ieno EN, Walker NJ, Saveliev AA, Smith GM. Mixed effects models and extensions in

ecology with R. Springer; 2009. 574 p.

Figure 1: **Heterogeneous heterogeneity outperforms constant heterogeneity for predicting**

plant native-exotic species richness relationships. Model selection for predictions (a) within

existing studies using stratified k-fold cross-validation, and (b) for new studies using leave-one-

group-out cross validation. Scale-dependent unexplained variation for (c) grain, and (d) extent.

Thick lines on (c) show median expectation for grain-size dependent unexplained variation from

model 1.2 ($log\ [\sigma_{ij}] = \beta_0^\sigma + \beta_1^\sigma X_{ij}$); thin lines show 100 draws from the posterior distribution to

visualise uncertainty. Violin plots (d) show the distribution of 1000 draws of the posterior

distribution of unexplained variation (σ) for each of the extent categories; boxplots show the

median (bar), 25% and 75% quantiles (box), whiskers show 1.5 times the interquartile range,

points show observations beyond 1.5 times the interquartile range; n = number of effect sizes for

each extent category (cases).

Figure 2: **Patch-scale species richness in habitat fragments is predicted best by models with**

heteroscedastic residual variation. Model selection for predictions (a) within existing studies

using stratified k-fold cross-validation, and (b) for predictions to new studies using leave-one-

group-out cross validation; (c) fragment size-dependent residual variation, and (d) study-level

residual variation grouped in taxon groups. Bold line on (c) shows median predicted average

relationship between fragment size and residual variation from model 2.5, thin lines show 100

draws from the posterior distribution. Density plots (d) for taxon groups show study-level

variation (1000 draws from the posterior) of residual variation from model 2.2 (i.e., $\beta_0^\sigma + \beta_{0i}^\sigma$

grouped into taxon groups); black triangle shows median σ for each taxon group, black dashed

line and surrounding shading are the overall mean (β_0^σ) and 95% credible interval.

Figure 1

 Figure 2

Appendix S1: A (case-study) comparison of two location-scale models for meta-analysis

My first extension to include heteroscedasticity for the meta-analysis case-study modelled

unexplained variation (i.e., the scale parameter, σ_{ij}) as a function of (log) grain size, which was

added to the known sampling error (s_{ij}) of the effect size. Specifically,

$$z_{ij} \sim N(\mu_{ij}, s_{ij} + \sigma_{ij}),$$

$$\mu_{ij} = \beta_0 + \beta_{0i} + \beta_1 X_i,$$

$$\beta_{0i} \sim N(0, \tau),$$

$$log(\sigma_{ij}) = \beta_0^\sigma + \beta_1^\sigma X_{ij}, \qquad \text{(Model 1.2)}$$

where X_{ij} is grain size on a log-scale for the jth case in study i; see Appendix S2 for the weakly

regularizing priors that were used. This model is atypical of meta-analyses, which normally only

include the known (or imputed if unknown) sampling error (s_{ij}) as the sole contributor to the

scale component of the likelihood. Accordingly, the recent location-scale meta-analytic models

instead include covariates for varying (random) intercepts in meta-regressions (e.g., Williams et

al. 2021, Rodriguez et al. 2022, Viechtbauer & Lopez-Lopez 2022). To compare the two

approaches I fit a model that includes grain size as a predictor the case-level varying intercept.

Specifically, as each case had a grain, and some studies had multiple cases (i.e., more than one

grain size), I modelled case-level variation (ω_j) as a linear function of (log) grain size (X_{ij}) with a

log-link to ensure positive values for ω:

$$z_{ij} \sim N(\mu_{ij}, s_{ij}),$$

$$\mu_{ij} = \beta_0 + \beta_{0i} + \beta_{0ij} + \beta_1 X_i,$$

$$\beta_{0i} \sim N(0, \tau),$$

$$\beta_{0ij} \sim N(0, \omega_{ij}),$$

$$log(\omega_{ij}) = \beta_0^\omega + \beta_1^\omega X_{ij}. \qquad \text{(Model 1.2a)}$$

I fit this model with similar weakly regularizing priors to that of model 1.2:

$$\beta_0 \sim N(0.3, 1),$$

$$\beta_0^\omega, \beta_1^\omega, \beta_1 \sim N(0, 1),$$

$$\tau \sim N(0, 1).$$

Model 1.2a had good convergence (all Rhats < 1.01), and was able to reproduce the observed

effect sizes well (Appendix S1: Fig. S1).

639
640   **Figure S1:** Posterior predictions from model 1.2a were able to reproduce the observed effect
641   sizes relatively accurately.
642

643   The two models (1.2 and 1.2a) predicted similar relationships between unexplained variation and

644   grain size (Appendix S1: Fig. S2), and had indistinguishable estimates for shared parameters

645   (Appendix S1: Fig. S3). Model 1.2 and 1.2a had similar performance when making predictions

646   within existing studies (Appendix S1: Fig. S4a), however model 1.2 was favored when making

647   predictions to new studies (Appendix S1: Fig. S4b).

648
649 **Figure S2:** Models 1.2 and 1.2a predicted very similar relationships between unexplained
650 variation and grain size.
651

653
654   **Figure S3**: All models produced qualitative similar estimates for shared parameters (β_0, β_1, τ).
655

657
658 **Figure S4: Heterogeneous heterogeneity outperforms constant heterogeneity for predicting**
659 **plant native-exotic species richness relationships.** Model selection for predictions (a) within
660 existing studies using stratified k-fold cross-validation, and (b) for new studies using leave-one-
661 group-out cross validation.
662
663 **References**
664

Rodriguez, J. E., D. R. Williams, and P. Bürkner. 2023. Heterogeneous heterogeneity by default:

Testing categorical moderators in mixed-effects meta-analysis. British Journal of

Mathematical and Statistical Psychology 76:402–433.

Viechtbauer, W., and J. A. López-López. 2022. Location-scale models for meta-analysis.

Research Synthesis Methods 13:697–715.

Williams, D. R., J. E. Rodriguez, and P.-C. Bürkner. 2021. Putting variation into variance:

modeling between-study heterogeneity in meta-analysis.

672
673

Appendix S2

Case-study one: Simulation-based calibration and supplemental figures

I used Simulation-Based Calibration (Talts et al. 2020, Modrák et al. 2023) to check

whether models could recover known parameter values with reasonable coverage (i.e., the

probability that a constructed [e.g., credible] interval contains the true value). Briefly, this

involves: (1) fitting a model to the empirical data; (2) using the same model with priors informed

by the fit to the empirical data to simulate many new (fake) datasets with the same size, shape

and structure as the empirical data (i.e., the same number of observations, and the same number

of groups for each level in the hierarchical structure); (3) refitting the model to each simulated

data set; and, (4) calculating SBC diagnostics and plotting results of the model fits to simulated

data to check for reasonable coverage (Talts et al. 2020, Modrák et al. 2023).

Model 1.1

I begin case study one by reproducing the main result from Peng et al. (2019) using a

multilevel linear meta-regression model for grain size that assumes the effect sizes, z_{ij}, are

normally distributed with known within-case variance (s_{ij}^2) and constant between-study variance

(τ^2), which can be expressed as:

$$z_{ij} \sim N(\mu_{ij}, s_{ij}),$$

$$\mu_{ij} = \beta_0 + \beta_{0i} + \beta_{0ij} + \beta_1 X_{ij},$$

$$\beta_{0i} \sim N(0, \tau),$$

$$\beta_{0ij} \sim N(0, \omega),$$

where cases (j) are nested within studies (i), and have among-case (within study) variance, ω^2;

study-level heterogeneity, β_{0i}, has variance τ^2, and varies around the overall linear relationship

with intercept, β_0, slope β_1, and predictor X_i (here the natural logarithm of grain size in study i).

The model was fit to empirical data with weakly regularizing priors:

698
$$\beta_0 \sim N(0.3, 1),$$

699
$$\beta_1 \sim N(0, 1),$$

700
$$[\tau, \omega] \sim N(0, 1), \qquad \text{(Model S1.1)}.$$

701   This model had good convergence (all Rhat < 1.01), and showed a reasonable fit to the empirical

702   data (Appendix S2: Fig. S1).

704
705   **Figure S1**: Posterior predictive check for model 1.1 fit to the empirical data (*z*-scores).
706

The parameter estimates from the fit of model S1.1 to empirical data (β₀: 0.13 [95% credible

interval: 0.01 – 0.24]; β₁: 0.04 [95% credible interval: 0.02 – 0.05]; τ: 0.34 [95% credible

interval: 0.25 – 0.43]; ω: 0.3 [95% credible interval: 0.24 – 0.35]) were used to inform the

following priors:

$$\beta_0 \sim N(0.13, 0.06),$$

$$\beta_1 \sim N(0.04, 0.02),$$

$$\tau \sim N(0, 0.25),$$

$$\omega \sim N(0, 0.25),$$

which were combined with model S1.1 to simulate (fake) data sets. To ensure as much realism as

possible in the simulated data, each simulated data set retained the characteristics of the

empirical data: 204 observations distributed across 101 studies (with the same balance, i.e.,

effect sizes per study, and the known standard errors $[s_{ij}]$ from the empirical data). I refit model

S1.1 to each simulated data set, and identified models fit to fake data with poor diagnostics (e.g.,

divergent transitions and Rhats > 1.05), likely due to the simulated data have extreme values for

the simulated data; I report the number of simulated data sets used to calculate simulation

diagnostics in figure captions.

Simulation-based calibration for model 1.1 showed that the rank statistics were

approximately uniformly distributed (Appendix S2: Fig. S2), and that the coverage of parameters

was reasonable (Appendix S2: Fig. S3); known parameters were approximately recovered

(Appendix S2: Fig. S4).

Figure S2: The posterior ranks of the prior draws were approximately normally distributed for the parameters of interest in model S1.1. Results are shown for n = 95 simulated data sets.

728
729
730

Figure S3: Model S1.1 had reasonable coverage for the parameters of interest. Results are shown
for n = 95 simulated data sets.

Figure S4: Model S1.1 was largely able to recover known parameter values. Results are shown
for n = 95 simulated data sets.

Model 1.2

The second model in case study one extended model S1.1 to include an additional parameter

(σ_{ij}) for residual variation as a function of grain size:

$$z_{ij} \sim N(\mu_{ij}, s_{ij} + \sigma_{ij}),$$

$$\mu_{ij} = \beta_0 + \beta_{0i} + \beta_1 X_i,$$

$$\beta_{0i} \sim N(0, \tau),$$

$$log(\sigma_{ij}) = \beta_0^\sigma + \beta_1^\sigma X_{ij},$$

where X_{ij} is grain size on a log-scale for the *j*th case in study *i*. I fit this model to the empirical

data with weakly regularizing priors:

$$\beta_0 \sim N(0.3, 1),$$

$$\beta_0^\sigma, \beta_1^\sigma, \beta_1 \sim N(0, 1),$$

$$\tau \sim N(0, 1), \qquad \text{(Model S1.2)}.$$

Model 1.2 had good convergence (all Rhats < 1.01), and showed a good fit to the empirical data

(Appendix S2: Fig. S5).

The parameter estimates from the fit of model S1.2 to empirical data (β_0: 0.13 [95%

credible interval: 0.01 – 0.24]; β_1: 0.04 [95% credible interval: 0.02 – 0.05]; τ: 0.33 [95%

credible interval: 0.24 – 0.42]; β_0^σ: -1.12 [95% credible interval: -1.4 – -0.86]; β_1^σ: -0.03 [95%

credible interval: -0.1 – 0.03]) were used to inform the following priors:

$$\beta_0 \sim N(0.13, 0.06),$$

$$\beta_1 \sim N(0.04, 0.02),$$

$$\tau \sim N(0, 0.2),$$

$$\beta_0^\sigma \sim N(-1.1, 0.2),$$

$$\beta_1^\sigma \sim N(-0.03, 0.03),$$

which were combined with model S1.2 to simulate (fake) data sets.

Figure S5: Posterior predictions from model 1.2 were largely consistent with the observed data.

Simulation-based calibration for model 1.2 showed that the rank statistics were approximately uniformly distributed (Appendix S2: Fig. S6), and that the coverage of parameters was reasonable (Appendix S2: Fig. S7); known parameters were approximately recovered, though with a fair amount of uncertainty for the intercept (β_0), the varying intercept for studies (τ), and the slope of relationship between unexplained variation and grain size (β_1^σ; Appendix S2: Fig. S8).

Figure S6: The posterior ranks of the prior draws were approximately normally distributed for the parameters of interest in model S1.2. Results are shown for n = 150 simulated data sets.

773
774
775
776

Figure S7: Model S1.2 had reasonable coverage for the parameters of interest. Results are shown
for n = 150 simulated data sets.

778
779
780

782 **Figure S8**: Model S1.2 was able to recover known parameter values, though with some
783 uncertainty (especially for β_0, τ, and β_1^{σ}). Results are shown for n = 150 simulated data sets.
784
785

Model 1.3

The third model replaced grain size as a predictor of unexplained variation with spatial

extent, which was estimated with indicator variables for each category of spatial extent [i.e., no

intercept, see main text]). I fit this model to the empirical data with the same weakly regularizing

priors as S1.2 (with the exception of β_0^{σ}, which was dropped from the model).

Model 1.3 had good convergence (all Rhats < 1.01), and showed a reasonable fit to the

empirical data (Appendix S2: Fig. S9).

Figure S9: Posterior predictions for model 1.3 showed reasonable fidelity to the empirical data.
795

The parameter estimates from the fit of model S1.3 to empirical data (β_0: 0.17 [95%

credible interval: 0.06 – 0.28]; β_1: 0.03 [95% credible interval: 0.02 – 0.04]; τ: 0.34 [95%

credible interval: 0.25 – 0.43]; β^{σ}_{0-10}: -0.79 [95% credible interval: -1.07 - -0.52]; β^{σ}_{10-100}: -1.58

[95% credible interval: -2.3 - -0.83]; $\beta^{\sigma}_{10^2-10^3}$: -1.03 [95% credible interval: -1.5 - -0.57];

$\beta^{\sigma}_{10^3-10^4}$: -2.09 [95% credible interval: -2.95- -1.39]; $\beta^{\sigma}_{10^4-10^5}$: -1.66 [95% credible interval:

-2.93 - -0.69]; $\beta^{\sigma}_{10^5-10^6}$: -1.53 [95% credible interval: -1.91 - -1.13]; $\beta^{\sigma}_{10^6-}$: -1.51 [95%

credible interval: -2.87- -0.38]) were used to inform the following priors:

$$\beta_0 \sim N(0.13, 0.05),$$

$$\beta_1 \sim N(0.04, 0.02),$$

$$\tau \sim N(0, 0.2),$$

$$\beta^{\sigma}_1 \sim N(-1,1),$$

which were combined with model S1.3 to simulate (fake) data sets.

Simulation-based calibration for model 1.3 showed that the rank statistics were

approximately uniformly distributed (Appendix S2: Fig. S10), and that the coverage of

parameters was reasonable (Appendix S2: Fig. S11); known parameters were approximately

recovered, though with a fair amount of uncertainty for the intercept (β_0), and the varying

intercept for studies (τ, Appendix S2: Fig. S12).

Figure S10: The posterior ranks of the prior draws were approximately normally distributed for
the parameters of interest in model S1.3. Results are shown for n = 176 simulated data sets.
816
817

818
819 **Figure S11**: Model S1.3 had reasonable coverage for the parameters of interest. Results are
820 shown for n = 176 simulated data sets.
821

822
823 **Figure S12**: Model S1.3 was able to recover known parameter values, though with some
824 uncertainty (most notably for β_0, and τ). Results are shown for n = 176 simulated data sets.
825

826 *Model 1.4*

827 The final model examined in the main text included study-level residual variation:

828
$$z_{ij} \sim N(\mu_{ij}, s_{ij} + \sigma_{ij}),$$

829
$$\mu_{ij} = \beta_0 + \beta_{0i} + \beta_1 X_i,$$

830
$$\beta_{0i} \sim N(0, \tau),$$

831
$$log(\sigma_{ij}) = \beta_0^\sigma + \beta_{0i}^\sigma,$$

832
$$\beta_{0i}^\sigma \sim N(0, \zeta),$$

833   where β_0^σ is the average residual variation (on a log-scale), and β_{0i}^σ is a normally distributed

834   study-level varying intercept with zero mean and ζ standard deviation. The model was fit to the

835   empirical data with weakly regularizing parameters:

$$\beta_0 \sim N(0.3, 1),$$

$$\beta_1, \beta_0^\sigma \sim N(0, 1),$$

$$[\tau, \zeta] \sim N(0, 1). \qquad \text{(Model S1.4)}$$

839   Model 1.4 had good convergence (all Rhats < 1.01), and showed a reasonable fit to the empirical

840   data, albeit with some very low predicted z-scores (Appendix S2: Fig. S13).

841
842 **Figure S13:** Posterior predictions from model 1.4 showed a good fit to the empirical data,
843 though it did predict some extremely low z-scores (not seen in the observed data).
844

845   The parameter estimates from the fit of model S1.4 to empirical data (β_0: 0.17 [95%

846   credible interval: 0.06 − 0.27]; β_1: 0.03 [95% credible interval: 0.02 − 0.05]; τ: 0.31 [95%

847   credible interval: 0.22 – 0.40]; β_0^σ: -1.65 [95% credible interval: -2.15 – -1.26]; ζ: 0.89 [95%

848   credible interval: 0.56 – 1.31]) were used to inform the following priors:

849
$$\beta_0 \sim N(0.17, 0.05),$$

850
$$\beta_1 \sim N(0.04, 0.02),$$

851
$$\tau \sim N(0, 0.2),$$

852
$$\beta_0^\sigma \sim N(-1.6, 0.25),$$

853
$$\zeta \sim N(0, 0.7),$$

which were combined with model S1.4 to simulate (fake) data sets.

Simulation-based calibration for model 1.4 showed that the rank statistics were

approximately uniformly distributed (Appendix S2: Fig. S14), and that the coverage of

parameters was reasonable (Appendix S2: Fig. S15). Finally, most of the known parameters were

approximately recovered, though with a fair amount of uncertainty for the intercept (β_0), and

particularly the varying intercept for study-level unexplained variation (ζ; Appendix S2: Fig.

S16). The relatively poor (and highly uncertain) estimate of study-level unexplained variation (ζ)

is likely due to the limited replication within studies (the median number of effect sizes per study

was one; 74 of 101 studies had a single effect size).

863
864 **Figure S14**: The posterior ranks of the prior draws were approximately normally distributed for
865 the parameters of interest in model S1.4. Results are shown for n = 108 simulated data sets.
866

867
Figure S15: Model S1.4 had reasonable coverage for the parameters of interest, though the
intercept (β_0) showed some discrepancies. Results are shown for n = 108 simulated data sets.
870

Figure S16: Model S1.3 was able to recover known parameter values, though with some uncertainty (most notably ζ). Results are shown for n = 108 simulated data sets.

References

Modrák, M., A. H. Moon, S. Kim, P. Bürkner, N. Huurre, K. Faltejsková, A. Gelman, and A. Vehtari. 2023. Simulation-based calibration checking for Bayesian computation: The choice of test quantities shapes sensitivity. Bayesian Analysis 1:1–28.

Peng, S., N. L. Kinlock, J. Gurevitch, and S. Peng. 2019. Correlation of native and exotic species richness: a global meta-analysis finds no invasion paradox across scales. Ecology 100:e02552.

Talts, S., M. Betancourt, D. Simpson, A. Vehtari, and A. Gelman. 2020, October 21. Validating Bayesian Inference Algorithms with Simulation-Based Calibration. arXiv.

Appendix S3

Case-study two: Simulation-based calibration and supplemental figures

I used Simulation-Based Calibration (Talts et al. 2020, Modrák et al. 2023) to check whether

models could recover known parameter values with reasonable coverage (i.e., the probability that

a constructed [e.g., credible] interval contains the true value). Briefly, for my purposes here this

involves: (1) fitting a model to the empirical data; (2) using the same model with priors informed

by the fit to the empirical data to simulate many new (fake) datasets with the same size, shape

and structure as the empirical data (i.e., the same number of observations, and the same number

of groups for each level in the hierarchical structure); (3) refitting the model to each simulated

data set; and, (4) calculating and plotting SBC diagnostics of model fits to simulated data to

check for reasonable coverage (Talts et al. 2020, Modrák et al. 2023).

Model 2.1

To start case-study two I refit the model fit by Chase et al. (2020) to effort-standardized for

species richness (S_{ij}) in fragment j from study i:

$$S_{ij} \sim lognormal(\mu_{ij}, \sigma)$$

$$\mu_{ij} = \beta_0 + \beta_{0i} + (\beta_1 + \beta_{1i})X_{ij},$$

$$[\beta_{0i}, \beta_{1i}]' \sim MVN(0, \boldsymbol{SRS}),$$

$$\boldsymbol{S} = \begin{bmatrix} \sigma_{0i} & 0 \\ 0 & \sigma_{1i} \end{bmatrix},$$

$$\boldsymbol{R} = \begin{bmatrix} 1 & \rho_{\sigma_{0i}\sigma_{1i}} \\ \rho_{\sigma_{0i}\sigma_{1i}} & 1 \end{bmatrix},$$

where X_{ij} is the fragment size on a (natural) log-scale, which was centered by subtracting the

overall mean from each observation before modelling; β_{0i} and β_{1i} are study-level departures from

the overall intercept and slope, respectively, drawn from a multivariate normal (*MVN*)

distribution with standard deviations, σ_{0i} and σ_{1i}, that allowed for correlations ($\boldsymbol{R}$ matrix)

between the varying intercepts and slopes. The model was fit to empirical data using weakly

regularizing priors:

910
$$\beta_0 \sim N(2.5, 1),$$

911
$$\beta_1 \sim N(0, 1),$$

912
$$[\sigma_{0i}, \sigma_{1i}, \sigma] \sim N(0, 1),$$

913
$$\boldsymbol{R} \sim LKJ(1). \qquad \text{(Model S2.1)}$$

914   This model had good convergence (all Rhat <= 1.01), and showed a reasonable fit to the

915   empirical data (Appendix S3: Fig. S1).

917   **Figure S1:** Posterior predictions from model 2.1 showed the model had a good ability to predict
918   the observed data.
919

The parameter estimates from the fit of model S2.1 to empirical data (β_0: 2.46 [95% credible

interval: 2.28 – 2.65]; β_1: 0.06 [95% credible interval: 0.04 – 0.07]; σ_{0i}: 1 [95% credible interval:

0.88 – 1.14]; σ_{1i}: 0.06 [95% credible interval: 0.04 – 0.08], and σ: 0.36 [95% credible interval:

0.34 – 0.37]) were used to inform the following priors:

$$\beta_0 \sim N(2.4, 0.1),$$

$$\beta_1 \sim N(0.06, 0.01),$$

$$\sigma_{0i} \sim N(1, 0.1),$$

$$\sigma_{1i} \sim N(0, 0.05),$$

$$\sigma \sim N(0.3, 0.05),$$

which were combined with model S2.1 to simulate (fake) data sets. To ensure as much realism as

possible in the simulated data, each simulated data set retained the characteristics of the

empirical data: 1509 observations distributed across 123 studies (with the same balance, i.e.,

fragment size distribution and fragments per study). I refit model S2.1 to each simulated data set,

and identified models fit to fake data with poor diagnostics (e.g., divergent transitions and Rhats

> 1.05), likely due to the simulated data have extreme values for the simulated data; I report the

number of simulated data sets used to calculate simulation diagnostics in figure captions.

Simulation-based calibration for model 2.1 showed that the rank statistics were

approximately uniformly distributed (Appendix S3: Fig. S2), and that the coverage of parameters

was reasonable (Appendix S3: Fig. S3); known parameters were approximately recovered

(Appendix S3: Fig. S4). Note, (here and in subsequent calibrations for model extensions) I do not

show parameters associated with the intercept (i.e., β_0, σ_{0i}) as these parameters are not readily

interpretable for this analysis. They correspond to the average richness (across all studies), and

study-level variation around the overall average. Richness (and other diversity measures) are not

readily comparable across this heterogeneous data compilation: richness in any given data set

will depend on the sample grain and size of the species pool (among other things).

945
946 **Figure S2:** The posterior ranks of the prior draws were approximately normally distributed for
947 the parameters of interest in model S2.1. Results are shown for n = 401 simulated data sets.

948
949 **Figure S3:** Model S2.1 had good coverage for the parameters of interest. Results are shown for n
950 = 401 simulated data sets.

951
952 **Figure S4:** Model S2.1 was able to recover known parameter values, though with considerable
953 uncertainty (with the exception of σ_1 and σ). Results are shown for n = 401 simulated data sets.
954

Model 2.2

The first heteroscedastic model is motivated similarly to the varying intercepts and slopes per

study, and allowed varying study-level residuals around an overall average residual standard

deviation. I estimated study-level residual variation independently of other varying study-level

parameters for the mean:

$$S_{ij} \sim lognormal(\mu_{ij}, \sigma_i)$$

$$\mu_{ij} = \beta_0 + \beta_{0i} + (\beta_1 + \beta_{1i})X_{ij},$$

$$[\beta_{0i}, \beta_{1i}]' \sim MVN(0, \boldsymbol{SRS}),$$

$$\boldsymbol{S} = \begin{bmatrix} \sigma_{0i} & 0 \\ 0 & \sigma_{1i} \end{bmatrix},$$

$$\boldsymbol{R} = \begin{bmatrix} 1 & \rho_{\sigma_{0i}\sigma_{1i}} \\ \rho_{\sigma_{0i}\sigma_{1i}} & 1 \end{bmatrix},$$

$$log(\sigma_i) = \beta_0^\sigma + \beta_{0i}^\sigma,$$

$$\beta_{0i}^\sigma \sim N(0, \sigma_{0i}^\sigma),$$

where β_0^σ is the overall average of residual variation (on a log-scale), and β_{0i}^σ are the varying

study-level departures (for the residual variation) drawn from a normal distribution with zero

mean and standard deviation σ_{0i}^σ. I fit the model with weakly regularizing priors:

$$\beta_0 \sim N(2.5, 1),$$

$$\beta_1, \beta_0^\sigma \sim N(0, 1),$$

$$[\sigma_{0i}, \sigma_{1i}, \sigma_{0i}^\sigma] \sim N(0, 1),$$

$$\mathbf{R} \sim LKJ(1). \qquad \text{(Model S2.2)}$$

This model had good convergence (all Rhat <= 1.01), and showed a reasonable fit to the

empirical data (Appendix S3: Fig. S5).

978
979   **Figure S5:** Posterior predictions from model 2.2 showed the model had a good ability to predict
980   the observed data.
981

982   The parameter estimates from the fit of model S2.2 to empirical data (β_0: 2.48 [95% credible

983   interval: 2.31 – 2.66]; β_1: 0.05 [95% credible interval: 0.04 – 0.07]; σ_{0i}: 1 [95% credible interval:

984   0.88 – 1.15]; σ_{1i}: 0.06 [95% credible interval: 0.04 – 0.07], β_0^σ: -1.23 [95% credible interval: -

985   1.32 – -1.13], and σ_{0i}^σ: 0.44 [95% credible interval: 0.36 – 0.53]) were used to inform the

986   following priors:

$$\beta_0 \sim N(2.5, 0.1),$$

$$\beta_1 \sim N(0.05, 0.01),$$

$$\sigma_{0i} \sim N(1, 0.1),$$

$$\sigma_{1i} \sim N(0.05, 0.05),$$

$$\beta_0^\sigma \sim N(-1.2, 0.1),$$

$$\sigma_{0i}^\sigma \sim N(0.4, 0.1).$$

993   Simulation-based calibration for model 2.2 showed that the rank statistics were

994   approximately uniformly distributed (Appendix S3: Fig. S6), and that the coverage of parameters

995   was reasonable (Appendix S3: Fig. S7); known parameters were approximately recovered

996   (Appendix S3: Fig. S8).

997
998 Figure S6: The posterior ranks of the prior draws were approximately normally distributed for
999 the parameters of interest in model S2.2. Results are shown for n = 185 simulated data sets.
1000

1001

Figure S7: Model S2.2 had good coverage for the parameters of interest. Results are shown for n
= 185 simulated data sets.

1002
1003
1004

1005
1006 **Figure S8**: Model S2.2 was able to recover known parameter values, though with considerable
1007 uncertainty (with the exception of σ_1). Results are shown for n = 185 simulated data sets.
1008

1009 *Model 2.3*

1010 Next, I extend this model to include (log) fragment size as a predictor of residual variation, and

1011 again allow it to varying between studies independently of varying study-level parameters for the

1012 mean:

1013
$$S_{ij} \sim lognormal(\mu_{ij}, \sigma_i)$$

1014
$$\mu_{ij} = \beta_0 + \beta_{0i} + (\beta_1 + \beta_{1i})X_{ij},$$

1015
$$[\beta_{0i}, \beta_{1i}]' \sim MVN(0, \mathbf{SRS}),$$

1016
$$\mathbf{S} = \begin{bmatrix} \sigma_{0i} & 0 \\ 0 & \sigma_{1i} \end{bmatrix},$$

1017
$$R = \begin{bmatrix} 1 & \rho_{\sigma_{0i}\sigma_{1i}} \\ \rho_{\sigma_{0i}\sigma_{1i}} & 1 \end{bmatrix},$$

1018
$$log(\sigma_i) = \beta_0^\sigma + \beta_{0i}^\sigma + (\beta_1^\sigma + \beta_{1i}^\sigma)X_i,$$

1019
$$[\beta_{0i}^\sigma, \beta_{1i}^\sigma]' \sim MVN(0, S^\sigma R^\sigma S^\sigma),$$

1020
$$S^\sigma = \begin{bmatrix} \sigma_{0i}^\sigma & 0 \\ 0 & \sigma_{1i}^\sigma \end{bmatrix},$$

1021
$$R^\sigma = \begin{bmatrix} 1 & \rho_{\sigma_{0i}^\sigma \sigma_{1i}^\sigma} \\ \rho_{\sigma_{0i}^\sigma \sigma_{1i}^\sigma} & 1 \end{bmatrix},$$

1022

1023   where β_0^σ is the overall average residual variation, and β_1^σ is the overall average slope of residual

1024   variation with fragment size; β_{0i}^σ and β_{1i}^σ are the varying study-level departures from the intercept

1025   and slope, respectively, and were drawn from a multivariate normal distribution with zero mean

1026   and standard deviation σ_{0i}^σ and σ_{1i}^σ, with correlations estimated in matrix R^σ. I fit the model with

1027   weakly regularizing priors:

1028
$$\beta_0 \sim N(2.5, 1),$$

1029
$$\beta_1, \beta_0^\sigma, \beta_1^\sigma \sim N(0, 1),$$

1030
$$[\sigma_{0i}, \sigma_{1i}] \sim N(0, 1),$$

1031
$$R \sim LKJ(1),$$

1032
$$[\sigma_{0i}^\sigma, \sigma_{1i}^\sigma] \sim N(0,1),$$

1033
$$R^\sigma \sim LKJ(1). \hspace{2cm} \text{(Model S2.3)}$$

1034   This model had good convergence (all Rhat <= 1.02), and showed a reasonable fit to the

1035   empirical data (Appendix S3: Fig. S9).

1036
1037 **Figure S9:** Posterior predictions from model 2.3 showed good fidelity to the observed data.
1038

The parameter estimates from the fit of model S2.3 to empirical data (β_0: 2.48 [95%

credible interval: 2.3 – 2.66]; β_1: 0.05 [95% credible interval: 0.03 – 0.06]; σ_{0i}: 1.01 [95%

credible interval: 0.89 – 1.14]; σ_{1i}: 0.06 [95% credible interval: 0.04 – 0.07]; β_0^{σ}: -1.25 [95%

credible interval: -1.35 – -1.15]; β_1^{σ}: -0.06 [95% credible interval: -0.09 – -0.03]; σ_{0i}^{σ}: 0.45 [95%

credible interval: 0.37 – 0.55], and σ_{1i}^{σ}: 0.06 [95% credible interval: 0.01 – 0.1]) were used to

inform the following priors:

$$\beta_0 \sim N(2.5, 0.1),$$

$$\beta_1 \sim N(0.05, 0.01),$$

$$\sigma_{0i} \sim N(1, 0.1),$$

$$\sigma_{1i} \sim N(0.05, 0.05),$$

$$\beta_0^{\sigma} \sim N(-1.2, 0.1),$$

$$\sigma_{0i}^{\sigma} \sim N(0.4, 0.1).$$

Simulation-based calibration for model 2.3 showed that the rank statistics were

approximately uniformly distributed (Appendix S3: Fig. S10), and that the coverage of

parameters was reasonable (Appendix S3: Fig. S11); known parameters were approximately

recovered (Appendix S3: Fig. S12).

1055

1056
1057   **Figure S10:** The posterior ranks of the prior draws were approximately normally distributed for
1058   the parameters of interest in model S2.3. Results are shown for n = 77 simulated data sets.

1059
Figure S11: Model S2.3 had good coverage for all parameters of interest. Results are shown for
1061   n = 77 simulated data sets.

1062

Figure S12: Model S2.3 was able to recover most parameters with reasonable accuracy, albeit with considerable uncertainty. Results are shown for n = 77 simulated data sets.

Model 2.4

The next model is the same as model 2.2 (i.e., varying study-level residuals), but allows the study-level residuals to covary with the other varying parameters for the mean:

$$S_{ij} \sim lognormal(\mu_{ij}, \sigma_i)$$

$$\mu_{ij} = \beta_0 + \beta_{0i} + (\beta_1 + \beta_{1i})X_{ij},$$

$$log(\sigma_i) = \beta_0^\sigma + \beta_{0i}^\sigma,$$

$$[\beta_{0i}, \beta_{1i}, \beta_{0i}^\sigma]' \sim MVN(0, \boldsymbol{SRS}),$$

$$\boldsymbol{S} = \begin{bmatrix} \sigma_{0i} & 0 & 0 \\ 0 & \sigma_{1i} & 0 \\ 0 & 0 & \sigma_{0i}^\sigma \end{bmatrix},$$

1074
$$R = \begin{bmatrix} 1 & \rho_{\sigma_{0i}\sigma_{1i}} & \rho_{\sigma_{0i}\sigma_{0i}^{\sigma}} \\ \rho_{\sigma_{0i}\sigma_{1i}} & 1 & \rho_{\sigma_{1i}\sigma_{0i}^{\sigma}} \\ \rho_{\sigma_{0i}\sigma_{0i}^{\sigma}} & \rho_{\sigma_{1i}\sigma_{0i}^{\sigma}} & 1 \end{bmatrix}.$$

The model was fit to empirical data with weakly regularizing priors:

$$\beta_0 \sim N(2.5, 1),$$

$$\beta_1, \beta_0^{\sigma} \sim N(0, 1),$$

$$[\sigma_{0i}, \sigma_{1i}, \sigma_{0i}^{\sigma}] \sim N(0, 1),$$

$$\mathbf{R} \sim LKJ(1). \qquad \text{(Model S2.4)}$$

This model had good convergence (all Rhat <= 1.02), and showed a reasonable fit to the

empirical data (Appendix S3: Fig. Sx).

1082

1083 **Figure S13:** Posterior predictions showed that model 2.4 was able to make predict the observed
1084 data well.

1085

The parameter estimates from the fit of model S2.4 to empirical data (β_0: 2.46 [95%

credible interval: 2.29 − 2.66]; β_1: 0.05 [95% credible interval: 0.04 − 0.07]; σ_{0i}: 1 [95% credible

interval: 0.89 – 1.13]; σ_{1i}: 0.06 [95% credible interval: 0.04 – 0.07], β_0^σ: -1.24 [95% credible

interval: -1.35 – -1.15], and σ_{0i}^σ: 0.44 [95% credible interval: 0.37 – 0.53]) were used to inform

the following priors:

$$\beta_0 \sim N(2.5, 0.1),$$

$$\beta_1 \sim N(0.05, 0.01),$$

$$\sigma_{0i} \sim N(1, 0.1),$$

$$\sigma_{1i} \sim N(0.05, 0.05),$$

$$\beta_0^\sigma \sim N(-1.2, 0.1),$$

$$\sigma_{0i}^\sigma \sim N(0.4, 0.1).$$

Simulation-based calibration for model 2.4 showed that the rank statistics were

approximately uniformly distributed (Appendix S3: Fig. S14), and that the coverage of

parameters was reasonable (Appendix S3: Fig. S15); known parameters were approximately

recovered (Appendix S3: Fig. S16).

1101
1102 **Figure S14:** The posterior ranks of the prior draws were approximately normally distributed for
1103 the parameters of interest in model S2.4. Results are shown for n = 181 simulated data sets.

Figure S15: Model S2.4 had good coverage for all parameters of interest. Results are shown for n = 181 simulated data sets.

1107
1108 **Figure S16:** Model S2.4 was able to recover known parameter values with reasonable accuracy.
1109 Results are shown for n = 181 simulated data sets.
1110

1111 *Model 2.5*

1112 The final model extended model 2.3 to allow for correlations between all of the varying study-

1113 level parameters (i.e., for the location and the scale):

1114
$$S_{ij} \sim lognormal(\mu_{ij}, \sigma_i)$$

1115
$$\mu_{ij} = \beta_0 + \beta_{0i} + (\beta_1 + \beta_{1i})X_{ij},$$

1116
$$log(\sigma_i) = \beta_0^\sigma + \beta_{0i}^\sigma + (\beta_1^\sigma + \beta_{1i}^\sigma)X_i,$$

1117
$$[\beta_{0i}, \beta_{1i}, \beta_{0i}^\sigma, \beta_{1i}^\sigma]' \sim MVN(0, \boldsymbol{SRS}),$$

1118
$$\boldsymbol{S} = \begin{bmatrix} \sigma_{0i} & 0 & 0 & 0 \\ 0 & \sigma_{1i} & 0 & 0 \\ 0 & 0 & \sigma_{0i}^\sigma & 0 \\ 0 & 0 & 0 & \sigma_{1i}^\sigma \end{bmatrix},$$

1119

$$\boldsymbol{R} = \begin{bmatrix} 1 & \rho_{\sigma_{0i}\sigma_{1i}} & \rho_{\sigma_{0i}\sigma_{0i}^\sigma} & \rho_{\sigma_{0i}\sigma_{1i}^\sigma} \\ \rho_{\sigma_{0i}\sigma_{1i}} & 1 & \rho_{\sigma_{1i}\sigma_{0i}^\sigma} & \rho_{\sigma_{1i}\sigma_{1i}^\sigma} \\ \rho_{\sigma_{0i}\sigma_{0i}^\sigma} & \rho_{\sigma_{1i}\sigma_{0i}^\sigma} & 1 & \rho_{\sigma_{0i}^\sigma\sigma_{1i}^\sigma} \\ \rho_{\sigma_{0i}\sigma_{1i}^\sigma} & \rho_{\sigma_{1i}\sigma_{1i}^\sigma} & \rho_{\sigma_{0i}^\sigma\sigma_{1i}^\sigma} & 1 \end{bmatrix}.$$

1120   I fit the model with weakly regularizing priors:

1121
$$\beta_0 \sim N(2.5, 1),$$

1122
$$\beta_1, \beta_0^\sigma, \beta_1^\sigma \sim N(0, 1),$$

1123
$$[\sigma_{0i}, \sigma_{1i}, \sigma_{0i}^\sigma, \sigma_{1i}^\sigma] \sim N(0, 1),$$

1124
$$\boldsymbol{R} \sim LKJ(1). \qquad\qquad \text{(Model S2.5)}$$

1125   This model had good convergence (all Rhat < 1.01), and showed a reasonable fit to the empirical

1126   data (Appendix S3: Fig. S17).

1127
1128   **Figure S17:** Model 2.5 made predictions that resembled the observed data closely.
1129

The parameter estimates from the fit of model S2.5 to empirical data (β_0: 2.47 [95%

credible interval: 2.3 – 2.66]; β_1: 0.05 [95% credible interval: 0.03 – 0.06]; σ_{0i}: 1.01 [95%

credible interval: 0.89 – 1.15]; σ_{1i}: 0.06 [95% credible interval: 0.04 – 0.07]; β_0^σ: -1.28 [95%

credible interval: -1.39 – -1.18]; β_1^σ: -0.06 [95% credible interval: -0.09 – -0.04]; σ_{0i}^σ: 0.46 [95%

credible interval: 0.39 – 0.56], and σ_{1i}^σ: 0.07 [95% credible interval: 0.03 – 0.11]) were used to

inform the following priors:

1136 $$\beta_0 \sim N(2.5, 0.1),$$

1137 $$\beta_1 \sim N(0.05, 0.01),$$

1138 $$\sigma_{0i} \sim N(1, 0.1),$$

1139 $$\sigma_{1i} \sim N(0.05, 0.05),$$

1140 $$\beta_0^\sigma \sim N(-1.2, 0.1),$$

1141 $$\sigma_{0i}^\sigma \sim N(0.4, 0.1).$$

1142   Simulation-based calibration for model 2.5 showed that the rank statistics were

1143   approximately uniformly distributed (Appendix S3: Fig. S18), and that the coverage of

1144   parameters was reasonable (Appendix S3: Fig. S19); known parameters were approximately

1145   recovered (Appendix S3: Fig. S20).

Figure S18: The posterior ranks of the prior draws were approximately normally distributed for the parameters of interest in model S2.5. Results are shown for n = 98 simulated data sets.

1150
Figure S19: Model S2.5 had good coverage for all parameters of interest. Results are shown for
n = 98 simulated data sets.
1153

1154
1155 **Figure S20:** Model S2.5 was able to recover known parameter values with reasonable accuracy,
1156 with the exception of the correlation ($\rho_{\sigma_1 \sigma_1^{\sigma}}$), which was estimated with high uncertainty. Results
1157 are shown for n = 98 simulated data sets.
1158
1159
1160
1161

1162

1163
1164 **Figure S21**: All models fit in case study two supported the ecosystem decay hypothesis.

1165
1166 **Figure S21:** Posterior distributions (density plots for 1000 posterior draws) showing correlations
1167 among the varying study-level parameters.
1168

Figure S22: The decline of residual variation (scale) with increasing patch size was greatest for studies with the strongest ecosystem decay. Each point shows the estimate of a study-level slope, with slopes for the location on the x-axis, and the scale on the y-axis.

References

Chase, J. M., S. A. Blowes, T. M. Knight, K. Gerstner, and F. May. 2020. Ecosystem decay

    exacerbates biodiversity loss with habitat loss. Nature 584:238–243.

Modrák, M., A. H. Moon, S. Kim, P. Bürkner, N. Huurre, K. Faltejsková, A. Gelman, and A.

    Vehtari. 2023. Simulation-based calibration checking for Bayesian computation: The choice

    of test quantities shapes sensitivity. Bayesian Analysis 1:1–28.

Talts, S., M. Betancourt, D. Simpson, A. Vehtari, and A. Gelman. 2020, October 21. Validating

    Bayesian Inference Algorithms with Simulation-Based Calibration. arXiv.