1 **SpaceExpress enables differential spatial transcriptomics with natural coordinate**
2 **systems**
3

4 Yeojin Kim[1], Abhishek Ojha[1], Alex Schrader[2], Juyeon Lee[2,3], Zijun Wu[1], Ian M Traniello[4],
5 Saumya Jain[5], Gene E Robinson[6], Hee Sun Han[2,6*], Sihai D Zhao[6,7*], Saurabh Sinha[1,8*].
6
1 The Wallace H. Coulter Department of Biomedical Engineering, Georgia Institute of
Technology, Atlanta, GA, 30332, USA
2 Department of Chemistry, University of Illinois Urbana-Champaign, Urbana, IL, 61801, USA
3 School of Biological and Environmental Studies, Millikin University, Decatur, IL 62522, USA
4 Lewis-Sigler Institute for Integrative Genomics, Princeton University, Princeton, NJ, 08544,
USA
5 School of Biological Sciences, Georgia Institute of Technology, Atlanta, GA, 30332, USA
6 Carl R. Woese Institute for Genomic Biology, University of Illinois Urbana-Champaign,
Urbana, IL, 61801, USA
7 Department of Statistics, University of Illinois Urbana-Champaign, Urbana, IL, 61820, USA
8 H. Milton Stewart School of Industrial & Systems Engineering, Georgia Institute of
Technology, Atlanta, GA, 30318, USA
19
20 * Corresponding Authors
21 Corresponding Authors-
22 1. Saurabh Sinha, Email- saurabh.sinha@bme.gatech.edu
23 2. Sihai Dave Zhao, Email- sdzhao@illinois.edu
24 3. Hee-Sun Han, Email- hshan@illinois.edu
25

ABSTRACT

Spatial transcriptomics (ST) technologies have enabled new explorations into the spatial organization of tissues and their functional implications. However, one of the most fundamental analyses – comparative analysis of spatial gene expression across phenotypes – remains a formidable challenge. We introduce SpaceExpress, a novel statistical tool for detecting phenotype-associated changes in spatial expression patterns. SpaceExpress employs a neural network to embed multiple ST samples in a common latent space, enabling robust cross-sample comparisons despite structural and technical variations. It then uses spline regression to test differential spatial expression of genes between conditions, identifying specific regions of the tissue where expression patterns diverge and quantifying the magnitude of those differences, with rigorous false discovery control and handling of multiple replicates per condition. It includes visualization tools to help interpret spatial pattern differences. We demonstrate the tool's effectiveness on synthetic and real ST datasets, revealing mechanistic insights into behavior- and development-related neurogenomic changes in honey bees and mice. Our work extends the highly influential paradigm of differential gene expression analysis to spatial omics.

INTRODUCTION

The rapid development of highly multiplexed spatially resolved omics technologies, including spatial transcriptomics (ST)[1,2], has opened new avenues of inquiry into the interplay of intercellular communication and intracellular gene regulatory mechanisms, as well as the spatio-temporal organization of tissues and its relationship to function. As ST technologies mature and costs decrease, research groups will collect ST measurements from many samples to understand how the fine global structure of spatial gene expression might be associated with phenotypes such as disease states and quantitative traits.

The field therefore needs tools for comparative ST analyses. These tools must be able to quantitatively describe associations between spatial gene expression and phenotypes in an interpretable way, as well as test for their statistical significance. This will be an important step in advancing from the largely descriptive nature of current ST analytics towards a more mechanistic understanding of how biological systems function. For non-spatial transcriptomics data, rigorous detection of differential[3] or trait-associated[4] gene expression has been central to mechanistic studies. In contrast, ST data call for a new comparative analysis, which we call "differential *spatial* expression" (DSE) analysis: to detect and describe alterations in the fine spatial distribution of gene expression across phenotypes. However, to the best of our knowledge, no such tools currently exist for spatial gene expression, posing a significant barrier to achieving the full potential of ST technologies. Most existing methods simply test for differential expression between spatial domains across ST samples[5], which can only capture coarse forms of differential spatial patterning.

DSE analysis is a challenging problem for both technical and conceptual reasons. Technically, ST samples can vary for reasons that are not of direct interest, e.g., natural variation in tissue structure or differences in sectioning. Conceptually, it is difficult to even define a quantitative construct that can flexibly describe the innumerable ways in which the spatial structure of gene

expression may associate with other traits. Available methods for describing spatial gene
expression patterns in a tissue[6] do not furnish a way to compare those patterns across tissues,
especially in the face of structural variations noted above. Simply aligning multiple ST samples[7,8]
or mapping cells across samples[9,10] does not provide a useful way to quantitatively describe
differential patterns or rigorously test for associations with phenotypic traits, because the
Cartesian coordinate system is not appropriate for tissues with complex geometric and topological
structures. Furthermore, relying on automated inter-tissue alignments[11]] makes the comparisons
vulnerable to the effects of misalignment.
78
We propose a novel approach to comparative analysis for spatial biology and use it to develop a
statistical tool for inferring cross-sample or phenotype-associated changes in spatial expression
patterns. Our new tool, called SpaceExpress, learns a new coordinate system that can most
naturally describe locations within the tissue reflecting its intrinsic biological structure. This natural
coordinate system is *shared* across analyzed samples, in the sense that biologically analogous
regions of different samples are assigned similar coordinates. We demonstrate the accuracy of
this common coordinate system through extensive evaluations on synthetic data and four different
ST datasets obtained by three different technologies of varying resolution. Included visualization
tools aid in interpreting the coordinate system. SpaceExpress directly compares spatial gene
expression patterns between ST samples from different conditions along each axis of this
coordinate system. It employs spline regression modeling to capture how expression along these
natural axes differs across phenotypes and to formulate a rigorous hypothesis testing framework
for differential spatial expression (DSE). The DSE concept introduced here is distinct from and
complementary to the widely used differential expression[12] (DE) strategy, which tests for changes
in average gene expression levels between groups (**Figure 1a**). We apply SpaceExpress to
compare three distinct spatial transcriptomic brain datasets (**Figure 1b**). First, we compare whole
brain sections and specific brain substructures between a forager and soldier honey bee, using
ST data that we generated, and identify statistically significant DSE genes that yield mechanistic
hypotheses about behavior-related neurogenomic changes. Second, we compare visual cortex
samples from normally reared and dark-reared mice, revealing spatial transcriptional patterns
related to sensory experience. Third, we analyze two groups of ST samples from the mouse
hypothalamic preoptic area, highlighting spatial expression differences associated with
aggressive behavior. In summary, we present a novel approach to multi-sample ST data analysis
capable of identifying spatial axes capturing the fine global structure shared by those samples,
and rigorously testing the differential spatial expression of genes.
104
105

RESULTS

Overview of SpaceExpress

SpaceExpress is the first approach that can rigorously test whether and describe how a gene's
spatial expression pattern differs between samples. More generally, it can test if spatial
differences in gene expression are associated with a trait across multiple spatial samples.

Furthermore, it can characterize these associations in a way that aligns with the tissue's inherent
biological organization. It is challenging to compare expression patterns along conventional x-, y-
and z-axes, since the correspondence between standard Cartesian coordinates and biologically
relevant spatial features – such as developmental axes or tissue layers – varies across ST
samples. SpaceExpress overcomes this fundamental limitation by learning a more natural
coordinate system for the tissue that enables more meaningful comparisons of spatial gene
expression patterns.

SpaceExpress first constructs a coordinate-free representation of the spatial locations of each
cell in each sample using nearest neighbor graphs. It then learns a d-dimensional vector
representation of each cell such that a pair of cells close (resp. far) in a sample's graph have
similar (resp. different) representations (**Figure 1c,d**, Methods). (The dimensionality d is user-
configurable and typically set to a small value.) This step involves training a neural network to
obtain a cell's d-dimensional representation as a function of its gene expression profile. This is
motivated by the insight that gene expression encodes spatial location, arising from the literature
on "positional information"[13]. The learnt dimensions constitute a natural coordinate system (set of
axes) that captures the fine global structure of the tissue as encoded in its transcriptome. This
reflects a key insight behind our approach: since gene expression, when properly normalized, is
comparable across samples, the learnt axes, produced by the same neural network for every
tissue sample, are automatically comparable as well. These axes are invariant to technical
variation such as rotation or scaling, adapt to complex tissue geometries, and can be easily
interpreted by visualizing the learnt coordinates of each cell in a tissue sample.

SpaceExpress uses its axes to characterize whether and how a gene is differentially spatially
expressed with respect to discrete or quantitative traits. For this, it projects all cells onto each
dimension of the shared coordinate system and models a gene's expression pattern as a
mathematical function of the dimension. A gene is defined as differentially spatially expressed
(DSE) with respect to a trait if this function varies across samples with different trait values (**Figure
1c**, Methods). Such an association is assessed in a statistical hypothesis testing framework.
Differences in mean expression (DE[12]) between samples are already accounted for in the model
design, ensuring that they do not confound the identification of DSE. It is possible for a gene to
be DSE for some dimensions but not others. This is a major advantage of SpaceExpress: it can
identify which intrinsic tissue structures are involved in a gene's DSE.

SpaceExpress learns natural tissue coordinate system across multiple samples

The d-dimensional vector representations learnt by SpaceExpress describe cellular locations
across multiple spatial tissue samples in a common coordinate system, assigning analogous cells
in different samples to similar coordinates, e.g., functional domains, specific positions on a
biological axis, etc., are assigned similar coordinates. Our first tests of this crucial capability
involved synthetic data sets. We synthesized an "early Drosophila embryo slice" with cells laid out
uniformly in an elliptical area, with a subset of genes varying in expression along the "anterior-
posterior" ("AP") axis and/or "dorso-ventral" ("DV") axis (**Figure 2a**, **Supplementary Table S1**);
these are the natural axes of this simple model tissue. We generated two such synthetic embryos
with identical cellular layouts, and also introduced inter-embryo differences in spatial patterns of
a subset (20%) of genes (Figure 2a, Methods, **Supplementary Figure S1**), to mimic inter-sample

biological differences expected in a comparative analysis scenario. The level of DSE of these
genes was a variable parameter (μ).

We trained SpaceExpress on the two embryos jointly to obtain a 4-dimensional coordinate system
and evaluated it using two complementary metrics: "inter-sample agreement" and "natural
coordinate fidelity". Inter-sample agreement assesses whether corresponding cells from different
samples (as per the "ground truth") are proximal in the learned coordinate system, i.e., it quantifies
the "shared" nature of the coordinate system (see Methods). Natural coordinate fidelity, by
contrast, measures how well spatial relationships within each sample are preserved in the learned
coordinate system, i.e., it assesses if the coordinate system truly reflects spatial organization (see
Methods). To benchmark performance, we compared SpaceExpress with two prominent ST tools
– STAGATE[14] and GraphST[15]. While neither method is designed to learn a natural coordinate
system, these were relevant comparators for this evaluation as they support multi-sample learning
of cellular representations for domain identification, by combining transcriptomic profiles of cells
with their spatial location.

Our benchmarking shows that as DSE strength (μ) in the synthetic data is increased, making the
task harder, SpaceExpress consistently maintains higher inter-sample agreement scores than
both comparators, with STAGATE showing a marked drop at higher μ settings (**Figure 2b**, left).
SpaceExpress also achieves the highest natural coordinate fidelity across all values of μ, as
indicated by scores near 1 (**Figure 2b,** right). In contrast, STAGATE and GraphST show lower
and more variable fidelity depending on μ, indicating that only SpaceExpress robustly captures
the true underlying spatial axis even under divergent expression patterns.

Visualization of the joint embedding using PCA further illustrates these findings (**Figure 2c**). The
left panels for each method depict the overlap of two synthetic samples in the learned coordinate
space, for the data synthesized with µ = 0.6. SpaceExpress achieves near-complete overlap
between samples, reflecting strong inter-sample agreement, while STAGATE and GraphST
embed the two samples to overlapping but somewhat distinct regions in the shared space. The
right panels show cells colored by their original anterior-posterior (AP) coordinate, allowing visual
assessment of spatial fidelity. Only SpaceExpress recovered a continuous AP gradient,
supporting that its learned coordinate system preserves biologically meaningful structure within
each sample. (Similar visualizations of the DV axis are shown in **Supplementary Figure S2**.) We
further tested robustness to spatial distortion, including rotation (30°), scaling (1.5x), and partial
truncation (removal of 20% of one side of the tissue), as well as combinations thereof (**Figure
2d**). For these experiments, we also introduced variation in spatial expression patterns using μ =
0.5. Across all seven perturbation settings, SpaceExpress retained the highest inter-sample
agreement and robust natural coordinate fidelity. In contrast, both STAGATE and GraphST were
more sensitive to distortion, especially under combinations involving rotation and partial
truncation.

To validate performance on real data, we applied joint training to three diverse ST datasets
spanning different biological contexts and experimental platforms: (i) MERFISH profiles of visual
cortex from dark-reared (DR) and normal-reared (NR) mice[16] (**Figures 2e,f**), representing distinct
experimental conditions; (ii) human dorso-lateral prefrontal cortex (DLPFC) samples from two

individuals profiled using Visium[17] (**Figures 2g,h**), representing biological replicates under the
same condition; and (iii) Stereo-seq data from two adjacent slices of a late-stage Drosophila
embryo[18] (**Figures 2i,j**), representing technical replicates from the same specimen. In each case,
SpaceExpress achieved the highest inter-sample agreement and natural coordinate fidelity
across methods.

These results demonstrate that SpaceExpress can reliably learn a shared natural coordinate
system for multiple ST samples, which is key to our goal of identifying DSE genes. The method
is robust to both signal heterogeneity and spatial distortions, and broadly applicable to
comparative analysis across biological conditions, replicates, or individuals. The learned
coordinate system captures salient aspects of tissue organization, whether it is in the form of axes
(synthetic embryos, visual cortex) or discrete regions (DLPFC, late embryo). *Note:* see
Supplementary Note 1 for additional evaluations of SpaceExpress-learnt coordinate systems, in
the context of single-sample analysis, and **Supplementary Note 2** for further assessment of the
model's generalizability, showing that SpaceExpress trained on one sample can reconstruct
spatial organization in unseen tissues based solely on gene expression.

SpaceExpress identifies and characterizes differentially spatially expressed genes

The second step of SpaceExpress involves a rigorous statistical procedure for detecting
differential spatial expression (DSE), achieved via nonparametric regression modeling and
significance testing with false discovery rate (FDR) control (**Figure 3a-d**, **Supplementary Figure
S3**, Methods). Once cells of all tissue samples have been embedded in a common d-dimensional
space, each gene is tested for DSE along each of the dimensions. This involves fitting a spline
regression model of a gene's expression in a cell and using likelihood ratio statistics to assess if
the contribution of the cell's coordinate value to the model is sample-specific (or in the case of
more than two samples, group-specific). FDR control is achieved using an empirical null fitting
procedure[19]. The first step of learning a shared coordinate system may be prone to being
compromised by DSE genes, since such genes, by definition, violate the premise of gene
expression encoding space in a "universal" manner. To address this, once DSE genes have been
identified, the neural network for mapping cells to a coordinate space is retrained without using
those genes, and the statistical testing procedure is repeated a second time. We justified the
individual steps of our testing procedure through an ablation analysis (**Figure 3e**).

We performed systematic tests of the sensitivity and precision of our approach using the synthetic
data sets comprising pairs of "early embryos", where 20 of 100 genes are made DSE in a
controlled manner, as presented in the previous section (Figure 2a, Methods). Figures 3a-d
illustrate how SpaceExpress identifies such a DSE gene. This exemplar gene exhibits a moderate
level of DSE (Figure 3b, top), manifested along the AP as well as DV axis (**Supplementary Figure
S4a,b**), and SpaceExpress-learnt dimensions capture these axes (Figures 3a,b, **Supplementary
Figure S5**). Figure 3c shows the gene's expression in individual cells plotted against the
dimension 1 coordinates, and the fitted spline regression model reveals the distinct expression
trends in the two embryos. This divergence is quantified in a high log likelihood ratio statistic,
which is then assigned an FDR of approximately 0. We evaluated the performance of
SpaceExpress across all 100 genes and across varying levels of DSE strength (μ), observing true
positive rates ~1 with false positive rates below 0.02 (**Figure 3f, Supplementary Data S1**) at

most values of μ. For instance, at $\mu = 0.6$, SpaceExpress shows a clear separation in FDR
between the two classes across all dimensions (**Figure 3g**). Notably, sensitivity remains high even
for subtle spatial changes ($\mu < 0.3$), enabling detection of fine-grained DSE that may be visually
indiscernible (Supplementary Figure S4c,d). In many cases the DSE was detected for the same
gene in multiple dimensions (**Supplementary Data S2**). Additionally, SpaceExpress also
identifies DSE genes that are missed by conventional differential expression (DE) analyses
(**Figure 3h**), and has a superior accuracy than the alignment-based River method[11]
(Supplementary Figure 31a).

To test DSE detection on real data, we applied SpaceExpress to four pairs of tissue samples of
mouse visual cortex under normal (NR) and dark-rearing (DR) conditions (**Figure 3i**, Methods).
This identified DSE genes in these four NR–DR comparisons, and 21 genes were significantly
DSE in three or more of the analyses. The significance threshold was set at a level not crossed
by any gene when tested for DSE between intra-condition replicates (NR-NR and DR-DR
comparisons), thus ensuring strong control over false positive. As a comparison point, we
implemented an alignment-based approach to DSE detection that does not utilize a shared
coordinate system, and repeated the above analysis to find only one gene to be consistently DSE
(in ≥ 3 of pairwise comparisons) at equivalent levels of false positive errors. (Also see
Supplementary Figure 31b for a comparison to the River method[11] for ranking DSE genes.) A
similar analysis of mouse hypothalamic preoptic area samples from naïve and pup-exposed mice
yielded 18 robust DSE genes versus seven with the alignment-based comparator (**Figure 3j,**
Methods). Thus, SpaceExpress achieves greater sensitivity in DSE detection with higher
consistency from multiple trial replicates compared to alignment-based methods. These results
provide objective evidence for the robustness of SpaceExpress in detecting DSE genes across
different biological conditions.

SpaceExpress pinpoints behavior-related changes in the honey bee brain spatial
transcriptome

We illustrate the power of SpaceExpress by using it to identify DSE patterns associated with
honey bee behavior. Honey bees form complex societies with specific castes responsible for
different tasks and previous research has revealed that neurotranscriptomic state is tightly linked
to a bee's occupation in the hive[20-22]. However, these transcriptomic states[20,21,23,24] have lacked
spatial resolution and it remains unclear whether behavioral differences between castes such as
soldiers and foragers are associated with regional variations in expression. Studying anatomical
locations responsible for this DSE may reveal principles of how complex behaviors emerge from
the spatial organization of cells in the bee brain.

To answer this question, we profiled the spatial expression of 130 behavior-related genes[23,24] in
similarly aged soldier and forager brains using our custom MERFISH platform and performed
spatial comparisons using SpaceExpress. For the forager brain, we intentionally used a section
that was only partially imaged (**Figure 4a**), to mimic the common challenges for differential spatial
analysis, including size/morphology variation and partially damaged tissues. **Figure 4b** visualizes
a four-dimensional natural coordinate system learned by SpaceExpress on the two brains.
Despite dramatic visual differences between the samples, each learned dimension successfully

captures anatomical structures previously characterized in honey bee brains, and assigns similar
coordinates to the same regions across both brains. Additionally, different embedding dimensions
distinguish different structures. For example, dimensions 2 and 4 clearly distinguish between
mushroom bodies (MB), the primary region responsible for learning[25], and the rest of the brain.
Dimension 1 differentiates regions within MB, which correspond to different subtypes of neurons
called Kenyon cells (**Supplementary Figure S6**). Dimension 4 differentiates between the central
complex (middle embedding values) and the optic lobes (low embedding values), which is
especially noteworthy because the central complex was not even imaged in the forager brain
sample. Additionally, comparisons with GraphST and STAGATE show that SpaceExpress
provides a more consistent alignment of these regions, underscoring its effectiveness in
generating a shared coordinate system for samples under different conditions (**Supplementary**
Figure S7).

Based on these embeddings, SpaceExpress found 27 unique DSE genes at an FDR of 0.005
across all four embedding dimensions (**Figure 4c**). Our spline model allows for different average
expression levels (see **Methods**), ensuring that these findings were not driven by overall
differential expression (DE) of a gene between the two samples. To confirm this, we tested for
overall differential expression of each gene using a Wilcoxon test, and found DE and DSE ranks
of genes to be uncorrelated (**Supplementary Figure S8**).

A unique advantage of the SpaceExpress framework is that its embedding visualizations can be
used along with its fitted spline models to interpret DSE patterns and derive biological insight. For
each DSE gene, SpaceExpress produces a consolidated report with figures showing the spatial
expression patterns in the samples, visualizations of the embedding dimension in which the DSE
was detected, and a plot of the best-fit spline models showing how expression varies with the
embedding dimension in each sample. We highlight a few of our results below.

Some genes show a similar spatial expression pattern across an embedding dimension between
the two phenotypes, except in a localized region. This is evident in the dimension 1 spline plots
for *hr38* (DSE FDR 1.8×10^{-11}), where the expression in the soldier differs from the forager
approximately between embedding values -7 and -4, where it shows a peak (**Figure 4e**). The
embedding visualization of dimension 1 (Figure 4b) shows that these embedding values roughly
correspond to the large Kenyon cells of the MB (Supplementary Figure S6), indicating that *hr38*
is more spatially localized to these cells in the soldier than in the forager (**Figure 4d**).
Corroborative of these findings, *in situ* hybridization has previously shown that spatial localization
of this immediate early gene within the MB varies with aggressive responsiveness in honey bees,
though overall expression levels do not change across aggressive or affiliative contexts[26].

Some genes' spatial expression patterns look different across most of the embedding dimension.
For example, the dimension 3 spline plots for *fasciclin-1* (DSE FDR 0.0) show that in the soldier,
the gene has higher expression for embedding values in the range $3 - 5$ (blue in Fig. 4b)
compared to values around -3 (green in Fig. 4b), while this trend is absent for the forager (**Figure**
4g). Given the embedding visualization of dimension 3 (Figure 4b), this suggests that *fasciclin-1*
is spatially more expressed near the optic lobes than large Kenyon cells of MB in the soldier, in
contrast to the pattern in the forager (**Figure 4f**). Interestingly, *fasciclin-1* is associated with
synaptic plasticity[27], which is important for behavioral differences.

We also found the circadian clock gene *per* to be DSE along dimension 4 (DSE FDR 2.6×10^{-4}),
and the associated visualizations (**Figure 4b,h,i)** reveal that its expression is concentrated in the
MB in both brains, but more so in the soldier. We anticipate substantial variation in timekeeping
strategies between soldiers and foragers: foraging behavior is gated by diurnal nectar and pollen
availability[28], whereas soldiers must always be prepared to launch an attack at the earliest
indication of a threat. Therefore, variation in circadian-associated localization may indicate a
mechanistic underpinning of phenotypic variation across soldiers and foragers.

SpaceExpress Reveals Experience-Dependent Transcriptomic Gradients in the Mouse
Visual Cortex

A common paradigm in transcriptomics is to identify molecular changes, e.g., DE genes, between
two groups of replicates representing different biological conditions or treatments. SpaceExpress
is the first tool to extend this influential paradigm to spatial transcriptomics in a statistically rigorous
manner. The main idea is to include random effects terms into the SpaceExpress spline
regression model (Methods).

We next applied SpaceExpress to map vision-dependent transcriptomic patterns with cellular
resolution in the mouse primary visual cortex. For this, we used a previously published MERFISH
dataset from postnatal day 28 (P28) mice[16] (**Figure 5a**). This dataset consisted of coronal
sections from normal-reared (NR) and dark-reared (DR) mice (two biological replicates per
condition). DR mice were raised under normal conditions, but moved to the dark at P21, which
marks the beginning of a critical period where development of the visual cortex is particularly
sensitive to visual deprivation[29]. For each animal, one anterior and one posterior slice were
profiled for a panel of 500 genes.

We conducted two separate analyses using SpaceExpress: one for the posterior V1 sections and
one for the anterior V1 sections, each using all four P28 datasets (NR – replicates NRa, NRb, and
DR – replicates DRa, DRb) to learn a shared, four-dimensional common coordinate system. In
the posterior V1, dimensions 1 and 2 formed smooth gradients that closely tracked cortical depth
along the pial-ventral axis, in all four replicates (**Figure 5b**). By contrast, in the anterior V1,
dimensions 2 and 3 exhibited inverse gradients, with coordinate values increasing toward the pial
surface (**Figure 5c**). These concordant embedding patterns confirm that SpaceExpress recovers
the intrinsic laminar organization of V1 across rearing conditions[30].

We next tested each gene for differential spatial expression (DSE) along each dimension,
performing this analysis separately for the posterior and anterior datasets. We used the multi-
replicate DSE analysis version of SpaceExpress (see Methods), which models variation across
biological replicates and controls for sample-to-sample variation and heterogeneity in cell type
composition. The number of DSE genes identified in each dimension for both analyses is
summarized in **Figure 5g**. Aggregating across all four dimensions, both anterior and posterior
analyses identified 26 DSE genes (**Figure 5d**), with 14 genes common to both regions. Notably,
nine of these shared genes are canonical immediate-early genes (IEGs) that are known to be
controlled by neuronal activity[31]. Using differential gene expression analysis, IEGs were
previously reported to be upregulated under dark-rearing conditions across several neuronal
subclasses in the visual cortex[16] (**Figures 5d,e**); however, their condition-related difference in

spatial pattern had not been quantified. To summarize their SpaceExpress-detected DSE pattern,
we plotted the normalized expression of the 9 DSE IEGs against cortical depth (**Figure 5e**). This
analysis revealed that IEGs are particularly upregulated in upper cortical layers in DR. This pattern
is exemplified by the IEGs Nr4a3 and Fosb (**Figure 5h-k**). Under DR, Nr4a3 shows upregulation
in low values of dimension 2 and in higher values of dimension 3 in posterior and anterior slices
respectively (**Figures 5h,i**). (Note that dimension 2 in posterior slices increases with cortical
depth, while dimension 3 in the anterior slices is anti-correlated with cortical depth.) Observed
changes in Fosb are also consistent with specific upregulation in upper layers of the cortex
(**Figures 5j,k**). As upper-layer cortical excitatory neurons have been suggested to be highly
susceptible to sensory deprivation[32], results from SpaceExpress are consistent with expected
roles for IEGs in mediating sensory input-dependent development of upper-layer cortical neurons
during the critical period.

SpaceExpress pinpoints behavior-related changes in the mouse hypothalamic preoptic
area spatial transcriptome

We next applied SpaceExpress to identify genes whose spatial expression patterns in the
hypothalamic preoptic area of virgin male mice differ based on pup-exposure, which is associated
with aggressive behavior. We analyzed MERFISH data from one control and one pup-exposed
mouse, collected by Moffitt et al.[33]. The four-dimensional natural coordinate system learned by
SpaceExpress (**Figure 6c**) captures meaningful tissue structure despite dramatic differences
between the samples. Roughly, dimension 1 identifies the dorsal-ventral axis, dimension 2
highlights a gradient that increases from the medial preoptic area (MPA) and fornix (fx) to the
periventricular zone (PV) and bed nucleus of the stria terminalis (BST), dimension 3 spans third
ventricle (V3) and fx on one extreme and MPA on the other, while dimension 4 distinguishes the
median preoptic nucleus (MPN) from the rest of the MPA. Importantly, these embeddings do not
simply cluster known brain regions, they also identify continuous gradients within each region.
For example, the dimension 3 embeddings exhibit a gradient within the V3 region. To support
these embedding interpretations, we transferred expression-based brain region annotations from
Li et al.[34] using the tool SLAT[8] (**Figure 6a,** Methods).

Based on these embeddings, SpaceExpress identified 41 unique DSE genes at an FDR of 0.005
(**Figure 6b**). These include the gene *Th* (tyrosine hydroxylase), which is DSE with FDR ~0 in
dimension 3. **Figure 6d** and **Supplementary Figure S9** show that the contrast in *Th* expression
between fx and V3 regions versus BST and MPN regions is stronger in the pup-exposed mouse
compared to the naïve mouse. Th is crucial for neurotransmitter production, including dopamine
and norepinephrine, which regulate emotional states including aggression[35,36]. V3 surrounds key
nuclei in the hypothalamus that regulate aggression[37] while fx is linked to emotional regulation
through connections to hypothalamus and hippocampus[38]. The fx- and V3-specific difference in
Th expression thus suggests catecholamine signaling as a contributor to aggression toward
pups[39]. A similar case of DSE is observed for the gene *Tac1*, which exhibits elevated expression
in the V3 region, especially in the pup-exposed mouse (**Figure 6e, Supplementary Figure S10**):
its expression along dimension 3 has opposing trends in the two samples (FDR ~ 0). *Tac1*
encodes "substance P", a neuropeptide that is a key modulator of neurotransmission in regions
involved in aggression such as V3[40], and our finding suggests that it may heighten aggression by

amplifying the hypothalamic response to stressful stimuli such as pup-exposure. Many other
genes well-known to be involved in aggression and parental behavior were also identified as DSE,
including *oxytocin*[41,42], *urocortin-3*[43], and *galanin*[44] (**Supplementary Figure S11**).

We also analyzed this dataset in the multi-replicate mode comparing three control and three pup-
exposed mice and found additional DSE genes, e.g., *Gad1*, that were not identified above
(**Supplementary Note 3**).

DISCUSSION

SpaceExpress enables the identification and biological interpretation of genes whose spatial
expression patterns correlate with phenotypic traits. SpaceExpress can compare the spatial
structure of gene expression across samples of highly organized tissue, capturing the fine spatial
features encoded in the tissue's transcriptome. This contrasts with existing tools that operate at
coarser resolutions such as spatial domains[5], which can be used to identify differential gene
expression in specific spatial domains across tissues[45-47]. It also differs from tools that study local
spatial contexts, such as spatial "niches"[48] and "microenvironments"[49], which are more suitable
for comparing tissues with repeating niche features such as tumor samples.

A possible alternative to SpaceExpress is to align tissue samples and compare gene expression
between aligned cells. However, this strategy faces technical challenges posed by structural
variations across tissue samples, which makes alignment difficult[50], and it is hard to scale beyond
pairwise comparisons (e.g., handling replicates) without treating one sample as a reference[51].
Our approach of leveraging positional information encoded in gene expression offers a more
robust solution. Furthermore, it has the added advantage that we can visualize expression along
the axes of a natural coordinate system adapted to intrinsic tissue structure and thus biologically
interpret the statistically detected DSE. A recent method called "River"[11] uses alignments between
ST samples to describe their cellular locations in a common reference frame, and then uses a
machine learning approach to identify any gene whose spatial pattern is informative of the
sample's phenotypic group. However, DSE detection by this strategy is prone to errors
propagated from the initial alignment step, which can be challenging in many real scenarios[52],
does not provide a rigorous statistical test of difference, and is based on the assumption that
every cell in one condition is distinguishable (in its spatial transcriptome) from every cell in the
other condition.
434
SpaceExpress, like other ST analysis tools, combines transcriptomic and spatial information to
obtain low-dimensional embeddings of cells. A unique property of these embeddings is that the
expression vector fully determines the embedding. This property enables the tool to embed cells
profiled in non-spatial single-cell experiments as well. It can thus be used for "spatial
reconstruction"[9,10] without explicitly mapping to an arbitrary reference sample as is done today.
The ability to generalize to unseen spatial data is also key to the model's robustness for learning
natural coordinate systems (**Supplementary Figure S12**).
442

443 The statistical methodology of DSE detection in SpaceExpress is capable of associating spatial
444 expression patterns with quantitative as well as discrete traits. We have described results for
445 binary traits but the underlying statistical framework allows quantitative traits to be accommodated
446 as well. The model is highly flexible and can optionally accommodate cell type information through
447 additional terms in the model (Methods).
448
An important direction of future research is to systematically explore model complexity via tuning
of hyperparameters, including the number of dimensions of the embedding space. Though the
embedding space is meant to capture space (typically two- or three-dimensional), there can be
value to using more than two embedding dimensions as each such dimension is meant to capture
a biologically meaningful spatial pattern and there may be more than two such patterns, just as
how the intrinsic dimension of a network object can be arbitrarily large.

METHODS

SpaceExpress Embeddings

The first step of SpaceExpress is to represent cells in a spatial transcriptomic (ST) dataset as low-dimensional vectors ("embedding" vectors or "embeddings"). Formally speaking, let the ST data set comprise a collection of cells C with each cell $c \in C$ being described by a gene expression vector $g(c)$ (of dimensionality G = number of genes) and a spatial location vector $x(c)$ of dimensionality two or three. Thus, the data set may be denoted by $D_{ST} = \{c, g(c), x(c)\}_{c \in C}$. Given D_{ST} and an integer d, the dimensionality of desired embedding vectors, SpaceExpress learns a function $F: R^G \to R^d$ that can map any given gene expression vector $g(c)$ to its d-dimensional embedding $p(c)$ such that for any two cells c_i, c_j in D_{ST}, the Euclidean distance between $p(c_i)$ and $p(c_j)$ reflects the spatial proximity between c_i and c_j. In this sense, the embedding function $F(g(c))$ is said to "capture space".

To make the previous definition precise, consider a k-nearest neighbor ("knn") graph whose nodes are all cells $c \in C$, and a directed edge from c_i to c_j is included if and only if cell c_j is among the k nearest neighbors of cell c_i based on their physical locations x. Here, k is a hyperparameter of the method, which is chosen as the smallest integer for which the graph is a connected graph, i.e., there is a path from every node to every other node. We now define the distance d_{ij} between any two cells c_i and c_j as the shortest path distance between their corresponding nodes in the knn graph. The function F introduced above is learnt so as to minimize the loss function $L = \sum_{c_i \in C, c_j \in C, i \neq j} \frac{1}{d_{ij}^2} \left(\left\| p(c_i) - p(c_j) \right\| - d_{ij} \right)^2$, where $p(c) = F(g(c))$ and $\|\cdot\|$ represents the L_2-norm of a vector. This loss function, henceforth called "KK" loss, is a special form of the Kamada-Kawai loss function used in network layout applications[53]. It is a sum over all pairs of cells (c_i, c_j) of the squared difference between d_{ij} (their spatial distance, as defined above) and $\left\| p(c_i) - p(c_j) \right\|$ (Euclidean distance between their embedding vectors), but with each term being weighted by $\frac{1}{d_{ij}^2}$, thus down-weighting errors made on more distant pairs of cells. The training of function F, also called the "model", can be done on one ST data set, as explained above, or on multiple data sets; in the latter case, the loss function used is the sum over the KK loss on each data set. Once trained, the model may be used to map the cells of any new data set, using only their expression vectors $g(c)$, to their corresponding embeddings $p(c)$, without requiring their spatial locations. SpaceExpress determines the optimal value of k by iteratively increasing k from 4 to 20 and selecting the smallest k that results in a fully connected graph. If no such k is found within this range, it first generates a graph with the smallest k that minimizes the number of connected components, then connects the physically closest pairs of nodes from different components to achieve full connectivity. Specifically, for n disconnected components, it adds n−1 edges to connect them: it finds the closest cell pair straddling two disconnected components and connects them with an added edge, repeating this process until all components are mutually connected.

When training on multiple ST datasets, SpaceExpress performs joint training to learn a shared
embedding function F that captures spatial relationships across all datasets. Suppose we
have n datasets. For each dataset, the same number of cell pairs are randomly selected to ensure
equal contribution from each dataset. The KK loss is computed separately for each dataset,
resulting in individual losses L_i for $i = 1, 2, ..., n$. The total loss function used for training is the
sum of the individual losses: $L = \sum_{i=1}^{n} L_i$. This approach allows the model to generalize across
different samples and conditions, effectively integrating multiple datasets into a unified embedding
space. In the implementation, during each training epoch, a fixed number of cell pairs from each
dataset are randomly sampled to compute the KK loss to prevent the model from being biased
toward larger datasets.

The function F that generates $p(c) = F\big(g(c)\big)$ is a neural network with an input layer comprising
G neurons (one for each gene's expression value), an output layer with d neurons and a single
hidden layer with h neurons (h is a hyperparameter), and two fully connected layers, for a total of
$O(Gh + dh)$ free parameters. Parameter optimization is implemented using Adam optimizer. The
full collection of $\binom{|C|}{2}$ cell pairs is partitioned at random into N_{mb} equal subsets and these
subsets are used as mini-batches in each epoch. Batch normalization is used as a form of
regularization, and though we explored the use of dropouts and L1/L2 regularization, these
features were not used in the final implementation as they did not yield significant improvement
on unseen data. The embedding dimension was fixed to 4 throughout all analyses, as we
observed minimal performance differences across a range of dimensions (4, 8, 16, 32). A detailed
evaluation of dimensionality effects is provided in **Supplementary Figure S13**.

Preprocessing: Each dataset was preprocessed by filtering out genes expressed in fewer than 3
cells and cells expressing fewer than 3 genes. The data was normalized to a target sum of 10,000
(each cell by total counts over all genes) and log transformed. To minimize the influence of
extreme values, gene expression values were capped at the 95th percentile. The user may
optionally select to use either all genes or only the most highly variable genes (HVGs); the number
of HVGs to use (n_{HVG}) is a configurable parameter. If the user selects to use HVG-based analysis,
SpaceExpress identifies the n_{HVG} highest variance genes in each sample, calculates the
intersection of these HVGs across all samples, and creates a limited version of each sample's
data that includes only these genes, and scales each gene in each sample to unit variance and
zero mean.

Synthetic data sets

"Early embryo slice" data set: Each such data set comprises 3,915 cells placed uniformly in an
elliptical area whose major and minor axes, called "anterior-posterior" (AP) or "dorso-ventral" (DV)
axes, have lengths of 100 and 50 units, respectively (Figure 2a). The transcriptomic data
synthesized spans 100 genes that fall into the following categories:

•   AP genes (20): these include two "core" AP genes AP1 and AP2, whose expression in a cell

with coordinates (x, y) is given by $e^{\frac{-x^2}{40^2}}$ and $e^{\frac{-(x-100)^2}{40^2}}$ respectively. Here, the major (AP) and

minor (DV) axes defined the x and y coordinates of a cell, with x ranging from zero at the left

(anterior) end of the major axis to one at the right (posterior) end, and y ranging from zero at

the top (dorsal) end of the minor axis to one at the bottom (ventral) end. An additional 18

"derivative" AP genes are assigned expression values according to the formula $\alpha_i \cdot AP_1 +$

$(1 - \alpha_i) \cdot AP_2$, i.e., a weighted combination of AP1 and AP2 expression profiles. Each

derivative gene uses a weight α_i chosen at random from the range. Note that all 20 AP genes

have an "AP" expression pattern, i.e., the expression is identical in all cells at a fixed x

coordinate (position along the AP axis).

•   DV genes (20): these are assigned expression patterns similarly to the AP genes except that

their expression varies along the DV axis and all cells with the same y coordinate have the

same expression value for a DV gene. There are two core DV genes – DV1 and DV2 – with

expression values given by $e^{\frac{-y^2}{20^2}}$ and $e^{\frac{-(y-50)^2}{20^2}}$ respectively, and 18 derivative DV genes whose

expression is a weighted combination of DV1 and DV2.

•   APDV genes (20): each such gene is defined by the formula $\frac{1}{2}(\gamma_{k0} \cdot AP_1 + (1 - \gamma_{k0}) \cdot AP_2 +$

$\gamma_{k1} \cdot DV_1 + (1 - \gamma_{k1}) \cdot DV_2)$ for $\gamma_k \sim \text{Unif}[0,1]$, i.e., sum of an AP and a DV expression pattern,

each being a weighted sum of the respective core genes.

•   Randomly expressed genes (40): For each of these genes, we assign an expression pattern

in each cell independently at random in the range $[0,1]$. These genes have no spatial pattern.

Once each gene's expression level is assigned in each cell, an expression "count" is sampled

from the Poisson distribution with parameter $1000e$.

Use of existing software

In this study, we utilized SEDR[54], GraphST[15], and STAGATE[14] as benchmark methods to

compare their embeddings with SpaceExpress. For SEDR, during the preprocessing step, if a

dataset contained fewer than 200 genes, we removed the PCA step used for dimensionality

reduction to 200. For STAGATE, we set each spot or cell to have six neighboring cells, following

the original paper's protocol. All other parameters were kept at their default settings as specified

by the original authors.

Evaluation of learned natural coordinate

We jointly trained SpaceExpress on two synthetic embryo samples to learn a shared

four-dimensional coordinate system. To evaluate this coordinate system, we defined two

complementary metrics: inter-sample agreement and natural coordinate fidelity. Inter-sample

agreement measures whether cells with known correspondences across samples occupy nearby

positions in the learned coordinate system ("embedding space"). Let $x_i^{(1)}$ and $x_j^{(2)}$ denote the true

spatial coordinates of cells i (in sample 1) and j (in sample 2), respectively, and let $p_i^{(1)}$, $p_j^{(2)}$ be

their learned embedding coordinates. This spatial coordinatesystem is defined along a biologically
meaningful axis common to both samples , specifically the anterior–posterior (AP) axis in the
synthetic embryo (Figure 2b,d), cortical depth in the mouse visual cortex (Figure 2e), or
histologically defined depth in the DLPFC samples (Figure 2g, Supplementary Figure S14). For
each cell i in sample 1, we find its nearest neighbor in the other sample in embedding space:

$$576 \qquad j_i^* = argmin_j \left\| p_i^{(1)} - p_j^{(2)} \right\|_2$$

577 We then compute a normalized spatial score

$$578 \qquad s_i = \begin{cases} max\left(0, 1 - \dfrac{\left\| x_i^{(1)} - x_{j^*}^{(2)} \right\|_2}{\tau}\right), & if \left\| p_i^{(1)} - p_{j^*}^{(2)} \right\|_2 \leq \overline{d}_p, \\ 0, & otherwise, \end{cases}$$

579 where $\overline{d}_p$ is the mean distance to the k-th nearest neighbor (k=10) in embedding space across
580 all cells, and τ is the mean physical distance to the k'-th nearest neighbor ($k' = \lfloor 0.5N \rfloor$) in true
581 spatial coordinates.

582 The overall inter-sample agreement is then

$$583 \qquad \text{Inter-Sample Agreement} = \frac{1}{N}\sum_{i=1}^{N} s_i$$

ranging from 0 to 1, with higher values indicating stronger agreement between samples. The
condition $\left\| p_i^{(1)} - p_{j^*}^{(2)} \right\|_2 \leq \overline{d}_p$ checks if the nearest neighbor of cell i in the other tissue sample is
within a small distance. If this condition is met, we examine the physical distance
$\left\| x_i^{(1)} - x_{j^*}^{(2)} \right\|_2$ between cell i and that nearest neighbor (such a distance is defined via the
correspondence between cells in the two tissue samples, known as part of ground truth). The
physical distance is converted to an "agreement" score by a linear transformation $1 - \dfrac{\left\| x_i^{(1)} - x_{j^*}^{(2)} \right\|_2}{\tau}$,
truncated at zero. This is the contribution of cell i to the overall inter-sample agreement. If the
condition $\left\| p_i^{(1)} - p_{j^*}^{(2)} \right\|_2 \leq \overline{d}_p$ is not met, we ignore cell i in calculating inter-sample agreement.
For the Drosophila embryo (Figure 2i), the coordinate system is defined by annotated spatial
domains, so agreement is measured as the fraction of cells whose nearest neighbor from the
other sample in embedding space belongs to the same domain.

Natural coordinate fidelity measures whether any single embedding axis closely aligns with the
true spatial organization. For each sample, let $P \in \mathbb{R}^{N \times D}$ be the embedding matrix (with N cells
and D dimensions), and let $z \in \mathbb{R}^D$ denote the known natural coordinate of each of the N cells.
Here, z is the position along the AP axis in the synthetic embryo (Figure 2b,d), cortical depth in

mouse visual cortex (Figure 2e), or histologically defined depth in DLPFC samples (Figure 2g,
Supplementary Figure S14. Natural coordinate fidelity is defined as:

$$\text{Natural coordinate fidelity} = \max_{d} |Corr(p_{:,d}, z)|$$

That is, we compute the absolute Pearson correlation between the natural coordinate of each cell
and its learned coordinate, for each dimension $d = 1, ..., D$ separately, and the take the maximum
of these correlations. This yields a value between 0 and 1, with higher values indicating that at
least one embedding axis aligns closely with true natural coordinate. For the Drosophila embryo
(Figure 2i), spatial organization is defined by annotated domains, so fidelity is measured as the
fraction of cells whose nearest neighbor in embedding space falls within the same domain.

Real data sets

We utilized SpaceExpress to analyze spatial transcriptomic datasets generated by various
platforms, including 10x Visium, Stereo-seq and MERFISH. The human dorsolateral prefrontal
cortex (DLPFC) dataset, generated using the 10x Visium platform, includes 12 sections from three
individuals[17]. This dataset contains 33,538 genes, with the number of spots per section ranging
from 3,460 to 4,789. The Drosophila embryo dataset, generated using the Stereo-seq platform[18],
includes 15 sections from the same embryo. For evaluation, we used three consecutive sections
(IDs 6 to 8), with each section containing between 815 and 1,272 spots and 13,668 genes. The
honey bee brain dataset was generated in-house using MERFISH, profiling 130 behavior-related
genes in one soldier and one forager brain. The dataset includes 11,373 cells for the soldier and
10,334 cells for the forager; additional details of data generation are provided in the Methods
section. The vision-dependent transcriptomic dataset of the mouse primary visual cortex[16],
generated using MERFISH, includes data from postnatal day 28 (P28) mice. It comprises two
samples from dark-reared (DR) mice and two from normal-reared (NR) mice, with each sample
providing data for both anterior and posterior regions, which are approximately 550 μm apart,
were profiled, yielding eight slices in total. This dataset contains 500 genes, with the number of
cells ranging from 2,333 to 8,768. The mouse brain hypothalamic preoptic dataset[33], which
consists of six samples, each with 4,908 to 5,998 cells and 155 genes. For 1:1 comparison
between naïve and pup-exposed samples, we used one male naïve sample (ID 5) and one male
pup-exposed sample (ID 36), both at bregma 0.16. For the analysis of multiple biological
replicates from different phenotype groups, we selected male naïve samples (IDs 5, 6, 7) and
male pup-exposed samples (IDs 34, 35, 36), all from the same bregma (0.16).

Synthetic data sets with differential spatial expression

To test the ability of SpaceExpress to learn "shared" coordinate systems, we synthesize a pair of
"early embryo slice" data sets that are both organized using the AP and DV axes as coordinate
axes, but exhibit a varying degree of divergence in that organization. Specifically, we follow the
procedure outlined above to generate expression data for all but the "APDV" category of genes.
Note that a gene in this category was assigned its expression value in a cell with coordinates

(x, y) according to the formula $1/2 \left(\gamma_{k0} \cdot AP_1 + (1 - \gamma_{k0}) \cdot AP_2 + \gamma_{k1} \cdot DV_1 + (1 - \gamma_{k1}) \cdot \right.$
$\left. DV_2 \right)$, where $\gamma_k \sim \text{Unif}[0,1]$, i.e., a random combination of an AP pattern and a DV pattern. To
simulate differential spatial expression of these genes, we now modify this procedure as follows:
in the first data set, we zero out the contribution of the AP pattern, so the new formula is $1/2 \left(\gamma_{k1} \cdot \right.$
$\left. DV_1 + (1 - \gamma_{k1}) \cdot DV_2 \right)$, meaning that such an "APDV" gene exhibits an exclusively DV pattern.
In the second data set, we weight the AP pattern by a constant $\mu \in \{0.1, 0.2, \ldots 0.9\}$, so the new
formula is $1/2 \left((\mu \cdot AP_1) + (\gamma_{k1} \cdot DV_1 + (1 - \gamma_{k1}) \cdot DV_2) \right)$ and the "APDV" genes exhibit both a
DV and an AP pattern with the strength of the latter being controlled by μ. All 20 APDV genes use
the same value of μ in a data set. Thus, a pair of data sets comprises one embryo where APDV
genes have zero AP pattern and one embryo where the AP pattern is weighted by a fixed value
of μ. In this way we can generate pairs of data sets exhibiting varying degrees of DSE, by
progressively increasing the value of μ.

Benchmarking SpaceExpress and an Alignment-Based Baseline method for DSE Detection

We implemented an alignment-based approach as a complementary method for DSE detection
that does not rely on learned coordinate systems. This approach aligns two ST tissue samples
using PASTE2[50] under default settings, with the overlap parameter set to the ratio of the number
of cells between the two samples. A gene's expression in aligned cell pairs (one cell from each
tissue) is then examined, and the correlation coefficient is calculated. A perfect correlation of +1
indicates no differential spatial expression, while smaller values denote greater DSE, with -1
representing the strongest DSE. This is repeated for every gene and produces a ranked list of
genes ordered by their DSE scores. On the other hand, SpaceExpress also produces a ranked
list of DSE genes ordered by their FDR values.

To compare SpaceExpress and the alignment-based method, we first used either method on pairs
of tissues from the same condition (intra-group comparison) to obtain their respective ranked lists
of DSE genes. For either list, the 5th percentile DSE score (FDR for SpaceExpress-DSE and
correlation for the alignment-based method) across all genes was noted. This was repeated for
two intra-group tissue pairs (see Figure 3i,j) and the average of the 5th percentile DSE scores in
these two analyses was designated as the DSE score threshold for the respective method. This
threshold was then applied to designate DSE genes in inter-group comparisons. (We assume that
intra-group comparisons provide a null model for DSE scores, so the threshold derived in this way
controls false positive errors at 5% for both methods.) To evaluate robustness, the number of
times each gene was designated as DSE across the four inter-group comparisons was noted. A
robust method should consistently detect shared DSE genes across multiple inter-group
comparisons. Additionally, River[11] was employed as a competing method to evaluate DSE
detection in comparison with SpaceExpress. For the prediction model in River, we defined each
cell's corresponding label as its condition group.

Differential Spatial Expression analysis between two samples

Goal: In this section, we describe how SpaceExpress assesses differential spatial expression
(DSE) of genes between two ST samples. We assume that the first step has learnt the shared
coordinate system of the two samples, by embedding each cell in a latent Euclidean space. For
each embedding dimension, SpaceExpress analyzes each gene separately and then combines
the results via appropriate multiple-hypothesis testing procedure. As a result, we get a set of DSE
genes for each embedding dimension.
683
684 *Pre-processing:* For a given gene, we log-transform the expression and then remove the cells
685 that are beyond 4 standard deviations away from the mean. We drop a gene from the analysis if
686 the transformed expression values are all zero in any group. We scale (without centering) the
687 transformed expression within each group to have unit variance.
688
Linear Models: As mentioned in the goal, the analysis involves regressing each gene on each
embedding dimension separately. For the sake of interpretation and statistical inference, we
decided to use a linear model to fit gene expression against a given embedding dimension.
However, a linear function of the embedding values will not be sufficient to capture the variation.
Therefore, we use a spline regression strategy, where we create spline basis functions based on
an embedding dimension. These spline basis functions are non-linear functions of the embedding
dimension. Consequently, we can regress gene expression as a linear function of these spline
basis functions. This approach allows us to retain the statistical interpretation advantages of a
linear model while achieving a fit that is a non-linear function of the embedding. The generation
of spline basis functions depends on a crucial parameter known as degrees of freedom. Higher
degrees of freedom correspond to more non-linearity. We generate natural cubic spline bases
with 300 degrees of freedom using the *ns* function from the *splines* package in R. For cell i, we
denote the transformed and scaled gene expression by Y_i, the generated spline basis vector by
X_i^d (d denotes the embedding dimension), and the group identity by G_i. We fit the following two
linear models using the *lm* package in R:
704

$$Y_i = \alpha + \beta^T X_i^d + \gamma G_i + \delta^T \left(G_i X_i^d \right) + \epsilon_i \text{ (full model: with interaction term),}$$

$$Y_i = \alpha + \beta^T X_i^d + \gamma G_i + \epsilon_i \text{ (null model: without interaction term).}$$

Here ϵ_i are assumed to be independent Gaussian errors with zero mean. The interaction term δ
represents the differential spatial expression between the two groups. Therefore, we test the
significance of δ via a likelihood ratio test (LRT) between full and the null models. Furthermore,
the γG_i term in the null model accounts for the mean expression difference between the two
groups, ensuring that the observed effects are not simply due to overall differential expression
(DE).

Multiple Hypothesis testing via Empirical Null method: For embedding dimension d, we collect
the LRT based chi-squared test statistic for each gene. We estimate the empirical null distribution
and then the false discovery rate (FDR) for each gene using the method described by Ren et al.[19].
As an implementation detail, the quantile parameter, q, for empirical null calculation that we use
is 0.75.

Multi-replicate Differential Spatial Expression analysis

720 Goal: In this section, we describe how SpaceExpress assesses differential spatial expression
721 (DSE) of genes across two groups with multiple replicates.
722
Pre-processing: For a given gene, we log-transform the expression and then remove the cells
that are beyond 4 standard deviations away from the mean. We drop a gene from the analysis if
the transformed expression values are all zero in any group. We scale (without centering) the
transformed expression within each group to have unit variance. Furthermore, we generate
natural cubic spline bases with 50 degrees of freedom using the *ns* function from the *splines*
package in R.
729
730 *Linear Mixed Effects Models:* For cell i in replicate j we denote the transformed and scaled gene
731 expression by Y_{ji}, the generated spline basis vector by X_{ji}^d (here d denotes the embedding
732 dimension), the group identity by G_{ji}, and the cell type of cell i by C_i. We fit the following linear
733 mixed effects model using *lmer* package in R:
734

$$Y_{ji} = \alpha + b_j + \beta^T X_{ji} + \gamma G_{ji} + \delta^T(X_{ji}G_{ji}) + \eta C_i + \epsilon_{ji} \text{ (full model: with interaction term),}$$

$$Y_{ji} = \alpha + b_j + \beta^T X_{ji} + \gamma G_{ji} + \eta C_i + \epsilon_{ji} \text{ (null model: without interaction term).}$$

Here b_j represents zero mean random intercept term. Other coefficients $\alpha, \beta, \gamma, \delta$, and η are
assumed to be fixed effects across replicates. Furthermore, ϵ_{ji} are assumed to be independent
Gaussian errors with zero mean. The interaction term $\delta^T(X_{ji}G_{ji})$ represents the differential
spatial expression between the two groups. Therefore, we test the significance of δ via a
likelihood ratio test between full and the null models. We can drop the C_i term and the coefficient
η if we have just one cell type in the data or we do not want to capture the effect of cell types on
the gene expression.

Multiple Hypothesis testing via Empirical Null method: For each embedding dimension, we collect
the LRT based chi-squared test statistic for each gene. We estimate the empirical null distribution
and then the false discovery rate (FDR) for each gene using the method described in Ren et al.[19].
As an implementation detail, the quantile parameter, q, for empirical null calculation that we use
is 0.7.

MERFISH data generation for honey bee brain

MERFISH sample preparation: Honey bee MERFISH samples were prepared using a slightly
modified published method[33]. Specifically, honey bee brains embedded in OCT were cryo-
sectioned at a thickness of 5-7 μm. To reduce spot density, samples were embedded in an
expandable poly-acrylamide gel using a previously published method[55].

MERFISH imaging: All images were acquired using a Zeiss Axiovert-200m widefield microscope
(Carl Zeiss AG) located in the IGB core imaging facility. The sample was placed into a flow cell
(Bioptechs, FCS2), filled with RNAse free 2x SSC, and connected to a lab built automated flow
system. Briefly, computer-controlled valves (Hamilton, MVP/4, 8-5 valve) are used to select which
solution was pulled across the sample by a computer controlled pump (Gilson, Minipuls 3). All
systems are controlled by a custom designed Python script that can communicate with the
microscope to start imaging or start flowing after an imaging round is done. In brief, a single round
of imaging involves staining with fluorescently labeled readout probes (10% (v/v) ethylene
carbonate (Sigma Aldrich), 0.1% Triton X-100 in 2x SSC, fluorescent DNA probe), washing with
readout wash buffer (10% (v/v) ethylene carbonate, 0.1% Triton X-100 in 2x SSC) to remove
unbound probes, and imaging buffer (5mM 3,4-dihydroxybenzoic acid (PCA; Sigma Aldrich), 2
mM trolox (Sigma Aldrich), 50 µM trolox quinone, 1:500 of recombinant protocatechuate 3,4-
dioxygenase (rPCO; OYC Americas), was flowed into the flow cell prior to imaging to reduce
photobleaching. A single quad band excitation filter (Chroma, ZET402/468/555/638x) and dichroic
(Chroma, ZT405/470/555/640rpc-UF1) were used to image all samples. Excitation was provided
by a 7 laser system (LDI WF, 89 North). Alexa Fluor 647 (Fisher scientific) labeled probes were
excited using a 647 nm laser (0.5 W) with a ET700/75m (Chroma) emission filter. Atto 565 (Atto
tec) labeled probes were excited using a 555 nm laser (1 W) with an ET610/75m (Chroma)
emission filter. Fiducial beads were imaged with a 405 nm laser (0.3 W) with a ET440/40m
emission filter. Samples were imaged with a 63x oil immersion objective (Carl Zeiss AG, 420782-
9900-000), and focus was maintained between imaging rounds using Definite Focus (Carl Zeiss
AG). After imaging is complete, a cleavage buffer (0.05 M TCEP HCl, adjusted to a pH of 7-7.2
using 1 N NaOH, in 2x SSC) was flowed across the sample to remove the fluorophores from the
probes. The cleavage buffer was washed away using RNAse free 2x SSC (0.5 mL/minute for 10
minutes). This process was repeated for a total of 8 rounds of imaging. PolyA probes were stained
after the final imaging round using the same method as described above.

MERFISH data processing: Individual FOVs were exported from czi format into 16 bit tiff format
using Zen (Carl Zeiss AG) using the image export method. Images then were reformatted into
image stacks by FOV and round. A modified copy of MERLIN[56] was used to decode MERFISH
spots. In brief, for each FOV, images from different rounds are aligned using fiducial beads that
were imaged in each round. Aligned images are then normalized, decoded, and identified spots
filtered using previously published methods[57]. Cell segmentation was done separately from
MERLIN using Cellpose[58] on PolyA and DAPI images for each FOV. To improve FOV alignment
to neighboring FOVs, the DAPI channel was used with the restitching function found in Zen (Edge
detection: on, minimal overlap: 5%, maximal shift: 15%, comparer: best, Global optimizer: best).
Using the aligned images, segmented cells that cross FOV boundaries were merged into single
cells, and global positions were generated for each spot. Spots are then assigned to cells based
on their spatial coordinates. Spots were then filtered to remove any spot smaller than 3 pixels in
size.

Obtaining neuroanatomical annotation of mouse brain

We obtained the neuroanatomical annotations of the hypothalamic preoptic area in mouse brains
based on the manual annotations in Li et al.[34]. These annotations were made in tissue sections
at Bregma -0.14 mm from animal 1, and included eight domains (V3: third ventricle; BST: bed
nuclei of the stria terminalis; fx: columns of the fornix; MPA: medial preoptic area; MPN: medial
preoptic nucleus; PV: periventricular hypothalamic nucleus; PVH: paraventricular hypothalamic
nucleus; and PVT: paraventricular nucleus of the thalamus). We used this annotated section as
reference and aligned cells from sections of naïve mice (IDs 5, 6, 7) and pup-exposed mice (IDs
34, 35, 36) at bregma 0.16mm using SLAT[8] with default settings. The domains from the reference
cells were assigned to the mapped cells.

DATA AVAILABILITY

Dorsolateral Prefrontal Cortex (DLPFC) Dataset: The 10x Visium data used in this study is available at https://github.com/LieberInstitute/spatialLIBD.git[17]. Drosophila Late-Stage Embryo Dataset: Stereo-seq data can be accessed at https://db.cngb.org/stomics/flysta3d/[18]. Mouse Hypothalamic Preoptic Area Dataset: MERFISH data was obtained from Moffitt et al.[33], available at https://datadryad.org/stash/dataset/doi:10.5061/dryad.8t8s248. Honeybee Dataset: Data is available at https://doi.org/10.13012/B2IDB-5536668_V1.

CODE AVAILABILITY

SpaceExpress is implemented in Python, and the code is available at https://github.com/YeojinKim220/SpaceExpress. The package utilizes the AnnData format and can be installed by following the instructions on the GitHub page. A tutorial demonstrating SpaceExpress with data from the mouse hypothalamic preoptic region is also provided[33].

ACKNOWLEDGEMENTS

We thank Anurendra Kumar for valuable discussion, and Seokjin Yeo and Sunny Sun for image segmentation, Marisa Asadian for help for imaging, and Adithya Karthik for assistance with alignment analysis. Funding: This work was supported by the National Institutes of Health (R35GM131819 to S.S., R35GM147420 to H.-S.H, R21HG013180 to S.D.Z. and H.-S.H.), Georgia Institute of Technology (Wallace H. Coulter Distinguished Faculty Chair: S.S.), and the European Union's Horizon 2020 Research and Innovation Program under ERC-2017-StG Grant Agreement 757583 (Brain2Bee; to Jennifer L Cook and Gene E. Robinson). Facilities: We acknowledge Core Facilities at the Carl R. Woese Institute for Genomic Biology for their microscope and staff support.

REFERENCES

1. Rao, A., Barkley, D., França, G.S. & Yanai, I. Exploring tissue architecture using spatial transcriptomics. *Nature* **596**, 211-220 (2021).
2. Marx, V. Method of the Year: spatially resolved transcriptomics. *Nature Methods* **18**, 9-14 (2021).
3. Squair, J.W. *et al.* Confronting false discoveries in single-cell differential expression. *Nature Communications* **12**, 5692 (2021).
4. Mancuso, N. *et al.* Integrating Gene Expression with Summary Association Statistics to Identify Genes Associated with 30 Complex Traits. *The American Journal of Human Genetics* **100**, 473-487 (2017).
5. Hao, S. *et al.* Cross-species single-cell spatial transcriptomic atlases of the cerebellar cortex. *Science* **385**, eado3927 (2024).
6. Yuan, Z. *et al.* Benchmarking spatial clustering methods with spatially resolved transcriptomics data. *Nature Methods* **21**, 712-722 (2024).
7. Zeira, R., Land, M., Strzalkowski, A. & Raphael, B.J. Alignment and integration of spatial transcriptomics data. *Nature Methods* **19**, 567-575 (2022).
8. Xia, C.-R., Cao, Z.-J., Tu, X.-M. & Gao, G. Spatial-linked alignment tool (SLAT) for aligning heterogenous slices. *Nature Communications* **14**, 7236 (2023).
9. Biancalani, T. *et al.* Deep learning and alignment of spatially resolved single-cell transcriptomes with Tangram. *Nature Methods* **18**, 1352-1362 (2021).
10. Martin, P.C., Kim, H., Lövkvist, C., Hong, B.W. & Won, K.J. Vesalius: high-resolution in silico anatomization of spatial transcriptomic data using image analysis. *Molecular systems biology* **18**, e11080 (2022).
11. Cui, Y. & Yuan, Z. Prioritizing perturbation-responsive gene patterns using interpretable deep learning. *Nature Communications* **16**, 6095 (2025).
12. Soneson, C. & Delorenzi, M. A comparison of methods for differential expression analysis of RNA-seq data. *BMC Bioinformatics* **14**, 91 (2013).
13. Dubuis, J.O., Tkačik, G., Wieschaus, E.F., Gregor, T. & Bialek, W. Positional information, in bits. *Proceedings of the National Academy of Sciences* **110**, 16301-16308 (2013).
14. Dong, K. & Zhang, S. Deciphering spatial domains from spatially resolved transcriptomics with an adaptive graph attention auto-encoder. *Nature Communications* **13**, 1739 (2022).
15. Long, Y. *et al.* Spatially informed clustering, integration, and deconvolution of spatial transcriptomics with GraphST. *Nature Communications* **14**, 1155 (2023).
16. Xie, F. *et al.* Spatial profiling of the interplay between cell type- and vision-dependent transcriptomic programs in the visual cortex. *Proceedings of the National Academy of Sciences* **122**, e2421022122 (2025).
17. Maynard, K.R. *et al.* Transcriptome-scale spatial gene expression in the human dorsolateral prefrontal cortex. *Nature Neuroscience* **24**, 425-436 (2021).
18. Wang, M. *et al.* High-resolution 3D spatiotemporal transcriptomic maps of developing Drosophila embryos and larvae. *Developmental Cell* **57**, 1271-1283.e4 (2022).
19. Ren, X., Wang, J., Liu, S. & Miecznikowski, J.C. Estimating empirical null distributions for Chi-squared and Gamma statistics with application to multiple testing in RNA-seq. (Department of Biostatistics, University at Buffalo).
20. Chandrasekaran, S. *et al.* Behavior-specific changes in transcriptional modules lead to distinct and predictable neurogenomic states. *Proceedings of the National Academy of Sciences* **108**, 18020-18025 (2011).
21. Shpigler, H.Y. *et al.* Deep evolutionary conservation of autism-related genes. *Proceedings of the National Academy of Sciences* **114**, 9653-9658 (2017).
22. Whitfield, C.W., Cziko, A.-M. & Robinson, G.E. Gene Expression Profiles in the Brain Predict Behavior in Individual Honey Bees. *Science* **302**, 296-299 (2003).

23.   Avalos, A. *et al.* Genomic regions influencing aggressive behavior in honey bees are
defined by colony allele frequencies. *Proceedings of the National Academy of Sciences*
117, 17135-17141 (2020).
24.   Traniello, I.M. *et al.* Single-cell dissection of aggression in honeybee colonies. *Nature*
Ecology & Evolution **7**, 1232-1244 (2023).
25.   Fahrbach, S.E. Structure of the mushroom bodies of the insect brain. *Annual review of*
entomology **51**, 209-232 (2006).
26.   Traniello, I.M., Chen, Z., Bagchi, V.A. & Robinson, G.E. Valence of social information is
encoded in different subpopulations of mushroom body Kenyon cells in the honeybee
brain. *Proc Biol Sci* **286**, 20190901 (2019).
27.   Parker, R. *et al.* Correlation of proteome-wide changes with social immunity behaviors
provides insight into resistance to the parasitic mite, Varroa destructor, in the honey bee
(Apis mellifera). *Genome Biology* **13**, R81 (2012).
28.   Moore, D. Honey bee circadian clocks: behavioral control from individual workers to
whole-colony rhythms. *Journal of Insect Physiology* **47**, 843-857 (2001).
29.   Gordon, J.A. & Stryker, M.P. Experience-Dependent Plasticity of Binocular Responses in
the Primary Visual Cortex of the Mouse. *The Journal of Neuroscience* **16**, 3274-3286
(1996).
30.   Tasic, B. *et al.* Adult mouse cortical cell taxonomy revealed by single cell transcriptomics.
Nature Neuroscience **19**, 335-346 (2016).
31.   Hrvatin, S. *et al.* Single-cell analysis of experience-dependent transcriptomic states in the
mouse visual cortex. *Nature Neuroscience* **21**, 120-129 (2018).
32.   Cheng, S. *et al.* Vision-dependent specification of cell types and function in the developing
cortex. *Cell* **185**, 311-327.e24 (2022).
33.   Moffitt, J.R. *et al.* Molecular, spatial, and functional single-cell profiling of the hypothalamic
preoptic region. *Science* **362**, eaau5324 (2018).
34.   Li, Z. & Zhou, X. BASS: multi-scale and multi-sample analysis enables accurate cell type
clustering and spatial domain detection in spatial transcriptomic studies. *Genome Biology*
23, 168 (2022).
35.   Daubner, S.C., Le, T. & Wang, S. Tyrosine hydroxylase and regulation of dopamine
synthesis. *Archives of Biochemistry and Biophysics* **508**, 1-12 (2011).
36.   Morgenroth, V.H., Boadle-Biber, M. & Roth, R.H. Tyrosine Hydroxylase: Activation by
Nerve Stimulation. *Proceedings of the National Academy of Sciences* **71**, 4283-4287
(1974).
37.   Falkner, A.L. *et al.* Hierarchical Representations of Aggression in a Hypothalamic-Midbrain
Circuit. *Neuron* **106**, 637-648.e6 (2020).
38.   Daitz, H.M. & Powell, T.P.S. STUDIES OF THE CONNEXIONS OF THE FORNIX
SYSTEM. *Journal of Neurology, Neurosurgery & Psychiatry* **17**, 75 (1954).
39.   Haller, J., Makara, G.B. & Kruk, M.R. Catecholaminergic involvement in the control of
aggression: hormones, the peripheral sympathetic, and central noradrenergic systems.
Neuroscience & Biobehavioral Reviews **22**, 85-97 (1997).
40.   He, Z.-X. *et al.* Substance P in the medial amygdala regulates aggressive behaviors in
male mice. *Neuropsychopharmacology* **49**, 1689-1699 (2024).
41.   Inada, K. *et al.* Plasticity of neural connections underlying oxytocin-mediated parental
behaviors of male mice. *Neuron* **110**, 2009-2023.e5 (2022).
42.   Nakahara, T.S. *et al.* Peripheral oxytocin injection modulates vomeronasal sensory activity
and reduces pup-directed aggression in male mice. *Scientific Reports* **10**, 19943 (2020).
43.   Autry, A.E. *et al.* Urocortin-3 neurons in the mouse perifornical area promote infant-
directed neglect and aggression. *eLife* **10**, e64680 (2021).
44.   Wu, Z., Autry, A.E., Bergan, J.F., Watabe-Uchida, M. & Dulac, C.G. Galanin neurons in the
medial preoptic area govern parental behaviour. *Nature* **509**, 325-330 (2014).

45. Vanrobaeys, Y. *et al.* Spatial transcriptomics reveals unique gene expression changes in
different brain regions after sleep deprivation. *Nature Communications* **14**, 7095 (2023).
46. Ospina, O.E. *et al.* Differential gene expression analysis of spatial transcriptomic
experiments using spatial mixed models. *Scientific Reports* **14**, 10967 (2024).
47. Qin, F., Luo, X., Cai, B., Xiao, F. & Cai, G. Spatial pattern and differential expression
analysis with spatial transcriptomic data. *bioRxiv*, 2023.07.06.547967 (2023).
48. Mason, K. *et al.* Niche-DE: niche-differential gene expression analysis in spatial
transcriptomics data identifies context-dependent cell-cell interactions. *Genome Biology*
25, 14 (2024).
49. Wu, Z. *et al.* Graph deep learning for the characterization of tumour microenvironments
from spatial protein profiles in tissue specimens. *Nature Biomedical Engineering* **6**, 1435-
1448 (2022).
50. Liu, X., Zeira, R. & Raphael, B.J. PASTE2: Partial Alignment of Multi-slice Spatially
Resolved Transcriptomics Data. *bioRxiv*, 2023.01.08.523162 (2023).
51. Clifton, K. *et al.* STalign: Alignment of spatial transcriptomics data using diffeomorphic
metric mapping. *Nature Communications* **14**, 8123 (2023).
52. Khan, M., Arslanturk, S. & Draghici, S. A comprehensive review of spatial transcriptomics
data alignment and integration. *Nucleic Acids Research* **53**(2025).
53. Kamada, T. & Kawai, S. An algorithm for drawing general undirected graphs. *Information*
Processing Letters **31**, 7-15 (1989).
54. Xu, H. *et al.* Unsupervised spatially embedded deep representation of spatial
transcriptomics. *Genome Medicine* **16**, 12 (2024).
55. Wang, G., Moffitt, J.R. & Zhuang, X. Multiplexed imaging of high-density libraries of RNAs
with MERFISH and expansion microscopy. *Scientific Reports* **8**, 4847 (2018).
56. Emanuel, G., Eichhorn, S.W. & Zhuang, X. MERlin - scalable and extensible MERFISH
analysis software. Vol. 0.1.6 (Zenodo, 2020).
57. Xia, C., Fan, J., Emanuel, G., Hao, J. & Zhuang, X. Spatial transcriptome profiling by
MERFISH reveals subcellular RNA compartmentalization and cell cycle-dependent gene
expression. *Proceedings of the National Academy of Sciences* **116**, 19490-19499 (2019).
58. Stringer, C., Wang, T., Michaelos, M. & Pachitariu, M. Cellpose: a generalist algorithm for
cellular segmentation. *Nature Methods* **18**, 100-106 (2021).

Figure Legends

Figure 1. Overview of SpaceExpress Framework and Evaluations of SpaceExpress cell
embeddings on synthetic data.

(a) Schematic contrasting conventional differential expression (DE) analysis, which tests for
changes in average gene expression levels between two samples, with differential spatial
expression (DSE) analysis, which tests for alterations in the spatial distribution of gene expression
across samples. In each scenario the left subpanel shows two idealized tissue regions colored by
expression intensity and the right subpanel shows boxplots of gene expression for Tissue 1 (blue)
and Tissue 2 (orange). Asterisks indicate significant differences and "ns" indicates no significant
spatial change. **(b)** Application of SpaceExpress to three spatial transcriptomic brain datasets
comparing two phenotypes. First, whole brain sections of honey bee workers in soldier and
forager roles are compared using custom MERFISH data. Second, mouse visual cortex samples
from normally reared and dark reared animals are compared. Third, samples from the mouse
hypothalamic preoptic area are compared between naïve males and males exposed to pups. **(c)**
SpaceExpress detects differential spatial expression between two tissue samples. In its first step
(bottom), SpaceExpress maps ("embeds") cells of each tissue sample in a common latent space
based on their locations as well as gene expression profiles. Dimensions of the learnt latent space
define a natural coordinate system shared by the samples, assigning similar coordinates to
analogous tissue structures. In step 2 (top), each gene G is analyzed for differential spatial
expression (DSE) between the tissue samples: spline regression is used to model the gene's
expression in a cell as a function of the cell's coordinate along a particular dimension, and a
statistical test is performed to assess whether this function differs between the samples, with the
results quantified as a false discovery rate (FDR). (This is repeated for each dimension.) The
same approach also applies to detecting a gene's DSE between two groups of samples. **(d)**
Schematic of step 1 of SpaceExpress. Spatial information in given tissue sample(s) is represented
by a nearest neighbor graph of cells, with distances between cells (e.g., d_{ij} between cells i and
j) defined as shortest path lengths between them and reflecting their physical proximity in the
tissue. Expression profile of each cell (g_i) is mapped by an encoder (neural network) to a low-
dimensional embedding, so that Euclidean distances between cells in the embedding space (e.g.,
$p_i - p_j$ V) approximate their spatial distances (e.g., d_{ij}). For simplicity, three dimensions are
shown in the embedding space.

Figure 2. SpaceExpress learns a shared coordinate system for multi-sample comparative
analysis. (a) Schematic of synthetic "early Drosophila embryo" data set, comprising two "tissue
samples", each with 3,915 cells arranged in an ellipse and gene expression patterns varying along
the anterior–posterior (AP) and dorso–ventral (DV) axes; a subset (20%) of genes exhibits inter-
sample differences and parameter μ controls DSE strength of these genes. **(b)** Inter-sample
agreement (left, assesses if the learnt coordinate system is "shared") and natural coordinate
fidelity (right, evaluates if the learnt coordinate preserves the original AP axis) for SpaceExpress,
STAGATE and GraphST across varying μ. Red dashed line indicates the maximum score of 1.

(c) PCA visualization of learnt coordinate system for data set at μ = 0.6. For each method evaluated, left panel shows cells colored by sample identity and right panel shows cells colored by ground-truth AP coordinate. **(d)** Evaluation of robustness to spatial distortions: rotation (30°), scaling (1.5×), partial truncation (20% removal) and their combinations. Bar plots show inter-sample agreement (left) and natural coordinate fidelity (right). Means across 10 replicates of synthetic data sets are shown, error bars indicate standard deviation. **(e-j)** Evaluations of shared coordinate system learnt for each of three real spatial transcriptomics datasets across diverse technologies and biological contexts: MERFISH profiles of normal-reared (NR) and dark-reared (DR) mouse visual cortex (e,f), Visium profiles of human DLPFC samples from two different individuals (g,h), and Stereo-seq profiles of adjacent slices of a Drosophila late-stage embryo (14-16h after egg laying) (i,j). Panels e,g,I (bottom) show agreement metrics – Inter-sample agreement (left) and natural coordinate fidelity (right) – for each method. Agreement score captures the extent to which learnt coordinate system preserves biologically meaningful correspondences between samples – based on cortical depth in the mouse visual cortex, and spatial domains in the DLPFC and Drosophila datasets. Panels f,h,j show PCA visualization of learnt coordinate systems. In each case, plot on left shows cells colored by sample identity and plot on right shows cells colored by ground-truth cortical depth / region annotation. In all cases, SpaceExpress achieves the highest inter-sample agreement and natural coordinate fidelity, demonstrating strong generalizability across platforms and tissue types.

Figure 3. Statistical inference of differential spatial expression (DSE) by SpaceExpress. (a-c) Schematic overview of DSE testing, illustrated with the synthetic early embryo data set. Cells from two synthetic tissue samples are embedded in a shared coordinate system (a) by SpaceExpress, and each gene is tested for sample-specific variation along each dimension. Panels (b) show an example gene whose expression pattern differs between Tissue 1 and Tissue 2 (top). Part of this difference manifests in the gene's expression variation along the AP axis, which is captured by SpaceExpress-learned dimension 1 (bottom). Panel c shows this difference in detail, with the gene's expression in each cell (of either tissue) plotted against the cell's coordinate value in dimension 1. The tissue-specific trends of this variation are visualized by the bold curves, which represent tissue-specific components of a spline regression model fit to all cells. The marked difference between these trends results in the gene being assigned an FDR ~0 for dimension 1; all the gene-dimension combinations are similarly tested and assigned FDR values (panel d). Finally, all significant genes are reported for each dimension. **(e)** False positive rate at varying levels of signal strength (μ), showing performance of the SpaceExpress-DSE pipeline compared to ablated versions that lack the second retraining step, utilize fewer knots in spline model (n = 10 compared to n = 300), or lack the empirical null fitting step. **(f)** True positive rate and false positive rate for DSE detection across increasing values of μ, demonstrating robust sensitivity even to small signal strengths (μ). **(g)** FDR values assigned by SpaceExpress to DSE (n = 20) and non-DSE (n = 80) genes for each of four learned dimensions, indicating clear separation of significant hits by SpaceExpress. The red lines mark significance thresholds DSE analysis (1e−3). Results are shown for data set at μ = 0.6. **(h)** Comparison of DSE and traditional DE results, showing SpaceExpress-assigned FDR and Wilcoxon p-value respectively. (Results are for μ = 0.6.) The data set includes no DE genes, by design; consequently, all genes show

nonsignificant p-values in DE analysis. In contrast, in DSE analysis using SpaceExpress with a
specific dimension, all 20 DSE genes are correctly detected without false positives. The red lines
mark significance thresholds for DE (p-value = 0.05) and DSE (FDR = 1e−3) analyses. **(i,j)**
Evaluation of robustness of DSE identified by SpaceExpress in MERFISH profiles of mouse visual
cortex under normal and dark-rearing conditions (i) and mouse hypothalamic preoptic region from
naive and pup-exposed animals (j). In each case, four inter-group two-sample comparisons (bold
arrows) and two intra-group comparisons (dashed arrows) were performed. DSE genes were
detected (with statistical significance scores) for each comparison, using SpaceExpress or an
alignment-based method. Bar plots show numbers of DSE genes detected (out of 500 genes
tested) in the four inter-group comparisons, at a significance level stronger than any seen in the
intra-group comparisons by the same method. Each bar also shows the number of genes
designated DSE in three or all of the four inter-group comparisons; see Methods for full details.

1063

Figure 4. **SpaceExpress identifies differential spatial expression of genes in honey bee**
brain. (a) Transcriptome map for forager and soldier honey bee brains. Each spot represents a
transcript (n= 11,373 and 10,334 respectively) with color representing gene identity. **(b)**
Visualization of SpaceExpress cell embeddings for the two honey bee brain samples, in all four
dimensions. Each point corresponds to an individual cell at its spatial location, colored by the
embedding value for each dimension. **(c)** Venn diagram showing the overlaps among sets of
genes exhibiting Differential Spatial Expression (DSE) in each of the four embedding dimensions
(Dim 1-4). **(d, f, h)** Spatial expression visualization of select DSE genes. The expression levels
of *hr38* (d), *fasciclin-1* (f), and *per* (h) are shown. Cells with higher expression levels of each gene
are colored in darker shades of red, revealing distinct patterns of spatial expression in different
brain regions. **(e, g, i)** Gene expression versus embedding dimension plots. These plots display
spline model fits for gene expression levels of *hr38* (e), *fasciclin-1* (g), and *per* (i) against the
embedding dimension in which DSE was detected as significant. See text for how to interpret
DSE based on gene-specific visualizations (d-i) and embedding dimension visualizations (b).

1078

Figure 5. SpaceExpress analysis reveals vision-dependent spatial gene expression
changes in the mouse visual cortex. (a) Schematic of experimental design. Primary visual
cortex (V1) samples were collected from normally reared (NR) and dark-reared (DR) mice at
postnatal day 28, including anterior and posterior slices (~550 μm apart). Each group includes
two biological replicates (NRa, NRb, DRa, DRb). **(b, c)** Visualization of neuronal subclass
annotation and SpaceExpress dimensions across (b) posterior and (c) anterior V1 samples. Top:
neuronal subclass annotation overlaid on the tissue sections, with arrows indicating the direction
of cortical depth. Bottom: Smoothed heatmaps generated using kernel density estimation,
displaying the learned natural coordinate values as continuous color gradients for each dimension.
(d) Venn diagram showing overlap of differentially spatially expressed (DSE) genes in anterior
and posterior analyses. Both regions yielded 26 DSE genes, with 14 shared genes, 9 of which
are immediate early genes (IEGs). **(e)** Normalized expression of the 9 shared IEGs plotted as a
function of cortical depth, averaged across replicates. DR samples show elevated expression
between 100–500 μm. **(f)** Beeswarm plot showing the FDR for each gene in posterior and anterior

analyses. FDR for each gene is calculated by the minimum FDR values across all dimensions.
IEGs are highlighted in orange. **(g)** Venn diagrams showing the number of DSE genes identified
per dimension in posterior (left) and anterior (right) analyses. **(h-k)** Differential spatial expression
of (h, i) Nr4a3 and (j, k) Fosb, two immediate early genes identified as DSE in both posterior and
anterior analyses. Each panel shows, from left to right, spline model fits and spatial expression
heatmaps for posterior or anterior samples. Spatial expression heatmaps are smoothed across
replicates using kernel density estimation.

1100

Figure 6. SpaceExpress identifies differential spatial expression of genes in mouse
hypothalamic preoptic **region related to pup-directed aggression. (a)** Neuroanatomical
annotation of cells in the hypothalamic preoptic region for naïve and a pup-exposed aggressive
mouse, obtained computationally by mapping the cells to a reference tissue sample with manual
annotations (Methods). Cells are color-coded according to their respective regional annotations,
including the third ventricle (V3), bed nuclei of the stria terminalis (BST), columns of the fornix
(fx), medial preoptic area (MPA), medial preoptic nucleus (MPN), periventricular hypothalamic
nucleus (PV), paraventricular hypothalamic nucleus (PVH), and paraventricular nucleus of the
thalamus (PVT). **(b)** Venn diagram illustrating the overlaps among sets of DSE genes across the
four embedding dimensions (Dim 1 to 4). **(c)** SpaceExpress embeddings visualized across four
dimensions for both the naïve and pup-exposed mouse brains. Each cell is represented by a
point, positioned according to its spatial location and colored by the corresponding embedding
value for each dimension. **(d, e)** Spatial expression visualization and gene expression versus
embedding dimension plots for select genes.   Shown are the expression levels (grey-red
colormaps) of *Th* (d) and *Slc17a6* (e) in both naïve and pup-exposed samples, alongside spline
model fits for gene expression along the respective embedding dimensions where significant DSE
was detected.

1118

1119

a Differentially Spatially Expressed Gene (DSE)

b

Honeybee Whole Brain (Our Data) — Soldier, Forager

Mouse Visual Cortex — Normal, Dark-rearing

Mouse Hypothalamic Preoptic Region — Normal, Pup-exposed

c Schematic of SpaceExpress

Learn Natural Coordinate System

Examine expression variation along each learnt dimension

Model Expression as Function of Dimension

Test of Difference; FDR = 0.001

Gene expression

d Learning Natural Coordinate System

X-Y coordinates — Nearest Neighbor Graph — Neural Network — Natural coordinates