<u>Title</u>: **Modeling Insights into Potential Mechanisms of Opioid-Induced Respiratory Depression**

within Medullary and Pontine Networks

<u>Authors</u>: Wendy L. Olsen[1,3*], John A. Hayes[1,2*], Dale Shuman[4], Kendall F. Morris[4], and Donald C.

Bolser[1,2]

<u>Author affiliations</u>:

[1]Breathing Research and Therapeutics (BREATHE) Center, University of Florida, Gainesville, FL

[2]Department of Physiological Sciences, University of Florida, Gainesville, FL

[3]Department of Rehabilitation Sciences, Appalachian State University, Boone NC

[4]Department of Molecular Pharmacology and Physiology, University of South Florida, Tampa, FL

* contributed equally

<u>ORCIDs</u>:

Wendy L. Olsen ORCID: 0000-0002-7607-8096

John A. Hayes ORCID: 0000-0003-2849-7672

Kendall F. Morris ORCID: 0000-0001-6582-1203

Donald C. Bolser ORCID: 0000-0002-4577-8954

<u>Corresponding author</u>: olsenwl@appstate.edu

Keywords: OIRD**,** opioids, glutamate release, fentanyl, mu-opioid receptors, mu-OR, µ-OR, modeling,

preBötzinger Complex, pons, medulla

<u>Funding</u>: Supported by NIH 1R01HL155721-01, 1R01HL163008, and T32HL134621.

<u>Table of Contents Category</u>: Computational Physiology and Modelling

<u>Author contributions</u>: Conception or design of the work: W.L.O. and D.C.B. Acquisition, analysis or

interpretation of data for the work: W.L.O. and J.A.H. Drafting the work or revising it critically for

important intellectual content: W.L.O., J.A.H., D.S., K.F.M., and D.C.B. All authors have read and

approved the final version of this manuscript and agree to be accountable for all aspects of the work in

ensuring that questions related to the accuracy or integrity of any part of the work are appropriately

investigated and resolved.

<u>Running title</u>: Mechanisms of Mammalian Opioid-Induced Respiratory Depression

ABSTRACT (246 of 250 words)

The opioid epidemic is a pervasive health issue and continues to have a drastic impact on the United

States. This is primarily because opioids cause respiratory suppression and the leading cause of death

in opioid overdose is respiratory failure (*i.e.*, opioid-induced respiratory depression, OIRD). Opioid

administration can affect the frequency and magnitude of inspiratory motor drive by activating μ-opioid

receptors that are located throughout the respiratory control network in the brainstem. This can

significantly affect ventilation and blunt CO_2 responsiveness, but the precise neural mechanisms that

suppress breathing are not fully understood. Previous research has suggested that opioids affect

medullary and pontine inspiratory neuron activity by disrupting upstream elements within this circuit.

Inspiratory neurons within this network exhibit synchrony consistent with shared excitation from other

neuron populations and recurrent mechanisms. One possible target for opioid suppression of

inspiratory drive are excitatory synapses. Reduced excitability of these synaptic elements may result in

disfacilitation and reduced synchrony among inspiratory neurons. Downstream effects of disfacilitation

may result in abnormal output from phrenic motoneurons resulting in distressed breathing. We tested

the plausibility of this hypothesis with a computational model of the respiratory network by targeting the

synaptic excitability in fictive medullary and pontine populations. The synaptic conductances were

systematically decreased while monitoring the overall respiratory motor pattern and aggregate firing

rates of subsets of cell populations. Simulations suggest that highly selective, rather than generalized,

actions of opioids on synapses within the inspiratory network may account for different observed

breathing mechanics.

INTRODUCTION

Opioids are considered the gold standard for the treatment of chronic and acute pain, when used appropriately. This is especially true in a perioperative and postoperative environment (Hill and Canals 2022; Palkovic et al. 2020). The risks associated with opioid use follow a dose-dependent response and has ensured that their availability remain restricted (Bateman et al. 2021; Hill and Canals 2022; Mattson et al. 2021). However, the opioid epidemic remains a clear and increasingly pervasive health crisis in the United States (Ramirez et al. 2021). In 2019, overdose deaths increased by more than 50.6% since 2013 (Mattson et al. 2021). This increase is largely attributed to the illicit distribution and use of fentanyl. The primary cause of death in opioid overdose is respiratory depression (Hill and Canals 2022; Ramirez et al. 2021). Thus, research has been dedicated to uncovering the mechanisms associated with opioid-induced respiratory-depression (OIRD) and uncovering neural pathways to reverse OIRD (Bateman et al. 2021; Ramirez et al. 2021).

Breathing

Multiple regions within the brain and brainstem can contribute to OIRD (Bateman et al. 2021; Lalley 2003; Ramirez et al. 2021). The brainstem is the primary hub that holds afferent and efferent neurons that drive breathing and breathing-related behaviors. These neurons work together to form connections that regulate respiration in mammals (Lindsey et al. 2012). These circuits control the drive to breathe, cardiorespiratory functioning, and maintain ventilation (Segers et al. 2015). Rhythmic respiratory activity is facilitated and modified by interactions between the bilaterally distributed respiratory control network located within the ventrolateral medulla (Baekey et al. 2004) and certain regions within the pons (Levitt et al. 2015; Varga et al. 2019). This interconnectedness of medullary and pontine neurons is essential for breathing (Lindsey et al. 2012).

The preBötzinger complex (preBötC), located within the medulla, has been reported to generate a coupled respiratory rhythm with the parafacial/retrotrapezoid regions (Bochorishvili et al. 2012; Mellen et al. 2003; Pattinson 2008). The parafacial/retrotrapezoid nuclei are involved in active expiration and central chemoreception. The Bötzinger complex maintains neurons that are active during expiration and is a major source of inhibition (Bateman et al. 2021; Varga et al. 2019), and these processes contribute to pacing respiration (Bateman et al. 2021). Inspiratory bursts generated by the preBötC network determine breathing frequency (Burgraff et al. 2021). Neurons experience a gradual increase in excitation that synchronizes to produce a coordinated population burst (Baertsch et al. 2019). Synaptic connections within the medulla coordinate these population bursts via excitatory and inhibitory networks to facilitate rhythmogenesis (Baertsch et al. 2019; Burgraff et al. 2021; Ramirez et al. 2021). Studies perturbing the preBötC with opioids, in particular, have reported apnea (Wenninger et al. 2004),

respiratory failure (Wenninger et al. 2004), and deficits in airway protection (Shen et al. 2022). For

example, in a series of studies conducted by Lalley and colleagues oscillation and burst patterns of

respiratory rhythm are slowed and decreased by doses of fentanyl (Lalley 2003; Lalley and Mifflin

2017).

The ventrolateral medulla is not the only region participating in respiratory rhythm generation.

Areas of the rostral pontine circuitry, that include the Kölliker-Fuse nucleus, the parabrachial complex,

and the locus coeruleus, also contribute to rhythmogenesis by delivering dense excitatory feedback to

the respiratory control network in the ventrolateral medulla (Pattinson 2008), stabilizing the respiratory

rate, and generating eupneic breathing patterns (Levitt et al. 2015; Varga et al. 2019). Researchers

found that these specific sites (*i.e.*, the Kölliker-Fuse nucleus, the parabrachial complex, and the locus

coeruleus) contribute to the generation of the three-phase normal, eupneic respiratory pattern:

inspiration, post-inspiration, and active expiration (Abdala et al. 2009). Additionally, the medulla

receives sensory information from the pontine network that affects respiration. Specifically, the nucleus

tractus solitarius receives afferent feedback from peripheral sites (*i.e.*, pulmonary stretch receptors,

bronchopulmonary C-fibers, and chemosensory information from carotid and aortic bodies) (Bateman et

al. 2021). Studies perturbing the pontine nuclei resulted in respiratory rate alterations (Bonis et al.

2010), ventilatory changes (Abdala et al. 2009), and tidal volume changes (Levitt et al. 2015).

Contributions from the pons and medullary areas form a spatially dynamic network that produces and

controls respiratory rhythm (Abdala et al. 2009; Baertsch et al. 2019; Segers et al. 2015).

Opioids and the Respiratory Control Network

Opioids alter breathing by activating μ-opioid receptors (μ-ORs) throughout the brainstem (Dahan et al.

2010). Pharmacological effects and adverse events are mediated by the presence of μ-ORs that are

found at multiple sites within the central and peripheral nervous system (Varga et al. 2019). In the

ventrolateral medulla, the preBötzinger complex is active during inspiration and has been found to be

extremely sensitive to opioid agonists (Pattinson 2008). Previous research has studied these effects

within intact animals and medullary slices. Results indicated that opioids have presynaptic and

postsynaptic effects that alter the excitability of brainstem respiratory neurons (Gray et al. 1999, 2001).

Lalley further investigated the effects of systemic administration of fentanyl by measuring intracellular

membrane potentials of respiratory bulbospinal, vagal, and propriobulbar neurons in anesthetized and

unanesthetized decerebrate cats (Lalley 2003). Lalley concluded that fentanyl had effects presynaptic

to respiratory motoneurons to depress neuronal activity. Additional rodent studies observed discrete,

rather than continual, stepwise depression in phrenic output and inspiratory neuron discharges by

opioids. These results are attributed to the effects on circuits upstream to inspiratory neurons within the

preBötzinger complex (Janczewski and Feldman 2006; Mellen et al. 2003). For example, applying the

opioid agonist DAMGO to the Kölliker-Fuse nucleus causes robust apneusis in a working heart–

brainstem preparation of rat (Levitt et al. 2015). However, how opioid agonists fully affect inspiratory

neurons within the respiratory control network remains unclear. What has been implicated across

several studies is that ventrolateral medullary and pontine circuitry, together, are affected by opioid

administration and this in turn affects respiration (Burgraff et al. 2021; Dahan et al. 2010; Janczewski

and Feldman 2006; Lalley 2003; Mellen et al. 2003).

The most common adverse effect of opioid use reported is respiratory depression, which in the

extreme can directly lead to death (Pattinson 2008). Previous researchers have used intracellular

recordings to measure membrane potentials of inspiratory neurons to better understand the inhibitory

effects within the respiratory control network (Gray et al. 1999, 2001; Lalley 2003; Lalley and Mifflin

2017; Mellen et al. 2003). Local application of opioid agonists affects the somatodendritic μ-ORs on

spatially confined presynaptic terminals while receptors in the broader region are left unaffected. This

phenomenon can be difficult to interpret when the pontine and medullary circuitry, specifically the

preBötzinger complex and the Kölliker-Fuse nucleus, reciprocally share sensory-motor information to

generate inspiratory bursts and respiratory patterns (Varga et al. 2019). Recently, Chou and coworkers

(Chou et al. 2024) disseminated findings from a computational model that described plausible

explanations for the observed variations in experimental responses to opioids. The group explains that

their model accounts for the fixed and dynamic excitatory/inhibitory μ-OR+ neurons, cellular

parameters, and network connections. They attribute discrete assigned randomness to these

parameters within the model that influence individual nodes. This small level of difference is sufficient to

introduce enough variance to explain the variances in experimental preparations. However,

understanding the neural network connections and respiratory mechanisms that are affected by opioids

is of widespread interest to researchers.

A joint neural-biomechanical computational model

Lalley and coworkers (Lalley 2003; Lalley and Mifflin 2017) have interpreted their results of suppressed

breathing and disfacilitation of discharge patterns as an attenuation of presynaptic excitability within the

pontomedullary circuitry. We have tested the plausibility of this hypothesis with a joint-neuromechanical

model of the respiratory network (Lindsey et al. 2012; O'Connor et al. 2012). This model uses an

integrate-and-fire neural network that drives deterministic equations that simulate human respiratory

mechanics (O'Connor et al. 2012). The first aim of the current study was to systematically and

individually decrease the strength of medullary inspiratory neuron connections within this joint neural-

biomechanical model to examine the overall respiratory output. The second aim was to systematically,

and individually, decrease the strength of pontine neuron connections within the same model. Model

and trial specifications are described in the Materials and Methods section.

MATERIALS AND METHODS

Network simulations

A joint neural biomechanical model (Lindsey et al. 2012; O'Connor et al. 2012), was applied to test the

hypothesis that decreasing the synaptic conductance of medullary and pontine inspiratory neurons

induces opioid-mediated respiratory breathing patterns. The neural components of the model were

derived from previously described respiratory network models of discrete integrate-and-fire neurons

after MacGregor (MacGregor 1987) and a hybridized bursting integrate-and-fire population based on

Hodgkin-Huxley equations (Breen et al. 2003). The latter was previously developed from a continuously

integrated model (Butera et al. 1999). The current model was derived from these equations paired with

in vivo data that enhanced the development. Ultimately, motor output from phrenic populations at each

time step (0.5 ms) within the model are calculated by counting the total spikes from all the cells in each

of two phrenic populations ($P0$ and $P1$) and dividing by the duration of a time step. Each population

accounts for different inspiratory burst activity within the medullary component of the model. The output

of the two can best be described by the equation:

174
$$\frac{0.3 \cdot P0 + 0.7 \cdot P1}{200}$$

175 Notably, there should be a maximum value of 1 for phrenic population recruitment. This equation

176 describes the model's representation of inspiratory output. The lumbar motor output is handled in a

177 similar way as the phrenic output with two lumbar motoneuronal populations ($L0$ and $L1$):

178
$$\frac{0.3 \cdot L0 + 0.7 \cdot L1}{80}$$

The biomechanical components of the model (Lindsey et al. 2012; O'Connor et al. 2012) were

developed from transdiaphragmatic pressure and diaphragm activation while controlling the

thoracoabdominal configuration (Cluzel et al. 2000; Grassino et al. 1978; Konno and Mead 1967; Song

et al. 2006).

The *uflsim* software package, version 1.0.36 (Lindsey et al. 2012; O'Connor et al. 2012) utilizes

a *Qt C++* cross-platform development framework written for Windows and Linux (source code may be

found here: https://github.com/jahayes-ns/uflsim with Windows binaries here:

https://github.com/jahayes-ns/uflsim/releases/download/neuroscience/uflsim_win_1.0.36.zip).

The program includes the functionality of the program *SYSTM11* (MacGregor 1987) used in

previous simulations of the respiratory network. The program allows neuron excitability to be modulated

by injected current, and elements designated as "fiber populations" external to the network can also be

used to represent transiently active afferent inputs to the network. A graphical user interface (*simbuild*)

was used to modify cell parameters and network structure while the resulting model files were

simulated using *simrun*. Simulations were run on 64-bit Intel-based computers under the Windows or

Linux operating systems. Python scripts were developed to produce large sets of *uflsim* networks (.snd

files) with varied parameters such as synaptic strengths between populations of neurons, so

simulations could be executed in batches and analyzed offline with figures produced using *Matplotlib*

(Hunter 2007). Network summary figures (Figs. 3D-E, 5, 6B-C, 8B-C, and 9B-C) were produced by

taking the mean±SEM of relevant features from 9 distinct runs of freshly produced networks derived

from a "trunk" network (Fig. 2). These 9 distinct networks were randomly generated with different

random seed values.

Neuron simulations

Single-cell neuron simulations of the Butera model (Butera et al. 1999) were performed using *XPPAUT*

(Ermentrout 2002). Parameter values were taken from the original study except for the changes

described in Results. Panels for the figures were produced using *Matplotlib* (Hunter 2007) and *pandas*

(team 2024).

RESULTS

Organization of a fictive bulbar and spinal respiratory network

We started with the network structure from a previously published respiratory model (O'Connor et al.

2012). This is comprised of 47 populations of neurons with up to 70-300 members each (9159 total

neurons) as well as 10 fiber sources that project to these neuronal populations (1800 total fibers). Each

population member had 50-200 axon terminals randomly distributed to members of its target

populations (average source→target were 100.1 terminals for each pair) with 175,394,850 total

terminals in the network. The identity of the neurons in each population are defined by three

characteristics: 1) their typical firing phenotype under eupneic conditions, 2) their general anatomical

location, and 3) their hypothesized, or experimentally identified, connectivity to other populations in the

model (Fig. 1). Typically, the prefix of the eupneic inspiratory-phasing neurons are "I-" and eupneic

expiratory-phasing neurons with "E-". After this "Aug", "Dec", suggests the predominant discharge

pattern during the respective phase as augmenting or decrementing spike rate consistent with

experimental phenotypes. "NRM" indicates that a population is not eupneic respiratory modulated.

Anatomical locations for the populations are sometimes specified parenthetically with "pons" or "raphe"

and the remainder are by default in the medulla. The exception to the latter is the "PHRENIC" and

"LUMBAR" populations of motoneurons and are meant to represent roughly the C4 and L1 levels of the
spinal cord and output to muscles.

Figure 1A highlights 17 of the key populations from this model network in the context of the
present study with gray boxes generally delineating the approximate anatomical location (pons,
medulla, and spinal cord) for the firing phenotypes. Figure 1B shows hypothesized lateral connections
within these structures, while Figure 1C and 1D show ascending and descending connections between
these structures, respectively.

Figure 2 shows the population activity of each of these populations where the core respiratory
rhythm-generating circuit is shown in Fig. 2A. Medullary interneurons (INs) periodically oscillate
bursting between the inspiratory (I) and expiratory phases (E) with the I-Driver neurons initiating the
cycles, I-Dec neurons following a similar firing pattern, and I-Aug neurons reciprocally inhibiting the
others. These medullary neurons project to bulbospinal premotoneurons that further project to cervical
motoneurons (Phrenic MNs) and lower spinal cord (Lumbar MNs) (Fig. 2B). Our biomechanical model
accounts for the activity from these MNs, as well as laryngeal MNs (not shown), to model airway
mechanics that drive lung inflation/deflation (Fig. 2C). Pulmonary stretch receptors (PSRs) then both
excite I-Aug and inhibit I-Dec neurons during inflation closing a feedback loop (Fig. 2D).

The relationship of I-Driver cellular properties to fictive breathing
The I-Driver neurons form the core kernel of the rhythm generator that produce the initial burst activity
that percolates through the inspiratory phase (Fig. 2A). A sub-spike threshold, slowly-inactivating
"persistent" sodium current (I_{NaP}) produces this augmenting activity during the late-expiratory phase
and is described by the following equation (modified from (Breen et al. 2003)):

250 $$I_{NaP}(V_m) = \bar{g}_{NaP} \cdot m_\infty(V_m) \cdot h_{NaP} \cdot (V_m - E_{Na})$$

Figure 3 illustrates the subthreshold activity of this I_{NaP} current. Fig. 3A shows the membrane voltage
trajectory (V_m) of an intrinsically bursting I-Driver-like neuron with the inactivation variable (h_{NaP}) in the
middle row and the I_{NaP} truncated to -50 pA in the bottom row to highlight the slowly de-inactivating
current between bursts of activity. E_{Na} is the Nernst reversal potential for sodium (+50 mV) while
$m_\infty(V_m)$ is the instantaneous voltage-dependent activation function for I_{NaP}.

Figure 3B shows the same simulated neuron with the maximum synaptic conductance ($\bar{g}_{NaP}$)
slightly decreased to a scaling factor of 95% (0.95x) which slows the bursting frequency. Further
decreasing the scaling factor to 90% (0.9x) resulted in a silent neuron that relaxes to a subthreshold
baseline V_m (not shown). In the same simulated neuron, returning to the original $\bar{g}_{NaP}$ but changing the
maximum time constant of NaP inactivation ($\tau_{h\text{-}NaP}$) to 50% (0.5x) results in a dramatic increase in the
bursting frequency in Fig. 3C.

For comparison to the more expansive network model, similar graded adjustments on 300 I-
Driver neurons from scaling factors 1.0x to 0.0x to $\bar{g}_{NaP}$ and $\tau_{h\text{-}NaP}$ led to changes in T_i (inspiratory
phase duration), T_e (expiratory phase duration), and T_{tot} (the sum of T_i and T_e, or full cycle period) and
is shown in Fig. 3D. Similar to the neuron model, changing $\bar{g}_{NaP}$ led to cessation of rhythm at relatively
high levels of scaling factor for $\bar{g}_{NaP}$ suggesting the importance of subthreshold I_{NaP} in this model to
initiate the population burst activity. Remarkably, in contrast to the single neuron model, decreasing $\tau_{h\text{-}}$
$_{NaP}$ increased both the expiratory phase (inter-burst interval) and consequently the full cycle period (T_{tot})
while having only a modest effect on T_i. This qualitatively shows the dramatic impact network
connectivity and synaptic properties can have on the overall production of rhythmic behavior in this
model brainstem, and we explore this in more detail below.
267
Generalized mechanisms for μ-OR-agonist influence on the neural control of respiration
In this study, we analyzed several distinct schemes by which μ-OR agonists may influence respiratory
activity (Fig. 4). The first are comprised of cellular effects that are conceptualized as affecting baseline
membrane properties through K^+-dominated leak channels and will be examined more closely
associated with Fig. 5. The key take-away from this mechanism is that, in the absence of active
membrane properties more dramatic than spike-generating currents, it would simply affect the
presynaptic spike rates of neurons and can be simulated by an adjustment in applied stimulus current
(I_{app}) (Fig. 4A). In contrast, mechanisms that influence connectivity strength could act through pre- or
postsynaptic mechanisms (Fig. 4B) and will be considered in the subsequent Results sections (Fig. 6-
10). For the purposes of this study, they are effectively the same mechanism and result in decreased
I_{syn} (synaptic current) given a uniform spike-rate between the two.
μ-OR agonists have been shown to directly cause Fig. 4A.iii and Fig. 4B.ii in some contexts
(Gray et al. 1999; Heinke et al. 2011; Ikoma et al. 2007; Jørgensen et al. 2022; Kim et al. 2024) and
Fig. 4B.i may be one mechanism of opioid tolerance (Gillis et al. 2020; Koch and Höllt 2008).
282
Alteration of excitability in populations of the upstream core network
We first started by examining the effects of biasing cellular excitability by altering I_{app} over the range
-10 to +10 pA, where the latter depolarizes neurons. There were 4 conditions, changes in ΔI_{app} on the
populations of: I-Drivers, I-Drivers + I-Augs, NRM-pons, and all neurons in the simulation for
comparison. The results are analogous to the situations demonstrated in Fig. 4Ai-iii.
There were 9 distinct runs of independently generated starting networks (ΔI_{app} = 0) for the 4
conditions. Changing ΔI_{app} > 0 for all neurons slows the respiratory rhythm (T_{tot}) as ΔI_{app} >> 0 but also
slows the rhythm slightly as inspiratory phase bursts (T_i) increase when ΔI_{app} < 0 (Fig. 5A). If ΔI_{app} < 0

falls too low the system loses respiratory activity. As this ceases at ΔI_{app} < ~-2 pA across all networks it
shows that the current system is very "stiff" and just on the precipice of cessation if the whole network is
seriously perturbed in the hyperpolarizing direction.

For I-Driver + I-Aug perturbations, there is a transient period as ΔI_{app} < 0 where the CV of T_{tot}, T_i,
and T_e, increase dramatically compared to similar I-Driver perturbations (Fig. 5A). Curiously, when I-
Driver population alone is manipulated the means of both T_i and T_e roughly track along the same
trajectories as I-Driver + I-Aug perturbations. This shows that the I-Aug population is contributing to
cycle-to-cycle stability of the respiratory rhythm.

We also modulated NRM-pons neurons to see how they influence overall respiratory activity.
While they are non-phasic, the stochasticity of this population's firing still influences activity in non-
intuitive, non-monotonic ways as a function of uniform ΔI_{app} (Fig. 5A). At ΔI_{app} > 0, breathing became
deeper, and lungs are inflated while at ΔI_{app} < 0 breaths are shallower (Fig. 5B). Similar trends were
found in the more targeted perturbations of I-Drivers and I-Drivers + I-Augs suggesting the NRM-pons
neurons are vicariously acting largely through these populations as the connectivity from NRM-pons
implies (Fig. 1D).

Intraplanar medullary synaptic sources affect rhythm generation
The essential elements of the respiratory network are found in the medullary region, so we examined
how modulating the maximal excitatory strength ($\bar{g}_{Excit}$) between planar connections in this structure
could influence activity (Fig. 6).

Fig. 6A illustrates the subset of anatomically (hypothesized) planar medullary connections from
Fig. 1A and 1B and we were focused on the role of excitatory connections. We scaled down the
synaptic strength (analogous to Fig. 4B) between the following populations of neurons: I-Drivers → I-
Drivers, I-Drivers → I-Augs, I-Drivers → I-Decs, I-Drivers → I-Drivers/I-Augs/I-Decs, and I-Augs → I-
Augs (Fig. 6A).

Scaling the strength of recurrent synapses in the I-Driver population (I-Drivers → I-Drivers)
resulted in relatively little change in T_{tot}, T_i, or T_e (Fig. 6B). Further, perturbation of the strength of these
recurrent synapses had little effect on phrenic amplitude, lung volume or peak inspiratory flow (Fig. 6C).

Reducing synaptic strength between the I-Driver and I-Aug populations increased T_{tot} by over
15% and that effect was primarily due to an increase in T_e of over 30% (Fig. 6B). There was little effect
on T_i by this perturbation. Further, there were linear reductions in both phrenic amplitude, lung volume,
and peak inspiratory flow (Fig. 6C). Figure 7A shows the spike-time histogram patterns of setting the
synaptic strength from I-Driver to I-Aug neurons to 0%. The Phrenic MNs lose robust temporal
coherence which explains the reduction in Flow and Lung Volume (compare to Fig. 2).

When synaptic strength between the I-Driver and I-Dec populations was reduced,

rhythmogenesis and inspiratory motor drive failed after a change between 60-80% (Fig. 6B, C). Figs.

7B and C show examples of firing rate records for medullary and bulbospinal neurons as well as

phrenic and abdominal motoneurons during reduction of synaptic strength to 80% (Fig. 7B), and 60%

(Fig. 7C) of control for I-Driver to I-Dec synapses.

Simultaneous reductions in the synaptic strength from I-Drivers to other I-Driver neurons, I-Aug

neurons and I-Dec neurons resulted in what appeared to be a synthesis of all changes induced by

perturbation of excitability for each of the individual populations alone (Fig. 6B, C). As such,

simultaneous reductions in synaptic strength by up to 55% increased T_{tot} and T_e and decreased phrenic

amplitude, lung volume and peak inspiratory flow (Fig. 6B, C). Large reductions in synaptic strength

resulted in simulated apnea.

We additionally decreased synaptic strength among recurrent synapses in the I-Aug population

alone. Unlike perturbation of synaptic strength among recurrent synapses within the I-Driver population;

this action lengthened both T_{tot} and T_i by 15-25% with no change in T_e (Fig. 6B). Further, phrenic

amplitude, lung volume, and peak inspiratory flow were also reduced in a linear manner (Fig. 6C).

Inhibitory influence of I-Dec hub neurons

Since the I-Dec population of neurons seems to have a dramatic effect on respiratory activity, we

decided to also examine the effects of the I-Dec synaptic connections (Fig. 8). This is novel in

comparison to the previous figures in that we were probing the influence of inhibitory synapses.

Perturbing all the inhibitory connections from the I-Dec population (Fig. 8A) causes the

respiratory behavior to drop (Fig. 8B/C) because these neurons are the hub of our system with

connections to 19 of the other 46 populations of neurons (10 inhibitory connections shown). In general,

as the maximal inhibitory strength ($\bar{g}_{Inhib}$) decrease the T_i, T_e, and T_{tot} get shorter and these quantities

get more regular (Fig. 8B), and the phrenic activity monotonically increases (Fig. 8C). When $\bar{g}_{Inhib}$ is

60% of control, Fig.8D demonstrates hyperpnea-like activity with intense inspiratory/expiratory activity

and large lung inflations/deflations before the rhythm goes out in Fig. 8E when $\bar{g}_{Inhib}$ drops to 55% and

lower.

Descending pontine synaptic sources affect burst patterning

Finally, we also looked at how perturbing $\bar{g}_{Excit}$ between NRM and I-Aug or I-Driver neurons affected

the overall breathing pattern. Decreasing the excitatory connections from the pontine synaptic sources

(Fig. 9A) led to disordered and inconsistent T_i, T_e, and $T_{tot.}$ production (Fig 9B). When $\bar{g}_{Excit}$ is blocked,

phrenic activity does increase for the I-Aug + I-Driver and I-Driver connections. However, it decreases

for the I-Aug connections and remains unchanged for the I-pons connections (Fig 9C). Blocking this

descending pontine transmission led to clustered breathing (Fig. 10).

DISCUSSION

The effects of opioids on respiratory function in experimental conditions is variable. In fact, little is

known about the variance of in-between subjects' effects of opioid use (Dunn et al. 2019, 2020). The

current study investigated the simulated responses of an opioid within a computational model of the

pontomedullary respiratory network to better understand the neural mechanisms contributing to OIRD.

The opioid epidemic remains a critical research priority. There are many effects that opioids

pose on respiratory function in experimental and clinical conditions (e.g., decreases or abnormal

function in chest wall compliance, tidal volume, respiratory rate, etc.). This has led to much

investigation that has focused on the impact of different opioid agonists on the upper airway in

respiratory depression (Skulsky et al. 2007). However, respiratory depression is a common deleterious

feature of opioid use. The current study investigated the simulated responses of activating μ-ORs within

a computational model of the pontomedullary respiratory network to better understand the neural

mechanisms contributing to OIRD. Since morphine, codeine, and similar drugs, have multiple side

effects beyond just activating μ-ORs (Dahan 2007; Simera et al. 2010; Tomazini Martins et al. 2018),

our model is best interpreted to most closely reproduce the highly specific μ-OR ligand fentanyl (Lalley

2003; Shen et al. 2022).

Simulated opioid effects on respiratory activity

To model these opioid effects, we decreased the synaptic strength across medullary neurons

associated with rhythmogenesis. Previous computational models have perturbed the connection

probabilities (Chou et al. 2024), time constants of central and peripheral chemoreceptors (Magosso et

al. 2004), and the polarization of sodium channels (Shevtsova et al. 2011) to assess the effects of

opioids on respiration specifically within fictive medullary neurons. The advantage of the current study,

with the employed joint neural network-biomechanical model, is that we examined these factors at a

biomechanical level and in the context of a broader brainstem neuronal network.

Overall, the strength of the network's connectivity is an important parameter that affects the

neural breathing patterns, and the model is generally inhibited when perturbed by opioids. While

decreasing the connection strength or explicitly hyperpolarizing member populations, breathing patterns

were significantly affected. OIRD is characterized by a decrease in respiratory rate and irregular

breathing frequencies and at high doses apnea. This has been attributed to opioids activating G-

coupled proteins through µ-ORs which hyperpolarize cells through G-protein-gated inward rectifying K^+

(GIRK) channels (Montandon et al. 2016a, 2016b).

Previous *in vivo* studies have reported decreased respiratory rates when opioids were directly

applied to the ventrolateral medulla or systemically injected (Montandon et al. 2016b, 2016a). For

example, when DAMGO (d-Ala2, *N*-MePhe4, Gly-ol]-enkephalin), was applied to the ventrolateral

medulla and presumably activating local µ-ORs, it reduced the respiratory rate of mice, but did not

affect the diaphragm amplitude in GIRK2$^{-/-}$ mice (Montandon 2022). In the same study, a moderate

intramuscular injection of fentanyl was provided to the GIRK2$^{-/-}$ mice, and only a slight depression in

diaphragm amplitude was observed. In a complementary study, systemic administration of fentanyl

reportedly decreased respiratory rate, yet had no effect on diaphragm amplitude (Montandon et al.

2016a, 2016b).

The administration of opioids has been shown to affect the burst duration of respiratory motor

units up to the point of respiratory arrest (Lalley 2006). As discussed above, one mechanism opioids

likely perturb respiratory patterns is through cell hyperpolarization, which in turn, affects the spiking

activity of respiratory neurons (Fig. 4A). In our simulations, modulating an injected bias current (I_{app})

within the core medullary populations (Fig. 5) affected spike burst durations (T_i) until the respiratory

activity was extinguished at larger hyperpolarizing I_{app}. Therefore, within our modeling efforts, modifying

the injected current qualitatively reproduces *in vivo* effects of OIRD.

Lalley (2003) investigated the intravenous effects of fentanyl in adult cats while recording

individual neurons. He reported prolonged discharges that induced tonic firing of bulbospinal expiratory

neurons (like our E-Aug-BS population) that were correlated with a reduced hyperpolarization of

synaptic drive potentials. Lalley suggested that this result may have been explained by the decrease in

the duration of the inspiratory phase observed at certain dose-responses of fentanyl (Lalley 2003). He

further interpreted lower doses of fentanyl to have a similar effect on vagal post-inspiratory

motoneurons which led to "sparse, low-frequency" discharges which suggests that fentanyl regulates

bulbospinal and motoneurons presynaptically at different dose-dependent responses (Lalley 2003;

Lalley and Mifflin 2017) similar to a combination of Fig. 4Aiii and Fig. 4Bii.

Fictive respiratory rhythm generation and patterning

An important consideration for this model is that the I-Driver population are fundamentally essential for

any kind of respiratory patterning under our simulated conditions. All members of that population burst

based on a slowly-inactivating persistent sodium current (I_{NaP}) (Breen et al. 2003; Butera et al. 1999)

which has been recently shown to be inessential for I-Driver-like activity (da Silva et al. 2023). For this

study, there is no salient difference in what bursting mechanism we choose for the I-Driver population
as we are interested in network effects as emphasized by the depiction in Fig. 3.

However, considering our model when $\bar{g}_{NaP}$ was altered, this led to the cessation of the network
rhythm (Fig. 3D). Essentially by decreasing I_{NaP} the cells broadly hyperpolarize, and the network
aborted the respiratory rhythm. However, as we showed in Figure 3, modulating the parameters
determining the qualities of $\bar{g}_{NaP}$ on individual I-Driver neurons has a dichotomous effect versus how
the more expansive multi-population network behaves.

This is consistent with previous findings: Phillips and Baertsch reported when lowering [K$^+$]
within their model (which hyperpolarizes neurons) it mimicked *in vitro* experiments by increasing the
presence of weaker "burstlet" rhythms (Phillips and Baertsch 2024). This is hypothesized to be driven
by preinspiratory spiking as opposed to intrinsic bursting within the network. Further, Baertsch and
coworkers (Baertsch et al. 2021) earlier reported that ventrolateral medullary excitatory postsynaptic
potential amplitudes were reduced in the presence of opioids, specifically by DAMGO, and that μ-OR
activation suppresses excitatory synapses from the preBötC neurons to their post-synaptic targets.
Thus, reducing excitability within the respiratory network (Del Negro et al. 2005).

Respiratory efforts and breathing are finely tuned and well-timed behaviors. They must be
executed with precision to prevent maladaptive consequences. These behaviors are primarily regulated
by the nucleus tractus solitarius and the ventral respiratory column. The presence of opioids has been
reported to adversely affect the coordination and execution of these crucial behaviors. The speed of the
network was altered when we manipulated $\tau_{h\text{-}NaP}$ in the I-Driver population which also affected burst
duration. Similar results have been reported by Baertsch and colleagues (Baertsch et al. 2021). These
motor unit spiking behaviors can be divided into three phases: refractory, percolation, and burst
phases. Researchers have reported that when fentanyl is administered, a neuron's discharge identity is
altered, causing changes in timing and frequency (Baertsch et al. 2021; Lalley and Mifflin 2017).
Changes in frequency and neuron identities are modulated by opioid membrane activity. Opioid activity,
within the ventral respiratory column, causes asynchronous activity, and, depending on the opioid
agonist, can cause varying effects on motoneurons. For example, Baertsch and Ramirez (Baertsch et
al. 2019) reported that substance P affected the percolation phase, also referred to as the recurrent
excitation phase, of the inspiratory rhythm by increasing the pre-inspiratory neurons' rate of firing by
activating a depolarizing current (Hayes and Del Negro 2007). Researchers have reported that when
fentanyl is administered, a neuron's discharge identity is altered, causing changes in timing and
frequency. Lalley and Mifflin (Lalley and Mifflin 2017) postulated that μ-opioid agonists directly affect
the controlling and timing of burst and oscillation patterns of bulbospinal and vagal motoneurons, which
also have a direct effect on respiratory muscle force. Specifically, they reported that a threshold dose (3

μg/kg) of intravenous fentanyl had an increase in burst frequencies, oscillation intensities, and a more

negative action potential threshold that had a direct effect on overall breathing pattern.

Physiological implications

One of the motivations for this kind of modeling study is that experimental studies of this kind are

currently unfeasible. Here, we are delineating neuronal populations of interest both anatomically and,

more importantly, functionally, because the respiratory network is distributed across much of the

brainstem (Segers et al. 2008). With advancing genetic technologies these avenues may be more

closely explored but challenges remain and may require higher-order intersectional approaches than

what is currently common.

In the case of anatomical specificity, approaches such as viral injections into specific locations

can partially address these issues (Liu et al. 2021; Varga et al. 2020). Especially, when expression from

these injections may be conditioned on specific gene promoters (Nectow and Nestler 2020). However,

specifying genetic tools to firing pattern is especially nebulous for the most part. A combination of ion

channel expression, endogenous Ca^{2+} buffers such as parvalbumin (Alheid et al. 2002), synaptic

partners, or constitutively expressed transcription factors (Bachmutsky et al. 2020; Sun et al. 2019),

may provide a means of intersectionally subdividing certain populations given a certain neuronal

population's "fingerprint" of multiple distinguishing criteria but that remains beyond the scope of the

present study.

It is important to note that several μ-opioid agonists have been investigated and reported

varying effects (Burgraff et al. 2023; Danaf et al. 2023; Hiranita et al. 2024; Neumueller et al. 2023). For

example, Burgraff, Baertsch, and Ramirez (2023) reported the effects of morphine and fentanyl on

breathing and airway stability. Specifically, the administration of fentanyl resulted in the disruption of

airflow that were resolved with a tracheostomy and/or the administration of salbutamol and adrenaline.

These airflow obstructions were absent during the administration of morphine (Burgraff et al. 2023).

Similar effects have been reported in emergency departments and within other hospital settings (e.g.,

perioperative and postoperative conditions) (Sutter et al. 2015; Wolf et al. 2020).

Opioids are clinically used for their analgesic effects perioperatively and postoperatively.

However, their use can lead to respiratory depression and the disfacilitation of airway protective

mechanisms. The early detection of respiratory suppression allows clinicians to make life-saving

decisions and avert the catastrophic consequences of OIRD. The overall results of our modeling efforts

indicate that the joint neural-biomechanical model employed in the current study demonstrated overall

inhibition, frequency alterations, spike burst changes, and timing changes, which are broadly supported

by the different perturbations observed in *in vivo* data that employs the μ-OR agonist fentanyl. The

proposed model is an excellent tool that lends itself to answering questions that persist within the opioid

crisis, specifically revolving around OIRD.

FIGURE LEGENDS

Figure 1. Connectivity among key elements of the model network. A, The core neuronal

populations interrogated in this study. Orange inverted arrow connections represent excitatory

connections while blue solid circle connections represent inhibitory connections. The graph is arranged

roughly corresponding to hypothesized anatomical location with top-tier rostral pontine populations,

middle-tier medullary populations, and spinal cord populations representing the motoneuronal output of

the neuronal network at the bottom tier. **B-D**, the planar (**B**), ascending (**C**), and descending (**D**)

subsets of these connections to improve clarity.

Figure 2. Fictive eupnea with active expiration. A, The core respiratory time course of activity by

classes of overlayed neuronal populations. The top are pontine interneurons (Pons INs), middle

medullary interneurons (Medulla INs), and bottom medullary bulbospinal premotor neurons

(Bulbospinal INs). **B**, Inspiratory motor output of the simulation as expressed as phrenic motoneuronal

activity (Phrenic MNs) while lumbar spinal motoneurons (Lumbar MNs) convey expiratory activity. **C**,

Simulated lung volume and flow at the mouth produced by the respiratory activity. **D**, Moving average of

lung pulmonary stretch receptors (Lung PSRs) activated by lung expansion. This vagal sensory

information feeds back into the core respiratory circuit continuously. Arrows indicate the feedforward

flow of information in the system.

Figure 3. Comparison of I_{NaP} at the neuronal and network level. A, top, In a simple model of a

single respiratory neuron, I_{NaP} can lead to periodic bursting in membrane potential (V_m). (middle), The

magnitude and kinetics of I_{NaP} are controlled by the inactivation gating variable h_{NaP} (bottom). **B**,

Decreasing $\bar{g}_{NaP}$ slows the burst frequency. **C**, Decreasing the rate h_{NaP} inactivates ($\tau_{h\text{-}NaP}$) increases

the burst frequency. **D**, Effects on the full network period (T_{tot}), burst duration (T_i), and expiratory phase

(T_e), when decreasing $\bar{g}_{NaP}$ and $\tau_{h\text{-}NaP}$ in the I-Driver population of neurons. CV is the coefficient of

variation. **E**, The same effects on motoneuronal output and biomechanical respiratory flow.

Figure 4. High-level comparison of possible cellular and synaptic effects of μOR-agonists.

Presynaptic neuronal (left spheres) spiking with neurotransmitter release (small circles) onto

postsynaptic neurons (right spheres). Clear boxes on the postsynaptic neurons represent transmitter

receptors. **A**, i.), Normal presynaptic spiking activity is equivalent to normal synaptic strength (1.0x $\bar{g}_{syn}$)

and ΔI_{app} = 0. ii.), Heightened presynaptic spiking activity is equivalent to normal synaptic strength and

ΔI_{app} > 0. iii.), Lower presynaptic spiking activity is equivalent to normal synaptic strength and ΔI_{app} < 0.

B, i.), Normal presynaptic spiking activity with normal neurotransmitter release but fewer postsynaptic

receptor targets is equivalent to ΔI_{app} = 0 and <1.0x $\bar{g}_{syn}$ but weaker synaptic strength from a

"postsynaptic effect". ii.), Normal presynaptic spiking activity with decrease in neurotransmitter release

but normal receptor targets is also equivalent to ΔI_{app} = 0 and <1.0x $\bar{g}_{syn}$ but weaker synaptic strength

from a "presynaptic effect".

Figure 5. Influence of ΔI_{app} on network and biomechanical behaviors. A, T_{tot} is total respiratory

period, T_i, inspiratory phase duration, and T_e, the expiratory phase duration. CV is the coefficient of

variation for the respective quantities. **B**, (top) Maximum phrenic motoneuron amplitude during the

inspiratory phase. (middle) Maximum lung volume during the inspiratory phase. (bottom) Inspiratory

flow. ΔI_{app} is in units of pA.

Figure 6. Influence of excitatory synaptic strength on network and biomechanical behaviors

through medullary planar connections. A, Illustration of the excitatory planar connections from Fig.

1B that are analyzed here. **B**, T_{tot} is respiratory period, T_i, inspiratory phase duration, and T_e, the

expiratory phase duration. CV is the coefficient of variation for the respective quantities. **C**, (top)

Maximum phrenic motoneuron amplitude during the inspiratory phase (arbitrary units) (middle)

Maximum lung volume during the inspiratory phase. (bottom) Inspiratory flow.

Figure 7. Example activity patterns of key planar excitatory connection perturbations. A, Spike-

time histogram patterns of the network with I-Driver output to I-Aug neurons at 0% of control. **B**, Spike-

time histogram patterns of the network with I-Driver output to I-Dec neurons at 80% of control

compared to 60% of control (**C**).

Figure 8. Influence of inhibitory synaptic strength on network and biomechanical behaviors

through I-Dec hub connections. A, Illustration of the inhibitory connections from I-Dec neurons in Fig.

1A that are collectively altered here. **B**, T_{tot} is respiratory period, T_i, inspiratory phase duration, and T_e,

the expiratory phase duration. CV is the coefficient of variation for the respective quantities. **C**, (top)

Maximum phrenic motoneuron amplitude during the inspiratory phase (arbitrary units). (middle)

Maximum lung volume during the inspiratory phase. (bottom) Inspiratory flow. **D**, The spike-time

histogram patterns of the network with I-Dec inhibitory output at 60% of control compared to 40% of

control (**E**).

Figure 9. Influence of excitatory synaptic strength on network and biomechanical behaviors

through descending pontine-medullary connections. A, Illustration of the descending excitatory

connections from pontine non-respiratory modulated neurons (NRM-pons), phase spanning neurons

(rIE-pons, cIE-pons, EI-pons), and inspiratory neurons (I-pons). **B**, T_{tot} is respiratory period, T_i,

inspiratory phase duration, and T_e, the expiratory phase duration. CV is the coefficient of variation for

the respective quantities. **C**, (top) Maximum phrenic motoneuron amplitude during the inspiratory phase

(arbitrary units). (middle) Maximum lung volume during the inspiratory phase. (bottom) Inspiratory flow.

Figure 10. Example activity patterns of descending excitatory connection perturbations from

NRM-pons neurons. The network pattern with NRM-pons excitatory output at 20% of control projecting

to I-Aug and I-Driver populations resulting in clustered-like bursts in inspiratory activity.

REFERENCES

Abdala APL, **Rybak IA**, **Smith JC**, **Zoccal DB**, **Machado BH**, **St-John WM**, **Paton JFR**. Multiple pontomedullary mechanisms of respiratory rhythmogenesis. *Respir Physiol Neurobiol* 168: 19–25, 2009.

Alheid GF, **Gray PA**, **Jiang MC**, **Feldman JL**, **McCrimmon DR**. Parvalbumin in respiratory neurons of the ventrolateral medulla of the adult rat. *J Neurocytol* 31: 693–717, 2002.

Bachmutsky I, **Wei XP**, **Kish E**, **Yackle K**. Opioids depress breathing through two small brainstem sites. *eLife* 9: e52694, 2020.

Baekey DM, **Morris KF**, **Nuding SC**, **Segers LS**, **Lindsey BG**, **Shannon R**. Ventrolateral medullary respiratory network participation in the expiration reflex in the cat. *J Appl Physiol* 96: 2057–2072, 2004.

Baertsch NA, **Bush NE**, **Burgraff NJ**, **Ramirez J-M**. Dual mechanisms of opioid-induced respiratory depression in the inspiratory rhythm-generating network. *eLife* 10: e67523, 2021.

Baertsch NA, **Severs LJ**, **Anderson TM**, **Ramirez J-M**. A spatially dynamic network underlies the generation of inspiratory behaviors. *Proc Natl Acad Sci* 116: 7493–7502, 2019.

Bateman JT, **Saunders SE**, **Levitt ES**. Understanding and countering opioid-induced respiratory depression. *Br J Pharmacol* bph.15580, 2021.

Bochorishvili G, **Stornetta RL**, **Coates MB**, **Guyenet PG**. Pre-Bötzinger complex receives glutamatergic innervation from galaninergic and other retrotrapezoid nucleus neurons. *J Comp Neurol* 520: 1047–1061, 2012.

Bonis JM, **Neumueller SE**, **Krause KL**, **Kiner T**, **Smith A**, **Marshall BD**, **Qian B**, **Pan LG**, **Forster HV**. Site-specific effects on respiratory rhythm and pattern of ibotenic acid injections in the pontine respiratory group of goats. *J Appl Physiol* 109: 171–188, 2010.

Breen BJ, **Gerken WC**, **Butera RJ**. Hybrid Integrate-and-Fire Model of a Bursting Neuron. *Neural Comput* 15: 2843–2862, 2003.

Burgraff NJ, **Baertsch NA**, **Ramirez J-M**. A comparative examination of morphine and fentanyl: unravelling the differential impacts on breathing and airway stability. *J Physiol* n/a, 2023.

Burgraff NJ, **Bush NE**, **Ramirez JM**, **Baertsch NA**. Dynamic Rhythmogenic Network States Drive Differential Opioid Responses in the *In Vitro* Respiratory Network. *J Neurosci* 41: 9919–9931, 2021.

Butera RJ, **Rinzel J**, **Smith JC**. Models of respiratory rhythm generation in the pre-Bötzinger complex. I. Bursting pacemaker neurons. *J Neurophysiol* 82: 382–397, 1999.

Chou GM, **Bush NE**, **Phillips RS**, **Baertsch NA**, **Harris KD**. Modeling Effects of Variable preBötzinger Complex Network Topology and Cellular Properties on Opioid-Induced Respiratory Depression and Recovery. *eNeuro* 11, 2024.

Cluzel P, **Similowski T**, **Chartrand-Lefebvre C**, **Zelter M**, **Derenne J-P**, **Grenier PA**. Diaphragm and Chest Wall: Assessment of the Inspiratory Pump with MR Imaging—Preliminary Observations. *Radiology* 215: 574–583, 2000.

Dahan A. Novel data on opioid effect on breathing and analgesia. *Semin Anesth Perioper Med Pain* 26: 58–64, 2007.

Dahan A, **Aarts L**, **Smith TW**. Incidence, Reversal, and Prevention of Opioid-induced Respiratory Depression. *Anesthesiology* 112: 226–238, 2010.

Danaf J, **da Silveira Scarpellini C**, **Montandon G**. βγ G-proteins, but not regulators of G-protein signaling 4, modulate opioid-induced respiratory rate depression. *Front Physiol* 14, 2023.

Del Negro CA, **Morgado-Valle C**, **Hayes JA**, **Mackay DD**, **Pace RW**, **Crowder EA**, **Feldman JL**. Sodium and Calcium Current-Mediated Pacemaker Neurons and Respiratory Rhythm Generation. *J Neurosci* 25: 446–453, 2005.

Dunn KE, **Barrett FS**, **Brands B**, **Marsh DC**, **Bigelow GE**. Individual differences in human opioid abuse potential as observed in a human laboratory study. *Drug Alcohol Depend* 205: 107688, 2019.

Dunn KE, **Weerts EM**, **Huhn AS**, **Schroeder JR**, **Tompkins DA**, **Bigelow GE**, **Strain EC**. Preliminary evidence of different and clinically meaningful opioid withdrawal phenotypes. *Addict Biol* 25: e12680, 2020.

Ermentrout B. *Simulating, Analyzing, and Animating Dynamical Systems: A Guide to XPPAUT for Researchers and*
Students. SIAM, 2002.
Gillis A, **Kliewer A**, **Kelly E**, **Henderson G**, **Christie MJ**, **Schulz S**, **Canals M**. Critical Assessment of G Protein-
Biased Agonism at the μ-Opioid Receptor. *Trends Pharmacol Sci* 41: 947–959, 2020.
Grassino A, **Goldman MD**, **Mead J**, **Sears TA**. Mechanics of the human diaphragm during voluntary contraction:
statics. *J Appl Physiol* 44: 829–839, 1978.
Gray PA, **Janczewski WA**, **Mellen N**, **McCrimmon DR**, **Feldman JL**. Normal breathing requires preBötzinger
complex neurokinin-1 receptor-expressing neurons. *Nat Neurosci* 4: 927–930, 2001.
Gray PA, **Rekling JC**, **Bocchiaro CM**, **Feldman JL**. Modulation of Respiratory Frequency by Peptidergic Input to
Rhythmogenic Neurons in the PreBötzinger Complex. *Science* 286: 1566–1568, 1999.
Hayes JA, **Del Negro CA**. Neurokinin Receptor-Expressing Pre-Bötzinger Complex Neurons in Neonatal Mice
Studied In Vitro. *J Neurophysiol* 97: 4215–4224, 2007.
Heinke B, **Gingl E**, **Sandkühler J**. Multiple Targets of μ-Opioid Receptor-Mediated Presynaptic Inhibition at
Primary Afferent Aδ- and C-Fibers. *J Neurosci* 31: 1313–1322, 2011.
Hill R, **Canals M**. Experimental considerations for the assessment of in vivo and in vitro opioid pharmacology.
Pharmacol Ther 230: 107961, 2022.
Hiranita T, **Ho NP**, **France CP**. Ventilatory Effects of Fentanyl, Heroin, and d-Methamphetamine, Alone and in
Mixtures in Male Rats Breathing Normal Air. *J Pharmacol Exp Ther* 388: 244–256, 2024.
Hunter JD. Matplotlib: A 2D Graphics Environment. *Comput Sci Eng* 9: 90–95, 2007.
Ikoma M, **Kohno T**, **Baba H**. Differential Presynaptic Effects of Opioid Agonists on Aδ- and C-afferent
Glutamatergic Transmission to the Spinal Dorsal Horn. *Anesthesiology* 107: 807–812, 2007.
Janczewski WA, **Feldman JL**. Distinct rhythm generators for inspiration and expiration in the juvenile rat:
Rhythm generators for inspiration and expiration. *J Physiol* 570: 407–420, 2006.
Jørgensen AB, **Rasmussen CM**, **Rekling JC**. μ-Opioid Receptor Activation Reduces Glutamate Release in the
PreBötzinger Complex in Organotypic Slice Cultures. *J Neurosci* 42: 8066–8077, 2022.
Kim HR, **Dey S**, **Sekerkova G**, **Martina M**. μ-Opioid Receptor Modulation of the Glutamatergic/GABAergic
Midbrain Inputs to the Mouse Dorsal Hippocampus. *J Neurosci* 44, 2024.
Koch T, **Höllt V**. Role of receptor internalization in opioid tolerance and dependence. *Pharmacol Ther* 117: 199–
206, 2008.
Konno K, **Mead J**. Measurement of the separate volume changes of rib cage and abdomen during breathing. *J*
Appl Physiol 22: 407–422, 1967.
Lalley PM. μ-Opioid receptor agonist effects on medullary respiratory neurons in the cat: evidence for
involvement in certain types of ventilatory disturbances. *Am J Physiol-Regul Integr Comp Physiol* 285:
R1287–R1304, 2003.
Lalley PM. Opiate slowing of feline respiratory rhythm and effects on putative medullary phase-regulating
neurons. *Am J Physiol-Regul Integr Comp Physiol* 290: R1387–R1396, 2006.
Lalley PM, **Mifflin SW**. Oscillation patterns are enhanced and firing threshold is lowered in medullary respiratory
neuron discharges by threshold doses of a μ-opioid receptor agonist. *Am J Physiol-Regul Integr Comp*
Physiol 312: R727–R738, 2017.
Levitt ES, **Abdala AP**, **Paton JFR**, **Bissonnette JM**, **Williams JT**. μ opioid receptor activation hyperpolarizes
respiratory-controlling Kolliker–Fuse neurons ¨ and suppresses post-inspiratory drive. *J Physiol Neurosci*
593: 4453–4469, 2015.
Lindsey BG, **Rybak IA**, **Smith JC**. Computational Models and Emergent Properties of Respiratory Neural
Networks. In: *Comprehensive Physiology*, edited by Terjung R. Wiley, p. 1619–1670.
Liu S, **Kim D-I**, **Oh TG**, **Pao GM**, **Kim J-H**, **Palmiter RD**, **Banghart MR**, **Lee K-F**, **Evans RM**, **Han S**. Neural basis of
opioid-induced respiratory depression and its rescue. *Proc Natl Acad Sci* 118: e2022134118, 2021.
MacGregor RJ. *Neural and Brain Modeling*. New York: Academic Press, 1987.

Magosso E, **Ursino M**, **vanOostrom JH**. Opioid-Induced Respiratory Depression: A Mathematical Model for
Fentanyl. *IEEE Trans Biomed Eng* 51: 1115–1128, 2004.
Mattson CL, **Tanz LJ**, **Quinn K**, **Kariisa M**, **Patel P**, **Davis NL**. Trends and Geographic Patterns in Drug and
Synthetic Opioid Overdose Deaths — United States, 2013–2019. *MMWR Morb Mortal Wkly Rep* 70: 202–
207, 2021.
Mellen NM, **Janczewski WA**, **Bocchiaro CM**, **Feldman JL**. Opioid-Induced Quantal Slowing Reveals Dual
Networks for Respiratory Rhythm Generation. *Neuron* 37: 821–826, 2003.
Montandon G. Chapter 12 - The pathophysiology of opioid-induced respiratory depression. In: *Handbook of
Clinical Neurology*, edited by Chen R, Guyenet PG. Elsevier, p. 339–355.
Montandon G, **Liu H**, **Horner RL**. Contribution of the respiratory network to rhythm and motor output revealed
by modulation of GIRK channels, somatostatin and neurokinin-1 receptors. *Sci Rep* 6: 32707, 2016a.
Montandon G, **Ren J**, **Victoria NC**, **Liu H**, **Wickman K**, **Greer JJ**, **Horner RL**. G-protein–gated Inwardly Rectifying
Potassium Channels Modulate Respiratory Depression by Opioids. *Anesthesiology* 124: 641–650, 2016b.
Nectow AR, **Nestler EJ**. Viral tools for neuroscience. *Nat Rev Neurosci* 21: 669–681, 2020.
Neumueller SE, **Buiter N**, **Hilbert G**, **Grams K**, **Taylor R**, **Desalvo J**, **Hodges GL**, **Hodges MM**, **Pan LG**, **Lewis SJ**,
Forster HV, **Hodges MR**. Effects of sub-lethal doses of fentanyl on vital physiologic functions and
withdrawal-like behaviors in adult goats. *Front Physiol* 14, 2023.
O'Connor R, **Segers LS**, **Morris KF**, **Nuding SC**, **Pitts T**, **Bolser DC**, **Davenport PW**, **Lindsey BG**. A Joint
Computational Respiratory Neural Network-Biomechanical Model for Breathing and Airway Defensive
Behaviors. *Front Physiol* 3, 2012.
Palkovic B, **Marchenko V**, **Zuperku EJ**, **Stuth EAE**, **Stucke AG**. Multi-Level Regulation of Opioid-Induced
Respiratory Depression. *Physiology* 35: 391–404, 2020.
Pattinson KTS. Opioids and the control of respiration. *Br J Anaesth* 100: 747–758, 2008.
Phillips RS, **Baertsch NA**. Interdependence of cellular and network properties in respiratory rhythm generation.
Proc Natl Acad Sci 121: e2318757121, 2024.
Ramirez J-M, **Burgraff NJ**, **Wei AD**, **Baertsch NA**, **Varga AG**, **Baghdoyan HA**, **Lydic R**, **Morris KF**, **Bolser DC**, **Levitt
ES**. Neuronal mechanisms underlying opioid-induced respiratory depression: our current understanding. *J
Neurophysiol* 125: 1899–1919, 2021.
Segers LS, **Nuding SC**, **Dick TE**, **Shannon R**, **Baekey DM**, **Solomon IC**, **Morris KF**, **Lindsey BG**. Functional
connectivity in the pontomedullary respiratory network. *J Neurophysiol* 100: 1749–1769, 2008.
Segers LS, **Nuding SC**, **Ott MM**, **Dean JB**, **Bolser DC**, **O'Connor R**, **Morris KF**, **Lindsey BG**. Peripheral
chemoreceptors tune inspiratory drive via tonic expiratory neuron hubs in the medullary ventral
respiratory column network. *J Neurophysiol* 113: 352–368, 2015.
Shen TY, **Poliacek I**, **Rose MJ**, **Musselwhite MN**, **Kotmanova Z**, **Martvon L**, **Pitts T**, **Davenport PW**, **Bolser DC**.
The role of neuronal excitation and inhibition in the pre-Bötzinger complex on the cough reflex in the cat. *J
Neurophysiol* 127: 267–278, 2022.
Shevtsova NA, **Manzke T**, **Molkov YI**, **Bischoff A**, **Smith JC**, **Rybak IA**, **Richter DW**. Computational modelling of 5-
HT receptor-mediated reorganization of the brainstem respiratory network. *Eur J Neurosci* 34: 1276–1291,
2011.
da Silva CA, **Grover CJ**, **Picardo MCD**, **Del Negro CA**. Role of NaV1.6-mediated persistent sodium current and
bursting-pacemaker properties in breathing rhythm generation. *Cell Rep* 42: 113000, 2023.
Simera M, **Poliacek I**, **Jakus J**. Central antitussive effect of codeine in the anesthetized rabbit. *Eur J Med Res* 15:
184, 2010.
Skulsky EM, **Osman NI**, **Baghdoyan HA**, **Lydic R**. Microdialysis Delivery of Morphine to the Hypoglossal Nucleus
of Wistar Rat Increases Hypoglossal Acetylcholine Release. *Sleep* 30: 566–573, 2007.
Song C, **Alijani A**, **Frank T**, **Hanna GB**, **Cuschieri A**. Mechanical properties of the human abdominal wall
measured in vivo during insufflation for laparoscopic surgery. *Surg Endosc* 20: 987–990, 2006.

Sun X, **Thörn Pérez C**, **Halemani D N**, **Shao XM**, **Greenwood M**, **Heath S**, **Feldman JL**, **Kam K**. Opioids modulate
an emergent rhythmogenic process to depress breathing. *eLife* 8: e50613, 2019.
Sutter ME, **Wintemute GJ**, **Clarke SO**, **Roche BM**, **Chenoweth JA**, **Gutierrez R**, **Albertson TE**. The Changing Use of
Intravenous Opioids in an Emergency Department. *West J Emerg Med* 16: 1079, 2015.
Tomazini Martins R, **Carberry JC**, **Gandevia SC**, **Butler JE**, **Eckert DJ**. Effects of morphine on respiratory load
detection, load magnitude perception, and tactile sensation in obstructive sleep apnea. *J Appl Physiol* 125:
393–400, 2018.
Varga AG, **Reid BT**, **Kieffer BL**, **Levitt ES**. Differential impact of two critical respiratory centres in opioid-induced
respiratory depression in awake mice. *J Physiol* 598: 189–205, 2019.
Varga AG, **Reid BT**, **Kieffer BL**, **Levitt ES**. Differential impact of two critical respiratory centres in opioid-induced
respiratory depression in awake mice. *J Physiol* 598: 189–205, 2020.
Wenninger JM, **Pan LG**, **Klum L**, **Leekley T**, **Bastastic J**, **Hodges MR**, **Feroah TR**, **Davis S**, **Forster HV**. Large lesions
in the pre-Bötzinger complex area eliminate eupneic respiratory rhythm in awake goats. *J Appl Physiol* 97:
1629–1636, 2004.
Wolf SJ, **Byyny R**, **Carpenter CR**, **Diercks DB**, **Gemme SR**, **Gerardo CJ**, **Godwin SA**, **Hahn SA**, **Hatten BW**, **Haukoos
JS**, **Hickey SM**, **Kaji A**, **Kwok H**, **Lo BM**, **Mace SE**, **Nazarian DJ**, **Promes SB**, **Shah KH**, **Shih RD**, **Silvers SM**,
Smith MD, **Thiessen MEW**, **Tomaszewski CA**, **Valente JH**, **Wall SP**, **Winger J**, **Cantrill SV**, **Hirshon JM**, **Mims
M**, **Schulz T**, **Hatten BW**, **Cantrill SV**, **Dubin JS**, **Ketcham EM**, **Runde DP**, **Wall SP**, **Wolf SJ**. Clinical Policy:
Critical Issues Related to Opioids in Adult Patients Presenting to the Emergency Department. *Ann Emerg
Med* 76: e13–e39, 2020.

Figure 1

A. Core Respiratory Rhythmogenic Circuit

Pons INs

- I-pons
- EI-pons
- rIE-pons/ cIE-pons

50 spikes/bin

Medulla INs

- I-Driver
- I-Dec
- E-Dec-P
- I-Aug
- E-Dec-T

50 spikes/bin

Bulbospinal INs

- I-Aug-BS
- E-Aug-BS

25 spikes/bin

B. Motor Output

Phrenic MNs — 25 spikes/bin

Lumbar MNs — 5 spikes/bin

C. Airway Mechanics

Lung Vol/ Flow

- Lung Vol
- Flow

1 unit

D. Pulmonary Feedback

Lung PSRs

1 s

Figure 2

Figure 3

A

presynaptic spikes   NT/sp

i. $\Delta I_{app} = 0$
      $1x\ g_{syn}$

ii. $\Delta I_{app} > 0$
      $1x\ g_{syn}$

iii. $\Delta I_{app} < 0$
      $1x\ g_{syn}$

cellular effects

B

i. $\Delta I_{app} = 0$ postsynaptic
      $<1x\ g_{syn}$ effects

ii. $\Delta I_{app} = 0$ presynaptic
      $<1x\ g_{syn}$ effects

Figure 4

Figure 5

Medullary synaptic sources
Planar excitatory connections

A

Medulla

NRM BotC E-Dec-P I-Driver I-Dec E-Aug E-Dec-T I-Aug

E-Aug-BS I-Aug-BS

B

I-Driver→I-Driver
I-Driver→I-Aug
I-Driver→I-Dec
I-Driver→
I-Aug / I-Dec / I-Driver
I-Aug→I-Aug

T_{tot} (s)
5.75
4.75

T_i (s)
5
2

T_e (s)
3.75
2.75

T_{tot} CV
1.0
0.0

T_i CV
1.0
0.0

T_e CV
1.0
0.0

Scaling Factor
1.0   0.0

Scaling Factor
1.0   0.0

C

Phrenic Amp.
25
10

Lung Vol.
1.3
1.1

Insp. Flow
-0.6
-1.2

Scaling Factor
1.0   0.0

Figure 6

Figure 7

A Inhibitory influence of I-Dec

Pons

NRM-pons | E-pons | I-pons | rIE-pons | EI-pons | cIE-pons

Medulla

NRM BotC | E-Dec-P | I-Driver | I-Dec | E-Aug | E-Dec-T | I-Aug

E-Aug-BS | I-Aug-BS

Spinal cord

Lumbar | Phrenic

B I-Dec→inhibitory

T_{tot} (s) | T_{tot} CV

T_i (s) | T_i CV

T_e (s) | T_e CV

Scaling Factor | Scaling Factor

C

Phrenic Amp.

Lung Vol.

Insp. Flow

Scaling Factor

D I-Dec→inhibitory = 40% (0.4x)

Medulla INs — I-Driver, I-Dec, E-Dec-P, I-Aug, E-Dec-T — 50 sp/bin

Bulbospinal INs — I-Aug-BS, E-Aug-BS — 25 sp/bin

Phrenic MNs — 25 sp/bin

Lumbar MNs — 5 sp/bin

Lung Vol/Flow — Lung Vol, Flow — 1 unit

Lung PSRs

1 s

E I-Dec→inhibitory = 20% (0.2x)

50 sp/bin

25 sp/bin

25 sp/bin

5 sp/bin

1 unit

1 s

Figure 8

A

Pontine synaptic sources
Descending excitatory connections

B

Figure 9

NRM-pons → I-Aug / I-Driver = 20% (0.2x)

Pons INs — I-pons, EI-pons, rIE-pons/cIE-pons — 50 spikes/bin

Medulla INs — I-Driver, I-Dec, E-Dec-P, I-Aug, E-Dec-T — 50 spikes/bin

Bulbospinal INs — I-Aug-BS, E-Aug-BS — 25 spikes/bin

Phrenic MNs — 25 spikes/bin

Lumbar MNs — 5 spikes/bin

Lung Vol/Flow — Lung Vol, Flow — 1 unit

1 s

Figure 10