1

Possible involvement of ghost introgressions in the striking diversity of V1R receptor genes in East African cichlids

Shunsuke Taki[1], Zicong Zhang[2], Mitsuto Aibara[1], Tatsuki Nagasawa[1], and Masato Nikaido[1*]

[1]School of Life Science and Technology, Institute of Science Tokyo, Tokyo, Japan

[2]Institute for the Advanced Study of Human Biology, Kyoto University, Kyoto, Japan

[*]Correspondence: Masato Nikaido, mnikaido@bio.titech.ac.jp

Abstract

Cichlids that have experienced adaptive radiation are genetically close but exhibit extreme ecological and morphological diversity, making them widely used as textbook examples to understand speciation mechanisms. V1R-type olfactory receptors are highly conserved at the amino acid sequence level among teleost fish and believed to play a fundamental role in reproduction. Our previous studies have reported the unexpected sequence diversity of V1Rs among some cichlid species, implying the possible involvement of V1Rs in their speciation. In this study, we extensively investigated the process of the evolutionary diversification of all six V1Rs (V1R1 to V1R6) by utilizing the genome data from 528 cichlid species, covering almost all lineages. In V1R2, two highly divergent alleles (1.17%: variant sites/CDS length) without recombination were shared and maintained among the cichlids of all East African Great Lakes. In V1R6, large number of highly variable alleles that could be derived from the multiple recombination of two highly divergent alleles (1.39%: variant sites/CDS length) were shared among the cichlids of Lake Victoria. Additionally, we also identified highly divergent alleles in V1R1 within the tribe Tropheini, and in both V1R3 and V1R6 within Trematocarini and Ectodini. Given that one of the two divergent alleles of these V1Rs emerged so rapidly during cichlid evolution, they are likely to be derived from introgression. However, the source lineages for those introgressions were not identified despite the extensive exploration implying that they have already been extinct. This work provides compelling support for the contribution of introgression to explain striking diversity of V1Rs in East African cichlids.

Keywords: adaptation, cichlids, ghost introgression, standing genetic variation, V1R

Introduction

Cichlids, belonging to the family Cichlidae, are a group of tropical freshwater fish that inhabit a wide range of regions from Central and South America to Africa, Madagascar, and India. The total number of species is estimated to be about 3,000, representing about 10% of all teleost fish (Kocher, 2004). Notably, about half of the extant cichlid species inhabit the East African Great Lakes (Lake Tanganyika, Lake Malawi, and Lake Victoria) (Salzburger et al., 2014), and these species have emerged through large-scale adaptive radiation in each lake (Kocher, 2004; Salzburger, 2018). These cichlids of each Lake are widely used as model organisms to investigate the mechanism of speciation because of their remarkable ecological and morphological diversity despite their genetic closeness. Cichlids are often reproductively separated by prezygotic isolation (Rometsch et al., 2020). Sexual selection and assortative mating contribute to speciation, particularly in environments where geographic isolation is less likely, such as lakes (Seehausen & Wagner, 2014).

Cichlids depend on multiple sensory systems for their reproductive behavior (Escobar-Camacho & Carleton, 2015). Research on the visual system is especially active, as cichlids exhibit a wide variety of body colors. Previous studies have shown that the visual system has diversified according to ecology and habitat, and that its environmental adaptation have contributed to speciation (Carleton & Yourick, 2020; Seehausen et al., 2008; Terai et al., 2006). The promotion of speciation through environmental adaptation of sensory systems is referred to as "sensory drive", which has been shown to play a significant role in the speciation of cichlids. The prominent example is the visual adaptation to the light environment (bule or red) through peak shift of long-wavelength-sensitive (lws) opsin together with the divergence of female preference for male nuptial coloration (bule or red) led to the speciation of Lake Victoria cichlids (Seehausen et al., 2008).

Cichlids use not only visual but also olfactory cues for species and individual

recognition (Keller-Costa et al., 2015; Santos et al., 2023). Behavioral experiments using

Lake Malawi cichlids have demonstrated that olfaction plays a role in conspecific recognition

(Plenderleith et al., 2005). The imprinting on olfactory cues through mouthbrooding have also

shown in Lake Victoria cichlids (Verzijden & ten Cate, 2007). In addition, compounds present

in the urine of Mozambique tilapia (*Oreochromis mossambicus*) act as pheromones,

influencing male behavior and the female reproductive system (Ashouri et al., 2023; Keller-

Costa et al., 2014). It is obvious that cichlids largely depend on olfaction, however, in contrast

to studies of the visual system, it remains unclear which olfactory receptor genes and alleles

contribute to species recognition and hence assortative mating.

Four types of olfactory receptor genes (OR, TAAR, V1R, V2R) have been identified

in teleost fish, all of them are expressed in the olfactory neurons (Calvo-Ochoa et al., 2021).

One of these, the V1R (vomeronasal type 1 receptor), which is also called ORA (olfactory

receptor class A related), is considered to detect pheromones in fish, as it is homologous to

the mammalian V1R, a known pheromone receptor (Del Punta et al., 2002; Pfister &

Rodriguez, 2005). Although the V1Rs are diverse in copy number and sequence in mammals,

they are highly conserved in teleost fish by sharing six copies of the V1R genes (V1R1 to

V1R6), of which the orthologs are also highly conserved at amino acid sequences level

(Saraiva & Korsching, 2007). Therefore, they likely to detect hormonal pheromones shared

across species and to contribute to the synchronization of gamete maturation and/or

spawning interactions (Ota et al., 2012; Policarpo et al., 2022; Saraiva & Korsching, 2007;

Zapilko & Korsching, 2016). The V1Rs in zebrafish has been shown to detect 4HPAA (4-

hydroxyphenylacetic acid) and bile acids (Behrens et al., 2014; Cong et al., 2019).

Additionally, V1R2 and V1R5 expressing neurons respond to male urine in cichlids

(Kawamura & Nikaido, 2022). Thus, V1Rs have great potential to contribute to speciation in

cichlids, however, it is not clear whether their genetic diversity is linked to speciation or

environmental adaptation, as has been shown in LWS.

The V1R genes of East African cichlids exhibit sequence-level diversity (Nikaido et

al., 2014). In our previous study, we found diverse alleles of V1R1, V1R3, and V1R6 in some

tribes of Lake Tanganyika cichlids. We also found two different alleles of V1R2, and revealed

that these are shared across species from different lakes. These alleles are separated by

more than 10 nonsynonymous substitutions, indicating the signature of functional

differentiation. Hence, V1Rs are ideal research targets for examining the possibility of

olfaction-mediated speciation of cichlids (Santos et al., 2023). However, the small number of

species analyzed hindered a complete understanding of the origin and formation process of

these divergent alleles.

In recent years, the release of whole-genome sequencing (WGS) data for more than

500 cichlid species has enabled comprehensive comparative genomic analyses across

species (Malinsky et al., 2018; McGee et al., 2020; Ronco et al., 2021). This allowed us to

extensively explore the V1R sequences to understand how the divergent alleles were

originated and maintained across lineages in morphologically and ecologically diverse

cichlids. In the present study, we conducted a comprehensive analysis of V1R allelic diversity

by using WGS data from 528 species and 908 samples of cichlids. Furthermore, we

investigated the potential impact of sequence-level differences on V1R function, particularly

ligand reception.

Materials and Methods

V1R gene sequences

We used short-read data from 28 tribes, 528 species, and 908 samples of cichlids registered

in the NCBI Sequence Read Archive (SRA) (Figure S1, Table S1). Low-quality reads were

removed using fastp v.0.23.2 (Chen et al., 2018). The filtered reads were mapped to the

reference genome (Supplementary Information) using bwa-mem2 v.2.2.1 (Vasimuddin et al.,

2019), and sorted and indexed with samtools v.1.15 (Danecek et al., 2021). The V1R gene

regions were extracted from the sorted BAM files using samtools v.1.15 (Danecek et al.,

2021). Variants were called from the extracted BAM files using GATK v.4.3.0 (McKenna et

al., 2010) with the HaplotypeCaller tool. The first round of hard filtering for SNPs and INDELs

was applied using GATK v.4.3.0 (McKenna et al., 2010) with the VariantFiltration tool and the

following parameters: SNPs (QD<2.0, QUAL<30.0, SOR>4.0, FS>60.0, MQ<40.0,

MQRankSum<-12.5, ReadPosRankSum<-8.0) and INDELs (QD<2.0, QUAL<30.0,

FS>200.0, SOR>10.0, ReadPosRankSum<-20.0). Based on the resulting VCF file, quality

recalibration of the BAM files was performed using GATK v.4.3.0 (McKenna et al., 2010) with

BaseRecalibrator and ApplyBQSR. The second round of hard filtering was conducted as in

the first round. The filtered VCF files were used to create V1R gene sequences in fasta format

using bcftools v.1.15 (Danecek et al., 2021). Sequences containing heterozygous variants

were phased using HapCUT2 v.1.3.3 (Edge et al., 2017). We built a custom pipeline from

downloading short-read data to getting V1R gene sequences (https://github.com/taki-

sh/RegionCall).

Phylogenetic analysis

To reduce the computational load of the phylogenetic analysis, we removed duplicated

sequences and then randomly extracted a total of 45 sequences from 337 V1R2 sequences,

creating five datasets (Figure 1, Figure S2-6). We performed multiple sequence alignment

for all of these datasets using MAFFT v.7.520 (Katoh & Standley, 2013) with the options: --

maxiterate 16 --localpair. Maximum likelihood trees were constructed using RAxML-NG

v.1.2.0 (Kozlov et al., 2019) with 1000 bootstrap replicates. The model used for phylogenetic

tree construction, HKY+G4, which was selected by Modeltest-NG v.0.1.7 (Darriba et al.,

2020). We used Dataset 1 in Figure 1, as Datasets 1 through 5 demonstrated similar

topological patterns (Figure S2-6). For V1R1 of Tropheini, we used 43 non-duplicated V1R1

sequences along with the outgroup *Oreochromis niloticus* V1R1 sequence, totaling 44

sequences (1000 bootstrap replicates, model: HKY+I). For V1R3 of Trematocarini, we used

11 non-duplicated V1R3 sequences along with the outgroup *Oreochromis niloticus* V1R1

sequence, for a total 12 sequences (1000 bootstrap replicates, model: HKY+I). For V1R3

and V1R6, we used 53 non-duplicated V1R3 sequences (100 bootstrap replicates, model

V1R3: HKY+G4, V1R6: GTR+G4). For visualization, we used the R packages: treeio (Wang

et al., 2020), ggtree (Yu et al., 2017), readr (Wickham H, Hester J, Bryan J, 2024), dplyr

(Wickham H, François R, Henry L, Müller K, Vaughan D, 2023), ggplot2 (Villanueva & Chen,

2019), ggmsa (Zhou et al., 2022), and iTOL v6 (Letunic & Bork, 2024). To calculate the

frequency of each allele of V1R1 and V1R2 across cichlid lineages, we used unphased

sequences (V1R1: 908 sequences, V1R2: 907 sequences). We used 1309 sequences for

the classification of V1R6 haplotypes. Additionally, to reveal the sequence diversity of V1R

genes in ray-finned fish, we used V1R sequences from 102 species of ray-finned fish

(Policarpo et al., 2022) and phylogenetic tree data (Chang et al., 2019; Rabosky et al., 2018).

We generated a pairwise similarity matrix for the 102 species using CIAlign v.1.0.17

(Tumescheit et al., 2022) and calculated the average similarity. We also calculated pairwise

dN/dS ratios using the CODEML program in PAML v.4.10.6 (Yang, 2007) and computed the

average value.

Selection analysis

Amino acid sequences translated using Aliview (Larsson, 2014), were aligned in MAFFT

v.7.520 (Katoh & Standley, 2013) with the options: --maxiterate 16 --localpair. This multiple

alignment of amino acid sequences was converted into a codon alignment using PAL2NAL

v.14 (Suyama et al., 2006). For branches where an accumulation of nonsynonymous

substitutions was observed, we estimated dN/dS ratios using branch model with the codeml

program in PAML v.4.10.7 (Yang, 2007) and performed a likelihood ratio test between the

foreground and background branches to examine the statistical significance for the elevation

of dN/dS in foreground branch. Clade I and Clade II were defined within V1R2. Using PAML,

the dN/dS ratio was calculated across all five datasets (Dataset 1 to 5, Figure S2-6), and

ancestral sequences were inferred for the ancestral and descendant nodes of each branch.

A comparison of the inferred ancestral sequences identified 11 nonsynonymous substitutions

common to all datasets, which were classified as mutations distinguishing Clade I and Clade

II. In V1R1, we also defined Clade I and Clade II based on the phylogenetic tree of the V1R1

gene in Tropheini (Figure S9). Similarly, we identified 11 nonsynonymous substitutions, which

were classified as mutations distinguishing Clade I and Clade II.

Protein modeling and ligand-binding predictions

We modeled 3D structures of V1R1, V1R2, and V1R6 using AlphaFold v.2.3 (Jumper et al.,

2021). Based on the most reliable structures, we estimated the coordinates of the ligand-

binding pockets using Deepsite (Jiménez et al., 2017) in PlayMolecule Viewer (Torrens-

Fontanals et al., 2024). Furthermore, we performed ligand docking simulations using Webina

v.1.0.5 (Kochnev et al., 2020) with the obtained coordinates. For these simulations, we used

4HPAA, a known ligand for zebrafish V1R2 (Behrens et al., 2014), and LCA (lithocholic acid),

a known ligand for zebrafish V1R1 and V1R6 (Cong et al., 2019). As a result, five relative

positions (poses) of the ligand in V1R1 (clade I: 3 poses, clade II: 2 poses), 18 poses in V1R2

(clade I: 9 poses, clade II: 9 poses), and 18 poses in V1R6 (ancestral allele: 9 poses, derived

allele: 9 poses) were predicted. Residues positioned within 5Å of all these poses were

designated as ligand-related sites. In addition, we predicted transmembrane regions using

DeepTMHMM v.1.0.39 (Hallgren et al., 2022). Visualization of the results was carried out

using ChimeraX v.1.6 (Goddard et al., 2018).

Recombination rate

We used the recombination rate of *Pundamilia nyererei* (Feulner et al., 2018) and the

genomic position of the V1R genes to relatively compare the recombination rates in the

regions surrounding the V1R genes. We obtained the genomic positions of the V1R genes

using BLASTN with the *Pundamilia nyererei* genome assembly (P_nyererei reference v2)

(Feulner et al., 2018) as the reference and *Oreochromis niloticus* V1R sequences (V1R1:

AB671405.1, V1R2: AB670433.1, V1R3: AB704581.1, V1R4: AB704618.1, V1R5:

AB704661.1, V1R6: AB704705.1 from NCBI) as the query. The genomic positions of the V1R

genes were identified as follows: V1R1 (chr13:8171366-8172331), V1R2 (chr13:8175817-

8176758), V1R3 (chr13:18527399-18530109), V1R4 (chr13:18531988-18532072), V1R5

(chr2:41585178-41586302), V1R6 (chr7:27107196-27108122).

Results

Two highly divergent alleles of V1R2 in cichlids of East African Great Lakes

We constructed a phylogenetic tree of V1R2 gene using sequences from 44 species selected

from all cichlid lineages (Figure 1). Because cichlids in the East African Great Lakes have

independently undergone adaptive radiation in each lake, species inhabiting the same lake

are closely related, with Lake Malawi and Lake Victoria each forming monophyletic groups

(Svardal et al., 2021). However, the topology of the V1R2 gene tree did not match the species

tree and was divided into clade I and clade II with maximal bootstrap probability (Figure 1).

Clade I included cichlids from the outgroup (India, Madagascar), African Riverine, Lake

Tanganyika, Lake Malawi, and Lake Victoria. Clade II included cichlids from Lake Tanganyika,

ancient Haplochromini, modern Haplochromini, Lake Malawi, Upper Nile lineage, Eastern

lineage, Congolese lineage, and Lake Victoria. Notably, both clade I and clade II contained

lineages from all of the three East African Great Lakes. Clade I and clade II were separated

by 11 amino acids (Figure 1, see Method). In addition, the dN/dS ratio of the branch was

significantly higher than that of the background (Table S2). This elevation of dN/dS indicates

that positive selection operated on the branch where clade II diverged from clade I.

We calculated the frequencies of the two major alleles (clade I and clade II) of V1R2 in

each cichlid lineage (Figure 2). In cichlids from Lamprologini and Ectodini, inhabiting Lake

Tanganyika, V1R2 alleles were fixed into clade II. In Haplochromini from Lake Malawi and

Lake Victoria, the alleles were not fixed into neither allele, becoming the mixtures of

homozygotes of clade I or clade II, as well as their heterozygotes. Despite the extensive

exploration, clade II was not observed in the lineages from India, Madagascar, New World,

or African Riverine. Interestingly, no recombination between the two alleles was detected in

any lineage in spite that the alleles co-exist as polymorphisms within population or species

with only one exception for Cyphotilapiini. The polymorphic sites separating clade I and clade

II in cichlids are highly conserved among non-cichlids ray-finned fish, of which V1R are all

classified into clade I (Figure S7).

Highly variable alleles of V1R6 with multiple recombination in cichlids of Lake Victoria

We next explored the V1R6 gene and constructed the phylogenetic tree, of which the

topology is consistent with the species tree (Figure 3A). Interestingly, we identified striking

number of different alleles that were separated by the combination of 13 variant sites in Lake

Victoria cichlids (Figure 3B). These variable alleles were shared among species as

polymorphisms. It is highly unlikely that the substitutions occur at the same site multiple times

independently in each allele. Therefore, we hypothesized that the striking number of different

alleles were derived from the recombination between two highly divergent alleles present in

the ancestral population in Lake Victoria (Figure 3C). Indeed, we calculated the

recombination rate in the genomic region surrounding V1R6 using the data from previous

study (Feulner et al., 2018), showing the higher recombination rate compared to other V1R

surrounding regions (Figure S8). Based on this hypothesis, we focused on the 13 variant

sites identified in Lake Victoria and classified the V1R6 alleles of the cichlids of all East

African Great Lakes and rivers. In cichlids inhabiting outside the Lake Victoria basin, we

found no apparent signature of recombination between highly divergent alleles. We then

reconstructed the ancestral sequences at the node marked with an arrowhead in Figure 3A

and defined it as "ancestral allele" (Figure 3B left top, dark blue nucleotide). Next, by

subtracting the nucleotides of "ancestral allele" (dark blue) from the 13 polymorphic sites in

Lake Victoria, we reconstructed a hypothetical "derived allele" (yellow), which could be a

counterpart sequence of the past recombination (Figure 3B right top). As a result, the alleles

found in Lake Victoria are divided into the "ancestral", "derived" and their "mixtures". Notably,

the cichlid of Upper Nile, Eastern lineage, and Congolese lineage also possess recombinant

alleles similar to those in Lake Victoria (Figure 3B). However, in spite of the extensive

exploration, the hypothetical "derived allele" was not identified in extant cichlids except for

Lake Victoria (e.g., haplotypes Lake Victoria 51, 52 in Figure 3B right bottom).

Highly divergent alleles of V1R1, V1R3, V1R6 in cichlids of particular tribes

We identified two highly divergent alleles named clade I and clade II, which are separated by

11 amino acids, in the V1R1 of Tropheini (Figure 4A, see Method). The monophyly of clade

I and clade II are supported by maximal bootstrap probability (Figure S9). Given that the

amino acid sequence of clade I of Tropheini is mostly similar with other cichlids, clade I is

defined as "ancestral" and II as "derived" alleles. Furthermore, the dN/dS ratio of the branch

was significantly higher than that of the background (Table S2). This elevation of dN/dS

indicates that positive selection operated on the branch where clade II diverged from clade I.

Next, we calculated the frequencies of the two major alleles (clade I and clade II) in the V1R1

of Tropheini. We found that 28% of samples in Tropheini possess clade I, while 65% possess

clade II (Figure 4B). We extensively explored the clade II sequences, but did not find them in

any cichlids other than Tropheini. The polymorphic sites separating clade I and clade II in

cichlids are highly conserved at clade I among ray-finned fish (Figure S10).

We also identified highly divergent allele in the V1R3 in cichlids of Ectodini and

Trematocarini. Seven nonsynonymous substitutions were observed on the branch

representing the common ancestor of Ectodini (Figure S11). The dN/dS ratio of this branch

was significantly higher than that of the background (Table S2). In Trematocarini, ten

nonsynonymous substitutions were observed on the branch representing the common

ancestor of Trematocarini, excluding *Trematocara unimaculatum* (Figure S12). The dN/dS

ratio of this branch was significantly higher than that of the background (Table S2). In V1R6,

we found the divergent alleles in Ectodini and Trematocarini (Figure S13) and the dN/dS ratio

of those clades (Ectodini: 2.69, Trematocarini: 1.61) were significantly higher than that of the

background (Table S2).

V1R2 allele sites are located near ligand binding pocket

To elucidate the functional impact of the amino acid substitutions observed between the two

major divergent alleles of V1R2, we predicted the 3D structure and ligand binding sites of

V1R2 (Figure 5A). First, we predicted the 3D structures for each allele using the amino acid

sequences of clade I and clade II. We then used 4HPAA, which is one promising candidate

of the ligand (Behrens et al., 2014; Kawamura & Nikaido, 2022), to estimate the ligand

binding sites and compared them between the alleles. As a result, we identified that eight out

of the 11 amino acid substitution sites (TM2: C62→Ser, ECL1: Q65→Pro, TM4: A145→Ser,

A146→Thr, ECL2: P159→Ser, H169→Leu, TM5: V185→Ile, TM6: D247→Glu) are located

close to the predicted ligand pocket (Figure 5B). Furthermore, four of these sites (TM4: A145

→Ser, A146→Thr, TM5: V185→Ile, TM6: D247→Glu) coincided with the ligand binding sites

(Figure 5B). The skewed amino acid substitutions of V1R2 towards the ligand binding sites

were statistically significant (Fisher's exact test: p=0.0389).

We also the same analysis of V1R1 and V1R6 (Figure S14A, S15A). In V1R1, we identified

that five out of the 11 amino acid substitution sites (TM2: V59→Gly, ECL2: I164→Leu, TM5:

G188→Val, TM7: S273→Thr, G280→Ser) are located close to the predicted ligand pocket

(Figure S14B). Furthermore, three of these sites (ECL2: I164→Leu, TM7: S273→Thr, G280

→Ser) coincided with the ligand binding sites (Figure S14B). However, the skewed amino

acid substitutions of V1R1 towards the ligand binding sites were not statistically significant

(Fisher's exact test: p=0.0922). In V1R6, we could not identify any amino acid substitution

sites that are located close to the predicted ligand pocket (Figure S15B).

Discussion

Allelic and functional diversity of V1R in the East African cichlids

We conducted large-scale analyses focusing on the diversity of V1R in cichlids, using WGS data from 528 species and 908 samples, which recently became available by the large scale comparative genomic studies (Malinsky et al., 2018; McGee et al., 2020; Ronco et al., 2021). As a result, we identified an accumulation of nonsynonymous substitutions and the signatures of the positive selection in V1R1, V1R2, V1R3, and V1R6 (Figure 6A). Notably, two highly divergent alleles were found in V1R1 and V1R2 of the cichlids of Tropheini (Fig. 4B) and those inhabiting the East African Great Lakes (Figure 2B), respectively. In addition, highly divergent alleles were found in V1R3 and V1R6 of the cichlids of Trematocarini and Ectodini (Figure S11-13). The results of the present study well supports previous findings from small scale species sampling (Nikaido et al., 2014).

Each of the V1R gene orthologs is highly conserved across fish species (Ota et al., 2012; Policarpo et al., 2022; Saraiva & Korsching, 2007; Zapilko & Korsching, 2016). The recently published V1R gene sequences of ray-finned fish (Policarpo et al., 2022) also exhibit high sequence similarity and low dN/dS ratios (Figure S16). Furthermore, the polymorphic sites separating the divergent alleles of V1R1 and V1R2 in cichlids are highly conserved among ray-finned fish (Figure S7, S10). These results suggest that the diversification of the V1R gene is unique to cichlids among ray-finned fish and are likely to be linked with their rapid adaptive radiation.

The prediction of the 3D structure of V1R2 and the estimation of the sites interacting with the ligand revealed that the amino acid residues distinct between clade I and II are located close to the ligand pocket. In addition, the amino acid substitutions were skewed towards the ligand binding sites, which was statistically significant (Figure 5B). Two amino

acid substitutions occurred in ECL2 (extracellular loop 2) (Figure 5B), which is involved in

ligand selectivity in class A GPCRs, a category that includes V1R (Nicoli et al., 2022). In the

bitter taste receptor T2R, which is phylogenetically close to V1R, ECL2 also contributes to

ligand binding (Xu et al., 2022). These lines of observation suggest that two divergent clades

of V1R2 are likely to be distinct each other in terms of ligand binding.

V1R6 has the lowest sequence similarity and the highest dN/dS ratio among six V1R

genes across all ray-finned fish (Figure S16). Hence, the pressure of negative selection

operated on V1R6 is weaker than other V1Rs not only in cichlids but also across ray-finned

fish. It is worth noting that V1R6 in Lake Victoria basin is highly variable with number of

different alleles due to the multiple recombination between divergent alleles (Figure 3B, 3C).

The recombination rate in the V1R6 region of Lake Victoria cichlids is notably high (Figure

S8), indicating that V1R6 may be located in a recombination hotspot. This is in sharp contrast

with V1R1 and V1R2, in which recombination is strictly suppressed. Thus, the structural

constraint on V1R6 seems to be weak, implying that the ligand specificity of V1R6 may be

lower than other V1Rs. In the 3D structural analysis, a ligand pocket of V1R1 and V1R2 was

predicted within the TM region, that is similar to typical class A GPCRs (Figure 5B, S14B). In

contrast, the ligand binding of V1R6 was predicted to be mediated by the ECL2 region (Figure

S15B). This suggests that V1R6 may be distinct with V1R1 and V1R2 in terms of functions

and ligand binding mechanisms. Further verification by considering the distinctness of the

predicted structural differences among V1Rs will provide insight into the function of V1Rs in

fish, which remain a topic of ongoing debate (Behrens et al., 2014; Cong et al., 2019; Saraiva

& Korsching, 2007).

The origin of striking V1R diversity through ghost introgression

It is of primary importance to understand how the remarkable diversity of V1Rs has been

originated and retained among the cichlids over the 10 million years of their evolution. We

first investigated the origin and process of V1R2 diversification by extensively examining the

existence of clade I and clade II alleles in each lineage of cichlids (Figure 2). As a result,

clade II was found in cichlid lineages inhabiting the East African Great Lakes, and was not

found in lineages that diverged prior to the Lake Tanganyika radiation. Specifically, the

cichlids of Lamprologini and Ectodini were fixed into clade II, and those of other Lake

Tanganyika tribes were fixed into clade I. The cichlids of Haplochromini were homozygotes

of clade I or clade II as well as the heterozygotes of them. The pattern of V1R2 allele

distribution suggest that the clade II suddenly appeared at the timing of radiation of Lake

Tanganyika cichlids around 10 million years ago. Given that the adaptive radiation in Lake

Tanganyika was so rapid, it is unlikely that 12 substitutions would accumulate step by step in

a single allele during that time, regardless of the operation of positive selection. Therefore, it

is parsimonious to assume that the clade II allele has already existed in some

populations/species prior to the radiation, and was introgressed into Lake Tanganyika

through hybridization.

Several examples of standing genetic variation (SGV), some of which date back to even

before the adaptive radiations in Lake Tanganyika (Nakamura et al., 2023), have been

identified in the genomes of East African cichlids (Nakamura et al., 2021; Svardal et al., 2021).

Clade I and II alleles of V1R2 well represents an example of such old SGV. The Melting-pot

hypothesis is one of the most plausible scenarios to explain the origin of old SGV (Nakamura

et al., 2023; Weiss et al., 2015). According to the Melting-pot hypothesis, several cichlid

populations/species, which were scattered in wetlands and small lakes, hybridized just before

the adaptive radiation in Lake Tanganyika. In case that each of the scattered populations

were fixed into clade I and clade II, the hybridization of them may provide the origin of SGV

of highly divergent V1R2. However, one of the two alleles comprising the SGV of V1R2, clade

II, were not found in spite of an extensive exploration of 46 basal lineages of riverine species

(Figure 2). The absence of clade II allele in the riverine species suggest that the transporter

lineage of clade II alleles have been already extinct.

In recent years, gene flow from extinct lineages (ghost introgression) has been detected

in various vertebrates from gobiid fish to elephants (Ai et al., 2015; Frei et al., 2022;

Gopalakrishnan et al., 2018; Kato et al., 2023; Palkopoulou et al., 2018; Zhang et al., 2019).

In cichlids, sudden emergence of clade II of V1R2 in Lake Tanganyika cichlids can be

explained by ghost introgression. In general, introgressed alleles are detected as highly

divergent compared with non-introgressed genomic region. However, neutral genomic

regions are often subject to recombination and drift after introgression, making it difficult to

detect the signature of introgression. In this regard, V1R2 is a good example to detect

introgression because recombination was strictly restricted. African riverine lineages that

existed prior to the adaptive radiation are referred to as "transporters" of genetic diversity

(Loh et al., 2013). Notably, gene flow from Steatocranini, a riverine lineage that diverged

before the adaptive radiation (Irisarri et al., 2018), and the appearance of clade II of the V1R2

occurred at identical time period. Thus, it is highly likely that the remarkable diversity of V1R2

alleles (clade II) derived from the introgression from an extinct African riverine lineage that

diverged prior to the adaptive radiation (ghost introgression) or from the unexamined lineages

(Figure 6B).

Our data on V1R6 suggest that remarkable number of different alleles derived from the

multiple times of recombination of two highly divergent alleles (ancestral and derived)

separated by 13 nucleotide substitutions was observed in cichlids of Lake Victoria basin

(Figure 3B). In Lake Victoria, the recombination of the LWS opsin gene was observed, which

was explained by the hybridization between the Congo and Upper Nile lineages (Meier et al.,

2017). In contrast, given that both the Congo and Upper Nile lineages also exhibit

recombinant sequences, recombination between ancestral and derived alleles of V1R6

occurred earlier than the hybridization event of the Congo and Upper Nile lineages. However,

none of the extant species in Haplochromini, except for Lake Victoria cichlids, has the derived

allele. Considering that the estimated mutation rate in cichlids is to be 3.5×10^{-9} (per site per

generation), it is unrealistic that 13 substitutions accumulate during the short period of the

formation of the cichlids of Lake Victoria basin (~400,000 years ago) (Johnson et al., 1996;

Salzburger et al., 2014). Therefore, the derived allele of V1R6 was also likely to be

introgressed from an extinct lineage (Figure 6B). Similar to the logics of V1R2 and V1R6, the

highly divergent alleles observed in V1R1 of Tropheini, as well as the in V1R3 and V1R6 in

Ectodini and Trematocarini, are also likely to have been derived from ghost introgression.

It is worth noting that the nucleotide sequence differences between the two alleles found

in V1R1, V1R2, and V1R6 were mostly all approximately higher than 1% (V1R1 = 1.14%,

V1R2 = 1.17%, V1R6 = 1.39%). It is possible to hypothesize that each divergent allele has

been introduced into the ancestral population through a single ghost introgression event

(Melting-pot), after which the derived alleles independently sorted to each lineage.

Alternatively, introgression events in each lineage may have led to the introduction of the

derived alleles in V1R1, V1R2, and V1R6 (Figure 6A, 6B).

Conclusion

In this study, through a comprehensive analysis of the V1R genes covering the most lineages

of East African cichlid species, we identified the large scale standing genetic variation that

could be explained by ghost introgression. It will be necessary to conduct the genome-wide

statistical analyses rather than at the gene level to examine the ghost introgression and

Melting-pot scenarios in further research. We also revealed differences in the structural

constraint and selection pressure among different V1Rs in cichlids, implying the functional

distinctness among them. This research provides important insights into the possibility of

V1R mediated speciation and the genetic source enabling the adaptive evolution of cichlids.

Acknowledgements

This study was supported by JSPS KAKENHI (20KK0167, 24K02074, 24KK0141) to MN.

Phylogenetic analyses were partially performed using the NIG supercomputer at ROIS

National Institute of Genetics.

References

Ai, H., Fang, X., Yang, B., Huang, Z., Chen, H., Mao, L., Zhang, F., Zhang, L., Cui, L., He,

W., Yang, J., Yao, X., Zhou, L., Han, L., Li, J., Sun, S., Xie, X., Lai, B., Su, Y., …

Huang, L. (2015). Adaptation and possible ancient interspecies introgression in pigs

identified by whole-genome sequencing. *Nature Genetics*, *47*(3), 217–225.

https://doi.org/10.1038/ng.3199

Ashouri, S., Da Silva, J. P., Canário, A. V. M., & Hubbard, P. C. (2023). Bile acids as putative

social signals in Mozambique tilapia (Oreochromis mossambicus). *Physiology &*

Behavior, *272*, 114378. https://doi.org/10.1016/j.physbeh.2023.114378

Behrens, M., Frank, O., Rawel, H., Ahuja, G., Potting, C., Hofmann, T., Meyerhof, W., &

Korsching, S. (2014). ORA1, a zebrafish olfactory receptor ancestral to all

mammalian V1R genes, recognizes 4-hydroxyphenylacetic acid, a putative

reproductive pheromone. *The Journal of Biological Chemistry*, *289*(28), 19778–

19788. https://doi.org/10.1074/jbc.M114.573162

Calvo-Ochoa, E., Byrd-Jacobs, C. A., & Fuss, S. H. (2021). Diving into the streams and

waves of constitutive and regenerative olfactory neurogenesis: insights from

zebrafish. *Cell and Tissue Research*, *383*(1), 227–253.

https://doi.org/10.1007/s00441-020-03334-2

Carleton, K. L., & Yourick, M. R. (2020). Axes of visual adaptation in the ecologically diverse

family Cichlidae. *Seminars in Cell & Developmental Biology*, *106*, 43–52.

https://doi.org/10.1016/j.semcdb.2020.04.015

Chang, J., Rabosky, D. L., Smith, S. A., & Alfaro, M. E. (2019). An r package and online

resource for macroevolutionary studies using the ray-finned fish tree of life. *Methods*

in Ecology and Evolution / British Ecological Society, *10*(7), 1118–1124.

https://doi.org/10.1111/2041-210X.13182

Chen, S., Zhou, Y., Chen, Y., & Gu, J. (2018). fastp: an ultra-fast all-in-one FASTQ

preprocessor. *Bioinformatics*, *34*(17), i884–i890.

https://doi.org/10.1093/bioinformatics/bty560

Cong, X., Zheng, Q., Ren, W., Chéron, J.-B., Fiorucci, S., Wen, T., Zhang, C., Yu, H.,

Golebiowski, J., & Yu, Y. (2019). Zebrafish olfactory receptors ORAs differentially

detect bile acids and bile salts. *The Journal of Biological Chemistry*, *294*(17), 6762–

6771. https://doi.org/10.1074/jbc.RA118.006483

Danecek, P., Bonfield, J. K., Liddle, J., Marshall, J., Ohan, V., Pollard, M. O., Whitwham, A.,

Keane, T., McCarthy, S. A., Davies, R. M., & Li, H. (2021). Twelve years of SAMtools

and BCFtools. *GigaScience*, *10*(2). https://doi.org/10.1093/gigascience/giab008

Darriba, D., Posada, D., Kozlov, A. M., Stamatakis, A., Morel, B., & Flouri, T. (2020).

ModelTest-NG: A New and Scalable Tool for the Selection of DNA and Protein

Evolutionary Models. *Molecular Biology and Evolution*, *37*(1), 291–294.

https://doi.org/10.1093/molbev/msz189

Del Punta, K., Leinders-Zufall, T., Rodriguez, I., Jukam, D., Wysocki, C. J., Ogawa, S., Zufall,

F., & Mombaerts, P. (2002). Deficient pheromone responses in mice lacking a cluster

of vomeronasal receptor genes. *Nature*, *419*(6902), 70–74.

https://doi.org/10.1038/nature00955

Edge, P., Bafna, V., & Bansal, V. (2017). HapCUT2: robust and accurate haplotype assembly

for diverse sequencing technologies. *Genome Research*, *27*(5), 801–812.

https://doi.org/10.1101/gr.213462.116

Escobar-Camacho, D., & Carleton, K. L. (2015). Sensory modalities in cichlid fish behavior.

Current Opinion in Behavioral Sciences, *6*, 115–124.

482 https://doi.org/10.1016/j.cobeha.2015.11.002

483 Feulner, P. G. D., Schwarzer, J., Haesler, M. P., Meier, J. I., & Seehausen, O. (2018). A Dense

484   Linkage Map of Lake Victoria Cichlids Improved the Pundamilia Genome Assembly

485   and Revealed a Major QTL for Sex-Determination. *G3* , *8*(7), 2411–2420.

486   https://doi.org/10.1534/g3.118.200207

487 Frei, D., De-Kayne, R., Selz, O. M., Seehausen, O., & Feulner, P. G. D. (2022). Genomic

488   variation from an extinct species is retained in the extant radiation following

489   speciation reversal. *Nature Ecology & Evolution*, *6*(4), 461–468.

490   https://doi.org/10.1038/s41559-022-01665-7

491 Goddard, T. D., Huang, C. C., Meng, E. C., Pettersen, E. F., Couch, G. S., Morris, J. H., &

492   Ferrin, T. E. (2018). UCSF ChimeraX: Meeting modern challenges in visualization

493   and analysis. *Protein Science: A Publication of the Protein Society*, *27*(1), 14–25.

494   https://doi.org/10.1002/pro.3235

495 Gopalakrishnan, S., Sinding, M.-H. S., Ramos-Madrigal, J., Niemann, J., Samaniego

496   Castruita, J. A., Vieira, F. G., Carøe, C., Montero, M. de M., Kuderna, L., Serres, A.,

497   González-Basallote, V. M., Liu, Y.-H., Wang, G.-D., Marques-Bonet, T., Mirarab, S.,

498   Fernandes, C., Gaubert, P., Koepfli, K.-P., Budd, J., … Gilbert, M. T. P. (2018).

499   Interspecific Gene Flow Shaped the Evolution of the Genus Canis. *Current Biology:*

500   *CB*, *28*(21), 3441-3449.e5. https://doi.org/10.1016/j.cub.2018.08.041

501 Hallgren, J., Tsirigos, K. D., Pedersen, M. D., Armenteros, J. J. A., Marcatili, P., Nielsen, H.,

502   Krogh, A., & Winther, O. (2022). DeepTMHMM predicts alpha and beta

503   transmembrane proteins using deep neural networks. In *bioRxiv* (p.

504   2022.04.08.487609). https://doi.org/10.1101/2022.04.08.487609

505 Irisarri, I., Singh, P., Koblmüller, S., Torres-Dowdall, J., Henning, F., Franchini, P., Fischer, C.,

Lemmon, A. R., Lemmon, E. M., Thallinger, G. G., Sturmbauer, C., & Meyer, A. (2018).

Phylogenomics uncovers early hybridization and adaptive loci shaping the radiation

of Lake Tanganyika cichlid fishes. *Nature Communications*, *9*(1), 3159.

https://doi.org/10.1038/s41467-018-05479-9

Jiménez, J., Doerr, S., Martínez-Rosell, G., Rose, A. S., & De Fabritiis, G. (2017). DeepSite:

protein-binding site predictor using 3D-convolutional neural networks. *Bioinformatics* ,

33(19), 3036–3042. https://doi.org/10.1093/bioinformatics/btx350

Johnson, T. C., Scholz, C. A., Talbot, M. R., Kelts, K., Ricketts, R. D., Ngobi, G., Beuning, K.,

Ssemmanda, I., I., & McGill, J. W. (1996). Late Pleistocene desiccation of lake

Victoria and rapid evolution of cichlid fishes. *Science (New York, N.Y.)*, *273*(5278),

1091–1093. https://doi.org/10.1126/science.273.5278.1091

Jumper, J., Evans, R., Pritzel, A., Green, T., Figurnov, M., Ronneberger, O., Tunyasuvunakool,

K., Bates, R., Žídek, A., Potapenko, A., Bridgland, A., Meyer, C., Kohl, S. A. A., Ballard,

A. J., Cowie, A., Romera-Paredes, B., Nikolov, S., Jain, R., Adler, J., … Hassabis, D.

(2021). Highly accurate protein structure prediction with AlphaFold. *Nature*,

596(7873), 583–589. https://doi.org/10.1038/s41586-021-03819-2

Kato, S., Arakaki, S., Nagano, A. J., Kikuchi, K., & Hirase, S. (2023). Genomic landscape of

introgression from the ghost lineage in a gobiid fish uncovers the generality of forces

shaping hybrid genomes. *Molecular Ecology*. https://doi.org/10.1111/mec.17216

Katoh, K., & Standley, D. M. (2013). MAFFT multiple sequence alignment software version

7: improvements in performance and usability. *Molecular Biology and Evolution*,

30(4), 772–780. https://doi.org/10.1093/molbev/mst010

Kawamura, R., & Nikaido, M. (2022). c-fos expression in the olfactory epithelium of the East

African cichlid (Haplochromis chilotes) in response to odorant exposure. *The Journal*

of Experimental Biology. https://doi.org/10.1242/jeb.244737

Keller-Costa, T., Canário, A. V. M., & Hubbard, P. C. (2015). Chemical communication in

cichlids: A mini-review. *General and Comparative Endocrinology*, *221*, 64–74.

https://doi.org/10.1016/j.ygcen.2015.01.001

Keller-Costa, T., Hubbard, P. C., Paetz, C., Nakamura, Y., da Silva, J. P., Rato, A., Barata, E.

N., Schneider, B., & Canario, A. V. M. (2014). Identity of a tilapia pheromone released

by dominant males that primes females for reproduction. *Current Biology: CB*, *24*(18),

2130–2135. https://doi.org/10.1016/j.cub.2014.07.049

Kocher, T. D. (2004). Adaptive evolution and explosive speciation: the cichlid fish model.

Nature Reviews. Genetics, *5*(4), 288–298. https://doi.org/10.1038/nrg1316

Kochnev, Y., Hellemann, E., Cassidy, K. C., & Durrant, J. D. (2020). Webina: an open-source

library and web app that runs AutoDock Vina entirely in the web browser.

Bioinformatics , *36*(16), 4513–4515. https://doi.org/10.1093/bioinformatics/btaa579

Kozlov, A. M., Darriba, D., Flouri, T., Morel, B., & Stamatakis, A. (2019). RAxML-NG: a fast,

scalable and user-friendly tool for maximum likelihood phylogenetic inference.

Bioinformatics , *35*(21), 4453–4455. https://doi.org/10.1093/bioinformatics/btz305

Larsson, A. (2014). AliView: a fast and lightweight alignment viewer and editor for large

datasets. *Bioinformatics (Oxford, England)*, *30*(22), 3276–3278.

https://doi.org/10.1093/bioinformatics/btu531

Letunic, I., & Bork, P. (2024). Interactive Tree of Life (iTOL) v6: recent updates to the

phylogenetic tree display and annotation tool. *Nucleic Acids Research*, *52*(W1),

W78–W82. https://doi.org/10.1093/nar/gkae268

Loh, Y.-H. E., Bezault, E., Muenzel, F. M., Roberts, R. B., Swofford, R., Barluenga, M., Kidd,

C. E., Howe, A. E., Di Palma, F., Lindblad-Toh, K., Hey, J., Seehausen, O., Salzburger,

W., Kocher, T. D., & Streelman, J. T. (2013). Origins of shared genetic variation in

African cichlids. *Molecular Biology and Evolution*, *30*(4), 906–917.

https://doi.org/10.1093/molbev/mss326

Malinsky, M., Svardal, H., Tyers, A. M., Miska, E. A., Genner, M. J., Turner, G. F., & Durbin,

R. (2018). Whole-genome sequences of Malawi cichlids reveal multiple radiations

interconnected by gene flow. *Nature Ecology & Evolution*, *2*(12), 1940–1955.

https://doi.org/10.1038/s41559-018-0717-x

Matschiner, M., Böhne, A., Ronco, F., & Salzburger, W. (2020). The genomic timeline of

cichlid fish diversification across continents. *Nature Communications*, *11*(1), 5895.

https://doi.org/10.1038/s41467-020-17827-9

McGee, M. D., Borstein, S. R., Meier, J. I., Marques, D. A., Mwaiko, S., Taabu, A., Kishe, M.

A., O'Meara, B., Bruggmann, R., Excoffier, L., & Seehausen, O. (2020). The

ecological and genomic basis of explosive adaptive radiation. *Nature*, *586*(7827), 75–

79. https://doi.org/10.1038/s41586-020-2652-7

McKenna, A., Hanna, M., Banks, E., Sivachenko, A., Cibulskis, K., Kernytsky, A., Garimella,

K., Altshuler, D., Gabriel, S., Daly, M., & DePristo, M. A. (2010). The Genome Analysis

Toolkit: a MapReduce framework for analyzing next-generation DNA sequencing data.

Genome Research, *20*(9), 1297–1303. https://doi.org/10.1101/gr.107524.110

Meier, J. I., Marques, D. A., Mwaiko, S., Wagner, C. E., Excoffier, L., & Seehausen, O. (2017).

Ancient hybridization fuels rapid cichlid fish adaptive radiations. *Nature*

Communications, *8*, 14363. https://doi.org/10.1038/ncomms14363

Nakamura, H., Aibara, M., Kajitani, R., Mrosso, H. D. J., Mzighani, S. I., Toyoda, A., Itoh, T.,

Okada, N., & Nikaido, M. (2021). Genomic Signatures for Species-Specific

Adaptation in Lake Victoria Cichlids Derived from Large-Scale Standing Genetic

Variation. *Molecular Biology and Evolution*, *38*(8), 3111–3125.

https://doi.org/10.1093/molbev/msab084

Nakamura, H., Aibara, M., & Nikaido, M. (2023). Ancient standing genetic variation facilitated

the adaptive radiation of Lake Victoria cichlids. *Genes & Genetic Systems*.

https://doi.org/10.1266/ggs.23-00024

Nicoli, A., Dunkel, A., Giorgino, T., de Graaf, C., & Di Pizio, A. (2022). Classification Model

for the Second Extracellular Loop of Class A GPCRs. *Journal of Chemical Information*

and Modeling, *62*(3), 511–522. https://doi.org/10.1021/acs.jcim.1c01056

Nikaido, M., Ota, T., Hirata, T., Suzuki, H., Satta, Y., Aibara, M., Mzighani, S. I., Sturmbauer,

C., Hagino-Yamagishi, K., & Okada, N. (2014). Multiple episodic evolution events in

V1R receptor genes of East-African cichlids. *Genome Biology and Evolution*, *6*(5),

1135–1144. https://doi.org/10.1093/gbe/evu086

Ota, T., Nikaido, M., Suzuki, H., Hagino-Yamagishi, K., & Okada, N. (2012). Characterization

of V1R receptor (ora) genes in Lake Victoria cichlids. *Gene*, *499*(2), 273–279.

https://doi.org/10.1016/j.gene.2012.03.002

Palkopoulou, E., Lipson, M., Mallick, S., Nielsen, S., Rohland, N., Baleka, S., Karpinski, E.,

Ivancevic, A. M., To, T.-H., Kortschak, R. D., Raison, J. M., Qu, Z., Chin, T.-J., Alt, K.

W., Claesson, S., Dalén, L., MacPhee, R. D. E., Meller, H., Roca, A. L., … Reich, D.

(2018). A comprehensive genomic history of extinct and living elephants.

Proceedings of the National Academy of Sciences of the United States of America,

115(11), E2566–E2574. https://doi.org/10.1073/pnas.1720554115

Pfister, P., & Rodriguez, I. (2005). Olfactory expression of a single and highly variable V1r

pheromone receptor-like gene in fish species. *Proceedings of the National Academy*

of Sciences of the United States of America, *102*(15), 5489–5494.

https://doi.org/10.1073/pnas.0402581102

Plenderleith, M., van Oosterhout, C., Robinson, R. L., & Turner, G. F. (2005). Female

preference for conspecific males based on olfactory cues in a Lake Malawi cichlid

fish. *Biology Letters*, *1*(4), 411–414. https://doi.org/10.1098/rsbl.2005.0355

Policarpo, M., Bemis, K. E., Laurenti, P., Legendre, L., Sandoz, J.-C., Rétaux, S., & Casane,

D. (2022). Coevolution of the olfactory organ and its receptor repertoire in ray-finned

fishes. *BMC Biology*, *20*(1), 195. https://doi.org/10.1186/s12915-022-01397-x

Rabosky, D. L., Chang, J., Title, P. O., Cowman, P. F., Sallan, L., Friedman, M., Kaschner, K.,

Garilao, C., Near, T. J., Coll, M., & Alfaro, M. E. (2018). An inverse latitudinal gradient

in   speciation   rate   for   marine   fishes.   *Nature*,   *559*(7714),   392–395.

https://doi.org/10.1038/s41586-018-0273-1

Rometsch, S. J., Torres-Dowdall, J., & Meyer, A. (2020). Evolutionary dynamics of pre- and

postzygotic reproductive isolation in cichlid fishes. *Philosophical Transactions of the*

Royal Society of London. Series B, Biological Sciences, *375*(1806), 20190535.

https://doi.org/10.1098/rstb.2019.0535

Ronco, F., Matschiner, M., Böhne, A., Boila, A., Büscher, H. H., El Taher, A., Indermaur, A.,

Malinsky, M., Ricci, V., Kahmen, A., Jentoft, S., & Salzburger, W. (2021). Drivers and

dynamics of a massive adaptive radiation in cichlid fishes. *Nature*, *589*(7840), 76–81.

https://doi.org/10.1038/s41586-020-2930-4

Salzburger, W. (2018). Understanding explosive diversification through cichlid fish genomics.

Nature Reviews. Genetics, *19*(11), 705–717. https://doi.org/10.1038/s41576-018-

0043-9

Salzburger, W., Van Bocxlaer, B., & Cohen, A. S. (2014). Ecology and Evolution of the African

Great Lakes and Their Faunas. *Annual Review of Ecology, Evolution, and*

Systematics, *45*(1), 519–545. https://doi.org/10.1146/annurev-ecolsys-120213-

091804

Santos, M. E., Lopes, J. F., & Kratochwil, C. F. (2023). East African cichlid fishes. *EvoDevo*,

14(1), 1. https://doi.org/10.1186/s13227-022-00205-5

Saraiva, L. R., & Korsching, S. I. (2007). A novel olfactory receptor gene family in teleost fish.

Genome Research, *17*(10), 1448–1457. https://doi.org/10.1101/gr.6553207

Seehausen, O., Terai, Y., Magalhaes, I. S., Carleton, K. L., Mrosso, H. D. J., Miyagi, R., van

der Sluijs, I., Schneider, M. V., Maan, M. E., Tachida, H., Imai, H., & Okada, N. (2008).

Speciation through sensory drive in cichlid fish. *Nature*, *455*(7213), 620–626.

https://doi.org/10.1038/nature07285

Seehausen, O., & Wagner, C. E. (2014). Speciation in Freshwater Fishes. *Annual Review of*

Ecology, Evolution, and Systematics, *45*(1), 621–651.

https://doi.org/10.1146/annurev-ecolsys-120213-091818

Suyama, M., Torrents, D., & Bork, P. (2006). PAL2NAL: robust conversion of protein

sequence alignments into the corresponding codon alignments. *Nucleic Acids*

Research, *34*(Web Server issue), W609-12. https://doi.org/10.1093/nar/gkl315

Svardal, H., Salzburger, W., & Malinsky, M. (2021). Genetic Variation and Hybridization in

Evolutionary Radiations of Cichlid Fishes. *Annual Review of Animal Biosciences*, *9*,

55–79. https://doi.org/10.1146/annurev-animal-061220-023129

Terai, Y., Seehausen, O., Sasaki, T., Takahashi, K., Mizoiri, S., Sugawara, T., Sato, T.,

Watanabe, M., Konijnendijk, N., Mrosso, H. D. J., Tachida, H., Imai, H., Shichida, Y.,

& Okada, N. (2006). Divergent selection on opsins drives incipient speciation in Lake

Victoria cichlids. *PLoS Biology*, *4*(12), e433.

https://doi.org/10.1371/journal.pbio.0040433

Torrens-Fontanals, M., Tourlas, P., Doerr, S., & De Fabritiis, G. (2024). PlayMolecule Viewer:

A toolkit for the visualization of molecules and other data. *Journal of Chemical*

Information and Modeling, *64*(3), 584–589. https://doi.org/10.1021/acs.jcim.3c01776

Tumescheit, C., Firth, A. E., & Brown, K. (2022). CIAlign: A highly customisable command

line tool to clean, interpret and visualise multiple sequence alignments. *PeerJ*, *10*,

e12983. https://doi.org/10.7717/peerj.12983

Vasimuddin, M., Misra, S., Li, H., & Aluru, S. (2019). Efficient Architecture-Aware Acceleration

of BWA-MEM for Multicore Systems. *2019 IEEE International Parallel and Distributed*

Processing Symposium (IPDPS), 314–324.

https://doi.org/10.1109/IPDPS.2019.00041

Verzijden, M. N., & ten Cate, C. (2007). Early learning influences species assortative mating

preferences in Lake Victoria cichlid fish. *Biology Letters*, *3*(2), 134–136.

https://doi.org/10.1098/rsbl.2006.0601

Villanueva, R. A. M., & Chen, Z. J. (2019). ggplot2: Elegant Graphics for Data Analysis (2nd

ed.). *Measurement: Interdisciplinary Research and Perspectives*, *17*(3), 160–167.

https://doi.org/10.1080/15366367.2019.1565254

Wang, L.-G., Lam, T. T.-Y., Xu, S., Dai, Z., Zhou, L., Feng, T., Guo, P., Dunn, C. W., Jones,

B. R., Bradley, T., Zhu, H., Guan, Y., Jiang, Y., & Yu, G. (2020). Treeio: An R package

for phylogenetic tree input and output with richly annotated and associated data.

Molecular Biology and Evolution, *37*(2), 599–603.

https://doi.org/10.1093/molbev/msz240

Weiss, J. D., Cotterill, F. P. D., & Schliewen, U. K. (2015). Lake Tanganyika--a "melting pot"

of ancient and young cichlid lineages (Teleostei: Cichlidae)? *PloS One*, *10*(4),

e0125043. https://doi.org/10.1371/journal.pone.0125043

Wickham H, François R, Henry L, Müller K, Vaughan D. (2023). *dplyr: A Grammar of Data*

Manipulation (1.1.4) [Computer software]. https://github.com/tidyverse/dplyr,

https://dplyr.tidyverse.org

Wickham H, Hester J, Bryan J. (2024). *readr: Read Rectangular Text Data* (2.1.5) [Computer

software]. https://github.com/tidyverse/readr, https://readr.tidyverse.org

Xu, W., Wu, L., Liu, S., Liu, X., Cao, X., Zhou, C., Zhang, J., Fu, Y., Guo, Y., Wu, Y., Tan, Q.,

Wang, L., Liu, J., Jiang, L., Fan, Z., Pei, Y., Yu, J., Cheng, J., Zhao, S., … Hua, T.

(2022). Structural basis for strychnine activation of human bitter taste receptor

TAS2R46. *Science*, *377*(6612), 1298–1304. https://doi.org/10.1126/science.abo1633

Yang, Z. (2007). PAML 4: phylogenetic analysis by maximum likelihood. *Molecular Biology*

and Evolution, *24*(8), 1586–1591. https://doi.org/10.1093/molbev/msm088

Yu, G., Smith, D. K., Zhu, H., Guan, Y., & Lam, T. T.-Y. (2017). Ggtree: An r package for

visualization and annotation of phylogenetic trees with their covariates and other

associated data. *Methods in Ecology and Evolution*, *8*(1), 28–36.

https://doi.org/10.1111/2041-210x.12628

Zapilko, V., & Korsching, S. I. (2016). Tetrapod V1R-like ora genes in an early-diverging ray-

finned fish species: the canonical six ora gene repertoire of teleost fish resulted from

gene loss in a larger ancestral repertoire. *BMC Genomics*, *17*, 83.

https://doi.org/10.1186/s12864-016-2399-6

Zhang, D., Tang, L., Cheng, Y., Hao, Y., Xiong, Y., Song, G., Qu, Y., Rheindt, F. E., Alström,

P., Jia, C., & Lei, F. (2019). "Ghost Introgression" As a Cause of Deep Mitochondrial

Divergence in a Bird Species Complex. *Molecular Biology and Evolution*, *36*(11),

2375–2386. https://doi.org/10.1093/molbev/msz170

Zhou, L., Feng, T., Xu, S., Gao, F., Lam, T. T., Wang, Q., Wu, T., Huang, H., Zhan, L., Li, L.,

Guan, Y., Dai, Z., & Yu, G. (2022). Ggmsa: A visual exploration tool for multiple

sequence alignment and associated data. *Briefings in Bioinformatics*, *23*(4).

https://doi.org/10.1093/bib/bbac222

Author Contributions

M.N. supervised the project. S.T. conducted all bioinformatics and evolutionary analyses. S.T.

and M.N. wrote the manuscript. Z.Z., M.A., and T.N. provided their expertise and commented

on the paper. All authors read and approved the final manuscript.

Tables and Figures

Figure 1 A phylogenetic tree and sequence alignment of the V1R2 gene in cichlids. Bootstrap

values are shown only for nodes with support values of 60 or higher. The sequences

are divided into two highly divergent alleles (clade I and clade II). Only diagnostic

amino acid sites between clade I and clade II were shown. Dots indicate identity with

the top sequence (clade I). The phase of each heterozygous allele in the individuals

is indicated in parentheses. Observed recombination-derived sequences, which are

exceptionally rare, have been excluded. Note that sequences from each lake

(Tanganyika, Malawi, Victoria) do not form monophyletic groups.

Figure 2 Frequencies of each allele of the V1R2 gene in each cichlid lineage. We calculated

the frequencies of clade I and clade II alleles in each lineage. Clade I are shown in

dark blue, clade II in yellow, heterozygotes in pink, and recombinants in grey. The

phylogenetic tree and divergence times are based on previous studies (Matschiner

et al., 2020; Meier et al., 2017; Nakamura et al., 2023; Ronco et al., 2021). Numbers

in parentheses next to lineage names show the number of samples. A star represents

the estimated times of acquisition of clade II.

Figure 3 Recombination events observed in the V1R6 gene of cichlids. (A) Phylogenetic tree of the V1R6 gene. Bootstrap values are shown only for nodes with support values of 60 or higher. (B) All haplotypes of the V1R6 gene. Only sites that are polymorphic in Lake Victoria lineage are displayed. Numbers in parentheses following lineage names show the number of samples. Asterisks above site numbers denote nonsynonymous substitutions. Ancestral and derived alleles at each site were defined based on the frequency of nucleotides across the sequences. Sites matching the ancestral allele are highlighted in dark blue, and those matching the derived allele are highlighted in yellow. (C) Schematic diagram of recombination.

Figure 4 Two highly divergent alleles observed in the V1R1 gene of Tropheini. (A) A

phylogenetic tree of the V1R1 gene. Bootstrap values are shown only for nodes with

support values of 60 or higher. (B) Frequencies of each allele of the V1R1 gene in

each cichlid lineage. Clade I are shown in dark blue, clade II in yellow, heterozygotes

in pink, and recombinants in grey. The phylogenetic tree and divergence times are

based on previous studies (Matschiner et al., 2020; Meier et al., 2017; Nakamura et

al., 2023; Ronco et al., 2021). Numbers in parentheses next to the lineage names

show the number of samples. A star represents the estimated timing of acquisition of

clade II.

Figure 5 3D structure of cichlid V1R2. (A) 3D structure of V1R2 viewed from the extracellular

side. Amino acids that differ between Clade I and Clade II are shown in dark blue.

The ligand used for analysis (4HPAA) is shown in yellow. (B) Predicted

transmembrane regions of cichlid V1R2. The transmembrane regions were predicted

using the sequence of Clade I. Dots indicate amino acids identical to those in Clade

I. The extracellular regions are shown in pink, transmembrane regions in blue, and

intracellular regions in yellow. TM: transmembrane; ECL: extracellular loop; ICL:

intracellular loop. Gray dots indicate predicted ligand-binding sites. Red squares

indicate ligand-binding sites that are also sites of substitution between alleles.

Mutations were found to occur at ligand-binding sites significantly more often than

expected by chance (Fisher's exact test: p = 0.0389).

Figure 6 Possible involvement of ghost introgressions in cichlids (A) Sequence diversity of

V1R genes in cichlids. Allele types of V1R genes are represented by colored dots for

each lineage. The phylogenetic tree and divergence times are based on previous

studies (Matschiner et al., 2020; Meier et al., 2017; Nakamura et al., 2023; Ronco et

al., 2021). Note that the derived allele (clade II) of V1R2 is shared across lineages,

whereas the derived sequences of V1R3 and V1R6 differ among lineages (asterisks).

(B) Predicted mechanisms of allele acquisition for V1R1, V1R2, and V1R6 genes.

Arrows indicate the most parsimonious estimated times of introgression.

Figure S1 The lineage names in this study were defined with reference to previous studies

(Matschiner et al., 2020; Meier et al., 2017; Nakamura et al., 2023; Ronco et al.,

2021). Haplochromini species that diverged before Tropheini are defined as "ancient

Haplochromini." Divergence times are expressed in millions of years (Mya).

```
*nonsynonymous = 13, synonymous = 2

 46 ATC (I) -> ATT (I)
 62 TGT (C) -> AGT (S)
 65 CAG (Q) -> CCG (P)
 66 ACA (T) -> TCA (S)
145 GCA (A) -> TCA (S)
146 GCT (A) -> ACT (T)
159 CCT (P) -> TCT (S)
164 AAC (N) -> GAC (D)
169 CAT (H) -> CTT (L)
185 GTC (V) -> ATC (I)
214 GTG (V) -> ATG (M)
218 CGT (R) -> CAT (H)
232 ACA (T) -> GCA (A)
247 GAT (D) -> GAA (E)
285 TCC (S) -> TCT (S)
```

Figure S2 A phylogenetic tree and sequence alignment of the V1R2 gene in cichlids (Dataset

1). The phylogenetic tree displays bootstrap values only for nodes with support

values of 60 or higher. The left shows nonsynonymous and synonymous substitution

sites at positions marked with an asterisk.

Figure S3 A phylogenetic tree and sequence alignment of the V1R2 gene in cichlids (Dataset

2). The phylogenetic tree displays bootstrap values only for nodes with support

values of 60 or higher. The left shows nonsynonymous and synonymous substitution

sites at positions marked with an asterisk.

```
*nonsynonymous = 14, synonymous = 5

 20 ATT (I) -> GTT (V)
 31 ACC (T) -> ACA (T)
 46 ATC (I) -> ATT (I)
 62 TGT (C) -> AGT (S)
 65 CAG (Q) -> CCG (P)
 66 ACA (T) -> TCA (S)
145 GCA (A) -> TCA (S)
146 GCT (A) -> ACT (T)
159 CCT (P) -> TCT (S)
164 AAC (N) -> GAC (D)
169 CAT (H) -> CTT (L)
170 GTG (V) -> GTA (V)
185 GTC (V) -> ATC (I)
214 GTG (V) -> ATG (M)
218 CGT (R) -> CAT (H)
232 ACA (T) -> GCA (A)
247 GAT (D) -> GAA (E)
274 TGC (C) -> TGT (C)
285 TCC (S) -> TCT (S)
```

Figure S4 A phylogenetic tree and sequence alignment of the V1R2 gene in cichlids (Dataset

3). The phylogenetic tree displays bootstrap values only for nodes with support

values of 60 or higher. The left shows nonsynonymous and synonymous substitution

sites at positions marked with an asterisk.

Figure S5 A phylogenetic tree and sequence alignment of the V1R2 gene in cichlids (Dataset

4). The phylogenetic tree displays bootstrap values only for nodes with support

values of 60 or higher. The left shows nonsynonymous and synonymous substitution

sites at positions marked with an asterisk.

Tree scale: 0.01

*nonsynonymous = 14, synonymous = 2

```
 38 TCG (S) -> CCG (P)
 46 ATC (I) -> ATT (I)
 62 TGT (C) -> AGT (S)
 65 CAG (Q) -> CCG (P)
 66 ACA (T) -> TCA (S)
 99 ACG (T) -> ACC (T)
145 GCA (A) -> TCA (S)
146 GCT (A) -> ACT (T)
156 GAC (D) -> GGC (G)
159 CCT (P) -> TCT (S)
169 CAT (H) -> CTT (L)
185 GTC (V) -> ATC (I)
214 GTG (V) -> ATG (M)
218 CGT (R) -> CAT (H)
232 ACA (T) -> GCA (A)
247 GAT (D) -> GAA (E)
```

Figure S6 A phylogenetic tree and sequence alignment of the V1R2 gene in cichlids (Dataset

5). The phylogenetic tree displays bootstrap values only for nodes with support

values of 60 or higher. The left shows nonsynonymous and synonymous substitution

sites at positions marked with an asterisk.

Figure S7 V1R2 alignment of 102 ray-finned fish species. The alignment displays sites

corresponding to the two alleles (clade I and clade II). Sites matching clade I are

highlighted in dark blue, those matching clade II are highlighted in yellow, and all

other sites are shown in grey.

Figure S8 Recombination rate (cM/Mb) per chromosome in *Pundamilia nyererei* from the

Lake Victoria lineage (Feulner et al., 2018). The positions of V1R genes are indicated

by yellow lines.

Figure S9 A phylogenetic tree of the V1R1 gene in Tropheini. Bootstrap values are shown

only for nodes with support values of 60 or higher. The sequences are divided into

two highly divergent alleles (clade I and clade II). Recombination-derived sequences

have been excluded.

Figure S10 V1R1 alignment of 102 ray-finned fish species. The alignment displays sites

corresponding to the two alleles (clade I and clade II). Sites matching clade I are

highlighted in dark blue, those matching clade II are highlighted in yellow, and all

other sites are shown in grey.

Figure S11 A phylogenetic tree of the V1R3 gene in cichlids. Nodes are marked with dots

based on bootstrap values: black for values of 90 or higher, grey for 80–89, and white

for 70–79. The inner circle represents tribes, while the outer circle indicates habitats.

Branches under positive selection are highlighted in red (p=0.0361, see Table S2).

The number of nonsynonymous and synonymous substitutions for the branch is

indicated as ns:s.

Figure S12 A phylogenetic tree and sequence alignment of the V1R3 gene in Trematocarini.

Bootstrap values are shown only for nodes with support values of 60 or higher.

Trematocara unimaculatum, which retains the ancestral sequence of V1R3, differ

significantly from those of other species. The amino acid alignment displays only the

sites with variations observed within Trematocarini. Dots indicate identity with the top

sequence. Positive selection was not detected in the red branch (p=0.0503, see Table

S2). The number of nonsynonymous and synonymous substitutions in the red branch

is indicated as ns:s.

Figure S13 A phylogenetic tree of the V1R6 gene in cichlids. Nodes are marked with dots

based on bootstrap values: black for values of 90 or higher, grey for 80–89, and white

for 70–79. The inner circle represents tribes, while the outer circle indicates habitats.

Branches under positive selection are highlighted in red (see Table S2).

Figure S14 3D structure of cichlid V1R1. (A) 3D structure of V1R1 viewed from the

extracellular side. Amino acids that differ between Clade I and Clade II are shown in

dark blue. The ligand used for analysis (LCA) is shown in yellow. (B) Predicted

transmembrane regions of cichlid V1R1. The transmembrane regions were predicted

using the sequence of Clade I. Dots indicate amino acids identical to those in Clade

I. The extracellular regions are shown in pink, transmembrane regions in blue, and

intracellular regions in yellow. TM: transmembrane; ECL: extracellular loop; ICL:

intracellular loop. Gray dots indicate predicted ligand-binding sites. Red squares

indicate ligand-binding sites that are also sites of substitution between alleles.

(Fisher's exact test: p = 0.0922)

865 Figure S15 3D structure of cichlid V1R6. (A) 3D structure of V1R6 viewed from the

866 extracellular side. Amino acids that differ between ancestral allele and derived allele

867 are shown in dark blue. The ligand used for analysis (LCA) is shown in yellow. (B)

868 Predicted transmembrane regions of cichlid V1R6. The transmembrane regions were

869 predicted using the sequence of ancestral allele. Dots indicate amino acids identical

870 to those in ancestral allele. The extracellular regions are shown in pink,

871 transmembrane regions in blue, and intracellular regions in yellow. TM:

872 transmembrane; ECL: extracellular loop; ICL: intracellular loop. Gray dots indicate

873 predicted ligand-binding sites.

Figure S16 Sequence similarity and dN/dS ratios of V1R genes among 102 ray-finned fish

species (A) Amino acid similarity of V1R genes. The average sequence similarity was

calculated for all pairwise combinations of the 102 species. Error bars represent

standard errors. (B) dN/dS ratios of V1R genes. The average dN/dS ratio was

calculated for all pairwise combinations of the 102 species. Error bars represent

standard errors.

Table S1 We used short-read data from 28 tribes, 528 species, and 908 samples of cichlids

registered in the NCBI Sequence Read Archive (SRA).

Table S2 The results of selection analyses and likelihood ratio tests.

V1R2

Model	np	lnL	p-value	omega
Null	66	-3334.240925		0.34224
Alternative	67	-3328.619231	0.000799	2.97063

V1R1

Model	np	lnL	p-value	omega
Null	61	-2086.808159		0.86632
Alternative	62	-2082.84435	0.004869	537.52213

V1R3

Model	np	lnL	p-value	omega
Null	208	-4602.837321		0.3905
Alternative	209	-4600.641708	0.036124	2.88326

V1R3 Trematocarini

Model	np	lnL	p-value	omega
Null	26	-1819.23301		0.50765
Alternative	27	-1817.317734	0.050326	2.13329

V1R6 Ectodini

Model	np	lnL	p-value	omega
Null	182	-5557.85997		0.54623
Alternative	183	-5553.243731	0.002378	2.6782

V1R6 Trematocarini

Model	np	lnL	p-value	omega
Null	182	-5557.85997		0.54591
Alternative	183	-5554.585293	0.010492	1.60894