Sex-specific mate choice and mate change based on MHC-IIB dissimilarity in a genetically monogamous seabird

Lucie Thel[1,2], Maxime Pineaux[1], Léa Ribeiro[1], Etienne Danchin[3,1], Scott A. Hatch[4], Pierrick Blanchard[1], Sarah Leclaire[1]

Authors details

[1] Centre de Recherche pour la Biodiversité et l'Environnement, UMR 5174 (CNRS, Université Paul Sabatier), 118 route de Narbonne, 31062 Toulouse, France.

[2] Department of Conservation Management, Faculty of Science, Nelson Mandela University, George, South Africa.

[3] Centre de Recherches sur la Cognition Animale, Centre de Biologie Intégrative, UMR 5169 (CNRS, Université de Toulouse Midi-Pyrénées), Toulouse, France.

[4] Institute for Seabird Research and Conservation, Anchorage, Alaska, USA.

Corresponding authors: Lucie Thel, lucie.mc.thel@gmail.com; Sarah Leclaire, sarah.leclaire@univ-tlse3.fr

Acknowledgments

We thank the Middleton fieldworkers who collected data, as well as the undergraduate students who helped with molecular analyses.

Authors' contributions

Lucie Thel: Conceptualization - Methodology - Formal analysis - Investigation - Writing: Original Draft - Writing: Review & Editing - Visualisation.

Maxime Pineaux: Data acquisition - Conceptualisation - Writing: Review & Editing.

Léa Ribeiro: Data acquisition.

Etienne Danchin: Writing: Review & Editing - Project administration - Funding acquisition.

Scott Hatch: Supervision - Project administration - Funding acquisition.

Pierrick Blanchard: Conceptualisation - Data Curation - Writing : Review & Editing.

Sarah Leclaire: Conceptualization - Methodology - Formal analysis - Resources - Data Curation - Writing: Original Draft - Writing: Review & Editing - Visualisation - Supervision - Project administration - Funding acquisition.

Ethics

The experiment was conducted under the approval of the U.S. Geological Survey Alaska Science Center, the University of Alaska Institutional Animal Care and Use Committees, and the Institut Paul-Emile Victor Ethical Committee, in accordance with United States laws and under permits from the U.S. Fish and Wildlife Service and the State of Alaska.

Conflict of interest

The authors declare that they have no conflict of interest.

Funding

This work was supported by the French Polar Institute Paul-Emile Victor (IPEV "Programme 1162" to ED and SL) and the Fondation Fyssen (to SL). This study is part of the "Laboratoire d'Excellence" (LABEX) entitled TULIP (ANR-10-LABX-41). LR was supported by an EUR TULIP Ph.D. grant obtained through SEVAB postgraduate school.

Data availability statement

Data available from the Zenodo Repository: https://doi.org/10.5281/zenodo.14523780.

Sex-specific mate choice and mate change based on MHC-IIB dissimilarity in a genetically monogamous seabird

Abstract

Among species reproducing sexually, mating strategies represent a major component of individual fitness. The Major Histocompatibility Complex (MHC) is an extremely diverse set of genes responsible for immunological recognition and defense against pathogens. Although MHC-dissimilarity between mates has been proposed to drive mate choice through increased offspring pathogen resistance, evidence is mixed, and evidence for other MHC-based mating strategies remains rare. We investigated whether MHC-IIB dissimilarity is associated with mate choice and mate change in the genetically monogamous black-legged kittiwake (*Rissa tridactyla*). We found that first-time male breeders are paired with females more MHC-IIB dissimilar to them than expected by chance. After a successful reproduction, the more males are MHC-IIB dissimilar to their mate, the less likely they are to change mate. When males change their mate, they tend to be more MHC-IIB dissimilar to the new mate compared to the previous mate. However, males are not more MHC-IIB dissimilar to their new mate than expected by chance. We did not find evidence for an effect of MHC-IIB dissimilarity on mating strategies when considering the female's perspective. Male mate choice should not be overlooked in species with bi-parental care in which the male performs a significant part of parental care.

Keywords: Black-legged kittiwake, *Rissa tridactyla*, Divorce, Reproduction, Major Histocompatibility Complex, Disassortative mate choice

Introduction

Mating strategies are the set of behavioural patterns used by individuals to attract, select, change or retain mates to promote individual fitness. Understanding their driving factors has long been a major topic in evolutionary biology ([1]; [2]; [3]; [4]). For instance, mate choice has repeatedly been shown to rely on traits that honestly reflect the quantity or quality of resources, the care or the genes that the mate can provide ([5]; [6]; [7]). In addition, in several species, individuals preferentially choose mates who are more genetically or behaviourally compatible to avoid inbreeding or to enhance behavioural coordination or task complementation between mates ([8]; [9]; [10]; [11]; [12]). However, individuals may face energetic, time or spatial constraints in choosing mates ([13]). In such circumstances, individuals are forced to choose within a limited pool of mates and to potentially accept suboptimal mates. A possible response to this limitation in iteroparous species is divorce, defined as one breeding individual changing partner for a new mate while the previous mate is still alive ([14]). Theoretical studies have suggested that individuals should divorce when pairing with a higher-quality or more compatible mate increases their lifetime reproductive success [15]. Accordingly, divorce is common after a reproductive failure ([16]; [17]; [18]), and usually leads to increased reproductive success ([19]; [20]; [17]). Like mate choice, divorce may be associated with the quality of the ornaments of the mate or the genetic or behavioural compatibility of the pair (e.g. [21]; [22]; [23]).

Concerning the genetic component of the pool of heritable factors that can affect mating strategies, the major histocompatibility complex (MHC, called HLA in humans) has been the focus of much research ([24]; [25]; [26]). The MHC, one of the most polymorphic sets of genes in vertebrates, encodes cell-surface proteins that are essential for immunological recognition and defense against pathogens. Because each MHC molecule can bind a limited set of antigens ([27]), higher MHC-diversity increases the number of antigens recognized,

thereby providing resistance to a wider range of parasites ([28]; [29]). Accordingly, several

studies have found that higher MHC-diversity increases health, reproductive success and

survival ([30]; [31]; [32]; [33]; [34]). Because the MHC allelic composition of the offspring

results from the combination of those of the parents, numerous studies have found evidence

for MHC-based mate choice ([35]; [36]; [37]; [38]; [39]; [40]; [41]). Although there is

evidence that the parental MHC-composition also influences other reproductive strategies

(e.g. cryptic female choice, [42]; extra-pair copulation, [43]), to our knowledge, a single

study on Seychelle warblers (*Acrocephalus sechellensis*) explored how parental MHC-

similarity is associated with divorce rate and found no evidence for such an association

([44]).

Here we tested whether mate choice and mate change (defined as the probability to

observe an individual with a new mate, whether the previous mate is seen alive or not)

depend on dissimilarity at MHC class-II B (hereafter MHC-IIB) in the black-legged kittiwake

(*Rissa tridactyla*), a long-lived genetically monogamous seabird with no extra-pair copulation

([45]). In kittiwakes, divorce is quite common (i.e., 15% - 50% of the pairs), and more

frequent among first-time breeders and after reproductive failure ([46]; [47]; [48]). Coulson

([46]) suggested therefore that some kinds of "incompatibility" between mates results in

unsuccessful breeding, thereby providing costs that select for divorce. In black-legged

kittiwakes, MHC-IIB similarity between mates results in reduced offspring MHC-IIB

diversity ([49]), which incurs costs for female chicks in terms of slower growth, poor tick

clearance and lower survival ([50]). We therefore hypothesised that, in this species, mating

strategies based on MHC-IIB dissimilarity between mates may have evolved to maximise the

MHC-IIB diversity of the offspring.

First, we expected individuals to preferentially choose an MHC-IIB dissimilar mate.

In several seabirds, including the black-legged kittiwake, young or inexperienced individuals

arrive later to the breeding ground than older or more experienced breeders, which may limit

the pool of potential partners ([51]; [52]; [53]). Inexperienced breeders may also be less

efficient in sampling mates and less able to make mating discrimination or to anticipate

alternative choices ([54]). For instance, in grey mouse lemurs (*Microcebus murinus*), young

females in their first mating season do not express mate choice, while experienced females

prefer MHC-IIB dissimilar males ([41]). The mate choice of first-time breeders may thus be

more constrained than that of experienced breeders. We therefore tested the existence of

MHC-IIB dependent mate choice in first-time breeders and experienced breeders separately.

Second, if mate change allows individuals paired with a suboptimal mate to increase the

MHC-IIB diversity of their future offspring, we expected the rate of mate change to increase

with MHC-IIB similarity between mates. Third, we expected that, after a mate change, the

new pair should be more MHC-IIB dissimilar than the previous one. In black-legged

kittiwakes, although both mates participate fairly equally in incubation and chick rearing,

sexes slightly differ in energetic investment in reproduction ([55]; [56]). We thus tested these

predictions separately in males and females to explore potential sex differences in mating

strategies.

Materials and methods

Study area and population

This study was conducted between 1996 and 2021 in a colony of black-legged kittiwakes

nesting in an abandoned U.S. Air Force radar tower on Middleton Island (59°26'N,

146°20'W), Gulf of Alaska, located approximately 100 km south of the mainland. Individuals

were tagged with a unique combination of coloured and metal bands for long-term

monitoring ([57]). About 400 nest sites were monitored on the tower each year. All these

sites were checked daily to record the identity of the individual(s) occupying the nests. For

each individual of the study, sex was determined, in order of priority, by: 1) molecular

assessment (n = 746 individuals, [58]); 2) behavioural observations (n = 413 individuals,

[59]); and 3) comparison of the relative size of both members of a pair, the largest bird being

considered male (n = 16 individuals). When only one individual of a pair was sexed, we

attributed the opposite sex to its mate (n = 112 individuals).

Like many populations monitored in the long term, our population has been subjected

to several treatments during the study period (e.g. supplemental feeding, brood manipulation,

feather clipping, use of anti-copulatory devices). These treatments can affect reproductive

success ([57]; [60]; [61]) and therefore pair stability. To limit any bias (e.g. artificial increase

of reproductive success, and thus decreased mate change probability due to supplemental

feeding), we removed pairs that were subjected to experiments from certain analyses (see

below).

We defined a pair as two individuals of opposite sex attending the same nest and

laying at least one egg during the reproductive season. Individuals usually remain and breed

in the same colony during their whole life ([46]). We thus defined the first mate choice as the

pair formed during the first observed breeding attempt. Each individual of the population was

involved in on average 4.57 ± 3.61 (SE) reproductive events (range = 1 - 21, Supporting

information 1) over the course of the study.

To study mate change, we focused on individuals that were observed with an

identified mate during at least one year and that re-mated with a new mate the next year,

whether its former mate was re-observed or not. In the studied population, the proportion of

pairs in which the two mates were both seen alive in the years after pair splitting was very

low (i.e. strict divorce; 6% of all pairs) compared to the proportion observed in other

populations of kittiwakes (25% in [46]; between 13 % and 47 % in [47]; 19.1 % on average

in [48]). On Middleton island, numerous kittiwakes nest on other man-made structures than

the tower ([57]). These sites were not monitored, except for occasional demographic surveys

for certain years. It therefore limited our ability to observe alive birds who changed nesting

sites, and thus to detect strict divorces. Therefore, we focused our analyses on mate change,

which was defined as the situation of an individual being seen with a different mate in year n

and n+1, whether the mate of the year n was observed or not in the studied population during

the year n+1 and subsequent. Our definition of mate change thus applies to a larger context

than strict divorce, and may include widowed birds. However, we also present analyses on

the probability of strict divorce in the supplementary material.

Molecular analysis

We focused on the MHC class II which exhibits stronger selection than the MHC class I in

non-passerines such as the black-legged kittiwake ([62]). Dissimilarity at MHC-IIB was

assessed following the methods described in Pineaux et al. ([50]; [63]). Briefly, DNA was

extracted from blood with the blood and tissue DNA extraction kit (Qiagen), and a 258-pb

fragment of exon 2 of MHC-IIB was amplified using specific primers (forward: 5′-

GCACGAGCAGGGTATTTCCA, and reverse: 5′-GTTCTGCCACACACTCACC).

Amplicons were then sequenced with an Illumina MiSeq platform, using the 2 x 300-bp

protocol (Fasteris SA, Switzerland). Amplicon sequences were analysed with ampliSaS, a

three-step pipeline that consists of read demultiplexing, unique sequence clustering and

erroneous sequence filtering (substitution errors, 1 %; indel errors, 0.001 %; minimum

frequency with respect to dominant, 33 %; minimum amplicon sequence frequency, 2.5 %,

[64]). We calculated the MHC-IIB functional distance between individuals based on amino

acid variations in the peptide-binding region (PBR) inferred from Leclaire et al. ([65]).

We described the chemical binding properties of each amino acid in the PBR with

Sandberg's five physicochemical descriptors (z-descriptors, [66]) and computed a Sandberg's

distance matrix between sequences using the DistCalc function of the MHCTools package in

R ([67]). Then, we calculated an UPGMA dendrogram of sequences (hclust function in R),

which represents clusters of functionally similar MHC-IIB sequences (hereafter,

"supertypes"). The MHC-IIB functional distance between a male and a female was estimated

using the tree distance (treedist function in the package vegan, [68]), which is found by

combining sequences in a common dendrogram and calculating how much of the dendrogram

height is shared and how much is unique. In addition, we calculated the difference in MHC-

IIB supertype sharing between mates by calculating the difference in supertype sharing

following the definition of Strandh et al. ([40]):

$$100 \times Nb_{supertypes\ pair} / (Nb_{supertypes\ female} + Nb_{supertypes\ male}),$$

where $Nb_{supertypes\ female}$ corresponds to the number of supertypes possessed by the female,

$Nb_{supertypes\ male}$ corresponds to the number of supertypes possessed by the male, and $Nb_{supertypes}$

$_{pair}$ corresponds to the total number of supertypes possessed by the female and the male

together (shared supertypes count as 1). This variable thus varies between 50 for individuals

sharing no supertype and 100 for individuals sharing all their supertypes. Overall, we

identified 15 supertypes, among which 3 were very common (possessed by more than 47 %

of the sequenced population), 6 were common, and 6 were rare (possessed by less than 7 %

of the sequenced population, Supporting information 2). The MHC-IIB functional distance

was highly correlated with the difference in MHC-IIB supertype sharing (Spearman

correlation test: $P < 0.01$, $\rho = 0.77$). We found similar results when studying MHC-IIB

functional distance and difference in MHC-IIB supertype sharing, except for one analysis

(i.e. the analysis on mate choice in first-time male breeders, see below).

Mate choice for MHC-IIB dissimilarity may not only be governed by evolutionary

processes that specifically target these genes, but may also be affected by mechanisms

operating at the level of the entire genome (e.g. inbreeding avoidance). To test whether

associations between MHC-IIB dissimilarity and mate choice might be driven by genome-

wide relatedness, we used RAD sequencing (n = 92 individuals) to estimate genome-wide

relatedness between individuals (Supporting information 3). We found that genome-wide

relatedness was weakly correlated with functional MHC-IIB similarity (Mantel test: r = 0.09,

P = 0.04), and not correlated with MHC-IIB supertype similarity (Mantel test: r = 0.04, P =

0.14). These associations were not significant when excluding the four individuals involved

in a first-degree family relationship (siblings or parents-chicks; Mantel tests: r = 0.05, P =

0.16 and r = 0.02, P = 0.33).

Statistical analysis

To test for potential differences between sexes, we conducted all analyses on males and

females separately (see sex-specific sample sizes and details of the analyses in Supporting

information 4). We performed all analyses using the R software (R Core Development Team

2023, version R 4.4.0). All datasets and code used in the analyses are available from the

Zenodo Repository (https://doi.org/10.5281/zenodo.14523780).

To test whether males and females selected a more MHC-IIB dissimilar mate than

under random mating, we used bootstrapping to generate random pairs, where we let each

individual choose randomly 10000 times between all available opposite-sex mates of the

same year, without replacement (i.e. one mate can be selected only once during a given year,

[69]). An individual was considered to be available for mating in year n if it was seen in the

colony during that year, unless it was observed with the same mate in both the previous year

n-1 and the current year n. We also ran the same analyses when considering all opposite-sex

mates of the same year, and found similar results (Supporting information 5). We calculated

p-values as the proportion of iterations in which the random MHC-IIB dissimilarity mean

was superior or equal to the observed MHC-IIB dissimilarity mean (one-tailed test). We

tested MHC-IIB based mate choice for: i) first-time breeders only (n = 487 males and n = 485

females), and ii) all the breeders of the population experiencing a mate change (n = 220

events in males and n = 233 events in females). Because of a lack of available mates of

known MHC-IIB composition to generate random re-matings for the females in 2004, we

excluded the 2004 data in the mate choice analysis (n = 13 reproductive events).

To test whether mate change probability (1 = mate change between the year n and

n+1, 0 = faithful pair between the year n and n+1) varied with the MHC-IIB dissimilarity of

the pair during the year n in males (n = 1104) and females (n = 1094), we used Generalised

Linear Mixed Models (GLMM) with a binomial distribution of error and a logit link (glmer

function of the lme4 package, [70]). We tested the effect of the MHC-IIB dissimilarity (fixed

effect) in two different models, after a reproductive success and after a reproductive failure.

Indeed, the MHC-IIB dissimilarity might influence the reproductive outcome, which is also

known to influence the probability of divorce in black-legged kittiwakes ([48]). We defined

the reproductive outcome as a binary variable recording the presence (i.e. "success") or

absence (i.e. "failure") of at least one chick fledged at the end of the reproductive season. In

the models, we set the nAGQ argument (number of points per axis for evaluating the adaptive

Gauss-Hermite approximation to the log-likelihood) of the glmer function to 0 to help for

convergence when necessary ([71]). We tested the effect of individual identity and year as

random intercepts with the ranef function of the lme4 package ([70]) and removed these

variables when they had no effect. We removed pairs that were subjected to experiments

from this analysis. As we used GLMM with binomial data, we provided the log odd (and its

standard error), which is equivalent to the coefficient of the slope in a linear model (i.e.

estimate in the model summary), i.e. the probability of an event in the log scale, which is the

transformed scale to satisfy the assumptions of a linear model for GLMM with binomial data.

When using mixed models, we calculated the confidence intervals on the fixed effects using

the predictInterval function of the merTools package (n iterations = 10000, [72]). We

conducted the same analysis when considering only the pairs for which both mates were seen

during the years n and n+1 (Supporting information 6). This dataset, although smaller than

the original one (n = 1022 events in males, and n = 1034 events in females) particularly

regarding the number of pair splitting (n = 50 (37 after breeding failure, 13 after breeding

success) versus 132 events in males, n = 62 (41 after breeding failure, 21 after breeding

success) versus 122 events in females), served as a refined dataset focusing on strict divorce.

However, as the individual monitoring focused mostly on individuals breeding in the tower,

this dataset might exclude individuals involved in a strict divorce but breeding elsewhere on

the island.

If individuals changed mate, we tested whether their new mate was more MHC-IIB

dissimilar to them compared to the previous one (n = 165 males, n = 179 females) using

Linear Models (LM). The response variable was the MHC-IIB dissimilarity gain, i.e. the

difference between the MHC-IIB dissimilarity to the previous mate and the MHC-IIB

dissimilarity to the new mate. Using the gain in a LM framework allowed us to estimate if the

difference between the MHC-IIB dissimilarity to the previous mate and the new mate was

significantly different from 0. Additionally, it allowed us to include individual and year as

random effects. Each individual possesses its own MHC-IIB composition, which can be rare

or common, and it is easier for an individual with a rare MHC-IIB composition to find a new

mate which is MHC-IIB dissimilar compared to an individual with a common MHC-IIB

composition. Depending on the year, the MHC-IIB composition of the individuals available

in the population might change and affect the ability to find MHC-IIB dissimilar mates.

These phenomena can create individual and year patterns, and the use of random effects

allows us to account for this potential source of variability. We tested the effect of individual

identity and year as random intercepts with the ranef function of the lme4 package ([70]) and

removed these variables when they had no effect. Similarly to the previous analysis, we used

two different models: after a reproductive failure and after a reproductive success. We used

the 0.5 quantile of a Quantile Regression Model (lqm function of the lqmm package, [73])

instead of LM when we encountered problems of non-normality and/or heteroscedasticity of

the residuals (tested with the simulateResiduals function of the DHARMa package, [74]).

Results

Analyses from the point of view of the males

First-time male breeders paired with a female which had a higher MHC-IIB supertype

sharing than expected by chance (P = 0.01). This result was not significant when considering

MHC-IIB functional distance (P = 0.08, Fig. 1).

Mate change probability varied differently with the MHC-IIB dissimilarity of the pair

depending on the reproductive output (Fig. 2). After a successful reproduction (n = 620

reproductive events), males were less likely to change mates if they were more MHC-IIB

dissimilar to the female (functional distance: log odd ± SE = -0.22 ± 0.09, P = 0.02;

difference in supertype sharing: log odd ± SE = -0.03 ± 0.01, P = 0.05). Mate change

probability was higher after a breeding failure (n = 484 reproductive events) than after a

breeding success (107 % increase in the probability of mate change compared to after a

breeding success). However, after a breeding failure, mate change probability did not vary

with the MHC-IIB dissimilarity of the pair (functional distance: log odd ± SE = 0.02 ± 0.06,

P = 0.73; difference in supertype sharing: log odd ± SE = 0.00 ± 0.01, P = 0.65). When

considering only pairs in which both mates were seen during the years n and n+1 (i.e., strict

divorce only), the MHC-IIB dissimilarity did not affect the probability of mate change,

neither after a reproductive failure (functional distance: log odd ± SE = 0.10 ± 0.10, P = 0.31;

difference in supertype sharing: log odd ± SE = 0.00 ± 0.01, P = 0.80; Supporting

information 6), nor after a reproductive success (functional distance: log odd $\pm$ SE = -0.20 $\pm$

0.15, P = 0.18; difference in supertype sharing: log odd $\pm$ SE = -0.02 $\pm$ 0.02, P = 0.28;

Supporting information 6).

When mate change occurred after a reproductive success, the MHC-IIB functional

distance between the male and the new female was higher compared to that between the male

and the previous female (mean $\pm$ SE = 0.51 $\pm$ 0.23, P = 0.03; difference in MHC-IIB

supertype sharing: P = 0.05; Fig. 3). In contrast, after a reproductive failure, there was no

difference in MHC-IIB functional distance between the new and the previous pair (mean $\pm$

SE = -0.23 $\pm$ 0.25, P = 0.35; Fig. 3; difference in MHC-IIB supertype sharing: P = 0.91).

However, males did not re-mate with a female who had a higher MHC-IIB functional

distance than expected under random mating, neither after a reproductive success (P = 0.31;

difference in MHC-IIB supertype sharing: P = 0.24) nor after a reproductive failure (P = 0.22,

Fig. 4; difference in MHC-IIB supertype sharing: P = 0.20).

Analyses from the point of view of the females

Females did not mate with a more MHC-IIB dissimilar male than expected under random

mating, neither for their first breeding attempt (functional distance: P = 0.33, Fig. 5a;

difference in supertype sharing: P = 0.20), nor when they were experienced breeders

(functional distance: P = 0.20; Fig. 5b; difference in supertype sharing: P = 0.08).

In females, mate change probability did not vary with the MHC-IIB dissimilarity of

the pair, neither after a reproductive failure (functional distance: P = 0.31; difference in

supertype sharing: P = 0.44), nor after a reproductive success (functional distance: P = 0.66;

difference in supertype sharing: P = 0.78). We found similar results when considering only

pairs for which both mates were seen in the years n and n+1 (Supporting information 6).

However, as expected, the probability to change mate was higher after a reproductive failure

than after a reproductive success (49 % increase in the probability of mate change compared

to after a breeding success). The difference in MHC-IIB dissimilarity between the new and

the previous male mate was not different from 0 (functional distance: mean $\pm$ SE = 0.03 $\pm$

0.17, P = 0.86; difference in supertype sharing: P = 0.95).

Discussion

In this study, we tested whether the mating strategies (i.e. mate choice and mate change) of

black-legged kittiwakes depend on the MHC-IIB dissimilarity between mates. We found sex-

specific patterns.

First-time male breeders were paired with more MHC-IIB dissimilar females than

expected by chance. Like several other species (e.g. [75]; [76]; [28]; [77]; [78]), including

birds ([79]; [80]), black-legged kittiwakes are able to recognize MHC-IIB dissimilarity using

body odors ([63]). In this species, the recognition of MHC-IIB dissimilarity might have

evolved as a mechanism to select MHC-IIB dissimilar mates to produce offspring with high

MHC-IIB dissimilarity ([49]), and therefore to increase female offspring growth, tick

clearance and survival ([50]). Similarly, in the domestic cat (*Felis catus*), mating with MHC-

IIB dissimilar mates increases offspring survival and thus probably parental fitness ([81]). In

contrast to first-time breeding males, we did not find evidence for MHC-IIB dissimilarity

based mate choice when considering first-time breeder females. Our analysis design does not,

however, allow us to determine whether these results are driven by males' choice or females'

choice. For instance, first-time male breeders may preferentially choose MHC-IIB dissimilar

females, or females may preferentially choose MHC-IIB dissimilar males when they have to

mate with a first-time male breeder. Similarly, in bluethroats (*Luscinia svecica*), only pairs

including females with a young male are more MHC-compatible than expected from random

pairing ([82]). In kittiwakes, the breeding experience is an important factor determining

reproductive success ([83]). Females paired with young inexperienced males might thus

compensate for the lack of experience of their mate by choosing MHC-IIB dissimilar males.

Males paired with a more MHC-IIB similar female had a higher probability of

changing mate in the next breeding season than males paired with a more MHC-IIB

dissimilar female. However, this result was significant only for pairs that reproduced

successfully. In pairs that failed to reproduce, mate change probability was higher than in

pairs that reproduced successfully, but it did not vary with the MHC-IIB dissimilarity

between mates. Hence, besides reinforcing the known role of breeding success on mate

change in black-legged kittiwakes ([47]; [84]; [48]), our results reveal the importance of

MHC-IIB dissimilarity in mate change in this species. However, we did not detect an effect

of MHC-IIB dissimilarity on mate change probability when considering females who were

observed changing mate, or when focusing on strict divorces. These results suggest that the

splitting pairs in which the males were seen alive in the studied population differ from those

in which the female or both mates were seen alive in the studied population after splitting.

Accordingly, in these former pairs, only 14 % of the females were seen alive in the studied

population. Low sample size coupled with potential sampling bias can affect the conclusions

of the analyses when studying the role of MHC-IIB composition in mating strategies (e.g. in

randomisation tests, [85]). Thus, the structure of our dataset limits our ability to conclude on

the absence of an effect of the MHC-IIB dissimilarity on the probability of divorce.

Nevertheless, in black-legged kittiwakes, females invest more than males in

reproduction ([55]; [56]), and they adjust their parental care to chick requirements more than

males ([86]; [87]). In pairs that reproduced successfully, females of MHC-IIB similar pairs

might have invested more energy in parental care than females of more MHC-IIB dissimilar

pairs to compensate for the deleterious effects of reduced MHC-IIB diversity of the offspring.

This higher investment rate might have reduced their survival at sea during the next winter

and explain the lower re-observation rate ([86]). It could eventually have forced their male

partner to mate with a new female. Non-exclusively, either mate might have chosen to

divorce. In kittiwakes, females are the dispersing sex ([88]) and they are less faithful to the

nesting site than males ([47]). Accordingly, in our dataset, females were less often re-

observed in the studied population after a mate change compared to males (n = 14 % of

females re-observed versus 21 % of males re-observed among splitting pairs). Therefore, the

females mated with a MHC-IIB similar male might have dispersed away from the tower, or

even from the colony, which forced the male to re-mate with a new partner. Under this latter

scenario, the reproductive success, as measured by the successful fledging of a chick, would

not be the only determinant of divorce in black-legged kittiwakes. For instance, not only the

presence of a fledgling, but also its quality (sensu [89]), which is shaped in part by its MHC-

IIB diversity and thus the MHC-IIB dissimilarity of the parents ([50]; [49]), might be a

determinant of the probability of female changing mates and nesting site. A previous study in

black-legged kittiwakes showed that MHC-IIB similar parents preferentially produce males

rather than females, probably to mitigate the cost of producing a female chick with low

MHC-IIB diversity ([49]).

Our study suggests that mate change might be another parental strategy to reduce the

deleterious effect of mating with a MHC-IIB similar mate. To which extent these potential

strategies translate into fitness benefits requires further studies. The lack of association

between MHC-IIB dissimilarity and mate change probability when considering females seen

alive in the studied population suggests that these females have changed mate for other

reasons than high MHC-IIB similarity. These reasons may include the death of the male, late

arrival of the male at its usual breeding site, or the intrusion of a third individual ousting the

male ([90]; [91]).

We found that although males who changed mate tended to pair with a new female

not more MHC-IIB dissimilar to them than a random female, the new female was more

MHC-IIB dissimilar to them than the previous one. Even though our results cannot ascertain

the existence of an active mate choice of males when remating, they suggest that even

random mate choice is sufficient to pair with a more MHC-IIB dissimilar female compared to

the previous one. Interestingly, we found that males involved in MHC-IIB similar pairs

tended to have very common MHC-IIB genotypes (Supporting information 2), thereby

probably limiting their possibility to mate with a female which is highly MHC-IIB dissimilar

to them.

The contrasting results between males and females likely illustrate sex-specific

mating strategies. In long-lived monomorphic species in which males and females equally

share parental care, such as the black-legged kittiwake ([61]), mate choice has been suggested

to be mutual ([92]). However, males and females may have different preference criteria ([93];

[94]). In the Leach's storm-petrel (*Oceanodroma leucorhoa*), males show MHC-IIB based

disassortative mate choice, whereas females' choice does not differ from random mating (

[95]). The intrinsic quality of the parent, such as its ability to provide good genes ([96]; [97]),

but also good parental care ([98]) or good-quality territory ([99]), may be more important

factor for one sex than the other. In kittiwakes, females invest more in reproduction than

males ([55]; [56]) and males may thus benefit from choosing a mate in good health or

condition more than females. Future studies are needed to investigate whether mate choice

based on MHC-IIB dissimilarity trades off with mate choice for other criteria and it varies

with sexes. For instance, in a cichlid fish, while genetic similarity is a more important criteria

for mate choice than body size for females, the opposite is true for males ([10]).

A previous study on the same population of black-legged kittiwakes showed that pairs

have a higher genetic dissimilarity (as measured by microsatellites) than expected by chance,

and that the resulting increased heterozygosity of their chicks leads to higher fitness ([100]). Although MHC-IIB disassortative mate choice may also participate in limiting inbreeding ([77]; [101]), here we detected only a weak correlation between genome-wide relatedness and MHC-IIB dissimilarity, which was driven by two pairs made of first-degree relatives. These results suggest that MHC-IIB based mating strategies detected in black-legged kittiwakes are probably driven by the MHC rather than resulting from inbreeding avoidance at the genome-wide scale.

In conclusion, we found evidence that MHC-IIB dissimilarity influences mate choice in the black-legged kittiwake and that low MHC-IIB dissimilarity between mates increases the probability for males to change mates in the future, even after a reproductive success. These results support the role of the benefits associated with high MHC-IIB dissimilarity in the evolution of mating strategies in this population ([50]). The selective pressures for higher MHC-IIB diversity, such as the level of parasite exposure depending on environmental conditions ([102]), as well as the balance between the benefits of a higher MHC-IIB diversity and those of other mating criteria, can however vary between species, population, ages or sexes, as suggested in the present study. In the latter case, when mate choice is mutual, a sexual conflict over mating strategies can arise ([103]) and may select for a rich repertoire of sex-specific behavioural adaptations ([104]). Our study paves the road for further studies of sex-specific mate choice in species with bi-parental care, a common trait in birds ([105]).

References

1. Talmon Y: Mate selection in collective settlements. American Sociological Review.

1964:491-508. DOI: 10.2307/2091199

2. Pomiankowski A: The costs of choice in sexual selection. Journal of theoretical Biology.

1987;128:195-218. DOI: 10.1016/s0022-5193(87)80169-8

3. Andersson M and Simmons LW: Sexual selection and mate choice. Trends in ecology \&

evolution. 2006;21:296-302. DOI: 10.1016/j.tree.2006.03.015

4. Rosenthal GG and Ryan MJ: Sexual selection and the ascent of women: Mate choice

research since Darwin. Science. 2022;375:eabi6308. DOI: 10.1126/science.abi6308

5. Hamilton WD and Zuk M: Heritable true fitness and bright birds: a role for parasites?.

Science. 1982;218:384-387. DOI: 10.1126/science.7123238

6. Andersson M and Iwasa Y: Sexual selection. Trends in ecology \& evolution. 1996;11:53-

58. DOI: 10.1016/0169-5347(96)81042-1

7. Dougherty LR: Meta-analysis reveals that animal sexual signalling behaviour is honest and

resource based. Nature Ecology \& Evolution. 2021;5:688-699. DOI: 10.1038/s41559-

021-01409-z

8. Tregenza T and Wedell N: Genetic compatibility, mate choice and patterns of parentage:

invited review. Molecular ecology. 2000;9:1013-1027. DOI: 10.1046/j.1365-

294x.2000.00964.x

9. Lihoreau M, Zimmer C and Rivault C: Mutual mate choice: when it pays both sexes to

avoid inbreeding. PloS one. 2008;3:e3365. DOI: 10.1371/journal.pone.0003365

10. Thünken T, Meuthen D, Bakker TC and Baldauf SA: A sex-specific trade-off between

mating preferences for genetic compatibility and body size in a cichlid fish with mutual

mate choice. Proceedings of the Royal Society B: Biological Sciences. 2012;279:2959-

2964. DOI: 10.1098/rspb.2012.0333

11. Ihle M, Kempenaers B and Forstmeier W: Fitness benefits of mate choice for

compatibility in a socially monogamous species. PLoS biology. 2015;13:e1002248. DOI:

10.1371/journal.pbio.1002248

12. McCully FR, Descamps S, Harris SM, Mckendrick F, Gillies N, Cornell SJ, Hatchwell BJ

and Patrick SC: Links between personality, reproductive success and re-pairing patterns

in a long-lived seabird. Ethology. 2023;129:686-700. DOI: 10.1111/eth.13405

13. Sullivan MS: Mate choice as an information gathering process under time constraint:

implications for behaviour and signal design. Animal Behaviour. 1994;47:141-151. DOI:

10.1006/anbe.1994.1016

14. Coulson J: The significance of the pair-bond in the Kittiwake. 1972;15:424-433.

15. McNamara JM and Forslund P: Divorce rates in birds: predictions from an optimization

model. The American Naturalist. 1996;147:609-640. DOI: 10.1086/285869

16. Dubois F and Cézilly F: Breeding success and mate retention in birds: a meta-analysis.

Behavioral Ecology and Sociobiology. 2002;52:357-364. DOI: 10.1007/s00265-002-

0521-z

17. Culina A, Radersma R and Sheldon BC: Trading up: the fitness consequences of divorce

in monogamous birds. Biological Reviews. 2015;90:1015-1034. DOI: 10.1111/brv.12143

18. Gousy-Leblanc M, Merkling T, Whelan S, Gaston AJ, Friesen VL and Elliott KH: Go

your own way? Reasons for divorce in a monogamous seabird. Animal Behaviour.

2023;204:13-24. DOI: 10.1016/j.anbehav.2023.07.007

19. Streif M and O ANNE E RASA: Divorce and its consequences in the common blackbird

Turdus merula. Ibis. 2001;143:554-560. DOI: 10.1111/j.1474-919x.2001.tb04882.x

20. Dreiss A and Roulin A: Divorce in the barn owl: securing a compatible or better mate

entails the cost of re-pairing with a less ornamented female mate. Journal of evolutionary

biology. 2014;27:1114-1124. DOI: 10.1111/jeb.12402

21. Jones IL and Montgomerie R: Mating and remating of least auklets (Aethia pusilla)

relative to ornamental traits. Behavioral Ecology. 1991;2:249-257. DOI:

10.1093/beheco/2.3.249

22. Fraser GS, Jones IL, Hunter FM and Cowen L: Mate switching patterns in crested auklets

(Aethia cristatella): the role of breeding success and ornamentation. Bird Behavior.

2004;16:7-12.

23. Hidalgo Aranzamendi N, Hall ML, Kingma SA, Sunnucks P and Peters A: Incest

avoidance, extrapair paternity, and territory quality drive divorce in a year-round

territorial bird. Behavioral Ecology. 2016:arw101. DOI: 10.1093/beheco/arw101

24. Penn DJ and Potts WK: The evolution of mating preferences and major histocompatibility

complex genes. The American Naturalist. 1999;153:145-164. DOI: 10.1086/303166

25. Sommer S: The importance of immune gene variability (MHC) in evolutionary ecology

and conservation. Frontiers in zoology. 2005;2:1-18. DOI: 10.1186/1742-9994-2-16

26. Winternitz J, Abbate J, Huchard E, Havliček J and Garamszegi L: Patterns of MHC-

dependent mate selection in humans and nonhuman primates: a meta-analysis. Molecular

ecology. 2017;26:668-688. DOI: 10.1111/mec.13920

27. Özer O and Lenz TL: Unique pathogen peptidomes facilitate pathogen-specific selection

and specialization of MHC alleles. Molecular Biology and Evolution. 2021;38:4376-

4387. DOI: 10.1093/molbev/msab176

28. Milinski M: The major histocompatibility complex, sexual selection, and mate choice.

Annu. Rev. Ecol. Evol. Syst.. 2006;37:159-186. DOI:

10.1146/annurev.ecolsys.37.091305.110242

29. Lenz TL: Computational prediction of MHC II-antigen binding supports divergent allele

advantage and explains trans-species polymorphism. Evolution. 2011;65:2380-2390.

DOI: 10.1111/j.1558-5646.2011.01288.x

30. Oliver MK, Telfer S and Piertney SB: Major histocompatibility complex (MHC)

heterozygote superiority to natural multi-parasite infections in the water vole (Arvicola

terrestris). Proceedings of the Royal Society B: Biological Sciences. 2009;276:1119-

1128. DOI: 10.1098/rspb.2008.1525

31. Dunn PO, Bollmer JL, Freeman-Gallant CR and Whittingham LA: MHC variation is

related to a sexually selected ornament, survival, and parasite resistance in common

yellowthroats. Evolution. 2013;67:679-687. DOI: 10.1111/j.1558-5646.2012.01799.x

32. Coker OM, OSAIYUWU OH and FATOKI AO: Major Histocompatibility Complex

(MHC) Diversity and its implications in human and wildlife health and Conservation.

Genetics \& Biodiversity Journal. 2023;7:1-11. DOI: 10.46325/gabj.v7i2.318

33. Fu M, Eimes JA and Waldman B: Divergent allele advantage in the MHC and amphibian

emerging infectious disease. Infection, Genetics and Evolution. 2023;111:105429. DOI:

10.1016/j.meegid.2023.105429

34. Ferreira D, San-Jose LM, Roulin A, Gaigher A and Fumagalli L: Limited associations

between MHC diversity and reproductive success in a bird species with biparental care.

Ecology and Evolution. 2024;14:e10950. DOI: 10.1002/ece3.10950

35. Landry C, Garant D, Duchesne P and Bernatchez L: Good genes as heterozygosity: the

major histocompatibility complex and mate choice in Atlantic salmon (Salmo salar).

Proceedings of the Royal Society of London. Series B: Biological Sciences.

2001;268:1279-1285. DOI: 10.1098/rspb.2001.1659

36. Radwan J, Babik W, Kaufman J, Lenz TL and Winternitz J: Advances in the evolutionary

understanding of MHC polymorphism. Trends in Genetics. 2020;36:298-311. DOI:

10.1016/j.tig.2020.01.008

37. Miller HC, Moore JA, Nelson NJ and Daugherty CH: Influence of major

histocompatibility complex genotype on mating success in a free-ranging reptile

population. Proceedings of the Royal Society B: Biological Sciences. 2009;276:1695-

1704. DOI: 10.1098/rspb.2008.1840

38. Agbali M, Reichard M, Bryjová A, Bryja J and Smith C: Mate choice for nonadditive

genetic benefits correlate with MHC dissimilarity in the rose bitterling (Rhodeus

ocellatus). Evolution. 2010;64:1683-1696. DOI: 10.1111/j.1558-5646.2010.00961.x

39. Juola FA and Dearborn DC: Sequence-based evidence for major histocompatibility

complex-disassortative mating in a colonial seabird. Proceedings of the Royal Society B:

Biological Sciences. 2012;279:153-162. DOI: 10.1098/rspb.2011.0562

40. Strandh M, Westerdahl H, Pontarp M, Canbäck B, Dubois M-P, Miquel C, Taberlet P and

Bonadonna F: Major histocompatibility complex class II compatibility, but not class I,

predicts mate choice in a bird with highly developed olfaction. Proceedings of the Royal

Society B: Biological Sciences. 2012;279:4457-4463. DOI: 10.1098/rspb.2012.1562

41. Huchard E, Baniel A, Schliehe-Diecks S and Kappeler PM: MHC-disassortative mate

choice and inbreeding avoidance in a solitary primate. Molecular Ecology.

2013;22:4071-4086. DOI: 10.1111/mec.12349

42. Løvlie H, Gillingham MA, Worley K, Pizzari T and Richardson DS: Cryptic female

choice favours sperm from major histocompatibility complex-dissimilar males.

Proceedings of the Royal Society B: Biological Sciences. 2013;280:20131296. DOI:

10.1098/rspb.2013.1296

43. Richardson DS, Komdeur J, Burke T and Von Schantz T: MHC based patterns of social

and extra pair mate choice in the Seychelles warbler. Proceedings of the Royal Society

B: Biological Sciences. 2005;272:759-767. DOI: 10.1098/rspb.2004.3028

44. Wright DJ, Brouwer L, Mannarelli M-E, Burke T, Komdeur J and Richardson DS: Social

pairing of Seychelles warblers under reduced constraints: MHC, neutral heterozygosity,

and age. Behavioral Ecology. 2016;27:295-303. DOI: 10.1093/beheco/arv150

45. Helfenstein F, Tirard C, Danchin E and Wagner RH: Low frequency of extra-pair

paternity and high frequency of adoption in black-legged kittiwakes. The Condor.

2004;106:149-155. DOI: 10.1650/7337

46. Coulson J: The influence of the pair-bond and age on the breeding biology of the

kittiwake gull Rissa tridactyla. Journal of Animal Ecology. 1966:269-279. DOI:

10.2307/2394

47. Naves L, Yves Monnat J and Cam E: Breeding performance, mate fidelity, and nest site

fidelity in a long-lived seabird: behaving against the current?. Oikos. 2006;115:263-276.

DOI: 10.1111/j.2006.0030-1299.14883.x

48. Mercier G, Yoccoz NG and Descamps S: Influence of reproductive output on divorce

rates in polar seabirds. Ecology and Evolution. 2021;11:12989-13000. DOI:

10.1002/ece3.7775

49. Pineaux M, Merkling T, Danchin E, Hatch SA, Leclaire S and Blanchard P: MHC-II

distance between parents predicts sex allocation decisions in a genetically monogamous

bird. Behavioral Ecology. 2022;33:245-251. DOI: 10.1093/beheco/arab130

50. Pineaux M, Merkling T, Danchin E, Hatch S, Duneau D, Blanchard P and Leclaire S: Sex

and hatching order modulate the association between MHC-II diversity and fitness in

early-life stages of a wild seabird. Molecular Ecology. 2020;29:3316-3329. DOI:

10.1111/mec.15551

51. Dittmann T and Becker PH: Sex, age, experience and condition as factors affecting

arrival date in prospecting common terns, Sterna hirundo. Animal Behaviour.

2003;65:981-986. DOI: 10.1006/anbe.2003.2128

52. Bosman DS, Vercruijsse HJ, Stienen EW, Vincx M and Lens L: Age of first breeding

interacts with pre-and post-recruitment experience in shaping breeding phenology in a

long-lived gull. PLoS One. 2013;8:e82093.

53. Campioni L, Dias MP, Granadeiro JP and Catry P: An ontogenetic perspective on

migratory strategy of a long-lived pelagic seabird: Timings and destinations change

progressively during maturation. Journal of Animal Ecology. 2020;89:29-43. DOI:

10.1111/1365-2656.13044

54. Burley NT and Foster VS: Variation in female choice of mates: condition influences

selectivity. Animal Behaviour. 2006;72:713-719. DOI: 10.1016/j.anbehav.2006.01.017

55. Fyhn M, Gabrielsen GW, Nordøy ES, Moe B, Langseth I and Bech C: Individual

variation in field metabolic rate of kittiwakes (Rissa tridactyla) during the chick-rearing

period. Physiological and Biochemical Zoology. 2001;74:343-355. DOI:

10.1086/320419

56. Moe B, Langseth I, Fyhn M, Gabrielsen GW and Bech C: Changes in body condition in

breeding kittiwakes Rissa tridactyla. Journal of Avian Biology. 2002;33:225-234.

57. Gill VA and Hatch SA: Components of productivity in black-legged kittiwakes Rissa

tridactyla: response to supplemental feeding. Journal of Avian Biology. 2002;33:113-

126. DOI: 10.1034/j.1600-048X.2002.330201.x

58. Merkling T, Leclaire S, Danchin E, Lhuillier E, Wagner RH, White J, Hatch SA and

Blanchard P: Food availability and offspring sex in a monogamous seabird: insights from

an experimental approach. Behavioral Ecology. 2012;23:751-758. DOI:

10.1093/beheco/ars023

59. Jodice P, Lanctot RB, Gill V, Roby DD and Hatch SA: Sexing adult black-legged

kittiwakes by DNA, behavior, and morphology. Waterbirds. 2000:405-415. DOI:

10.2307/1522177

60. White J, Wagner RH, Helfenstein F, Hatch SA, Mulard H, Naves LC and Danchin E:

Multiple deleterious effects of experimentally aged sperm in a monogamous bird.

Proceedings of the National Academy of Sciences. 2008;105:13947-13952. DOI:

10.1073/pnas.0803067105

61. Leclaire S, White J, Arnoux E, Faivre B, Vetter N, Hatch SA and Danchin É: Integument

coloration signals reproductive success, heterozygosity, and antioxidant levels in chick-

rearing black-legged kittiwakes. Naturwissenschaften. 2011;98:773-782. DOI:

10.1007/s00114-011-0827-7

62. Minias P, Pikus E, Whittingham LA and Dunn PO: A global analysis of selection at the

avian MHC. Evolution. 2018;72:1278-1293. DOI: 10.1111/evo.13490

63. Pineaux M, Blanchard P, Ribeiro L, Hatch SA and Leclaire S: A gull species recognizes

mhc-ii diversity and dissimilarity using odor cues. 2023:139-151. DOI: 10.1007/978-3-

031-35159-4_7

64. Sebastian A, Herdegen M, Migalska M and Radwan J: amplisas: a web server for

multilocus genotyping using next-generation amplicon sequencing data. Molecular

Ecology Resources. 2016;16:498-510.

65. Leclaire S, van Dongen WF, Voccia S, Merkling T, Ducamp C, Hatch SA, Blanchard P,

Danchin É and Wagner RH: Preen secretions encode information on MHC similarity in

certain sex-dyads in a monogamous seabird. Scientific Reports. 2014;4:1-6. DOI:

10.1038/srep06920

66. Sandberg M, Eriksson L, Jonsson J, Sjöström M and Wold S: New chemical descriptors

relevant for the design of biologically active peptides. A multivariate characterization of

87 amino acids. Journal of medicinal chemistry. 1998;41:2481-2491. DOI:

10.1021/jm9700575

67. Roved J, Hansson B, Stervander M, Hasselquist D and Westerdahl H: MHCtools--an R

package for MHC high-throughput sequencing data: Genotyping, haplotype and

supertype inference, and downstream genetic analyses in non-model organisms.

Molecular Ecology Resources. 2022;22:2775-2792. DOI: 10.1111/1755-0998.13645

68. Oksanen J, Kindt R, Legendre P, O?Hara B, Stevens MHH, Oksanen MJ and Suggests M:

The vegan package. Community ecology package. 2007;10:719.

69. Zhang B-y, Hu H-y, Song C-m, Huang K, Dunn DW, Yang X, Wang X-w, Zhao H-t,

Wang C-l, Zhang P and others: MHC-based mate choice in wild golden snub-nosed

monkeys. Frontiers in Genetics. 2020;11:609414. DOI: 10.3389/fgene.2020.609414

70. Bates D, Mächler M, Bolker B and Walker S: Fitting Linear Mixed-Effects Models Using

lme4. Journal of Statistical Software. 2015;67:1-48. DOI: 10.18637/jss.v067.i01

71. Breslow N: Whither pql?. 2004:1-22. DOI: 10.1007/978-1-4419-9076-1_1

72. Knowles JE and Frederick C: merTools: Tools for Analyzing Mixed Effect Regression

Models. 2024. Available from: https://github.com/jknowles/mertools

73. Geraci M: Linear Quantile Mixed Models: The lqmm Package for Laplace Quantile

Regression. Journal of Statistical Software. 2014;57:1-29. DOI: 10.18637/jss.v057.i13

74. Hartig F: DHARMa: Residual Diagnostics for Hierarchical (Multi-Level / Mixed)

Regression Models. 2024. Available from: http://florianhartig.github.io/DHARMa/

75. Wedekind C, Seebeck T, Bettens F and Paepke AJ: MHC-dependent mate preferences in

humans. Proceedings of the Royal Society of London. Series B: Biological Sciences.

1995;260:245-249. DOI: 10.1098/rspb.1995.0087

76. Grob B, Knapp L, Martin R and Anzenberger G: The major histocompatibility complex

and mate choice: inbreeding avoidance and selection of good genes. Experimental and

clinical immunogenetics. 1998;15:119-129. DOI: 10.1159/000019063

77. Penn DJ: The scent of genetic compatibility: sexual selection and the major

histocompatibility complex. Ethology. 2002;108:1-21. DOI: 10.1046/j.1439-

0310.2002.00768.x

78. Schubert N, Nichols HJ and Winternitz JC: How can the MHC mediate social odor via

the microbiota community? A deep dive into mechanisms. Behavioral Ecology.

2021;32:359-373. DOI: 10.1093/beheco/arab004

79. Leclaire S, Strandh M, Mardon J, Westerdahl H and Bonadonna F: Odour-based

discrimination of similarity at the major histocompatibility complex in birds.

Proceedings of the Royal Society B: Biological Sciences. 2017;284:20162466. DOI:

10.1098/rspb.2016.2466

80. Grieves L, Gloor G, Bernards M and MacDougall-Shackleton E: Songbirds show odour-

based discrimination of similarity and diversity at the major histocompatibility complex.

Animal behaviour. 2019;158:131-138. DOI: 10.1016/j.anbehav.2019.10.005

81. Erofeeva MN, Alekseeva GS, Kim MD, Sorokin PA and Naidenko SV: Inbreeding

coefficient and distance in MHC genes of parents as predictors of reproductive success in

domestic cat. Animals. 2022;12:165. DOI: 10.3390/ani12020165

82. Rekdal SL, Anmarkrud JA, Lifjeld JT and Johnsen A: Do female bluethroats without

extra-pair offspring have more MHC-compatible social mates?. Behavioral Ecology and

Sociobiology. 2023;77:36. DOI: 10.1007/s00265-023-03311-z

83. Cam E and Monnat J-Y: Apparent inferiority of first-time breeders in the kittiwake: the

role of heterogeneity among age classes. Journal of Animal Ecology. 2000;69:380-394.

DOI: 10.1046/j.1365-2656.2000.00400.x

84. Naves LC, Cam E and Monnat JY: Pair duration, breeding success and divorce in a long-

lived seabird: benefits of mate familiarity?. Animal Behaviour. 2007;73:433-444. DOI:

10.1016/j.anbehav.2006.10.004

85. Hoover B and Nevitt G: Modeling the importance of sample size in relation to error in

MHC-based mate-choice studies on natural populations. Integrative and Comparative

Biology. 2016;56:925-933. DOI: 10.1093/icb/icw105

86. Jacobsen K-O, Erikstad KE and Saether B-E: An experimental study of the costs of

reproduction in the kittiwake Rissa tridactyla. Ecology. 1995;76:1636-1642. DOI:

10.2307/1938164

87. Leclaire S, Helfenstein F, Degeorges A, Wagner RH and Danchin É: Family size and sex-

specific parental effort in black-legged kittiwakes. Behaviour. 2010:1841-1862. DOI:

10.1163/000579510x538872

88. Coulson J and Nève de Mévergnies G: Where do young kittiwakes Rissa tridactyla breed,

philopatry or dispersal. Ardea. 1992;80:187-197.

89. Wilson AJ and Nussey DH: What is individual quality? An evolutionary perspective.

Trends in ecology \& evolution. 2010;25:207-214. DOI: 10.1016/j.tree.2009.10.002

90. Jeschke JM, Wanless S, Harris MP and Kokko H: How partnerships end in guillemots

Uria aalge: chance events, adaptive change, or forced divorce?. Behavioral Ecology.

2007;18:460-466. DOI: 10.1093/beheco/arl109

91. Gilsenan C, Valcu M and Kempenaers B: Difference in arrival date at the breeding site

between former pair members predicts divorce in blue tits. Animal Behaviour.

2017;133:57-72. DOI: 10.1016/j.anbehav.2017.09.004

92. Johnstone RA, Reynolds JD and Deutsch JC: Mutual mate choice and sex differences in

choosiness. Evolution. 1996;50:1382-1391. DOI: 10.2307/2410876

93. Bergstrom CT and Real LA: Toward a theory of mutual mate choice: Lessons from two-

sided matching. 2000.

94. Aquiloni L and Gherardi F: Mutual mate choice in crayfish: large body size is selected by

both sexes, virginity by males only. Journal of Zoology. 2008;274:171-179. DOI:

10.1111/j.1469-7998.2007.00370.x

95. Hoover B, Alcaide M, Jennings S, Sin SYW, Edwards SV and Nevitt GA: Ecology can

inform genetics: Disassortative mating contributes to MHC polymorphism in Leach s

storm-petrels (Oceanodroma leucorhoa). Molecular Ecology. 2018;27:3371-3385. DOI:

10.1111/mec.14801

96. Neff BD and Pitcher TE: Genetic quality and sexual selection: an integrated framework

for good genes and compatible genes. Molecular ecology. 2005;14:19-38. DOI:

10.1111/j.1365-294x.2004.02395.x

97. Puurtinen M, Ketola T and Kotiaho JS: The good-genes and compatible-genes benefits of

mate choice. The American Naturalist. 2009;174:741-752. DOI: 10.1086/606024

98. Moody AT, Wilhelm SI, Cameron-MacMillan ML, Walsh CJ and Storey AE: Divorce in

common murres (Uria aalge): relationship to parental quality. Behavioral Ecology and

Sociobiology. 2005;57:224-230. DOI: 10.1007/s00265-004-0856-8

99. Hasegawa M, Arai E, Watanabe M and Nakamura M: Female mate choice based on

territory quality in barn swallows. Journal of ethology. 2012;30:143-150. DOI:

10.1007/s10164-011-0307-8

100. Mulard H, Danchin E, Talbot SL, Ramey AM, Hatch SA, White JF, Helfenstein F and

Wagner RH: Evidence that pairing with genetically similar mates is maladaptive in a

monogamous bird. BMC Evolutionary Biology. 2009;9:1-12. DOI: 10.1186/1471-2148-

9-147

101. Huang W, Pilkington JG and Pemberton JM: Patterns of MHC-dependent sexual

selection in a free-living population of sheep. Molecular Ecology. 2021;30:6733-6742.

DOI: 10.1101/2020.11.18.387332

102. Liu P, Li G, Zhao N, Liu Q, Liu X, Song X, Shi X, Lun X, Zhang L, Wang J and others:

Climate heterogeneity, season variation, and sexual dimorphism modulate the association

between MHC II diversity and parasite variation in striped hamster. Integrative Zoology.

2023. DOI: 10.1111/1749-4877.12791

103. Parker GA: Sexual conflict over mating and fertilization: an overview. Philosophical

Transactions of the Royal Society B: Biological Sciences. 2006;361:235-259. DOI:

10.1098/rstb.2005.1785

104. Sæther SA, Fiske P and Kålås JA: Male mate choice, sexual conflict and strategic

allocation of copulations in a lekking bird. Proceedings of the Royal Society of London.

Series B: Biological Sciences. 2001;268:2097-2102. DOI: 10.1098/rspb.2001.1745

105. Whittingham LA and Dunn PO: Male parental care and paternity in birds. 2001:257-

298. DOI: 10.1007/978-1-4615-1211-0_5

Figures

Figure 1: Frequency distribution of difference in MHC-IIB supertype sharing means between

first time male breeders and their female mate calculated from 10000 bootstrap

randomizations of potential male-female pairs out of the sample studied. The red vertical line

is the observed MHC-IIB functional distance. The blue vertical line is the upper 95 %

quantile of the simulated distribution.

Figure 2: Probability of mate change in males according to the MHC-IIB functional distance

with their female mate, when the pair reproduced successfully (in green) or not (in yellow).

The yellow and green dots represent the raw observations, and the yellow and green lines

represent the predictions of the model. The yellow and green shaded areas represent the 95 %

confidence intervals on the fixed effects.

Figure 3: Change in MHC-IIB functional distance between mates in year n+1 compared to

year n after the male changed mate following a reproductive success or a reproductive failure.

The horizontal black lines represent the medians, the red dots represent the means, the boxes

represent the interquartile ranges (IQR) of the distribution, the whiskers extend to the

minimum and maximum values within 1.5 times the IQR and the open dots represent the

values beyond these limits.

Figure 4: Frequency distribution of MHC-IIB functional distance means between male

breeders and their female mate calculated from 10000 bootstrap randomizations of potential

male-female pairs out of the sample studied when the previous pair had: a) a reproductive

success, b) a reproductive failure. The red vertical line is the observed MHC-IIB functional

distance. The blue vertical line is the upper 95 % quantile of the simulated distribution.

Figure 5: Frequency distribution of MHC-IIB functional distance means between: a) first

time female breeders, b) experienced female breeders and their male mate calculated from

10000 bootstrap randomizations of potential male-female pairs out of the sample studied. The

red vertical line is the observed MHC-IIB functional distance. The blue vertical line is the

upper 95 % quantile of the simulated distribution.