Does MHC-IIB dissimilarity between partners drive mate choice and divorce in a genetically monogamous seabird?

Lucie Thel[1,2], Maxime Pineaux[1], Léa Ribeiro[1], Etienne Danchin[1,3], Shannon Whelan[4], Scott A. Hatch[4], Pierrick Blanchard[1], Sarah Leclaire[1]

Authors details

[1] Université de Toulouse, Toulouse INP, CNRS, IRD, CRBE, Toulouse, France.

[2] Department of Conservation Management, Faculty of Science, Nelson Mandela University, George, South Africa.

[3] Centre de Recherches sur la Cognition Animale, Centre de Biologie Intégrative, UMR 5169 (CNRS, Université de Toulouse Midi-Pyrénées), Toulouse, France.

[4] Institute for Seabird Research and Conservation, Anchorage, Alaska, USA.

Corresponding authors: Lucie Thel, lucie.mc.thel@gmail.com; Sarah Leclaire, sarah.leclaire@univ-tlse3.fr

Acknowledgments

We thank the Middleton fieldworkers who collected data, as well as the undergraduate students who helped with molecular analyses.

Authors' contributions

Conceptualisation: LT, MP, PB, SL; Methodology: LT, MP, PB, SL; Data Curation: LT, PB, SL; Formal Analyses: LT, SL; Investigation: LT; Writing: Original Draft: LT, SL; Writing:

Review & Editing: LT, MP, ED, PB, SL; Visualisation: LT, SL; Data acquisition: MP, LR; Project administration: ED, SW, SH, SL; Funding acquisition: ED, SH, SL; Supervision: PB, SL; Resources: SL.

Ethics

The experiment was conducted under the approval of the U.S. Geological Survey Alaska Science Center, the University of Alaska Institutional Animal Care and Use Committees, and the Institut Paul-Emile Victor Ethical Committee, in accordance with United States laws and under permits from the U.S. Fish and Wildlife Service and the State of Alaska.

Conflict of interest

The authors declare that they have no conflict of interest.

Funding

This work was supported by the French Polar Institute Paul-Emile Victor (IPEV "Programme 1162" to ED and SL) and the Fondation Fyssen (to SL). This study is part of the "Laboratoire d'Excellence" (LABEX) entitled TULIP (ANR-10-LABX-41). LR was supported by an EUR TULIP Ph.D. grant obtained through SEVAB postgraduate school.

Data availability statement

Data available from the Zenodo Repository: https://doi.org/10.5281/zenodo.14509255.

Does MHC-IIB dissimilarity between partners drive mate choice

and divorce in a genetically monogamous seabird?

Abstract

Among species reproducing sexually, mating strategies represent a major component of

individual fitness. The Major Histocompatibility Complex (MHC) is an extremely diverse set

of genes responsible for immunological recognition and defence against pathogens. Although

MHC dissimilarity between mates has been proposed to drive mate choice through increased

offspring pathogen resistance, evidence is mixed. In addition, support for other MHC based

mating strategies, such as divorce, is rare. We investigated whether MHC-IIB dissimilarity

(MHC-IIB functional distance and difference in MHC-IIB supertype sharing) is associated

with mate choice and divorce probability in the genetically monogamous black-legged

kittiwake (*Rissa tridactyla*). We found that first-time male breeders, as well as divorced

males, were paired with a female with a greater difference in MHC-IIB supertype sharing to

them than expected under random mating. We did not find evidence for mate choice based

MHC-IIB dissimilarity when considering females. In addition, divorce probability did not

significantly vary with the MHC-IIB dissimilarity of the pair, and after a divorce, the newly

formed pair was not more MHC-dissimilar than the original pair. Our results pave the way to

a better understanding of the complex role of MHC dissimilarity in mating decisions of

species displaying mutual choice and biparental care.

Keywords: Black-legged kittiwake, *Rissa tridactyla*, Divorce, Reproduction, Major

Histocompatibility Complex, Disassortative mate choice

Introduction

Mating strategies are the set of behavioural tactics used by individuals to attract, select, change or retain mates to promote individual fitness. Understanding their driving factors has long been a major topic in evolutionary biology [1;2]. For instance, mate choice has repeatedly been shown to rely on traits that honestly reflect the quantity or quality of resources, the care or the genes that the mate can provide [3;4;5]. In addition, in several species, individuals preferentially choose mates which are more genetically or behaviourally compatible to avoid inbreeding or to enhance behavioural coordination or task complementation between mates [6;7;8;9;10]. However, individuals may face energetic, time or spatial constraints in choosing mates [11]. Under such circumstances, individuals are forced to choose from a limited pool of potential mates and thus may accept suboptimal partners [12]. A possible response to this limitation in iteroparous species is divorce, defined as one breeding individual changing partner for a new mate while the original mate is still alive [13]. Theoretical studies have suggested that individuals should divorce when pairing with a higher-quality or more compatible mate improves their reproductive success [14]. Accordingly, divorce is common after a reproductive failure [15;16;17], and usually leads to increased reproductive success [18;19]. Like mate choice, divorce may be associated with the quality of the ornaments of the mate, or the behavioural or genetic compatibility of the pair (e.g. [20;21]).

Concerning the genetic component of the pool of heritable factors that can affect mating strategies, the major histocompatibility complex (MHC, called HLA in humans) has been the focus of much research [22;23;24]. The MHC, one of the most polymorphic sets of genes in vertebrates, encodes cell-surface proteins that are essential for immunological recognition and defence against pathogens. Because each MHC molecule can bind a limited set of antigens [25], higher MHC diversity increases the number of antigens recognized,

thereby providing resistance to a wider range of parasites [26;27]. Accordingly, several

studies have found that higher MHC diversity increases health, reproductive success and

survival [28;29]. Because the MHC allelic composition of the offspring results from the

combination of those of the parents, multiple studies have found evidence for mate choice

maximising MHC dissimilarity between mates (e.g. [30;31;32;33]). However, high MHC

diversity can also increase the risk of autoimmune diseases and reduce the T cell receptors

diversity [34;35]. In a few species, intermediate MHC diversity has thus been shown to be

optimum for fitness, and to be an important factor in mate choice (e.g. [36;37;38;39]).

Although there is evidence that the MHC composition or diversity of the parents also

influences other reproductive strategies (e.g. cryptic female choice, [40]; extra-pair

copulation, [41]), to our knowledge, a single study on Seychelle warblers (*Acrocephalus*

sechellensis) explored the association between parental MHC similarity and divorce rate and

found no evidence [42].

Here we tested whether mate choice and divorce depend on dissimilarity at MHC

class-II B (hereafter MHC-IIB dissimilarity) in the black-legged kittiwake (*Rissa tridactyla*),

a long-lived genetically monogamous seabird with no extra-pair copulation [43]. In black-

legged kittiwakes, divorce is fairly common (i.e. 15 % to 50 % of the pairs divorce each

year), and more frequent among first-time breeders and after reproductive failure [44;45;46]

[10]. Coulson [44] suggested therefore that some kinds of "incompatibility" between mates

result in unsuccessful breeding, thereby providing costs that select for divorce. In black-

legged kittiwakes, individuals can recognize MHC-IIB dissimilarity using body odours [47],

and MHC-IIB similarity between mates results in reduced offspring MHC-IIB diversity [48],

which incurs costs for female chicks such as slower growth, poor tick clearance efficiency

and lower survival [49]. We therefore hypothesised that, in this species, mating decisions

based on MHC-IIB dissimilarity between mates may have evolved to maximise the MHC-IIB

diversity of the offspring. However, the association between MHC diversity and fitness has

never been explored in adults in this species, so one cannot exclude that intermediate MHC

diversity is optimum when considering lifetime fitness. For all our analyses, we thus also

explored the intermediate optimum hypothesis that predicts that individuals should mate with

partners with intermediate MHC dissimilarity [36].

First, we tested whether individuals chose a mate according to MHC dissimilarity

(Test 1). In several seabirds, including the black-legged kittiwake, young or inexperienced

individuals arrive later to the breeding ground than older or more experienced breeders

[50;51;52;53], which may increase the temporal constraints and limit the pool of potential

mates. Inexperienced breeders may also be less efficient in sampling mates and less able to

make mating discrimination or to anticipate alternative choices [54]. For instance, in grey

mouse lemurs (*Microcebus murinus*), young females in their first mating season do not

express mate choice, while experienced females prefer MHC-IIB dissimilar males [55]. The

mate choice of first-time breeders may thus be more constrained than that of experienced

breeders. We therefore tested the existence of MHC-IIB dependent mate choice in first-time

breeders (Test 1.1) and experienced breeders experiencing a divorce (Test 1.2). Second, if

divorcing allows individuals paired with a suboptimal mate to optimise the MHC-IIB

diversity of their future offspring, we expected the rate of divorce to depend on the MHC-IIB

dissimilarity between mates (Test 2). Third, we expected that, after a divorce, the new pair

should be more MHC-IIB compatible (maximised or intermediate dissimilarity) than the

original one (Test 3). In black-legged kittiwakes, although both mates participate fairly

equally in incubation and chick rearing, sexes slightly differ in energetic investment in

reproduction [56;57]. We thus tested these predictions separately in males and females to

explore potential sex differences in mating decisions.

Materials and methods

Study area and population

This study was conducted between 1996 and 2023 in a colony of black-legged kittiwakes nesting in an abandoned U.S. Air Force radar tower on Middleton Island (59°26'N, 146°20'W), Gulf of Alaska, located approximately 100 km south of the mainland. Individuals were tagged with a unique combination of coloured and metal bands for long-term monitoring [58]. About 400 nest sites were monitored on the tower each year. All the nests were checked daily to record the identity of the present individual(s). For each individual of the study, sex was determined, in order of priority, by: 1) molecular assessment (n = 840 individuals; [59]); 2) behavioural observations (n = 438 individuals; [60]); 3) comparison of the relative size of both members of a pair, the largest bird being considered male (n = 6 individuals); 4) when only one individual of a pair was sexed, we attributed the opposite sex to its mate (n = 214 individuals).

Like many populations monitored in the long term, our population has been subjected to several treatments during the study period (e.g. supplemental feeding, brood manipulation, feather clipping, use of anti-copulatory devices). These treatments can affect reproductive success [58;61] and therefore pair stability. To limit any bias (e.g. artificial increase of reproductive success, and thus decreased divorce probability due to supplemental feeding), we excluded pairs that were subjected to experiments from the analysis on the probability of divorce (Test 2).

We defined a pair as two individuals of opposite sex involved in a reproductive event, i.e. attending the same nest and laying at least one egg during a given reproductive season. Individuals usually remain and breed in the same colony during their whole life [44]. We thus defined the first mate choice as the pair formed during the first observed breeding attempt. Each individual was involved in on average 4.75 ± 3.74 (SE) reproductive events (range = [1;

21]; Supporting information 1) over the course of the study. To study divorces, we focused

on individuals that were observed with an identified mate during at least one year and that re-

mated with a new mate the next year, the original mate being re-observed at least once during

the following years [10].

We performed all analyses using the R software [62]. All datasets and code used in

the analyses are available from the Zenodo Repository

(https://doi.org/10.5281/zenodo.14509255).

Molecular analyses

We focused on the MHC class II which exhibits stronger selection than the MHC class I in

non-passerine birds such as the black-legged kittiwake [63]. Dissimilarity at MHC-IIB was

assessed following the methods described in Pineaux et al. [49;47]. Briefly, DNA was

extracted from blood with the blood and tissue DNA extraction kit (Qiagen), and a 258-pb

fragment of exon 2 of MHC-IIB was amplified using specific primers (forward: 5′-

GCACGAGCAGGGTATTTCCA, and reverse: 5′-GTTCTGCCACACACTCACC).

Amplicons were then sequenced with an Illumina MiSeq platform, using the 2 x 300-bp

protocol (Fasteris SA, Switzerland). Amplicon sequences were analysed with ampliSaS, a

three-step pipeline that consists of read demultiplexing, unique sequence clustering and

erroneous sequence filtering (substitution errors: 1 %; indel errors: 0.001 %; minimum

frequency with respect to dominant: 33 %; minimum amplicon sequence frequency: 2.5 %;

[64]). We calculated the MHC-IIB functional distance between individuals based on amino

acid variations in the peptide-binding region (PBR) inferred from Leclaire et al. [65].

We described the chemical binding properties of each amino acid in the PBR with

Sandberg's five physicochemical descriptors (z-descriptors, [66]) and computed a Sandberg's

distance matrix between sequences using the *DistCalc* function of the *MHCTools* package

[67]. Then, we calculated an UPGMA dendrogram of sequences (*hclust* function), which

represents clusters of functionally similar MHC-IIB sequences (hereafter, "supertypes"). The

MHC-IIB functional distance between a male and a female was estimated using the tree

distance (*treedist* function in the *vegan* package [68]), which is found by combining

sequences in a common dendrogram and calculating how much of the sum of the branch

length is shared and how much is unique. In addition, we calculated the difference in MHC-

IIB supertype sharing between mates similarly to Strandh et al. [69]:

$$100 \times [1 - Nb_{supertypes\ pair} / (Nb_{supertypes\ female} + Nb_{supertypes\ male})],$$

where $Nb_{supertypes\ female}$ corresponds to the number of supertypes possessed by the female,

$Nb_{supertypes\ male}$ corresponds to the number of supertypes possessed by the male, and $Nb_{supertypes}$

$_{pair}$ corresponds to the total number of supertypes possessed by the female and the male

together (where shared supertypes count as 1). This index thus varies between 0 for

individuals sharing no supertype and 100 for individuals sharing all their supertypes. The

MHC-IIB functional distance was highly correlated with the difference in MHC-IIB

supertype sharing (Spearman correlation test: $\rho = 0.79$, $P < 0.0001$, $n = 959$ pairs; Supporting

information 2).

Mate choice for MHC-IIB dissimilarity may not only be governed by evolutionary

processes that specifically target these genes, but may also be affected by mechanisms

operating at the level of the entire genome (e.g. inbreeding avoidance). To test whether the

associations between MHC-IIB dissimilarity and mate choice may be driven by genome-wide

relatedness, we used RAD sequencing ($n = 254$ individuals) to estimate genome-wide

relatedness between individuals. We found that genome-wide relatedness was correlated

neither with functional MHC-IIB similarity (Mantel test: $r = 0.03$, $P = 0.09$), nor with MHC-

IIB supertype sharing (Mantel test: $r = 0.03$, $P = 0.17$; Figure S3.1).

 Statistical analyses

To test the association between mate choice and MHC-IIB dissimilarity (Test 1), we

used bootstrapping to generate random pairs, where we let each individual choose randomly

10000 times between all available opposite-sex mates of the same year, without replacement

(i.e. one mate can be selected only once during a given year; [70;71]). To test for the

maximisation hypothesis, we calculated p-values as the proportion of iterations in which the

random MHC-IIB dissimilarity mean was greater than the observed MHC-IIB dissimilarity

mean, plus the proportion of iterations in which the random MHC-IIB dissimilarity mean was

smaller than the symmetrical (relative to the mean of random MHC-IIB dissimilarity means)

of the observed MHC-IIB dissimilarity mean (two-tailed test, following the method described

in [55]). To test for the intermediate optimum hypothesis, we used a similar procedure, but

instead of comparing the observed and random means of MHC-IIB dissimilarity, we

compared the observed and random variance in MHC-IIB dissimilarity (following the method

described in [71]). In the case of a mate choice for intermediate MHC-IIB dissimilarity, we

expect the observed and random means of MHC-IIB dissimilarity between partners to be

similar. However, because individuals are expected to prefer a mate that allows them to reach

the intermediate MHC-IIB dissimilarity optimum, we expect the observed variance in MHC-

IIB dissimilarity to be smaller than the random variance. We tested MHC-IIB based mate

choice for first-time breeders (n = 534 females and n = 554 males; Test 1.1) and for breeders

experiencing a divorce (n = 113 divorces in females and n = 83 divorces in males; Test 1.2)

separately.

To test whether divorce probability varied with the MHC-IIB dissimilarity of the pair

during the year n (Test 2), we defined a binary response variable as 1 = divorce between the

year n and n+1, and 0 = no divorce between the year n and n+1. We tested females (n = 1244)

and males (n = 1232) separately. We used Generalised Linear Mixed Models (GLMM) with a

binomial distribution of error and a logit link (*glmer* function of the *lme4* package; [72]). To

test for the intermediate optimum hypothesis, we included the quadratic term of the MHC-IIB

dissimilarity [73]. If this variable did not have a significant effect (i.e. $P > 0.05$), we removed

the quadratic term and then tested for the maximisation hypothesis with the linear term of the

MHC-IIB dissimilarity. The reproductive outcome is also known to influence the probability

of divorce in black-legged kittiwakes [46];10]. We thus defined the reproductive outcome as

a binary variable recording the presence (i.e. "success") or absence (i.e. "failure") of at least

one chick fledged at the end of the reproductive season and included this variable as an

additive effect. In the models, we set the *nAGQ* argument (number of points per axis for

evaluating the adaptive Gauss-Hermite approximation to the log-likelihood; [74]) of the

glmer function to 0 to help for convergence, when necessary. We tested the effect of

individual identity and year as random intercepts with the *ranef* function of the *lme4* package

[72], and removed these variables when they had no effect. We removed pairs that were

subjected to experiments from this analysis. As we used GLMM with binomial data, we

provided the log odd (and its standard error), which is equivalent to the coefficient of the

slope in a linear model (i.e. estimate in the model summary). It corresponds to the probability

of an event in the log scale, which is the transformed scale to satisfy the assumptions of a

linear model for GLMM with binomial data. For convenience, we also provided the odd ratio,

which corresponds to the exponential of the log odd. An odd ratio of 1 means no effect,

whereas an odd ratio > 1 (< 1, resp.) corresponds to an increase (decrease, resp.) of the

response variable when the explanatory variable (here, reproductive outcome or MHC-IIB

dissimilarity) increases by 1 unit. We calculated the confidence intervals from the model on

the fixed effects using the *predictInterval* function of the *merTools* package (n iterations =

10000; [75]).

To test whether, after divorce, the new mate was more MHC-IIB dissimilar than the

original mate (i.e. maximisation hypothesis), we used Linear Mixed Models (LMM; Test 3).

The response variable was the MHC-IIB dissimilarity gain, i.e. the difference between the

MHC-IIB dissimilarity to the original mate and the MHC-IIB dissimilarity to the new mate.

Using the gain in a linear modelling framework allowed us to include individual and year as

random effects. The MHC-IIB composition of the individuals available in the population

might vary with year, thereby potentially affecting the ability to find MHC-IIB dissimilar

mates. The use of random effects allowed us to account for this potential source of

variability. We tested the effect of individual identity and year as random intercepts with the

ranef function of the *lme4* package [72], and removed these variables when they had no

effect. We tested the intermediate optimum hypothesis by comparing the variance of the

MHC-IIB dissimilarity between the pool of original pairs and the pool of new pairs with a

Fisher exact test (or its non-parametric equivalent, the Levene test, when necessary). In the

case of mate choice following the intermediate hypothesis, we expect the variance of the

MHC-IIB dissimilarity in the new pair to be lower than the variance of the MHC-IIB

dissimilarity in the original pair.

Power analyses

We conducted test-specific power analyses to assess the robustness of our results. When

studying the MHC-IIB dissimilarity with the new mate compared to a random mate (Test 1.1

and 1.2), we evaluated the likelihood of error as detailed in Hoover et al. [70;76]. For

example, for males involved in divorces, we resampled between 10 and 83 males (the actual

number of individuals in this dataset) and conducted the same analysis as described

previously ($n_{boots} = 500$ instead of 10000 for reasons of computational limitations). We then

assigned a "1" to a significant result and a "0" to a non-significant result for every sample

size tested, and used a GLM with a binomial distribution of error and a logit link (*glmer*

function of the *lme4* package; [72]) to describe the relationship between sample size and

likelihood of error.

When assessing the probability of divorce according to MHC-IIB dissimilarity (Test

2), our dataset was large enough (> 1000 individuals), but largely unbalanced between

faithful and divorcing pairs (approx. 20 times less divorced than faithful pairs in both females

and males). We thus randomly resampled the faithful pairs to obtain the same sample size as

for divorced pairs and adjusted the same models (n_{boots} = 1000) as described previously on the

balanced dataset using GLMs. We then estimated the probability to obtain a significant effect

for each explanatory variable (i.e. quadratic and linear terms of MHC-IIB dissimilarity, and

reproductive outcome).

When assessing the gain in MHC-IIB dissimilarity between the original and the new

pair after a divorce (Test 3), we calculated the effect sizes for our analysis using Cohen's d

statistic [77].

Results

Mate choice in first-time breeders (Test 1.1)

Most of the first-time breeders mated with first-time breeders (62 % when considering males

and 65 % when considering females). For their first breeding attempt, males tended to mate

with a female which had a significantly greater difference in supertype sharing than expected

under random mating (P = 0.05; Table S4.1; Figure 1c). The probability to find a significant

result with the available sample size (n = 554 mate choice events) was 0.40 (Table S4.1;

Figure 1d). This effect was however not significant when considering the functional distance

between mates (P = 0.30; Probability to find a significant result = 0.03; Figure S4.1c and

S4.1d). Females did not mate with a male which had a significantly greater difference in

supertype sharing or greater functional distance to them than expected under random mating

(functional distance: P = 0.89; difference in supertype sharing: P = 0.72; Table S4.1; Figure

1a and Figure S4.1a). Neither females (functional distance: P = 0.98; difference in supertype

sharing: P = 0.88) nor males (functional distance: P = 0.75; difference in supertype sharing: P

= 0.49) mated with a significantly more intermediately dissimilar mates than expected under

random mating (Table S4.1; Figure S4.2 and S4.3).

Probability of divorce (Test 2)

In both sexes, the probability of divorce was significantly lower after a reproductive success

than after a reproductive failure (all $P \leq 0.01$; Probability to find a significant result with a

balanced dataset > 0.80; Table S5.1). However, the probability of divorce did not

significantly vary with the MHC-IIB dissimilarity of the pair (Figure S5.2), when considering

either females (linear term: functional distance: P = 0.58 and difference in supertype sharing:

P = 0.74; quadratic term: functional distance: P = 0.18 and difference in supertype sharing: P

= 0.46; Table S5.1) or males (linear term: functional distance: P = 0.68 and difference in

supertype sharing: P = 0.54; quadratic term: functional distance: P = 0.64 and difference in

supertype sharing: P = 0.81; Table S5.1). The probability to find a significant effect of MHC-

IIB dissimilarity on the probability of divorce was consistently < 0.20 with a balanced dataset

(Table S5.1).

Mate choice in divorced individuals (Tests 3 and 1.2)

Divorced individuals re-mated mainly with experienced breeders (63 % when considering

males and 71 % when considering females). The difference in MHC-IIB dissimilarity

between the new pair and the original pair was not different from 0, either in females or

males (Test 3; all $P > 0.10$; Table S6.1). The effect sizes revealed by Cohen's d statistics were

very low (all Cohen's d < 0.20; Table S6.1).

However, after a divorce, males were paired with a female which had a significantly

greater difference in supertype sharing to them than expected under random mating (Test 1.2:

$P = 0.04$; Probability to find a significant result $= 0.78$; Table S7.1; Figure 2c & 2d). Females

did not re-mate with a male which had a significantly greater difference in supertype sharing

than expected under random mating (Test 1.2: $P = 0.06$; Probability to find a significant result

$= 0.23$; Table S7.1; Figure 2a). These effects were not significant when considering

functional distance (males: $P = 0.09$ and females: $P = 0.62$; Table S7.1; Figure S7.1).

When considering either males or females, after a divorce, the new pair was not more

intermediately dissimilar than the original pair (Test 3: all $P > 0.10$; Table S6.1) or than a

random pair (Test 1.2: all $P > 0.10$; Table S7.1).

Discussion

In this study, we tested whether mating decisions (i.e. mate choice and divorce) of black-

legged kittiwakes vary with the MHC-IIB dissimilarity between mates. We found that first-

time male breeders (Test 1.1) as well as divorced males (and to a lower extent, divorced

females; Test 1.2) were paired with a mate more MHC-IIB dissimilar to them than expected

under random mating. However, we did not detect an association between the MHC-IIB

dissimilarity of the pair and the probability of divorce.

First-time male breeders, as well as divorced males, were more MHC-IIB dissimilar

to their female partner than expected under random mating. These results may suggest active

mate choice for MHC-IIB dissimilarity. Like several other species (e.g. [78;79;26;80]),

including birds [81;82], black-legged kittiwakes are able to recognize MHC-IIB dissimilarity

using body odours [47]. In this species, the recognition of MHC-IIB dissimilarity might have

evolved as a mechanism to select MHC-IIB dissimilar mates to produce offspring with high

MHC-IIB diversity [48], and thus with higher early-life performance [49]. Similarly, in the

domestic cat (*Felis catus*), mating with MHC-IIB dissimilar mates increases offspring

survival and thus probably parental fitness [83]. We detected MHC-based mate choice when

MHC-IIB dissimilarity was measured based on difference in supertype sharing, but not when

based on functional distance. The causes of these discrepancies are unknown but are

consistent with the literature. For instance, in blue petrels (*Halobaena caerulea*), yellow-

rumped flycatchers (*Ficedula zanthopygia*) and little auks (*Alle alle*), evidence for MHC-

based mating strategies are stronger when considering supertype or allele sharing than when

considering functional distances ([69;84;85]; but see [55] for the opposite pattern).

In contrast to results when considering males, we did not detect MHC-IIB

disassortative mate choice when considering first-time female breeders. In addition, we

detected only a tendency for MHC-IIB disassortative mate choice when considering divorced

females ($P = 0.06$, with 23 % likelihood of obtaining a significant result). Our analysis

design, however, does not clearly allow us to determine whether the MHC-IIB disassortative

mate choice detected when considering males is driven by males' choice or females' choice.

For instance, a MHC-IIB disassortative mate choice detected in divorced males may suggest

that divorced males preferentially choose MHC-IIB dissimilar females or that females

preferentially select MHC-IIB dissimilar males when they have to choose amongst divorced

males. Similarly, a lack of MHC-IIB disassortative mate choice detected when considering

first-time female breeders may be due to first-time female breeders not choosing a male

partner based on MHC-IIB dissimilarity or to males not choosing a MHC-IIB dissimilar

females when they have to choose amongst first-time female breeders.

The contrasting results between males and females might however illustrate sex-

specific mating strategies. Similarly, in Leach's storm-petrels (*Oceanodroma leucorhoa*),

while males show MHC-IIB disassortative mate choice, females' choice does not differ from

random mating [76]. In long-lived monomorphic species in which males and females equally

share parental care, such as the black-legged kittiwake [86;87;88], mate choice has been

suggested to be mutual [12]. However, the trade-off between the benefits obtained through

MHC disassortative mate choice and those obtained through mate choice based on the

intrinsic quality of the partner, such as its ability to provide good genes [89;90], good

parental care [91] or good-quality territory [92], can vary between individuals [93;89],

including between sexes. It may thus result in males and females having different preference

criteria for mate choice [94;95]. Sex-differences in the relative influence of MHC

dissimilarity vs. other traits on mating preference might be due to sex-specific variability in

either of these traits or to sex-specific variations in the direct benefits of these other traits to

the chooser. For instance, in mice, MHC dissimilarity is a more important criteria for mate

choice than scent-marking rate, only when variability in MHC dissimilarity between mates is

large [93]. In a cichlid fish (*Pelvicachromis taeniatus*), body size is a more important criteria

for mate choice than genetic compatibility for males than for females, which may be due to

the crucial role of female body size on fecundity [8].

When using Monte Carlo simulations to test mate choice, Hoover et al. [70] showed

that a sample size of 500 pairs may be necessary to confidently reject the null hypothesis

when the effect size is small (see also [73]), small effect sizes being common when studying

MHC-based mate choice [73]. In the present study, the sample size of first-time breeding

males was ca. 500 individuals, and we found a $P = 0.05$ in males, with 60 % likelihood to

find a non-significant result. This suggests at best a low effect size for mate choice based on

MHC-IIB dissimilarity in first-time male breeders. In contrast, the sample size of divorced

males was largely smaller (i.e. only 83 males), but we found a significant MHC-IIB

disassortative mate choice ($P = 0.04$) with 22 % likelihood to find a non-significant result,

suggesting a much larger effect size in divorced males than in first-time male breeders. In

several species, the level of mate selectivity is influenced by age [96;55;97;98]. In kittiwakes,

higher selectivity in divorced individuals compared to first-time breeders may be related to

their earlier arrival on the nesting site [50], which may increase both decision time and the

pool of potential partners and thereby mating decision accuracy [99]. However, like for the

sex-specific variations detected in the mate choice analysis, we cannot determine whether the

higher selectivity for MHC-IIB dissimilar partners detected when considering divorcees than

when considering first-time breeders is due to divorced individuals being more selective in

choosing MHC-IIB dissimilar males or individuals being more selective in choosing MHC-

IIB dissimilar mates when they have to choose amongst divorced partners. For instance, in

bluethroats (*Luscinia svecica*), females seem to select males based on MHC-related traits

only when paired with young males [39]. In black-legged kittiwakes, in the context of a

balance between selecting mates for MHC-IIB dissimilarity vs. other individual quality traits,

the lower inter-individual variability in individual quality in experienced breeders than in

first-time breeders [100], may explain the higher selectivity for MHC-IIB dissimilarity when

the pool of potential partners are divorcees. Clearly, while our study provides invaluable

insight into the MHC-IIB disassortative mating strategy of black-legged kittiwakes, further

studies are needed to determine which sex and age-groups are more selective or selected

based on MHC-IIB dissimilarity, and why.

In accordance with previous studies in black-legged kittiwakes [45;101;46;10], we

found that the probability of divorce was greater in pairs that failed to fledge a chick than in

pairs that reproduced successfully. However, it does not significantly vary with the MHC-IIB

dissimilarity between mates. Also, after a divorce, individuals do not mate with a partner

more MHC-IIB dissimilar to them than their original partner. Together, these results suggest

that in black-legged kittiwakes, MHC-IIB dissimilarity between the male and the female is

not a driving factor of divorce. In the present study, the proportion of divorce was much

lower than the proportions observed in other populations of kittiwakes (annual mean ± SD =

4.4 ± 2.6 % in our study *vs.* 25 % in [44]; between 13 % and 47 % in [45]; 19.1 % on average

in [46]). On Middleton island, numerous black-legged kittiwakes nest on other man-made

structures than the tower [58]. These sites were not monitored, except for occasional

demographic surveys for certain years. It may therefore have limited our ability to observe

birds that changed nesting sites between breeding season, and thus to detect all divorces,

leading to a low sample size that might have affected our results.

We did not find any support for the intermediate optimum hypothesis in mating

decisions in the black-legged kittiwake. This result is consistent with Pineaux et al. [49],

where no benefits of intermediate MHC-IIB diversity were detected in kittiwake chicks. In

this species, the benefits of a larger MHC-IIB repertoire associated with increased resistance

to wider range of pathogens may outcompete the costs associated with increased risks of

autoimmune diseases and reduced T cell receptors diversity [35;34].

A previous study on the same population of black-legged kittiwakes showed that pairs

have a higher genetic dissimilarity (as measured by microsatellites) than expected under

random mating, and that the resulting increased heterozygosity of their chicks leads to higher

fitness [102]. Although MHC-IIB disassortative mate choice may also participate in limiting

inbreeding [79;103], in the present study we detected no significant correlation between

genome-wide relatedness and MHC-IIB dissimilarity. As observed in other species (e.g. [30]

[32;104]), these results suggest that the MHC-IIB dissimilarity-based mate choice in black-

legged kittiwakes is probably driven by the MHC rather than resulting from inbreeding

avoidance at the genome-wide scale.

In conclusion, we found evidence that black-legged kittiwakes choose partners based

on MHC-IIB dissimilarity. This finding supports the role of the benefits associated with high

MHC-IIB dissimilarity in the evolution of mating decisions in this population. It is consistent

with a previous study showing that, in this species, sex-allocation strategy varies with the

MHC-IIB dissimilarity between parents [48]. In addition, our study suggests that mate choice

for MHC-IIB dissimilar partners varies with sex and age or experience of the chooser or the

selected mate. When the relative benefits of MHC disassortative mate choice vary with sex

and mate choice is mutual, a sexual conflict over mating strategies can arise and may select

for a rich repertoire of sex-specific behavioural adaptations [105;8]. Our study paves the road

for further studies of sex-specific mate choice in species with bi-parental care, a common trait

in birds [106].

References

References

[1] Andersson M., Simmons L.W. 2006 Sexual selection and mate choice. *Trends in ecology*

& evolution **21**, 296-302. (doi:10.1016/j.tree.2006.03.015)

[2] Rosenthal G.G., Ryan M.J. 2022 Sexual selection and the ascent of women: Mate choice

research since Darwin. *Science* **375**, eabi6308. (doi:10.1126/science.abi6308)

[3] Hamilton W.D., Zuk M. 1982 Heritable true fitness and bright birds: a role for parasites?

Science **218**, 384-387. (doi:10.1126/science.7123238)

[4] Andersson M., Iwasa Y. 1996 Sexual selection. *Trends in ecology & evolution* **11**, 53-58.

(doi:10.1016/0169-5347(96)81042-1)

[5] Dougherty L.R. 2021 Meta-analysis reveals that animal sexual signalling behaviour is

honest and resource based. *Nature Ecology & Evolution* **5**, 688-699.

(doi:10.1038/s41559-021-01409-z)

[6] Tregenza T., Wedell N. 2000 Genetic compatibility, mate choice and patterns of

parentage: invited review. *Molecular ecology* **9**, 1013-1027. (doi:10.1046/j.1365-

294x.2000.00964.x)

[7] Lihoreau M., Zimmer C., Rivault C. 2008 Mutual mate choice: when it pays both sexes to

avoid inbreeding. *PloS one* **3**, e3365. (doi:10.1371/journal.pone.0003365)

[8] Thünken T., Meuthen D., Bakker T.C., Baldauf S.A. 2012 A sex-specific trade-off

between mating preferences for genetic compatibility and body size in a cichlid fish

with mutual mate choice. *Proceedings of the Royal Society B: Biological Sciences*

279, 2959-2964. (doi:10.1098/rspb.2012.0333)

[9] Ihle M., Kempenaers B., Forstmeier W. 2015 Fitness benefits of mate choice for

compatibility in a socially monogamous species. *PLoS biology* **13**, e1002248.

(doi:10.1371/journal.pbio.1002248)

[10] McCully F.R., Descamps S., Harris S.M., Mckendrick F., Gillies N., Cornell S.J.,

Hatchwell B.J., Patrick S.C. 2023 Links between personality, reproductive success

and re-pairing patterns in a long-lived seabird. *Ethology* **129**, 686-700.

(doi:10.1111/eth.13405)

[11] Sullivan M.S. 1994 Mate choice as an information gathering process under time

constraint: implications for behaviour and signal design. *Animal Behaviour* **47**, 141-

151. (doi:10.1006/anbe.1994.1016)

[12] Johnstone R.A., Reynolds J.D., Deutsch J.C. 1996 Mutual mate choice and sex

differences in choosiness. *Evolution* **50**, 1382-1391. (doi:10.2307/2410876)

[13] Choudhury S. 1995 Divorce in birds: a review of the hypotheses. *Animal Behaviour* **50**,

413-429. (doi:10.1006/anbe.1995.0256)

[14] McNamara J.M., Forslund P. 1996 Divorce rates in birds: predictions from an

optimization model. *The American Naturalist* **147**, 609-640. (doi:10.1086/285869)

[15] Dubois F. Cézilly F. 2002 Breeding success and mate retention in birds: a meta-analysis.

Behavioral Ecology and Sociobiology **52**, 357-364. (doi:10.1007/s00265-002-0521-z)

[16] Culina A., Radersma R., Sheldon B.C. 2015 Trading up: the fitness consequences of

divorce in monogamous birds. *Biological Reviews* **90**, 1015-1034.

(doi:10.1111/brv.12143)

[17] Gousy-Leblanc M., Merkling T., Whelan S., Gaston A.J., Friesen V.L., Elliott K.H.

2023 Go your own way? Reasons for divorce in a monogamous seabird. *Animal*

Behaviour **204**, 13-24. (doi:10.1016/j.anbehav.2023.07.007)

[18] Streif M., Rasa O.A.E. 2001 Divorce and its consequences in the common blackbird

Turdus merula. *Ibis* **143**, 554-560. (doi:10.1111/j.1474-919x.2001.tb04882.x)

[19] Dreiss A., Roulin A. 2014 Divorce in the barn owl: securing a compatible or better mate

entails the cost of re-pairing with a less ornamented female mate. *Journal of*

evolutionary biology **27**, 1114-1124. (doi:10.1111/jeb.12402)

[20] Jones I.L., Montgomerie R. 1991 Mating and remating of least auklets (*Aethia pusilla*)

relative to ornamental traits. *Behavioral Ecology* **2**, 249-257.

(doi:10.1093/beheco/2.3.249)

[21] Fraser G.S., Jones I.L., Hunter F.M., Cowen L. 2004 Mate switching patterns in crested

auklets (*Aethia cristatella*): the role of breeding success and ornamentation. *Bird*

Behavior **16**, 7-12.

[22] Penn D.J., Potts W.K. 1999 The evolution of mating preferences and major

histocompatibility complex genes. *The American Naturalist* **153**, 145-164.

(doi:10.1086/303166)

[23] Sommer S. 2005 The importance of immune gene variability (MHC) in evolutionary

ecology and conservation. *Frontiers in zoology* **2**, 1-18. (doi:10.1186/1742-9994-2-

16)

[24] Winternitz J., Abbate J., Huchard E., Havliček J., Garamszegi L. 2017 Patterns of MHC-

dependent mate selection in humans and nonhuman primates: a meta-analysis.

Molecular ecology **26**, 668-688. (doi:10.1111/mec.13920)

[25] Özer O., Lenz T.L. 2021 Unique pathogen peptidomes facilitate pathogen-specific

selection and specialization of MHC alleles. *Molecular Biology and Evolution* **38**,

4376-4387. (doi:10.1093/molbev/msab176)

[26] Milinski M. 2006 The major histocompatibility complex, sexual selection, and mate

choice. *Annual Reviews of Ecology, Evolution and Systematics* **37**, 159-186.

(doi:10.1146/annurev.ecolsys.37.091305.110242)

[27] Lenz T.L. 2011 Computational prediction of MHC II-antigen binding supports divergent

allele advantage and explains trans-species polymorphism. *Evolution* **65**, 2380-2390.

(doi:10.1111/j.1558-5646.2011.01288.x)

[28] Oliver M.K., Telfer S., Piertney S.B. 2009 Major histocompatibility complex (MHC)

heterozygote superiority to natural multi-parasite infections in the water vole

(*Arvicola terrestris*). *Proceedings of the Royal Society B: Biological Sciences* **276**,

1119-1128. (doi:10.1098/rspb.2008.1525)

[29] Dunn P.O., Bollmer J.L., Freeman-Gallant C.R., Whittingham L.A. 2012 MHC variation

is related to a sexually selected ornament, survival, and parasite resistance in common

yellowthroats. *Evolution* **67**, 679-687. (doi:10.1111/j.1558-5646.2012.01799.x)

[30] Landry C., Garant D., Duchesne P., Bernatchez L. 2001 Good genes as heterozygosity:

the major histocompatibility complex and mate choice in Atlantic salmon (*Salmo*

salar). *Proceedings of the Royal Society of London. Series B: Biological Sciences*

268, 1279-1285. (doi:10.1098/rspb.2001.1659)

[31] Agbali M., Reichard M., Bryjová A., Bryja J., Smith C. 2010 Mate choice for

nonadditive genetic benefits correlate with MHC dissimilarity in the rose bitterling

(*Rhodeus ocellatus*). *Evolution* **64**, 1683-1696. (doi:10.1111/j.1558-

5646.2010.00961.x)

[32] Juola F.A., Dearborn D.C. 2012 Sequence-based evidence for major histocompatibility

complex-disassortative mating in a colonial seabird. *Proceedings of the Royal Society*

B: Biological Sciences **279**, 153-162. (doi:10.1098/rspb.2011.0562)

[33] Strandh M., Lannefors M., Bonadonna F., Westerdahl H. 2011 Characterization of MHC

class I and II genes in a subantarctic seabird, the blue petrel, *Halobaena caerulea*

(Procellariiformes). *Immunogenetics* **63**, 653-666. (doi:10.1007/s00251-011-0534-8)

[34] Radwan J., Babik W., Kaufman J., Lenz T.L., Winternitz J. 2020 Advances in the

evolutionary understanding of MHC polymorphism. *Trends in Genetics* **36**, 298-311.

(doi:10.1016/j.tig.2020.01.008)

[35] Migalska M., Sebastian A., Radwan J. 2019 Major histocompatibility complex class I

diversity limits the repertoire of T cell receptors. *Proceedings of the National*

Academy of Sciences **116**, 5021-5026. (doi:10.1073/pnas.1807864116)

[36] Wegner K.M., Kalbe M., Kurtz J., Reusch T.B., Milinski M. 2003 Parasite selection for

immunogenetic optimality. *Science* **301**, 1343-1343. (doi:10.1126/science.1088293)

[37] Forsberg L.A., Dannewitz J., Petersson E., Grahn M. 2007 Influence of genetic

dissimilarity in the reproductive success and mate choice of brown trout - females

fishing for optimal MHC dissimilarity. *Journal of evolutionary biology* **20**, 1859-

1869. (doi:10.1111/j.1420-9101.2007.01380.x)

[38] Baratti M., Dessi-Fulgheri F., Ambrosini R., Bonisoli-Alquati A., Caprioli M., Goti E.,

Matteo A., Monnanni R., Ragionieri L., Ristori E. and others. 2012 MHC genotype

predicts mate choice in the ring-necked pheasant *Phasianus colchicus*. *Journal of*

Evolutionary Biology **25**, 1531-1542. (doi:10.1111/j.1420-9101.2012.02534.x)

[39] Rekdal S.L., Anmarkrud J.A., Lifjeld J.T., Johnsen A. 2023 Do female bluethroats

without extra-pair offspring have more MHC-compatible social mates? *Behavioral*

Ecology and Sociobiology **77**, 36. (doi:10.1007/s00265-023-03311-z)

[40] Løvlie H., Gillingham M.A., Worley K., Pizzari T., Richardson D.S. 2013 Cryptic

female choice favours sperm from major histocompatibility complex-dissimilar

males. *Proceedings of the Royal Society B: Biological Sciences* **280**, 20131296.

(doi:10.1098/rspb.2013.1296)

[41] Richardson D.S., Komdeur J., Burke T., Von Schantz T. 2005 MHC based patterns of

social and extra pair mate choice in the Seychelles warbler. *Proceedings of the Royal*

Society B: Biological Sciences **272**, 759-767. (doi:10.1098/rspb.2004.3028)

[42] Wright D.J., Brouwer L., Mannarelli M.-E., Burke T., Komdeur J., Richardson D.S.

2016 Social pairing of Seychelles warblers under reduced constraints: MHC, neutral

heterozygosity, and age. *Behavioral Ecology* **27**, 295-303.

(doi:10.1093/beheco/arv150)

[43] Helfenstein F., Tirard C., Danchin E., Wagner R.H. 2004 Low frequency of extra-pair

paternity and high frequency of adoption in black-legged kittiwakes. *The Condor* **106**,

149-155. (doi:10.1650/7337)

[44] Coulson J. 1966 The influence of the pair-bond and age on the breeding biology of the

kittiwake gull *Rissa tridactyla*. *Journal of Animal Ecology* **35**, 269-279.

(doi:10.2307/2394)

[45] Naves L., Monnat Y.J., Cam E. 2006 Breeding performance, mate fidelity, and nest site

fidelity in a long-lived seabird: behaving against the current? *Oikos* **115**, 263-276.

(doi:10.1111/j.2006.0030-1299.14883.x)

[46] Mercier G., Yoccoz N.G., Descamps S. 2021 Influence of reproductive output on

divorce rates in polar seabirds. *Ecology and Evolution* **11**, 12989-13000.

(doi:10.1002/ece3.7775)

[47] Pineaux M., Blanchard P., Ribeiro L., Hatch S.A., Leclaire S. (2023) A gull species

recognizes mhc-ii diversity and dissimilarity using odor cues. *Symposium of Chemical*

Signals in Vertebrates, 139-151.

[48] Pineaux M., Merkling T., Danchin E., Hatch S.A., Leclaire S., Blanchard P. 2022 MHC-

II distance between parents predicts sex allocation decisions in a genetically

monogamous bird. *Behavioral Ecology* **33**, 245-251. (doi:10.1093/beheco/arab130)

[49] Pineaux M., Merkling T., Danchin E., Hatch S., Duneau D., Blanchard P., Leclaire S.

2020 Sex and hatching order modulate the association between MHC-II diversity and

fitness in early-life stages of a wild seabird. *Molecular Ecology* **29**, 3316-3329.

(doi:10.1111/mec.15551)

[50] Coulson J., White E. 1958 The effect of age on the breeding biology of the Kittiwake

Rissa tridactyla. *Ibis* **100**, 40-51. (doi:10.1111/j.1474-919x.1958.tb00364.x)

[51] Dittmann T., Becker P.H. 2003 Sex, age, experience and condition as factors affecting

arrival date in prospecting common terns, *Sterna hirundo*. *Animal Behaviour* **65**, 981-

986. (doi:10.1006/anbe.2003.2128)

[52] Bosman D.S., Vercruijsse H.J., Stienen E.W., Vincx M., Lens L. 2013 Age of first

breeding interacts with pre-and post-recruitment experience in shaping breeding

phenology in a long-lived gull. *PLoS One* **8**, e82093.

(doi:10.1371/journal.pone.0082093)

[53] Campioni L., Dias M.P., Granadeiro J.P., Catry P. 2020 An ontogenetic perspective on

migratory strategy of a long-lived pelagic seabird: Timings and destinations change

progressively during maturation. *Journal of Animal Ecology* **89**, 29-43.

(doi:10.1111/1365-2656.13044)

[54] Burley N.T., Foster V.S. 2006 Variation in female choice of mates: condition influences

selectivity. *Animal Behaviour* **72**, 713-719. (doi:10.1016/j.anbehav.2006.01.017)

[55] Huchard E., Baniel A., Schliehe-Diecks S., Kappeler P.M. 2013 MHC-disassortative

mate choice and inbreeding avoidance in a solitary primate. *Molecular Ecology* **22**,

4071-4086. (doi:10.1111/mec.12349)

[56] Fyhn M., Gabrielsen G.W., Nordøy E.S., Moe B., Langseth I., Bech C. 2001 Individual

variation in field metabolic rate of kittiwakes (*Rissa tridactyla*) during the chick-

rearing period. *Physiological and Biochemical Zoology* **74**, 343-355.

(doi:10.1086/320419)

[57] Moe B., Langseth I., Fyhn M., Gabrielsen G.W., Bech C. 2002 Changes in body

condition in breeding kittiwakes *Rissa tridactyla*. *Journal of Avian Biology* **33**, 225-

234. (doi:10.1034/j.1600-048X.2002.330304.x)

[58] Gill V.A., Hatch S.A. 2002 Components of productivity in black-legged kittiwakes *Rissa*

tridactyla: response to supplemental feeding. *Journal of Avian Biology* **33**, 113-126.

(doi:10.1034/j.1600-048X.2002.330201.x)

[59] Merkling T., Leclaire S., Danchin E., Lhuillier E., Wagner R.H., White J., Hatch S.A.,

Blanchard P. 2012 Food availability and offspring sex in a monogamous seabird:

insights from an experimental approach. *Behavioral Ecology* **23**, 751-758.

(doi:10.1093/beheco/ars023)

[60] Jodice P., Lanctot R. B., Gill V., Roby D.D., Hatch S.A. 2000 Sexing adult black-legged

kittiwakes by DNA, behavior, and morphology. *Waterbirds* **23**, 405-415.

(doi:10.2307/1522177)

[61] White J., Wagner R.H., Helfenstein F., Hatch S.A., Mulard H., Naves L.C., Danchin E.

2008 Multiple deleterious effects of experimentally aged sperm in a monogamous

bird. *Proceedings of the National Academy of Sciences* **105**, 13947-13952.

(doi:10.1073/pnas.0803067105)

[62] R Core Team (2024) R: A Language and Environment for Statistical Computing.

[63] Minias P., Pikus E., Whittingham L.A., Dunn P.O. 2018 A global analysis of selection at

the avian MHC. *Evolution* **72**, 1278-1293. (doi:10.1111/evo.13490)

[64] Sebastian A., Herdegen M., Migalska M., Radwan J. 2016 amplisas: a web server for

multilocus genotyping using next-generation amplicon sequencing data. *Molecular*

Ecology Resources **16**, 498-510. (doi:10.1111/1755-0998.12453)

[65] Leclaire S., van Dongen W.F., Voccia S., Merkling T., Ducamp C., Hatch S.A.,

Blanchard P., Danchin E., Wagner R.H. 2014 Preen secretions encode information on

MHC similarity in certain sex-dyads in a monogamous seabird. *Scientific Reports* **4**,

1-6. (doi:10.1038/srep06920)

[66] Sandberg M., Eriksson L., Jonsson J., Sjöström M., Wold S. 1998 New chemical

descriptors relevant for the design of biologically active peptides. A multivariate

characterization of 87 amino acids. *Journal of medicinal chemistry* **41**, 2481-2491.

(doi:10.1021/jm9700575)

[67] Roved J., Hansson B., Stervander M., Hasselquist D., Westerdahl H. 2022 MHCtools–an

R package for MHC high-throughput sequencing data: Genotyping, haplotype and

supertype inference, and downstream genetic analyses in non-model organisms.

Molecular Ecology Resources **22**, 2775-2792. (doi:10.1111/1755-0998.13645)

[68] Oksanen J., Kindt R., Legendre P., O'Hara B., Stevens M.H.H., Oksanen M.J., Suggests

M. 2007 The vegan package. *Community ecology package* **10**, 719.

[69] Strandh M., Westerdahl H., Pontarp M., Canbäck B., Dubois M.P., Miquel C., Taberlet

P., Bonadonna F. 2012 Major histocompatibility complex class II compatibility, but

not class I, predicts mate choice in a bird with highly developed olfaction.

Proceedings of the Royal Society B: Biological Sciences **279**, 4457-4463.

(doi:10.1098/rspb.2012.1562)

[70] Hoover B., Nevitt G. 2016 Modeling the importance of sample size in relation to error in

MHC-based mate-choice studies on natural populations. *Integrative and Comparative*

Biology **56**, 925-933. (doi:10.1093/icb/icw105)

[71] Zhang B., Hu H., Song C., Huang K., Dunn D.W., Yang X., Wang X., Zhao H., Wang

C., Zhang P. and others. 2020 MHC-based mate choice in wild golden snub-nosed

monkeys. *Frontiers in Genetics* **11**, 609414. (doi:10.3389/fgene.2020.609414)

[72] Bates D., Mächler M., Bolker B., Walker S. 2015 Fitting Linear Mixed-Effects Models

Using lme4. *Journal of Statistical Software* **67**, 1-48. (doi:10.18637/jss.v067.i01)

[73] Kamiya T., O'dwyer K., Westerdahl H., Senior A., Nakagawa S. 2014 A quantitative

review of MHC-based mating preference: The role of diversity and dissimilarity.

Molecular ecology **23**, 5151-5163. (doi:10.1111/mec.12934)

[74] Breslow N. 2004 Whither pql? *Proceedings of the Second Seattle Symposium in*

Biostatistics, 1-22. (doi:10.1007/978-1-4419-9076-1_1)

[75] Knowles J.E., Frederick C. 2024 merTools: Tools for Analyzing Mixed Effect

Regression Models.

[76] Hoover B., Alcaide M., Jennings S., Sin S.Y.W., Edwards S.V., Nevitt G.A. 2018

Ecology can inform genetics: Disassortative mating contributes to MHC

polymorphism in Leach s storm-petrels (*Oceanodroma leucorhoa*). *Molecular*

Ecology **27**, 3371-3385. (doi:10.1111/mec.14801)

[77] Sullivan G.M., Feinn R. 2012 Using effect size - or why the P value is not enough.

Journal of graduate medical education **4**, 279-282. (doi:10.4300/JGME-D-12-

00156.1)

[78] Wedekind C., Seebeck T., Bettens F., Paepke A.J. 1995 MHC-dependent mate

preferences in humans. *Proceedings of the Royal Society of London. Series B:*

Biological Sciences **260**, 245-249. (doi:10.1098/rspb.1995.0087)

[79] Penn D.J. 2002 The scent of genetic compatibility: sexual selection and the major

histocompatibility complex. *Ethology* **108**, 1-21. (doi:10.1046/j.1439-

0310.2002.00768.x)

[80] Schubert N., Nichols H.J., Winternitz J.C. 2021 How can the MHC mediate social odor

via the microbiota community? A deep dive into mechanisms. *Behavioral Ecology* **32**,

359-373. (doi:10.1093/beheco/arab004)

[81] Leclaire S., Strandh M., Mardon J., Westerdahl H., Bonadonna F. 2017 Odour-based

discrimination of similarity at the major histocompatibility complex in birds.

Proceedings of the Royal Society B: Biological Sciences **284**, 20162466.

(doi:10.1098/rspb.2016.2466)

[82] Grieves L., Gloor G., Bernards M., MacDougall-Shackleton E. 2019 Songbirds show

odour-based discrimination of similarity and diversity at the major histocompatibility

complex. *Animal behaviour* **158**, 131-138. (doi:10.1016/j.anbehav.2019.10.005)

[83] Erofeeva M.N., Alekseeva G.S., Kim M.D., Sorokin P.A., Naidenko S.V. 2022

Inbreeding coefficient and distance in MHC genes of parents as predictors of

reproductive success in domestic cat. *Animals* **12**, 165. (doi:10.3390/ani12020165)

[84] E Mingju S.X., Wang L., Yang Y., Wei X., Yu J., Gong Y., Wang H. and others. 2021

Mate choice for major histocompatibility complex (MHC) complementarity in the

Yellow-rumped Flycatcher (*Ficedula zanthopygia*). *Avian Research* **12**, 1-9.

(doi:10.1186/s40657-021-00261-w)

[85] Wojczulanis-Jakubas K., Hoover B., Jakubas D., Fort J., Grémillet D., Gavrilo M.,

Zieliňska S., Zagalska-Neubauer M. 2024 Diversity of major histocompatibility

complex of II B gene and mate choice in a monogamous and long-lived seabird, the

Little Auk (*Alle alle*). *Plos one* **19**, e0304275. (doi:10.1371/journal.pone.0304275)

[86] Coulson J., Johnson M. 1993 The attendance and absence of adult Kittiwakes *Rissa*

tridactyla from the nest site during the chick stage. *Ibis* **135**, 372-378.

(doi:10.1111/j.1474-919x.1993.tb02108.x)

[87] Leclaire S., Helfenstein F., Degeorges A., Wagner R.H., Danchin E. 2010 Family size

and sex-specific parental effort in black-legged kittiwakes. *Behaviour* **147**, 1841-

1862. (doi:10.1163/000579510x538872)

[88] Leclaire S., White J., Arnoux E., Faivre B., Vetter N., Hatch S.A., Danchin E. 2011

Integument coloration signals reproductive success, heterozygosity, and antioxidant

levels in chick-rearing black-legged kittiwakes. *Naturwissenschaften* **98**, 773-782.

(doi:10.1007/s00114-011-0827-7)

[89] Neff B.D., Pitcher T.E. 2005 Genetic quality and sexual selection: an integrated

framework for good genes and compatible genes. *Molecular ecology* **14**, 19-38.

(doi:10.1111/j.1365-294x.2004.02395.x)

[90] Puurtinen M., Ketola T., Kotiaho J.S. 2009 The good-genes and compatible-genes

benefits of mate choice. *The American Naturalist* **174**, 741-752. (doi:10.1086/606024)

[91] Moody A.T., Wilhelm S.I., Cameron-MacMillan M.L., Walsh C.J., Storey A.E. 2005

Divorce in common murres (*Uria aalge*): relationship to parental quality. *Behavioral*

Ecology and Sociobiology **57**, 224-230. (doi:10.1007/s00265-004-0856-8)

[92] Hasegawa M., Arai E., Watanabe M., Nakamura M. 2012 Female mate choice based on

territory quality in barn swallows. *Journal of ethology* **30**, 143-150.

(doi:10.1007/s10164-011-0307-8)

[93] Roberts S.C., Gosling L.M. 2003 Genetic similarity and quality interact in mate choice

decisions by female mice. *Nature genetics* **35**, 103-106. (doi:10.1038/ng1231)

[94] Bergstrom C.T., Real L.A. 2000 Toward a theory of mutual mate choice: Lessons from

two-sided matching.

[95] Aquiloni L., Gherardi F. 2008 Mutual mate choice in crayfish: large body size is selected

by both sexes, virginity by males only. *Journal of Zoology* **274**, 171-179.

(doi:10.1111/j.1469-7998.2007.00370.x)

[96] Mautz B.S., Sakaluk S.K. 2008 The effects of age and previous mating experience on

pre-and post-copulatory mate choice in female house crickets (*Acheta domesticus L.*).

Journal of Insect Behavior **21**, 203-212. (doi:10.1007/s10905-008-9120-9)

[97] Baxter C.M., Barnett R., Dukas R. 2015 Effects of age and experience on male mate

choosiness. *Ethology* **121**, 353-363. (doi:10.1111/eth.12344)

[98] Reding L., Cummings M.E. 2018 Rational mate choice decisions vary with female age

and multidimensional male signals in swordtails. *Ethology* **124**, 641-649.

(doi:10.1111/eth.12769)

[99] Chittka L., Skorupski P., Raine N.E. 2009 Speed–accuracy tradeoffs in animal decision

making. *Trends in ecology & evolution* **24**, 400-407. (doi:10.1016/j.tree.2009.02.010)

[100] Cam E., Monnat J.Y. 2000 Apparent inferiority of first-time breeders in the kittiwake:

the role of heterogeneity among age classes. *Journal of Animal Ecology* **69**, 380-394.

(doi:10.1046/j.1365-2656.2000.00400.x)

[101] Naves L.C., Cam E., Monnat J.Y. 2007 Pair duration, breeding success and divorce in a

long-lived seabird: benefits of mate familiarity? *Animal Behaviour* **73**, 433-444.

(doi:10.1016/j.anbehav.2006.10.004)

[102] Mulard H., Danchin E., Talbot S.L., Ramey A.M., Hatch S.A., White J.F., Helfenstein

F., Wagner R. H. 2009 Evidence that pairing with genetically similar mates is

maladaptive in a monogamous bird. *BMC Evolutionary Biology* **9**, 1-12.

(doi:10.1186/1471-2148-9-147)

[103] Huang W., Pilkington J.G., Pemberton J.M. 2021 Patterns of MHC-dependent sexual

selection in a free-living population of sheep. *Molecular Ecology* **30**, 6733-6742.

(doi:10.1101/2020.11.18.387332)

[104] Burger D., Thomas S., Aepli H., Dreyer M., Fabre G., Marti E., Sieme H., Robinson

M.R., Wedekind C. 2017 Major histocompatibility complex-linked social signalling

affects female fertility. *Proceedings of the Royal Society B: Biological Sciences* **284**,

20171824. (doi:10.1098/rspb.2017.1824)

[105] Parker G.A. 2006 Sexual conflict over mating and fertilization: an overview.

Philosophical Transactions of the Royal Society B: Biological Sciences **361**, 235-259.

(doi:10.1098/rstb.2005.1785)

[106] Whittingham L.A., Dunn P.O. (2001) *Male parental care and paternity in birds*. In:

Current ornithology, Springer.

Figures

Figure 1: **a-c)** Frequency distribution of means of random difference in MHC-IIB supertypes sharing between first-time breeders and their mate, calculated from 10000 bootstrap randomizations of potential pairs: a) when considering females, c) when considering males. The red vertical line is the observed mean of the difference in MHC-IIB supertypes sharing. The black vertical line corresponds to the mean of the simulated distribution, the black dashed lines are the 2.5 and 97.5 % quantiles of the simulated distribution showing the 95 % confidence intervals. **b-d)** Likelihood of a type II error according to the sample size: a) when considering females, b) when considering males. The grey dots represent the raw observations from the simulations at each sample size, the line represents the predictions of the model, the shaded area represents the 95 % confidence intervals around the predictions of the model.

Figure 2: **a-c)** Frequency distribution of means of random difference in MHC-IIB supertypes sharing between divorced breeders and their new mate, calculated from 10000 bootstrap randomizations of potential pairs, when considering a) divorced females and c) divorced males. The red vertical line is the observed mean of the difference in MHC-IIB supertypes sharing between the focal divorced individuals and their real new partner. The black vertical line corresponds to the mean of the simulated distribution, and the black dashed lines are the 2.5 and 97.5 % quantiles of the simulated distribution showing the 95 % confidence intervals. **b-d)** Probability of obtaining a significant result according to the sample size, when considering b) females and d) males. The grey dots represent the raw observations from the simulations at each sample size, the line represents the predictions of the model, the shaded area represents the 95 % confidence intervals around the predictions of the model.

775   Figure 1

    Figure 2