Interindividual variability in flower pickiness
by foraging bumblebees

Mélissa Armand[1], Christoph Beckenbauer[1], Aurore Avarguès-Weber[2], Mathieu Lihoreau[2] and Tomer J. Czaczkes[1,3]

[1]Animal Comparative Economics Laboratory, Department of Zoology and Evolutionary Biology, University of Regensburg, Germany

[2] Research Center on Animal Cognition (CRCA), Center for Integrative Biology (CBI); CNRS, Toulouse University, France

[3]Animal Cognition and Ecology Laboratory, Applied Zoology & Animal Ecology, Institute of Biology, Freie Universität Berlin

Abstract

Pollinators navigate complex and heterogeneous "flower markets", where floral resources vary in quality, availability, and spatial distribution. Bumblebees, as generalist foragers, visit numerous flowers during their foraging bouts, yet the factors influencing their flower choices and the individual differences in foraging behaviour remain poorly understood. Here, we tested how bees adjust their foraging in response to different reward structures. *Bombus terrestris* workers completed three foraging bouts in two artificial flower environments: one simulating a favourable environment with patches alternating high- and low-quality flowers (40% vs. 20% w/w sucrose solution), and the other a challenging environment with high-quality flowers alongside unrewarded ones (40% w/w sucrose solution vs. plain water). We hypothesised that bees would improve their foraging efficiency in both environments, but more rapidly in the more extreme one, where the greater reward difference creates stronger pressure to learn quickly. In both conditions, bees increased their sucrose intake per unit time over bouts. We also observed consistent differences in flower selectivity among individuals: in the favourable environment, bees that first visited high-quality flowers focused on them and avoided low-quality ones (became "picky"), while bees that first visited low-quality flowers kept visiting both types. Despite these differences, bees across environments and pickiness levels all reached similar sucrose intake rates by the third foraging bout, either by becoming more selective, collecting more sucrose solution, or reducing time spent foraging. These findings highlight the adaptability of bee foraging and suggest that early flower experiences may contribute to lasting individual differences in foraging behaviour.

Significance statement

Bumblebees are highly efficient pollinators of wild plants and commercial crops, yet their foraging behaviour varies notably among individuals. Understanding why bees differ in their foraging decisions is crucial, as it affects how they collect resources. We show that individual *Bombus terrestris* workers rapidly adapt to flower patches with different reward qualities. Remarkably, a bee's first flower visits have lasting effects: individuals that started with high-quality flowers consistently favoured them and avoided low-quality ones, while those that started with low-quality flowers continued to visit both types over time. Despite these differences, bees reached similar nectar intake rates after just three foraging trips, regardless of how selective they were. Our findings show that bees can quickly adjust their foraging behaviour, and that early experiences play a key role in how they exploit resources across different environments.

Introduction

Pollinators such as bees, bats, and birds forage in diverse and dynamic "flower markets", where collecting food efficiently is a complex task. Optimal Foraging Theory (OFT) posits that animals forage in ways that optimise net energy gain, i.e., by maximising energy intake per unit of time (Pyke, 1984; Schmid-Hempel et al., 1985; Pyke and Starr, 2021). In heterogeneous environments, there is no single way to improve foraging efficiency; animals may forage faster, select higher-quality resources, or collect larger quantities. Optimising foraging may thus be particularly challenging for generalist bees like the central-place foraging bumblebees, which can exploit hundreds of flowers each day with rates of 10 to 20 flower visits per minute (Heinrich, 1979a). Yet, bees vary considerably in how they exploit floral resources (Woodgate et al., 2016). Despite extensive research on bee foraging (see review by Sommer et al., 2022), the causes of individual differences in foraging behaviour remain unclear. Although environmental variability likely affects bee foraging, how it contributes to consistent differences between individuals is not well understood.

The natural foraging environments of bees are highly dynamic, with nectar availability and quality fluctuating across flowers and throughout the day (Goulson, 2003). For example, aging flowers can change colour and reduce nectar rewards (Weiss, 1991; Willmer, 2011). Many flowers lack nectar entirely, with up to 70% on a single plant being nectarless (Real and Rathcke, 1988; Cresswell, 1990; Thakar et al., 2003). Nectar quality also varies widely, with sugar concentrations ranging from 10% to 80% w/w (Kevan and Baker, 1983). In addition to environmental variability, bees face internal constraints that influence their foraging decisions. Wing wear, for instance, can lead bumblebees to favour denser floral patches to minimise flight effort (Foster and Cartar, 2011). Body size also plays a role: larger bees carry more nectar and tend to forage more efficiently (Heinrich, 1979a; Goulson et al., 2002; Spaethe and Weidenmüller, 2002). Experience shapes foraging as well, with experienced bees making faster, more

direct flights (Osborne et al., 2013; Woodgate et al., 2016), although learning ability varies among
individuals (Evans and Raine, 2014). Given the high energetic cost of foraging (Darveau et al., 2014;
Vaudo et al., 2016), bees must balance environmental variability and internal constraints to optimise
efficiency.

Foraging bees typically collect nectar from multiple flower types over their lifetimes (Hagbery and Nieh,
2012; Russell et al., 2017), and their floral choices vary across environments and individuals. Naïve bees
often show innate preferences for violet or blue flowers (Gumbert, 2000; Raine and Chittka, 2007), larger
flowers that are easier to locate (Ohara and Higashi, 1994; Spaethe et al., 2001), and dense flower patches
(Makino and Sakai, 2007). Bumblebees also exhibit flower constancy, the tendency to forage
predominantly on a single flower species (Chittka et al., 1999), though they do so more flexibly than
honeybees (Osborne and Williams, 2001; Gegear and Laverty, 2004). However, both innate preferences
and constancy can be overridden with experience (Gumbert, 2000; Makino and Sakai, 2007). While bees
generally prefer higher sugar concentrations (Whitney et al., 2008), they do not always choose nectar-rich
flowers. For instance, Abrol (2006) showed that honeybees favoured plants with high flowering density
and lower nectar concentration over patches with fewer, nectar-rich flowers. Similarly, smaller-size
bumblebees invested equal effort in memorising flowers with high and low sugar concentration (Frasnelli
et al., 2021). Preferences for nectar volume have received less attention, and findings are inconsistent
(Menzel and Erber, 1972; Waddington and Gottlieb, 1990; Lihoreau et al., 2011), although bees appeared
more responsive to changes in nectar concentration than volume (Cnaani et al., 2006). Floral choices are
also shaped by bees' energetic demands, as they adjust foraging to meet both individual needs and those
of the colony (Molet et al., 2008; Vaudo et al., 2016; Hendriksma et al., 2019).

Depending on environmental conditions, bees may use different foraging "tactics" to navigate floral
resources. In simpler flower patches, they often follow a nearest-neighbour rule, moving sequentially to
the closest unvisited flower until resources are depleted (Saleh and Chittka, 2007; Ohashi et al., 2007). In
denser or more aggregated patches, bees adopt a near-far search rule, making short flights between
rewarding flowers and staying longer in nectar-rich areas, but leaving quickly or taking longer flights
when rewards decline (Dukas and Real, 1993; Chittka et al., 1997; Cartar, 2004). At larger spatial scales,
bees alternate between exploration and exploitation flights to establish foraging routes (Kembro et al.,
2019). A common pattern is traplining: repeatedly visiting flowers in a fixed sequence to maximise
rewards while avoiding already depleted sites (Ohashi et al., 2007; Lihoreau et al., 2010). Bees gradually
refine these routes to balance shorter paths with prioritising high-reward flowers (Reynolds et al., 2013).
Still, foraging patterns in heterogeneous environments are highly variable (Lihoreau et al., 2012), and
individual bees differ in patch fidelity, the exploration–exploitation balance, and the duration and
frequency of foraging bouts (Woodgate et al., 2016). We propose that environmental conditions shape
lasting individual differences in flower choice and foraging behaviour.

Here, we conducted a lab-based experiment to test how bees improve foraging efficiency in
heterogeneous flower environments. We examined how the quality difference between good and poor
flowers within patches influences the foraging behaviour of *Bombus terrestris* workers. Individual bees
completed three consecutive foraging bouts in two distinct artificial flower environments. The first
simulated a favourable environment, with patches containing alternating high- and low-quality flowers
("High vs. Low" environment; 40% vs. 20% w/w sucrose solution). The second represented a challenging
environment, with patches of rewarding flowers alongside unrewarded ones ("High vs. Water"
environment; 40% w/w sucrose solution vs. plain water). We hypothesised that environmental context
would shape bees' subsequent foraging behaviour. Specifically, we predicted that the more extreme
conditions in the High vs. Water environment would allow bees to learn flower quality more rapidly and
improve foraging efficiency faster than in the High vs. Low environment. Nonetheless, we expected that
bees in both environments would similarly increase their energy intake rate with experience, consistent
with optimal foraging theory (Pyke, 1984; Schmid-Hempel et al., 1985).

Material and Methods

Bees

Six commercial *B. terrestris* colonies (Koppert, The Netherlands) were maintained under
controlled laboratory conditions (22–24°C, 14:10 light:dark cycle; lights on from 6:30 a.m. to 8:30 p.m.)
with ad libitum pollen. Colonies were housed in wooden nestboxes connected to individual flight arenas
(60 x 50 x 35 cm) via transparent tubes fitted with shutters (see apparatus in **Fig. 1**). Between
experimental sessions (i.e., pre-training), bees had unrestricted access to six artificial flowers providing ad
libitum 20% w/w sucrose solution through a damp cotton mesh soaked in solution and protruding through
a central hole. Bees that foraged regularly were individually marked on the thorax with uniquely
coloured, numbered tags and selected for the experiment.

A total of 52 bees from four colonies were tested over 20 days (October–November 2021; see
Supplement S1 for colony details). Additionally, six bees from two other colonies were tested in January
2024 in a complementary experiment to determine whether individuals collected the entire sucrose
solution drop from the artificial flowers or only a portion of it. We conducted this control to validate
whether sucrose solution consumption could be accurately estimated based on reward collection. This
setup mirrored the main experiment but reversed flower colours and included weighing flowers before
and after each foraging bout. Results showed that bees typically either fully collected the sucrose solution
or rejected the flower without drinking (see **Supplement S2** for detailed analysis).

Artificial flowers

Each flower consisted of a transparent plastic cup (38 mm diameter, 70 mm height) topped
with a coloured laminated paper disk of the same diameter. Flowers used during the experiment were
either blue (peak reflectance ≈ 450 nm) or yellow (peak reflectance ≈ 530–700 nm; see **Supplement**
S3 for reflectance curves). Sucrose solution was pipetted onto a small white dot in the centre of each
flower to aid the bees in locating the rewards (Heuschen et al., 2005; see **Fig. 1**). During pre-training,
flowers were made of two halves, one blue and one yellow, following the method of Raine and Chittka
(2008). These bicoloured flowers ensured bees associated sucrose rewards equally with both colours,
which were later used as distinct flower types in the experiment.

Figure 1: Top view of the scaled experimental setup. The 3D model, created in Blender, illustrates the
layout of pattern A of the flower environment (see **Supplement S5** for all layout patterns). Flower
patches were positioned 15–50 cm from the arena entrance and mounted on transparent white circular

plastic bases (15 cm diameter, 0.5 cm thick), with flowers within each patch spaced 1.5 cm apart. Flowers

were removed and rearranged after each foraging bout using the arena's side doors.

Experiment

Individual bees completed three consecutive foraging bouts on a flower environment consisting

of five dimorphic flower patches, each with six flowers arranged in a circle of alternating blue and yellow

colours. This ensured that neighbouring flowers were always of different colours. Patches were arranged

using four distinct layout patterns (see **Supplement S5**), and each bee experienced a random sequence of

three different patterns. This pseudo-random design prevented reliance on spatial memory while keeping

layouts consistent across bees.

In the main experiment, yellow flowers consistently offered high-concentration sucrose solution to

counteract bees' innate preference for blue (Gumbert, 2000) and avoid a colour bias. To further test the

robustness of the setup, colour assignments were reversed in the complementary experiment. Bees were

tested on one of two flower environment types:

(1) High vs. Low flower environment (N = 37 bees from 6 colonies): For 31 bees, yellow

flowers contained 14 µL of concentrated sucrose solution (40% w/w) and blue flowers 14 µL of diluted

sucrose solution (20% w/w). For the six bees in the complementary experiment, the colours were

reversed. This simulated a favourable environment where all flowers were rewarding but varied in

quality. We expected bees in this environment to be less selective in their flower choices and to show

slower improvement in foraging efficiency compared to the environment with more extreme differences

in flower rewards described below.

(2) High vs. Water flower environment (N = 15 bees from 3 colonies): Yellow flowers

contained 14 µL of concentrated sucrose solution (40% w/w), and blue flowers 14 µL of plain water (tap

water). This represented a more challenging environment, with only half of the flowers offering a reward.

We expected bees in this environment to be highly selective and show rapid improvement in foraging

efficiency.

In both environments, each flower type provided a total of 210 µL of sucrose solution per bout, enough

for a bee to fill its crop by visiting a single flower type. This volume exceeds the average crop capacity of

bumblebee foragers (120–180 µL; Lihoreau et al., 2010). To prevent chemical marks from influencing

flower choices, all flower surfaces were wiped clean with a paper towel moistened with 70% ethanol

between foraging bouts (Goulson et al., 2000; Saleh et al., 2007).

After the third training bout, bees completed a final test bout in which the arena contained two

monochromatic flower patches, one yellow and one blue, both unrewarded (plain water). The patches

were positioned against the back wall, facing the arena entrance, and the bee's first flower visit was
recorded as an indicator of its flower type preference.

Data collection and analysis

Each foraging bout was video-recorded using a camera (Sony HDR CX405) mounted above the
flight arena. Bee behaviour was analysed using an event-logging software (BORIS, v8.6), recording: (1)
foraging bout duration (i.e., time from entering the arena to returning to the nest), (2) flower visits (bee
landing on a flower), (3) flower revisits (bee landing on a previously visited flower within the same bout),
and (4) reward collection (bee licking a drop of sucrose solution for more than five seconds). Based on
video analysis, bees were classified as either rejecting the flower or accepting and fully consuming the
sucrose drop (see **Supplement S6** for video examples). This categorisation was supported by the
complementary experiment, in which flowers were weighed before and after each foraging bout (see
Supplement S2 for methodology and results). Bees that accepted a flower overwhelmingly consumed the
entire drop, regardless of whether it was high- or low-concentration sucrose.

Data processing was performed with Python (v3.11, Python Software Foundation, 2023) using the
following libraries: *pandas* (McKinney, 2010) for data structuring, *seaborn* (Waskom, 2021) and
Matplotlib (Hunter, 2007) for data visualisation, and *Scikit-learn* (Pedregosa et al., 2011) for data
clustering. Statistical analyses were carried out with R (v4.1, R Core Team 2022) using the packages
glmmTMB (Brooks et al., 2017) for Generalised Linear Mixed Models (GLMMs) and *emmeans* (Lenth,
2020) for post-hoc comparisons. Model residuals were evaluated using the *DHARMa* package (Hartig,
2020).

Datasets from the main and complementary experiments, where flower colours associated with reward
were reversed, were merged for analysis. To account for potential effects of flower colour, experiment
type (main vs. complementary) was included as a fixed effect in all models. Individual bees were treated
as random effects nested within colonies to account for within-colony variation. All statistical analyses
and datasets are available on Zenodo (https://doi.org/10.5281/zenodo.16418595).

Bee clustering

205   During initial analysis of bees' foraging behaviour in the High vs. Low flower environment, we
206   observed substantial dispersion in the proportions of visits to high-quality flowers, collection of high-
207   concentration sucrose, and rejection of low-quality flowers, suggesting notable interindividual variability
208   in flower pickiness. To assess this variation, we applied K-means clustering (k = 2) to the bees based on
209   three behavioural variables measured during the final foraging bout:

1. **Proportion of visits to high-quality flowers**: number of visits to high-quality flowers divided by total visits. A visit is defined as a landing on a flower, regardless of whether the bee collected or rejected the sucrose solution;

2. **Proportion of high-concentration sucrose solution collected**: number of high-quality flowers collected (i.e., solution depleted) divided by total flowers collected;

3. **Proportion of low-concentration sucrose solution flowers rejected**: number of non-empty low-quality flowers visited but not collected (i.e., bees inspected but rejected the sucrose solution), divided by total visits to low-quality flowers.

Each variable was standardised (converted to z-scores by subtracting the mean and dividing by the standard deviation) and clustering was applied directly to these standardised values, identifying two groups of bees with distinct foraging behaviours (**Fig. 2B;** see detailed analysis in **Supplement S1**):

A. **High pickiness** cluster (N = 28 bees): bees in this group predominantly visited high-quality flowers, collected a high proportion of high-concentration sucrose, and strongly rejected low-concentration flowers

B. **Low pickiness** cluster (N = 9 bees): bees visited high- and low-quality flowers more evenly, collected lower proportions of high-concentration sucrose, and often accepted low-quality flowers instead of rejecting them.

Estimation of absolute sucrose intakes

Absolute sucrose intake per foraging bout was inferred from video analysis, based on the number of high- and low-quality flowers collected and the theoretical sucrose content of 20% and 40% w/w sucrose solutions. These calculations assume that bees consumed the entire sucrose drop upon collection, a behaviour confirmed in our complementary experiment. Calculations were based on the following values (Hofmann, 1977):

1. **20% w/w sucrose solution**: With a density of 1.081 g/mL, a 14 µL drop weighs approximately 15.134 mg, comprising 3.027 mg of sucrose and 12.106 mg of water.

2. **40% w/w sucrose solution**: With a density of 1.176 g/mL, a 14 µL drop weighs approximately 16.464 mg, comprising 6.588 mg of sucrose and 9.882 mg of water.

Results

We combined data from the main and complementary experiments (the latter reversed colour–reward pairings) for analysis. Despite the reversal, bees showed similar behavioural trends. Naïve bees tended to visit yellow flowers first: 60.9% (18/28) in the main experiment and 67% (4/6) in the complementary

experiment. To test whether an innate preference for yellow might have influenced foraging behaviour across bouts, we included "experiment type" (main vs. complementary) as a fixed effect in all models. This variable had no significant effect, indicating that flower colour did not drive the observed behavioural patterns (see **Supplement S2** for model details).

1. Foraging efficiency in the favourable vs. challenging environments

We first tested whether environmental context influenced how bees adjusted their foraging behaviour. We predicted that bees in the challenging environment (40% vs. water) would learn to distinguish flower quality faster than those in the favourable environment (40% vs. 20%), and that bees in both environments would improve their foraging efficiency with experience.

(A) Estimated sucrose intake rate

Bees in both treatments significantly increased their estimated sucrose intake rate across bouts (GLMM, χ^2 = 84.23, df = 2, p < 0.0001; N = 52 bees), consistent with expectations from Optimal Foraging Theory. Estimated intake rates did not differ between environments at any bout (post-hoc Tukey tests, challenging vs. favourable environment, bout 1: 10.8 mg/min ± 1.19 vs. 12.1 mg/min ± 0.82, z-ratio = 0.93, p = 0.35; bout 2: 17.3 ± 1.69 vs. 15.9 ± 1.01, z-ratio = -0.88, p = 0.38; bout 3: 18.8 ± 1.81 vs. 19.2 ± 1.17, z-ratio = 0.24, p = 0.81; **Fig. 2B**).

(B) Proportion of visits to high-quality flowers

Bees in the challenging environment became more selective over time, visiting high-reward flowers more frequently than those in the favourable environment (GLMM, treatment: χ^2 = 9.13, df = 1, p = 0.0025). There was no difference in bout 1 (post-hoc Tukey test, challenging vs. favourable environment: 58.7% ± 6.0% vs. 52.4% ± 4.4%, t-ratio = -1.18, p = 0.24; N = 52 bees), but significant differences emerged in bout 2 (81.5% ± 6.0% vs. 61.0% ± 4.4%, t-ratio = -3.83, p < 0.001) and bout 3 (84.1% ± 6.0% vs. 70.1% ± 4.4%, t-ratio = -2.61, p = 0.010; **Fig. 2C**).

2. Bees in the favourable environment rapidly increased foraging efficiency

We then examined whether bees foraging on patches with both high- and low-quality flowers favoured the high-concentration sucrose solution or collected solution from both flower types. Over successive foraging bouts, bees significantly increased: (1) their proportion of visits to high-quality flowers, calculated as the number of visits to high-quality flowers divided by total visits (post-hoc Tukey test, bout 1 vs. bout 3: 50.5% ± 4.2% vs. 68.3% ± 4.2%, t-ratio = -6.20, p < 0.0001; N = 37 bees; **Fig. 2A**); (2) their proportion of high-concentration sucrose solution collected, calculated as the number of high-quality flowers collected divided by total flowers collected (post-hoc Tukey test, bout 1 vs. bout 3: 55.6% ± 4.6% vs. 87.7% ± 4.7%, t-ratio = -10.55, p < 0.0001; N = 37 bees; **Fig. 2A**), and (3) their

proportion of low-concentration sucrose solution rejected, calculated as the number of non-depleted low-
quality flowers visited and not collected, divided by total visits to low-quality flowers (post-hoc Tukey
test, bout 1 vs. bout 3: 29.4% ± 7.0% vs. 74.0% ± 6.2%, t-ratio = -5.51, $p < 0.0001$; N = 37 bees; **Fig.
2A**).

(A) Bees varied in their selectivity towards high-quality flowers

The K-means clustering analysis identified two distinct bee clusters: High pickiness (N = 28 bees;
23 from the main experiment and 5 from the complementary experiment) and Low pickiness (N = 9 bees;
8 from the main experiment and 1 from the complementary experiment).

These clusters reflect marked differences in foraging behaviour (**Fig. 2D**). Bees in the High pickiness
cluster visited high-reward flowers in 75% of visits on average, collected 97.8% high-concentration
sucrose, and rejected 98.1% of low-quality flowers. In contrast, bees in the Low pickiness cluster visited
high-reward flowers in only 51.8% of visits, collected 57.3% high-concentration sucrose, and rejected
only 36.6% of low-quality flowers. These proportions illustrate clear behavioural differences between the
two groups.

(B) Bees' degree of pickiness was correlated with the quality of the first flowers visited

Bees in the High pickiness cluster tended to have initially visited high-quality flowers more often
than those in the Low pickiness cluster (post-hoc Tukey test, High pickiness cluster vs. Low pickiness
cluster: 60.1% ± 16.4% vs. 27.6% ± 19.4%; z = 1.48, p = 0.138, N = 37 visits; **Fig. 2E**). This effect
became statistically significant when considering only the first two flower visits (post-hoc Tukey test,
High pickiness cluster vs. Low pickiness cluster: 61% ± 8.13% vs. 28.2% ± 11.36%; z = 2.19, p = 0.028,
N = 74 visits; **Fig. 2E**). This finding suggests that early exposure to high-quality flowers may drive bees
to become more selective, whereas initial visits to low-quality flowers might reduce selectivity in
subsequent choices.

**3. Bees across environments and clusters improved foraging efficiency via different
trade-offs**

Next, we examined the foraging behaviour of bees across foraging bouts, comparing flower
environment (High vs. Low and High vs. Water) and clusters (High pickiness and Low pickiness). All
groups significantly increased their estimated sucrose intake per unit of time with each successive bout
(GLMM: χ^2 = 90.76, df = 2, $p < 0.0001$, N = 52 bees). This improvement rate was consistent across
groups at the same foraging stage (GLMM: χ^2 = 0.77, df = 2, p = 0.68). We analysed the foraging
behaviour of each bee group in more detail:

(A) Low pickiness bees

Low pickiness bees maintained consistent foraging durations across bouts (post-hoc Tukey test,
bout 1 vs. bout 3: 3.89 min ± 0.34 vs. 3.17 min ± 0.45, z-ratio = 1.01, p = 0.57; **Fig. 3A**). They increased
the number of flowers collected (post-hoc Tukey test, bout 1 vs. bout 3: 10.8 flowers ± 1.02 vs. 12.8
flowers ± 1.18, z-ratio = -2.70, p = 0.019; **Fig. 3B**) but did not significantly increase the proportion of
high-concentration sucrose solution collected (post-hoc Tukey test, bout 1 vs. bout 3: 39.1% of high-
quality flowers collected ± 6.74 vs. 55.2% ± 6.57, z-ratio = -1.81, p = 0.17; **Fig. 3C**).

The bees reduced revisits to previously depleted flowers (post-hoc Tukey test, bout 1 vs. bout 3: 7.57
revisits ± 1.87 vs. 4.7 revisits ± 1.23, z-ratio = 2.63, p = 0.02; **Fig. 3D**). Overall, bees from this cluster
showed an increase in estimated sucrose intake across bouts (post-hoc Tukey test, bout 1 vs. bout 3: 48.1
mg ± 5.07 vs. 64.2 mg ± 6.58, z-ratio = -4.47, p < 0.0001; **Fig. 3E**). In the final test, 55.6% of bees first
visited the high-quality flower patch (**Fig. 3F**).

(B) High pickiness bees

By contrast, High pickiness bees decreased their foraging duration over bouts (post-hoc Tukey
test, bout 1 vs. bout 3: 6.39 min ± 0.56 vs. 3.89 min ± 0.34, z-ratio = 6.18, p < 0.0001; **Fig. 3A**). They
decreased the number of flowers collected (post-hoc Tukey test, bout 1 vs. bout 3: 13.2 flowers ± 0.76 vs.
11 flowers ± 0.64, z-ratio = 5.12, p < 0.0001; **Fig. 3B**) while strongly shifting their collection of sucrose
solution towards high-quality flowers (post-hoc Tukey test, bout 1 vs. bout 3: 61.6% of high-quality
flowers collected ± 3.59 vs. 97.6% ± 1.01, z-ratio = -7.18, p < 0.0001; **Fig. 3C**).

The bees reduced flower revisits (post-hoc Tukey test, bout 1 vs. bout 3: 12.73 revisits ± 1.99 vs. 9.88
revisits ± 1.57, z-ratio = 3.26, p = 0.003; **Fig. 3D**), while their estimated sucrose intakes remained stable
across bouts (post-hoc Tukey test, bout 1 vs. bout 3: 68 mg ± 4.27 vs. 71.8 mg ± 4.5, z-ratio = -1.69, p =
0.21; **Fig. 3E**). In the final test, 92.6% of bees first visited the high-quality flower patch (**Fig. 3F**).

(C) Bees in the High vs. Water flower environment

In this flower environment, where only high-concentration sucrose solution was available, bees
nearly halved their foraging duration over successive bouts (post-hoc Tukey test, bout 1 vs. bout 3: 6.74
min ± 0.83 vs. 3.89 min ± 0.34, z-ratio = 5.96, p < 0.0001; **Fig. 3A**) while maintaining a steady number of
flowers collected (post-hoc Tukey test, bout 1 vs. bout 3: 10.8 flowers ± 0.91 vs. 10.1 flowers ± 0.86, z-
ratio = 1.25, p = 0.42; **Fig. 3B**).

Bees reduced flower revisits by nearly threefold (post-hoc Tukey test, bout 1 vs. bout 3: 24.6 revisits ±
4.95 vs. 9.01 revisits ± 1.9, z-ratio = 10.64, p < 0.0001; **Fig. 3D**), while their estimated sucrose intakes
remained consistent across bouts (post-hoc Tukey test, bout 1 vs. bout 3: 71.5 mg ± 6.26 vs. 66.5 mg ±
5.85, z-ratio = 1.61, p = 0.24; **Fig. 3E**). In the final test, 86.7% of bees first visited the high-quality flower
patch (**Fig. 3F**).

Figure 2: Foraging efficiency across foraging bouts. A) Box plots of the proportions of bees (1) visiting high-quality flowers, (2) collecting high-quality flowers, and (3) rejecting low-quality flowers across foraging bouts (N = 37 bees); **B)** Box plots of estimated sucrose intake rates (mg/min) in each flower environment; **C)** Box plots of the proportion of visits to high-quality flowers in each environment. Boxes show the interquartile range (IQR), horizontal lines indicate the median, and whiskers extend to the smallest and largest values within 1.5×IQR. **D)** Bar plots of the mean proportions of variables: (1) visits to high-quality flowers, (2) high-quality flowers collected, and (3) low-quality flowers rejected for each pickiness cluster; **E)** Bar plots of the proportions of high-quality flowers visited during the first flower visits for each bee cluster. Error bars represent standard errors of the mean. Statistical significance is indicated as ******* ($p < 0.001$), ****** ($p < 0.01$), ***** ($p < 0.05$), and **n.s.** ($p > 0.05$, post-hoc Tukey tests).

Figure 3: Foraging behaviour across environments and pickiness levels. A) Foraging duration per foraging bout (minutes); **B)** Number of flowers collected per bout; **C)** Proportion of high-concentration sucrose solution collected per bout; **D)** Number of flowers revisited per bout; **E)** Estimated sucrose intake per foraging bout (mg), and **F)** Proportion of bees whose first visit was to the high-quality flower in the final test (N = 52 bees). Each boxplot shows the median (centre line), the interquartile range (IQR), and whiskers extending to the smallest and largest values within 1.5×IQR. Error bars represent standard errors of the predicted probabilities. In panel C, probabilities for the High vs. Water environment are equal to 1 in each bout because only high-quality flowers were available. Statistical significance is indicated as ******* ($p < 0.001$), ****** ($p < 0.01$), ***** ($p < 0.05$), and **n.s.** ($p > 0.05$, post-hoc Tukey tests).

Discussion

This study investigated how differences between high- and low-quality flowers in mixed-quality patches influence bees' flower choices, addressing a gap in our understanding of interindividual variability in bumblebee foraging behaviour. We found that foragers rapidly improved their efficiency across successive bouts, reaching similar sucrose intake rates by the third bout regardless of flower environment type, whether the environment was "favourable" (containing both high- and low-quality flowers) or "challenging" (where only half the flowers were rewarding). This finding supports our prediction and aligns with Optimal Foraging Theory (OFT), which posits that nectarivores should maximise their net energy intake rate (Heinrich, 1979a; Waddington and Holden, 1979; Pyke and Starr, 2021). Additionally, we observed substantial interindividual variability in foraging behaviour, particularly in flower selectivity: some bees strongly preferred high-quality flowers, while others were less selective. This is consistent with previous findings of high individual variability in bumblebee foraging behaviour in heterogeneous environments (Saleh and Chittka, 2007; Lihoreau et al., 2011; Woodgate et al., 2016).

A key prediction of our study was that bees in the challenging environment would more quickly learn to distinguish high-quality flowers, due to the stronger contrast between flower types. This was supported by their increased visits to high-quality flowers in foraging bouts 2 and 3, compared to bees in the favourable environment. While this may reflect faster learning, bees in the favourable environment may have intentionally continued visiting and collecting both flower types, as both offered some reward (see next section). Notably, bees in the challenging environment were even more selective than high-pickiness bees in the favourable environment, suggesting that harsher conditions may have indeed accelerated learning. Nevertheless, bees in both environments ultimately achieved similar estimated sucrose intake rates in each bout, highlighting their behavioural flexibility in adapting to different environments.

Bees in the favourable environment showed marked individual differences in how they improved foraging efficiency. High-pickiness bees collected smaller volumes of more concentrated sucrose, prioritising quality over quantity. In contrast, low-pickiness bees increased the total volume collected, regardless of

concentration. Although both groups learned and improved over successive bouts, these differences in
foraging behaviours remained consistent over time. This mirrors findings by Chittka et al. (2003), where
individual bees consistently differed in speed-accuracy trade-offs throughout learning. Similar
behavioural trade-offs have been reported in other studies, with bees balancing speed and accuracy (Burns
and Dyer, 2008) or balancing reward value against foraging effort (Lihoreau et al., 2011; Pattrick et al.,
2023). Such individual variability may be adaptive in unpredictable environments, where resource quality
and availability vary across space and time. Moreover, it can enhance colony performance by diversifying
foraging roles (Muller and Chittka, 2008; Holland et al., 2021). Supporting this, Hagbery and Nieh (2012)
found that generalist bumblebees, those not specialising strictly in nectar or pollen, adjusted their foraging
behaviour flexibly in response to changes in colony needs.

Interestingly, most bees did not increase their estimated absolute sucrose intake across bouts; only low-
pickiness bees showed an increase. Bees in this cluster had lower estimated intakes during their first two
foraging bouts compared to both high-pickiness bees and bees foraging in the challenging environment
with only one rewarding flower type. For high-pickiness bees, their focus on high-quality flowers likely
constrained their intake over time. In our setup, only half the flowers in each patch were rewarding,
increasing the likelihood of revisiting depleted flowers as the best rewards were collected. Previous
studies have shown that bees do not visit all flowers in a patch (Goulson, 2000; Hemingway et al., 2024),
and according to the optimal departure rule, pollinators should leave a patch when their reward intake rate
declines (Cresswell, 1990; Goulson, 2000).

However, we found that naïve bees, regardless of group, did not collect substantially less sucrose during
their first foraging bout, despite longer foraging times and frequent revisits to empty flowers. This
suggests that bees did not abandon patches based on declining reward rates; if they had followed an
optimal departure rule, we would expect lower sucrose intakes in early bouts. That said, our setup offered
no alternative foraging sites, so bees may have continued exploring simply because there was nowhere
else to go. Their behaviour likely reflects exploration of a novel environment, similar to the orientation
flights seen in wild bumblebees (Woodgate et al., 2016).

Surprisingly, high-pickiness bees in the favourable environment collected over 15% fewer flowers in their
final bout compared to their first, while maintaining stable sucrose intake. This pattern suggests that bees
may return to the nest after reaching a fixed energetic threshold, rather than filling their crop or foraging
for a set duration. It also supports the idea that bees may return with partially empty crops (see Schmid-
Hempel et al., 1985). However, explicit testing is necessary to confirm this fixed energetic target
hypothesis.

One result worth further consideration is that all bee groups in our study improved their foraging
efficiency at a similar rate, reaching comparable estimated sucrose intake rates at each bout. This suggests
that bees may be capable of compensating for less favourable initial conditions by adjusting their foraging

behaviour. However, this finding should be interpreted with caution, as we did not directly measure sucrose intake but instead inferred it through video analysis and calculations (refer to Data collection and analysis section). Importantly, our estimates may underestimate intake from low-quality flowers, which were often probed for less than a second and thus classified as non-collected. Yet, Lechantre et al. (2021) demonstrated that bumblebees can imbibe up to 2 µL of solution per second, suggesting that even very brief visits may still result in meaningful reward intake.

Our study also raises questions about what shapes individual foraging behaviours, especially flower-quality pickiness. In the favourable environment of high- and low-quality flowers, most bees quickly became highly selective for the high-quality flowers. Notably, pickiness was associated with bees' initial flower choices, which were most often to high-concentration sucrose flowers. While this pattern may reflect reward-driven learning, it is also possible that innate colour preferences influenced flower choices, as most high-concentration flowers in the main experiment were yellow. To test this, we reversed the colour-reward associations in a complementary experiment and included experiment type (main vs. complementary: yellow vs. blue as the high-quality flower) as a fixed effect in all statistical models. This variable had no significant effect on foraging behaviours, suggesting that the observed pickiness was primarily driven by reward quality rather than colour.

Previous studies have shown that bees tend to remain constant to flower types offering high rewards but explore alternative options when rewards are low (Chittka et al., 1999; Grüter and Ratnieks, 2011; Nityananda and Chittka, 2020). This could explain why bees in our study strongly favoured high-quality flowers after initial exposure, whereas those first encountering low-quality flowers displayed more variability in their flower choices. Although flower constancy was long considered a cognitive limitation (Waser, 1986; Chittka et al., 1999), more recent work suggests it can function as an adaptive strategy (Gegear and Thomson, 2004). Supporting this, recent model simulations have shown that colonies composed of highly flower-constant individuals exploited only 30–50% of available flower species, whereas colonies of indiscriminate foragers could exploit nearly all species (Grüter et al., 2024).

Can pickiness be solely attributed to the quality of a bee's first few flower visits? The moderate strength of the correlation suggests that other factors likely influenced pickiness as well. For instance, bees vary in their sucrose sensitivity thresholds, which may drive differences in preference for reward quality (Page et al., 2006; Riveros & Gronenberg, 2010). Additionally, differences in learning ability among bumblebees (Raine et al., 2006) might explain why bees in the low pickiness cluster did not favour the high-quality patch in the final test. While all bees in our study increased their visits to high-quality flowers over successive bouts, they did so at different rates. Would every bee have eventually become highly selective in our setup? Adjusting foraging behaviour can be energetically costly, especially in dynamic environments (DeWitt et al., 1998). A promising research direction would be to manipulate initial

conditions to test whether, and when, bees converge on the same optimal foraging behaviour. This could provide insight into the limits of behavioural flexibility.

It is worth noting that most high-concentration flowers in our experiment were yellow. We had expected bees to favour blue flowers on their first visit, given well-documented innate preferences (Gumbert, 2000; Raine & Chittka, 2007). However, over two-thirds of initial visits were to yellow flowers. This unexpected bias may be due to yellow flowers reflecting more light across a broader wavelength range, making them appear brighter overall (see reflectance curves in **S3**). Brightness is known to strongly influence bee preferences (Sletvold et al., 2016) and may explain this initial preference, even though blue flowers in our setup had higher spectral purity and saliency in the white arena—traits typically associated with greater attractiveness to bees (Lunau, 1990; Goulson, 2000).

While our findings suggest that bees aimed to optimise energy intake, not all energetic costs of foraging were accounted for. For example, flight costs were minimal in our setup, as flowers within patches were close enough for bees to walk between them, potentially conserving energy. Energy expenditure also likely varied with the weight of the sucrose loads carried by each bee (Wolf et al., 1989; Combes et al., 2020). Although bees learned quickly in our simple flower environments over just a few bouts, natural environments are far more complex, with less predictable and more subtle differences between high- and low-reward flowers. Our small flight arena also represented a "local" foraging scale (Sommer et al., 2022), where bees likely encountered floral cues almost immediately (Heinrich, 2004)—a contrast to natural foraging conditions. Still, our findings highlight the rapid adaptability of bumblebee foraging behaviour and point to promising future research directions: exploring how environmental conditions, initial experience, and foraging behaviour interact could provide new insights into the ecological success of bumblebees across diverse and dynamic habitats.

Supplementary Material

S1 presents the statistical analysis of the main experiment, while **S2** provides the statistical analysis of the complementary experiment. **S3** includes the reflectance curves of the artificial flowers, and **S4** contains the reflectance dataset. **S5** shows the layout patterns of the flower environments. **S6** features video examples of bees foraging during the experiment. **S7** and **S8** contain the datasets for the main and complementary experiments, respectively. All statistical analyses and datasets are publicly available on Zenodo (https://doi.org/10.5281/zenodo.16418595).

Acknowledgments

We thank Dorian Champelovier for his help with data collection, and the anonymous reviewers for their thoughtful feedback which helped improve the manuscript.

Funding

M. A. was supported by an ERC Starting Grant to T. J. C. [H2020-EU.1.1. #948181]. M. L. was supported by an ERC Consolidator Grant [H2020-EU.1.1. #101002644]. T. J. C. was supported by a Heisenberg Fellowship from the Deutsche Forschungsgemeinschaft [#462101190].

References

Abrol DP (2006) Foraging behaviour of bees as influenced by quality and quantity of rewards from flowers. Journal of Asia-Pacific Entomology 9:145–148. https://doi.org/10.1016/S1226-8615(08)60285-X

Brooks ME, Kristensen K, van Benthem KJ, et al (2017) glmmTMB balances speed and flexibility among packages for zero-inflated generalized linear mixed modeling. The R journal 9:378–400. https://doi.org/10.3929/ethz-b-000240890

Burns JG, Dyer AG (2008) Diversity of speed-accuracy strategies benefits social insects. Current Biology 18:R953–R954. https://doi.org/10.1016/j.cub.2008.08.028

Cartar RV (2004) Resource tracking by bumble bees: responses to plant-level differences in quality. Ecology 85:2764–2771. https://doi.org/10.1890/03-0484

Chittka L, Dyer AG, Bock F, Dornhaus A (2003) Bees trade off foraging speed for accuracy. Nature 424:388–388. https://doi.org/10.1038/424388a

Chittka L, Gumbert A, Kunze J (1997) Foraging dynamics of bumble bees: correlates of movements within and between plant species. Behavioral Ecology 8:239–249. https://doi.org/10.1093/beheco/8.3.239

Chittka L, Thomson JD, Waser NM (1999) Flower constancy, insect psychology, and plant evolution. Naturwissenschaften 86:361–377. https://doi.org/10.1007/s001140050636

Cnaani J, Thomson JD, Papaj DR (2006) Flower choice and learning in foraging bumblebees: effects of variation in nectar volume and concentration. Ethology 112:278–285. https://doi.org/10.1111/j.1439-0310.2006.01174.x

Combes SA, Gagliardi SF, Switzer CM, Dillon ME (2020) Kinematic flexibility allows bumblebees to increase energetic efficiency when carrying heavy loads. Science Advances 6:eaay3115. https://doi.org/10.1126/sciadv.aay3115

Cresswell JE (1990) How and why do nectar-foraging bumblebees initiate movements between inflorescences of wild bergamot Monarda fistulosa (Lamiaceae)? Oecologia 82:450–460. https://doi.org/10.1007/BF00319785

Darveau C-A, Billardon F, Bélanger K (2014) Intraspecific variation in flight metabolic rate in the

bumblebee *Bombus impatiens*: repeatability and functional determinants in workers and drones. Journal

of Experimental Biology 217:536–544. https://doi.org/10.1242/jeb.091892

DeWitt TJ, Sih A, Wilson DS (1998) Costs and limits of phenotypic plasticity. Trends in Ecology &

Evolution 13:77–81. https://doi.org/10.1016/S0169-5347(97)01274-3

Dukas R, Real LA (1993) Effects of recent experience on foraging decisions by bumble bees. Oecologia

94:244–246. https://doi.org/10.1007/BF00341323

Evans LJ, Raine NE (2014) Changes in learning and foraging behaviour within developing bumble bee

(*Bombus terrestris*) colonies. PLOS ONE 9:e90556. https://doi.org/10.1371/journal.pone.0090556

Foster DJ, Cartar RV (2011) Wing wear affects wing use and choice of floral density in foraging bumble

bees. Behavioral Ecology 22:52–59. https://doi.org/10.1093/beheco/arq160

Frasnelli E, Robert T, Chow PKY, et al (2021) Small and large bumblebees invest differently when

learning about flowers. https://doi.org/10.1016/j.cub.2020.11.062']

Gegear RJ, Laverty TM (2004) Effect of a colour dimorphism on the flower constancy of honey bees and

bumble bees. Can J Zool 82:587–593. https://doi.org/10.1139/z04-029

Gegear RJ, Thomson JD (2004) Does the flower constancy of bumble bees reflect foraging economics?

Ethology 110:793–805. https://doi.org/10.1111/j.1439-0310.2004.01010.x

Goulson D (2000) Why do pollinators visit proportionally fewer flowers in large patches? Oikos 91:485–

492. https://doi.org/10.1034/j.1600-0706.2000.910309.x

Goulson D (2003) Bumblebees: Their behaviour and ecology. Oxford University Press

Goulson D, Peat J, Stout JC, et al (2002) Can alloethism in workers of the bumblebee, *Bombus terrestris*,

be explained in terms of foraging efficiency? Animal Behaviour 64:123–130.

https://doi.org/10.1006/anbe.2002.3041

Goulson D, Stout JC, Langley J, Hughes WOH (2000) Identity and function of scent marks deposited by

foraging bumblebees. J Chem Ecol 26:2897–2911. https://doi.org/10.1023/A:1026406330348

Grüter C, Ratnieks FLW (2011) Flower constancy in insect pollinators: adaptive foraging behaviour or

cognitive limitation? Communicative & Integrative Biology 4:633–636. https://doi.org/10.4161/cib.16972

Grüter C, Segers FHID, Hayes L (2024) Extensive loss of forage diversity in social bees owing to flower

constancy in simulated environments. Proceedings of the Royal Society B: Biological Sciences

291:20241036. https://doi.org/10.1098/rspb.2024.1036

Gumbert A (2000) Color choices by bumble bees (*Bombus terrestris*): innate preferences and generalization after learning. Behav Ecol Sociobiol 48:36–43. https://doi.org/10.1007/s002650000213

Hagbery J, Nieh JC (2012) Individual lifetime pollen and nectar foraging preferences in bumble bees. Naturwissenschaften 99:821–832. https://doi.org/10.1007/s00114-012-0964-7

Hartig F (2020) DHARMa: residual diagnostics for hierarchical (multi-level/mixed) regression models. R package version 03 3:

Heinrich B (1979a) "Majoring" and "minoring" by foraging bumblebees, *Bombus vagans*: an experimental analysis. Ecology 60:245–255. https://doi.org/10.2307/1937652

Heinrich B (1979b) Resource heterogeneity and patterns of movement in foraging bumblebees. Oecologia 40:235–245. https://doi.org/10.1007/BF00345321

Heinrich B (2004) Bumblebee economics. Harvard University Press

Hemingway CT, Pimplikar S, Muth F (2024) Wild bumblebees use both absolute and relative evaluation when foraging. Behavioral Ecology 35:arad112. https://doi.org/10.1093/beheco/arad112

Hendriksma HP, Toth AL, Shafir S (2019) Individual and colony level foraging decisions of bumble bees and honey bees in relation to balancing of nutrient needs. Front Ecol Evol 7: https://doi.org/10.3389/fevo.2019.00177

Heuschen B, Gumbert A, Lunau K (2005) A generalised mimicry system involving angiosperm flower colour, pollen and bumblebees' innate colour preferences. Plant Syst Evol 252:121–137. https://doi.org/10.1007/s00606-004-0249-5

Hofmann G (1977) Iscotables: a handbook of data for biological and physical scientists. Instrumentation Specialties Company

Holland JG, Nakayama S, Porfiri M, et al (2021) Body size and behavioural plasticity interact to influence the performance of free-foraging bumble bee colonies. Insects 12:236. https://doi.org/10.3390/insects12030236

Hunter JD (2007) Matplotlib: a 2D graphics environment. Computing in Science & Engineering 3:90–95. https://doi.org/10.1109/MCSE.2007.55

Kembro JM, Lihoreau M, Garriga J, et al (2019) Bumblebees learn foraging routes through exploitation–exploration cycles. Journal of The Royal Society Interface 16:20190103. https://doi.org/10.1098/rsif.2019.0103

Kevan PG, Baker HG (1983) Insects as flower visitors and pollinators. Annu Rev Entomol 28:407–453. https://doi.org/10.1146/annurev.en.28.010183.002203

Lechantre A, Draux A, Hua H-AB, et al (2021) Essential role of papillae flexibility in nectar capture by bees. Proceedings of the National Academy of Sciences 118:e2025513118. https://doi.org/10.1073/pnas.2025513118

Lenth RV (2020) Emmeans: estimated marginal means, aka least-squares means. R Package Version 1.5.3. CRAN

Lihoreau M, Chittka L, Raine NE (2011) Trade-off between travel distance and prioritization of high-reward sites in traplining bumblebees. Functional Ecology 25:1284–1292. https://doi.org/10.1111/j.1365-2435.2011.01881.x

Lihoreau M, Chittka L, Raine NE (2010) Travel optimization by foraging bumblebees through readjustments of traplines after discovery of new feeding locations. The American Naturalist 176:744–757. https://doi.org/10.1086/657042

Lihoreau M, Raine NE, Reynolds AM, et al (2012) Radar Tracking and Motion-Sensitive Cameras on Flowers Reveal the Development of Pollinator Multi-Destination Routes over Large Spatial Scales. PLOS Biology 10:e1001392. https://doi.org/10.1371/journal.pbio.1001392

Lunau K (1990) Colour saturation triggers innate reactions to flower signals: Flower dummy experiments with bumblebees. J Comp Physiol A 166:827–834. https://doi.org/10.1007/BF00187329

Lunau K, Wacht S, Chittka L (1996) Colour choices of naive bumble bees and their implications for colour perception. J Comp Physiol A 178:477–489. https://doi.org/10.1007/BF00190178

Makino TT, Sakai S (2007) Experience changes pollinator responses to floral display size: from size-based to reward-based foraging. Functional Ecology 21:854–863. https://doi.org/10.1111/j.1365-2435.2007.01293.x

McKinney W (2010) Data structures for statistical computing in python. Austin, Texas, pp 56–61

Menzel R, Erber J (1972) The influence of the quantity of reward on the learning performance in honeybees. Behaviour 41:27–42. https://doi.org/10.1163/156853972X00176

Molet M, Chittka L, Stelzer RJ, et al (2008) Colony nutritional status modulates worker responses to foraging recruitment pheromone in the bumblebee Bombus terrestris. Behav Ecol Sociobiol 62:1919–1926. https://doi.org/10.1007/s00265-008-0623-3

Muller H, Chittka L (2008) Animal personalities: the advantage of diversity. Current Biology 18:R961–R963. https://doi.org/10.1016/j.cub.2008.09.001

Nityananda V, Chittka L (2020) Reward value is more important than physical saliency during bumblebee visual search for multiple rewarding targets. 2020.10.01.322172

Ohara M, Higashi S (1994) Effects of inflorescence size on visits from pollinators and seed set of Corydalis ambigua (Papaveraceae). Oecologia 98:25–30. https://doi.org/10.1007/BF00326086

Ohashi K, Thomson JD, D'Souza D (2007) Trapline foraging by bumble bees: IV. Optimization of route geometry in the absence of competition. Behavioral Ecology 18:1–11. https://doi.org/10.1093/beheco/arl053

Osborne JL, Smith A, Clark SJ, et al (2013) The ontogeny of bumblebee flight trajectories: from naïve explorers to experienced foragers. PLOS ONE 8:e78681. https://doi.org/10.1371/journal.pone.0078681

Osborne JL, Williams IH (2001) Site constancy of bumble bees in an experimentally patchy habitat. Agriculture, Ecosystems & Environment 83:129–141. https://doi.org/10.1016/S0167-8809(00)00262-0

Page RE, Scheiner R, Erber J, Amdam GV (2006) The development and evolution of division of labor and foraging specialization in a social insect (*Apis mellifera* L.). In: Current Topics in Developmental Biology. Academic Press, pp 253–286

Pattrick JG, Symington HA, Federle W, Glover BJ (2023) Bumblebees negotiate a trade-off between nectar quality and floral biomechanics. iScience 26:. https://doi.org/10.1016/j.isci.2023.108071

Pedregosa F, Varoquaux G, Gramfort A, et al (2011) Scikit-learn: machine learning in python. Journal of Machine Learning Research 12:2825–2830

Pyke GH (1984) Optimal foraging theory: a critical review. Annual Review of Ecology and Systematics 15:523–575

Pyke GH, Starr CK (2021) Optimal foraging theory. In: Starr CK (ed) Encyclopedia of Social Insects. Springer International Publishing, Cham, pp 677–685

Raine NE, Chittka L (2008) The correlation of learning speed and natural foraging success in bumble-bees. Proceedings of the Royal Society B: Biological Sciences 275:803–808. https://doi.org/10.1098/rspb.2007.1652

Raine NE, Chittka L (2007) The adaptive significance of sensory bias in a foraging context: floral colour preferences in the bumblebee *Bombus terrestris*. PLOS ONE 2:e556. https://doi.org/10.1371/journal.pone.0000556

Real L, Rathcke BJ (1988) Patterns of individual variability in floral resources. Ecology 69:728–735. https://doi.org/10.2307/1941021

Reynolds AM, Lihoreau M, Chittka L (2013) A simple iterative model accurately captures complex trapline formation by bumblebees across spatial scales and flower arrangements. PLOS Computational Biology 9:e1002938. https://doi.org/10.1371/journal.pcbi.1002938

Riveros AJ, Gronenberg W (2010) Brain allometry and neural plasticity in the bumblebee bombus occidentalis. brain behavior and evolution 75:138–148. https://doi.org/10.1159/000306506

Russell AL, Buchmann SL, Papaj DR (2017) How a generalist bee achieves high efficiency of pollen collection on diverse floral resources. Behavioral Ecology 28:991–1003. https://doi.org/10.1093/beheco/arx058

Saleh N, Chittka L (2007) Traplining in bumblebees (*Bombus impatiens*): a foraging strategy's ontogeny and the importance of spatial reference memory in short-range foraging. Oecologia 151:719–730. https://doi.org/10.1007/s00442-006-0607-9

Saleh N, Scott AG, Bryning GP, Chittka L (2007) Distinguishing signals and cues: bumblebees use general footprints to generate adaptive behaviour at flowers and nest. Arthropod-Plant Interactions 1:119–127. https://doi.org/10.1007/s11829-007-9011-6

Schmid-Hempel P, Kacelnik A, Houston AI (1985) Honeybees maximize efficiency by not filling their crop. Behav Ecol Sociobiol 17:61–66. https://doi.org/10.1007/BF00299430

Sletvold N, Trunschke J, Smit M, et al (2016) Strong pollinator-mediated selection for increased flower brightness and contrast in a deceptive orchid. Evolution 70:716–724. https://doi.org/10.1111/evo.12881

Sommer J, Rao V, Sprayberry J (2022) Deconstructing and contextualizing foraging behavior in bumble bees and other central place foragers. Apidologie 53:32. https://doi.org/10.1007/s13592-022-00944-3

Spaethe J, Tautz J, Chittka L (2001) Visual constraints in foraging bumblebees: Flower size and color affect search time and flight behavior. Proceedings of the National Academy of Sciences 98:3898–3903. https://doi.org/10.1073/pnas.071053098

Spaethe J, Weidenmüller A (2002) Size variation and foraging rate in bumblebees (*Bombus terrestris*). Insectes soc 49:142–146. https://doi.org/10.1007/s00040-002-8293-z

Thakar JD, Kunte K, Chauhan AK, et al (2003) Nectarless flowers: ecological correlates and evolutionary stability. Oecologia 136:565–570. https://doi.org/10.1007/s00442-003-1304-6

Vaudo AD, Patch HM, Mortensen DA, et al (2016) Macronutrient ratios in pollen shape bumble bee (*Bombus impatiens*) foraging strategies and floral preferences. Proceedings of the National Academy of Sciences 113:E4035–E4042. https://doi.org/10.1073/pnas.1606101113

Waddington KD, Gottlieb N (1990) Actual vs perceived profitability: A study of floral choice of honey bees. J Insect Behav 3:429–441. https://doi.org/10.1007/BF01052010

Waddington KD, Holden LR (1979) Optimal foraging: on flower selection by bees. The American Naturalist 114:179–196. https://doi.org/10.1086/283467

Waser NM (1986) Flower constancy: definition, cause, and measurement. The American Naturalist 127:593–603. https://doi.org/10.1086/284507

Waskom M (2021) seaborn: statistical data visualization. JOSS 6:3021. https://doi.org/10.21105/joss.03021

Weiss MR (1991) Floral colour changes as cues for pollinators. Nature 354:227–229. https://doi.org/10.1038/354227a0

Whitney HM, Dyer A, Chittka L, et al (2008) The interaction of temperature and sucrose concentration on foraging preferences in bumblebees. Naturwissenschaften 95:845–850. https://doi.org/10.1007/s00114-008-0393-9

Willmer P (2011) Pollination and floral ecology. Princeton University Press

Wolf ThJ, Schmid-Hempel P, Ellington CP, Stevenson RD (1989) Physiological correlates of foraging efforts in honey-bees: oxygen consumption and nectar load. Functional Ecology 3:417–424. https://doi.org/10.2307/2389615

Woodgate JL, Makinson JC, Lim KS, et al (2016) Life-long radar tracking of bumblebees. PLOS ONE 11:e0160333. https://doi.org/10.1371/journal.pone.0160333