The TUTase URT1 regulates the transcriptome of seeds and their primary dormancy

Jackson Peter[1,3], Jeanne Roignant[1,3], Sebastian Sacharowski[2], Elodie Ubrig[1], Benjamin Lefèvre[1], Szymon Swiezewski[2], Dominique Gagliardi[1], Hélène Zuber[1,*]

[1]Institut de biologie moléculaire des plantes, CNRS, Université de Strasbourg, Strasbourg, France.

[2] Laboratory of Seeds Molecular Biology, Institute of Biochemistry and Biophysics Polish Academy of Sciences, Warsaw, Poland

[3]These authors contributed equally.

[*]Correspondence:

helene.zuber@ibmp-cnrs.unistra.fr

+33367155381

Short title

Uridylation regulates the seed transcriptome

Keywords

Uridylation, URT1, RNA degradation, Seed maturation, Primary dormancy

Abstract

RNA uridylation is a pervasive mechanism that regulates the degradation of eukaryotic mRNAs. In Arabidopsis, uridylation influences mRNA decay both by favoring 5' to 3' degradation and by preventing excessive deadenylation. Yet, the significance of mRNA uridylation during plant development remains largely unknown. Here, we adapted FLEP-seq2, a method based on nanopore sequencing, to generate a comprehensive inventory of mRNA uridylation events in different Arabidopsis tissues. We also evaluated the respective contribution of the two known Arabidopsis uridylyltransferases, URT1 and HESO1, in mRNA uridylation. Our transcriptome-wide analysis showed that, URT1 is the main enzyme responsible for mRNA uridylation, in all analyzed tissues, while HESO1 can marginally uridylate mRNAs. Importantly, our results revealed the singularity of mRNA uridylation pattern in seeds and the dual function of URT1-dependent uridylation in shaping the transcriptome during seed maturation. We propose that during seed maturation, URT1-dependent uridylation promotes the elimination of unnecessary translation-associated mRNAs, by facilitating their degradation, while also promoting the accumulation of mRNAs associated with the maturation program, by hindering their deadenylation. In line with its important function in shaping the seed transcriptome, our study also identifies URT1 as a novel regulator of seed dormancy. Overall, our study reveals the biological relevance of mRNA uridylation during seed maturation.

Introduction

mRNA uridylation is an abundant post-transcriptional modification that is conserved in most eukaryotes, from fission yeast to plants and humans (De Almeida et al. 2018; Warkocki et al. 2018; Zigáčková and Vaňáčová 2018; Yu and Kim 2020). The addition of uridines (U) to the 3' end of mRNAs is catalyzed by Terminal UridylylTransferases (TUTases) such as TUT4/TUT7 in mammals (Lim et al. 2014; Thomas et al. 2015; Morgan et al. 2017; Chang et al. 2018) or Cid1 in fission yeast (Rissland and Norbury 2009; Grochowski et al. 2024). In Arabidopsis, UTP:RNA URIDYLYLTRANSFERASE 1 (URT1) is the main enzyme responsible for mRNA uridylation (Sement et al. 2013; Zuber et al. 2016). Yet, the residual uridylation detected in the *urt1* null mutation indicates that Arabidopsis contains at least a second enzyme able to uridylate mRNAs. HEN1 SUPPRESSOR1 (HESO1), the second prominent TUTase characterized in Arabidopsis,

is involved in the uridylation of small RNAs and 5'-cleavage fragments resulting from cleavage by the RNA-induced silencing complex (RISC) (Ren et al. 2012, 2014, 2023; Zhao et al. 2012; Zuber et al. 2018; Vigh et al. 2024) and shared common RNA substrates with URT1 including some microRNAs (Tu et al. 2015; Wang et al. 2015), some 5'-cleavage fragments from RISC-cleavage (Zuber et al. 2018; Vigh et al. 2024) and the polyadenylated viral RNA of TuMV (Joly et al. 2023). While its role in mRNA uridylation has not been investigated yet, HESO1 is a good candidate as the alternative TUTase for mRNA uridylation. In eukaryotes, the main function of mRNA uridylation is to assist mRNA decay. In fission yeast or human cells, mRNA uridylation was described to facilitate the degradation of mRNAs from 5' to 3', by recruiting the decapping machinery and thus triggering the decay by 5'-3' exoribonuclease 1 (XRN1), or alternatively from 3' to 5' by promoting the activity of Dis3-like protein 2 (Dis3L2) or the RNA exosome (Chowdhury et al. 2007; Song and Kiledjian 2007; Rissland and Norbury 2009; Malecki et al. 2013; Lim et al. 2014). In Arabidopsis, mRNA uridylation also participates in mRNA decay pathways. URT1 is part of a protein network comprising the deadenylation complex CCR4-NOT and other translation repressors/decapping activators. Notably, URT1 directly interacts with DECAPPING 5 (DCP5), a protein that promotes mRNA decapping and participates in translation inhibition. Moreover, recent results based on direct RNA sequencing (DRS) using Oxford Nanopore Technologies (ONT) revealed a global role of URT1 in shaping the poly(A) tail length of mRNAs. URT1-mediated uridylation was shown to prevent the accumulation of excessively deadenylated mRNAs, likely by both favoring their 5' to 3' degradation and hindering excessive deadenylation. Importantly, this function protects plants against the biogenesis of illegitimate siRNAs that silence endogenous mRNAs and strongly impact plant growth and development (Scheer et al. 2021). Altogether, these results reflect the complexity of the molecular mechanisms associated with mRNA uridylation in plants and of its consequences on mRNA fate.

In mammals, mRNA uridylation has been shown to be essential for oocyte maturation and fertility (Morgan et al. 2017). In Arabidopsis, functional mRNA uridylation and degradation by the cytosolic RNA exosome are required to prevent that a subset of mRNAs associated with photosynthesis enters the siRNA pathway, highlighting the physiological significance of RNA uridylation (Wang et al. 2022). Yet, the regulatory roles

of mRNA uridylation during a plant development process remains elusive and so far, knowledge about mRNA uridylation has mainly been collected in flowers and in seedlings. A comprehensive gene-to-gene analysis of mRNA uridylation profiles and of the contribution of URT1 and HESO1 in different plant tissues and stages is still lacking. FLEP-seq2, a Nanopore-based cDNA sequencing method, was recently developed to measure poly(A) tail sizes of mRNAs in a transcriptome-wide manner (Jia et al. 2022). This long-read sequencing method has the significant advantage to circumvent an inherent bias of Illumina sequencing toward poly(A) size estimation and to avoid purifying polyadenylated mRNAs. It is theoretically possible to use FLEP-seq2 to detect and analyze 3' terminal added non-adenosine nucleotides and therefore uridylation. In the present study, we developed an additional bioinformatic module to analyze the composition and the length of mRNA non-adenosine tails and we showed that FLEP-seq2 can be used to accurately characterize U-tails. Using FLEP-seq2, we generated an atlas of mRNA U-tails in different plant tissues, including in developing, dry mature and germinating seeds for wild-type plants and *urt1* and *heso1* single or double mutants. The most striking observation was a strong accumulation of uridylated mRNAs, coupled to a lengthening of U-tails, specifically detected in seeds. Moreover, we showed that URT1 was the main contributor to this uridylation and that its loss has a prominent impact on the transcriptome of dry seeds and on seed primary dormancy. Finally, our findings revealed the importance of mRNA uridylation dual roles during a developmental process: during seed maturation, URT1-mediated promotes the downregulation of mRNAs encoding translation-associated proteins, while it facilitates the accumulation of mRNAs associated with the seed maturation program.

Results

mRNAs of dry mature seeds are highly uridylated

To first assess the accuracy of FLEP-seq2 to analyze uridines added to the RNA 3' end, we sequenced synthetic DNA fragments harboring a 10-nt poly(A) tail followed by 0 to 6 thymidines (Ts). We improved the original version of the FLEP-seq2 analysis pipeline (Jia et al. 2022) by including an additional module to analyze the composition and the length of non-adenosine tails (https://github.com/jackson-peter/FLEPseq2). Our results showed

that 91 to 96% of sequences corresponding to DNA fragments with one or several Ts were correctly tagged as uridylated RNAs while the technical background, *i.e.* percentage of sequences mistakenly tagged as uridylated RNAs, was lower than 1% (Supplementary Figure 1A, Supplementary Data Set S1). The number of Ts was more accurately called for shorter than for longer tails. For the synthetic DNA fragment containing 6 Ts, this number was correctly estimated for 59% of sequences and underestimated at 5 for 28% of sequences. Importantly, the number of Ts was accurately estimated (>90%) for synthetic DNA fragments containing 1 to 3 Ts (Supplementary Figure 1B, Supplementary Data Set S1). This accuracy is important because the vast majority of uridylated mRNAs detected in Arabidopis contain from 1 to 3 Us (Sement et al. 2013; Zuber et al. 2016). Altogether these results demonstrate good performances and accuracy of our pipeline for quantifying levels of uridylation and differentiating short and long U-tails from FLEP-seq2 data.

To perform a comprehensive analysis of mRNA U-tailing in different Arabidopsis plant tissues, we took advantage of FLEP-seq2 datasets from (Jia et al. 2022) and we analyzed uridylation profiles for cytosolic mRNAs. Our results showed consistent overall percentages of uridylation across replicates ranging from 5 to 6% in most tissues (Figure 1A). Interestingly, the highest level of uridylation was found in dry seeds for which uridylation rate reached 9% on average. This represents an increase of 70% when compared to flowers. We then calculated the percentage of uridylation per mRNA and plotted the distribution of percentages for each replicate and tissue (Figure 1B, Supplementary Figure S1C). The intergenic distribution showed higher uridylation rates in seeds regardless of the mRNA population, whether calculated for all mRNAs (Figure 1B), for those detected in all tissues or for those specifically expressed in dry seeds (Supplementary Figure S1C). Therefore, the overall high uridylation rate of mRNAs cannot be explained by the specificities of the seed transcriptome. Interestingly, we also observed a lengthening of U-tails in dry seeds. In all tissues except seeds, U-tails were mainly short, composed of 1 or 2 Us with less than 26% of tails longer than 2 Us (Figure 1C). This result confirms previous results obtained in flowers and in leaves using alternative methods (Sement et al. 2013). By contrast in dry seeds, the proportion of long U-tails substantially

increased with 55% of U-tails longer than 2 Us. These findings highlight both an accumulation and a lengthening of U-tails in dry seeds as compared to other tissues.

Uridylation preferentially targets mRNAs regulated during seed maturation

We then wondered whether some genomic characteristics were associated with a high or low mRNA uridylation rates and in particular whether some of these features could be specific to dry seeds. In dry seeds, highly uridylated mRNAs had both a significant lower number of introns and a shorter CDS length, while weakly uridylated mRNAs had a higher number of intron and a longer CDS length (Figure 1D). Intriguingly, the relationship between CDS length and uridylation is opposite in tissues other than seeds with highly uridylated mRNAs having longer CDS length in all other tissues except in pollen (Figure 1D). We then performed a gene ontology analysis and looked for biological processes significantly enriched among highly uridylated mRNAs. While in all tissues except pollen, gene ontologies associated with DNA transcription were significantly overrepresented among the 10% most uridylated mRNAs (hereafter called TOP10 uridylated mRNAs), the biological processes associated with response to water deprivation, wounding and bacterium were specifically enriched in dry seeds (Supplementary Figure S1D and Supplementary Data S1). Of note, the most enriched gene ontology among TOP10 uridylated mRNAs in seeds was the cellular response to hypoxia (fdr: 1,92E-17, Fold Enrichment: 4.4) which was also significantly enriched but to a lesser extent in pollen (fdr: 0.02, Fold Enrichment: 3.6). Among the most uridylated mRNAs in dry seeds, we found many key genes of the seed maturation program, such as regulators of the abscisic acid pathway (HIGHLY ABA-INDUCED PP2C GENE 1, 2 and 3; ABI five-binding protein 1 and 2), the chlorophyll degradation (STAY-GREEN 1 and 2) or regulators of seed dormancy (DELAY OF GERMINATION 1 (DOG1); SLEEPY1). We expanded this gene ontology analysis to mRNAs, which were not necessarily the most uridylated but which showed the highest increase in uridylation levels in dry seeds when compared to other tissues. To identify these mRNAs, we plotted an heatmap showing for each mRNA the variation patterns of uridylation percentages between tissues (Supplementary Figure S1E). Out of the 1987 mRNAs detected with at least 50 reads in all tissues, the 869 mRNAs belonging to the cluster 1 showed the highest increase in the uridylation level in dry seeds compared to other tissues with an average 2.2-fold increase. Biological processes enriched for

cluster 1 differed from those enriched among TOP10 uridylated mRNAs and were associated with ribosome (FDR: 2.82-22, Fold Enrichment: 1.9) and translation (FDR: 8.32E-23, Fold Enrichment: 1.9, Supplementary Data Set S1). In line with this clustering analysis, the intergenic distribution confirmed that the uridylation rate of most mRNAs associated with translation substantially increase in dry seeds compared to all other tissues (Supplementary Figure 1 F). Interestingly, the seed maturation was previously associated with a gradual decrease in translation at the end of the seed maturation (Bai et al. 2023). Altogether, the FLEP-seq2 data revealed that uridylation preferentially targets major classes of mRNAs that are associated with important and highly regulated processes of the seed maturation.

Distinct processes explain high uridylation rates in developing and dry seeds

In order to confirm our analysis based on results from (Jia et al. 2022) datasets and to obtain further information about the dynamic of uridylation profiles in seeds, we produced FLEP-seq2 libraries from three biological replicates for different seed stages: developing, dry mature and germinating seeds (Figure 2A). For developing seeds, we analyzed a pool of developing stages from embryogenesis to seed filling (~ from 5 to 14 days after flowering, before the onset of desiccation). We analyzed both isolated seeds and siliques containing the seeds. For germinating seeds, we analyzed seeds imbibed in water for 24 hours. For comparison, we produced libraries from flowers. Our analysis first confirmed previous observations: both bulk and intergenic percentages of uridylation were two-fold higher in dry seeds (bulk uridylation of 11% on average) than in flowers (bulk uridylation of 5% on average) (Figure 2B and C, Supplementary Data Set S2). This increased uridylation rate was not restricted to seed specific mRNAs (Supplementary Figure 2A), and U-tails were longer in dry seeds with on average 60% of U-tails containing more than 2 Us compared to 23% in flowers (Figure 2D). Moreover, as previously observed, highly uridylated mRNAs in dry seeds have a lower number of introns and a shorter CDS length (Supplementary Figure 2B). Interestingly, our new analysis also highlighted changes of mRNA uridylation rates and profiles during seed development and germination. In developing seeds or siliques, the mRNA uridylation rate reached a similar, albeit a little lower, level (9.4-9.9%) as compared with the uridylation rate observed in dry seeds (Figure 2B and C). By contrast in germinating seeds, this rate decreased to 5% on

average, similar to what was observed in flowers. This observation implies a highly dynamic regulation of mRNA uridylation rates during the first stages of seed germination. The length of U-tails was also modulated during development and germination with U-tails being longer in dry seeds compared to developing seeds (25% of U-tails longer than 2 Us in developing seeds) and shortened again in germinating seeds (only 16% of U-tails longer than 2 Us in developing seeds). Finally, the relationship between the CDS length and the uridylation level of mRNAs differs between dry seeds and developing or germinating seeds: CDS of highly uridylated mRNAs were significantly shorter in dry seeds but longer in both developing and germinating seeds (Supplementary Figure 2B). These results imply that the preferential targets of mRNA uridylation vary during seed development or germination. Interestingly, the proportion of mRNAs with a short CDS and a low number of introns is higher among mRNAs that are specifically expressed at the later stages of seed maturation (Supplementary Figure 2C, based on (Le et al. 2010)). These results indicate that mRNAs that accumulate during the late stages of seed maturation are highly uridylated.

We have previously shown in Arabidopsis that uridylation targets mostly oligoadenylated mRNAs with (A)-tail shorter than 25As (Sement et al. 2013; Zuber et al. 2016). Therefore, the high uridylation rates observed in developing seeds and in dry seeds could reflect changes in poly(A) tail populations and could result from an increased proportion of A-tail shorter than 25As. To test this possibility, we analyzed poly(A) size distributions for both non-uridylated and uridylated tails from 10 to 200 As (Figure 3). We also compared the proportion of mRNAs with no or very short tail (tail shorter than 10 As, hereafter called non-polyadenylated mRNAs) to the proportion of mRNAs with tails between 10 and 25 As (called hereafter oligoadenylated mRNAs) or longer than 25 As (Figure 3). In their study, Jia et al. (Jia et al. 2022) reported tissue-specific regulation of poly(A) tail length with patterns in dry seeds and pollen that differ from other tissues. In line with their results, our analysis confirmed the singularity of the poly(A) profile in dry seeds: the two main peaks of the poly(A) distribution in dry seeds were shifted toward longer tails, from ~22 and ~41-46 As in flowers or germinating seeds, to ~29 and ~58 As in dry seeds (Figure 3A). Yet, the most striking difference was observed in developing seeds and siliques for which there was a massive shift of poly(A) tail size profiles toward

shorter tails compared to other stages (Figure 3A). Our results revealed a strong accumulation of mRNAs with A-tails shorter than 25 As, a small size shift of the first peak to ~20 nt and a disappearance of the second one. This poly(A) tail size variation was not specific to a subset of mRNAs but instead affected most mRNAs (Figure 4). Altogether, our results reflected important modulations of poly(A) tail sizes during seed development and germination: for the majority of mRNAs, the proportion of oligoadenylated mRNAs was very high in developing seeds, dropped during seed maturation and then increased again at an intermediary level during seed germination (Figure 3 and 4). Actually, the increased proportion of oligoadenylated mRNAs in developing seeds also explained the increased uridylation rate observed at this stage. Indeed, uridylation rates were similar between developing seeds and flowers, when only these oligoadenylated poly(A) tails were taken into account for the calculation of percentages (Figure 3B). Importantly, uridylation rates were higher in dry seeds when compared to other tissues regardless of the poly(A) tail population (Figure 3B) and both short and long uridylated poly(A) tails accumulated (Figure 3A). Altogether, our results revealed two molecular scenarios leading to the accumulation of U-tails in developing and dry seeds. In developing seeds, the higher uridylation rates of mRNAs is likely governed by a stage-specific regulation of poly(A) size distributions. In dry seeds, both the increased uridylation rate and the lengthening of U-tails could reflect changes in the degradation processes during maturation such as increased activity of TUTases and/or a slowdown in the degradation of uridylated mRNAs.

Respective contributions of URT1 and HESO1 to mRNA uridylation

While we have previously shown that URT1 is the main TUTase responsible for mRNA uridylation in Arabidopsis leaves and flowers (Sement et al. 2013; Zuber et al. 2016), the contribution to mRNA uridylation of HESO1, the second TUTase characterized in Arabidopsis, has not been studied so far. To apprehend the respective contributions of each TUTase to mRNA uridylation, we compared FLEP-seq2 data from three biological replicates for WT plants (same libraries as described above), *urt1-1*, *heso1-4* and *heso1-4 urt1-1* mutants and for four different tissues: developing siliques, dry mature seeds, germinating seeds and flowers. In total, 48 libraries were analyzed. For each tissue and genotype, we calculated the global uridylation percentage and the uridylation percentages per mRNA if detected with at least 50 reads in all three replicates and four genotypes

(Figure 5A, Supplementary Data Set S2). Our results confirmed the preponderant role of URT1 in mRNA uridylation: for all tissues, both global and intergenic percentages of uridylation strongly decreased in *urt1-1* when compared to WT.

HESO1 contributed only modestly to mRNA uridylation because percentages of uridylation were similar in WT and *heso1-4*, and only further decreased in *heso1-4 urt1-1* when compared to *urt1* by an additional 3.8-8.4%. To identify potential mRNAs preferentially uridylated by HESO1, we performed a statistical analysis and compared mRNAs differentially uridylated between WT and *urt1-1* and *heso1-4* mutants and between *urt1-1* and *heso1-4 urt1-1* mutants (Figure 5B and C, Supplementary Data Set S2). As expected, the highest number of mRNAs with a significant decrease in uridylation rate was found in *urt1-1* and *heso1-4 urt1-1* (Figure 5B) and 93 to 99% of mRNAs with a significant decrease in *urt1-1* were also differentially uridylated in *heso1-4 urt1-1* (Figure 5C). We identified only a few mRNAs showing a significant decrease in *heso1-4* when compared to WT, from 2 in dry seeds to 33 in flowers, or in *heso1-4 urt1-1* when compared to *urt1-1*, from 22 in imbibed seeds to 57 in developing siliques. In addition, we detected a few mRNAs with an increased uridylation rate in *heso1-4* when compared to WT. For example, in dry seeds, 47 mRNAs showed a significant increase in uridylation (Figure 5B). Yet, there was almost no overlap of these gene lists across the different tissues (Supplementary Figure 3A) and no gene ontology or genomic feature was enriched in mRNAs differentially uridylated in *heso1-4* (Supplementary Figure 3B). While URT1 and HESO1 were previously shown to sequentially uridylate some microRNAs (Tu et al. 2015), our analysis did not show any changes in the length of mRNA U-tails in *heso1* mutant and did not allow demonstrating a similar sequential uridylation for mRNAs (Supplementary Figure 3C). In *heso1-4 urt1-1*, the uridylation percentage median ranged from 0.95 in flowers to 1.7% in dry seeds when considering all reads. When considering only polyadenylated mRNAs (≥ 10 As), for a better comparison with the synthetic DNA fragments previously analyzed, the uridylation percentage median ranged from 0.7% in flowers to 1.1% in imbibed seeds, (Supplementary Data Set S2). These uridylation levels were in the same range as the uridylation background measured for the non-uridylated synthetic DNA fragment (0.6%) (Supplementary Data Set S1). These results suggest that most of the residual uridylation observed in the *heso1-4 urt1-1* likely corresponds to

technical background. Altogether, our results showed that URT1 is the main enzyme responsible for the bulk mRNA uridylation regardless of the tissue. HESO1 can participate in mRNA uridylation but that its overall contribution remains modest and limited to a small subset of mRNAs. The existence of a third unidentified TUTase uridylating mRNAs remains elusive at this stage.

URT1 has a major impact on dry seed transcriptome

The accumulation and the lengthening of mRNA U-tails in dry seeds suggests that uridylation could play an important role in the regulation of gene expression during seed maturation. To evaluate the contribution of both TUTases on the mRNA steady state level in seeds, we produced total RNA-seq libraries from dry seeds of WT, *urt1-1*, *heso1-4* and *heso1-4 urt1-1* plants. For comparison, we produced in parallel RNA-seq from flowers. We performed a differential gene expression analysis to identify mRNAs that are differentially accumulated in mutant seeds and flowers when compared to WT (Figure 6, Supplementary Data Set S3 and S4). Interestingly, the stronger impact of URT1 inactivation on the transcriptome was observed in dry seeds, with 298 and 326 mRNAs differentially accumulated in *urt1-1* and *heso1-4 urt1-1*, respectively. In flowers, only 32 and 46 mRNAs were differentially accumulated in *urt1-1* and *heso1-4 urt1-1*. These results are in line with the strong uridylation rate observed in seeds and highlight the contribution of URT1-dependent uridylation in shaping the transcriptome during seed maturation. Unlike URT1, HESO1 had a stronger impact in flowers with 118 mRNAs differentially accumulated against only 7 in dry seeds. This weaker impact of HESO1 on dry seed transcriptome can be possibly explained by the different expression profiles of the two TUTases during seed development. Indeed, whereas *URT1* is strongly expressed in seeds throughout their development with an increased mRNA accumulation at late maturation stages, the expression of *HESO1* strongly decreased at the end of seed development (Supplementary Figure 4).

To search for potential cellular or metabolic processes associated with mRNA uridylation in seeds, we performed gene ontology analysis for the different lists of differentially accumulated mRNAs. Our analysis revealed a strong overrepresentation of gene ontologies associated with ribosome and translation among mRNAs upregulated in both *urt1* and *heso1 urt1* dry seeds (Supplementary Data Set S3). Using publicly available

RNA-seq data from Col-0 plants (Klepikova et al. 2016), we then analyzed the expression profile during WT seed development of mRNAs found to be down or up -regulated in *urt1-1* and *heso1-4 urt1-1* seeds (Figure 6B, Supplementary Data Set S5). Interestingly, we observed an opposite trend of expression between mRNAs found to be up and down regulated by URT1 inactivation: while the majority of down-regulated mRNAs strongly accumulate at late developmental stages (mRNAs from clusters 5 and 7; 101/141), the majority of mRNAs up-regulated in *urt1-1* are weakly expressed at the same stages in WT (mRNAs from clusters 1,3-6; 194/300). Of note, most of the up-regulated mRNAs associated with cytosolic ribosomes (62/65) are weakly expressed at the end of seed maturation in a WT context. Thus, URT1-mediated uridylation participates in the reduction of the accumulation of these mRNAs during late seed maturation.

URT1 inactivation hinders the deadenylation of mRNAs that accumulate during seed maturation

We have previously shown using nanopore Direct RNA sequencing that URT1 shapes poly(A) tail size distributions with a clear shift in the distribution toward short oligo(A) tails in *urt1-1* flowers. Our data showed that URT1-dependent uridylation of mRNAs prevents the accumulation of excessively deadenylated mRNAs, likely by both favoring their 5' to 3' degradation and hindering deadenylation (Scheer et al. 2021). Our results here confirmed the previous findings: in flowers of *urt1-1*, we observed a clear shift in the global poly(A) tail size distribution compared to WT with an increased accumulation of oligoadenylated tails with sizes centered around 22 As (Figure 7A). While this accumulation of oligoadenylated tails was also clearly visible in imbibed seeds and developing seeds, no change was observed in dry seeds. Yet in all tissues, we observed an increased accumulation of mRNAs with tails shorter than 10 As in *urt1-1* mutant. In flowers, the proportion of non-polyadenylated tails was moderately increased from 10% in WT to 13% in *urt1-1*. In dry seeds, the increase was more important when compared to other tissues with a proportion going from 12% in WT to 20% in *urt1-1*. These results again reflect the specificity of uridylation-associated processes in maturing seeds, in which the *urt1-1* mutation did not result in a global accumulation of the oligoadenylated mRNAs but rather triggered an accumulation of excessively deadenylated mRNAs. These results also imply that the function of URT1-mediated uridylation in preventing the accumulation of

excessively deadenylated mRNAs could be particularly critical in seeds. Of note, the *heso1-4* mutation did not have a robust impact on the poly(A) tail profiles, regardless of the tissue, and the *heso1-4 urt1-1* double mutation did not worsen the *urt1-1* effect. These results show that HESO1-mediated uridylation of mRNAs is not able to compensate for the absence of URT1 to modulate poly(A) tail size and that both TUTases have distinct functions as previously reported (Zuber et al. 2016).

To identify the most impacted mRNAs in terms of poly(A) shortening in dry seeds, we computed, for each mRNA detected with at least 50 reads, a poly(A) shortening score that corresponds to the difference of the percentage of non-polyadenylated tails between *urt1-1* and WT dry seeds (Supplementary Data Set S2). These poly(A) shortening scores showed a good correlation with mRNA uridylation: the most uridylated mRNAs also showed the highest scores (Supplementary Figure 5A). Using this poly(A) shortening score, we identified the 93 most impact mRNAs with an average shortening score exceeding 20%. For these mRNAs, the first peak of the poly(A) distribution profile (centered around 20 As in WT) disappeared in *urt1-1* seeds and the proportion of non-polyadenylated tails went on average from 20 to 47% (Figure 7B and C). Interestingly, these 93 mRNAs were significantly enriched in mRNAs encoding proteins related to cellular response to hypoxia and abscisic acid signaling pathway (Supplementary Data Set S2) and most of them (89/93) were genes strongly up-regulated at the end of seed maturation (Supplementary Figure 5B). Moreover, among these 93 most impacted mRNAs, eight also showed a significant decrease in total mRNA accumulation in *urt1-1*. This overlap is relatively weak but it shows that for some mRNAs, the poly(A) shortening is coupled with a decrease steady state level of total mRNA. To investigate more carefully a link between the poly(A) shortening and the mRNA steady-state level, we compared intergenic poly(A) tail profiles for mRNAs down or up-regulated in *urt1-1* or *heso1-4 urt1-1* seeds (Figure 7D and E). Our results showed that mRNAs, which show a significant decrease in steady-state mRNA level, were also the most impacted in terms of poly(A) tail size compared to up-regulated or other mRNAs: for mRNAs down-regulated in *urt1-1* dry seeds, the 20 As centered peak of the distribution profile was absent in *urt1-1* dry seeds (Figure 7D) and the median proportion of non-polyadenylated tails went from 16-20% in WT to 32-39% in *urt1-1* dry seeds (Figure 7E). The down-regulated mRNAs were also

highly uridylated compared to up-regulated or other mRNAs (Supplementary Figure 5C). These results show that the poly(A) shortening caused by URT1 defect can trigger a decrease in the mRNA steady-state level at least for some mRNAs.

Altogether our results reveal the diverse consequences of mRNA uridylation during seed maturation: while URT1-mediated uridylation may promote the deregulation of mRNAs encoding translation-associated proteins at the end of seed maturation, it may hinder the deadenylation of mRNAs associated with the seed maturation program, thereby facilitating their accumulation.

URT1 inactivation alters primary dormancy levels of seeds

The important contribution of URT1-dependent uridylation in shaping the seed transcriptome prompted us to investigate a possible impact of URT1 inactivation on seed germination. We first monitored the germination kinetic of after-ripened seeds, in which primary dormancy is alleviated. We also applied different duration of accelerated aging treatment to investigate a potential impact of URT1 inactivation on seed longevity. Our results did not reveal any impact of URT1 inactivation on the germination kinetic of after-ripened seeds and the final germination rates or the germination speed remained unchanged in *urt1-1* mutant even after accelerated aging treatments (Supplementary Figure 6). We then conducted germination assays on freshly harvested seeds, which usually show high levels of dormancy, and we monitored the percentages of germination after 7 days. These assays revealed a significant decrease in primary dormancy in *urt1-1* mutant when compared to WT (Figure 8A). This weaker dormancy of *urt1-1* freshly harvested seeds was repeatedly observed for seeds harvested from individual plants and from individual batches of plants (at least four plants for four individual batches, Figure 8A). Interestingly, *DOG1* a regulator of seed dormancy, whose protein level was shown to highly correlate with dormancy (Nakabayashi et al. 2012), was among the most uridylated mRNAs and also among the most impacted mRNAs in terms of poly(A) shortening in *urt1-1* seeds (Figure 8B). While the level of total *DOG1* mRNA was not significantly altered in *urt1-1* seeds, the proportion of non-polyadenylated mRNAs significantly increased from 24% in WT to 47% in *urt1-1* seeds (Figure 8C and D). The weaker dormancy levels in *urt1-1* seeds could be a consequence of a reduction of the pool of *DOG1* polyadenylated mRNAs in *urt1-1* seeds. The weaker dormancy could also be explained by the increased

accumulation in *urt1-1* seeds of translation-associated mRNAs, as an up-regulation of genes encoding protein of the translational machinery was previously linked to a loss of dormancy (Cadman et al. 2006; Nelson and Steber 2017; Buijs et al. 2020). In any case, our results highlighted the importance of URT1-mediating uridylation in the regulation of primary dormancy.

Discussion

Developmental transitions that occur in the seed imply extensive changes of gene expression, at both transcript and protein levels, which have been documented thoroughly in Arabidopsis during seed development (Le et al. 2010; Belmonte et al. 2013; Bai et al. 2023) or germination (Nakabayashi et al. 2005; Narsai et al. 2011; Basbouss-Serhal et al. 2015; Bai et al. 2017). While transcriptional regulation has been well studied, the post-transcriptional mechanisms involved during seed development and germination remain less understood, especially those related to RNA degradation. mRNA uridylation is a widespread post-transcriptional modification that influences mRNA decay. In the present work, we optimized the data analysis of the nanopore sequencing-based FLEP-seq2 method developed by (Jia et al. 2022) to generate a first inventory of mRNA uridylation events in different Arabidopsis tissues, including developing and germinating seeds. Our results showed overall high levels of mRNA uridylation in developing and in dry seeds, higher than in all other analyzed tissues. In developing seeds, the high uridylation rates of mRNAs were associated with a strong regulation of poly(A) tail size and likely resulted from the massive accumulation of oligoadenylated mRNAs that are the preferential targets of mRNA uridylation (Sement et al. 2013; Zuber et al. 2016). By contrast in dry seeds, high uridylation rates were independent of a poly(A) tail size regulation and reflected either an increased activity of URT1 and/or a decrease degradation of uridylated mRNA during maturation. *URT1* mRNA strongly accumulates at the end of seed maturation suggesting that the uridylation activity could be particularly high at this stage. Yet, the lengthening of U-tails observed specifically in dry seeds could also reflect a lower degradation activity. Indeed, a similar lengthening was previously shown when 5' to 3' or 3' to 5' degradation was impaired (Sement et al. 2013; Lim et al. 2014), but not when URT1 was overexpressed (Scheer et al. 2021). The maturation of orthodox seeds is characterized by a gradual decrease in water content coupled with a strong reduction of the metabolic

activity. Therefore, the accumulation of long U-tails in dry seeds could reflect the gradual shutdown during late seed maturation of the enzymatic degradation pathways, which would be spatiotemporally uncoupled from uridylation processes for reasons yet to be investigated. While the uridylation level of mRNAs was maximal in dry seeds, it was reduced during germination after 24 hours of imbibition in water. The size of U-tails was also shorter in imbibed seeds. This quantitative and qualitative change in uridylation profiles of mRNAs likely reflects the resumption of degradation processes during early germination stages.

These high mRNA uridylation levels in the seed were coupled with a greater impact of URT1 inactivation on the transcriptome of dry seeds as compared to flowers (the present study) or leaves (Sement et al. 2013). These results revealed the importance of URT1-dependent uridylation in the regulation of gene expression during seed maturation, which therefore constitutes a relevant model to dissect the multiple consequences of mRNA uridylation on RNA metabolism. The housekeeping function of mRNA uridylation in Eukaryotes is to facilitate mRNA degradation. In plants, we have previously shown that mRNA uridylation prevents excessive deadenylation thereby favoring the polarity of degradation from 5' to 3'. This dual function of mRNA uridylation is not limited to plants as a recent study shows that a defect in uridylation also leads to an excessive deadenylation in fission yeast (Grochowski et al. 2024). Our results here highlighted the respective contribution of the dual functions of URT1-dependent uridylation in regulating distinct subsets of the transcriptome during seed maturation. On one side, we showed that URT1-mediated uridylation promotes the deregulation of translation-associated mRNAs during seed maturation. Interestingly, translation-associated mRNAs also exhibited a significantly higher level of uridylation in seeds compared to other tissues showing that these mRNAs are important targets of uridylation at this developmental stage. Translation is developmentally regulated in maturing seeds: the overall abundance of ribosomes increases during the initial phase of seed maturation, but then decreases during seed desiccation, indicating a general decline in translation (Bai et al. 2023). Therefore URT1-mediated uridylation could participate in the down-regulation of the translation by facilitating the elimination of translation-associated mRNAs that become unnecessary when the seed enters a quiescent state. On the other side, we showed that URT1-

mediated uridylation facilitates the accumulation of a hundred of mRNAs associated with the seed maturation program, likely by protecting them from an excessive deadenylation. In the absence of URT1, the decrease accumulation of these mRNAs was coupled with an important shortening of their poly(A) tail size: the population of mRNAs with tails centered around 20 As decreased in favor of a significant accumulation of mRNAs with very short tails. This distribution peak centered around 20 As may reflect the fingerprint of one Poly(A) Binding protein (PABP). Indeed, even if PABPs was shown to occupy ~ 27 As (Baer and Kornberg 1983), the minimal A-tail size for high affinity binding *in vitro* is 12 As (Sachs et al. 1987). Moreover, Arabidopsis PABPs were shown to bind mRNAs with tails as short than 16 As, as well as to uridylated mRNAs (Zuber et al. 2016). In fact, our previous findings revealed that URT1-mediated uridylation can repair deadenylated ends to restore a binding site for PABP (Zuber et al. 2016), supporting the idea that in seeds URT1-mediated uridylation could facilitate the binding of the PABP to mRNAs associated with the maturation program. Dry seeds are characterized by the accumulation of many mRNAs at the end of maturation, called stored mRNAs or long-lived mRNAs (Sajeev et al. 2019; Sano et al. 2020). Some of these stored mRNAs were shown to serve as templates for de novo translation during germination (Rajjou et al. 2004; He et al. 2011; Sano et al. 2020), while the others may be reminiscent of the maturation program, thereby reflecting the transcriptomic regulations taking place on the mother plant. While the mechanisms underlying the storage of these mRNAs in dry seeds remain largely unknown, processing bodies or stress granules, biomolecular condensates balancing mRNA degradation, storage and translation (for reviews in plants Kearly et al., 2024; Solis-Miranda et al., 2023), were proposed to serve as a storage location and to act as a protective armor (Sajeev et al. 2019). Interestingly, both URT1 and PABP are components of plant stress granules (Sement et al. 2013; Solis-Miranda et al. 2023). We can therefore speculate that URT1 and PABP act synergically during late stages of seed maturation to facilitate the storage of mRNA in the seed stress granules.

URT1-mediated uridylation shapes the transcriptome during seed maturation. But what are the physiological consequences of this URT1 function? The present work identified URT1 as a novel positive regulator of the primary dormancy. The dormancy is an important seed agronomical trait that is defined by the state in which seeds are unable

to germinate despite having favorable environmental conditions. While secondary dormancy is induced when imbibed non-dormant seeds are exposed to adverse environmental conditions, primary dormancy is established during seed development (for reviews about dormancy see Chahtane et al., 2017; Née et al., 2017; Sajeev et al., 2024). A previous study already highlighted the importance of RNA degradation pathways in the regulation of primary dormancy. In their work, the authors showed primary dormancy phenotypes for two major components of the 5' to 3' decay pathways, the decapping enhancer VARICOSE (VCS) and the main 5' to 3' exoribonuclease EXORIBONUCLEASE4 (XRN4), with *vcs* and *xrn4* mutant exhibiting reduced and higher dormancy level, respectively (Basbouss-Serhal et al. 2017). Explaining these opposite phenotypes, the authors identified subsets of mRNAs specifically targeted by VCS or XRN4 that positively regulate the germination or on the contrary are involved in the maintenance of dormancy, respectively (Basbouss-Serhal et al. 2017). The varying influence of these two 5'-3' decay factors show the complexity of the interplay between RNA degradation and the regulation of the primary dormancy. The regulatory role of URT1 could occur during seed maturation, when primary dormancy is established, or during seed imbibition, when dormant seeds arrest the germination program. The high uridylation rate of mRNAs in dry seeds and its drop as soon as 24 hours after imbibition support the first hypothesis. Moreover, the transcriptomic changes in *urt1-1* dry seeds could also explained the reduced depth of primary dormancy triggered by URT1 inactivation. Indeed, our transcriptomic analysis highlighted a reduced accumulation of polyadenylated mRNAs for DOG1 and for genes related to the ABA signaling pathway and an increased expression of translation-associated mRNAs in *urt1-1* seeds. Altogether, our results suggest that of URT1-mediated uridylation could positively regulate the establishment of the primary dormancy during seed maturation. The depth of dormancy is controlled by the environmental conditions experienced by the mother plant, for example by the temperature or by the nitrate availability (Klupczyńska and Pawłowski 2021). A better understanding of how plants perceive and signal these environmental factors to modulate the depth of seed dormancy is a major challenge from an agronomic perspective. In the future, it would be particularly interesting to explore the contribution of mRNA uridylation

and RNA decay pathways in shaping the seed transcriptome in response to these environmental signals and to investigate its influence on the regulation of dormancy.

Methods

Plant material and growth condition

The *Arabidopsis thaliana* plants used in this work are of Columbia accession (Col-0). Arabidopsis mutants analyzed in this study are T-DNA insertion lines described previously: *urt1-1* (SALK_087647C) (Sement et al. 2013) and *heso1-4* (GK-369H06-017072) (Joly et al. 2023; Vigh et al. 2024). Arabidopsis plants were grown on soil in long-day conditions with a 16 h light/8 h darkness photoperiod in a greenhouse at 21/18 °C. The flowers (unopened flower buds), developing siliques (pool of siliques from ~5 to 14 DAF before visible yellowing is apparent) and dry mature seeds for RNAseq and FLEP-seq2 were harvested from a minimum of five plants each for three biological replicates. Samples were harvested two weeks after the onset of flowering for flowers and the first series of developing siliques, three weeks after the onset of flowering for the second series of siliques, and finally eight weeks after the onset of flowering for dry mature seeds, when all siliques of the plants were dry. For the first series of developing siliques, developing seeds were also isolated from their encapsulating siliques using liquid nitrogen and dry ice as described in (Bates et al. 2013). 24-hour imbibed seeds were obtained after imbibition of the dry mature seeds for 24 hours in milliQ water (16 h-day at 21°C / 8 h-night at 18°C cycles).

Germination test and accelerating aging treatment for after-ripened seeds

After-ripened seeds were harvested 7 to 8 weeks after the onset of flowering. Accelerating aging treatment of seeds were then performed according to (Rajjou et al. 2008). All seeds were equilibrated for three days at 85% relative humidity (20°C) in a constant climate chamber. Day-0 controls were immediately dried back for three days at 32% relative humidity (20°C) and stored at 4°C. Aging treatment was then done by storing the seeds for two, four or seven days at 85% relative humidity (40°C). After each treatment, seeds were dried back for three days at 32% relative humidity (20°C) and stored at 4°C. The germination tests were then performed by placing plates under constant light (20°C) and

germination was scored every day for four days. Source data for germination assays are shown in Supplementary Data Set S6.

Primary dormancy assays

Freshly harvested seeds used for primary dormancy assays were obtained from plants grown in a greenhouse under a long-day photoperiod at 22°C. Importantly, the dry, mature seeds were collected when flower buds were still on the main stem. Dormancy assays were repeated four times for four independent batches of plants. Seeds were collected from four to six individual plants for each independent experiment and each genotype. Approximately 100 to 200 seeds for each plant were sown on blue germination paper (Hoffman Manufacturing Inc.) supported by a layer of thick fabric saturated with water. Plates were placed in a greenhouse under a long-day photoperiod at 22°C and germination was scored after seven days. Source data for germination assays are shown in Supplementary Data Set S6.

Oligonucleotides.

Oligonucleotides used in this study are listed in Supplementary Data Set S7.

RNA isolation

Total RNA from seeds (dry, imbibed and developing seeds) and siliques was extracted according to the protocol (Meng and Feldman 2010) except for the last precipitation performed at -80 °C for 1 hour. RNA from flowers was extracted with TRI Reagent (EUROMEDEX, TR118-100) following the manufacturer's protocol. Total RNA was then purified using the RNeasy MinElute Cleanup kit (Qiagen, 74204).

Library preparation and nanopore sequencing

FLEP-seq2 libraries were prepared from 3 µg of total RNA as described in (Long et al. 2021; Jia et al. 2022). Briefly, 3 µg of RNA were ribodepleted using the Arabidopsis thaliana riboPOOL kit (siTOOLSs BIOTECH, dp-K024-000008) and then ligated to 50 pmol of a 5'-riboadenylated DNA oligonucleotide (Universal miRNA cloning linker, NEB, Supplementary Data Set 7) in a 20 µl reaction containing 20 U of T4 RNA Ligase 2 truncated KQ (NEB, M0373S), 1X T4 RNA Ligase Reaction Buffer (NEB, 50 mM Tris-HCl pH 7.5, 10 mM MgCl2, 1 mM DTT), 25% PEG8000 and 40 U RNAseOUT (Life Technologies, 10777019). Ligation reactions were incubated for 10 hours at 16°C. The product was cleaned up using RNA Clean & Concentrator-5 kit (ZYMO, R1013). The cDNA

synthesis was then performed in two steps. First, the ligated RNAs were incubated with 2 pmol of a custom reverse transcriptase primer (FLEP_seq2_RT, Supplementary Data Set 7) and 10 nmol of dNTPs, incubated at 65°C for 5 min, and placed on ice for 2 min. Then, 1X Maxima RT Buffer (Thermo Fisher, EP0752), 40 U RNaseOUT, 20 pmol of Strand-Switching Primer (SSPI or SSPII, ONT, SQK-PCB111.24) and 200 U Maxima H Minus Reverse Transcriptase (Thermo Fisher, EP0752) were added to the reaction in a 20 µl final volume. Samples were incubated in a thermal cycler at 42°C for 90 min 4°C; 10 cycles at 50°C for 2 min and 42°C for 2 min; and a final step at 85°C for 5 min. cDNA libraries were amplified for 14 cycles in a 25 µl reaction containing 1X PrimeSTAR GXL Buffer (TaKaRa, R050A), 5 nmol of each dNTP, 7.5 pmol of each barcoded primer of the PCR-cDNA Barcoding kit (ONT, SQK-PCB111.24) and 0.625 U of PrimeSTAR GXL DNA polymerase (TaKaRa, R050A). The PCR conditions were as follows: one step at 98°C for 30 s; 14 cycles at 98°C for 10 s, 65°C for 15 s and 68°C for 10 min; a final step at 68°C for 5 min. The PCR products were then incubated with 20 U of Exonuclease I (NEB, M0293) at 37°C for 15 min, followed by a step at 80°C for 15 min. For each sample, 4-6 PCR reactions were carried out and subsequently pooled. The PCR products were cleaned up with three successive rounds of purifications with NucleoMag NGS Clean-up and Size Select kit (Macherey-Nagel, 744970.50) using bead ratios of 0.8X, 0.55X and 0.55X, respectively. Libraries were quantified using the Quant-iT HS DNA assay (Invitrogen, Q33120) and QuBit fluorometer, and the size and the quality were checked using the Agilent High Sensitivity DNA Kit and the Agilent 2100 Bioanalyzer System. Finally, after adding nanopore adapter (Rapid Adapter T, SQK-PCB109/SQK-PCB111), libraries were sequenced for ~72 hours on a MinION or a P2 PromethION 2 Solo device using a FLO-MIN106 (R9.4.1) or a FLO-PRO002 (R9.4.1) flowcell, respectively.

Computational analysis for FLEP-seq2

Raw signal FAST5 files were basecalled using Guppy (Version 6.3.8+d9e0f64) with default parameters using the high accuracy model (-c dna_r9.4.1_450bps_hac). The reads with a mean quality score inferior to 7 were filtered out. The remaining reads were mapped to the *Arabidopsis thaliana* TAIR10 genome using Minimap2 (2.26-r1175) with the Long-read spliced alignment preset enabled (-ax splice) without keeping secondary alignments (--secondary=no). The detection of the 3' ends of the transcripts and the

determination of their length have been performed using a pipeline adapted from (Jia et al. 2022) and available on Github (https://github.com/jackson-peter/FLEPseq2). In particular, the pipeline includes an additional step to analyze the added 3' end U-tail. This additional script, coded in Python 3, extracts and analyzes the composition and the length of the sequence between the end of the detected poly(A) tail and the beginning of the 3' adapter. This sequence was referred to as an additional tail. Additional tails containing at least 70% of Ts were considered as uridylated. All calculations and graphs of the present study were performed for annotated cytoplasmic mRNAs, based on TAIR10. Falsely or ambiguously annotated RNAs were filtered out (see a list of filtered out gene identifiers in Supplementary Data Set S8). Bulk uridylation percentages were calculated using the total number of sequences as denominator. For intergenic uridylation percentages, uridylation percentages were calculated for all mRNAs detected with at least 50 reads in each replicate and boxplots show the distribution of these percentages across mRNAs for each sample and condition. To plot the distribution of poly(A) tail sizes in Figure 3A and in Figure 7A, percentages of reads according to poly(A) tail size (1-nt bin) were calculated using the total number of sequences as denominator. To plot the intergenic distribution of poly(A) tail sizes in Figure 7B and 7C, percentages of reads according to poly(A) tail size (2-nt bin) were calculated for each mRNA detected with at least 50 reads in each replicate and poly(A) profiles show the median (line), the first and the third quartile (grey shadow) of the percentages across mRNAs. For heatmaps, percentages of uridylation and percentages of reads according to poly(A) tail sizes (5nt-bin) in Supplementary Figure S1 and Figure 4, respectively, were calculated for each mRNAs detected with at least 50 reads in each replicate and tissues and were z-score rescaled.

Source data for figures are available in Supplementary Data Set S2. Statistical analysis results are available in Supplementary Data Set S9. FLEP-seq2 results are also available in RNAvis, a web application developed in-house using Rshiny, accessible at https://nebula.ibmp.unistra.fr/RNAvis/.

Analysis of synthetic DNA fragments harboring 0 to 6 Ts.

Synthetic DNA fragments harboring a 10-nt poly(A) tail followed by 0, 1, 2, 3 or 6 Ts were synthesized as gBlocks Gene Fragments (Integrated DNA Technologies, Supplementary Data Set S7). gBlocks Gene Fragments were cloned in pGEM-T Easy (Promega) using

the TA cloning method according to manufacturer's instructions. Plasmids were purified using the Nucleospin Plasmid EasyPure kit (Macherey-Nagel) and their sequences verified by Sanger sequencing. Each validated cloned DNA fragment was then PCR-amplified from 10 ng of purified plasmid for 14 cycles using the PrimeSTAR GXL DNA polymerase (TaKaRa, R050A) and barcoded primers from the PCR-cDNA Barcoding kit (ONT, SQK-PCB111.24) as described for FLEP-seq2 library preparation. 1.25 fmol of each library were then sequenced alongside other samples on a P2 PromethION 2 Solo for ~72 hours (Flow Cell FLO-PRO002, R9.4.1). Computational analysis of these synthetic DNA fragments was performed as described for the FLEP-seq2 library using the sequence of synthetic DNA fragments as the reference for mapping. Source data for figures are available in Supplementary Data Set S1.

Uridylation analysis of FLEP-seq2 data from (Jia et al. 2022)

Uridylation analysis was performed from mapping .bam files available on China National Center for Bioinformation with accession OMIX881 using a homemade Python script (Extract_unmappedseq_final3.py), available on Github (https://github.com/jackson-peter/FLEPseq2) The script extracts the unmapped sequence found at the 3' end of RNAs from the soft-clipped sequences of the mapping files, then searches for the nanopore adapter sequence and finally extracts the poly(A) tail and the eventual other added nucleotides. Additional tails containing at least 70% of Ts were considered as uridylated. As previously, all calculations were then performed for annotated cytoplasmic mRNAs, based on TAIR10 and ambiguously annotated RNAs were filtered out (see a list of filtered out gene identifiers in Supplementary Data Set S8). Source data for figures are available in Supplementary Data Set S1. Statistical analysis results are available in Supplementary Data Set S9.

Preparation of RNA-seq libraries

RNAseq libraries were generated from three biological replicates of flowers and after-ripened seeds using the same RNAs as those analyzed by FLEP-seq2. Total RNA was treated with DNase I (Thermo Fisher Scientific, EN0521) and purified using the RNeasy MinElute Clean-up (Qiagen, 74204). RNA-seq libraries were then produced from 500 ng of DNase-treated RNAs using the TruSeq Stranded Total RNA Library Prep Plant (Illumina, 20020610) according to manufacturer's instructions. Sequencing was

performed on an Illumina NextSeq 2000 system by the GenomEast platform (paired-end sequencing, 2 × 50 bases).

Computational analysis for RNAseq

Reads have been mapped against the TAIR10 genome using hisat2 (v.2.2.1), with the following options: -p 4 -k 20 --max-intronlen 2000 --rna-strandness R. Read counts were then extracted for each representative transcript using FeatureCounts (v.2.0.6) and the Araport11 annotation ("Araport11_GTF_genes_transposons.Jul2023.gtf") by specifying "gene" as feature type (-t option) and "gene_id" as attribute type (-g option). Differential expression analysis was performed in R (v.4.2.2) using the DESeq2 package (Love et al. 2014) (v1.38.3).

Statistical analysis

All statistical analysis were performed were using R (v. 4.2.2) on RStudio (v.2023.12.1+402). Global percentages (Figure 1A and C, Figure 2B and D, Figure 3B, Figure 5A and B, Figure 7A and Figure 8A-C), were statistically compared using the R package car (v3.1-2) applying a generalized linear model for proportions with a quasibinomial distribution. To test the difference between distributions (Figure 1B, Figure 2C, Figure 5A, Figure 7C and E, Figure S1C, Figure S1F, Figure S2A, Figure S3B, Figure S5C), medians were calculated for each replicate and samples and a linear regression (R package stats, v3.6.1) was applied, as the Shapiro-Wilk test did not show evidence of non-normality. To compare FLEP-seq2 data with genomic features in Figure 1D and in Figure S2B, the statistical analysis was performed using Pairwise Wilcoxon Rank Sum Tests with data considered as unpaired (non-parametric test, two-tailed). The multcomp R package (1.4-26) with Tukey contrasts was used for multiple comparison post hoc tests and the calculation of adjusted p-values. Differential expression analysis of RNAseq data was performed using the DESeq2 package (Love et al. 2014) (v.1.38.3). For all statistical analyses, an adjusted p-value of 0.05 (Benjamini-Hochberg) was defined as the threshold of significance except for the analysis performed in Figure 1D and Figure S2B for which the p-value threshold was set at 0.0001, due to the high number of observations. Detailed results of the statistical analysis are provided in Supplementary Data Sets S3 and S4 (RNAseq analysis) and Supplementary Data Set S9 for all other tests.

Other analyses

All plots were generated using R (v. 4.2.2) and the R package ggplot2 (v. 3.5.1) on RStudio (v.2023.12.1+402), except for heatmaps that were generated using the R pheatmap R package (v. 1.0.12). Boxplots shown in the paper displays the median, first and third quartiles (lower and upper hinges), the largest value within 1.5 times the interquartile range above the upper hinge (upper whisker) and the smallest value within 1.5 times the interquartile range below the lower hinge (lower whiskers). Publicly available RNAseq data from (Klepikova et al. 2016) (http://travadb.org) were used to analyze gene expression during seed development (Figure 6B, Figure S4 and Figure S5B). For heatmap in Figure 6B and Figure S5B, normalized read count (TMM, EdgeR) of each mRNA were divided by the maximum value of expression level among development stages, so all values vary from 0 to 1. As a reference for seed development stage, expression of mRNAs shown to be expressed at specific seed stage were also analyzed using RNAseq data from (Klepikova et al. 2016). All gene ontology analyses performed in the present paper were performed using DAVID (Huang et al. 2009b, 2009a), using an adjusted p-value of 0.05 (FDR) as threshold of significance.

Accession Numbers

NGS datasets generated during this study have been deposited in NCBI's Gene Expression Omnibus (Edgar et al. 2002) and are accessible through the GEO Series accession number GSE284704. Sequence data for the genes mentioned in this article can be found in the Arabidopsis Information Resource (TAIR, https://www.arabidopsis.org/) under the following accession numbers: AT2G45620 (*URT1*), AT2G39740 (*HESO1*), *DOG1* (AT5G45830).

List of supplementary data files

Supplementary Figure S1. FLEP-seq2 to analyze U-tailing in different Arabidopsis plant tissues. Supports Figure 1.

Supplementary Figure S2. Changes of U-tail profiles during seed development and germination. Supports Figure 2.

Supplementary Figure S3. Analysis of mRNAs differentially uridylated in *urt1-1*, *heso1-4* and *heso1-4 urt1-1*. Supports Figure 5.

Supplementary Figure S3. Analysis of mRNAs differentially uridylated in *urt1-1*, *heso1-4* and *heso1-4 urt1-1*. Supports Figure 5.

Supplementary Figure S4. Expression of *URT1* and *HESO1* during seed development. Supports Figure 6.

Supplementary Figure S6. Impact of URT1 inactivation on the germination of after-ripened seeds. Supports Figure 8.

Supplementary Data Set S1. Source data related to spike-in and uridylation analysis of FLEP-seq data from Jia et al. 2022.

Supplementary Data Set S2. Source data related to FLEP-seq data generated in the present study.

Supplementary Data Set S3. Source data related to RNAseq analysis in seeds.

Supplementary Data Set S4. Source data related to RNAseq analysis in flowers.

Supplementary Data Set S5. Expression data from publicly available RNAseq data from Klepikova et al., 2016.

Supplementary Data Set S6. Source data related to germination assays.

Supplementary Data Set S7. List of oligonucleotides and synthetic DNA fragment used in this study.

Supplementary Data Set S8: List of AGI filtered out of the FLEP-seq2 analysis.

Supplementary Data Set S9: Source data for statistical analyses.

Acknowledgments

This research was funded by Centre National de la Recherche Scientifique (CNRS) and by a research grant from the French National Research Agency ANR-22-CE12-0002. This work of the Interdisciplinary Thematic Institute IMCBio, as part of the ITI 2021-2028 program of the University of Strasbourg, CNRS and Inserm, was supported by IdEx Unistra (ANR-10-IDEX-0002), and by SFRI-STRAT'US project (ANR 20-SFRI-0012) and EUR IMCBio (ANR-17-EURE-0023) under the framework of the French Investments for the Future Program. Nanopore sequencing was performed at the IBMP Gene Expression Analysis Platform. The RNA-seq sequencing was performed by the GenomEast platform, a member of the "France Génomique" consortium (ANR-10-INBS-0009). The authors thank Abdelmalek Alioua and Sandrine Koechler from IBMP Gene Expression Analysis Platform for help in Nanopore sequencing and RNA-seq libraries preparation,

respectively, and David Pfliegler of the IBMP bioinformatics core facility for help in bioinformatic analysis. The authors thank Anthony Gobert for critical reading.

Author Contributions

H.Z. and D.G. acquired funding; H.Z. conceived and supervised the research; J.P. conceived and developed analytical pipelines; J.R. performed experiments with the assistance of S.Sa., E.U., B.L.; J.P. and H.Z. analyzed data; H.Z., J.P., J.R., S. Sa, S.Sw, D.G. discussed the data. H.Z. drafted the paper. D.G. J.P., J.R., S. Sa, S.Sw edited and revised the paper.

References

Baer BW and Kornberg RD. The protein responsible for the repeating structure of cytoplasmic poly(A)-ribonucleoprotein. J Cell Biol. 1983:**96**(3):717–721. https://doi.org/10.1083/jcb.96.3.717

Bai B, Peviani A, van der Horst S, Gamm M, Snel B, Bentsink L, and Hanson J. Extensive translational regulation during seed germination revealed by polysomal profiling. New Phytol. 2017:**214**(1):233–244. https://doi.org/10.1111/nph.14355

Bai B, Schiffthaler B, van der Horst S, Willems L, Vergara A, Karlström J, Mähler N, Delhomme N, Bentsink L, and Hanson J. SeedTransNet: a directional translational network revealing regulatory patterns during seed maturation and germination. J Exp Bot. 2023:**74**(7):2416–2432. https://doi.org/10.1093/jxb/erac394

Basbouss-Serhal I, Pateyron S, Cochet F, Leymarie J, and Bailly C. 5' to 3' mRNA Decay Contributes to the Regulation of Arabidopsis Seed Germination by Dormancy. Plant Physiol. 2017:**173**(3):1709–1723. https://doi.org/10.1104/pp.16.01933

Basbouss-Serhal I, Soubigou-Taconnat L, Bailly C, and Leymarie J. Germination Potential of Dormant and Nondormant Arabidopsis Seeds Is Driven by Distinct Recruitment of Messenger RNAs to Polysomes. Plant Physiol. 2015:**168**(3):1049–1065. https://doi.org/10.1104/pp.15.00510

Bates PD, Jewell JB, and Browse J. Rapid separation of developing Arabidopsis seeds from siliques for RNA or metabolite analysis. Plant Methods. 2013:**9**(1):9. https://doi.org/10.1186/1746-4811-9-9

Belmonte MF, Kirkbride RC, Stone SL, Pelletier JM, Bui AQ, Yeung EC, Hashimoto

M, Fei J, Harada CM, Munoz MD, et al. Comprehensive developmental profiles of gene activity in regions and subregions of the Arabidopsis seed. Proc Natl Acad Sci. 2013:**110**(5):E435–E444. https://doi.org/10.1073/pnas.1222061110

Buijs G, Vogelzang A, Nijveen H, and Bentsink L. Dormancy cycling: translation-related transcripts are the main difference between dormant and non-dormant seeds in the field. Plant J. 2020:**102**(2):327–339. https://doi.org/10.1111/tpj.14626

Cadman CSC, Toorop PE, Hilhorst HWM, and Finch-Savage WE. Gene expression profiles of Arabidopsis Cvi seeds during dormancy cycling indicate a common underlying dormancy control mechanism. Plant J. 2006:**46**(5):805–822. https://doi.org/10.1111/j.1365-313X.2006.02738.x

Chahtane H, Kim W, and Lopez-Molina L. Primary seed dormancy: a temporally multilayered riddle waiting to be unlocked. J Exp Bot. 2017:**68**(4):857–869. https://doi.org/10.1093/jxb/erw377

Chang H, Yeo J, Kim J, Kim H, Lim J, Lee M, Kim HH, Ohk J, Jeon H-Y, Lee H, et al. Terminal Uridylyltransferases Execute Programmed Clearance of Maternal Transcriptome in Vertebrate Embryos. Mol Cell. 2018:**70**(1):72-82.e7. https://doi.org/10.1016/j.molcel.2018.03.004

Chowdhury A, Mukhopadhyay J, and Tharun S. The decapping activator Lsm1p-7p–Pat1p complex has the intrinsic ability to distinguish between oligoadenylated and polyadenylated RNAs. RNA. 2007:**13**(7):998–1016. https://doi.org/10.1261/rna.502507

De Almeida C, Scheer H, Zuber H, and Gagliardi D. RNA uridylation: a key posttranscriptional modification shaping the coding and noncoding transcriptome. WIREs RNA. 2018:**9**(1):e1440. https://doi.org/10.1002/wrna.1440

Edgar R, Domrachev M, and Lash AE. Gene Expression Omnibus: NCBI gene expression and hybridization array data repository. Nucleic Acids Res. 2002:**30**(1):207–210. https://doi.org/10.1093/nar/30.1.207

Grochowski M, Lipińska-Zubrycka L, Townsend S, Golisz-Mocydlarz A, Zakrzewska-Płaczek M, Brzyżek G, Jurković B, Świeżewski S, Ralser M, and Małecki M. Uridylation regulates mRNA decay directionality in fission yeast. Nat Commun. 2024:**15**(1):8359. https://doi.org/10.1038/s41467-024-50824-w

He D, Han C, Yao J, Shen S, and Yang P. Constructing the metabolic and regulatory

pathways in germinating rice seeds through proteomic approach. PROTEOMICS. 2011:**11**(13):2693–2713. https://doi.org/10.1002/pmic.201000598

Huang DW, Sherman BT, and Lempicki RA. Bioinformatics enrichment tools: paths toward the comprehensive functional analysis of large gene lists. Nucleic Acids Res. 2009a:**37**(1):1–13. https://doi.org/10.1093/nar/gkn923

Huang DW, Sherman BT, and Lempicki RA. Systematic and integrative analysis of large gene lists using DAVID bioinformatics resources. Nat Protoc. 2009b:**4**(1):44–57. https://doi.org/10.1038/nprot.2008.211

Jia J, Lu W, Liu B, Fang H, Yu Y, Mo W, Zhang H, Jin X, Shu Y, Long Y, et al. An atlas of plant full-length RNA reveals tissue-specific and monocots–dicots conserved regulation of poly(A) tail length. Nat Plants. 2022:**8**(9):1118–1126. https://doi.org/10.1038/s41477-022-01224-9

Joly AC, Garcia S, Hily J-M, Koechler S, Demangeat G, Garcia D, Vigne E, Lemaire O, Zuber H, and Gagliardi D. An extensive survey of phytoviral RNA 3′ uridylation identifies extreme variations and virus-specific patterns. Plant Physiol. 2023:**193**(1):271–290. https://doi.org/10.1093/plphys/kiad278

Kearly A, Nelson ADL, Skirycz A, and Chodasiewicz M. Composition and function of stress granules and P-bodies in plants. Semin Cell Dev Biol. 2024:**156**:167–175. https://doi.org/10.1016/j.semcdb.2022.11.008

Klepikova A V., Kasianov AS, Gerasimov ES, Logacheva MD, and Penin AA. A high resolution map of the Arabidopsis thaliana developmental transcriptome based on RNA-seq profiling. Plant J. 2016:**88**(6):1058–1070. https://doi.org/10.1111/tpj.13312

Klupczyńska EA and Pawłowski TA. Regulation of Seed Dormancy and Germination Mechanisms in a Changing Environment. Int J Mol Sci. 2021:**22**(3):1357. https://doi.org/10.3390/ijms22031357

Le BH, Cheng C, Bui AQ, Wagmaister JA, Henry KF, Pelletier J, Kwong L, Belmonte M, Kirkbride R, Horvath S, et al. Global analysis of gene activity during Arabidopsis seed development and identification of seed-specific transcription factors. Proc Natl Acad Sci U S A. 2010:**107**(18):8063–8070. https://doi.org/10.1073/pnas.1003530107

Lim J, Ha M, Chang H, Kwon SC, Simanshu DK, Patel DJ, and Kim VN. Uridylation by TUT4 and TUT7 Marks mRNA for Degradation. Cell. 2014:**159**(6):1365–1376.

https://doi.org/10.1016/j.cell.2014.10.055

Long Y, Jia J, Mo W, Jin X, and Zhai J. FLEP-seq: simultaneous detection of RNA polymerase II position, splicing status, polyadenylation site and poly(A) tail length at genome-wide scale by single-molecule nascent RNA sequencing. Nat Protoc. 2021:**16**(9):4355–4381. https://doi.org/10.1038/s41596-021-00581-7

Love MI, Huber W, and Anders S. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. Genome Biol. 2014:**15**(12):550. https://doi.org/10.1186/s13059-014-0550-8

Malecki M, Viegas SC, Carneiro T, Golik P, Dressaire C, Ferreira MG, and Arraiano CM. The exoribonuclease Dis3L2 defines a novel eukaryotic RNA degradation pathway. EMBO J. 2013:**32**(13):1842–1854. https://doi.org/10.1038/emboj.2013.63

Meng L and Feldman L. A rapid TRIzol-based two-step method for DNA-free RNA extraction from Arabidopsis siliques and dry seeds. Biotechnol J. 2010:**5**(2):183–186. https://doi.org/10.1002/biot.200900211

Morgan M, Much C, DiGiacomo M, Azzi C, Ivanova I, Vitsios DM, Pistolic J, Collier P, Moreira PN, Benes V, et al. mRNA 3′ uridylation and poly(A) tail length sculpt the mammalian maternal transcriptome. Nature. 2017:**548**(7667):347–351. https://doi.org/10.1038/nature23318

Nakabayashi K, Bartsch M, Xiang Y, Miatton E, Pellengahr S, Yano R, Seo M, and Soppe WJJ. The Time Required for Dormancy Release in Arabidopsis Is Determined by DELAY OF GERMINATION1 Protein Levels in Freshly Harvested Seeds. Plant Cell. 2012:**24**(7):2826–2838. https://doi.org/10.1105/tpc.112.100214

Nakabayashi K, Okamoto M, Koshiba T, Kamiya Y, and Nambara E. Genome-wide profiling of stored mRNA in Arabidopsis thaliana seed germination: epigenetic and genetic regulation of transcription in seed. Plant J. 2005:**41**(5):697–709. https://doi.org/10.1111/j.1365-313X.2005.02337.x

Narsai R, Law SR, Carrie C, Xu L, and Whelan J. In-Depth Temporal Transcriptome Profiling Reveals a Crucial Developmental Switch with Roles for RNA Processing and Organelle Metabolism That Are Essential for Germination in Arabidopsis. Plant Physiol. 2011:**157**(3):1342–1362. https://doi.org/10.1104/pp.111.183129

Née G, Xiang Y, and Soppe WJ. The release of dormancy, a wake-up call for seeds to

germinate. Curr Opin Plant Biol. 2017:**35**:8–14. https://doi.org/10.1016/j.pbi.2016.09.002

Nelson SK and Steber CM. Transcriptional mechanisms associated with seed dormancy and dormancy loss in the gibberellin-insensitive sly1-2 mutant of Arabidopsis thaliana. PLOS ONE. 2017:**12**(6):e0179143. https://doi.org/10.1371/journal.pone.0179143

Rajjou L, Gallardo K, Debeaujon I, Vandekerckhove J, Job C, and Job D. The Effect of α-Amanitin on the Arabidopsis Seed Proteome Highlights the Distinct Roles of Stored and Neosynthesized mRNAs during Germination. Plant Physiol. 2004:**134**(4):1598–1613. https://doi.org/10.1104/pp.103.036293

Rajjou L, Lovigny Y, Groot SPC, Belghazi M, Job C, and Job D. Proteome-Wide Characterization of Seed Aging in Arabidopsis: A Comparison between Artificial and Natural Aging Protocols. Plant Physiol. 2008:**148**(1):620–641. https://doi.org/10.1104/pp.108.123141

Ren G, Chen X, and Yu B. Uridylation of miRNAs by HEN1 SUPPRESSOR1 in Arabidopsis. Curr Biol. 2012:**22**(8):695–700. https://doi.org/10.1016/j.cub.2012.02.052

Ren G, Xie M, Zhang S, Vinovskis C, Chen X, and Yu B. Methylation protects microRNAs from an AGO1-associated activity that uridylates 5' RNA fragments generated by AGO1 cleavage. Proc Natl Acad Sci. 2014:**111**(17):6365–6370. https://doi.org/10.1073/pnas.1405083111

Ren Y, Ma X, Song B, Yang X, Chen Y, Yu Y, Chen X, Mo B, and Wang X. HEN1 SUPPRESSOR1 stabilizes polymerase IV RNAs via uridylation in Arabidopsis. Plant Physiol. 2023:**193**(1):186–189. https://doi.org/10.1093/plphys/kiad297

Rissland OS and Norbury CJ. Decapping is preceded by 3' uridylation in a novel pathway of bulk mRNA turnover. Nat Struct Mol Biol. 2009:**16**(6):616–623. https://doi.org/10.1038/nsmb.1601

Sachs AB, Davis RW, and Kornberg RD. A single domain of yeast poly(A)-binding protein is necessary and sufficient for RNA binding and cell viability. Mol Cell Biol. 1987:**7**(9):3268–3276. https://doi.org/10.1128/mcb.7.9.3268-3276.1987

Sajeev N, Bai B, and Bentsink L. Seeds: A Unique System to Study Translational Regulation. Trends Plant Sci. 2019:**24**(6):487–495. https://doi.org/10.1016/j.tplants.2019.03.011

Sajeev N, Koornneef M, and Bentsink L. A commitment for life: Decades of unraveling

the molecular mechanisms behind seed dormancy and germination. Plant Cell. 2024:**36**(5):1358–1376. https://doi.org/10.1093/plcell/koad328

Sano N, Rajjou L, and North HM. Lost in Translation: Physiological Roles of Stored mRNAs in Seed Germination. Plants. 2020:**9**(3):347. https://doi.org/10.3390/plants9030347

Scheer H, de Almeida C, Ferrier E, Simonnot Q, Poirier L, Pflieger D, Sement FM, Koechler S, Piermaria C, Krawczyk P, et al. The TUTase URT1 connects decapping activators and prevents the accumulation of excessively deadenylated mRNAs to avoid siRNA biogenesis. Nat Commun. 2021:**12**(1):1–17. https://doi.org/10.1038/s41467-021-21382-2

Sement FM, Ferrier E, Zuber H, Merret R, Alioua M, Deragon J-M, Bousquet-Antonelli C, Lange H, and Gagliardi D. Uridylation prevents 3′ trimming of oligoadenylated mRNAs. Nucleic Acids Res. 2013:**41**(14):7115–7127. https://doi.org/10.1093/nar/gkt465

Solis-Miranda J, Chodasiewicz M, Skirycz A, Fernie AR, Moschou PN, Bozhkov PV, and Gutierrez-Beltran E. Stress-related biomolecular condensates in plants. Plant Cell. 2023:**35**(9):3187–3204. https://doi.org/10.1093/plcell/koad127

Song M-G and Kiledjian M. 3′ Terminal oligo U-tract-mediated stimulation of decapping. RNA. 2007:**13**(12):2356–2365. https://doi.org/10.1261/rna.765807

Thomas MP, Liu X, Whangbo J, McCrossan G, Sanborn KB, Basar E, Walch M, and Lieberman J. Apoptosis Triggers Specific, Rapid, and Global mRNA Decay with 3′ Uridylated Intermediates Degraded by DIS3L2. Cell Rep. 2015:**11**(7):1079–1089. https://doi.org/10.1016/j.celrep.2015.04.026

Tu B, Liu L, Xu C, Zhai J, Li S, Lopez MA, Zhao Y, Yu Y, Ramachandran V, Ren G, et al. Distinct and Cooperative Activities of HESO1 and URT1 Nucleotidyl Transferases in MicroRNA Turnover in Arabidopsis. PLoS Genet. 2015:**11**(4):e1005119. https://doi.org/10.1371/journal.pgen.1005119

Vigh ML, Thieffry A, Arribas-Hernández L, and Brodersen P. Arabidopsis terminal nucleotidyl transferases govern secondary siRNA production at distinct steps. 2024:2024.05.27.596008. https://doi.org/10.1101/2024.05.27.596008

Wang X, Kong W, Wang Y, Wang J, Zhong L, Lao K, Dong X, Zhang D, Huang H, Mo

B, et al. Uridylation and the SKI complex orchestrate the Calvin cycle of photosynthesis through RNA surveillance of TKL1 in Arabidopsis. Proc Natl Acad Sci. 2022:**119**(38):e2205842119. https://doi.org/10.1073/pnas.2205842119

Wang X, Zhang S, Dou Y, Zhang C, Chen X, Yu B, and Ren G. Synergistic and Independent Actions of Multiple Terminal Nucleotidyl Transferases in the 3' Tailing of Small RNAs in Arabidopsis. PLOS Genet. 2015:**11**(4):e1005091. https://doi.org/10.1371/journal.pgen.1005091

Warkocki Z, Liudkovska V, Gewartowska O, Mroczek S, and Dziembowski A. Terminal nucleotidyl transferases (TENTs) in mammalian RNA metabolism. Philos Trans R Soc B Biol Sci. 2018:**373**(1762):20180162. https://doi.org/10.1098/rstb.2018.0162

Yu S and Kim VN. A tale of non-canonical tails: gene regulation by post-transcriptional RNA tailing. Nat Rev Mol Cell Biol. 2020:**21**(9):542–556. https://doi.org/10.1038/s41580-020-0246-8

Zhao Y, Yu Y, Zhai J, Ramachandran V, Dinh TT, Meyers BC, Mo B, and Chen X. The Arabidopsis Nucleotidyl Transferase HESO1 Uridylates Unmethylated Small RNAs to Trigger Their Degradation. Curr Biol. 2012:**22**(8):689–694. https://doi.org/10.1016/j.cub.2012.02.051

Zigáčková D and Vaňáčová Š. The role of 3' end uridylation in RNA metabolism and cellular physiology. Philos Trans R Soc B Biol Sci. 2018:**373**(1762):20180171. https://doi.org/10.1098/rstb.2018.0171

Zuber H, Scheer H, Ferrier E, Sement FM, Mercier P, Stupfler B, and Gagliardi D. Uridylation and PABP cooperate to repair mRNA deadenylated ends in Arabidopsis. Cell Rep. 2016:**14**(11):2707–2717. https://doi.org/10.1016/j.celrep.2016.02.060

Zuber H, Scheer H, Joly A-C, and Gagliardi D. Respective Contributions of URT1 and HESO1 to the Uridylation of 5' Fragments Produced From RISC-Cleaved mRNAs. Front Plant Sci. 2018:**9**:1438. https://doi.org/10.3389/fpls.2018.01438

Figure 1: Comprehensive analysis of mRNA U-tailing in different Arabidopsis plant tissues. FLEPseq2 data from Jia et al. 2022 were re-analyzed with a pipeline adapted to analyze U-tails. Data are shown for two replicates. Bulk (A) and intergenic (B) percentages of uridylation. Intergenic percentages were calculated for mRNAs detected with at least 50 reads for each replicate. C, % of tails according to the the number of Us. Individual points and the bar plot show individual replicates and the average, respectively. The average percentage of tail > 2 Us are indicated above each bar plot. D, Feature analysis for the 10% and 25% most and less uridylated mRNAs or for other mRNAs. Each boxplot displays the median, first and third quartiles (lower and upper hinges), the largest value within 1.5 times the interquartile range above the upper hinge (upper whisker) and the smallest value within 1.5 times the interquartile range below the lower hinge (lower whiskers). Significantly different values (adjusted p-value < 0.05) are labelled by different letters.

A

Two weeks after flowering

1) Flower
Unopened flower buds

2) Dev. seed series 1
Isolated developing seeds**

3) Dev. silique series 1
Developing siliques with seeds**

Three weeks after flowering

4) Dev. silique series 2
Developing siliques with seeds**

Eight weeks after flowering

5) Dry seed

6) Imbibed seed
Seeds imbibed 24h in water

**Pool of developing stages
(from ~5 to 14 DAF)

Seed development

Figure 2: Dynamic of uridylation levels and profiles in seeds. A. Developmental stages used for the preparation of FLEP-seq2 libraries. All samples were harvested for three biological replicates from a minimum of five plants each. B-C. Bulk (B) and intergenic (C) percentages of uridylation. Intergenic percentages were calculated for mRNAs detected with at least 50 reads for each replicate. D. % of tails according to the number of Us. Individual points and the bar plot show individual replicates and the average, respectively. The average percentage of tails > 2 Us is indicated above each bar plot. For all plots, significantly different values (adjusted p-value < 0.05) are labelled by different letters.

Figure 3: Modulation of poly(A) tail size in seeds. FLEP-seq2 data from developing seeds and siliques (containing seeds), dry seeds and 24h-imbibed seeds. Data are shown for three replicates. A, Distribution of poly(A) tail sizes from 10 to 200 As for non-uridylated (left panel) and uridylated (right panel) reads. Percentages were calculated using the total number of sequences as denominator. Bar plots next to each distribution show the percentage of reads with no or short A-tail (<10As), with oligoadenylated A-tail between 10 and 25 As, or with poly(A) tail longer than 25As. B, Percentage of uridylation according to A-tail sizes. Percentages were calculated using the total number of sequences of each A-tail population. For all plots of the figure, individual points show the three replicates while color-coded area and bar plots show the average of all replicates.

Figure 4: Changes of poly(A) tail size profiles during seed development and germination. Heatmap showing the modulations of poly(A) tail sizes during seed development and germination. % of reads according to poly(A) tail sizes (5nt-bin) were calculated using the total number of sequences as denominator and were z-score rescaled. Transcripts were clustered based on their similarity of polyadenylation profiles (k=6).

Figure 5: Contribution of URT1 and HESO1 to mRNA uridylation in Arabidopsis. FLEP-seq2 data from developing seeds and siliques (containing seeds), dry seeds and 24h-imbibed seeds for WT, *urt1-1*, *heso1-4* and *heso1-4 urt1-1*. Data are shown for three replicates. Bulk (top panel) and intergenic (lower panel) percentages of uridylation. Intergenic percentages were calculated for mRNAs detected with at least 50 reads for each replicate and genotype. Each boxplot displays the median, first and third quartiles (lower and upper hinges), the largest value within 1.5 times the interquartile range above the upper hinge (upper whisker) and the smallest value within 1.5 times the interquartile range below the lower hinge (lower whiskers). Significantly different values (adjusted p-value < 0.05) are labelled by different letters. B, Proportion of mRNAs differentially uridylated between mutants and WT, or between the *heso1-4 urt1-1* and *urt1-1* (adjusted p-value < 0.05, generalized linear model for proportion, quasibinomial distribution). mRNAs with a significant increase and decrease percentage are shown in the top and lower panel, respectively. Number of mRNAs are indicated above the bars. C, Venn diagrams showing the intersection between the different lists of mRNAs showing a decreased uridylation percentage.

Figure 6: Impact of URT1 and HESO1 inactivation on flower and seed transcriptome. A, Differential expression analysis in flowers and dry seeds of WT, *urt1-1*, *heso1-4* and *heso1-4 urt1-1*. RNAseq data were produced from three biological replicates. Bar plots show the number of significantly differentially regulated transcripts (adjusted p-value < 0.05, DEseq2) and venn diagrams show the intersection between the different lists. B, Heatmap showing the expression pattern, in a WT context, of transcripts down and up-regulated by URT1 inactivation during seed development, from unfertilized ovules to dry seeds. Expression data are publicly available RNAseq data from Klepikova et al., 2016. As a reference of the different seed development stages, the small heatmap shows the expression pattern during seed development of stage-specific mRNAs (see description in Supplemental Dataset S5). For each mRNA, normalized read count (TMM, EdgeR) were divided by the maximum value of expression level among development stages, so all values vary from 0 to 1. Transcripts were clustered based on their similarity of expression profiles (k=8). Transcripts associated to the cytoplasmic ribosome gene ontology are highlighted by black rectangles.

Figure 7: URT1-mediated uridylation triggers a massive accumulation of short A-tails in seeds. FLEP-seq2 data from developing seeds and siliques (containing seeds), dry seeds and 24h-imbibed seeds. Data are shown for three replicates. A, Distribution of poly(A) tail sizes from 10 to 200 As in WT, *urt1-1*, *heso1-4* and *heso1-4 urt1-1*. Percentages were calculated using the total number of sequences as denominator. Bar plots bellow each distribution show the percentage of reads with no or short A-tail (<10As) or with oligoadenylated A-tail between 10 and 25 As. Individual points show the three replicates while color-coded area or bar plot show the average of all replicates. B and C, Comparison of the poly(A) tail sizes between WT and *urt1-1* for the most and the less impacted mRNAs in term of poly(A) shortening (score >20% or < 2%, respectively). B, Intergenic distribution of poly(A) tail sizes from 10 to 200 As. C, Intergenic proportion of reads with no or short A-tail (<10As) or with oligoadenylated A-tail between 10 and 25 As. D and E, Comparison of the poly(A) tail sizes between WT and *urt1-1* for mRNAs down or up-regulated regulated by *urt1-1* mutation. D, Intergenic distribution of poly(A) tail sizes from 10 to 200 As. E, Intergenic proportion of reads with no or short A-tail (<10As) or with oligoadenylated A-tail between 10 and 25 As. In B-E, percentages were calculated using the total number of sequences per mRNA as denominator. In B and D, the lines and the grey shadow show the median and the first and third quartiles of percentages across mRNAs, respectively. Significantly different values (adjusted p-value < 0.05) are labelled by different letters.

Figure 8: URT1 inactivation alters primary dormancy level. A, Level of primary dormancy of WT and *urt1-1* freshly harvested seeds. Germination assays were repeated four times for four independent batch of seeds. The bar plots show the % of germinated seeds after seven days. Individual points show the percentages for seeds harvested from individual plants and bar plots show the average. B, % of uridylation of the *DOG1* mRNA in dry seeds of WT, *urt1-1*, *heso1-4* and *heso1-4 urt1-1* based on FLEP-seq2 data. C, % of non-polyadenylated mRNAs (< 10As) of the *DOG1* mRNA in dry seeds of WT, *urt1-1*, *heso1-4* and *heso1-4 urt1-1* based on FLEP-seq2 data. D, Relative expression of *DOG1* (normalized to WT) in dry seeds of WT, *urt1-1*, *heso1-4* and *heso1-4 urt1-1* based on RNAseq data. Individual points show individual biological replicates, and bar plots show the average. For all graphs, significantly different values (adjusted p-value < 0.05) are labelled by different letters.