1 Kinase Plasticity in Response to Vandetanib Enhances Sensitivity to Tamoxifen in Estrogen
2 Receptor Positive Breast Cancer
3

Rasha T. Kakati MD[1], Austin A. Whitman MS[1], Santiago Haase PhD[1], Attila T. Szenasi PhD[1],
Christine HNC Thai BS[1,] Elizabeth Brunk PhD[1,2], Denis O. Okumu PhD[2], Michael P. East PhD[2],
Charles M. Perou PhD[1,3,4,5], Gary L. Johnson, PhD[1,2], and Philip M. Spanheimer MD[1,4,6]

[1]Lineberger Comprehensive Cancer Center, University of North Carolina, Chapel Hill, NC
[2]Department of Pharmacology, University of North Carolina, Chapel Hill, NC
[3]Department of Pathology and Laboratory Medicine, University of North Carolina at Chapel
Hill, Chapel Hill, NC
[4]Department of Genetics, University of North Carolina, Chapel Hill, NC
[5]Computational Medicine Program, University of North Carolina, Chapel Hill, NC
[6]Department of Surgery, University of North Carolina, Chapel Hill, NC

14
15 Running title: Vandetanib ER+ Breast Cancer
16
17 Keywords: Breast cancer, endocrine resistance, tyrosine kinase inhibition, functional proteomics,
18 targeted therapeutics
19

Financial Support: This work was supported in part by the National Institutes of Health grants
K08CA280388, R37CA292075, P30CA016086, and a Solid Tumor Research Scholars Grant
from Gilead Sciences. CMP was supported by funds from the NCI breast SPORE program (P50-
CA058223).

Correspondence should be addressed to:
Philip M. Spanheimer MD
Lineberger Comprehensive Cancer Center
University of North Carolina at Chapel Hill
170 Manning Drive, Suite 1149
Chapel Hill, NC 27599-7213 USA
Phone: 919-966-5221; Fax: 919-966-8806
E-mail: Philip_Spanheimer@med.unc.edu

33
34
35 Disclosure of conflict of interests: C.M.P is an equity stockholder and consultant of BioClassifier
36 LLC; C.M.P is also listed as an inventor on patent applications for the Breast PAM50 Subtyping
37 assay. The remaining authors declare no potential conflicts of interest.
38
39

40   **ABSTRACT**
41
Resistance to endocrine therapy (ET) is common in estrogen receptor (ER) positive breast

cancer. Multiple studies have demonstrated that upregulation of MAPK signaling pathways

contributes to ET resistance. Herein we show that vandetanib treatment enhances sensitivity to

ET in ET-sensitive and -resistant ER+ breast cancer models. Vandetanib treatment alters the

gene expression program of ER+ breast cancer cells resulting in a less proliferative and more

estrogen responsive Luminal-A like character. Tyrosine kinase network reprogramming was

assessed using multiplexed kinase inhibitor beads-mass spectrometry (MIB/MS) assay to identify

adaptive resistance mechanisms to vandetanib treatment, including upregulation of HER2

activity. Co-treatment to inhibit HER2 with lapatinib enhanced sensitivity to vandetanib,

demonstrating biologic activity of HER2 upregulation. Using a CRISPR knockout model, we

demonstrate that vandetanib effects are partially mediated by RET receptor tyrosine kinase.

Finally, we use our operating room-to-laboratory assay that measures drug response in individual

primary tumor cells in short term cultures to demonstrate conserved gene expression changes,

including increased HER2 activity signatures, in vandetanib treated cells, and identify features

associated with vandetanib response. These results support future investigation of RET targeting

strategies considering reprogrammed networks, such as activated HER2, in patients with ET

resistant ER+ breast cancer.

SIGNIFICANCE

Vandetanib enhances sensitivity to tamoxifen in ER+ breast cancer cells by programing kinase
signaling networks, which can be used to select patients most likely to respond and develop more
efficacious co-treatment strategies.
64

INTRODUCTION

Approximately 75% of breast cancers fall within the estrogen receptor (ER)+/luminal subtypes(1-3). Over 40,000 women are expected to die from breast cancer each year and most of these will have ER+ disease(4). Most patients with ER+ breast cancer will be treated with endocrine therapy (ET) targeting ER. However, resistance to ET occurs in about 1/3 of early cancers and develops over time in all metastatic ER+ cancers(5-7). Targeted inhibitors to PI3K, mTOR, and CDK4/6 have shown efficacy in patients with ET resistant disease(8-11). However, responses are sometimes short lived and there are a significant number of patients in need of durable, effective therapy.

Multiple studies have demonstrated that activation of ERK/MAPK pathways is an important mechanism by which ER+ breast tumors resist ET(12-18). Gain of function mutations in ERK/MAPK nodes or upstream receptor tyrosine kinase (RTK) are seen in ET-resistant tumors, although these are only present in about 13% of cases(12). The rearranged during transfection (RET) receptor tyrosine kinase is expressed in approximately 50% of ER+ breast tumors and is more highly expressed in ET resistant tumors(19-22). Expression of RET and co-expression of its activating ligand glial derived neurotropic factor (GDNF) is associated with transciptional signatures of ERK/MAPK activation and worse outcomes in human ER+ breast tumors.(23) Further, RET is overexpressed in ET-resistant breast cancer organoids that contain gain of function ESR1 (encoding the estrogen receptor) mutations and inhibiting RET suppresses tumor growth(24).

Our work has shown that tyrosine kinase inhibitors (TKIs), such as vandetanib and

sunitinib, that inhibit RET reduce proliferation and enhance sensitivity to ET in ER+ breast

cancer cell lines. Further, dual treatment with vandetanib and tamoxifen reduced breast tumor

xenograft growth greater than either agent alone(25, 26). In RET+ primary breast tumors, we

showed that *in* vitro treatment with sunitinib reduces phosho-ERK.(27) These findings

support the use of vandetanib to reduce ERK/MAPK activation and improve response to

endocrine therapy.

Despite promise in the pre-clinic, clinical trials with vandetanib have had

disappointing results. However, the design of these trials failed to account for potentially

predictive biomarkers including ER and RET or only evaluated vandetanib as a

monotherapy. For example, a phase II trial of single-agent vandetanib in 46 patients with

metastatic breast cancer did not demonstrate a benefit(28), but did not report receptor status,

making it difficult to interpret the extent to which this trial included patients that mechanistically

would be predicted to respond. Our work, including the findings presented herein, suggests

that vandetinib may provide a clinical benefit for some patients. More recent TKI trials with

more specific inclusion criteria have shown more promising results and have renewed interest in

TKI for ET resistant ER+ breast cancer. A phase II trial of cabozantinib, in metastatic

ER+/HER2- was positive by objective response rate(29), as was a phase Ib/II trial of lenvatinib

with letrozole(30). More precise biomarkers are needed to identify patients most likely to

respond to TKI therapy. To develop more precise biomarkers and elucidate bypass resistance

mechanisms, we sought to investigate how vandetanib alters kinase signaling pathways in ET-

sensitive and ET-resistant models of ER+/HER2- breast cancer.

METHODS

Cell Lines

We obtained ER+ breast cancer cell lines parental MCF7, tamoxifen-resistant MCF7

(MCF7-TAM), exemestane-resistant MCF7 (MCF7-EXE), parental T47D, and tamoxifen-

resistant T47D (T47D-TAM), from the American Type Culture Collection (ATCC). Cell lines

were cultured using Dulbecco's Modified Eagle Medium (DMEM) (Gibco) with 10% fetal

bovine serum (Gibco) and 1% penicillin-streptomycin (Life Technologies) at 37°C and 5% CO2.

MCF7-TAM and T47D-TAM were maintained by chronic treatment with 1 μM of tamoxifen and

MCF7-EXE with 0.1 μM of exemestane.

Chemicals and Treatment

Vandetanib (SelleckChem) and z-4-Hydroxytamoxifen (4OH-Tam) (SelleckChem) was

dissolved in dimethyl sulfoxide (DMSO). GDNF (R&D Systems) was dissolved in phosphate-

buffered saline (PBS) and treatments were done at a final concentration of 25 ng/mL. Pralsetinib

was obtained from Chemietek, and lapatinib from Medchem Express LLC. Treatments were

performed for 24 hours unless otherwise specified.

Cell Viability Assays

Cells were plated on opaque-walled 96-well plate at a density of 4,000 cells per well in

technical triplicates and allowed to adhere overnight prior to the addition of the specified

inhibitors. Viability was assessed using the CellTiter-Glo (CTG) Luminescent Cell Viability

Assay (Promega). Plates were read using a Synergy 2 microplate reader. Samples were averaged

over biological triplicates. Statistical analysis and plotting were performed using GraphPad

Prism 9.

Western Blots

Cells were lysed in Pierce RIPA buffer containing Halt protease inhibitor cocktail and

PhosStop phosphatase inhibitor (Thermo Scientific). Proteins were resolved by SDS

polyacrylamide gel electrophoresis, transferred to PVDF membrane, and blocked with 10%

BSA, then incubated overnight with the indicated primary antibody in blocking buffer.

Membranes were washed three times in tris buffered saline containing 0.1% tween-20 then then

incubated for 1 hour with HRP-conjugated secondary antibody. Blots were developed using a

Chemidoc-touch digital imager and band intensity quantified using ImageJ

(https://imagej.net/ij/index.html).The antibodies used for Western blot included: p44/42 MAPK

(ERK1/2) (Cell Signaling Technologies), Phospho-p44/42 MAPK (Erk1/2) (Cell Signaling

Technologies), RET (Abcam), and β-Actin (SantaCruz).

RET CRISPR Knockout

An sgRNA was designed to target exon 3 of RET (AGCACTCCCGCTCACGGTC),

cloned into the pSpCas9(BB)-2A-Puro(PX459) V2.0 (Addgene) plasmid, and verified by Sanger

sequencing. MCF-7 cells were cultured in antibiotic free media and grown to a confluency of

70%-90%. Cells were then transfected with sgRNA plasmid using FuGENE 4K Transfection

Reagent according the manufacturer's recommendations. After 72 hours, cells were passaged

onto 15 cm plates at a low density and selected with 1 μg/mL puromycin (Gibco) for 3 days. Cell

colonies were then isolated by pipet, and DNA was isolated using the Quick-DNA Miniprep Kit

(Zymo Research). The target sequence was then amplified by PCR and then sequenced for

verification of a frameshift mutation. Both RET alleles were found to have frameshift mutations

resulting in the introduction of premature stop codons.

mRNA Sequencing and Data Analysis

Cells were treated with 10 µM vandetanib or vehicle for 24 hours in biological triplicates

and total RNA was isolated (Qiagen). mRNA sequencing libraries were made using TruSeq

RNA Library Prep Kit (Illumina) and sequenced using the Illumina NextSeq2000. Plots and

statistical analyses were performed using RStudio 2022.07.1 Build 554.

Differential gene expression (DGE) was determined using the DESeq2 R package using

raw mRNA-seq counts from STAR/Salmon. Genes with less than 10 counts across all samples

were omitted before analysis and counts were filtered and normalized to the upper quartile using

NOISeq. Significant differentially expressed genes were defined by using a Log2FC (log2 fold

change) cutoff of 0.5 for upregulation and -0.5 for downregulation, with a false discovery rate

(FDR) < 0.05. Volcano plots of all genes were plotted using the EnhancedVolcano R package.

Enrichment of gene sets was then investigated through hypergeometric analysis of an

unranked list of upregulated or downregulated elements using the enricher function of the

clusterProfiler package. Gene sets from MSigDB were first compiled and screened for keywords

of interest pertaining to breast cancer, including estrogen and ER signaling.

METABRIC and SCAN-B databases were obtained for survival analysis and filtered for

ER+/HER2- patients treated with ET. A median-centroid approach was implemented to assign a

gene set score per patient. Kaplan-Meier (KM) plots were then computed using the survminer

and survival packages.

MIB/MS and Data Analysis

Cells were treated with vandetanib at its IC90 for 48 hours. Cells were harvested by

scraping and cell pellets obtained for proteomic analysis using the MIB/MS technology(31, 32).

Scaled label free quantification (LFQ) values were obtained and a correlation plot of samples

was computed to assess homogeneity of triplicates and heterogeneity of treatment conditions for

quality control using ggplot2. Differential protein expression (DPE) was analyzed using the

limma R package using scaled LFQ values. Significant differentially expressed proteins were

defined using a Log2FC (log2 fold change) cutoff of 0.5 for upregulation and -0.5 for

downregulation, along with a p-value < 0.05.

Primary ER+/HER2- Human Tumor Cells

A primary tumor sample was obtained from a patient with ER+/HER2- invasive ductal

carcinoma. The tumor was processed fresh as previously described(33). In brief, tumor tissue

was transported immediately to the laboratory after surgical resection, dissociated into a single

cell suspension, and treated with vandetanib (10 µM) or control media for 12 hours. After

treatment, single cell RNA sequencing libraries were made using the 10x Genomics Chromium

Next GEM Single Cell 3' v3.1 and sequenced on the NextSeq2000 (Illumina) with pair-end

sequencing.

Single cell RNA sequencing analysis

Reads were aligned and counted using Cellranger count function with default parameters.

The counts and features matrixes were used to generate Seurat objects. The Seurat objects

corresponding to each treatment group were merged with Seurat's function merge(), and low

quality cells were removed. Doublets were removed using scDblFinder

(https://pubmed.ncbi.nlm.nih.gov/35814628/) and cells were further filtered with the following

parameters: nFeature_RNA > 200 & nFeature_RNA < 6000 & percent.mt < 10.

Cell types were annotated on the combined Seurat using SingleR using the cell type atlas

HumanPrimaryCellAtlasData() accessed with the package celldex. InferCNV was used on the

combined Seurat to predict copy number alterations, defining the immune cells

("T_cells","Macrophage","DC","Monocyte","NK_cell") as reference cells (cells without CNVs).

The tumor cells were identified using the copy number heatmap per cell.

Differential analysis of gene expression was performed on the tumor cells to compare

vandetanib-treated versus non-treated cells using findmarkers(). Gene Set Enrichment Analysis

was then performed using the list of differentially expressed genes, ranking the genes according

to log2(FC)/FDR, and the ranked gene lists were used in the GSEA software

(https://pubmed.ncbi.nlm.nih.gov/16199517/)   (https://pubmed.ncbi.nlm.nih.gov/12808457/).

Gene signatures were scored using AddModuleScore_UCell() from the UCell package.

We used the package NMF to build a non-negative matrix factorization (NMF) model of the

tumor cells using a list of genes differentially expressed upon vandetanib treatment in MCF7

cells from this study. The Seurat was subset to contain only the above-mentioned genes, and the

NMF model was fitted to 8 factors. Heatmaps were built for each treatment group to assess

enrichment of cells to different NMF factors. Each cell was assigned to a unique NMF according

to the NMF with maximum coefficient value for that cell, and the percentage of cells belonging

to each NMF factor was calculated for each treatment group. DE analysis was performed

comparing cells in certain NMF factors.

Study Approval

215   Patient samples were obtained after written informed consent under a protocol approved

216   by the institutional review board at the University of North Carolina (IRB#22-1888) in

217   accordance with the US Common Rule.

Sex as a Biologic Variable

219   Breast cancer occurs primarily in women and so primary tumor cells were obtained from

220   a female patient with breast cancer.

Statistics

222   All experiments were performed with at least 3 biological or technical replicates. Small

223   sample number comparisons were analyzed with an unpaired t test with Welch's correction for

224   non-equal SD. Linear regressions and nonlinear regression curves were compared by statistical

225   tests on the parameters of the curves using the extra sum of squares F test. Experiments with

226   large sample numbers were analyzed by Wilcoxon test.

Data Availability

Single cell and bulk RNA sequencing data was made available at the Sequence Read

Archive (SRA) under BioProject PRJNA1177387. Mass spectrometry protein sequencing results

are included in this manuscript as Supplementary Table 1.

Code Availability

All original code has been deposited on the Github, and it is publicly available at the

Github repository at https://github.com/santiagohaase/Spanheimer-lab-Kinase-Plasticity-in-

Response-to-Vandetanib-Enhances-Sensitivity-to-Tamoxifen

RESULTS

Vandetanib Inhibits ERK Activation and Viability in ER+ Breast Cancer Cells

We previously showed that vandetanib treatment decreases ERK activation in MCF7

breast cancer cells. To investigate if vandetanib similarly decreases ERK activation in ET-

resistant ER+ breast cancer cells we used MCF7 and two ET-resistant derivatives, MCF7-TAM

(tamoxifen) and MCF7-EXE (exemestane) which had been cultured long-term under drug

treatment to develop resistance. pERK/ERK was reduced in MCF7 with vandetanib treatment (-

81%, p<0.001). Similarly, pERK/ERK was reduced in MCF-TAM with vandetanib treatment (-

78%, p=0.025) and MCF7-EXE (-78%, p=0.01), Fig. 1A. Dose response curves demonstrated an

IC50 for vandetanib of 8.5 μM in MCF7, 10.4 μM in MCF7-TAM, and 15.9 μM in MCF7-EXE

(Fig. 1B). Similar vandetanib sensitivities were seen in T47D and T47D-TAM (Supplementary

Fig. 1)

Vandetanib Sensitizes ER+ Breast Cancer Cells to Tamoxifen

Because we had previously shown interaction between vandetanib and ER downstream

signaling, we tested if co-treatment with vandetanib enhanced antiproliferative effects of

tamoxifen. The IC50 of the active metabolite of tamoxifen, 4OH-Tam, was 10.7 μM in MCF7,

15.8 μM in MCF7-TAM, and 14.1 μM in MCF7-EXE (Fig. 1C) demonstrating decreased

sensitivity to tamoxifen with chronic ET treatment.Treatment with 10 μM vandetanib resulted in

left shift of the 4OH-Tam sensitivity curve in all 3 cell lines with the IC50 for 4OH-Tam of 3.8

μM in MCF7, 4.8 μM in MCF7-TAM, and 4.8 μM in MCF7-EXE. This demonstrates that cells

are more sensitive to tamoxifen in the presence of vandetanib. To confirm synergy, we calculated

the combination index (CI) of a combined treatment of 5 μM vandetanib with 2 μM 4OH-Tam

(Fig. 1D). When compared to the individual treatments (i.e., Tamoxifen or vandetanib alone) the

combination index was 0.47 for MCF7-TAM, 0.55 for MCF7-EXE and 0.59 for MCF7,

indicating increased synergy in ET-chronically treated cells.

Vandetanib Regulated Genes in ET Sensitive and Resistant ER+ Breast Cancer

To delineate the molecular basis for how ER+ breast cancer cells respond to vandetanib

treatment resulting in increased sensitivity to tamoxifen, we performed mRNA sequencing after

treatment with vandetanib (10 μM) compared to vehicle. In MCF7, downregulated genes were

enriched for MAPK signaling and proliferative signatures, Fig. 2A, consistent with loss of pERK

observed in Fig. 1A. Recently derived KRAS/ERK activity signatures were also depleted in

treated cells (34). Upregulated genes were enriched for signatures of estrogen response and

sensitivity to tamoxifen consistent with sensitization of MCF7 cells to tamoxifen (Fig. 1D). Up-

regulated and down-regulated gene sets in MCF7-TAM (Fig. 2B) and MCF7-EXE (Fig. 2C)

showed similar depletion of proliferative and MAPK associated signatures and enrichment of

estrogen and tamoxifen response signatures. Similar transcriptional effects of vandetanib were

seen in T47D and T47D-TAM (Supplementary Fig. 2).

Because estrogen response signatures were enriched and proliferative signatures reduced

with vandetanib treatment, which are hallmark features of the indolent and ET responsive

Luminal A molecular subtype, we analyzed the effect of vandetanib on correlation to genes that

define the LumA subtype on the PAM50 assay (LumA centroid). This analysis demonstrated

vandetanib treatment resulted in a stronger LumA correlation supporting that treatment is

inducing a more indolent and ET responsive transcriptional phenotype (Fig. 2D).

To elucidate common and unique vandetanib regulated genes we determined overlap

across the parental and ET resistant MCF7 cell lines (Fig. 2E). Notably, significant commonly

upregulated genes included *SELENBP1* and *BAG1*, which are associated with reduced

proliferation, improved survival, and increased ER transcriptional activity in ER+ breast

tumors(35-37). In the common downregulated genes, *AURKA* is associated with poor survival in

ER+ breast cancer patients(38). Reduced expression of *ECT2* is associated with abrogation of

MAPK signaling(39). Finally, several key regulators of cell cycle progression associated with

higher grade, worse outcomes, and poor response to endocrine therapy in ER+ breast cancer are

commonly downregulated including *CCND1*, *TOP2A*, and *CCNB1*.(40-43)

To test if signatures of vandetanib regulated genes are associated with prognosis in

patients with ER+/HER2- breast cancer treated with endocrine therapy, we obtained survival

annotated transcriptional data from METABRIC(44) and SCAN-B(45). Tumors were stratified

by median gene set score using median thresholding. METABRIC patients with low expression

of genes that were downregulated by vandetanib had significantly improved OS (HR=0.75,

p<0.0001) and RFS (HR=0.60, p<0.0001). Similarly, SCAN-B ER+ breast cancer patients with

low expression of vandetanib downregulated genes demonstrated significantly lower mortality

compared to high signature score (HR=0.87, p=0.034), Fig. 2F. These findings support that

downregulation of genes that are repressed by vnadetanib is associated with better outcomes

across datasets. To determine whether vandetanib treatment is reducing expression of genes that

drive an ET- resistant phenotype, we obtained gene expression data from patients with

ER+/HER2- breast cancer profiled on ET and clinically annotated as sensitive or resistant(13).

Resistant tumors had significantly higher expression of the vandetanib downregulated signature

compared to sensitive tumors (4.3 vs 3.8, p<0.001), Fig. 2G. Cumulatively, these findings

support that in ER+ breast cancer, vandetanib treatment promotes gene expression changes

associated with lower proliferation, increased ET response, and improved survival.

Functional Signatures of Vandetanib Response

Vandetanib alters kinase signaling which is regulated at the protein level. We used

multiplex inhibitor bead enrichment coupled to mass spectrometry (MIB/MS) to investigate

reprogramming of kinase signaling networks in response to vandatinib (31). In this technique,

kinases are enriched by incubating cell extracts with type 1 kinase inhibitors immobilized

on Sepharose beads. Each cell line was treated with the IC90 of vandetanib or vehicle for 48

hours, Fig. 3A. We first investigated the effects of vandetanib on enrichment and depletion of

known vandetanib targets(46). RET, EGFR, ABL1, RIP2K, PTK6, and DDR1 had similar

magnitude of depletion in all 3 models, despite large differences in baseline enrichment between

MCF7 and the ET-resistant derivatives, Fig. 3B. Similar results were seen for T47D and T47D-

TAM (Supplementary Fig. 3). These results demonstrate that several distinct target proteins may

be mediating vandetanib effects in ER+ breast cancer models, that effects of treatment do not

appear to be proportional to baseline expression, and that the relative level of expression of a

target alone may not be a good biomarker of response.

Enriched and depleted kinase proteins by cell line are shown in Fig. 3C. Overlap analysis

(Fig. 3D) showed that similar to transcriptional findings, kinome dynamics were heterogenous

across cell lines. There were 12 common upregulated and 22 downregulated kinase proteins.

Common upregulated genes included AURKA, PLK1, PIP4K2A, which promote proliferation

and aggressive tumors(47-49), as well as JAK1 and TYK2 which are associated with immune

infiltrates and are upstream of AKT signaling(50, 51). Common depleted proteins with

vandetanib treatment included proliferation promoting cell cycle regulators CDK1, CKD2, and

PKMYT1. Depletion of MAP2K2 (MEK2) as well as upstream receptor tyrosine kinases

EPHA1, EPHB4, DDR1, and EGFR suggest downregulation of the MAPK signaling pathway in

response to vandetanib treatment consistent with Western blots (Fig. 1A). Gene set enrichment

analysis demonstrated a common upregulation of the PI3K/AKT pathway and downregulation of

the ERK cascade including transciprtional and phosphoproteomic signatures (52), Fig. 3E.

Pharmacologic Inhibition of Adaptive Resistance Kinases Enhances Vandetanib Efficacy

Upregulation of functional kinases may identify common mechanisms of adaptive bypass

resistance so activity levels of druggable kinases(24) were assessed by treatment (Fig. 4A). To

determine whether enriched druggable targets may be mediating adaptive resistance to

vandetanib we performed co-treatment assays on the principle that inhibiting mediators of

adaptive resistance should enhance sensitivity to vandetanib. Because ERBB2/HER2 was

enriched in all 3 MCF7 cell lines, we co-treated cells with vandetanib and the FDA approved

HER2 small molecule inhibitor lapatinib. Treatment with lapatinib resulted in significantly

augmented vandetanib reduction in cell viability in all 3 cell lines, (Fig. 4B), demonstrating the

phenotypic significance of upregulated protein activity.

RET Mediates Vandetanib Sensitivity

Functional kinases that are downregulated after vandetanib treatment may be mediating

the effects of vandetanib. Based on our prior work showing a role for RET in mediating

vandetanib response and in driving poor outcomes and ET resistance in ER+ breast cancer we

explored the role of RET signaling in vandetanib sensitivity. As expected from the MIB/MS

data, transient RET antagonism with the selective RET inhibitor BLU667 resulted in reduced

sensitivity to vandetanib in MCF7 in dose dependent fashion. BLU667 also reduced vandetanib

sensitivity in MCF7-TAM, but not in the low RET expressing MCF7-EXE (Fig. 5A). These

findings demonstrate that baseline activity, as well redcuced activity with treatment, are

important for mediators of response.

We next sought to clearly elucidate the role of RET In ER+ breast cancer cells by

generating a CRISPR knockout in MCF7 cells (RET KO). Western blot confirmed knockout of

RET protein (Fig. 5B). We used our CRISPR KO model to determine the impact of RET KO on

sensitivity to vandetanib. Supporting a role for RET as a mediator of vandetanib response, RET

KO cells were significantly less sensitive to vandetanib compared to WT (Fig. 5C). Based on the

vandetanib synergy assay we hypothesized that RET KO would sensitize cells to tamoxifen, but

unexpectedly observed that KO cells were less sensitive (Fig. 5D).

Because decreased sensitivity to tamoxifen with RET KO was observed we next sought

to explore the molecular basis for this unexpected finding by performing mRNA sequencing and

comparing RET KO and WT cells. We assessed ESR1 (encoding ER) expression in the mRNA

data and found that RET KO cells had reduced expression of ESR1 (log2FC -1.27, padj <0.001),

Fig. 5E. Reduced ER expression could account for reduced sensitivity to tamoxifen in cells that

have stable KD of RET. We next performed unbiased analysis of differentially regulated genes

with RET KO (Fig. 5F). Hallmark gene sets for estrogen response and KRAS signaling were

depleted with RET KO, as were genes previously associated with RET knockout. RET KO

resulted in lower proliferation score compared to WT. Supportive reduced proliferative

signature, RET KO cells had significantly reduced growth at 48 and 72 hours relative to WT

cells (Fig. 5G). Cumulatively these findings support that RET is a mediator of vandetanib

response and drives a proliferative phenotype, but that complex RET-ER interactions may evolve

over time, and this may impact sensitivity to endocrine therapy.

Vandetanib Response at Single Cell Resolution in Primary ER+ Breast Cancer

We used our previously validated *ex vivo* short term drug treatment platform coupled to

single cell RNA sequencing(33) to assess how primary breast tumor cells respond to vandetanib

treatment. Specifically, we obtained a primary ER+/HER2- breast cancer, dissociated into single

cells, treated with control of vandetanib containing media for 12 hours and assayed by 10x

scRNAseq. Uniform manifold projection (UMAP) was performed using Seurat v3(53). Cell

populations were annotated using known transcriptional profiles and canonical markers (Fig. 6A)

and by treatment group (Fig. 6B). Malignant (tumor) cells were distinguished from non-tumor

epithelial cells as cells having significant inferred copy number alterations by InferCNV(54).

Differentially expressed genes were identified using a false discovery rate (FDR) cutoff of <

0.05 in the iCNV+ tumor cell compartment with vandetanib treatment compared to control,

Fig. 6C. Enriched and depleted pathways in vandetanib treated tumor cells relative to control

treated were determined, Fig. 6D. Of note, as seen in cell lines, ESR1 targets and ERBB2

pathway signaling were enriched in vandetanib treated cells (Fig. 6E).

We tested the overlapping downregulated genes after vandetanib treatment across the 3

MCF7 cell lines (Fig. 3) and these genes were also depleted in vandetanib treated primary tumor

cells (Fig. 6E). Next LumA correlation score was assigned to each tumor cell using a validated

single cell PAM50 classifier(55). Treated tumor cells had a stronger correlation to the LumA

centroid (median score treated 0.259 vs median score untreated 0.244, p= 0.001, Wilcoxon rank-

sum test). Thus, as with cell lines, short term vandetanib treatment alters the transcriptional

profile of ER+/HER2- primary tumor cells towards a more indolent and estrogen responsive

phenotype.

To explore features of vandetanib response within primary tumor cells we used non-

negative matrix factorization (NMF) to elucidate features that were enriched and depleted in

vandetanib treated cells (Supplementary Fig. 4A). This analysis yielded metagene modules that

were enriched in vandetanib treated samples (module 1) and depleted in vandetanib treated cells

(Module 6) (Supplementary Fig. 4B). We next identified features that may be associated with

vandetanib response by comparing module 1 and 6 gene expression profiles. Module 1

(vandetanib resistant) had expected high expression of genes that increase with vandetanib

treatment, consistent with reduced response. Module 6 was enriched for MYC targets, which is known to support tamoxifen resistance, but may support a transcriptional profile of vandetanib sensitivity (Supplementary Fig. 4C). Module 1 was also enriched for MTORC signaling which we identified in cells lines as an adaptive vandetanib resistance mechanism. Interestingly, module 1 was enriched for hallmark estrogen response genes and our previously generated single cell tamoxifen response score(33) supporting that cells with poor vandetanib response have enhanced tamoxifen sensitivity (Supplementary Fig. 4D). Lastly, we looked at our human tumor derived signatures of high GDNF/RET expressing human ER+-breast tumors(23). Genes that were depleted in GDNF/RET high human tumors were depleted in vandetanib sensitive cells, supporting that signature as a biomarker of response (Supplementary Fig. 4E).

DISCUSSION

Resistance to endocrine therapy remains a common problem and a significant driver of

poor patient outcomes despite advances in understanding of how ER+ breast tumors respond to

therapy. The identification of activation in MAPK pathways as a driver of resistance to ET has

generated significant interest in targeting this pathway to enhance ET response. However,

clinical trials have demonstrated inconsistent and disappointing results regarding response to

tyrosine kinase inhibitors for non HER2-amplified tumors. These findings demonstrate the need

for improved understanding of how intracellular signaling pathways are altered by treatment to

determine biomarkers of patients most likely to respond and clinically actionable kinase targeting

strategies in MAPK-activated ET resistant breast tumors.

We found that response to vandetanib, measured by IC50 or by depletion of RET activity,

was largely the same across the cell lines tested, despite large differences in baseline RET

expression, and this was also true for the other direct vandetanib targets. These findings

demonstrate that single protein expression may not be an effective biomarker of response and

signatures of activity may be more informative of response than single gene expression. This

concept has been shown in HER2 targeting therapies where expression of the HER2E gene

expression pattern is more predictive of response than HER2 expression alone(56). We recently

generated a signature of RET active ER+ breast tumors(23) using treatment naïve samples from

The Cancer Genome Atlas (TCGA), and that low expression of genes that were depleted in

GDNF/RET high tumors was associated with vandetanib sensitivity, suggesting that this may be

a biomarker to select patients most likely to respond.

RET partially mediated effects of vandetanib, but stable knockout of RET unexpectedly

resulted in reduced, rather than augmented, sensitivity to tamoxifen. ESR1 expression was found

to be reduced after RET knockout and this could account for decreased sensitivity to tamoxifen.

While ER has 2 binding sites in the RET promoter that activate RET expression (57), no

reciprocal mechanism by which RET regulates ER is known. Prior work has shown that TKI

treatment reprograms transcriptional networks in HER2-amplified (58) and triple negative breast

cancer(32) and similar remodeling over time may reduce ESR1 expression with loss of RET

activity. Alternatively, because RET expression and ER-activity are correlated, RET KO may

select for cells that are less dependent on RET signaling, which may also be cells that have lower

baseline ER-activity. This finding could help explain for negative clinical trials of RET inhibitors

with anti-ER therapy if tumor cell subpopulations that evolve resistance to RET inhibitors are

also less dependent on ER-activty, and therfore less responsive to ET. Further study is needed to

model emerging resistance to RET and ER inhibitors and track relative dependencies on RET

and ER within emerging resistant tumor cell subpopulations.

Kinase signaling networks are plastic and inhibiting one node can lead to adaptive bypass

through another pathway. We were able to identify mediators of adaptive resistance to

vandetanib treatment and augment response with dual treatment. Up-regulation of HER2 activity

was common to all cell lines and primary tumor cells and targeting HER2 with lapatinib

enhanced sensitivity to vandetanib. Activating HER2 signaling could open a number of

efficacious therapeutic options for patients. While HER2 activity was increased, there was no

change in HER2 RNA expression, and this highlights the need for assessment at the level of

protein activity to determine adaptive bypass mechanisms.

While HER2 upregulation was a common adaptive bypass pathway, we also identified

cell line specific enriched proteins. This could be a result of distinct adaptations necessary to

resist exemestane compared to tamoxifen or could represent distinct alternative paths to resist

vandetanib treatment. How these adaptive mechanisms are conserved or unique in human tumors

will be essential to understand the therapeutic implications of these co-treatment targets. Specific

adaptive bypass pathways may be predictable from pretreatment gene expression or chromatin

accessibility states, or tumor level assessment of response to therapy may be needed to identify

tumor specific bypass mechanisms and cotreatment vulnerabilities. To this end, our group has

recently developed a method for whole kinome characterization on limited sample (such as a

core needle biopsy), which could make on treatment assessment of adaptive response and

targeted co-treatment strategies feasible.(59)

In summary, we demonstrate that ER+ breast cancer cells become more sensitive to ET

with vandetanib treatment. Vandetanib results in broad reprogramming of gene expression and

kinase signaling which results in a lower proliferative, more estrogen responsive luminal A-like

state. Early adaptive response, including upregulation of HER2, to vandetanib identified co-

treatment strategies to enhance response. Future trials will need to utilize signature-based

biomarkers of response with attention to tumor specific upregulated adaptive bypass pathways as

cotreatment strategies to improve efficacy.

ACKNOWLEDGEMENTS

Dr Philip Spanheimer is supported by the National Institutes of Health grants R37CA292075, K08CA280388, and a Solid Tumor Research Scholars Grant from Gilead Sciences. CMP was supported by funds from the NCI breast SPORE program (P50-CA058223). This work was additionally supported in part by P30CA016086 UNC Lineberger Comprehensive Cancer Center Core Support Grant.

FIGURES

489
Figure 1. (A) Treatment of MCF-7 and tamoxifen resistant MCF7 (MCF7-TAM) and exemestane resistant MCF7
(MCF7-EXE) reduces pERK/ERK in all 3 cell lines. (B) Vandetanib dose dependent reduction in cell viablilty in
MCF7, MCF7-TAM, and MCF7-EXE. (C) Treatment with Vandetanib results in a left shift of the tamoxifen
sensitivity curve and reduction in IC50 in all 3 cell lines. (D) Synergy plots demonstrate synergistic activity of
tamoxifen and vandetanib.*p<0.05, ***p<0.01 by t-test with Welch correction and analysis of the slope difference
in the nonlinear regression model.

496
Figure 2. Heatmap and enrichment analysis of differentially expressed genes in (A) MCF7, (B) MCF7-TAM, (C)
MCF7-EXE. KRAS activity signatures were depleted in all 3 cell lines by vandetanib treatment. (D) Vandetanib
treatment increases association with the more indolent, endocrine therapy responsive Luminal A subtype, measured
by correlation to Luminal A defining genes (centroid). (E) Overlap of up- and down- regulated genes by cell line.
(F) Signatures of overlapping up-and down regulated genes are associated with recurrence free and overall survival
in ER+ breast cancer patients in METARIC and SCAN-B studies and (G) downregulated genes are enriched in
clinically ET-resistant ER+ tumors. *p<0.05, **p<0.01, ***p<0.001, ****p<0.0001 by t-test with Welch correction
for Wilcoxon test.

505
Figure 3. (A) Multiplex inhibitor bead mass spectrometry assay work flow. (B) Effects of vandetanib on enrichment
and depletion of known vandetanib targets. Baseline expression of vandetanib targets in tamoxifen and exemestane
resistant cell lines compared to WT MCF7. (C) Heatmap showing enriched and depleted proteins after vandetanib
treatment. (D) Overlap analysis of up- and downregulated proteins between the 3 cell lines when treated with
vandetanib. (E) Pathway analysis of the downregulation of the ERK cascade and upregulation of the PI3K/AKT
pathway.

Figure 4. (A) Heatmap showing enrichment levels of druggable kinases of sensitive and resistant MCF7 cell lines before and after vandetanib treatment. Of note, ERBB2 (encoding HER2) was enriched in all 3 cell lines after treatment and RET was dpeleted in MCF7 and MCF7-TAM. (B) Sensitivity plots of co-treated cells with vandetanib and HER2 inhibitor lapatinib demonstrating enhanced lapatinib sensitivity with vandetanib treatment. *p<0.05, ****p<0.0001 by t-test with Welch correction.

519
Figure 5. (A) Co-treatment with vandetanib and the selective RET inhibitor BLU667 reduced vandetanib
sensitivitiy in MCF7 and MCF7-TAM, but not MCF7-EXE with RET inhibition consistent with activity depletion in
Fig4. (B) Western blot demonstrating CRISPR knockout of RET resulting in >90% reduction in RET protein levels.
(C) RET knockout cells have reduced sensitivity to vandetanib. (D) RET knockout cells have reduced sensitivity to
tamoxifen. (E) ESR1 RNA levels encoding the estrogen receptor are decreased in RET KO cells, consistent with
reduced sensitivity to tamoxifen. (F) Heatmap demonstrating differentially expressed genes with RET KO in MCF7.
Enrichment analysis demonstrates reduction in hallmark estrogen response and RAS signaling in RET KO cells.
RET KO cells have lower proliferation score. (G) RET knockout cells have reduced proliferation compared to WT.

(. *p<0.05, **p<0.01, ***p<0.001, ****p<0.0001 by t-test with Welch correction for the bar graphs, and Wilcoxon
test.

530
Figure 6. (A) Single-cell RNA sequencing uniform manifold projection (UMAP) annotated by cell type within
a primary ER+ human breast tumor. (B) Tumor cells were computationally isolated as having significant
inferred copy number changes and plotted by treatment condition. (C) Heatmap showing differentially
expressed genes compared between vandetanib treated vs non-treated tumor cells. (D) Enriched and depleted
pathways of vandetanib treated vs non-treated tumor cells. (E) ERBB2 activity signature score was increased
in vandetanib treated tumor cells as seen in cell lines. Down regulated genes in vandetanib treated cell lines
are depleted in vandetanib treated primary tumor cells, and LumA correlation increased in vandetanib treated
tumor cells as was seen in cell lines, and depleted KRAS activity signature demonstrates on target effects.
Significant signatures filtered by < 0.1 FDR on GSEA analysis and significant genes filtered by adjusted p
value <0.05 (Wilcoxon test).

REFERENCES

1. Nabholtz JM, Tonkin K, Smylie M, Au HJ, Lindsay MA, and Mackey J. Chemotherapy of breast cancer: are the taxanes going to change the natural history of breast cancer? *Expert Opin Pharmacother.* 2000;1(2):187-206.

2. Mouridsen H, Gershanovich M, Sun Y, Perez-Carrion R, Boni C, Monnier A, et al. Phase III study of letrozole versus tamoxifen as first-line therapy of advanced breast cancer in postmenopausal women: analysis of survival and update of efficacy from the International Letrozole Breast Cancer Group. *J Clin Oncol.* 2003;21(11):2101-9.

3. Bachelot T, Bourgier C, Cropet C, Ray-Coquard I, Ferrero JM, Freyer G, et al. Randomized phase II trial of everolimus in combination with tamoxifen in patients with hormone receptor-positive, human epidermal growth factor receptor 2-negative metastatic breast cancer with prior exposure to aromatase inhibitors: a GINECO study. *J Clin Oncol.* 2012;30(22):2718-24.

4. Giaquinto AN, Sung H, Newman LA, Freedman RA, Smith RA, Star J, et al. Breast cancer statistics 2024. *CA Cancer J Clin.* 2024.

5. Harvey JM, Clark GM, Osborne CK, and Allred DC. Estrogen receptor status by immunohistochemistry is superior to the ligand-binding assay for predicting response to adjuvant endocrine therapy in breast cancer. *J Clin Oncol.* 1999;17(5):1474-81.

6. Jeselsohn R, Barry WT, Migliaccio I, Biagioni C, Zhao J, De Tribolet-Hardy J, et al. TransCONFIRM: Identification of a Genetic Signature of Response to Fulvestrant in Advanced Hormone Receptor-Positive Breast Cancer. *Clin Cancer Res.* 2016;22(23):5755-64.

7. Finn RS, Martin M, Rugo HS, Jones S, Im SA, Gelmon K, et al. Palbociclib and Letrozole in Advanced Breast Cancer. *N Engl J Med.* 2016;375(20):1925-36.

8. Baselga J, Campone M, Piccart M, Burris HA, 3rd, Rugo HS, Sahmoud T, et al. Everolimus in postmenopausal hormone-receptor-positive advanced breast cancer. *N Engl J Med.* 2012;366(6):520-9.

9. Piccart M, Hortobagyi GN, Campone M, Pritchard KI, Lebrun F, Ito Y, et al. Everolimus plus exemestane for hormone-receptor-positive, human epidermal growth factor receptor-2-negative advanced breast cancer: overall survival results from BOLERO-2dagger. *Ann Oncol.* 2014;25(12):2357-62.

10. Andre F, Ciruelos E, Rubovszky G, Campone M, Loibl S, Rugo HS, et al. Alpelisib for PIK3CA-Mutated, Hormone Receptor-Positive Advanced Breast Cancer. *N Engl J Med.* 2019;380(20):1929-40.

11. Turner NC, Slamon DJ, Ro J, Bondarenko I, Im SA, Masuda N, et al. Overall Survival with Palbociclib and Fulvestrant in Advanced Breast Cancer. *New Engl J Med.* 2018;379(20):1926-36.

12. Razavi P, Chang MT, Xu G, Bandlamudi C, Ross DS, Vasan N, et al. The Genomic Landscape of Endocrine-Resistant Advanced Breast Cancers. *Cancer Cell.* 2018;34(3):427-38 e6.

13.   Xia Y, He X, Renshaw L, Martinez-Perez C, Kay C, Gray M, et al. Integrated DNA and
RNA Sequencing Reveals Drivers of Endocrine Resistance in Estrogen Receptor-Positive
Breast Cancer. *Clin Cancer Res.* 2022;28(16):3618-29.
14.   Arpino G, Wiechmann L, Osborne CK, and Schiff R. Crosstalk between the estrogen
receptor and the HER tyrosine kinase receptor family: molecular mechanism and clinical
implications for endocrine therapy resistance. *Endocr Rev.* 2008;29(2):217-33.
15.   Meijer D, Sieuwerts AM, Look MP, van Agthoven T, Foekens JA, and Dorssers LC.
Fibroblast growth factor receptor 4 predicts failure on tamoxifen therapy in patients with
recurrent breast cancer. *Endocr Relat Cancer.* 2008;15(1):101-11.
16.   Bergqvist J, Elmberger G, Ohd J, Linderholm B, Bjohle J, Hellborg H, et al. Activated
ERK1/2 and phosphorylated oestrogen receptor alpha are associated with improved
breast cancer survival in women treated with tamoxifen. *Eur J Cancer.* 2006;42(8):1104-
12.
17.   Kirkegaard T, Witton CJ, McGlynn LM, Tovey SM, Dunne B, Lyon A, et al. AKT
activation predicts outcome in breast cancer patients treated with tamoxifen. *J Pathol.*
2005;207(2):139-46.
18.   Shoman N, Klassen S, McFadden A, Bickis MG, Torlakovic E, and Chibbar R. Reduced
PTEN expression predicts relapse in patients with breast carcinoma treated by tamoxifen.
Mod Pathol. 2005;18(2):250-9.
19.   Morandi A, Martin LA, Gao Q, Pancholi S, Mackay A, Robertson D, et al. GDNF-RET
signaling in ER-positive breast cancers is a key determinant of response and resistance to
aromatase inhibitors. *Cancer Res.* 2013;73(12):3783-95.
20.   Boulay A, Breuleux M, Stephan C, Fux C, Brisken C, Fiche M, et al. The Ret receptor
tyrosine kinase pathway functionally interacts with the ERalpha pathway in breast cancer.
Cancer Res. 2008;68(10):3743-51.
21.   Plaza-Menacho I, Morandi A, Robertson D, Pancholi S, Drury S, Dowsett M, et al.
Targeting the receptor tyrosine kinase RET sensitizes breast cancer cells to tamoxifen
treatment and reveals a role for RET in endocrine resistance. *Oncogene.*
2010;29(33):4648-57.
22.   Spanheimer PM, Woodfield GW, Cyr AR, Kulak MV, White-Baer LS, Bair TB, et al.
Expression of the RET proto-oncogene is regulated by TFAP2C in breast cancer
independent of the estrogen receptor. *Ann Surg Oncol.* 2013;20(7):2204-12.
23.   Kakati RT, Kim H, Whitman A, and Spanheimer PM. High expression of the RET
receptor tyrosine kinase and its ligand GDNF identifies a high-risk subset of estrogen
receptor positive breast cancer. *Breast Cancer Res Treat.* 2023.
24.   Gou X, Kim BJ, Anurag M, Lei JT, Young MN, Holt MV, et al. Kinome reprogramming
is a targetable vulnerability in ESR1 fusion-driven breast cancer. *Cancer Res.* 2023.
25.   Spanheimer PM, Cyr AR, Gillum MP, Woodfield GW, Askeland RW, and Weigel RJ.
Distinct pathways regulated by RET and estrogen receptor in luminal breast cancer
demonstrate the biological basis for combination therapy. *Ann Surg.* 2014;259(4):793-9.

26. Spanheimer PM, Park JM, Askeland RW, Kulak MV, Woodfield GW, De Andrade JP, et
al. Inhibition of RET increases the efficacy of antiestrogen and is a novel treatment
strategy for luminal breast cancer. *Clin Cancer Res.* 2014;20(8):2115-25.
27. Spanheimer PM, Lorenzen AW, De Andrade JP, Kulak MV, Carr JC, Woodfield GW, et
al. Receptor Tyrosine Kinase Expression Predicts Response to Sunitinib in Breast
Cancer. *Ann Surg Oncol.* 2015;22(13):4287-94.
28. Miller KD, Trigo JM, Wheeler C, Barge A, Rowbottom J, Sledge G, et al. A multicenter
phase II trial of ZD6474, a vascular endothelial growth factor receptor-2 and epidermal
growth factor receptor tyrosine kinase inhibitor, in patients with previously treated
metastatic breast cancer. *Clin Cancer Res.* 2005;11(9):3369-76.
29. Xu J, Higgins MJ, Tolaney SM, Come SE, Smith MR, Fornier M, et al. A Phase II Trial
of Cabozantinib in Hormone Receptor-Positive Breast Cancer with Bone Metastases.
Oncologist. 2020.
30. Lim JSJ, Wong ALA, Ow SGW, Ngoi NYL, Chan GHJ, Ang YLE, et al. Phase Ib/II dose
expansion study of lenvatinib combined with letrozole in post-menopausal women with
hormone receptor positive breast cancer. *Clin Cancer Res.* 2022.
31. Duncan JS, Whittle MC, Nakamura K, Abell AN, Midland AA, Zawistowski JS, et al.
Dynamic reprogramming of the kinome in response to targeted MEK inhibition in triple-
negative breast cancer. *Cell.* 2012;149(2):307-21.
32. Zawistowski JS, Bevill SM, Goulet DR, Stuhlmiller TJ, Beltran AS, Olivares-Quintero
JF, et al. Enhancer Remodeling during Adaptive Bypass to MEK Inhibition Is Attenuated
by Pharmacologic Targeting of the P-TEFb Complex. *Cancer Discov.* 2017;7(3):302-21.
33. Kim H, Whitman AA, Wisniewska K, Kakati RT, Garcia-Recio S, Calhoun BC, et al.
Tamoxifen Response at Single Cell Resolution in Estrogen Receptor-Positive Primary
Human Breast Tumors. *Clin Cancer Res.* 2023.
34. Klomp JA, Klomp JE, Stalnecker CA, Bryant KL, Edwards AC, Drizyte-Miller K, et al.
Defining the KRAS- and ERK-dependent transcriptome in KRAS-mutant cancers.
Science. 2024;384(6700):eadk0775.
35. Zhang S, Li F, Younes M, Liu H, Chen C, and Yao Q. Reduced selenium-binding protein
1 in breast cancer correlates with poor survival and resistance to the anti-proliferative
effects of selenium. *PLoS One.* 2013;8(5):e63702.
36. Chen R, Wang X, Fu J, Liang M, and Xia T. High FLT3 expression indicates favorable
prognosis and correlates with clinicopathological parameters and immune infiltration in
breast cancer. *Front Genet.* 2022;13:956869.
37. Papadakis ES, Reeves T, Robson NH, Maishman T, Packham G, and Cutress RI. BAG-1
as a biomarker in early breast cancer prognosis: a systematic review with meta-analyses.
Br J Cancer. 2017;116(12):1585-94.
38. Kahl I, Mense J, Finke C, Boller AL, Lorber C, Gyorffy B, et al. The cell cycle-related
genes RHAMM, AURKA, TPX2, PLK1, and PLK4 are associated with the poor
prognosis of breast cancer patients. *J Cell Biochem.* 2022;123(3):581-600.

39. Yi M, Zhang D, Song B, Zhao B, Niu M, Wu Y, et al. Increased expression of ECT2
predicts the poor prognosis of breast cancer patients. *Exp Hematol Oncol.*
2022;11(1):107.
40. Lundberg A, Lindstrom LS, Li J, Harrell JC, Darai-Ramqvist E, Sifakis EG, et al. The
long-term prognostic and predictive capacity of cyclin D1 gene amplification in 2305
breast tumours. *Breast Cancer Res.* 2019;21(1):34.
41. Montalto FI, and De Amicis F. Cyclin D1 in Cancer: A Molecular Connection for Cell
Cycle Control, Adhesion and Invasion in Tumor and Stroma. *Cells.* 2020;9(12).
42. An X, Xu F, Luo R, Zheng Q, Lu J, Yang Y, et al. The prognostic significance of
topoisomerase II alpha protein in early stage luminal breast cancer. *BMC Cancer.*
2018;18(1):331.
43. Ding K, Li W, Zou Z, Zou X, and Wang C. CCNB1 is a prognostic biomarker for ER+
breast cancer. *Med Hypotheses.* 2014;83(3):359-64.
44. Curtis C, Shah SP, Chin SF, Turashvili G, Rueda OM, Dunning MJ, et al. The genomic
and transcriptomic architecture of 2,000 breast tumours reveals novel subgroups. *Nature.*
2012;486(7403):346-52.
45. Saal LH, Vallon-Christersson J, Hakkinen J, Hegardt C, Grabau D, Winter C, et al. The
Sweden Cancerome Analysis Network - Breast (SCAN-B) Initiative: a large-scale
multicenter infrastructure towards implementation of breast cancer genomic analyses in
the clinical routine. *Genome Med.* 2015;7(1):20.
46. Klaeger S, Heinzlmeir S, Wilhelm M, Polzer H, Vick B, Koenig PA, et al. The target
landscape of clinical kinase drugs. *Science.* 2017;358(6367).
47. Zhu L, Lu Z, and Zhao H. Antitumor mechanisms when pRb and p53 are genetically
inactivated. *Oncogene.* 2015;34(35):4547-57.
48. Emerling BM, Hurov JB, Poulogiannis G, Tsukazawa KS, Choo-Wing R, Wulf GM, et
al. Depletion of a putatively druggable class of phosphatidylinositol kinases inhibits
growth of p53-null tumors. *Cell.* 2013;155(4):844-57.
49. O'Connell CE, and Vassilev A. Combined Inhibition of p38MAPK and PIKfyve
Synergistically Disrupts Autophagy to Selectively Target Cancer Cells. *Cancer Res.*
2021;81(11):2903-17.
50. Zhang Q, Sturgill JL, Kmieciak M, Szczepanek K, Derecka M, Koebel C, et al. The role
of Tyk2 in regulation of breast cancer growth. *J Interferon Cytokine Res.* 2011;31(9):671-
7.
51. Chen B, Lai J, Dai D, Chen R, Li X, and Liao N. JAK1 as a prognostic marker and its
correlation with immune infiltrates in breast cancer. *Aging (Albany NY).*
2019;11(23):11124-35.
52. Klomp JE, Diehl JN, Klomp JA, Edwards AC, Yang R, Morales AJ, et al. Determining
the ERK-regulated phosphoproteome driving KRAS-mutant cancer. *Science.*
2024;384(6700):eadk0850.
53. Stuart T, Butler A, Hoffman P, Hafemeister C, Papalexi E, Mauck WM, 3rd, et al.
Comprehensive Integration of Single-Cell Data. *Cell.* 2019;177(7):1888-902 e21.

54.   Puram SV, Tirosh I, Parikh AS, Patel AP, Yizhak K, Gillespie S, et al. Single-Cell
704   Transcriptomic Analysis of Primary and Metastatic Tumor Ecosystems in Head and Neck
705   Cancer. *Cell.* 2017;171(7):1611-24 e24.
706   55.   Wu SZ, Al-Eryani G, Roden DL, Junankar S, Harvey K, Andersson A, et al. A single-cell
707   and spatially resolved atlas of human breast cancers. *Nat Genet.* 2021;53(9):1334-47.
708   56.   Tanioka M, Fan C, Parker JS, Hoadley KA, Hu Z, Li Y, et al. Integrated Analysis of
709   RNA and DNA from the Phase III Trial CALGB 40601 Identifies Predictors of Response
710   to Trastuzumab-Based Neoadjuvant Chemotherapy in HER2-Positive Breast Cancer. *Clin*
711   *Cancer Res.* 2018;24(21):5292-304.
712   57.   Stine ZE, McGaughey DM, Bessling SL, Li S, and McCallion AS. Steroid hormone
713   modulation of RET through two estrogen responsive enhancers in breast cancer. *Hum*
714   *Mol Genet.* 2011;20(19):3746-56.
715   58.   Angus SP, Stuhlmiller TJ, Mehta G, Bevill SM, Goulet DR, Olivares-Quintero JF, et al.
716   FOXA1 and adaptive response determinants to HER2 targeted therapy in TBCRC 036.
717   *NPJ Breast Cancer.* 2021;7(1):51.
718   59.   East MP, Sprung RW, Okumu DO, Olivares-Quintero JF, Joisa CU, Chen X, et al.
719   Quantitative proteomic mass spectrometry of protein kinases to determine dynamic
720   heterogeneity of the human kinome. *bioRxiv.* 2024.