Kinase Plasticity in Response to Vandetanib Enhances Sensitivity to Tamoxifen and Identifies
Co-Treatment Strategies in Estrogen Receptor Positive Breast Cancer

Rasha T. Kakati MD[1], Austin A. Whitman MS[1], Santiago Haase PhD[1], Attila T. Szenasi PhD[1],
Christine HNC Thai BS[1,] Elizabeth Brunk PhD[1,2], Denis O. Okumu PhD[2], Michael P. East PhD[2],
Charles M. Perou PhD[1,3,4,5], Gary L. Johnson, PhD[1,2], and Philip M. Spanheimer MD[1,4,6]

[1]Lineberger Comprehensive Cancer Center, University of North Carolina, Chapel Hill, NC
[2]Department of Pharmacology, University of North Carolina, Chapel Hill, NC
[3]Department of Pathology and Laboratory Medicine, University of North Carolina at Chapel
Hill, Chapel Hill, NC
[4]Department of Genetics, University of North Carolina, Chapel Hill, NC
[5]Computational Medicine Program, University of North Carolina, Chapel Hill, NC
[6]Department of Surgery, University of North Carolina, Chapel Hill, NC

Running title: Vandetanib ER+ Breast Cancer

Keywords: Breast cancer, endocrine resistance, tyrosine kinase inhibition, functional proteomics,
targeted therapeutics

Financial Support: This work was supported in part by the National Institutes of Health grants
K08CA280388, R37CA292075, P30CA016086, and a Solid Tumor Research Scholars Grant
from Gilead Sciences. CMP was supported by funds from the NCI breast SPORE program (P50-
CA058223).

Correspondence should be addressed to:
Philip M. Spanheimer MD
Lineberger Comprehensive Cancer Center
University of North Carolina at Chapel Hill
170 Manning Drive, Suite 1149
Chapel Hill, NC 27599-7213 USA
Phone: 919-966-5221; Fax: 919-966-8806
E-mail: Philip_Spanheimer@med.unc.edu

Disclosure of conflict of interests: C.M.P is an equity stockholder and consultant of BioClassifier
LLC; C.M.P is also listed as an inventor on patent applications for the Breast PAM50 Subtyping
assay. The remaining authors declare no potential conflicts of interest.

ABSTRACT

Resistance to endocrine therapy (ET) is common in estrogen receptor (ER) positive breast cancer. Multiple studies have demonstrated that upregulation of MAPK signaling pathways contributes to ET resistance. Herein we show that vandetanib treatment enhances sensitivity to ET in ET-sensitive and -resistant ER+ breast cancer models. Using a CRISPR knockout model, we demonstrate that vandetanib effects are partially mediated by RET receptor tyrosine kinase. Vandetanib treatment alters the gene expression program of ER+ breast cancer cells resulting in a less proliferative and more estrogen responsive Luminal-A like character. Tyrosine kinase network reprogramming was assessed using multiplexed kinase inhibitor beads-mass spectrometry (MIB/MS) assay to identify adaptive resistance mechanisms to vandetanib treatment, which identified upregulation of HER2 activity. Co-treatment to inhibit HER2 with lapatinib enhanced sensitivity to vandetanib, demonstrating biologic activity of HER2 upregulation. Finally, we use our operating room-to-laboratory assay that measures drug response in individual primary tumor cells in short term cultures to demonstrate conserved gene expression changes, including increased HER2 activity signatures, in vandetanib treated cells, and identify features associated with vandetanib response. These results support future investigation of RET targeting strategies considering reprogrammed networks, such as activated HER2, in patients with ET resistant ER+ breast cancer.

INTRODUCTION

Approximately 75% of breast cancers fall within the estrogen receptor (ER)+/luminal subtypes[1-3]. About 42,250 women are expected to die from breast cancer in 2024 and most of these will have ER+ disease[4]. Most patients with ER+ breast cancer will be treated with endocrine therapy (ET) targeting ER. However, resistance to ET occurs in about 1/3 of early cancers and develops over time in all metastatic ER+ cancers[5-7]. Targeted inhibitors to PI3K, mTOR, and CDK4/6 have shown efficacy in patients with ET resistant disease[8-11]. However, responses are sometimes short lived and there are a significant number of patients in need of durable, effective therapy.

Multiple studies have demonstrated that activation of ERK/MAPK pathways is an important mechanism by which ER+ breast tumors resist ET[12-18]. Gain of function mutations in ERK/MAPK nodes or upstream receptor tyrosine kinase (RTK) are seen in ET-resistant tumors, although these are only present in about 13% of cases[12]. The rearranged during transfection (RET) receptor tyrosine kinase is expressed in approximately 50% of ER+ breast tumors and is more highly expressed in ET resistant tumors[19-22]. Expression of RET and co-expression of its activating ligand glial derived neurotropic factor (GDNF) is associated with transciptional signatures of ERK/MAPK activation and worse outcomes in human ER+ breast tumors.[23] Further, RET is overexpressed in ET-resistant breast cancer organoids that contain gain of function ESR1 (encoding the estrogen receptor) mutations and inhibiting RET suppresses tumor growth[24].

Our work has shown that tyrosine kinase inhibitors (TKIs), such as vandetanib and

sunitinib, that inhibit RET reduce proliferation and enhance sensitivity to ET in ER+ breast cancer

cell lines and xenografts.(25, 26) In tamoxifen resistant ER+/HER2+ BT-474 cells, inhibiting

RET increased sensitivity to tamoxifen and reduced ERK activation.(25) Similarly, we have

shown that *in* vitro treatment with sunitinib reduces phosho-ERK in RET+ primary human

breast tumors.(27) These findings support the use of vandetanib to reduce ERK/MAPK

activation and improve response to endocrine therapy.

Despite promise in the pre-clinic, clinical trials with vandetanib have had

disappointing results. However, the design of these trials failed to account for potentially

predictive biomarkers including ER and RET or only evaluated vandetanib as a monotherapy.

For example, a phase II trial of single-agent vandetanib in 46 patients with metastatic breast cancer

did not demonstrate a benefit(28), but the study did not report receptor status or select for ER+

patients, which makes it difficult to interpret the extent to which this trial included patients that

mechanistically would be predicted to respond. Our work, including the findings presented

herein, suggests that vandetinib may provide a clinical benefit for some patients. More recent

TKI trials with more specific inclusion criteria have shown more promising results and have

renewed interest in TKI for ET resistant ER+ breast cancer. A phase II trial of cabozantinib, in

metastatic ER+/HER2- was positive by objective response rate(29), as was a phase Ib/II trial of

lenvatinib with letrozole(30). More precise biomarkers are needed to identify patients most likely

to respond to TKI therapy. To develop more precise biomarkers and elucidate bypass resistance

mechanisms, we sought to investigate how vandetanib alters kinase signaling pathways in ET-

sensitive and ET-resistant models of ER+/HER2- breast cancer.

RESULTS

Vandetanib Inhibits ERK Activation and Viability in ER+ Breast Cancer Cells

We previously showed that vandetanib treatment decreases ERK activation in MCF7 breast cancer cells. To investigate if vandetanib similarly decreases ERK activation in ET-resistant ER+ breast cancer cells we used MCF7 and two ET-resistant derivatives, MCF7-TAM (tamoxifen) and MCF7-EXE (exemestane) which had been cultured long-term under drug treatment to develop resistance. pERK/ERK was reduced in MCF7 with vandetanib treatment (-81%, $p<0.001$). Similarly, pERK/ERK was reduced in MCF-TAM with vandetanib treatment (-78%, $p=0.025$) and MCF7-EXE (-78%, $p=0.01$), Fig. 1A. Dose response curves demonstrated an IC50 for vandetanib of 8.5 μM in MCF7, 10.4 μM in MCF7-TAM, and 15.9 μM in MCF7-EXE (Fig. 1B). Similar vandetanib sensitivities were seen in T47D and T47D-TAM (Supplementary Fig. 1)

Vandetanib Sensitizes ER+ Breast Cancer Cells to Tamoxifen

Because we had previously shown interaction between vandetanib and ER downstream signaling, we tested if co-treatment with vandetanib enhanced antiproliferative effects of tamoxifen. The IC50 of the active metabolite of tamoxifen, 4OH-Tam, was 10.7 μM in MCF7, 15.8 μM in MCF7-TAM, and 14.1 μM in MCF7-EXE (Fig. 1C) demonstrating decreased sensitivity to tamoxifen with chronic ET treatment.Treatment with 10 μM vandetanib resulted in left shift of the 4OH-Tam sensitivity curve in all 3 cell lines with the IC50 for 4OH-Tam of 3.8 μM in MCF7, 4.8 μM in MCF7-TAM, and 4.8 μM in MCF7-EXE. This demonstrates that cells are more sensitive to tamoxifen in the presence of vandetanib. To confirm synergy, we calculated the combination index (CI) of a combined treatment of 5 μM vandetanib with 2 μM 4OH-Tam

(Fig. 1D). When compared to the individual treatments (i.e., Tamoxifen or vandetanib alone) the

combination index was 0.47 for MCF7-TAM, 0.55 for MCF7-EXE and 0.59 for MCF7, indicating

increased synergy in ET-chronically treated cells.

RET Enhances Proliferation and Partially Mediates Vandetanib Sensitivity in MCF7 Cells

To better elucidate the role of RET in ER+ breast cancer cells, we performed a CRISPR

knockout in MCF7 cells (RET KO). Western blot confirmed knockout of RET protein (Fig. 2A).

RET KO also resulted in an unexpected reduction in ER protein and increased p-ERK. For

unbiased assessment of transcriptional effects of RET KO in MCF7, we performed mRNA

sequencing. Differentially regulated genes are shown in (Fig. 2B) including reduced ESR1

transcription in RET KO cells (log2FC -1.27, padj <0.001) compared to wild-type (WT) as was

seen by Western blot. Hallmark gene sets for estrogen response and KRAS signaling were depleted

with RET KO, as was a signature of RET knockout. RET KO resulted in lower proliferation score

and a gene expression profile that is more closely associated with the indolent, estrogen responsive

Luminal A centroid compared to WT. RET KO cells had significantly reduced growth at 48 and

72 hours relative to WT cells (Fig. 2C).

We used this model to determine the impact of RET KO on sensitivity to vandetanib. RET

KO cells were significantly less sensitive to vandetanib compared to WT (Fig. 2D). Similarly,

transient RET antagonism with the selective RET inhibitor BLU667 resulted in reduced sensitivity

to vandetanib in WT MCF7. BLU667 also reduced vandetanib sensitivity in MCF7-TAM, but not

in the low RET expressing MCF7-EXE (Supplemental Fig. 2). Based on the vandetanib synergy

assay we hypothesized that RET KO would sensitize cells to tamoxifen, but observed that KO cells

were less sensitive, likely due to the unexpected reduction in ER expression in KO cells and

reduced hallmark estrogen response gene expression profile. Cumulatively these results

demonstrate that RET is a partial mediator of vandetanib response in ER+ breast cancer cell line

models.

Vandetanib Regulated Genes in ET Sensitive and Resistant ER+ Breast Cancer

To understand more comprehensively how ER+ breast cancer cells are responding to

vandetanib treatment, we performed mRNA sequencing after treatment with Vandetanib (10 μM)

compared to vehicle. In MCF7, downregulated genes were enriched for MAPK signaling and

proliferative signatures, Fig. 3A, consistent with loss of pERK observed in Fig. 1A. Upregulated

genes were enriched for signatures of estrogen response and sensitivity to tamoxifen consistent

with sensitization of MCF7 cells to tamoxifen (Fig. 1D). Up-regulated and down-regulated gene

sets in MCF7-TAM (Fig. 3B) and MCF7-EXE (Fig. 3C) showed similar depletion of proliferative

and MAPK associated signatures and enrichment of estrogen and tamoxifen response signatures.

Similar transcriptional effects of vandetanib were seen in T47D and T47D-TAM (Supplementary

Fig. 3).

Because estrogen response signatures were enriched and proliferative signatures reduced

with vandetanib treatment, which are features of the Luminal A molecular subtype, we analyzed

the effect of vandetanib on correlation to the LumA centroid (Fig. 3D). This analysis demonstrated

vandetanib treatment resulted in a stronger LumA correlation supporting that treatment is inducing

a more indolent and ET responsive transcriptional phenotype.

To elucidate common and unique vandetanib regulated genes we determined overlap across

the parental and ET resistant MCF7 cell lines. This demonstrated that most genes with altered

expression in response to vandetanib were cell line specific, although the enrichment analysis

indicated regulation of biologic processes is more conserved across the 3 cell lines. Notably,

significant commonly upregulated genes included *SELENBP1* and *BAG1*, which are associated

with reduced proliferation, improved survival, and increased ER transcriptional activity in ER+

breast tumors(31-33). In the common downregulated genes, *AURKA* is associated with poor

survival in ER+ breast cancer patients(34). Reduced expression of *ECT2* is associated with

abrogation of MAPK signaling(35). Finally, several key regulators of cell cycle progression

associated with higher grade, worse outcomes, and poor response to endocrine therapy in ER+

breast cancer are commonly downregulated including *CCND1*, *TOP2A*, and *CCNB1*.(36-39)

To test if signatures of vandetanib regulated genes are associated with prognosis in patients

with ER+/HER2- breast cancer treated with endocrine therapy, we obtained survival annotated

transcriptional data from METABRIC(40) and SCAN-B(41). Tumors were stratified by median

gene set score using median thresholding. METABRIC patients with low downregulated gene set

score had significantly improved OS (HR=0.75, p<0.0001) and RFS (HR=0.60, p<0.0001).

Similarly, SCAN-B ER+ breast cancer patients with low downregulated gene set scores

demonstrated significantly lower mortality compared to high signature score (HR=0.87, p=0.034),

Fig. 3E. To determine whether vandetanib treatment is reducing expression of genes that drive an

ET- resistant phenotype, we obtained gene expression data from patients with ER+/HER2- breast

cancer profiled on ET and clinically annotated as sensitive or resistant(13). Resistant tumors had

significantly higher expression of the vandetanib downregulated signature compared to sensitive

tumors (4.3 vs 3.8, p<0.001), Fig. 3F. Cumulatively, these findings support that in ER+ breast

cancer, vandetanib treatment promotes gene expression changes associated with lower

proliferation, increase ET response, and improved survival.

Functional Signatures of Vandetanib Response

We used multiplex inhibitor bead enrichment coupled to mass spectrometry (MIB/MS) to

investigate changes to the functional kinome in response to vandatinib at the protein leve(42). In

this technique, kinases are enriched by incubating cell extracts with type 1 kinase inhibitors

immobilized on Sepharose beads. Each cell line was treated with the IC90 of vandetanib or

vehicle for 48 hours, Fig. 4A. We first investigated the effects of vandetanib on enrichment and

depletion of known vandetanib targets(43). RET, EGFR, ABL1, RIP2K, PTK6, and DDR1 had

similar magnitude of depletion in all 3 models, despite large differences in baseline enrichment

between MCF7 and the ET-resistant derivatives, Fig. 4B. Similar results were seen for T47D and

T47D-TAM (Supplementary Fig. 4). These results demonstrate that several distinct target proteins

may be mediating vandetanib effects in ER+ breast cancer cells and that effects of treatment do

not appear to be proportional to baseline expression, and that the relative level of expression of a

target alone may not be a good biomarker of response.

Enriched and depleted kinase proteins by cell line are shown in Fig. 4C. Overlap analysis

(Fig. 4D) showed that similar to transcriptional findings, kinome dynamics were heterogenous

across cell lines. There were 12 common upregulated and 22 downregulated kinase proteins.

Common upregulated genes included AURKA, PLK1, PIP4K2A, which promote proliferation and

aggressive tumors(44-46), as well as JAK1 and TYK2 which are associated with immune

infiltrates and are upstream of AKT signaling(47, 48). Common depleted proteins with vandetanib

treatment included proliferation promoting cell cycle regulators CDK1, CKD2, and PKMYT1.

Depletion of MAP2K2 (MEK2) as well as upstream receptor tyrosine kinases EPHA1, EPHB4,

DDR1, and EGFR suggest downregulation of the MAPK signaling pathway in response to

vandetanib treatment consistent with Western blots (Fig. 1A). Gene set enrichment analysis

demonstrated a common downregulation of the ERK cascade and upregulation of the PI3K/AKT

pathway, Fig. 4E.

Pharmacologic Inhibition of Adaptive Resistance Kinases Enhances Vandetanib Efficacy

Upregulation of functional kinases may identify common mechanisms of adaptive bypass

resistance so expression levels of druggable kinases(24) were assessed by treatment (Fig. 5A). To

determine whether enriched druggable targets may be mediating adaptive resistance to vandetanib

we performed co-treatment assays on the principle that inhibiting mediators of adaptive resistance

should enhance sensitivity to vandetanib. Because ERBB2/HER2 was enriched in all 3 MCF7 cell

lines, we co-treated cells with vandetanib and the FDA approved HER2 small molecule inhibitor

lapatinib. Treatment with lapatinib resulted in significantly augmented vandetanib reduction in cell

viability in all 3 cell lines, (Fig. 5B).

Vandetanib Response at Single Cell Resolution in Primary ER+ Breast Cancer

We used our previously validated *ex vivo* short term drug treatment platform coupled to

single cell RNA sequencing(49) to assess how primary breast tumor cells respond to vandetanib

treatment. Specifically, we obtained a primary ER+/HER2- breast cancer, dissociated into single

cells, treated with control of vandetanib containing media for 12 hours and assayed by 10x

scRNAseq. Uniform manifold projection (UMAP) was performed using Seurat v3(50). Cell

populations were annotated using known transcriptional profiles and canonical markers (Fig. 6A)

and by treatment group (Fig. 6B). Malignant (tumor) cells were distinguished from non-tumor

epithelial cells as cells having significant inferred copy number alterations by InferCNV(51).

Differentially expressed genes were identified using a false discovery rate (FDR) cutoff of <

0.05 in the iCNV+ tumor cell compartment with vandetanib treatment compared to control,

Fig. 6C. Enriched and depleted pathways in vandetanib treated tumor cells relative to control

treated were determined, Fig. 6D. Of note, as seen in cell lines, ESR1 targets and ERBB2 pathway

signaling were enriched in vandetanib treated cells (Fig. 6E).

We tested the overlapping downregulated genes after vandetanib treatment across the 3

MCF7 cell lines (Fig. 3) and these genes were also depleted in vandetanib treated primary tumor

cells (Fig. 6E). Next LumA correlation score was assigned to each tumor cell using a validated

single cell PAM50 classifier(52). Treated tumor cells had a stronger correlation to the LumA

centroid (median score treated 0.259 vs median score untreated 0.244, p= 0.001, Wilcoxon rank-

sum test). Thus, short term vandetanib treatment alters the transcriptional profile of ER+/HER2-

primary tumor cells towards a more indolent and estrogen responsive phenotype.

To explore features of vandetanib response within primary tumor cells we used non-

negative matrix factorization (NMF) to elucidate features that were enriched and depleted in

vandetanib treated cells (Supplementary Fig. 5A). This analysis yielded metagene modules that

were enriched in vandetanib treated samples (module 1) and depleted in vandetanib treated cells

(Module 6) (Supplementary Fig. 5B). We next identified features that may be associated with

vandetanib response by comparing module 1 and 6 gene expression profiles. Module 1 (vandetanib

resistant) had expected high expression of genes that increase with vandetanib treatment,

consistent with reduced response. Module 6 was enriched for MYC targets, which is known to

support tamoxifen resistance, but may support a transcriptional profile of vandetanib sensitivity

(Supplementary Fig. 5C). Module 1 was also enriched for MTORC signaling which we identified

in cells lines as an adaptive vandetanib resistance mechanism. Interestingly, module 1 was

enriched for hallmark estrogen response genes and our previously generated single cell tamoxifen

response score(49) supporting that cells with poor vandetanib response have enhanced tamoxifen

sensitivity (Supplementary Fig. 5D). Lastly, we looked at our human tumor derived signatures of

high GDNF/RET expressing human ER+-breast tumors(23). Genes that were depleted in

GDNF/RET high human tumors were depleted in vandetanib sensitive cells, supporting that

signature as a biomarker of response (Supplementary Fig. 5E).

DISCUSSION

Resistance to endocrine therapy remains a common problem and a significant driver of poor patient outcomes despite advances in understanding of how ER+ breast tumors respond to therapy. The identification of activation in MAPK pathways as a driver of resistance to ET has generated significant interest in targeting this pathway to enhance ET response. However, clinical trials have demonstrated inconsistent and disappointing results regarding response to tyrosine kinase inhibitors for non HER2-amplified tumors. These findings demonstrate the need for improved understanding of how intracellular signaling pathways are altered by treatment to determine biomarkers of patients most likely to respond and clinically actionable kinase targeting strategies in MAPK-activated ET resistant breast tumors.

We found that response to vandetanib, measured by IC50 or by depletion of RET activity, was largely the same across the cell lines tested, despite large differences in baseline RET expression, and this was also true for the other direct vandetanib targets. These findings demonstrate that expression of a protein that is a target for a TKI alone may not be an effective biomarker of response. Signatures of activity may be more informative of response than single gene expression. In fact, HER2 targeting therapeutics, expression of the HER2E gene expression pattern is more predictive of response than HER2 expression alone(53). We recently generated a signature of RET active ER+ breast tumors(23) using treatment naïve samples from The Cancer Genome Atlas (TCGA), and the low expression of genes that were depleted in GDNF/RET high tumors was associated with vandetanib sensitivity, suggesting that this may be a biomarker to select patients most likely to respond.

Prior work has shown that reprogramming of tyrosine kinase signaling networks can be

underscored by changes in gene expression. Transcriptional changes with vandetanib treatment

did identify potential mediating transcription factors. Specifically, altered MYC gene regulatory

function determines kinase reprogramming in response to MEK inhibition in triple negative breast

cancer(54), and FOXA1 binding determines reprogramming in HER2-directed therapy inhibited

HER2-amplified breast cancer(55). How ER+/HER2- breast cancer cells alter gene regulation to

impact remodeling of kinase signaling, both with emerging resistance to ET, and with TKI such

as vandetanib, are important mechanistic questions for further study.

Kinase signaling networks are plastic and inhibiting one node commonly leads to adaptive

bypass through another pathway. We were able to identify mediators of adaptive resistance to

vandetanib treatment and augment response with dual treatment. Up-regulation of HER2 activity

was common to all cell lines and primary tumor cells and targeting HER2 with lapatinib enhanced

sensitivity to vandetanib. Activating HER2 signaling could open a number of efficacious

therapeutic options for patients. While HER2 activity was increased, there was no change in HER2

RNA expression, and this highlights the need for assessment at the level of protein activity to

determine adaptive bypass mechanisms.

While HER2 upregulation was a common adaptive bypass pathway, we also identified cell

line specific enriched proteins. This could be a result of distinct adaptations necessary to resist

exemestane compared to tamoxifen or could represent distinct alternative paths to resist vandetanib

treatment. How these adaptive mechanisms are conserved or unique in human tumors will be

essential to understand the therapeutic implications of these co-treatment targets. Specific adaptive

bypass pathways may be predictable from pretreatment gene expression or chromatin accessibility

states, or tumor level assessment of response to therapy may be needed to identify tumor specific

bypass mechanisms and cotreatment vulnerabilities. To this end, our group has recently developed

a method for whole kinome characterization on limited sample (such as a core needle biopsy),

which could make on treatment assessment of adaptive response and targeted co-treatment

strategies feasible.(56)

In summary, we demonstrate that ER+ breast cancer cells become more sensitive to ET

with vandetanib treatment. Vandetanib results in broad reprogramming of gene expression and

kinase signaling which results in a lower proliferative, more estrogen responsive luminal A-like

state. Early adaptive response, including upregulation of HER2, to vandetanib identified co-

treatment strategies to enhance response. Future trials will need to utilize signature-based

biomarkers of response with attention to tumor specific upregulated adaptive bypass pathways as

cotreatment strategies to improve efficacy.

METHODS

Cell Lines

We obtained ER+ breast cancer cell lines parental MCF7, tamoxifen-resistant MCF7

(MCF7-TAM), exemestane-resistant MCF7 (MCF7-EXE), parental T47D, and tamoxifen-

resistant T47D (T47D-TAM), from the American Type Culture Collection (ATCC). Cell lines

were cultured using Dulbecco's Modified Eagle Medium (DMEM) (Gibco) with 10% fetal bovine

serum (Gibco) and 1% penicillin-streptomycin (Life Technologies) at 37°C and 5% CO_2. MCF7-

TAM and T47D-TAM were maintained by chronic treatment with 1 μM of tamoxifen and MCF7-

EXE with 0.1 μM of exemestane.

Chemicals and Treatment

Vandetanib (SelleckChem) and z-4-Hydroxytamoxifen (4OH-Tam) (SelleckChem) was

dissolved in dimethyl sulfoxide (DMSO). GDNF (R&D Systems) was dissolved in phosphate-

buffered saline (PBS) and treatments were done at a final concentration of 25 ng/mL. Pralsetinib

was obtained from Chemietek, and lapatinib from Medchem Express LLC. Treatments were

performed for 24 hours unless otherwise specified.

Cell Viability Assays

Cells were plated on opaque-walled 96-well plate at a density of 4,000 cells per well in

technical triplicates and allowed to adhere overnight prior to the addition of the specified

inhibitors. Viability was assessed using the CellTiter-Glo (CTG) Luminescent Cell Viability Assay

(Promega). Plates were read using a Synergy 2 microplate reader. Samples were averaged over

biological triplicates. Statistical analysis and plotting were performed using GraphPad Prism 9.

Western Blots

Cells were lysed in Pierce RIPA buffer (Thermo Scientific) containing Halt protease inhibitor cocktail (Thermo Scientific) and PhosStop phosphatase inhibitor (Thermo Scientific). Proteins were resolved by SDS polyacrylamide gel electrophoresis, transferred to PVDF membrane, and blocked with 10% BSA, then incubated overnight with the indicated primary antibody in blocking buffer. Membranes were washed three times in tris buffered saline containing 0.1% tween-20 then then incubated for 1 hour with HRP-conjugated secondary antibody. Blots were developed using a Chemidoc-touch digital imager and band intensity quantified using ImageJ (https://imagej.net/ij/index.html).The antibodies used for Western blot included: p44/42 MAPK (ERK1/2) (Cell Signaling Technologies), Phospho-p44/42 MAPK (Erk1⁄2) (Cell Signaling Technologies), RET (Abcam), EGFR (Cell Signaling Technologies), and β-Actin (SantaCruz).

RET CRISPR Knockout

An sgRNA was designed to target exon 3 of RET (AGCACTCCCGCTCACGGTC), cloned into the pSpCas9(BB)-2A-Puro(PX459) V2.0 (Addgene) plasmid, and verified by Sanger sequencing. MCF-7 cells were cultured in antibiotic free media and grown to a confluency of 70%-90%. Cells were then transfected with sgRNA plasmid using FuGENE 4K Transfection Reagent according the manufacturer's recommendations. After 72 hours, cells were passaged onto 15 cm plates at a low density and selected with 1 μg/mL puromycin (Gibco) for 3 days. Cell colonies were then isolated by pipet, and DNA was isolated using the Quick-DNA Miniprep Kit (Zymo Research). The target sequence was then amplified by PCR and then sequenced for

verification of a frameshift mutation. Both RET alleles were found to have frameshift mutations

resulting in the introduction of premature stop codons.

mRNA Sequencing and Data Analysis

Cells were treated with 10 µM vandetanib or vehicle for 24 hours in biological triplicates

and total RNA was isolated (Qiagen). mRNA sequencing libraries were made using TruSeq RNA

Library Prep Kit (Illumina) and sequenced using the Illumina NextSeq2000. Plots and statistical

analyses were performed using RStudio 2022.07.1 Build 554.

Differential gene expression (DGE) was determined using the DESeq2 R package using

raw mRNA-seq counts from STAR/Salmon. Genes with less than 10 counts across all samples

were omitted before analysis and counts were filtered and normalized to the upper quartile using

NOISeq. Significant differentially expressed genes were defined by using a Log2FC (log2 fold

change) cutoff of 0.5 for upregulation and -0.5 for downregulation, with a false discovery rate

(FDR) < 0.05. Volcano plots of all genes were plotted using the EnhancedVolcano R package.

Enrichment of gene sets was then investigated through hypergeometric analysis of an

unranked list of upregulated or downregulated elements using the enricher function of the

clusterProfiler package. Gene sets from MSigDB were first compiled and screened for keywords

of interest pertaining to breast cancer, including estrogen and ER signaling.

METABRIC and SCAN-B databases were obtained for survival analysis and filtered for

ER+/HER2- patients treated with ET. A median-centroid approach was implemented to assign a

gene set score per patient. Kaplan-Meier (KM) plots were then computed using the survminer and

survival packages.

MIB/MS and Data Analysis

Cells were treated with vandetanib at its IC90 for 48 hours. Cells were harvested by

scraping and cell pellets obtained for proteomic analysis using the MIB/MS technology(42, 54).

Scaled label free quantification (LFQ) values were obtained and a correlation plot of samples was

computed to assess homogeneity of triplicates and heterogeneity of treatment conditions for

quality control using ggplot2. Differential protein expression (DPE) was analyzed using the limma

R package using scaled LFQ values. Significant differentially expressed proteins were defined

using a Log2FC (log2 fold change) cutoff of 0.5 for upregulation and -0.5 for downregulation,

along with a p-value < 0.05.

Primary ER+/HER2- Human Tumor Cells

A primary tumor sample was obtained from a patient with ER+/HER2- invasive ductal

carcinoma. The tumor was processed fresh as previously described(49). In brief, tumor tissue was

transported immediately to the laboratory after surgical resection, dissociated into a single cell

suspension, and treated with vandetanib (10 µM) or control media for 12 hours. After treatment,

single cell RNA sequencing libraries were made using the 10x Genomics Chromium Next GEM

Single Cell 3' v3.1. Libraries were sequenced on the NextSeq2000 (Illumina) with pair-end

sequencing.

Single cell RNA sequencing analysis

Reads were aligned and counted using Cellranger count function with default parameters.

The counts and features matrixes were used to generate Seurat objects. The Seurat objects

corresponding to each treatment group were merged with Seurat's function merge(), and low

quality cells were removed. Doublets were removed using scDblFinder

(https://pubmed.ncbi.nlm.nih.gov/35814628/) and cells were further filtered with the following

parameters: nFeature_RNA > 200 & nFeature_RNA < 6000 & percent.mt < 10.

Cell types were annotated on the combined Seurat using SingleR using the cell type atlas

HumanPrimaryCellAtlasData() accessed with the package celldex. InferCNV was used on the

combined Seurat to predict copy number alterations, defining the immune cells

("T_cells","Macrophage","DC","Monocyte","NK_cell") as reference cells (cells without CNVs).

The tumor cells were identified as cells with several CNVs, using the copy number heatmap per

cell.

Differential analysis of gene expression was performed on the tumor cells to compare

vandetanib-treated versus non-treated cells using findmarkers(). Gene Set Enrichment Analysis

was then performed using the list of differentially expressed genes, ranking the genes according to

log2(FC)/FDR, and the ranked gene lists were used in the GSEA software

(https://pubmed.ncbi.nlm.nih.gov/16199517/)   (https://pubmed.ncbi.nlm.nih.gov/12808457/).

Gene signatures were scored using AddModuleScore_UCell() from the UCell package.

We used the package NMF to build a non-negative matrix factorization (NMF) model of the tumor

cells using a list of genes differentially expressed upon vandetanib treatment in MCF7 cells from

this study. The Seurat was subset to contain only the above-mentioned genes, and the NMF model

was fitted to 8 factors. Heatmaps were built for each treatment group to assess enrichment of cells

to different NMF factors. Each cell was assigned to a unique NMF according to the NMF with

maximum coefficient value for that cell, and the percentage of cells belonging to each NMF factor

was calculated for each treatment group. DE analysis was performed comparing cells in certain

NMF factors.

Statistics

Study Approval

424   Patient samples were obtained after written informed consent under a protocol approved

425   by the institutional review board at the University of North Carolina (IRB#22-1888) in accordance

426   with the US Common Rule.

Sex as a Biologic Variable

428   Breast cancer occurs primarily in women and so primary tumor cells were obtained from a

429   female patient with breast cancer.

Statistics

431   All experiments were performed with at least 3 biological or technical replicates. Small

432   sample number comparisons were analyzed with an unpaired t test with Welch's correction for

433   non-equal SD. Linear regressions and nonlinear regression curves were compared by statistical

434   tests on the parameters of the curves using the extra sum of squares F test. Experiments with large

435   sample numbers were analyzed by Wilcoxon test.

Data Availability

437   Single cell and bulk RNA sequencing data was made available at the Sequence Read

438   Archive (SRA) under BioProject PRJNA1177387. Mass spectrometry protein sequencing results

439   are included in this manuscript as Supplementary Table 1.

Code Availability

441   All original code has been deposited on the Github, and it is publicly available at the Github

442   repository at https://github.com/santiagohaase/Spanheimer-lab-Kinase-Plasticity-in-Response-to-

443   Vandetanib-Enhances-Sensitivity-to-Tamoxifen

ACKNOWLEDGEMENTS

Dr Philip Spanheimer is supported by the National Institutes of Health grants R37CA292075, K08CA280388, and a Solid Tumor Research Scholars Grant from Gilead Sciences. CMP was supported by funds from the NCI breast SPORE program (P50-CA058223). This work was additionally supported in part by P30CA016086 UNC Lineberger Comprehensive Cancer Center Core Support Grant.

 FIGURES

Vandetanib dose-response curves

Tamoxifen dose-response curves

465
Figure 1. (A) Treatment of wild-type MCF-7 and tamoxifen resistant MCF7 (MCF7-TAM) and exemestane
resistant MCF7 (MCF7-EXE) reduces pERK/ERK in all 3 cell lines. (B) Vandetanib dose dependent reduction in
cell viablilty in MCF7, MCF7-TAM, and MCF7-EXE. (C) Treatment with Vandetanib results in a left shift of the
tamoxifen sensitivity curve and reduction in IC50 in all 3 cell lines. (D) Synergy plots demonstrate synergistic
activity of tamoxifen and vandetanib.*$p<0.05$, ***$p<0.01$ by t-test with Welch correction and analysis of the slope
difference in the nonlinear regression model (B, C)

472
Figure 2. (A) Western blot demonstrating CRISPR knockout of RET resulting in >90% reduction in RET protein
levels. There was no significant change in pERK/ERK. (B) Heatmap demonstrating differentially expressed genes
with RET KO in MCF7. Enrichment analysis demonstrates reduction in hallmark estrogen response and RAS signaling
in RET KO cells. RET KO cells have lower proliferation score and higher Luminal A correlation. (D) RET knockout
cells have reduced sensitivity to vandetanib. Transient inhibition of RET via co-treatment with the selective RET
inhibitor BLU667 reduces sensitivity to vandetanib in a dose dependent fashion in MCF7 cells. RET KO cells have
reduced sensitivity to tamoxifen. (E) Enrichment analysis demonstrates reduction in hallmark estrogen response and
RAS signaling in RET KO cells. *p<0.05, **p<0.01, ***p<0.001, ****p<0.0001 by t-test with Welch correction for
the bar graphs, and Wilcoxon test for the heatmap in (B).

482
Figure 3. Heatmap and enrichment analysis of differentially expressed genes in (A) MCF7, (B) MCF7-TAM, (C)
MCF7-EXE. (D) Vandetanib treatment increases the Luminal A centroid correlation. (E) Overlap of up- and down-
regulated genes by cell line. (F) Signatures of overlapping up-and down regulated genes are associated with recurrence
free and overall survival in ER+ breast cancer patients in METARIC and SCAN-B studies and (F) downregulated
genes are enriched in clinically ET-resistant ER+ tumors. *p<0.05, **p<0.01, ***p<0.001, ****p<0.0001 by t-test
with Welch correction for (D) and Wilcoxon test (D).

489
Figure 4. (A) Multiplex inhibitor bead mass spectrometry assay work flow. (B) Effects of vandetanib on enrichment
and depletion of known vandetanib targets. Baseline expression of vandetanib targets in tamoxifen and exemestane
resistant cell lines compared to WT MCF7. (C) Heatmap showing enriched and depleted proteins after vandetanib
treatment. (D) Overlap analysis of up- and downregulated proteins between the 3 cell lines when treated with
vandetanib. (E) Pathway analysis of the downregulation of the ERK cascade and upregulation of the PI3K/AKT
pathway.

496
497 **Figure 5.** (A) Heatmap showing enrichment levels of druggable kinases of sensitive and resistant MCF7 cell lines
498 before and after vandetanib treatment. (B) Sensitivity plots of co-treated cells with vandetanib and HER2 inhibitor
499 lapatinib demonstrating enhanced lapatinib sensitivity with vandetanib treatment. (C) Sensitivity plots of co-
500 treatment with vandetanib and FGFR4 inhibitor BLU991 show selective sensitization in MCF7-EXE. *p<0.05,
501 ***p<0.001, ****p<0.0001 by t-test with Welch correction. (B-C)
502
503
504

505
Figure 6. (A) Single-cell RNA sequencing uniform manifold projection (UMAP) annotated by cell type within
a primary ER+ human breast tumor. (B) Tumor cells were computationally isolated as having significant
inferred copy number changes and plotted by treatment condition. (C) Heatmap showing differentially
expressed genes compared between vandetanib treated vs non-treated tumor cells. (D) Enriched and depleted
pathways of vandetanib treated vs non-treated tumor cells. (E) ERBB2 (encoding HER2) expression was
increased in vandetanib treated tumor cells as was ERBB2 activity signature score. A signature of down
regulated genes in vandetanib treated cell lines is depleted in vandetanib treated primary tumor cells, and
LumA correlation increased in vandetanib treated tumor cells. Significant signatures filtered by < 0.1 FDR on
GSEA analysis (C) and significant genes filtered by adjusted p value <0.05 (Wilcoxon test).

REFERENCES

1. Nabholtz JM, Tonkin K, Smylie M, Au HJ, Lindsay MA, and Mackey J. Chemotherapy of breast cancer: are the taxanes going to change the natural history of breast cancer? *Expert Opin Pharmacother.* 2000;1(2):187-206.

2. Mouridsen H, Gershanovich M, Sun Y, Perez-Carrion R, Boni C, Monnier A, et al. Phase III study of letrozole versus tamoxifen as first-line therapy of advanced breast cancer in postmenopausal women: analysis of survival and update of efficacy from the International Letrozole Breast Cancer Group. *J Clin Oncol.* 2003;21(11):2101-9.

3. Bachelot T, Bourgier C, Cropet C, Ray-Coquard I, Ferrero JM, Freyer G, et al. Randomized phase II trial of everolimus in combination with tamoxifen in patients with hormone receptor-positive, human epidermal growth factor receptor 2-negative metastatic breast cancer with prior exposure to aromatase inhibitors: a GINECO study. *J Clin Oncol.* 2012;30(22):2718-24.

4. Giaquinto AN, Sung H, Newman LA, Freedman RA, Smith RA, Star J, et al. Breast cancer statistics 2024. *CA Cancer J Clin.* 2024.

5. Harvey JM, Clark GM, Osborne CK, and Allred DC. Estrogen receptor status by immunohistochemistry is superior to the ligand-binding assay for predicting response to adjuvant endocrine therapy in breast cancer. *J Clin Oncol.* 1999;17(5):1474-81.

6. Jeselsohn R, Barry WT, Migliaccio I, Biagioni C, Zhao J, De Tribolet-Hardy J, et al. TransCONFIRM: Identification of a Genetic Signature of Response to Fulvestrant in Advanced Hormone Receptor-Positive Breast Cancer. *Clin Cancer Res.* 2016;22(23):5755-64.

7. Finn RS, Martin M, Rugo HS, Jones S, Im SA, Gelmon K, et al. Palbociclib and Letrozole in Advanced Breast Cancer. *N Engl J Med.* 2016;375(20):1925-36.

8. Baselga J, Campone M, Piccart M, Burris HA, 3rd, Rugo HS, Sahmoud T, et al. Everolimus in postmenopausal hormone-receptor-positive advanced breast cancer. *N Engl J Med.* 2012;366(6):520-9.

9. Piccart M, Hortobagyi GN, Campone M, Pritchard KI, Lebrun F, Ito Y, et al. Everolimus plus exemestane for hormone-receptor-positive, human epidermal growth factor receptor-2-negative advanced breast cancer: overall survival results from BOLERO-2dagger. *Ann Oncol.* 2014;25(12):2357-62.

10. Andre F, Ciruelos E, Rubovszky G, Campone M, Loibl S, Rugo HS, et al. Alpelisib for PIK3CA-Mutated, Hormone Receptor-Positive Advanced Breast Cancer. *N Engl J Med.* 2019;380(20):1929-40.

11. Turner NC, Slamon DJ, Ro J, Bondarenko I, Im SA, Masuda N, et al. Overall Survival with Palbociclib and Fulvestrant in Advanced Breast Cancer. *New Engl J Med.* 2018;379(20):1926-36.

12. Razavi P, Chang MT, Xu G, Bandlamudi C, Ross DS, Vasan N, et al. The Genomic Landscape of Endocrine-Resistant Advanced Breast Cancers. *Cancer Cell.* 2018;34(3):427-38 e6.

13. Xia Y, He X, Renshaw L, Martinez-Perez C, Kay C, Gray M, et al. Integrated DNA and
RNA Sequencing Reveals Drivers of Endocrine Resistance in Estrogen Receptor-Positive
Breast Cancer. *Clin Cancer Res.* 2022;28(16):3618-29.
14. Arpino G, Wiechmann L, Osborne CK, and Schiff R. Crosstalk between the estrogen
receptor and the HER tyrosine kinase receptor family: molecular mechanism and clinical
implications for endocrine therapy resistance. *Endocr Rev.* 2008;29(2):217-33.
15. Meijer D, Sieuwerts AM, Look MP, van Agthoven T, Foekens JA, and Dorssers LC.
Fibroblast growth factor receptor 4 predicts failure on tamoxifen therapy in patients with
recurrent breast cancer. *Endocr Relat Cancer.* 2008;15(1):101-11.
16. Bergqvist J, Elmberger G, Ohd J, Linderholm B, Bjohle J, Hellborg H, et al. Activated
ERK1/2 and phosphorylated oestrogen receptor alpha are associated with improved
breast cancer survival in women treated with tamoxifen. *Eur J Cancer.* 2006;42(8):1104-
12.
17. Kirkegaard T, Witton CJ, McGlynn LM, Tovey SM, Dunne B, Lyon A, et al. AKT
activation predicts outcome in breast cancer patients treated with tamoxifen. *J Pathol.*
2005;207(2):139-46.
18. Shoman N, Klassen S, McFadden A, Bickis MG, Torlakovic E, and Chibbar R. Reduced
PTEN expression predicts relapse in patients with breast carcinoma treated by tamoxifen.
Mod Pathol. 2005;18(2):250-9.
19. Morandi A, Martin LA, Gao Q, Pancholi S, Mackay A, Robertson D, et al. GDNF-RET
signaling in ER-positive breast cancers is a key determinant of response and resistance to
aromatase inhibitors. *Cancer Res.* 2013;73(12):3783-95.
20. Boulay A, Breuleux M, Stephan C, Fux C, Brisken C, Fiche M, et al. The Ret receptor
tyrosine kinase pathway functionally interacts with the ERalpha pathway in breast cancer.
Cancer Res. 2008;68(10):3743-51.
21. Plaza-Menacho I, Morandi A, Robertson D, Pancholi S, Drury S, Dowsett M, et al.
Targeting the receptor tyrosine kinase RET sensitizes breast cancer cells to tamoxifen
treatment and reveals a role for RET in endocrine resistance. *Oncogene.*
2010;29(33):4648-57.
22. Spanheimer PM, Woodfield GW, Cyr AR, Kulak MV, White-Baer LS, Bair TB, et al.
Expression of the RET proto-oncogene is regulated by TFAP2C in breast cancer
independent of the estrogen receptor. *Ann Surg Oncol.* 2013;20(7):2204-12.
23. Kakati RT, Kim H, Whitman A, and Spanheimer PM. High expression of the RET
receptor tyrosine kinase and its ligand GDNF identifies a high-risk subset of estrogen
receptor positive breast cancer. *Breast Cancer Res Treat.* 2023.
24. Gou X, Kim BJ, Anurag M, Lei JT, Young MN, Holt MV, et al. Kinome reprogramming
is a targetable vulnerability in ESR1 fusion-driven breast cancer. *Cancer Res.* 2023.
25. Spanheimer PM, Cyr AR, Gillum MP, Woodfield GW, Askeland RW, and Weigel RJ.
Distinct pathways regulated by RET and estrogen receptor in luminal breast cancer
demonstrate the biological basis for combination therapy. *Ann Surg.* 2014;259(4):793-9.

26.   Spanheimer PM, Park JM, Askeland RW, Kulak MV, Woodfield GW, De Andrade JP, et
597   al. Inhibition of RET increases the efficacy of antiestrogen and is a novel treatment
598   strategy for luminal breast cancer. *Clin Cancer Res.* 2014;20(8):2115-25.

599   27.   Spanheimer PM, Lorenzen AW, De Andrade JP, Kulak MV, Carr JC, Woodfield GW, et
600   al. Receptor Tyrosine Kinase Expression Predicts Response to Sunitinib in Breast
601   Cancer. *Ann Surg Oncol.* 2015;22(13):4287-94.

602   28.   Miller KD, Trigo JM, Wheeler C, Barge A, Rowbottom J, Sledge G, et al. A multicenter
603   phase II trial of ZD6474, a vascular endothelial growth factor receptor-2 and epidermal
604   growth factor receptor tyrosine kinase inhibitor, in patients with previously treated
605   metastatic breast cancer. *Clin Cancer Res.* 2005;11(9):3369-76.

606   29.   Xu J, Higgins MJ, Tolaney SM, Come SE, Smith MR, Fornier M, et al. A Phase II Trial
607   of Cabozantinib in Hormone Receptor-Positive Breast Cancer with Bone Metastases.
608   *Oncologist.* 2020.

609   30.   Lim JSJ, Wong ALA, Ow SGW, Ngoi NYL, Chan GHJ, Ang YLE, et al. Phase Ib/II dose
610   expansion study of lenvatinib combined with letrozole in post-menopausal women with
611   hormone receptor positive breast cancer. *Clin Cancer Res.* 2022.

612   31.   Zhang S, Li F, Younes M, Liu H, Chen C, and Yao Q. Reduced selenium-binding protein
613   1 in breast cancer correlates with poor survival and resistance to the anti-proliferative
614   effects of selenium. *PLoS One.* 2013;8(5):e63702.

615   32.   Chen R, Wang X, Fu J, Liang M, and Xia T. High FLT3 expression indicates favorable
616   prognosis and correlates with clinicopathological parameters and immune infiltration in
617   breast cancer. *Front Genet.* 2022;13:956869.

618   33.   Papadakis ES, Reeves T, Robson NH, Maishman T, Packham G, and Cutress RI. BAG-1
619   as a biomarker in early breast cancer prognosis: a systematic review with meta-analyses.
620   *Br J Cancer.* 2017;116(12):1585-94.

621   34.   Kahl I, Mense J, Finke C, Boller AL, Lorber C, Gyorffy B, et al. The cell cycle-related
622   genes RHAMM, AURKA, TPX2, PLK1, and PLK4 are associated with the poor
623   prognosis of breast cancer patients. *J Cell Biochem.* 2022;123(3):581-600.

624   35.   Yi M, Zhang D, Song B, Zhao B, Niu M, Wu Y, et al. Increased expression of ECT2
625   predicts the poor prognosis of breast cancer patients. *Exp Hematol Oncol.*
626   2022;11(1):107.

627   36.   Lundberg A, Lindstrom LS, Li J, Harrell JC, Darai-Ramqvist E, Sifakis EG, et al. The
628   long-term prognostic and predictive capacity of cyclin D1 gene amplification in 2305
629   breast tumours. *Breast Cancer Res.* 2019;21(1):34.

630   37.   Montalto FI, and De Amicis F. Cyclin D1 in Cancer: A Molecular Connection for Cell
631   Cycle Control, Adhesion and Invasion in Tumor and Stroma. *Cells.* 2020;9(12).

632   38.   An X, Xu F, Luo R, Zheng Q, Lu J, Yang Y, et al. The prognostic significance of
633   topoisomerase II alpha protein in early stage luminal breast cancer. *BMC Cancer.*
634   2018;18(1):331.

635   39.   Ding K, Li W, Zou Z, Zou X, and Wang C. CCNB1 is a prognostic biomarker for ER+
636   breast cancer. *Med Hypotheses.* 2014;83(3):359-64.

40. Curtis C, Shah SP, Chin SF, Turashvili G, Rueda OM, Dunning MJ, et al. The genomic
and transcriptomic architecture of 2,000 breast tumours reveals novel subgroups. *Nature.*
2012;486(7403):346-52.
41. Saal LH, Vallon-Christersson J, Hakkinen J, Hegardt C, Grabau D, Winter C, et al. The
Sweden Cancerome Analysis Network - Breast (SCAN-B) Initiative: a large-scale
multicenter infrastructure towards implementation of breast cancer genomic analyses in
the clinical routine. *Genome Med.* 2015;7(1):20.
42. Duncan JS, Whittle MC, Nakamura K, Abell AN, Midland AA, Zawistowski JS, et al.
Dynamic reprogramming of the kinome in response to targeted MEK inhibition in triple-
negative breast cancer. *Cell.* 2012;149(2):307-21.
43. Klaeger S, Heinzlmeir S, Wilhelm M, Polzer H, Vick B, Koenig PA, et al. The target
landscape of clinical kinase drugs. *Science.* 2017;358(6367).
44. Zhu L, Lu Z, and Zhao H. Antitumor mechanisms when pRb and p53 are genetically
inactivated. *Oncogene.* 2015;34(35):4547-57.
45. Emerling BM, Hurov JB, Poulogiannis G, Tsukazawa KS, Choo-Wing R, Wulf GM, et
al. Depletion of a putatively druggable class of phosphatidylinositol kinases inhibits
growth of p53-null tumors. *Cell.* 2013;155(4):844-57.
46. O'Connell CE, and Vassilev A. Combined Inhibition of p38MAPK and PIKfyve
Synergistically Disrupts Autophagy to Selectively Target Cancer Cells. *Cancer Res.*
2021;81(11):2903-17.
47. Zhang Q, Sturgill JL, Kmieciak M, Szczepanek K, Derecka M, Koebel C, et al. The role
of Tyk2 in regulation of breast cancer growth. *J Interferon Cytokine Res.* 2011;31(9):671-
7.
48. Chen B, Lai J, Dai D, Chen R, Li X, and Liao N. JAK1 as a prognostic marker and its
correlation with immune infiltrates in breast cancer. *Aging (Albany NY).*
2019;11(23):11124-35.
49. Kim H, Whitman AA, Wisniewska K, Kakati RT, Garcia-Recio S, Calhoun BC, et al.
Tamoxifen Response at Single Cell Resolution in Estrogen Receptor-Positive Primary
Human Breast Tumors. *Clin Cancer Res.* 2023.
50. Stuart T, Butler A, Hoffman P, Hafemeister C, Papalexi E, Mauck WM, 3rd, et al.
Comprehensive Integration of Single-Cell Data. *Cell.* 2019;177(7):1888-902 e21.
51. Puram SV, Tirosh I, Parikh AS, Patel AP, Yizhak K, Gillespie S, et al. Single-Cell
Transcriptomic Analysis of Primary and Metastatic Tumor Ecosystems in Head and Neck
Cancer. *Cell.* 2017;171(7):1611-24 e24.
52. Wu SZ, Al-Eryani G, Roden DL, Junankar S, Harvey K, Andersson A, et al. A single-cell
and spatially resolved atlas of human breast cancers. *Nat Genet.* 2021;53(9):1334-47.
53. Tanioka M, Fan C, Parker JS, Hoadley KA, Hu Z, Li Y, et al. Integrated Analysis of
RNA and DNA from the Phase III Trial CALGB 40601 Identifies Predictors of Response
to Trastuzumab-Based Neoadjuvant Chemotherapy in HER2-Positive Breast Cancer. *Clin*
Cancer Res. 2018;24(21):5292-304.

54.   Zawistowski JS, Bevill SM, Goulet DR, Stuhlmiller TJ, Beltran AS, Olivares-Quintero
678   JF, et al. Enhancer Remodeling during Adaptive Bypass to MEK Inhibition Is Attenuated
679   by Pharmacologic Targeting of the P-TEFb Complex. *Cancer Discov.* 2017;7(3):302-21.
680   55.   Angus SP, Stuhlmiller TJ, Mehta G, Bevill SM, Goulet DR, Olivares-Quintero JF, et al.
681   FOXA1 and adaptive response determinants to HER2 targeted therapy in TBCRC 036.
682   *NPJ Breast Cancer.* 2021;7(1):51.
683   56.   East MP, Sprung RW, Okumu DO, Olivares-Quintero JF, Joisa CU, Chen X, et al.
684   Quantitative proteomic mass spectrometry of protein kinases to determine dynamic
685   heterogeneity of the human kinome. *bioRxiv.* 2024.