1   Kinase Plasticity with Vandetanib Treatment Enhances Sensitivity to Tamoxifen in Estrogen
2   Receptor Positive Breast Cancer
3

Austin A. Whitman[1]†, Rasha T. Kakati[1†], Santiago Haase[1†], Christine HNC Thai[1,] Attila T.
Szenasi[1], Elizabeth C. Brunk[1,2], Denis O. Okumu[2], Michael P. East[2], Sarahi G. Molina[3],
Adriana S. Beltran[1,3], Jesse R. Raab[1,3], Lisa A Carey[1,5], Charles M. Perou[1,3,4,6], Gary L.
Johnson[1,2], and Philip M. Spanheimer[1,3,7*]

[1]Lineberger Comprehensive Cancer Center, University of North Carolina, Chapel Hill, NC,
USA
[2]Department of Pharmacology, University of North Carolina, Chapel Hill, NC, USA
[3]Department of Genetics, University of North Carolina, Chapel Hill, NC, USA
[4]Department of Pathology and Laboratory Medicine, University of North Carolina at Chapel
Hill, Chapel Hill, NC, USA
[5]Department of Medicine, University of North Carolina, Chapel Hill, NC, USA
[6]Computational Medicine Program, University of North Carolina, Chapel Hill, NC, USA
[7]Department of Surgery, University of North Carolina, Chapel Hill, NC, USA
†AAW, RTK, and SH contributed equally to this manuscript

Running title: Vandetanib ER+ Breast Cancer
19
20   Keywords: Breast cancer, endocrine resistance, tyrosine kinase inhibition, functional
21   proteomics, targeted therapeutics
22

Financial Support: This work was supported in part by the National Institutes of Health grants
K08CA280388 (PMS), R37CA292075 (PMS), P30CA016086 (PMS, LAC, CMP),
P50CA058223 (LAC,CMP), a Young Investigator Award from the Society of Surgical
Oncology (PMS), Susan G. Komen Foundation Grant (LAC, CMP), Breast Cancer Research
Foundation Grant (LAC, CMP)

Correspondence should be addressed to:
Philip M. Spanheimer MD
Lineberger Comprehensive Cancer Center
University of North Carolina at Chapel Hill
170 Manning Drive, Suite 1149
Chapel Hill, NC 27599-7213 USA
Phone: 919-966-5221; Fax: 919-966-8806
E-mail: Philip_Spanheimer@med.unc.edu

Disclosure of conflict of interests: C.M.P is an equity stockholder and consultant of
BioClassifier LLC; C.M.P is also listed as an inventor on patent applications for the Breast
PAM50 Subtyping assay. The remaining authors declare no potential conflicts of interest.

41
42

ABSTRACT

Resistance to endocrine therapy (ET) is common in estrogen receptor-positive (ER+) breast

cancer. Multiple studies have demonstrated that upregulation of MAPK signaling pathways

contributes to ET resistance. Herein we show that vandetanib treatment suppresses MAPK

signaling and enhances sensitivity to ET across ET-sensitive and ET-resistant ER+ cell lines

and patient derived organoids. Vandetanib treatment reprograms transcription toward a less

proliferative, more estrogen responsive, Luminal-A like state by enriching ERα chromatin

binding at canonical estrogen response elements. Multiplexed kinase inhibitor beads-mass

spectrometry (MIB/MS) revealed kinase network reprogramming, including upregulation of

PI3K and HER2 activity, as shared adaptive resistance mechanisms to vandetanib treatment.

Co-treatment with the HER2 inhibitor lapatinib, further enhanced sensitivity to vandetanib.

Using an operating room-to-laboratory short-term *ex-vivo* assay coupled to single-cell RNA

sequencing, we demonstrate conserved gene expression changes in primary tumor cells,

including increased HER2 activity signatures, following vandetanib treatment. Vandetanib

sensitivity signatures were generated from cell line and primary human tumor cells which

correlate with vandetanib sensitivity in ER+ patient-derived organoid and xenograft models.

Future clinical trials of vandetanib in ER+ breast cancer should include rationally designed

co-treatments based on adaptive resistance pathways, including HER2, and evaluate response

signatures as biomarkers predicting patients most likely to benefit.

SIGNIFICANCE

Vandetanib enhances sensitivity to tamoxifen in ER+ breast cancer by reprograming ERα

gene regulation and kinase signaling networks which define gene expression signatures

associated with response and identify targetable adaptive resistance pathways.

INTRODUCTION

Approximately 75% of breast cancers are estrogen receptor-positive (ER+) and fall within the luminal subtypes (1-3). Most patients with ER+ breast cancer receive endocrine therapy (ET) targeting ER. However, resistance to ET occurs in about 1/3 of early cancers and eventually develops in all metastatic ER+ cancers(4-6). Targeted inhibitors to PI3K, mTOR, and CDK4/6 have shown efficacy in patients with ET-resistant disease (7-10), but responses are often short lived, and many patients remain in need of durable, effective therapy.

Activation of the ERK/MAPK pathway is a well-established mechanism by which ER+ breast tumors resist ET (11-17). Gain-of-function mutations in ERK/MAPK nodes or upstream receptor tyrosine kinase (RTK) are enriched in ET-resistant tumors, although these changes occur in only ~13% of cases (11). ERK/MAPK can be activated by upstream recpetors, including the RET receptor tyrosine kinase, which is expressed in ~50% of ER+ breast tumors and is more highly expressed in ET-resistant tumors (18-21). Co-expression of RET and its activating ligand, glial derived neurotropic factor (GDNF), is associated with transciptional signatures of ERK/MAPK activation and with worse outcomes in human ER+ breast tumors (22). Further, RET is overexpressed with ESR1 (encoding ER) activating mutations, and inhibiting RET in ESR1 mutant models suppresses tumor growth (23).

Our prior work showed that tyrosine kinase inhibitors (TKIs) that inhibit RET, such as vandetanib and sunitinib, reduce proliferation and enhance ET sensitivity in ER+ breast cancer cell lines (24, 25). In RET-positive primary breast tumors, *in vitro* TKI treatment reduced ERK activity (26). These findings support a role for TKIs to inhibit ERK/MAPK activation to improve response to ET.

Despite encouraging preclinical data, clinical trials with vandetanib have had

disappointing results. However, these trials failed to account for potentially predictive

biomarkers such as ER and RET or evaluated vandetanib only as a monotherapy. For

example, a phase II trial of single-agent vandetanib in 46 patients with metastatic breast

cancer did not demonstrate a benefit (27), but did not report receptor status, limiting the

interpretation of whether this trial included patients that mechanistically would be predicted

to respond. Our work, including the findings presented herein, suggests that vandetanib

may provide a benefit for select patients. Recent TKI trials enrolling only ER+ patients

have shown promise (28, 29), renewing interest in TKI and highlighted the need for

biomarkers to identify patients most likely to respond to TKI therapy. To develop biomarkers

of response and to elucidate bypass resistance mechanisms, we sought to investigate how

vandetanib remodels gene expression and kinase signaling pathways in ER+/HER2- breast

cancer.

MATERIALS AND METHODS

Cell Lines

We obtained ER+ breast cancer cell lines parental MCF7 (RRID: CVCL_0031), tamoxifen-resistant MCF7 (MCF7-TAM) (RRID:CVCL_1D43), exemestane-resistant MCF7 (MCF7-EXE) (RRID:CVCL_M436), parental T47D (RRID:CVCL_0195), and tamoxifen-resistant T47D (T47D-TAM) (RRID:CVCL_1D36), from the American Type Culture Collection (ATCC). Cell lines were cultured using Dulbecco's Modified Eagle Medium (DMEM) (Gibco) with 10% fetal bovine serum (Gibco) and 1% penicillin-streptomycin (Life Technologies) at 37°C and 5% CO_2. MCF7-TAM and T47D-TAM were maintained by chronic treatment with 1 μM of 4OH-Tam and MCF7-EXE with 0.1 μM of exemestane.

Chemicals and Treatment

Vandetanib (SelleckChem) and z-4-Hydroxytamoxifen (4OH-Tam) (SelleckChem) were dissolved in dimethyl sulfoxide (DMSO). Pralsetinib was obtained from Chemietek, and lapatinib from Medchem Express LLC. Treatments were performed for 24 hours unless otherwise specified.

Cell Viability Assays

Cells were plated in opaque-walled 96-well plates at a density of 4,000 cells per well in technical triplicates and allowed to adhere overnight before the addition of the specified inhibitors. Viability was assessed using the CellTiter-Glo (CTG) Luminescent Cell Viability Assay (Promega). Luminescence was measured using a Synergy 2 microplate reader. Samples were averaged across biological triplicates. Statistical analysis and plotting were performed using GraphPad Prism 9.

Western Blots

Cells were lysed in Pierce RIPA buffer containing Halt protease inhibitor cocktail and PhosStop phosphatase inhibitor (Thermo Scientific). Proteins were resolved by SDS-polyacrylamide gel electrophoresis, transferred to PVDF membrane, and blocked with 10% BSA in PBS, then incubated overnight with the indicated primary antibody in blocking buffer. Membranes were washed three times in Tris-buffered saline containing 0.1% tween-20 then incubated for 1 hour with HRP-conjugated secondary antibodies. Blots were developed using a ChemiDoc-touch digital imager (Bio-Rad) and band intensity quantified using ImageJ (https://imagej.net/ij/index.html). The antibodies included: p44/42 MAPK (ERK1/2) (Cell Signaling Technology Cat# 9107, RRID:AB_10695739), Phospho-p44/42 MAPK (Erk1/2) (Cell Signaling Technology Cat# 75165, RRID:AB_2728835), RET (Abcam Cat# 3454-1, RRID:AB_10639232), and β-Actin (Santa Cruz Biotechnology Cat# sc-47778, RRID:AB_626632).

Patient Derived Organoid Derivation

Samples were obtained from patients with breast cancer who were undergoing surgical resection. This was performed under an IRB-approved protocol with written informed consent to generate new patient-derived breast cancer models. Patient-derived organoids (PDOs) were established from human tumor tissue samples as previously described (30). Briefly, tumor tissues were finely minced and digested enzymatically with a collagenase–hyaluronidase mixture (STEMCELL Technologies, Cat. No. 07912) for 2–5 h at 37 °C to release viable cell clusters. Red blood cells were removed using ACK Lysing Buffer (Gibco, Cat. No. A10492-01). The resulting cell clusters were resuspended in growth factor–reduced Matrigel (Corning, Cat. No. 356231) and plated as 30 µL domes in prewarmed 6-well plates. After Matrigel polymerization, domes were overlaid with breast cancer organoid (BCO) medium composed of Advanced DMEM/F-12 (Gibco, Cat. No. 12634010)

supplemented with 1X GlutaMAX (Gibco, Cat. No. 35050061), 10 mM HEPES (Gibco, Cat.

No. 15630080), 1X Penicillin–Streptomycin (Gibco, Cat. No. 15140122), 50 µg/mL

Primocin (InvivoGen, Cat. No. ant-pm-05), 1X B27 Supplement (Gibco, Cat. No. 17504001),

5 mM Nicotinamide (Thermo Fisher Scientific, Cat. No. 128271000), 1.25 mM N-acetyl-L-

cysteine (MilliporeSigma, Cat. No. A9165), 250 ng/mL R-spondin-3 (PeproTech, Cat. No.

120-44-100UG), 5 nM Heregulin β1 (PeproTech, Cat. No. 100-03-100UG), 100 ng/mL

Noggin (PeproTech, Cat. No. 120-10C-100UG), 20 ng/mL FGF-10 (PeproTech, Cat. No.

100-26-100UG), 5 ng/mL FGF-7/KGF (PeproTech, Cat. No. 100-19-100UG), 5 ng/mL EGF

(PeproTech, Cat. No. AF-100-15-100UG), 500 nM A-83-01 (Selleck Chemicals, Cat. No.

S7692), and 500 nM SB202190 (Selleck Chemicals, Cat. No. S1077). Y-27632 (Selleck

Chemicals, Cat. No. S1049; 5 µM) was included for the first three days of culture to enhance

cell survival. Culture medium was refreshed every 3–4 days.

PDOs were passaged every 2–4 weeks. Matrigel domes were collected, mechanically

disrupted, and incubated with TrypLE™ Express (Gibco, Cat. No. 12604013) for 7–10 min at

37 °C. The enzyme was neutralized with soybean trypsin inhibitor (Gibco, Cat. No.

17075029). Cells were then washed with Advanced DMEM/F-12, resuspended in cold

Matrigel, and replated to generate new organoids. After each passage, dissociated cells were

plated for immunofluorescence.

Immunofluorescence Staining

PDOs were dissociated, and single cells were seeded in 96-well plates for 5–24 h.

Cells were fixed with 4% paraformaldehyde (Electron Microscopy Sciences, Cat. No. 15710)

in PBS for 15 min at room temperature, then washed three times with PBS. Samples were

permeabilized with 0.3% Triton™ X-100 (MilliporeSigma, Cat. No. T8787) in PBS for 15

min. Non-specific binding was blocked with 5% bovine serum albumin (BSA;

MilliporeSigma, Cat. No. A9647) in PBS for 1–2 h at room temperature. Primary antibodies

were diluted in 1% BSA/0.1% Triton X-100/PBS and incubated overnight at 4 °C: anti-

Estrogen Receptor α (ERα; Abcam, Cat. No. ab32063, RRID: AB_732249) and anti-ERα,

clone SP1 (Thermo Fisher Scientific, Cat. No. RM-9101-S, RRID: AB_720919), each at

1:200. The following day, cells were washed three times (5 min each) in 1% BSA/0.1%

Triton X-100/PBS and incubated for 1 h at room temperature in the dark with Alexa Fluor™–

conjugated secondary antibodies (Thermo Fisher Scientific, Cat. Nos. A-21236 and A-21244,

RRIDs: AB_2535805 and AB_2535812, respectively) at 1:500 dilution in blocking buffer.

Nuclei were counterstained with DAPI (1:5000 from a 10 µg/mL stock; Thermo Fisher

Scientific, Cat. No. D1306) in PBS. After three final PBS washes, cells/organoids were

maintained in PBS or mounting medium and imaged using an EVOS™ Auto fluorescence

microscope. Secondary-only controls were included to verify signal specificity.

Drug Treatment and Viability Assay

For drug treatment, organoids were dissociated into single cells by mechanical

disruption of Matrigel domes, followed by incubation with TrypLE™ Express (Gibco, Cat.

No. 12604013) for 8 min at 37 °C. The enzyme was neutralized, and cells were pelleted by

centrifugation at $300 \times g$ for 5 min, then resuspended in ice-cold growth factor–reduced

Matrigel (Corning, Cat. No. 356231). For each well, 5,000 viable cells were seeded in 10 µL

Matrigel and plated in the center of a white-walled 384-well tissue culture plate to form a

dome. Plates were incubated at 37 °C for 15–30 min to allow Matrigel polymerization, then

40 µL of prewarmed BCO medium was gently added to each well. Drug treatments were

initiated at 72 h after seeding. Organoids were exposed to varying concentrations of

tamoxifen or vandetanib, either alone or in combination, with vehicle control containing

dimethyl sulfoxide (DMSO; Sigma-Aldrich, Cat. No. D2650). Each treatment was performed

in technical and biologic triplicate. Cell viability was assessed 72 h post-treatment using the

CellTiter-Glo 3D Cell Viability Assay (Promega, Cat. No. G9682, RRID: SCR_006724).

Plates were shaken on an orbital shaker for 30 min at room temperature. Luminescence was

measured using a microplate reader (BMG Labtech CLARIOstar), and background signal

from Matrigel-only wells was subtracted.

RET CRISPR Knockout

Two sgRNAs targeting exon 3 of RET (5'-ACTGGCACTCGCCCTCACGA-3' and

5'-GGGGTCGGTTCTCCCGAATG-3'), according to UCSC Genome Browser, were cloned

into the pSpCas9(BB)-2A-Puro(PX459) V2.0 plasmid (Addgene) and verified by Sanger

sequencing. MCF-7 cells were cultured in antibiotic-free media and grown to 70-90%

confluency. Cells were then transfected with sgRNA plasmids using FuGENE 4K

Transfection Reagent according the manufacturer's recommendations. After 72 hours, cells

were passaged onto 15-cm plates at a low density and selected with 1 μg/mL puromycin

(Gibco) for 3 days. Cell colonies were then isolated and genomic DNA was purified using the

Quick-DNA Miniprep Kit (Zymo Research). The targeted region was then amplified by PCR

and sequenced confirming frameshift mutations resulting in the introduction of premature

stop codons in both sgRNA-targeted alleles.

Synergy Analysis

Dose–response curves for tamoxifen, vandetanib, and vandetanib plus 0.5 μM tamoxifen

were fitted using a four□parameter logistic model (4PL). From each curve, we extracted the

drug concentration required to reach 50% inhibition.

Combination Index (CI50)

$CI50 = (D1 / Dx1) + (D2 / Dx2)$

D1 = vandetanib concentration in the combination

223 Dx1 = vandetanib concentration alone

224 D2 = fixed tamoxifen dose (0.5 μM)

225 Dx2 = tamoxifen concentration alone at 50% inhibition

226 CI < 1 indicates synergy, CI ≈ 1 additivity, CI > 1 antagonism.

227 Sensitization Ratio (SR50)

228 SR50 = Dx1 / D1

229 SR > 1 indicates increased vandetanib potency in the presence of tamoxifen.

230 **mRNA Sequencing and Data Analysis**

231  Cells were treated with 10 µM vandetanib or vehicle for 24 hours in biological

232 triplicates and total RNA was isolated (Qiagen). mRNA sequencing libraries were made

233 using TruSeq RNA Library Prep Kit (Illumina) and sequenced using the Illumina

234 NextSeq2000. Plots and statistical analyses were performed using RStudio 2022.07.1 Build

235 554. Differential gene expression (DGE) was determined using the DESeq2 R package using

236 raw mRNA-seq counts from STAR/Salmon. Genes with less than 10 counts across all

237 samples were omitted before analysis and counts were filtered and normalized to the upper

238 quartile using NOISeq. Significant differentially expressed genes were defined by using a

239 Log2FC cutoff of +/- 0.5 and a false discovery rate (FDR) < 0.05. Enrichment of gene sets

240 was investigated through hypergeometric analysis of an unranked list of upregulated or

241 downregulated elements using the enricher function of the clusterProfiler package.

242 METABRIC and SCAN-B databases were obtained for survival analysis and filtered for

243 ER+/HER2- patients treated with ET. A median-centroid approach was implemented to

244 assign a gene set score per patient. Kaplan-Meier (KM) plots were then computed using the

245 survminer and survival packages.

246 **Cleavage Under Targets and Release Using Nuclease (CUT&RUN) and Analysis**

Cut and Run was performed with the EpiCypher Cutana CUT&RUN Kit Version 3

according to the manufacturer's manual v3.5. Cells were treated with 10 µM vandetanib or

vehicle for 24 h, and cell counts were quantified using the Countess. 500,000 viable cells

were bound to ConA beads per reaction, and 0.5 ug (or manufacturer's recommendation) of

antibody was added to each sample overnight. pAG-MNase was then used to cleave DNA,

which was extracted and used to generate libraries following the EpiCypher Cutana

CUT&RUN Library Prep Kit. Libraries were sequenced on the NextSeq2000 (Illumina)

platforms with paired-end sequencing and dual indexing, according to the manufacturer's

protocol.

Cut and Run sequencing results were processed through the Nextflow DSL2

implementation of the Raab Lab CUT&RUN processing pipeline (https://github.com/raab-

lab/cut-n-run). The pipeline produced peak filtered calls and BAM coverage files, which were

used to run differential peak binding between vandetanib versus vehicle treated samples for

ERα, FOXA1 and H3K4me3 separately. For this, a union peak set was generated from the

vehicle and vandetanib samples (with 2 replicates each) using bedtools merge function (31).

Peak count with IgG subtraction was performed on the union peakset by subtracting

the corresponding IgG sequencing controls for Vehicle and Vandetanib-treated samples.

Counts were library-size normalized with DiffBind dba.normalize(), and differential peak

analysis was performed with DiffBind dba.analyze() with EdgeR (32). Gene annotations were

added to the differential peak analysis using ChIPseeker annotatePeak()(33, 34). To analyze

peak gene signature enrichment, the binding fold changes for all the peaks associated with a

signature were computed (vandetanib vs vehicle) and a Wilcoxon test was performed to

determine significant differential deposition for the given signature.

To illustrate the global shifts in ERα, FOXA1 and H3K4me3 deposition, we extracted the Diffbind EdgeR top upregulated and downregulated differential peaks (vandetanib vs vehicle) and generated peak profile heatmaps from normalized coverage plots using deepTools computeMatrix and plotHeatmap functions (35). For significant genes and peaks, coverages plots were produced using library size normalized coverage plots with Gviz GeneRegionTrack (36). To illustrate peak gene distribution, we used the CnRAP (Cut & Run Analysis Pipeline) step 4 script, which generates peak gene distribution images from peak annotations using a modified CUT&RUN analysis pipeline to identify transcription factor binding sites in human cell lines (37). The CUT&RUN quality assessment through peak profile heatmaps, transcriptional start site distribution and peak gene distribution are in Fig 5E-G.

MIB/MS and Data Analysis

Cells were treated with vandetanib at its IC90 for 48 h. Cells were harvested by scraping and cell pellets obtained for proteomic analysis using the MIB/MS technology (38, 39). Scaled label free quantification (LFQ) values were obtained and a correlation plot of samples was computed to assess homogeneity of triplicates and heterogeneity of treatment conditions for quality control using ggplot2. Differential protein expression (DPE) was analyzed using the limma R package using scaled LFQ values. Significant differentially expressed proteins were defined using a Log2FC (log2 fold change) cutoff of 0.5 for upregulation and -0.5 for downregulation, along with a p-value < 0.05.

Primary ER+/HER2- Human Tumor Cells

A primary tumor sample was obtained from a patient with ER+/HER2- invasive ductal carcinoma. The tumor was processed fresh as previously described (40). In brief, tumor tissue was transported to the laboratory immediately after surgical resection,

dissociated into a single cell suspension, and treated with vandetanib (10 µM) or control

media for 12 hours. After treatment, single cell RNA sequencing libraries were made using

the 10x Genomics Chromium Next GEM Single Cell 3' v3.1 and sequenced on the

NextSeq2000 (Illumina) with pair-end sequencing.

Single Cell RNA Sequencing Analysis

Reads were aligned and counted using Cellranger count function with default

parameters. The counts and features matrixes were used to generate Seurat objects. The

Seurat objects corresponding to each treatment group were merged with Seurat's function

merge (), and low quality cells were removed. Doublets were removed using scDblFinder

(41) and cells were filtered: nFeature_RNA > 200 & nFeature_RNA < 6000 & percent.mt <

10. Cell types were annotated on the combined Seurat object using SingleR using the cell

type atlas HumanPrimaryCellAtlasData () accessed with the package celldex. InferCNV was

used   to   predict   copy   number   alterations,   defining   the   immune   cells

("T_cells","Macrophage","DC","Monocyte","NK_cell") as reference.

Differential analysis of gene expression was performed on the tumor cells to compare

vandetanib-treated versus non-treated cells using findmarkers (). GSEA was then performed

using the list of differentially expressed genes, ranking the genes according to log2(FC)/FDR,

and the ranked gene lists were used in the GSEA software (42). Gene signatures were scored

using AddModuleScore_UCell() from the UCell package.

We used the NMF R package to build a non-negative matrix factorization (NMF)

model of the tumor cells using a list of genes differentially expressed upon vandetanib

treatment in MCF7 cells from this study. The Seurat object was subset to contain only the

above-mentioned genes, and the NMF model was fitted to 8 factors. Heatmaps were

generated for each treatment group to assess enrichment of cells to different NMF factors.

Each cell was assigned to a unique NMF according to the NMF with maximum coefficient

value for that cell, and the percentage of cells belonging to each NMF factor was calculated

for each treatment group. DE analysis was performed comparing cells in certain NMF

factors.

Patient Derived Xenografts

Two ER+ patient-derived xenografts (PDX) were obtained as a generous gift from Dr.

Alana Welm (43). PDXs were orthotopically injected into athymic mice that had been

implanted with estrogen releasing pellets (Innovative Research of America, NE-121). When

tumors were 0.5 cm in greatest dimension, five mice were randomized to each of the

following treatments: control gavage, tamoxifen (50 mg/kg daily), or vandetanib (25 mg/kg

daily). Tumor size was measured three times per week using calipers. After 6 weeks, mice

were sacrificed and tumors were collected for mRNA sequencing.

Statistics

All experiments were performed with at least 3 biological or technical replicates.

Small sample number comparisons were analyzed with an unpaired t test with Welch's

correction for non-equal SD. Linear regressions and nonlinear regression curves were

compared by statistical tests on the parameters of the curves using the extra sum of

squares F test. Experiments with large sample numbers were analyzed by Wilcoxon test.

Data Availability

Single cell and bulk RNA sequencing data was made available at the Sequence Read

Archive (SRA) under BioProject PRJNA1177387. Mass spectrometry protein sequencing

results are included in this manuscript as Supplementary Table 1.

Code Availability

All original code has been deposited on the Github, and it is publicly available at the

Github repository at https://github.com/santiagohaase/Spanheimer-lab-Kinase-Plasticity-in-

Response-to-Vandetanib-Enhances-Sensitivity-to-Tamoxifen

RESULTS

Vandetanib Inhibits ERK Activation and Viability in ER+ Breast Cancer Cells

Vandetanib inhibits receptor tyrosine kinases (RTK) including RET and EGFR which are upstream activators of ERK/MAPK (Fig. 1A). We first investigated if vandetanib inhibited the ERK/MAPK pathway using MCF7 and two ET-resistant derivatives, MCF7-TAM (tamoxifen) and MCF7-EXE (exemestane), which had been cultured long-term under drug treatment to develop resistance. Vandetanib treatment reduced both the cellular protein levels of ERK and levels of activating phosphorylation of ERK (Fig. 1B). To account for changes in total ERK, we calculated the ratio of pERK/ERK signal intensity, which was reduced in all 3 cell lines with vandetanib treatment. Dose-response curves demonstrated dose IC50 values for vandetanib of 8.5 μM in MCF7, 10.4 μM in MCF7-TAM, and 15.9 μM in MCF7-EXE (Fig. 1C). Similar vandetanib sensitivities were seen in T47D and T47D-TAM (Fig. S1).

Vandetanib Sensitizes ER+ Breast Cancer Cells to Tamoxifen

We next tested if co-treatment with vandetanib enhanced the antiproliferative effects of tamoxifen in ET-resistant cells. The IC50 of the active metabolite of tamoxifen, 4OH-Tam, was 10.7 μM in MCF7, 15.8 μM in MCF7-TAM, and 14.1 μM in MCF7-EXE (Fig. 1D) demonstrating decreased sensitivity to tamoxifen with chronic ET. Treatment with 10 μM vandetanib resulted in a left shift of the 4OH-Tam sensitivity curve in all 3 cell lines with the IC50 for 4OH-Tam of 3.8 μM in MCF7, 4.8 μM in MCF7-TAM, and 4.8 μM in MCF7-EXE. We produced an additive effect isobole line by connecting the individual tamoxifen and vandetanib concentrations that caused the survival percentage equal to the drug combination. The point indicating the combination of 2 μM of vantetanib with 5 μM of tamoxifen is below

the additive isobole line, demonstrating a synergic effect for the MCF7, MCF7-TAM, and

MCF7-EXE (Fig. 1E).

Vandetanib Response in Breast Cancer Patient Derived Organoids

We derived a panel of primary breast tumor patient-derived organoids (PDOs) to

investigate the impact of vandetanib across a diverse group of translationally relevant breast

cancer models. Patient and tumor characteristics are listed with intrinsic subtype assigned

from RNA-seq (Fig. S2A). Of the 6 ER+/HER2- PDOs, 2 were LumA subtype, 1 was LumB,

1 was HER2E, and 2 were basal-like subtype, recapitulating the molecular heterogeneity of

ER+ disease. The TNBC derived PDO (T4) had a basal molecular subtype. PDOs displayed a

range of sensitivity to tamoxifen (Fig. S2B), with ER-negative models, including the

organoid derived from the TNBC cell line Sum149, having the lowest sensitivity. PDOs were

further validated for expression of the estrogen receptor (ER), which displayed heterogeneity

in ER expression within and across PDO, and the epithelial marker EPCAM (Fig. S2C). PDO

displayed a range of sensitivity to vandetanib, with greater sensitivity in ER+ PDOs (Fig. 1F).

To analyze the effect of the drug combination, we calculated the combination index (CI) and

sensitization index (SR) at 50% of survival. To represent both the interaction (CI) and the

magnitude of sensitization (SR), we plotted SR versus CI for each cell model at the 50%

effect level. Low CI and high SR values indicate strong synergy. Synergy analysis

demonstrated that 5 ER+ PDOs displayed synergy with dual treatment of vandetanib and

tamoxifen, and this included 3 of 4 luminal intrinsic subtype (Fig. 1G). These results show a

high vandetanib-tamoxifen synergy rate in PDOs that retain ER expression and luminal

transcriptional profile.

RET Regulates ERK Activity and Vandetanib Response in ER+ Breast Cancer Cells

Based on our prior work using a transient knockdown model showing a role for RET

in mediating vandetanib response and driving poor outcomes and ET resistance in ER+ breast

cancer (22), we explored the role of RET in vandetanib sensitivity. Transient RET

antagonism with the selective RET inhibitor BLU667 resulted in reduced sensitivity to

vandetanib in MCF7 cells in dose-dependent fashion (Fig. 2A). BLU667 also reduced

vandetanib sensitivity in MCF7-TAM (Fig. S3A), but not in the low RET expressing MCF7-

EXE (Fig. S3B).

To precisely elucidate the role of RET in ER+ breast cancer cells using a more robust

experimental model than prior work transient siRNA knockdowns, we generated CRISPR

knockouts of RET in MCF7 cells (RET KO#1 and RETKO#2). Western blot confirmed

knockout of RET protein (Fig. 2B). Both knockouts had reduced pERK levels compared to

wild type (WT), demonstrating that RET is a key regulator of this pathway. RET KO cells

also had reduced ER protein levels. Further supporting a key role for RET as a mediator of

vandetanib response, both KO lines were significantly less sensitive to vandetanib compared

to WT (Fig. 2C). Based on the vandetanib synergy assay we hypothesized that RET KO

would sensitize cells to tamoxifen, however, KO cells were less sensitive (Fig. 2C), likely

because of reduced ER expression.

We further explored the molecular basis of this unexpected finding by comparing

gene expression of RET KO vs WT cells using RNA sequencing. Differentially expressed

gene analysis revealed many genes significantly regulated by RET expression (Fig. 2D) with

RET KO. Hallmark gene sets for KRAS signaling and RET knockout were lower with RET

KO (Fig. 2E), as expected. Consistent with reduced sensitivity to tamoxifen, estrogen

response genes were reduced, indicating that RET KO cells have reduced ER activity. RET

KO resulted in lower proliferation scores compared to WT (Fig. 2F) and RET KO cells had

significantly reduced growth vs WT. Cumulatively, these findings support that RET promotes

an ERK-driven proliferative phenotype and mediates vandetanib response, but stable

knockout is associated with reduced expression of ER and estrogen response genes.

Vandetanib Induces Low-Proliferative and Estrogen Responsive Gene Expression

To delineate the transcriptional basis for how vandetanib enhances tamoxifen

sensitivity in ER+ breast cancer cells, we performed mRNA sequencing after treatment with

vandetanib (10 μM) compared to vehicle. Heatmaps of DEGs are shown for MCF7 (Fig. 3A),

MCF7-TAM (Fig. 3B), and MCF7-EXE (Fig. 3C). In MCF7, downregulated genes were

enriched for MAPK signaling and proliferative signatures (Fig. 3D), consistent with loss of

pERK observed in Fig. 1A. Upregulated genes are enriched for signatures of estrogen

response and sensitivity to tamoxifen, consistent with observed sensitization to tamoxifen

(Fig. 1D). Up-regulated and down-regulated gene sets in MCF7-TAM (Fig. 3E) and MCF7-

EXE (Fig. 3F) showed similar depletion of proliferative and MAPK signatures and

enrichment of estrogen and tamoxifen response signatures. RAS signaling signatures were

depleted with vandetanib treatment in all 3 cell lines (Fig. 3G). Similar transcriptional effects

were seen in T47D and T47D-TAM (Fig. S4).

Enriched estrogen response signatures and reduced proliferation are hallmarks of the

ET responsive Luminal A (LumA) molecular subtype that is associated with the most

favorable patient outcomes(44). We analyzed the effect of vandetanib on genes that define

the LumA subtype on the PAM50 assay (LumA centroid). This analysis demonstrated that

vandetanib treatment resulted in a stronger LumA correlation supporting that treatment

induced a more LumA-like indolent and ET responsive transcriptional phenotype (Fig. 3H).

To elucidate conserved molecular actions of vandetanib we determined overlap across

the parental and ET-resistant MCF7 cell lines (Fig. 2I). Notably, significant commonly

upregulated genes, including *SELENBP1* and *BAG1*, which are associated with reduced

proliferation and improved survival in ER+ breast tumors (45-47). In the common

downregulated genes, *AURKA* is associated with poor survival in ER+ breast cancer patients

(48). Reduced expression of *ECT2* is associated with abrogation of MAPK signaling (49).

Finally, several key regulators of cell cycle progression associated with worse outcomes and

poor response to endocrine therapy in ER+ breast cancer are downregulated, including

CCND1, *TOP2A*, and *CCNB1* (Fig. 2J).

To test if the conserved vandetanib down-regulated gene set is associated with

prognosis in patients with ER+/HER2- breast cancer treated with endocrine therapy, we

obtained survival annotated transcriptional data from METABRIC(50) and SCAN-B(51).

Tumors were stratified by gene set score using median thresholding. METABRIC patients

with low expression of genes that were downregulated by vandetanib had significantly

improved OS (HR=0.75, p<0.0001, Fig. 3K) and RFS (HR=0.60, p<0.0001, Fig. 3L).

Similarly, SCAN-B ER+ breast cancer patients with low expression of vandetanib

downregulated genes demonstrated significantly lower mortality compared to high signature

score (HR=0.87, p=0.034) (Fig. 3M). We next obtained gene expression data from patients

with ER+/HER2- breast cancer profiled on ET and clinically annotated as sensitive or

resistant (12). Resistant tumors had significantly higher expression of the vandetanib

downregulated signature compared to sensitive tumors (4.3 vs 3.8, p<0.001, Fig. 3N).

Cumulatively, these findings support that in ER+ breast cancer, vandetanib treatment

promotes gene expression changes associated with lower proliferation, increased ET

response, and improved survival.

Vandetanib Treatment Enriches ERα Chromatin Occupancy at E2 Response Genes

Because vandetanib treatment resulted in enriched estrogen response signatures, we

performed CUT&RUN to evaluate how vandetanib alters ERα chromatin binding. We

performed peak differential analysis followed by gene peak annotation, comparing control

and vandetanib treated cells. Taking the annotated peaks for each gene, we calculated the

differential mean peak binding signals for gene signatures of interest. As expected, we found

that vandetanib treatment increases binding of ERα to genes associated with estrogen activity,

as well as genes associated with PI3K, STAT2 and ERBB2 signaling with vandetanib

treatment (Fig. 4A). Genes with depleted ERα binding are associated with cell cycle

transition, TNF and NOTCH pathway, among others.

To further characterize the changes on ERα occupancy induced by vandetanib, we

also assessed chromatin occupancy of FOXA1, a pioneer factor that mediates ERα binding to

chromatin sites and ERα co-localizing protein, and for H3K4me3, a histone mark that

characterizes transcriptional active sites and active promotor regulatory elements. FOXA1

showed increased occupancy to genes containing ERα binding sites but decreased occupancy

to gene signatures indicating ERα activity in vandetanib-treated cells (Fig. S5A) and this was

similar for H3K4me3 marks (Fig. S5B).

To assess the interaction of ERα, FOXA1 and H3K4me3 and the effect on expression

of ERα-response genes, we analyzed the concurrent peak occupancy with mRNA expression

in vandetanib-treated cells versus control (Fig. 4B). Generally, genes that gain ERα

occupancy also gain FOXA1 and H3K4me3 occupancy, and this is associated with increased

gene expression. GSEA of the genes with increased ERα, FOXA1 and H3K4me3 binding and

transcriptional activity on vandetanib-treated cells showed these genes are associated with

estrogen response (Fig. 4C). This reinforces that vandetanib leads to enriched ERα and

FOXA1 chromatin activity on key estrogen response genes. Furthermore, we found increased

ERα binding to ESR1 transcriptional start site and gene body peaks, indicating that

vandetanib promotes a positive auto-regulatory loop in ESR1 expression (Fig. 4D). We next

looked at enriched H3K4me3 sites with vandetanib treatment filtering for known promotor

activating loci to assess transcription factor motifs enriched at those sites. Both ER and

FOXA1 motifs were enriched at enhancer sites supporting a central role in those proteins

mediating the gene regulatory effects of vandetanib (Fig. 4E).

Functional Signatures of Vandetanib Response

Using multiplex inhibitor bead enrichment coupled to mass spectrometry (MIB/MS),

we investigated the reprogramming of kinase signaling networks in response to vandetanib

(38). In this technique, kinases are enriched by incubating cell extracts with type 1 kinase

inhibitors immobilized on Sepharose beads (Fig. 5A). Vandetanib treatment caused

depletion of known vandetanib targets (52) in all 3 models (Fig. 5B), despite large baseline

differences in activity, (Fig. 5C). Similar results were seen for T47D and T47D-TAM (Fig.

S6).

Enriched and depleted proteins by cell line are shown in Fig. 5D. Overlap analysis

showed that kinome dynamics were heterogenous. Common upregulated proteins included

AURKA, PLK1, PIP4K2A, which promote proliferation (53-55). Common depleted proteins

with vandetanib treatment included proliferation, promoting cell cycle regulators CDK1,

CDK2, and PKMYT1. Depletion of MAP2K2 (MEK2) supports downregulation of the

MAPK signaling pathway in response to vandetanib treatment, consistent with Fig. 1A. Gene

set enrichment analysis similarly demonstrated downregulation of the ERK cascade (56) with

largely consistent depletion of specific node proteins (Fig. 5E). PI3K/AKT pathway and

ERBB2 signaling were also upregulated. Interestingly these were two pathways identified

with enriched ERα chromatin occupancy (Fig. 4A), demonstrating the transcriptional basis

for protein signaling network adaptation.

Inhibition of Adaptive Resistance Pathways Enhances Vandetanib Efficacy

Upregulated functional kinases may identify mechanisms of adaptive bypass

resistance, so we assessed activity of druggable kinases (23) by treatment (Fig. 5F). We

performed co-treatment assays on the principle that inhibiting mediators of adaptive

resistance should enhance sensitivity to vandetanib. Because ERBB2/HER2 was enriched in

all 3 MCF7 cell lines, we co-treated cells with vandetanib and the FDA approved HER2

small molecule inhibitor lapatinib. Treatment with lapatinib resulted in significantly

augmented vandetanib response measured by reduction in cell viability in all 3 cell lines,

(Fig. 5G), validating increased reliance on HER2-activity in vandetanib treated cells.

Vandetanib Response at Single Cell Resolution in Primary ER+ Breast Cancer

We used our previously validated primary tumor *ex vivo* short-term drug treatment

platform coupled to single-cell RNA sequencing (40), to assess how primary ER+ breast

tumor cells respond to vandetanib treatment. Tumor cells were treated with control or

vandetanib containing media for 12 hours and assayed by scRNAseq. Uniform manifold

projection (UMAP) was performed using Seurat v3 (57). Cell populations were annotated

using known transcriptional profiles and canonical markers (Fig. 6A) and by treatment group

for tumor cells (Fig. 6B). Malignant (tumor) cells were distinguished from non-tumor

epithelial cells as cells having significant inferred copy number alterations by InferCNV

(58). Differentially expressed genes were identified using a false discovery rate (FDR)

cutoff of < 0.05 in the iCNV+ tumor cell compartment with vandetanib treatment

compared to control (Fig. 6C). Enriched and depleted pathways in vandetanib treated tumor

cells relative to control treated were determined (Fig. 6D). As was seen in cell lines, estrogen

response genes and signatures associated with tamoxifen sensitivity were enriched.

To demonstrate the in-human relevance of the cell line findings, we examined salient

features seen in cell lines in these primary tumor cells. First, we tested the overlapping

downregulated genes after vandetanib treatment across the 3 MCF7 cell lines (Fig. 3J) and

these genes were lower in vandetanib treated primary tumor cells (Fig. 6E). Depletion of

KRAS signature supports on-target effects. Next, LumA correlation score was assigned to

each tumor cell using a validated single cell PAM50 classifier (59) and treated tumor cells

had a stronger correlation to LumA. Thus, as with cell lines, short-term vandetanib treatment

alters the transcriptional profile of ER+/HER2- primary tumor cells towards a more indolent

and estrogen responsive phenotype. Lastly, ERBB2 signaling was enriched supporting that

primary human tumor cells also induce vulnerability to HER2 inhibitors in response to

vandetanib treatment.

To explore features of vandetanib response we used non-negative matrix factorization

(NMF) (Fig. S7A). This analysis yielded metagene modules that were enriched in vandetanib

treated samples (module 1) and depleted in vandetanib treated cells (Module 6) (Fig. S7B).

We next identified features that may be associated with vandetanib response by comparing

modules 1 and 6 gene expression profiles. Module 1 (vandetanib resistant) had expected high

expression of genes that increase with vandetanib treatment, consistent with poor response.

Module 6 was enriched for MYC targets, which is known to support tamoxifen resistance, but

may support a transcriptional profile of vandetanib sensitivity (Fig. S7C). Interestingly,

module 1 was enriched for hallmark estrogen response genes and our previously generated

single cell tamoxifen response score (40). Therefore, cells with poor vandetanib response

may have enhanced tamoxifen sensitivity (Fig. S7D) supporting dual treatment to target

distinct cell types within the tumor. Lastly, we looked at our human tumor derived signatures

of high GDNF/RET expressing human ER+-breast tumors (22). Genes downregulated in

GDNF/RET high human tumors had low expression in vandetanib sensitive cells, supporting

RET activity as a key feature of responsive cells (Fig. S7E).

559   Vandetanib in ER+/HER2- Breast Cancer Patient Derived Xenografts

Mice implanted with ER+/HER2- patient derived xenografts were treated with control

or vandetanib gavage. Tumor FFPE sections showed expected morphology and expression of

ER and lack of HER2 expression (Fig. 7A). For HCI011 vandetanib significantly reduced

tumor growth at 6 weeks (Fig. 7B). HCI05 tumors had no reduction in growth with

vandetanib, demonstrating a vandetanib sensitive (HCI011) and resistant (HCI05) model. As

expected, the sensitive model (HCI011) had higher expression of genes associated with RET

activity in human ER+ tumors, supporting that RET activity is also a key feature of

vandetanib response in PDX (Fig. 7C). Further, high expression of the vandetanib

downregulated genes in MCF7 was also seen in HCI011. Lastly, NMF1 module (Fig. S7),

which are genes highly expressed in vandetanib resistant human tumor cells were expressed

more highly in the resistant HCI05 PDX.

571   Vandetanib Response Signatures in PDX and PDO

We next used our PDO mRNA sequencing and sensitivity data (IC50) to test

performance of derived signatures on predicting vandetanib sensitivity in human tumor

derived models. We tested if 1) our TCGA human tumor derived signature of RET activity

(GDNF/RET up TCGA Signature) (22), 2) cell line derived signature of down-regulated

genes with vandetanib treatment in MCF7 (Fig. 3J)., and 3) a single cell human tumor

vandetanib response score of the difference between sensitive cell genes (NMF6) and

resistant cell genes (NMF1) (Fig. S7B/C) were associated with sensitivity to vandetanib. We

found that both the MCF7 down signature and the human tumor single cell response score

were inversely correlated with IC50 of ER+ PDO demonstrating that a high signature score is

associated with more sensitive PDO (Fig. 7D).

DISCUSSION

Resistance to endocrine therapy (ET) is a significant driver of poor patient outcomes, despite advances in understanding how ER+ breast tumors respond to therapy. The identification of MAPK pathway activation as a driver of resistance to ET has generated significant interest in targeting this pathway to enhance ET response, but this has not translated to the clinic. This highlights the need for clinically actionable kinase targeting strategies in MAPK-activated, ET-resistant breast tumors and biomarkers of patients most likely to respond.

Kinase signaling networks can be plastic and inhibiting one node can lead to adaptive bypass through another pathway. We identified PI3K and HER2 as potential mediators of adaptive resistance to vandetanib treatment, and targeting HER2 with lapatinib enhanced sensitivity to vandetanib. ERα occupancy was enriched at genes mediating PI3K and ERBB2 signaling upon vandetanib treatment, suggesting that ER rewiring may contribute to the adaptive kinase landscape. Activating HER2 could open a number of efficacious therapeutic options for patients. While HER2 activity was increased, there was no change in HER2 RNA expression, highlighting the need for protein activity level assessment to identify rationale cotreatments. To this end, our group has recently developed a method for comprehensive-kinome characterization on limited sample, such as a core needle biopsy, which could enable on-treatment assessment of adaptive responses and facilitate real-time, patient specific interactive cotreatment regimen design (60).

RET mediates effects of vandetanib and we found that RET-regulated genes are a key marker of response to vandetanib in patient derived models, but CRISPR knockout of RET unexpectedly resulted in reduced, rather than enhanced, sensitivity to tamoxifen. ER expression was reduced after RET knockout, as were ER induced genes, and this likely

accounts for decreased sensitivity to tamoxifen. While ER has 2 binding sites in the RET

promoter that activate RET expression (61), no reciprocal mechanism by which RET

regulates ER is known. RET expression and ER-activity are correlated, RET knockout may

select for cells that have lower baseline ER-activity by enriching populations that are less

dependent on RET signaling. Further studies are needed to model emerging resistance to RET

and ER inhibitors and to track relative dependencies on RET and ER within emerging

resistant tumor cell subpopulations.

Our human tumor single cell analysis reinforces key *in vitro* findings. RET activity is

a key component sensitivity to vandetanib in ER+ breast tumors. Cells categorized within the

vandetanib sensitive metagene module have low expression of genes that are downregulated

in GDNF/RET high human tumors, indicating that vandetanib sensitivity is associated with

our previously described high RET-activity phenotype (22). Importantly, we show that

vandetanib response is not homogeneous within tumors. Instead, distinct sensitive and

resistant populations can be resolved at single-cell resolution. Resistant cells express features

associated with tamoxifen response, consistent with a conceptual model where the population

that persists under vandetanib remains sensitive to ET. In contrast, a vandetanib sensitive

cells were enriched for MYC targets and programs linked to tamoxifen resistance, consistent

with a vandetanib-sensitive but ET-refractory state. These results indicate that within

heterogenous human tumors, vandetanib preferentially eliminates a MYC-high, ET-poorly

responsive compartment, while a more estrogen-responsive, tamoxifen-sensitive

compartment persists.

Similarly in PDX we identified a vandetanib-sensitive and a resistant ER+/HER2–

tumor. The sensitive PDX expressed a RET activity signature consistent with high RET-

activity phenotype, as well as the vandetanib-sensitivity gene programs identified in cell

lines, whereas genes defining the vandetanib-resistant human tumor cells were more highly

expressed in the resistant model. These results show that the same vandetanib-linked

transcriptional states identified in cell lines and primary tumor cells stratify response in

orthotopic PDX models, indicating that these states reflect true biological vulnerabilities

rather than model-specific features, and that these signatures could be used to as to predict

response to vandetanib.

In summary, we demonstrate that ER+ breast cancer cells become more sensitive to

ET with vandetanib treatment. Vandetanib enriches ER binding site occupancy at canonical

E2 response genes altering gene expression and kinase signaling resulting in a lower

proliferative, more estrogen-responsive luminal A-like state with an induced vulnerability to

HER2 inhibitors. We define response signatures that strongly correlate with sensitive and

resistant patient derived organoid and xenograft models. These results support that

vandetanib treatment could benefit select patients with ER+ breast cancer and that future

trials should evaluate response signatures as predictive biomarkers.

REFERENCES
645

1.   Nabholtz JM, Tonkin K, Smylie M, Au HJ, Lindsay MA, and Mackey J.
Chemotherapy of breast cancer: are the taxanes going to change the natural history of
breast cancer? *Expert Opin Pharmacother.* 2000;1(2):187-206.
2.   Mouridsen H, Gershanovich M, Sun Y, Perez-Carrion R, Boni C, Monnier A, et al.
Phase III study of letrozole versus tamoxifen as first-line therapy of advanced breast
cancer in postmenopausal women: analysis of survival and update of efficacy from
the International Letrozole Breast Cancer Group. *J Clin Oncol.* 2003;21(11):2101-9.
3.   Bachelot T, Bourgier C, Cropet C, Ray-Coquard I, Ferrero JM, Freyer G, et al.
Randomized phase II trial of everolimus in combination with tamoxifen in patients
with hormone receptor-positive, human epidermal growth factor receptor 2-negative
metastatic breast cancer with prior exposure to aromatase inhibitors: a GINECO
study. *J Clin Oncol.* 2012;30(22):2718-24.
4.   Harvey JM, Clark GM, Osborne CK, and Allred DC. Estrogen receptor status by
immunohistochemistry is superior to the ligand-binding assay for predicting response
to adjuvant endocrine therapy in breast cancer. *J Clin Oncol.* 1999;17(5):1474-81.
5.   Jeselsohn R, Barry WT, Migliaccio I, Biagioni C, Zhao J, De Tribolet-Hardy J, et al.
TransCONFIRM: Identification of a Genetic Signature of Response to Fulvestrant in
Advanced Hormone Receptor-Positive Breast Cancer. *Clin Cancer Res.*
2016;22(23):5755-64.
6.   Finn RS, Martin M, Rugo HS, Jones S, Im SA, Gelmon K, et al. Palbociclib and
Letrozole in Advanced Breast Cancer. *N Engl J Med.* 2016;375(20):1925-36.
7.   Baselga J, Campone M, Piccart M, Burris HA, 3rd, Rugo HS, Sahmoud T, et al.
Everolimus in postmenopausal hormone-receptor-positive advanced breast cancer. *N
Engl J Med.* 2012;366(6):520-9.
8.   Piccart M, Hortobagyi GN, Campone M, Pritchard KI, Lebrun F, Ito Y, et al.
Everolimus plus exemestane for hormone-receptor-positive, human epidermal growth
factor receptor-2-negative advanced breast cancer: overall survival results from
BOLERO-2dagger. *Ann Oncol.* 2014;25(12):2357-62.
9.   Andre F, Ciruelos E, Rubovszky G, Campone M, Loibl S, Rugo HS, et al. Alpelisib
for PIK3CA-Mutated, Hormone Receptor-Positive Advanced Breast Cancer. *N Engl J
Med.* 2019;380(20):1929-40.
10. Turner NC, Slamon DJ, Ro J, Bondarenko I, Im SA, Masuda N, et al. Overall
Survival with Palbociclib and Fulvestrant in Advanced Breast Cancer. *New Engl J
Med.* 2018;379(20):1926-36.
11. Razavi P, Chang MT, Xu G, Bandlamudi C, Ross DS, Vasan N, et al. The Genomic
Landscape of Endocrine-Resistant Advanced Breast Cancers. *Cancer Cell.*
2018;34(3):427-38 e6.
12. Xia Y, He X, Renshaw L, Martinez-Perez C, Kay C, Gray M, et al. Integrated DNA
and RNA Sequencing Reveals Drivers of Endocrine Resistance in Estrogen Receptor-
Positive Breast Cancer. *Clin Cancer Res.* 2022;28(16):3618-29.
13. Arpino G, Wiechmann L, Osborne CK, and Schiff R. Crosstalk between the estrogen
receptor and the HER tyrosine kinase receptor family: molecular mechanism and
clinical implications for endocrine therapy resistance. *Endocr Rev.* 2008;29(2):217-
33.

14.   Meijer D, Sieuwerts AM, Look MP, van Agthoven T, Foekens JA, and Dorssers LC.
Fibroblast growth factor receptor 4 predicts failure on tamoxifen therapy in patients
with recurrent breast cancer. *Endocr Relat Cancer.* 2008;15(1):101-11.

15.   Bergqvist J, Elmberger G, Ohd J, Linderholm B, Bjohle J, Hellborg H, et al.
Activated ERK1/2 and phosphorylated oestrogen receptor alpha are associated with
improved breast cancer survival in women treated with tamoxifen. *Eur J Cancer.*
2006;42(8):1104-12.

16.   Kirkegaard T, Witton CJ, McGlynn LM, Tovey SM, Dunne B, Lyon A, et al. AKT
activation predicts outcome in breast cancer patients treated with tamoxifen. *J Pathol.*
2005;207(2):139-46.

17.   Shoman N, Klassen S, McFadden A, Bickis MG, Torlakovic E, and Chibbar R.
Reduced PTEN expression predicts relapse in patients with breast carcinoma treated
by tamoxifen. *Mod Pathol.* 2005;18(2):250-9.

18.   Morandi A, Martin LA, Gao Q, Pancholi S, Mackay A, Robertson D, et al. GDNF-
RET signaling in ER-positive breast cancers is a key determinant of response and
resistance to aromatase inhibitors. *Cancer Res.* 2013;73(12):3783-95.

19.   Boulay A, Breuleux M, Stephan C, Fux C, Brisken C, Fiche M, et al. The Ret receptor
tyrosine kinase pathway functionally interacts with the ERalpha pathway in breast
cancer. *Cancer Res.* 2008;68(10):3743-51.

20.   Plaza-Menacho I, Morandi A, Robertson D, Pancholi S, Drury S, Dowsett M, et al.
Targeting the receptor tyrosine kinase RET sensitizes breast cancer cells to tamoxifen
treatment and reveals a role for RET in endocrine resistance. *Oncogene.*
2010;29(33):4648-57.

21.   Spanheimer PM, Woodfield GW, Cyr AR, Kulak MV, White-Baer LS, Bair TB, et al.
Expression of the RET proto-oncogene is regulated by TFAP2C in breast cancer
independent of the estrogen receptor. *Ann Surg Oncol.* 2013;20(7):2204-12.

22.   Kakati RT, Kim H, Whitman A, and Spanheimer PM. High expression of the RET
receptor tyrosine kinase and its ligand GDNF identifies a high-risk subset of estrogen
receptor positive breast cancer. *Breast Cancer Res Treat.* 2023.

23.   Gou X, Kim BJ, Anurag M, Lei JT, Young MN, Holt MV, et al. Kinome
reprogramming is a targetable vulnerability in ESR1 fusion-driven breast cancer.
Cancer Res. 2023.

24.   Spanheimer PM, Cyr AR, Gillum MP, Woodfield GW, Askeland RW, and Weigel RJ.
Distinct pathways regulated by RET and estrogen receptor in luminal breast cancer
demonstrate the biological basis for combination therapy. *Ann Surg.* 2014;259(4):793-
9.

25.   Spanheimer PM, Park JM, Askeland RW, Kulak MV, Woodfield GW, De Andrade
JP, et al. Inhibition of RET increases the efficacy of antiestrogen and is a novel
treatment strategy for luminal breast cancer. *Clin Cancer Res.* 2014;20(8):2115-25.

26.   Spanheimer PM, Lorenzen AW, De Andrade JP, Kulak MV, Carr JC, Woodfield GW,
et al. Receptor Tyrosine Kinase Expression Predicts Response to Sunitinib in Breast
Cancer. *Ann Surg Oncol.* 2015;22(13):4287-94.

27.   Miller KD, Trigo JM, Wheeler C, Barge A, Rowbottom J, Sledge G, et al. A
multicenter phase II trial of ZD6474, a vascular endothelial growth factor receptor-2
and epidermal growth factor receptor tyrosine kinase inhibitor, in patients with
previously treated metastatic breast cancer. *Clin Cancer Res.* 2005;11(9):3369-76.

28. Xu J, Higgins MJ, Tolaney SM, Come SE, Smith MR, Fornier M, et al. A Phase II
Trial of Cabozantinib in Hormone Receptor-Positive Breast Cancer with Bone
Metastases. *Oncologist.* 2020.
29. Lim JSJ, Wong ALA, Ow SGW, Ngoi NYL, Chan GHJ, Ang YLE, et al. Phase Ib/II
dose expansion study of lenvatinib combined with letrozole in post-menopausal
women with hormone receptor positive breast cancer. *Clin Cancer Res.* 2022.
30. Gupta G, Wang Q, Ellington S, Guerra P, Gharibpoor F, Simpson D, et al. FANCA
Deficiency Induces Oncogenic R-Loop Dependent Synthetic Lethality with PARP1
Inhibitors. *Res Sq.* 2025.
31. Quinlan AR, and Hall IM. BEDTools: a flexible suite of utilities for comparing
genomic features. *Bioinformatics.* 2010;26(6):841-2.
32. Ross-Innes CS, Stark R, Teschendorff AE, Holmes KA, Ali HR, Dunning MJ, et al.
Differential oestrogen receptor binding is associated with clinical outcome in breast
cancer. *Nature.* 2012;481(7381):389-93.
33. Wang Q, Li M, Wu T, Zhan L, Li L, Chen M, et al. Exploring Epigenomic Datasets
by ChIPseeker. *Curr Protoc.* 2022;2(10):e585.
34. Yu G, Wang LG, and He QY. ChIPseeker: an R/Bioconductor package for ChIP peak
annotation, comparison and visualization. *Bioinformatics.* 2015;31(14):2382-3.
35. Ramirez F, Ryan DP, Gruning B, Bhardwaj V, Kilpert F, Richter AS, et al.
deepTools2: a next generation web server for deep-sequencing data analysis. *Nucleic
Acids Res.* 2016;44(W1):W160-5.
36. Hahne F, and Ivanek R. Visualizing Genomic Data Using Gviz and Bioconductor.
Methods Mol Biol. 2016;1418:335-51.
37. Kong NR, Chai L, Tenen DG, and Bassal MA. A modified CUT&RUN protocol and
analysis pipeline to identify transcription factor binding sites in human cell lines.
STAR Protoc. 2021;2(3):100750.
38. Duncan JS, Whittle MC, Nakamura K, Abell AN, Midland AA, Zawistowski JS, et al.
Dynamic reprogramming of the kinome in response to targeted MEK inhibition in
triple-negative breast cancer. *Cell.* 2012;149(2):307-21.
39. Zawistowski JS, Bevill SM, Goulet DR, Stuhlmiller TJ, Beltran AS, Olivares-
Quintero JF, et al. Enhancer Remodeling during Adaptive Bypass to MEK Inhibition
Is Attenuated by Pharmacologic Targeting of the P-TEFb Complex. *Cancer Discov.*
2017;7(3):302-21.
40. Kim H, Whitman AA, Wisniewska K, Kakati RT, Garcia-Recio S, Calhoun BC, et al.
Tamoxifen Response at Single Cell Resolution in Estrogen Receptor-Positive Primary
Human Breast Tumors. *Clin Cancer Res.* 2023.
41. Germain PL, Lun A, Garcia Meixide C, Macnair W, and Robinson MD. Doublet
identification in single-cell sequencing data using scDblFinder. *F1000Res.*
2021;10:979.
42. Subramanian A, Tamayo P, Mootha VK, Mukherjee S, Ebert BL, Gillette MA, et al.
Gene set enrichment analysis: a knowledge-based approach for interpreting genome-
wide expression profiles. *Proc Natl Acad Sci U S A.* 2005;102(43):15545-50.
43. Guillen KP, Fujita M, Butterfield AJ, Scherer SD, Bailey MH, Chu Z, et al. A human
breast cancer-derived xenograft and organoid platform for drug discovery and
precision oncology. *Nat Cancer.* 2022;3(2):232-50.
44. Sorlie T, Perou CM, Tibshirani R, Aas T, Geisler S, Johnsen H, et al. Gene expression
patterns of breast carcinomas distinguish tumor subclasses with clinical implications.
P Natl Acad Sci USA. 2001;98(19):10869-74.

45.   Zhang S, Li F, Younes M, Liu H, Chen C, and Yao Q. Reduced selenium-binding
protein 1 in breast cancer correlates with poor survival and resistance to the anti-
proliferative effects of selenium. *PLoS One.* 2013;8(5):e63702.
46.   Chen R, Wang X, Fu J, Liang M, and Xia T. High FLT3 expression indicates
favorable prognosis and correlates with clinicopathological parameters and immune
infiltration in breast cancer. *Front Genet.* 2022;13:956869.
47.   Papadakis ES, Reeves T, Robson NH, Maishman T, Packham G, and Cutress RI.
BAG-1 as a biomarker in early breast cancer prognosis: a systematic review with
meta-analyses. *Br J Cancer.* 2017;116(12):1585-94.
48.   Kahl I, Mense J, Finke C, Boller AL, Lorber C, Gyorffy B, et al. The cell cycle-
related genes RHAMM, AURKA, TPX2, PLK1, and PLK4 are associated with the
poor prognosis of breast cancer patients. *J Cell Biochem.* 2022;123(3):581-600.
49.   Yi M, Zhang D, Song B, Zhao B, Niu M, Wu Y, et al. Increased expression of ECT2
predicts the poor prognosis of breast cancer patients. *Exp Hematol Oncol.*
2022;11(1):107.
50.   Curtis C, Shah SP, Chin SF, Turashvili G, Rueda OM, Dunning MJ, et al. The
genomic and transcriptomic architecture of 2,000 breast tumours reveals novel
subgroups. *Nature.* 2012;486(7403):346-52.
51.   Saal LH, Vallon-Christersson J, Hakkinen J, Hegardt C, Grabau D, Winter C, et al.
The Sweden Cancerome Analysis Network - Breast (SCAN-B) Initiative: a large-
scale multicenter infrastructure towards implementation of breast cancer genomic
analyses in the clinical routine. *Genome Med.* 2015;7(1):20.
52.   Klaeger S, Heinzlmeir S, Wilhelm M, Polzer H, Vick B, Koenig PA, et al. The target
landscape of clinical kinase drugs. *Science.* 2017;358(6367).
53.   Zhu L, Lu Z, and Zhao H. Antitumor mechanisms when pRb and p53 are genetically
inactivated. *Oncogene.* 2015;34(35):4547-57.
54.   Emerling BM, Hurov JB, Poulogiannis G, Tsukazawa KS, Choo-Wing R, Wulf GM,
et al. Depletion of a putatively druggable class of phosphatidylinositol kinases inhibits
growth of p53-null tumors. *Cell.* 2013;155(4):844-57.
55.   O'Connell CE, and Vassilev A. Combined Inhibition of p38MAPK and PIKfyve
Synergistically Disrupts Autophagy to Selectively Target Cancer Cells. *Cancer Res.*
2021;81(11):2903-17.
56.   Klomp JE, Diehl JN, Klomp JA, Edwards AC, Yang R, Morales AJ, et al.
Determining the ERK-regulated phosphoproteome driving KRAS-mutant cancer.
Science. 2024;384(6700):eadk0850.
57.   Stuart T, Butler A, Hoffman P, Hafemeister C, Papalexi E, Mauck WM, 3rd, et al.
Comprehensive Integration of Single-Cell Data. *Cell.* 2019;177(7):1888-902 e21.
58.   Puram SV, Tirosh I, Parikh AS, Patel AP, Yizhak K, Gillespie S, et al. Single-Cell
Transcriptomic Analysis of Primary and Metastatic Tumor Ecosystems in Head and
Neck Cancer. *Cell.* 2017;171(7):1611-24 e24.
59.   Wu SZ, Al-Eryani G, Roden DL, Junankar S, Harvey K, Andersson A, et al. A single-
cell and spatially resolved atlas of human breast cancers. *Nat Genet.* 2021;53(9):1334-
47.
60.   East MP, Sprung RW, Okumu DO, Olivares-Quintero JF, Joisa CU, Chen X, et al.
Quantitative proteomic mass spectrometry of protein kinases to determine dynamic
heterogeneity of the human kinome. *bioRxiv.* 2024.

61.   Stine ZE, McGaughey DM, Bessling SL, Li S, and McCallion AS. Steroid hormone
831   modulation of RET through two estrogen responsive enhancers in breast cancer. *Hum*
832   *Mol Genet.* 2011;20(19):3746-56.
833
834

835 **ACKNOWLEDGEMENTS:** We would like to thank Dr Alana Welm at the University of
836 Utah for generously providing the patient derived xenograft models.
837
Funding:
National Institutes of Health grant R37CA292075 (PMS)
National Institutes of Health grant K08CA280388 (PMS)
Gilead Sciences Solid Tumor Research Scholars Grant (PMS)
Society of Surgical Oncology Young Investigator Award (PMS)
National Institutes of Health grant P30CA016086 (PMS, CMP, LAC)
National Institutes of Health grant P50CA058223 (CMP, LAC)
Susan G. Komen Foundation Grant (LAC, CMP)
Breast Cancer Research Foundation Grant (LAC, CMP)
847
Author Contributions:
Conceptualization: RTK, GLJ, PMS
Methodology: AAW, ECB, DOO, MPE, ASB, JRR
Investigation: AAW, RTK, SH, CHNCT, ATS, DOO, MPE, ASB, JRR
Visualization: SH, ECB, JRR
Funding Acquisition: LAC, CMP, PMS
Supervision: GLJ, PMS
Writing – original draft: RTK, SH, PMS
Writing – review and editing: all authors
857
858 **Competing interests:** C.M.P is an equity stockholder and consultant of BioClassifier LLC;
859 C.M.P is also listed as an inventor on patent applications for the Breast PAM50 Subtyping
860 assay. The remaining authors declare no potential conflicts of interest.
861
Data and Materials Availability: Single cell and bulk RNA sequencing data was made
available at the Sequence Read Archive (SRA) under BioProject PRJNA1177387. Mass
spectrometry protein sequencing results are included in this manuscript as Supplementary
Table 1. All original code has been deposited and is publicly available in the Github
repository https://github.com/santiagohaase/Spanheimer-lab-Kinase-Plasticity-in-Response-
to-Vandetanib-Enhances-Sensitivity-to-Tamoxifen

FIGURES

Figure 1. Vandetanib reduces ERK activity and is synergistic with Tamoxifen in endocrine sensitive and resistant breast cancer cells and Patient Derived Organoids (A) Schematic of vandetanib target kinases and downstream RAS-RAF-ERK signaling cascade. (B) Treatment of MCF-7 and tamoxifen resistant MCF7 (MCF7-TAM) and exemestane resistant MCF7 (MCF7-EXE) reduces pERK/ERK in all 3 cell lines. *$p<0.05$, ***$p<0.01$ by t-test with Welch correction (C) Vandetanib causes a dose dependent reduction in cell viability in MCF7, MCF7-TAM, and MCF7-EXE. (D) Treatment with Vandetanib results in a left shift of the tamoxifen sensitivity curve and reduction in IC50 in all 3 cell lines. (E) Isobologram plots showing predicted additive lines and regions of synergy (red) or antagonism (blue) for vandetanib–tamoxifen combinations demonstrate synergy in all 3 cells lines. CI = Combination Index (<1 indicates synergic interaction) (F) Patient derived breast cancer organoids show a range of vandetanib sensitiivty. (G) In total, 5 PDO showed synergy between vandetanib and tamoxifen, 2 had an additive effect, and 1 PDO and an organoid derived from the TNBC cell line SUM159 were antagonistic.

Figure 2. Vandetanib effects are partially mediated by the RET receptor (A) Co-treatment with vandetanib and the selective RET inhibitor BLU667 reduced vandetanib sensitivity in MCF7(B) Western blot demonstrating CRISPR knockout of RET resulting in >90% reduction in RET protein levels. RET knockout resulted in reduced pERK and ER expression. (C) RET knockout cells have reduced sensitivity to vandetanib and tamoxifen. (D) Heatmap showing top up- and down-regulated genes in RET KO cells. (E) Gene Set Enrichment Analysis demonstrates reduction in hallmark RAS signaling and RET activity in RET KO cells. RET KO cells also show decreased expression of estrogen response genes consistent with reduced sensitivity to tamoxifen. (F) RET KO cells have lower proliferation score and reduced proliferation compared to WT. (. *p<0.05, **p<0.01, ***p<0.001, ****p<0.0001 by t-test with Welch correction for the bar graphs, and Wilcoxon test (A-C) and for statistical test on k (growth constant) parameter from lineal regression model (F).

Figure 3. Vandetanib regulates gene expression resulting in a low proliferation, estrogen responsive Luminal-A like state. Heatmap and enrichment analysis of differentially expressed genes between vandetanib-treated vs control cells in (A) MCF7, (B) MCF7-TAM, (C) MCF7-EXE. Enriched and depleted gene signatures in (D) MCF7, (E) MCF7-TAM, and (F) MCF7-EXE. (G) KRAS activity signatures were depleted in all 3 cell lines by vandetanib treatment supporting on target effects. (H) Vandetanib treatment increases association with the more indolent, endocrine therapy responsive Luminal A subtype, measured by correlation to Luminal A defining genes (centroid). Overlap of up- (I) and down- regulated (J) genes by cell line. (K) Signatures of down regulated gene signature are associated with overall survival and recurrence free survival (L) in ER+ breast cancer patients in METARIC and in (M) SCAN-B studies. (N) The downregulated gene set is enriched in clinically ET-resistant ER+ tumors. *p<0.05, **p<0.01, ***p<0.001, ****p<0.0001 by t-test with Welch correction for Wilcoxon test.

Figure 4. Enriched estrogen response signatures with vandetanib treatment correlate with enriched ERα binding site occupancy. (A) Heatmap showing gene expression signatures with enriched and depleted ERα promotor binding site occupancy. (B) Heatmap of estrogen response genes graphically demonstrating that changes in gene expression with vandetanib treatment correspond with changes in ERα promoter binding. Genes with increased expression also show enriched FOXA1 occupancy and promoter activity (H3K4me3). (C) GSEA for enriched estrogen activity genes demonstrate canonical E2 response signatures (D) Vandetanib results in increased ER binding to its own promotor representing a forward driven feedback mechanism. (E) H3K4me3 peaks filtered for known transcription driving promoter binding sites demonstrate enriched ER and FOXA1 binding motifs in upregulated sites demonstrating a central role for these transcription factors in changes in gene expression with vandetanib treatment.

Figure 5. Functional Changes in Protein Signaling Networks with Vandetanib Treatment Identifies Targetable Adaptive Resistance Via Upregulation of HER2 activity (A) Multiplex inhibitor bead mass spectrometry assay workflow. (B) Vandetanib resulted in depletion of known vandetanib targets with similar magnitude observed across cell lines. (C) Baseline expression of vandetanib targets in tamoxifen and exemestane resistant cell lines compared to WT MCF7. (D) Heatmap showing enriched and depleted proteins after vandetanib treatment. (E) ERK pathway proteins were downregulated and PI3K upregulated with vandetanib treatment in all 3 cell lines. (F) Heatmap showing enrichment levels of druggable kinases of sensitive and resistant MCF7 cell lines before and after vandetanib treatment. Of note, ERBB2 (encoding HER2) was enriched in all 3 cell lines after treatment and RET was depleted in MCF7 and MCF7-TAM. (G) Sensitivity plots of MCF7 cells co-treated with vandetanib and HER2 inhibitor lapatinib demonstrating enhanced sensitivity with vandetanib treatment. *p<0.05, ****p<0.0001, t-test with Welch correction.

Figure 6. Primary ER+/HER2- breast tumor cells demonstrate conserved transcriptional effects with vandetanib treatment (A) Single-cell RNA sequencing uniform manifold projection (UMAP) annotated by cell type within a primary ER+ human breast tumor. (B) Tumor cells were computationally isolated as having significantly inferred copy number changes and plotted by treatment condition. (C) Heatmap showing differentially expressed genes between vandetanib treated vs non-treated tumor cells. (D) Enriched and depleted pathways of vandetanib treated vs non-treated tumor cells. (E) ERBB2 activity signature score was increased in vandetanib treated tumor cells as seen in cell lines. Down regulated genes in vandetanib treated cell lines are depleted in vandetanib treated primary tumor cells as were KRAS activity signature demonstrating on target effects. LumA correlation increased in vandetanib treated tumor cells as was seen in cell lines. Significant genes filtered by adjusted p value <0.05 (Wilcoxon test) and significant signatures filtered by < 0.1 FDR on GSEA analysis.

Figure 7. Vandetanib signatures predict sensitivity in patient derived models. (A) H+E slides demonstrate histology (H+E) and strongly ER+, weakly PR+, and HER2- status for two PDX models. (B) Five mice for each PDX were randomized to treatment with control or vandetanib (50 mg/kg daily) by oral gavage. HCI011 had reduced growth with vandetanib and HCI05 did not. ****p<0.0001 by Linear mixed-effects model growth slope test after 20 days post treatment (C) The vandetanib sensitive HCI011 had higher baseline expression of vandetanib down-regulated genes and the vandetanib resistant HCI05 had higher expression of our primary tumor NMF6 derived resistance signature, supporting the utility of these signatures to identify tumors that are responsive or resistant to vandetanib. *p<0.05 t-test with Welch correction. (D) PDO signature correlation with response to vandetanib measured by IC50 and vandetanib-tamoxifen synergy. A score of the difference between our human tumor NMF derived vandetanib resistance and vandetanib sensitive signatures most strongly correlated with response to vandetanib (high score indicated resistance (high IC50)).