Prefrontal gamma oscillations and fear extinction learning require early postnatal interneuron-oligodendroglia communication

Fabrice Plaisier[1,2], Cristóbal Ibaceta-González[2], Hasni Khelfaoui[2], Adrianne E S Watson[3], Corinne Poilbout[2,6], Anastassia Voronova[3,4], María Cecilia Angulo[2,5]

1 Inovarion SAS, 251 rue Saint Jacques, 75005 Paris, France

2 Université Paris Cité, Institute of Psychiatry and Neuroscience of Paris (IPNP), INSERM U1266, "Team: Interactions between neurons and oligodendroglia in myelination and myelin repair", F-75014 Paris, France

3 Department of Medical Genetics, Faculty of Medicine and Dentistry, University of Alberta, Edmonton, Alberta, Canada

4 Department of Cell Biology, Faculty of Medicine and Dentistry, University of Alberta, Edmonton, Alberta, Canada

5 GHU PARIS psychiatrie & neurosciences, F-75014 Paris, France

6 Present address: Inserm, UMR-S 1180, "Team: Signaling and cardiovascular pathophysiology", Université Paris-Saclay, 91400 Orsay, France.

Corresponding author:

María Cecilia Angulo, Institute of Psychiatry and Neuroscience of Paris (IPNP), INSERM 1266, 102, rue de la Santé, 75014 Paris, FRANCE. Tel: 33-1-40789243. e-mail address: maria-cecilia.angulo@u-paris.fr

Abstract

Emerging evidence links oligodendrocyte (OL) lineage cells and myelin to cognitive processes, yet the role of myelination in shaping neuronal networks critical for cognitive tasks remains unknown. We demonstrate that early postnatal GABAergic signaling between interneurons and oligodendrocyte precursor cells (OPCs) is crucial for myelination of parvalbumin (PV) interneurons, which facilitates *in vivo* low-gamma oscillations in the medial prefrontal cortex (mPFC) and supports fear extinction learning. Disruption of this signaling resulted in PV interneuron dysmyelination, decreased low-gamma power, and impaired tone fear extinction. These deficits were specific to PV interneuron dysmyelination, as overall mPFC myelination, high-gamma oscillations and context fear extinction remained unaffected. Increasing PV interneuron activity or enhancing myelination did not reverse the deficits, indicating the long-term consequences of these early myelination impairments. Our findings reveal critical roles of OPC GABAergic signaling in PV interneuron myelination and mPFC circuit maturation, with lasting impacts on gamma rhythms and cognitive function.

Introduction

Oligodendrocytes (OLs) are specialized glial cells of the central nervous system (CNS) that produce myelin, a lipid-rich substance that envelops axons, facilitates faster impulse propagation and supplies energy. Beyond their cellular functions as myelinating glia, several lines of evidence highlight the pivotal role of OL lineage cells in the underpinnings of cognitive processes[1]. Behavioral interventions, such as fear conditioning tasks or working memory training in mice, enhance both oligodendrogenesis and myelination, while genetic models impeding OL differentiation result in compromised fear memory recall and spatial memory consolidation[2–4]. In humans, the correlation between white matter changes and cognitive functions further supports the link between OLs, myelin plasticity and cognition[1,5].

While cognitive processes rely on the orchestrated coordination of neuronal networks and the resulting brain oscillations, the contribution of glial cells to this orchestration remains underexplored. Recent advances in mathematical modeling have provided insights into the potential role of myelin in promoting temporal synchronization among neuronal ensembles[6]. This theoretical framework is consistent with *in vivo* experiments conducted on the auditory system of dysmyelinated mouse models characterized by abnormal myelin formation, which exhibited delayed action potentials and loss of temporal precision[7]. However, the conversion of OL-induced neuronal synchronization into brain oscillation activities is still poorly understood. For instance, cuprizone demyelinated mice exhibit an enhanced rather than a decreased cortical theta power during phases of quiet wakefulness[8]. Moreover, besides myelin, the generation of new OLs themselves is crucial for coupling spindle oscillations in the prefrontal cortex with sharp wave ripple oscillations in the hippocampus following learning[2]. While these initial studies offer some insight, the influence of OL lineage cells on distinct oscillation frequencies that dictate cognitive performance remains elusive.

Cortical oscillations result from the dynamic interactions of excitatory and inhibitory neurons,

with gamma oscillations being of particular importance for cognitive function[9]. Notably, gamma-band

oscillations result from the fast synchronization of excitatory neuronal activity, a phenomenon largely

modulated by parvalbumin (PV) interneurons that provide a robust synaptic inhibition[10–12]. Interestingly,

PV interneurons share a unique association with OL lineage cells, arising from common embryonic

origins[13]. During postnatal development, these interneurons form *bona fide* GABAergic synapses with

OL precursor cells (OPCs) which disappear after the second postnatal week in mice[14–16]. Recent evidence

highlights the critical role of early communication between PV interneurons and OPCs *via* GABA

receptors in the myelination and maturation of PV interneurons, as well as in shaping cortical inhibitory

circuit function and mouse behaviour[17,18]. Moreover, PV interneurons constitute the largest myelinated

population of GABAergic interneurons in the cortex[19,20]. Considering the relationship between PV

interneurons, OL lineage cells and myelin, we propose that early postnatal interactions between PV

interneurons and OPCs play a crucial role in establishing prefrontal synchronization and oscillations in

adulthood, especially in the gamma-frequency range, which could have significant implications for

cognitive function.

To address our hypothesis, we genetically disrupted $GABA_A$ receptor-mediated synapses on

OPCs during early postnatal development and subjected adult mice to a fear conditioning task while

recording oscillations in the infralimbic (IL) region of the medial prefrontal cortex (mPFC), a key area

for processing emotional and cognitive information. Our results demonstrate that early postnatal

disruption of OPC GABAergic synaptic signalling results in impairments in tone fear extinction learning

which arise from a reduction in low-gamma frequency power during tone fear extinction sessions. These

functional deficits are associated with PV interneuron dysmyelination and an excitation/inhibition (E/I)

imbalance in the adult mPFC. Efforts to restore the altered phenotype in transgenic mice by local

chemogenetic-mediated enhancement of PV interneuron activity in the mPFC or by *in vivo* infusion of

the pro-myelinating compound fractalkine (FKN) failed to recover learning and memory deficits.

Altogether, our findings reveal the critical role of early interneuron-OPC synaptic signalling in shaping

mPFC circuits and associated fear extinction learning and memory, while also highlighting the long-

lasting and irreversible impact of early OL-mediated disruptions on cognitive deficits.

 Results

Early disruption of GABAergic synapses of OPCs impairs tone fear extinction

To investigate the potential impact of early postnatal interneuron-OPC interactions on cognitive

functioning in adulthood, we used the *NG2cre$^{ERT2+/-}$;Gcamp3$^{f/f}$;γ2$^{f/f}$* mice (γ2$^{f/f}$ mice) in which transient

γ2 subunit-mediated GABAergic synapses of OPCs are inactivated by tamoxifen injections administered

from postnatal day 3 (P3) to P5[21] (Supplementary Fig. 1a; see Online Methods). In agreement with our

previous reports which demonstrated recombination at the γ2 locus and a partial reduction in OPC evoked

GABAergic postsynaptic currents (I$_{GABA}$) in the somatosensory cortex[17,21], our current results reveal a

near-complete abolishment of I$_{GABA}$ in layer 5 OPCs within the IL region of acute mPFC slices of γ2$^{f/f}$

mice at P9-P13 (Supplementary Fig. 1b, 1c). This finding indicates a more pronounced impairment of

interneuron-OPC synaptic signaling in the mPFC compared to the somatosensory cortex. After

confirming the early functional inactivation of OPC GABAergic synapses in this cortical region in γ2$^{f/f}$

mice, we next investigated whether this developmental synaptic alteration would affect mPFC-dependent

cognitive functions in adulthood. To assess this, we evaluated four-month-old γ2$^{f/f}$ mice using a fear

conditioning task, a widely used paradigm to investigate mPFC-dependent processes such as learning,

memory, emotional regulation, and perception[22].

To evaluate learning and memory processes associating environmental cues with aversive

experiences, adult mice subjected to 4-OHT injections at P3-P5, underwent a behavioral test where a 30

s conditioned tone stimulus was paired with a foot shock delivered during the last 2 s of the tone in a

specific context (Fig. 1a). In the conditioning session, adult control and γ2$^{f/f}$ mice exhibited similar fear

memory acquisition, albeit slightly faster in γ2$^{f/f}$ mice (Fig. 1b). No significant differences were found

between the two groups during the context extinction session (Fig. 1c). However, in the first tone

extinction session, although the baseline level (BL, first trial block) of fear expression were comparable

in control and γ2[f/f] mice (Fig. 1d), the control mice showed faster and more extensive fear extinction

learning by the end of the protocol (Fig. 1d). To test whether this deficit in fear extinction learning was

persistent, we performed a second tone extinction session on the following day (extinction retrieval). We

found that, in this session, control mice exhibited significantly lower baseline values and reduced

freezing levels compared to the first tone extinction session, indicating successful extinction retrieval

(Fig. 1e). The tone extinction learned in the first session was therefore retained and successfully retrieved

over time in control mice. In contrast, γ2[f/f] mice showed persistently high baseline values and a weak

extinction, with freezing levels remaining similar to those observed in the first tone extinction session

(Fig. 1e), indicating that these mice failed to improve or consolidate fear extinction learning between

sessions. Of note, no significant sex differences were observed between male and female mice in both

control and γ2[f/f] groups (Supplementary Fig. 2a-h).

Given that the deficit was specific to tone extinction and not observed in context extinction, we

considered whether this could be due to auditory impairments. To rule out this possibility, we assessed

auditory function (Supplementary Fig. 3a, 3b). Both groups exhibited comparable startle responses to

random noise of varying dB levels, indicating no significant differences in auditory sensitivity

(Supplementary Fig. 3c, 3d). Building on these findings, we further analyzed the extinction learning

deficit in γ2[f/f] mice throughout different stages of the retrieval session (Supplementary Fig. 2i). The

extinction learning deficit in γ2[f/f] mice was predominant at early and intermediate stages, but it was also

evident at the late stage (Fig. 1f), showing their impaired ability to acquire and consolidate an extinction

memory.

Our results indicate that γ2[f/f] mice exhibit reduced fear learning and memory capabilities,

particularly in tone fear extinction learning, while auditory perception remains intact. These results

suggest that the disruption of OPC GABAergic synaptic signaling during the early postnatal period leads

to long-term cognitive impairments in adulthood, specifically affecting processes related to learning and

memory, without impacting basic sensory functions like auditory sensitivity.

PV interneuron dysmyelination in the mPFC after early interneuron-OPC synapse disruption

The observed cognitive deficits prompted us to investigate whether early disruption of

interneuron-OPC synapses results in persistent myelination deficits, particularly in PV interneurons

known for their highly plastic myelination[23]. Previous findings in the somatosensory cortex demonstrated

that γ2$^{f/f}$ mice exhibit myelination defects in PV interneurons during juvenile stages[17]. Here, we extended

this analysis to the mPFC in adulthood to determine whether these abnormalities persist and specifically

affect the myelination of PV interneurons, the primary source of transient GABAergic synaptic input

onto OPCs[15,16]. Since fear conditioning experiments revealed specific deficits in tone fear extinction in

γ2$^{f/f}$ mice, we focused on the IL region of the mPFC, a critical region for fear extinction memory

processes[24]. We found that γ2$^{f/f}$ mice displayed normal overall myelin coverage and intensity of MBP

staining in IL cortical layers (Fig. 2a-c). We then analyzed the length of individual myelin segments, or

internodes, along PV$^+$ axons in the same region. This was achieved by reconstructing high resolution

images of axonal segments expressing MBP, PV and the axonal marker SMI-312 (Fig. 2d).

Measurements revealed a statistically significant increase in the mean internode length of PV$^+$, but not

PV$^-$, axons of γ2$^{f/f}$ mice compared to controls (Fig. 2e, 2f). The myelination defect observed in PV$^+$ axons

was not accompanied by alterations in either the cell density or distribution of PV interneurons across

mPFC layers, nor in the total interneuron population identified by GAD67 expression (Supplementary

Fig. 4). Furthermore, we found no differences in the cell density or proportion of Olig2$^+$/CC1$^-$ OPCs and

Olig2$^+$/CC1$^+$ OLs between control and γ2$^{f/f}$ mice (Supplementary Fig. 5). These results suggest that the

early disruption of γ2-mediated synapses in OPCs impacts predominantly PV interneuron myelination,

without affecting interneuron survival or the maturation of OL lineage cell in the mPFC.

Although OPCs form GABAergic synapses only transiently during postnatal development and

these synapses are absent in adulthood[14,15,25], myelination defects in PV interneurons persists into

adulthood of $\gamma2^{f/f}$ mice. Notably, these alterations are specific to PV interneurons as we found no

significant changes in the overall mPFC myelination pattern, length of internodes of PV$^-$ axons, OL

lineage cell densities, PV interneuron survival or the distribution of interneurons within cortical layers.

Early interneuron-OPC synapse disruption irreversibly impairs fear extinction, even with

enhanced myelination and PV interneuron activity in the mPFC

The persistent myelination and cognitive impairments observed in $\gamma2^{f/f}$ mice raised the question

of whether these deficits could be corrected through targeted interventions. To this end, we explored two

distinct strategies aimed at addressing the underlying structural and behavioral abnormalities: enhancing

PV interneuron activity and promoting myelination.

Previous research has demonstrated that using a modified form of the human M3 muscarinic

(hM3Dq) receptor, activated by injections of the inert clozapine metabolite clozapine-N-oxide (CNO)

for 14 days, can specifically promote activity-dependent myelination of PV interneurons in the adult

mPFC[26]. Building on this finding, we explored whether targeting selectively PV interneurons with a

chemogenetic approach could effectively address dysmyelination and fear learning deficits (Fig. 3a). We

performed bilateral intracranial injections into the IL region of the mPFC in $\gamma2^{f/f}$ mice using either pAAV-

S5E2-dTom-nlsdTom (dTom) or pAAV-S5E2-hM3D(Gq)-P2A-dTomato (hM3Dq-dTom) viral vectors[27]

(Fig. 3a, 3b). Approximately 80% of cells infected by these viral constructs were identified as PV$^+$

interneurons in this region (Fig. 3b, 3c). Our results demonstrate that twice-daily in vivo injections of

CNO over two weeks in hM3Dq-dTom-injected $\gamma2^{f/f}$ mice did not induce any significant changes in either

the coverage or mean intensity of the MBP fluorescence signal compared to those in dTom-injected

control mice (Fig. 3c-e). Furthermore, we did not detect any reduction of PV interneuron internode length

in hM3Dq-dTom in $\gamma 2^{f/f}$ mice (Fig. 3g), suggesting that increased PV interneuron activity in adulthood

does not restore aberrant internode length. To examine whether enhancing PV interneuron activity *in*

vivo, independent of the effect on interneuron dysmyelination, ameliorates cognitive deficits, we

conducted the fear conditioning task in CNO-treated control and $\gamma 2^{f/f}$ mice. As expected, no differences

were observed in the acquisition or context sessions (Supplementary Fig. 6a, 6b). Consistent with the

lack of effect on PV interneuron internode length, we observed no significant differences between the

two groups during the fear acquisition phase or the context and tone extinction sessions between the two

groups (Fig. 3f-i, Supplementary Fig. 6a, 6b).

Our second strategy consisted in enhancing myelination in young adult mice, a developmental

period when myelination is still actively progressing in the mPFC and cognitive deficits associated with

neurodevelopmental disorders are often recognized due to the emergence of visible symptoms[1]. We

reasoned that targeting this critical period might offer greater potential for modifying myelination and

improving cognitive deficits. We focused on fractalkine (FKN), a well-studied chemokine known for its

role in microglial function[28], but also shown to promote OPC differentiation into mature OLs during

cortical development through its release from GABAergic interneurons[29]. Recent studies demonstrate

that FKN can promote myelination under both healthy and demyelinating conditions when administered

exogenously[30,31]. We thus exploited the pro-myelinating potential of FKN to examine whether its

exogenous application in mice could compensate for PV interneuron dysmyelination in the mPFC and

mitigate the associated learning and memory deficits observed during tone fear extinction. To this end,

we performed *in vivo* FKN infusions into the lateral ventricles of P42-46 mice over a 25-28-day period

(Fig. 4a). Our results revealed that *in vivo* FKN infusions in $\gamma 2^{f/f}$ mice increased both the coverage and

mean intensity of MBP fluorescent signal in the mPFC of infused mice compared to vehicle-treated mice,

indicating an enhanced myelination (Fig. 4b-d). However, measurements of internode length of

$PV^+/SMI-312^+$ axons in the same animals showed no difference between the two groups, suggesting that

this intervention is insufficient to repair PV interneuron-specific myelin abnormalities in γ2$^{f/f}$ mice (Fig.

4e). To assess whether FKN-induced myelination enhancement could still result in cognitive

improvements, we conducted the fear conditioning task in γ2$^{f/f}$ mice treated with either vehicle or FKN

for comparison. As expected, no differences were observed in the acquisition or context sessions

(Supplementary Fig. 6c, 6d). However, no improvements were observed in the tone extinction or

extinction retrieval sessions either (Fig. 4f, 4g), indicating that enhancing myelination during ongoing

mPFC myelination fails to rectify PV interneuron dysmyelination or improve the associated learning and

memory deficits.

Our findings show the irreversible cognitive impairments resulting from early disruption of OPC

GABAergic synapses. Despite targeted interventions aimed at increasing PV interneuron activity or

enhancing myelination, we observed no improvements in PV interneuron internode length and fear

extinction learning and memory. Once early interneuron-OPC interactions are compromised during a

critical developmental window, subsequent efforts to reverse structural and functional deficits become

particularly challenging. The early postnatal period, particularly the first two postnatal weeks[15], is

therefore essential for ensuring proper cognitive development and function.

Excitation-inhibition imbalance in the mPFC after early interneuron-OPC synapse disruption

The long-lasting cognitive impairments observed in γ2$^{f/f}$ mice, particularly during tone fear

extinction, suggest that these deficits may be caused by underlying neurophysiological alterations.

Therefore, we specifically tested the local excitation-inhibition (E/I) balance in the mPFC because this

mechanism is involved in regulating cognitive processes such as learning, memory, and emotional

regulation[9,11]. This hypothesis is further supported by the known role of mPFC PV interneurons in

regulating the E/I balance by modulating excitatory inputs, which are necessary for higher-order

cognitive functions[32]. To test the E/I balance in the IL region of adult γ2$^{f/f}$ mice, we performed patch-

clamp recordings of layer 2/3 pyramidal cells in acute slices, while stimulating in layer V with an

extracellular electrode (Fig. 5a). The stimulation elicited both excitatory (eEPSCs) and inhibitory

(eIPSCs) postsynaptic currents in mature IL circuits (Fig. 5a). Our results revealed that, three months

after inactivating the γ2 subunit-mediated GABAergic synapses in OPCs, γ2$^{f/f}$ mice exhibited a

significant reduction in the mean amplitudes of eIPSCs, but not eEPSCs, in pyramidal neurons (Fig. 5a-

c). Consequently, the E/I ratio appeared higher in γ2$^{f/f}$ mice, causing a strong E/I imbalance (Fig. 5d).

Since the proper regulation of E/I balance is essential for normal cognitive processing, these findings

point to a critical impairment in mPFC functioning due to early interneuron-OPC synapse disruption.

Specific alterations in gamma oscillations during fear extinction, not exploration

The E/I balance in the mPFC, which is disrupted in γ2$^{f/f}$ mice, plays a fundamental role in

generating gamma oscillations (30-90 Hz), high-frequency neural rhythms which support prefrontal-

dependent cognitive processing. These oscillations arise from the rapid synchronization of excitatory

activity, with PV interneurons playing a central role by providing perisomatic inhibition onto pyramidal

cell ensembles, which fine-tunes network discharges[9,11,12]. Deficits in E/I balance and PV interneuron

myelination could potentially disrupt gamma oscillations, impacting cognitive functions. To investigate

this possibility, we recorded local field potentials (LFPs) from the IL region of the mPFC in freely moving

mice using an implanted tetrode (Fig. 6a), of which the placement was confirmed postmortem through

histological analysis (Fig. 6b). After 10 days of recovery post-implantation, mice were first habituated to

wearing the recording cable in their home cage for three days. Then, they were exposed to an open field

environment and later subjected to the fear conditioning task, during which LFP signals were recorded

(Fig. 6a, 6c). We analyzed these signals using fast Fourier transformation (FFT) to generate spectrograms

and power spectra allowing us to extract different oscillatory activities during the behavioral tasks (Fig.

6c-f, 6i, 6j).

In vivo LFP recordings during the open field test detected both theta (4-12 Hz) and gamma (30-

90 Hz) oscillations in the mPFC of control and $\gamma2^{f/f}$ mice (Fig. 6c). Contrary to our hypothesis that

disrupted myelination would impair gamma oscillations, there were no significant differences in theta or

gamma oscillatory activity between the two groups, regardless of whether the mice were in the center or

periphery of the arena (Fig. 6e-h). These results are consistent with the absence of significant differences

in exploratory behavior between control and $\gamma2^{f/f}$ mice, including distance traveled, time spent,

immobility time in the center and periphery of the arena, as well as the number of crossings or average

speed (Supplementary Fig. 7a-f). These findings indicate that deficits in PV interneuron myelination did

not result in alterations in either neural activity of IL region or exploratory behavior under the conditions

of the open field task.

The consequences of PV interneuron dysmyelination in the mPFC may be task-specific,

potentially impacting complex processes rather than general locomotor or exploratory behaviors. These

same implanted mice were therefore subjected to the fear conditioning protocol which is designed to

assess more cognitively demanding processes (Fig. 6a). *In vivo* LFP recordings were performed during

the tone extinction retrieval session, when more pronounced behavioral differences between control and

$\gamma2^{f/f}$ mice were observed (Fig. 1e). During this session, we optimized recordings by using a Faraday cage

and minimizing external noise. Analysis of the oscillatory activity revealed no significant differences in

the relative power of theta oscillations between the two groups during tone presentation and inter-trial

intervals (ITI) (Fig. 6i-k). However, $\gamma2^{f/f}$ mice exhibited a significant reduction in gamma oscillations

throughout both periods, suggesting a specific disruption of neural dynamics at this frequency range (Fig.

6i-j, 6l). These results indicate that early postnatal disruption of interneuron-OPC signaling leads to

persistent PV interneuron dysmyelination and E/I imbalance in the mPFC, which in turn drive alterations

in gamma oscillations and the subsequent impairments in fear extinction learning observed in $\gamma2^{f/f}$ mice.

The absence of significant gamma oscillation changes in the open field, in contrast to the

pronounced reduction observed during fear extinction retrieval, suggests that PV interneuron

dysmyelination predominantly affects higher cognitive processes, without necessarily disrupting basal

neural activity or basic behaviors like exploration or locomotion.

Specific disruption in low-gamma oscillations during fear extinction retrieval

Gamma oscillations can be subdivided into low and high frequency ranges, each contributing

distinctly to cognitive processes. High-gamma frequencies have been associated with encoding

processes, whereas low-gamma frequencies support retrieval in human episodic memory[33]. To

investigate whether the differences in gamma oscillations observed during fear extinction retrieval

between control and $\gamma2^{f/f}$ mice were frequency-specific, we conducted a detailed analysis of both low-

gamma (30-45 Hz) and high-gamma (55-90 Hz) power. Our results revealed a significant reduction in

the relative power of low-gamma oscillations in $\gamma2^{f/f}$ mice during both tone presentation and ITI of fear

extinction retrieval, while high-gamma oscillations showed no significant differences (Fig. 7a, 7b). These

results suggest that deficits in neuronal processing in mPFC of $\gamma2^{f/f}$ mice are primarily associated with

alterations in low-gamma oscillations.

Gamma oscillations are known to couple strongly with theta oscillations during cognitive

processes, playing a key role in coordinating network-wide communication[34]. Our analyses reveal that

the strength of the theta phase-low-gamma amplitude and theta phase-high-gamma amplitude coupling

increased during tone presentation compared to the ITI in both control and $\gamma2^{f/f}$ mice (Supplementary

Fig. 8). This increase was observed across the entire session as well as at the different stages within the

session (Supplementary Fig. 8), indicating a robust network activation and an ability in both groups to

discriminate the tone stimuli. This strong theta–gamma coupling serves as a hallmark of effective

cognitive processing, especially since no significant changes in coupling were observed between the

center and periphery of the arena in the open field task (Supplementary Fig. 7g, 7h). However, despite

the clear increase in coupling during tone presentation compared to ITI, there were no significant

differences between control and γ2$^{f/f}$ mice in the tone/ITI coupling ratio, either for low-gamma or high-

gamma oscillations (Supplementary Fig. 8c, 8g). The lack of coupling defects shows that the coordination

of broad network interactions remains intact, despite a decrease in low-gamma power.

To further dissect the temporal dynamics of low-gamma power alterations, we analyzed the

relative power of low-gamma oscillations at different stages of the retrieval session: early, intermediate,

and late (Fig. 7c, 7d). We found that low-gamma power was significantly reduced during the early and

intermediate stages in γ2$^{f/f}$ mice compared to controls (Fig. 7c), while it remained unaffected during the

late stage (Fig. 7d). These early and intermediate reductions in low-gamma power correspond with the

higher freezing levels observed in γ2$^{f/f}$ mice at these stages (Fig. 1f), suggesting a temporal relationship

between low-gamma disruption and behavioral deficits. In contrast, high-gamma activity remained

similar between control and γ2$^{f/f}$ mice throughout all three stages of the retrieval session (Fig. 7d), further

supporting the conclusion that the observed alterations are specific to the low-gamma frequency range.

In summary, our findings suggest that low-gamma oscillations in the mPFC are more critical than

high gamma oscillations in supporting fear extinction retrieval. The reduction in low-gamma power,

particularly during the early and intermediate stages of the retrieval session in γ2$^{f/f}$ mice, points to its

essential role in enabling neuronal network activity required for the successful fear extinction learning

and memory. This reduction, alongside the lack of changes in theta-gamma coupling, is consistent with

an intracortical mPFC network dysfunction, resulting from PV interneuron impairments which are key

for intracortical projections.

Discussion

This study reveals that the early postnatal period is a critical window for interneuron-oligodendroglia interactions, during which disruption of OPC GABAergic synaptic signaling in γ2[f/f] mice leads to persistent impairments in local neuronal network function and cognitive performance in the adulthood. Specifically, deficits in learning and memory during tone fear extinction are associated with disrupted low-gamma oscillations in the IL region of the mPFC. These alterations arise from early dysfunctions in interneuron-OPC signaling, leading to impaired PV interneuron myelination and disrupted neuronal E/I balance. Our findings show the necessity of this developmental phase for oligodendroglia and PV interneuron myelination to sustain adult low-gamma oscillations that are critical for tone fear extinction learning and memory. Overall, this study emphasizes the importance of the early postnatal PV interneuron myelination process in maturing inhibitory circuits and its role in maintaining cortical low-gamma rhythms essential for specific cognitive functions.

Defects in OL lineage cells and myelination are increasingly associated with a variety of cognitive deficits[1]. For example, social isolation, whether occurring immediately after weaning or during adulthood, results in hypomyelination of the mPFC[35,36], while early maternal separation impairs emotional responses and object recognition in adult mice, linked to premature OPC differentiation in this cortical region[37]. Proper myelination is also critical for emotional regulation and memory processes, including fear conditioning and extinction, as disruptions in *de novo* OL production or myelin formation prolongs fear responses and hampers the consolidation of contextual fear memories[2,3]. However, while cognitive impairments are often attributed to pronounced hypomyelination or demyelination, the impact of subtle, cell-type-specific dysmyelination, such as that in PV interneurons, remains largely unexplored. PV interneurons play a key role in regulating gamma oscillations and maintaining the E/I balance, both of which are essential for coordinated neuronal network function and cognitive performance[9,11,32]. Our findings suggest that even mild myelination defects in PV interneurons during critical developmental

windows can lead to persistent alterations in inhibitory circuitry, impairing cognitive function. The

alterations in low-gamma oscillations observed in our study, without significant changes in high-gamma

oscillations, suggest that specific deficits in PV interneurons can selectively impact distinct gamma

frequency bands. This cell-specific dysmyelination could represent a previously underestimated

mechanism underlying cognitive deficits and emotional dysregulations present in neurodevelopmental

and psychiatric disorders, where disruptions in PV interneuron function and gamma oscillations are

commonly implicated[1]. Our findings open the possibility that subtle myelination abnormalities in PV

interneurons may contribute to the E/I imbalance and network dysfunctions characteristic of conditions

such as schizophrenia and autism spectrum disorders[1]. The contribution of oligodendroglia to these

conditions has likely been underestimated and deserves further investigation.

Recent studies in healthy mice demonstrate that behavioral training and stimuli can induce subtle

changes in myelination. Motor-skill learning induces myelin remodelling, including the elongation of

nodes of Ranvier[38], highlighting the importance of a fine-tuned myelination during learning processes.

Furthermore, the length of nodes of Ranvier is modulated in distinct ways by interventions like repetitive

transcranial magnetic stimulation or spatial learning[39]. Despite these insights into activity-dependent

delicate myelination changes, the role of oligodendroglia and myelin in modulating neuronal network

activity and cortical oscillations during the execution of cognitive processes in health and disease is

poorly understood. A recent study using cuprizone-treated mice has shown that prolonged impairments

in social and spatial working memory, as well as in cognitive flexibility, persist after remyelination. These

impairments correlate with long-term desynchronized activity between the hippocampus and the mPFC,

characterized by lasting disruptions in gamma synchronization and ripple/slow oscillation coupling, a

condition that develops during the demyelination phase and is only partially restored during

remyelination[40]. Furthermore, fiber photometry has revealed that animals unable to form new myelin

exhibit a lack of the time-dependent modulation in mPFC neuronal activity seen in control animals,

which initially show a decrease and then a progressive increase in activity during transitions from

mobility to freezing[3]. Active oligodendrogenesis and myelination are therefore required for the proper

evolution of prefrontal dynamics associated with fear memory recall. Moreover, the disruption of *de novo*

production of OLs immediately before contextual fear conditioning also impairs the coupling between

spindle oscillations in the prefrontal cortex and sharp wave ripple oscillations in the hippocampus,

independent of myelin deficits[2]. Therefore, both OL production alterations and myelin deficiencies are

linked to impaired neuronal responses in brain regions activated during cognitive tasks.

While the recognition of the involvement of oligodendroglia in cognition is growing, the

mechanisms by which these cells influence specific neuronal subtypes and network dynamics during the

execution of a cognitive task remain elusive. Our results demonstrate that early disruption of OPC

GABAergic signaling results in a selective dysmyelination of PV interneurons, without major changes

in overall oligodendrogenesis or myelination, and altered low-gamma oscillation activity during tone fear

extinction. This defect does not uniformly impact neuronal networks or cognitive processes, as high-

gamma oscillations remained unaffected, and no impairments were observed in fear acquisition, context

fear extinction, or exploratory behavior in the open field. The specific effects observed in our study may

be explained by the fact that myelination characteristics such as patterns, myelin content, internode and

node lengths vary across specific neuronal identities and for a specific cell type in different brain

areas[20,41], which can lead to different consequences. This selective effect may arise from differences in

myelination requirements in mPFC subregions, with the IL region depending on low-gamma oscillations

critical for extinction memory, while the prelimbic (PL) region, involved in fear learning, may rely on

different oscillatory patterns and thus be less affected by PV interneuron dysmyelination. In any case,

our findings emphasize the critical role of precise myelination patterns of specific neuronal subtypes in

maintaining proper network dynamics during distinct cognitive functions. While neuronal contributions

to cognitive processes are undisputable, our study positions oligodendroglia and myelination as pivotal

elements driving the construction of prefrontal-dependent low-gamma oscillations necessary for a

specific cognitive process. In addition to oligodendroglia, other types of glial cells, such as astrocytes,

also play a significant role in regulating neuronal network dynamics. Recent findings suggest that

GABAergic signaling to astrocytes, although non-synaptic in this case, also influences low-gamma

oscillations without impacting high-gamma oscillations[42], further strengthening the importance of glial-

neuronal interactions in maintaining cognitive functions. These findings highlight the importance of glial

cells in fine-tuning network oscillations important for higher-order cognitive functions.

While slow oscillations like theta oscillations result from the coordinated activity of widely

distributed neuronal ensembles, fast-frequency oscillations like gamma oscillations are considered a local

phenomenon, involving smaller tissue volumes[1]. Cortical PV interneurons contribute to various

frequency oscillations, but their role in gamma oscillations is crucial. This is due to their ability to

discharge action potentials at high frequencies and their extensive local synaptic connectivity with

surrounding excitatory neurons, which facilitates rapid and synchronized neuronal activity essential for

gamma oscillations[10–12]. Our study revealed that PV interneuron dysmyelination in $\gamma 2^{f/f}$ mice are

associated with a reduced inhibitory drive causing an E/I imbalance and alterations of low-gamma

oscillations in the IL region during tone fear extinction retrieval. A critical involvement of PV

interneurons in low-gamma activity during cognitive processes is supported by a previous study showing

a rescue of low-gamma oscillation alterations and social deficits by the optogenetic stimulation of PV

interneurons at 40 Hz in a model of autism spectrum disorders characterized by a Neuregulin-3

mutation[43]. Moreover, the optogenetic activation of PV interneurons at 30 Hz allows for the preservation

of gamma frequency oscillations in control, but not in mice with widespread demyelination[8]. In these

models as well as in $\gamma 2^{f/f}$ mice, a decreased PV interneuron firing rate is also observed[8,17,43]. Our findings

uniquely highlight the role of PV interneuron myelination in sustaining low-gamma oscillations during

specific cognitive tasks, aligning with previous studies that explore the function of these interneurons in

cortical processing.

PV interneuron myelination remains particularly plastic throughout life, making these neurons

more susceptible to myelin deficiencies. For instance, monocular deprivation results in experience-

dependent adaptive myelin remodeling specifically in cortical PV interneurons, but not in glutamatergic

projection neurons[23]. Furthermore, myelination defects occur specifically in PV interneurons of the

mPFC in a rat model displaying schizophrenia-like behaviors[44], and demyelination induces selective

vulnerability of GABAergic interneurons in both progressive multiple sclerosis and a mouse model of

local cortical demyelination[45]. In a transgenic mouse, mild hypomyelination due to the loss of

oligodendrocyte ErbB receptor signaling results in a decreased PV expression and a reduction in

inhibitory but not excitatory synaptic connections, causing an E/I imbalance in L2/3 neurons of the

auditory cortex[46]. Thus, PV interneuron myelination contributes to the pathological mechanisms

underlying various neurological disorders, making it a novel and attractive pharmacological target[47].

Interestingly, it was previously shown that chemogenetic activation of hippocampal PV interneurons of

adult mice subjected to a modified fear conditioning protocol with severe electric foot shocks reduced

fear generalization and restored myelination of PV interneuron axons[48]. Nevertheless, altering early

interneuron-oligodendroglia interactions, myelination, and neuronal plasticity during critical periods can

generate lasting cognitive and behavioral effects that may be challenging to reverse. In our study, targeted

interventions to increase PV interneuron activity or enhance global myelination in $\gamma2^{f/f}$ mice did not repair

the defect in PV interneuron internode length or lead to improvements in fear extinction memory. This

suggests that, once early interneuron-OPC interactions are compromised, subsequent attempts to enhance

myelination or promote PV interneuron activity are ineffective in reversing the resulting cognitive

deficits. The critical developmental window for these interactions, occurring transiently in the first two

postnatal weeks[15], appears crucial for establishing proper cognitive development and function.

Preserving the integrity of transient interneuron-OPC interactions during this period appears essential to

prevent long-lasting and irreversible cognitive deficits. It should be noted that most psychiatric

treatments begin after the onset of symptoms, which often occur well after these critical periods, making

it challenging to reverse such early-established myelination deficits. Additional evidence suggests that

early but transient dysfunctional events of interneurons can be deleterious to cortical function in

adulthood. Kessaris et al. (2021)[49] demonstrated that an aberrant overproduction of PV interneurons

during embryogenesis has lasting effects that persist into critical periods of cortical development, despite

the normalization of cell density postnatally. This transient excess of interneurons during early

development results in behavioral abnormalities in adulthood, indicating that the timing of interneuron

clearance is essential for the proper maturation of neural circuits that underlie adult behavior. In humans,

early life adversity correlates with increased OPC-mediated perineuronal nets production around PV

interneurons in the ventromedial PFC; as perineuronal nets regulate the closure of developmental

windows of cortical plasticity, this potentially limits cortical circuit neuroplasticity and promotes

maladaptive behaviors[50]. Together, these findings emphasize the challenges in reversing cognitive

deficits once developmental processes are disrupted. An interesting line for future research in myelin

biology would be to explore how abnormalities in PV interneurons and OL lineage synergize, potentially

increasing the risk of neurodevelopmental and psychiatric disorders.

 Methods

Transgenic mice and induction of Cre-mediated recombination

All experiments adhered the European Union and institutional guidelines for the care and use of

laboratory animals. Ethical approval was obtained from the French ethical committee for animal care of

university Paris Descartes (Committee N°CEEA34) and the Ministry of National Education and Research

(Project No: 202010131827131). The experimental approach involved *NG2cre^{ERT2+/-};Gcamp3^{f/f};γ2^{f/f}*

mice (γ2^{f/f} mice) to inactivate γ2-GABA$_A$ receptor-mediated synapses in OPCs while the

NG2^{creERT2+/-};Gcamp3^{f/f} mice were used as controls[17,21]. Genotyping was conducted by PCR using

primers specific to the different alleles, and Cre expression was induced from P3 to P5 by daily

intraperitoneal injections of 0.2 mg of 4-OHT (Tocris Bioscience) as previously described[21]. Animals

were housed in controlled conditions with 12 h light/dark cycle, ad libitum access to food and water, and

in average ambient temperature of 21°C and 45% humidity. Both females and males were used at

different postnatal stages from P10 to P134.

Behavioral assays

Anxiety and locomotion were assessed in a 50 cm × 50 cm x 45 cm open field arena, where mice were

placed for 10 minutes under a 50 lx illumination. An infrared camera monitored animal movements and

the ViewPoint behavioral analysis software processed the data (version 5.31.0.8, ViewPoint, Behavior

Technology, France). The fraction of time spent in an outer area of 2500 cm^2 over an inner square area

of 900 cm^2 was calculated. The open field test was used to assess locomotion by measuring distance

traveled, velocity, time spent in the inner (center) or outer (periphery) area, and number of crossings in

the center area.

The fear conditioning paradigm consisted of four consecutive days of behavioral testing in a

chamber of 30 cm × 25 cm × 25 cm made of two aluminum sidewalls, acrylic front and rear walls and a

floor of metal bars of 3.2 mm diameter separated by 7.9 mm. An infrared camera, controlled by a

computerized system interface, monitored behavior and freezing (percentage of time of immobility)

using the Videofreeze software (version 2.7.2.117, MED Associates, St. Albans, VT, USA). On day-1

(fear conditioning), mice were individually placed into a dark conditioning chamber, odorized with 1%

carvone (mint odor). After a baseline period of 3 min, animals received five pairings of an auditory

conditional stimulus (CS; 30 s, 5 kHz, 85 dB) with the unconditioned stimulus (US) applied at the end of

the tone (2 s, 0.5 mA foot-shock) with a 2 min ITI period. On day-2 (contextual fear extinction), mice

were introduced in the same conditioning chamber, odorized with carvone, and allowed to explore for 18

min without presentation of CS or US. On day-3 (tone fear extinction), mice were placed in chambers

with a modified context and odor (0.5% isopentyl acetate; banana odor). After a baseline period of 3 min,

the CS alone was presented 40 times (30 s), separated by 5 s ITIs. On day-4 (tone fear extinction

retrieval), mice were subjected to the same protocol of day 3. The open-field arena and fear conditioning

chambers box were wiped with 70% ethanol and air-dried between mice.

Freezing behavior was automatically quantified in VideoFreeze software. Animal movement was

analyzed with the default settings (motion threshold: 18 arbitrary units, sample rate: 30 frames per

second, linear method, minimum freeze duration: 30 frames), with freezing defined as immobility lasting

at least 1 s. Freezing responses were calculated across intertrial-intervals (ITIs) on day 1, averaged across

periods of 2 minutes on day 2, and those on day 3 and day 4 were calculated using the average freezing

response to five consecutive CS (5-trial blocks). The tone fear extinction retrieval session (day 4) was

divided into 3 stages (early, intermediate and late), each composed of 13 CS presentation and 13 ITIs for

analysis of the behavioral and *in vivo* electrophysiology data. In this case, the last CS and ITI was

excluded from the analysis. Since no sex differences were observed between male and female mice in

both control and γ2^{f/f} groups (Supplementary Fig. 2a-h), data from both sexes were combined for each

group.

Surgical implantation of the microdrive

A custom microdrive was fashioned using a 10 ml syringe cut to a length of 1.5 cm. This allows the

Neuralynx EIB-16 headstage to be glued to the inside, serving as a base for a mini-Faraday cage. The

cage was made by sticking a sheet of copper foil adhesive tape (noll electronic, Germany) around the

syringe and grounding it with a silver-soldered wire (Ø 203.2 µm, ref. 782000, A-M systems). The conical

bottom of the microdrive is made by cutting the bottom of a 10 ml Falcon® tube. This conical shape

increased the surface in contact with the skull and eased the application of dental cement. The center of

the conical base is drilled to glue a plastic guide for introducing the tetrode composed of 4 tungsten wires

(Ø 50.8/101.6 µm) insulated with Teflon (ref. 795500, A-M systems). The tail of the tetrode along with

the silver ground and reference wires were gently inserted into channels holes on the EIB-16-QC

(Neuralynx), and pinned by small golden pins (Ref. 31-0603, Neuralynx). The fully assembled

microdrive weighed less than 3g.

520   For chronic implantation of the microdrive, adult control and γ2^{f/f} mice were deeply anaesthetized

with isoflurane (4% for induction and 1.5% for maintenance, Vetflurane, Virbac). Local analgesics,

including a solution of lidocaïne (Laocaïne, 7 mg/kg, Intervet-MSD), buprenorphine (BUPRECARE, 0,1

mg/kg; 2 mL/kg; Axience) and an ophtalmic gel (Ocrygel, DOMES PHARMA FR) were administered

prior to or during the surgery. After fixing the animal in the stereotaxic frame (Kopf Instruments), the

skin above the skull was disinfected with 70% ethanol and betadine solution (Vétédine, Vetoquinol,

France) before incision. A hole was stereotaxically drilled to reach the IL region of mPFC using

coordinates −0.4 mm mediolateral, +1.70 mm anterior-posterior relative to bregma. One additional hole

in the interparietal region was performed to insert a screw overlying the cerebellum where the reference

wire and the soldered silver ground wire were connected together. Then, the screw was sealed with a

light-cure, cyanoacrylate-based adhesive (Loctite 4310) to ensure stability. Finally, the microdrive was

fixed to the skull using resin-modified glass ionomer cement (Ketac Cem Plus Automix 3M) and acrylic

dental cement (Pattern Resin LS, GC America Inc). The surface of the exposed brain was covered with

saline (0.9% NaCl) for the duration of the experiment. The skin was sutured (Mersilene, EH7147H,

Ethicon) and a saline solution at 37°C was administered subcutaneously (100 to 200 µL) to rehydrate the

animal. Finally, mice were recovered from anesthesia in an environment at 37°C before they returned to

their home cages.

At the end of experimental procedures, mice were deeply anesthetized using an overdose of a

solution of Ketamine (180 mg/kg, Clorketam 1000, Vetoquinol, France) and Xylazine (10 mg/kg,

Rompum, Elanco France), and electrolytic lesions were made to make the electrode implantation site

more visible. A current of 12 µA for 30 seconds was applied under gas anesthesia (isoflurane) through

the implanted tetrode *via* the EIB-16-QC interface and the NanoZ impedance measurement system

(Neuralynx, USA). Mice were then perfused with phosphate buffer saline (PBS) followed by 0.15 M

phosphate buffer (PB; pH ≈ 7.3), containing 4% paraformaldehyde (PFA). Brains were kept 2 h in PFA,

rinse three times in PB and incubated in sucrose 30% at 4°C until sinking for cryoprotection. Then, brains

were embedded with a film of OCT by gently rolling them (Microm Microtech), frozen in dry ice and

stored at -70°C. Coronal slices were cut with a cryostat (20 µm; Leica CM3050 S Cryostat), stained with

DAPI and mounted. Sections were visualized using an epifluorescence microscope (Zeiss Axioplan 2) to

confirm the recording location. Animals with the tetrode not clearly positioned in the mPFC were

excluded from the analysis.

In vivo **electrophysiological recordings**

Recordings were performed 7-15 days after surgery by using Neuralynx 16 channels preamplifier and 7 m tether with its data acquisition software Cheetah (Neuralynx, USA). LFPs were recorded at 30 kHz sampling rate (Neuralynx, USA). Animals were habituated to experimenter handling and to recording sessions in their home cage for 15 min twice a day during 3 days; recording duration were 10 min in the open-field and 27 min on day-4 of the fear conditioning test.

The analysis of mPFC oscillations were carried out with custom-made programs written in Python3 (Spyder). All scripts used in this study are freely available at Angulo's lab gitlab repository (https://gitlab.com/d5674/c_ibaceta/psd-volt). The mPFC LFP signals were band-pass filtered from 1 Hz to 120 Hz (fifth-order IR Butterworth filter), 50 Hz notch filtered (Notch width 0.1 Hz) to minimize AC 50 Hz noise. For LFP analysis, the data was downsampled from 30 kHz to 1 kHz. Then, we performed a power spectrum analysis (PSD) using the Welch method from SciPy Python and Neurodsp library[51]. The parameters used were: 2048 FFTs, Hann window and 1024 overlapping points. Following the PSD calculation, we normalized the data obtained during the open-field exploration by averaging the values across all frequencies (from 1Hz to 120Hz). For fear extinction retrieval, we normalized the data by averaging the values of frequencies ranging from 0Hz to 120Hz during the baseline period (3 min). We defined three frequency bands, theta (4-12Hz), low gamma (30-45Hz) and high gamma (55-90Hz), and obtained the PSD for each band range. For 3D plots depicting PSD over time and frequency, we used Neurodsp library[51].

The PAC analysis was conducted using the Hilbert Transform, extracting the phase of Theta and the amplitude of both low-gamma and high-gamma, with a FIT filter to isolate the three frequencies (using SciPy Python library). For the analysis during tone fear extinction, we considered three stages: baseline, sound and ITI. We further subdivided the sound and ITI in three phases based on trial numbers, according to their trials: early (from 1 to 13), intermediate (from 14 to 26) and late (from 27 to 39). We calculated PSD and PAC for each of these phases.

In vivo **fractalkine (FKN) infusions**

In vivo FKN infusions were performed by adapting the protocol described by Watson et al. (2021)[30]. On the day of the surgery, FKN was dissolved in 0.2% BSA-PBS (75 µg/ml; Recombinant Soluble Mouse CX3CL1/Fractalkine, R&D systems) or BSA in PBS (0.2%, Jackson ImmunoResearch). P42-P46 $\gamma2^{f/f}$ mice were deeply anesthetized with isoflurane (4% for induction and 1.5% for maintenance, Vetflurane, Virbac). The surgical procedure and mouse recovery were similar to that described above. For the local delivery of these compounds, a hole was stereotaxically drilled to implant a canula, containing either FKN or BSA, into the right lateral ventricle using coordinates −1.000 mediolateral, −0.300 anterior-posterior, −2.500 dorsoventral relative to bregma. Finally, the micro-osmotic pump (model 1004, Alzet) was attached via a catheter to the cannula and placed in a subcutaneous pocket between the shoulder blades. Infusions were administered at a flow rate of 0.11 µL per hour for 25-28 days. In total, each mouse received 200 ng of FKN per day, a dose that promoted myelination in adult murine CNS[30,31].

Viral injections and CNO protocol

Viral-mediated specific labeling of PV interneurons in the mPFC were performed by adapting the protocol described by Stedehouder et al., 2018[26]. Adult $\gamma2^{f/f}$ mice were used for viral injections, receiving either pAAV-S5E2-dTom-nlsdTom for controls (Addgene, #135630) or pAAV-S5E2-Gq-P2A-dTomato (Addgene, #135635) for DREADD stimulation[27]. The surgical procedure and mouse recovery were similar to that described above. Viral particles were injected into the mPFC through a 10 µl gastight syringe (model 1701, Hamilton) using a motorized micropump (Legato 130, KD Scientific) and silica needle (Ø 75/150 µm, PHYMEP), using coordinates −0.4 mm mediolateral, +1.70 mm anterior-posterior, −2.9 mm dorsoventral relative to bregma. Four weeks after surgery, all mice received intraperitoneal

injections of CNO (Clozapine N-oxide (CNO) dihydrochloride, 1 mg/kg; Hello Bio) twice daily

(approximately at 9:00 h and 18:00 h) for 14 consecutive days.

Patch-clamp recordings on acute mPFC slices

Patch-clamp experiments were performed in the IL region of the mPFC using coronal slices of 300 µm-

thick at P9-P13 for OPCs (Supplementary Fig. 1) and P92-P124 for layer 2/3 pyramidal neurons (Fig.

5a-d). Recorded cells were visualized with an Orca Flash 4.0 LT CMOS digital camera (Hamamatsu) and

Micro-Manager software (version 1.4.21[52]) under an Olympus BX51 microscope equipped with a 40X

fluorescent water-immersion objective. Excitation light to detect fluorescent OPCs was provided by the

CoolLed pE-300 ultra-system (Scientifica, UK) and 470 and 525 nm filters for excitation and emission

wavelengths, respectively.

Patch-clamp recordings were performed at RT using an extracellular solution containing (in mM):

126 NaCl, 2.5 KCl, 1.25 NaH_2PO_4, 26 $NaHCO_3$, 20 glucose, 5 pyruvate, 3 $CaCl_2$ and 1 $MgCl_2$ (95%

O_2, 5% CO_2). Glutamatergic and GABAergic postsynaptic currents of layer 2/3 pyramidal neurons and

fluorescent layer 5 OPCs were evoked at −70 mV and 0 mV, respectively by a monopolar electrode (glass

pipette) placed in layer V (100 ms pulse, 10 V; Iso-Stim 01D, npi electronic GmbH, Tamm, Germany),

with a CsMeS-based intracellular solution containing (in mM): 125 $CsCH_3SO_3H$, 5.4-aminopyridine,

10 tetraethylammonium chloride, 0.2 EGTA, 0.5 $CaCl_2$, 2 $MgCl_2$, 10 HEPES, 2 Na_2-ATP, 0.2 Na-GTP

and 10 Na_2-phosphocreatine (pH ≈ 7.3). The rate of stimulation was 0.2 Hz for pyramidal cells and 0.1

Hz for OPCs. Recordings were made without series resistance (R_S) compensation. R_S was monitored

during recordings and cells showing a change of more than 30% were discarded. Potentials were

corrected for a junction potential of −10 mV. Acquisition was obtained using Multiclamp 700B and

pClamp version 10.7 software, filtered at 4 kHz and digitized at 20 kHz. Evoked postsynaptic currents

were analyzed off-line using pClamp version 10.7 software (Molecular Devices).

Immunostaining and quantification

For PV and GAD-67 immunolabelling, mice were perfused and their brains were frozen in isopentane at

-70°C and cut into coronal slices with a cryostat (20 µm; Leica CM3050 S Cryostat). The slices were

then fixed in 4% paraformaldehyde (PFA) for 10 min with 0.1 M phosphate buffer (PB) at 4°C, washed

first with PBS and then with tris buffered saline (TBS). Next, they were placed in a blocking buffer with

50 mM TBS, 5% goat serum, 0.4% bovine serum albumin (BSA) and 0.1% Triton x100 for 2h at RT, and

incubated with primary antibodies in a 5% serum, 10% BSA, 10% Triton and 50 mM TBS solution for

three nights in a cold chamber (4°C). Primary antibodies were rabbit polyclonal anti-PV (1:500; ref. PV-

27, Swant) and mouse monoclonal anti-GAD67 (1:200; ref. MAB5406, Sigma-Aldrich). Slices were

washed three times in 50 mM TBS and 0.1% Triton x100 before they were incubated with secondary

antibodies with 10% BSA, 10% Triton and 50 mM TBS at 4°C for two nights. We used goat anti-rabbit

DyLight-405 (ref. 35551; Thermofisher Scientific) and goat anti-mouse Alexa Fluor-546 (ref. A-11030,

Thermofisher Scientific) at 4°C for two nights at 1:200. Slices were washed three times with the same

previous washing buffer and mounted with Vectashield mounting medium (ref. H-1000, Vectorlabs).

For other immunostainings, mice were perfused with PBS followed by 0.15 M phosphate buffer

(PB; pH ≈ 7.3), containing 4% paraformaldehyde (PFA). Brains were kept 2 h in PFA, rinsed three times

in PB and incubated in sucrose 30% at 4°C until sinking for cryoprotection. Then, they were embedded

with a film of OCT (Microm Microtech) by rolling them gently, frozen in dry ice and stored at -70°C.

Coronal slices were cut with a cryostat (20 µm; Leica CM3050 S Cryostat), permeabilized with 0.3%

triton X-100 and 4% Normal Goat Serum (NGS) overnight and incubated with primary antibodies. For

internode length analysis, the slices were incubated with anti-PV and anti-SMI-312 for four nights before

the incubation with the respective secondary antibodies during 2 hours at room temperature. Then, they

were first incubated with anti-MBP for two nights and then with its secondary antibody for 2 hours at

room temperature. For quantification of OPC and OL densities, the slices were incubated with anti-Olig2

and anti-CC1 for four nights and then with its secondary antibody for 2 hours at room temperature.

Primary antibodies used for immunohistochemistry were rabbit polyclonal anti-PV (1:500; ref. PV-27,

Swant), mouse anti-SMI-312 (1:200; ref. 837901, BioLegend), rat monoclonal anti-MBP (1:200; clone

26; ref. AB7349, Abcam), rabbit polyclonal anti-Olig2 (1:400; ref. AB9610, Merck Millipore), mouse

monoclonal anti-APC (CC1; 1:100; clone CC-1; ref. OP80, Merck Millipore). Slices with primary

antibodies were washed three times in PBS and incubated in secondary antibodies coupled, respectively,

to goat anti-rabbit DyLight-405 (ref. 35551, Thermofisher Scientific), goat anti-mouse Alexa Fluor-546

(ref. A-11030, Thermofisher Scientific), goat anti-rat Alexa Fluor-633 (ref. A-21094, Thermofisher

Scientific) and goat anti-mouse DyLight-633 (ref. GTX76787; Genetex) at RT at 1:200.

For marker[+] cell analysis, confocal images were acquired using 63x oil objectives and, for

internode length and MBP analysis, confocal image stacks of 15-20 µm depth were acquired using 93x

oil objectives with an SP8 Leica confocal microscope and the LAS X software (version 3.5.7.23225).

Images were then analyzed blind using the Fiji/ImageJ2 framework (NIH, version 1.53i[53]). Cells were

counted from layer 1 to 6 in the IL region of mPFC with ROI manager tool (3-4 images per animal). Each

image containing 20 z-sections of 1 µm each. Cells that were partially cut from the image were excluded

in three of the six sides of the cube to prevent border effects. PV[+,] GAD-67[+], Olig2[+] and CC1[+] cell

densities were calculated as the number of cells per volume (mm^3). For the analysis of cell distributions,

the x coordinate of each cell was extracted from FIJI/ImageJ2 and its position was normalized with

respect to the total length of the layers of the mPFC, which can vary from one slice to another. Then, we

calculated the frequency of cells in the normalized mPFC length with a binning of 0.1. To quantify MBP

fluorescence coverage and intensity, we applied a mask generated from a binary (black & white) image

in FIJI. This mask was created to specifically isolate regions with MBP signal, allowing for targeted

analysis of fluorescence exclusively in these areas. To measure the length of internodes, semi-automatic

3D reconstructions of SMI-312$^+$/PV$^+$/MBP$^+$ segments were obtained using Simple Neurite Tracer plugin

in Fiji/ImageJ2 framework[17,54]. Only internodes flanked by two nude axonal regions were considered for

analysis.

Statistics

All data are expressed as mean ± SEM. In the dot plots, individual mice (N) are represented by closed

dots while repeated measures (n) are shown as open dots. Statistical analysis were performed using

GraphPad Prism (version 10; GraphPad Software Inc., USA), and JASP (version 0.16.4; JASP Team,

2022, Amsterdam, The Netherlands). Each group of data was first subjected to Shapiro-Wilk normality

test. According to the data structure and variance equality, two-group comparisons were performed using

a two-tailed Student's *t* test, a Welch's t test or a Mann–Whitney U test for unpaired samples. We also

used linear mixed models with likelihood ratio tests to account for repeated measures within each animal.

Multiple group comparisons were performed using the parametric two-way ANOVA followed by a post-

hoc Sidak, Tukey's or Fishers Least Significant Difference (LSD) multiple comparison tests, as well as

mixed-effects model followed by a Sidak's multiple comparison test for situations where data were not

normality distributed. Differences were considered significant when $P < 0.05$.

Acknowledgements

We thank the NeurImag platform, Phenobrain and the animal facility of IPNP and their funding sources (Fédération pour la Recherche Médicale, Fondation Leducq). We would also thank Ludivine Therreau from Phenobrain as well as Beata Turanska and Julie Cognet in the team for their technical assistance, Steven A. Kushner and Karim Benchenane for insightful discussions on the project, Najate Benamer, Marie Vidal, Yana Kibalnyk, Philippe Faure and Elise Bousseyrol for help at the preliminary phase of the study. M.C.A. is a CNRS (Centre National de la Recherche Scientifique) investigator.

Funding

A.V and M.C.A. were supported by an Era-Net Neuron grant (CIHR and ANR under the frame of Neuron Cofund, No. 161466 and No. R19068KK for A.V. and M.C.A, respectively). A.E.S.W. was supported by CIHR CGSD. A.V. was supported by CIHR grants (PS 166120 and PJT 195940), MS Canada grant (1249541), the Scottish Rite Charitable Foundation of Canada grant as well as Canada Research Chair in Neural Stem Cell Biology and Sloan Research Fellowship in Neuroscience. M.C.A. was supported by grants from Fondation pour la Recherche Médicale (FRM, EQU202103012626 and MND202310017891), ANR (ANR CoLD: ANR-20-CE16–0001-01; ANR Myelex: ANR-21-CE37–0020-01), Fondation de France (00147199/WB-2023-50772) and Fédération pour la Recherche sur le Cerveau.

Author contributions

F.P. conducted the majority of the experiments and performed data analysis of the study. C.I-G. developed the Python code for in vivo electrophysiological analysis and contributed to data analysis. C.P. carried out behavioral experiments and immunostainings. A.E.S.W. and A.V. performed and assisted with FKN infusions. H.K. performed patch-clamp recordings of OPCs and virus injections. M.C.A. supervised the

project, contributed to data analysis, and drafted the initial version of the manuscript. All authors edited

and approved the manuscript.

Data and code availability

The data generated and analyzed during this study are included in the manuscript and extended data; they

are available from the corresponding author upon reasonable request. Psd-volt Python scripts are hosted

at https://gitlab.com/d5674/c_ibaceta/psd-volt and published under a Free Software GNU GPLv3 license.

Declaration of interests

None.

 Figure Legends

Figure 1. Inactivation of GABAergic synapses in OPCs disrupts tone fear extinction

a. Mice were subjected to a four-day fear conditioning protocol consisting of four sessions: a conditioning

session (five 2s-foot shocks delivered at the end of each 30s tone), a contextual extinction session (18

min), a tone extinction session (40 tones of 30s each with 5s inter-trial interval), and a tone extinction

retrieval session (40 tones of 30s each with 5s inter-trial interval). **b-e.** Comparison of freezing behavior

(percentage of time spent immobile) between control (dark blue) and $\gamma2^{f/f}$ (light blue) mice during each

session. No major differences were observed between the two groups during conditioning (**b**) and context

extinction (**c**) sessions. However, during both tone extinction (**d**) and tone extinction retrieval (**e**)

sessions, $\gamma2^{f/f}$ mice exhibited significantly higher freezing levels across all 5-trial blocks compared to

controls. Note that baseline (BL) freezing was also higher in $\gamma2^{f/f}$ mice during the extinction retrieval

session. **f.** Comparison of freezing behavior between control and $\gamma2^{f/f}$ mice at early, intermediate and late

stages of the extinction retrieval session (averaged freezing of 13 tones for each stage see Supplementary

Fig. 2i). *$p<0.05$, **$p<0.01$, ***$p<0.001$, ****$p<0.0001$, two-way ANOVA test followed by post-hoc

Sidak multiple comparisons test. Data are represented as mean±s.e.m (control mice: N=18 and $\gamma2^{f/f}$ mice:

N=23).

Figure 2. Inactivation of GABAergic synapses of OPCs impairs PV interneuron internode length

in the mPFC

a. Confocal images of MBP in the IL region of the mPFC of control (left) and $\gamma2^{f/f}$ (right) mice. Scale

bar: 50 µm. **b, c.** Dot plots of MBP fluorescence coverage (**b**) and intensity (**c**) of control (dark blue) and

$\gamma2^{f/f}$ (light blue) mice. ns: not significant; linear mixed models with likelihood ratio tests applied to

account for repeated measures (open dots) within each animal (closed dots). Data are presented as

mean±s.e.m. **d.** Confocal images of PV (red), MBP (green), and SMI-312 (blue) labeling, illustrating

myelinated internodes of PV$^+$ axons, identified by bare axonal regions on each side, in control and γ2$^{f/f}$

mice (white arrowheads). Note that the internode of PV$^+$ axons in γ2$^{f/f}$ mice appeared longer compared

to those of control mice. Scale bar: 10 µm. **e,f.** Dot plots of internode lengths of PV$^+$ (**e**) and PV$^-$ (**f**) axons

in control (dark blue) and γ2$^{f/f}$ (light blue) mice. *p<0.05, ns: not significant; linear mixed models with

likelihood ratio tests applied to account for repeated measures (open dots) within each animal (closed

dots). Data are represented as mean±s.e.m.

Figure 3. Chemogenetic-induced increase in PV interneuron activity does not rescue fear behavior

a. Schematic of viral injections and CNO administration protocol for PV interneuron activation in the

mPFC. The γ2$^{f/f}$ mice received bilateral injections of either pAAV-S5E2-dTom-nlsd (dTom) or pAAV-

S5E2-hM3Dq(Gq)-P2A-dTomato (hM3Dq-dTom) virus targeting PV interneurons in the mPFC at

around P90. After one month, the same mice received twice-daily intraperitoneal injections of CNO for

14 consecutive days. The day following the final CNO injection, they were subjected to the fear

conditioning protocol (d1–d4). **b.** Confocal image of a coronal section showing the restricted expression

of hM3Dq-dTom virus on the IL region of the mPFC. Scale bar: 500 µm. **c.** Confocal images of

dTom$^+$/PV$^+$ neurons in the IL region. A dTom$^+$/PV$^+$ cell is shown (*Inset*, arrow); a dTom$^+$/PV$^-$ cells is

indicated (arrowhead). Dot plots (right) of the percentage of dTom$^+$/PV$^+$ cells in both dTom-injected

(white) and hM3Dq-dTom-injected (red) γ2$^{f/f}$ mice. ns: not significant; linear mixed models with

likelihood ratio tests applied to account for repeated measures (open dots) within each animal (closed

dots). Data are represented as mean±s.e.m. Scale bar: 50 µm and 10 µm. **d.** Confocal images of MBP in

the IL region of the mPFC of dTom-injected (left) and hM3Dq-dTom-injected (right) γ2$^{f/f}$ mice. **e, f.** Dot

plots of MBP fluorescence coverage (**e**) and intensity (**f**) of the IL region of dTom-injected (white) and

hM3Dq-dTom-injected (red) γ2$^{f/f}$ mice. ns: not significant; linear mixed models with likelihood ratio

tests applied to account for repeated measures (open dots) within each animal (closed dots). Data are

presented as mean±s.e.m. **g.** Dot plots of internode lengths of PV$^+$ axons in dTom-injected (white) and

hM3Dq-dTom-injected (red) γ2$^{f/f}$ mice. ns: not significant; linear mixed models with likelihood ratio tests

applied to account for repeated measures (open dots) within each animal (closed dots). Data are

represented as mean±s.e.m. **h-i.** Comparison of freezing behavior between dTom-injected (white) and

hM3Dq-dTom-injected (red) γ2$^{f/f}$ mice during tone extinction (**h**) and tone extinction retrieval (**i**)

sessions. No significant differences were observed between the two groups; two-way ANOVA test

followed by Tukey's multiple comparison test. Data are represented as mean±s.e.m (red mice: N=13 and

hM3Dq-dTom mice: N=12).

Figure 4. *In vivo* FKN infusions increase myelination, but does not rescue fear behavior

a. Schematic of the *in vivo* infusion protocol. Vehicle (VC) or FKN was continuously infused into the

lateral ventricle of γ2$^{f/f}$ mice at P42-P46 for 25-28 days (D25-D28) using a micro-osmotic pump. These

mice were subjected to the fear conditioning protocol (d1-d4) during the last days of infusion. **b.** Confocal

images of MBP in the IL region of the mPFC of VC-infused (left) and FKN-infused (right) γ2$^{f/f}$ mice at

P70-74. Note the increased myelination of FKN-infused γ2$^{f/f}$ mouse. Scale bar: 50 µm. **c, d.** Dot plots of

MBP fluorescence coverage (**c**) and intensity (**d**) of the IL region of VC-infused (white) and FKN-infused

(yellow) γ2$^{f/f}$ mice. ***p<0.001, ****p<0.0001; linear mixed models with likelihood ratio tests applied

to account for repeated measures (open dots) within each animal (closed dots). Data are presented as

mean±s.e.m. **e.** Dot plots of internode lengths of PV$^+$ axons in VC-infused (white) and FKN-infused

(yellow) γ2$^{f/f}$ mice. ns: not significant; linear mixed models with likelihood ratio tests applied to account

for repeated measures (open dots) within each animal (closed dots). Data are represented as mean±s.e.m.

f, g. Comparison of freezing behavior between VC-infused (white) and FKN-infused (yellow) γ2$^{f/f}$ mice

during tone extinction (**f**) and tone extinction retrieval (**g**) sessions. No significant differences were

observed between the two groups; two-way ANOVA test followed by post-hoc Sidak multiple

comparisons test. Data are represented as mean±s.e.m (VC-infused mice: N=6 and FKN-infused mice:

N=8).

Figure 5. Inactivation of GABAergic synapses of OPCs impairs inhibition in the mPFC

a. eIPSCs (top trace) and eEPSCs (bottom trace) in layer 2/3 pyramidal neurons held at 0 mV and −70

mV, respectively, upon extracellular stimulation of layer 5 of the infralimbic (IL) region of acute slices

of the mPFC (*inset*, left) in control (dark blue) and $\gamma 2^{f/f}$ (light blue) mice at P100. Stimulation artifacts

blanked, stimulation time indicated (black arrowheads). **b, c.** Dot plots of eIPSCs (**b**) and eEPSCs (**c**) of

layer 2/3 pyramidal neurons evoked by extracellular stimulation in layer 5 in control (dark blue) and $\gamma 2^{f/f}$

mice (light blue) (n=15 cells from 9 mice and n=14 cells from 7 mice for control and $\gamma 2^{f/f}$ mice,

respectively). *p<0.05, ns: not significant; two-tailed Mann–Whitney U test. **d.** Dot plots of E/I ratio

calculated from eEPSCs and eIPSCs. ***p<0.001; two-tailed Mann–Whitney U test.

Figure 6. Disruption of gamma oscillations during fear extinction retrieval, but not exploration

a. Schematic of the protocol for conducting *in vivo* recordings in behaving mice. Surgery to implant the

microdrive containing the tetrode was performed at around P115. Recordings of LFPs were performed

7–15 days-post-surgery during open-field exploration and fear extinction retrieval. **b.** The tetrode was

implanted in the IL region of the mPFC (left). Brain slice of a recorded mouse (middle) after a DAPI

staining (blue) and showing the correct location of the tetrode (right, arrowhead). Scale bar, 1 mm. fmi,

forceps minor of the corpus callosum; CPu, Caudate-Putamen; PL, prelimbic; IL, infralimbic. **c.**

Representative raw LFP signal from the mPFC (top), used to extract the theta (4-12 Hz, middle) and

gamma (30-90 Hz, bottom) frequency components following band-pass filtering. **d.** Color-coded

spectrogram from a mouse LFP recording, showing theta (4-12 Hz) and gamma (30-90 Hz) oscillation

powers during fear extinction retrieval. **e,f, i,j.** Normalized (Norm.) power spectra of mPFC recordings

from the center (C) and periphery (P) of the open-field (OF) arena (**e,f**), as well as during tone and inter-

trial interval (ITI) phases of the fear extinction retrieval session (**i,j**) in control (dark blue) and $\gamma2^{f/f}$ (light

blue) mice. Theta (θ) and gamma (γ) frequency bands are indicated. Power spectra are averaged of all

animals, and the curves and the shaded areas indicate mean±s.e.m. **g,h,k,l.** Dot plots showing the

normalized (Norm.) theta (**g, k**) and gamma (**h,l**) oscillation powers for the center (C) and periphery (P)

of the open-field arena (**g, h**), and for tones and inter-trial intervals (ITI) during the fear extinction

retrieval session (**k, l**) in control (dark blue) and $\gamma2^{f/f}$ (light blue) mice. *p<0.05, ns: not significant; two-

tailed Student's *t* test. All data are represented as mean±s.e.m (control mice: N=5 and $\gamma2^{f/f}$ mice: N=5).

Figure 7. Disruption of low-Gamma oscillations during fear extinction retrieval in $\gamma2^{f/f}$ mice

a,b. Dot plots of the relative power of low-gamma and high-gamma oscillations during tone (**a**) and inter-

trial interval (ITI) (**b**) in control (dark blue) and $\gamma2^{f/f}$ (light blue) mice. **p<0.01, ns: not significant; two-

way ANOVA test followed by post-hoc Sidak multiple comparisons test. **c,d.** Comparisons of the

normalized (Norm.) power of low-gamma (**c**) and high-gamma (**d**) oscillations during tone (left) or inter-

trial interval (ITI; right) in control and $\gamma2^{f/f}$ mice at early, intermediate and late stages of the extinction

retrieval session (average responses during 13 tones for each stage, see Supplementary Fig. 2i).

Individual animal responses are shown by faint traces. *p<0.05, **p<0.01, ns: not significant; two-way

ANOVA test followed by a post-hoc Sidak multiple comparisons test. All data are represented as

mean±s.e.m (control mice: N=5 and $\gamma2^{f/f}$ mice: N=5).

References

1. Khelfaoui, H., Ibaceta-Gonzalez, C. & Angulo, M. C. Functional myelin in cognition and
neurodevelopmental disorders. *Cell Mol Life Sci* **81**, 181 (2024).

2. Steadman, P. E. *et al.* Disruption of Oligodendrogenesis Impairs Memory Consolidation in Adult
Mice. *Neuron* **105**, 150-164.e6 (2020).

3. Pan, S., Mayoral, S. R., Choi, H. S., Chan, J. R. & Kheirbek, M. A. Preservation of a remote fear
memory requires new myelin formation. *Nat Neurosci* **23**, 487–499 (2020).

4. Shimizu, T. *et al.* Oligodendrocyte dynamics dictate cognitive performance outcomes of working
memory training in mice. *Nat Commun* **14**, 6499 (2023).

5. Kaller, M. S., Lazari, A., Blanco-Duque, C., Sampaio-Baptista, C. & Johansen-Berg, H. Myelin
plasticity and behaviour-connecting the dots. *Curr Opin Neurobiol* **47**, 86–92 (2017).

6. Pajevic, S., Plenz, D., Basser, P. J. & Fields, R. D. Oligodendrocyte-mediated myelin plasticity and its
role in neural synchronization. *Elife* **12**, e81982 (2023).

7. Moore, S. *et al.* A role of oligodendrocytes in information processing. *Nat Commun* **11**, 5497 (2020).

8. Dubey, M. *et al.* Myelination synchronizes cortical oscillations by consolidating parvalbumin-
mediated phasic inhibition. *Elife* **11**, e73827 (2022).

9. Jung, F. & Carlén, M. Neuronal oscillations and the mouse prefrontal cortex. *Int Rev Neurobiol* **158**,
337–372 (2021).

10.   Bartos, M., Vida, I. & Jonas, P. Synaptic mechanisms of synchronized gamma oscillations in
inhibitory interneuron networks. *Nat Rev Neurosci* **8**, 45–56 (2007).

11.   Buzsáki, G. & Wang, X.-J. Mechanisms of gamma oscillations. *Annual review of neuroscience*
35, 203–225 (2012).

12.   Hijazi, S., Smit, A. B. & Van Kesteren, R. E. Fast-spiking parvalbumin-positive interneurons in
brain physiology and Alzheimer's disease. *Mol Psychiatry* (2023) doi:10.1038/s41380-023-02168-y.

13.   Benamer, N., Vidal, M. & Angulo, M. C. The cerebral cortex is a substrate of multiple interactions

between GABAergic interneurons and oligodendrocyte lineage cells. *Neurosci Lett* **715**, 134615

(2020).

14.   Vélez-Fort, M., Maldonado, P. P., Butt, A. M., Audinat, E. & Angulo, M. C. Postnatal Switch

from Synaptic to Extrasynaptic Transmission between Interneurons and NG2 Cells. *J Neurosci* **30**,

6921–6929 (2010).

15.   Orduz, D. *et al.* Interneurons and oligodendrocyte progenitors form a structured synaptic network

in the developing neocortex. *eLife* **4**, e06953 (2015).

16.   Orduz, D. *et al.* Developmental cell death regulates lineage-related interneuron-oligodendroglia

functional clusters and oligodendrocyte homeostasis. *Nat Commun* **10**, 4249 (2019).

17.   Benamer, N., Vidal, M., Balia, M. & Angulo, M. C. Myelination of parvalbumin interneurons

shapes the function of cortical sensory inhibitory circuits. *Nat Commun* **11**, 5151 (2020).

18.   Fang, L.-P. *et al.* Impaired bidirectional communication between interneurons and

oligodendrocyte precursor cells affects social cognitive behavior. *Nat Commun* **13**, 1394 (2022).

19.   Micheva, K. D. *et al.* A large fraction of neocortical myelin ensheathes axons of local inhibitory

neurons. *eLife* **5**, e15784 (2016).

20.   Stedehouder, J. *et al.* Fast-spiking Parvalbumin Interneurons are Frequently Myelinated in the

Cerebral Cortex of Mice and Humans. *Cereb Cortex* **27**, 5001–5013 (2017).

21.   Balia, M., Benamer, N. & Angulo, M. C. A specific GABAergic synapse onto oligodendrocyte

precursors does not regulate cortical oligodendrogenesis. *Glia* **65**, 1821–1832 (2017).

22.   Maren, S., Phan, K. L. & Liberzon, I. The contextual brain: implications for fear conditioning,

extinction and psychopathology. *Nat Rev Neurosci* **14**, 417–428 (2013).

23.   Yang, S. M., Michel, K., Jokhi, V., Nedivi, E. & Arlotta, P. Neuron class-specific responses govern

adaptive myelin remodeling in the neocortex. *Science* **370**, eabd2109 (2020).

24.   Sung, Y. & Kaang, B.-K. The Three Musketeers in the Medial Prefrontal Cortex: Subregion-

specific Structural and Functional Plasticity Underlying Fear Memory Stages. *Exp Neurobiol* **31**, 221–

231 (2022).

25.   Balia, M. *et al.* Postnatal Down-Regulation of the GABAA Receptor γ2 Subunit in Neocortical

NG2 Cells Accompanies Synaptic-to-Extrasynaptic Switch in the GABAergic Transmission Mode.

Cereb Cortex **25**, 1114–1123 (2015).

26.   Stedehouder, J., Brizee, D., Shpak, G. & Kushner, S. A. Activity-Dependent Myelination of

Parvalbumin Interneurons Mediated by Axonal Morphological Plasticity. *J Neurosci* **38**, 3631–3642

(2018).

27.   Vormstein-Schneider, D. *et al.* Viral manipulation of functionally distinct interneurons in mice,

non-human primates and humans. *Nat Neurosci* **23**, 1629–1636 (2020).

28.   Camacho-Hernández, N. P. & Peña-Ortega, F. Fractalkine/CX3CR1-Dependent Modulation of

Synaptic and Network Plasticity in Health and Disease. *Neural Plast* **2023**, 1–17 (2023).

29.   Voronova, A. *et al.* Migrating Interneurons Secrete Fractalkine to Promote Oligodendrocyte

Formation in the Developing Mammalian Brain. *Neuron* **94**, 500-516.e9 (2017).

30.   Watson, A. E. S. *et al.* Fractalkine signaling regulates oligodendroglial cell genesis from SVZ

precursor cells. *Stem Cell Reports* **16**, 1968–1984 (2021).

31.   De Almeida, M. M. A. *et al.* Fractalkine enhances oligodendrocyte regeneration and

remyelination in a demyelination mouse model. *Stem Cell Reports* **18**, 519–533 (2023).

32.   Ferguson, B. R. & Gao, W.-J. PV Interneurons: Critical Regulators of E/I Balance for Prefrontal

Cortex-Dependent Behavior and Psychiatric Disorders. *Front Neural Circuits* **12**, 37 (2018).

33.   Griffiths, B. J. *et al.* Directional coupling of slow and fast hippocampal gamma with neocortical

alpha/beta oscillations in human episodic memory. *Proc Natl Acad Sci USA* **116**, 21834–21842 (2019).

34. Tamura, M., Spellman, T. J., Rosen, A. M., Gogos, J. A. & Gordon, J. A. Hippocampal-prefrontal

theta-gamma coupling during performance of a spatial working memory task. *Nat Commun* **8**, 2182

(2017).

35. Makinodan, M., Rosen, K. M., Ito, S. & Corfas, G. A critical period for social experience-

dependent oligodendrocyte maturation and myelination. *Science* **337**, 1357–1360 (2012).

36. Liu, J. *et al.* Impaired adult myelination in the prefrontal cortex of socially isolated mice. *Nat*

Neurosci **15**, 1621–1623 (2012).

37. Teissier, A. *et al.* Early-life stress impairs postnatal oligodendrogenesis and adult emotional

behaviour through activity-dependent mechanisms. *Mol Psychiatry* **25**, 1159–1174 (2020).

38. Bacmeister, C. M. *et al.* Motor learning drives dynamic patterns of intermittent myelination on

learning-activated axons. *Nat Neurosci* **25**, 1300–1313 (2022).

39. Cullen, C. L. *et al.* Periaxonal and nodal plasticities modulate action potential conduction in the

adult mouse brain. *Cell Reports* **34**, 108641 (2021).

40. Mercier, O. *et al.* Transient demyelination causes long-term cognitive impairment, myelin

alteration and network synchrony defects. *Glia* **72**, 960–981 (2024).

41. Call, C. L. & Bergles, D. E. Cortical neurons exhibit diverse myelination patterns that scale

between mouse brain regions and regenerate after demyelination. *Nat Commun* **12**, 4767 (2021).

42. Mederos, S. *et al.* GABAergic signaling to astrocytes in the prefrontal cortex sustains goal-

directed behaviors. *Nat Neurosci* **24**, 82–92 (2021).

43. Cao, W. *et al.* Gamma Oscillation Dysfunction in mPFC Leads to Social Deficits in Neuroligin 3

R451C Knockin Mice. *Neuron* **97**, 1253-1260.e7 (2018).

44. Maas, D. A. *et al.* Interneuron hypomyelination is associated with cognitive inflexibility in a rat

model of schizophrenia. *Nat Commun* **11**, 2329 (2020).

45.   Zoupi, L. *et al.* Selective vulnerability of inhibitory networks in multiple sclerosis. *Acta*

Neuropathol **141**, 415–429 (2021).

46.   Borges, B. C., Meng, X., Long, P., Kanold, P. O. & Corfas, G. Loss of oligodendrocyte ErbB

receptor signaling leads to hypomyelination, reduced density of parvalbumin-expressing interneurons,

and inhibitory function in the auditory cortex. *Glia* **71**, 187–204 (2023).

47.   Stedehouder, J. & Kushner, S. A. Myelination of parvalbumin interneurons: a parsimonious locus

of pathophysiological convergence in schizophrenia. *Mol Psychiatry* **22**, 4–12 (2017).

48.   Yamada, J. *et al.* Altered regulation of oligodendrocytes associated with parvalbumin neurons in

the ventral hippocampus underlies fear generalization in male mice. *Neuropsychopharmacol* **48**,

1668–1679 (2023).

49.   Magno, L. *et al.* Transient developmental imbalance of cortical interneuron subtypes presages

long-term changes in behavior. *Cell Rep* **35**, 109249 (2021).

50.   Tanti, A. *et al.* Child abuse associates with increased recruitment of perineuronal nets in the

ventromedial prefrontal cortex: a possible implication of oligodendrocyte progenitor cells. *Mol*

Psychiatry **27**, 1552–1561 (2022).

51.   Cole, S., Donoghue, T., Gao, R. & Voytek, B. NeuroDSP: A package for neural digital signal

processing. *JOSS* **4**, 1272 (2019).

52.   D. Edelstein, A. *et al.* Advanced methods of microscope control using µManager software. *JBM*

1, 1 (2014).

53.   Schindelin, J. *et al.* Fiji: an open-source platform for biological-image analysis. *Nat Methods* **9**,

676–682 (2012).

54.   Schneider, C. A., Rasband, W. S. & Eliceiri, K. W. NIH Image to ImageJ: 25 years of image

analysis. *Nat Methods* **9**, 671–675 (2012).

a

pAAV-S5E2-dTom-nlsdTom
or
pAAV-S5E2-hM3Dq(Gq)-P2A-dTomato

$\gamma 2^{f/f}$

$\gamma 2^{f/f}$

Age	P3 P4 P5	P90	D28	14 days	D42	d1 d2 d3 d4

4-OHT injection D0 virus bilateral injection CNO injections (I.P. twice daily) fear conditioning

b

hM3Dq-dTom

c

dTom
PV

ns

dTom⁺ PV⁺ cells / dTom⁺ cells (%) → $dTom^+ PV^+ cells / dTom^+ cells (\%)$

dTom   hM3Dq-dTom

d

dTOM   hM3Dq-dTom

IL   MBP   MBP

e

ns

MBP coverage (μm^2)

dTom   hM3Dq-dTom

f

ns

MBP intensity (a.u.)

dTom   hM3Dq-dTom

g

ns

Internode length of PV⁺ axons (μm)

dTom   hM3Dq-dTom

h

Tone extinction

dTom   hM3Dq-dTom

Freezing (%)

BL 1 2 3 4 5 6 7 8
5-trial blocks

i

Extinction retrieval

dTom   hM3Dq-dTom

Freezing (%)

BL 1 2 3 4 5 6 7 8
5-trial blocks

a

γ2^f/f

Age   P3 P4 P5   P42-46   d1 d2 d3 d4
         ↑ ↑ ↑   D0   D25-28
      4-OHT   pump   VC or FKN   fear conditioning
      injection   implantation   infusion

b

VC   FKN

MBP   MBP

IL

c

MBP coverage (μm²)

****

VC   FKN

d

MBP intensity (a.u.)

***

VC   FKN

e

Internode length of PV⁺ axons (μm)

ns

VC   FKN

f Tone extinction

Freezing (%)

○ VC   ● FKN

BL 1 2 3 4 5 6 7 8
5-trial blocks

g Extinction retrieval

Freezing (%)

○ VC   ● FKN

BL 1 2 3 4 5 6 7 8
5-trial blocks