Prefrontal gamma oscillations and fear extinction learning require early postnatal interneuron-oligodendroglia communication

Fabrice Plaisier[1,2], Cristóbal Ibaceta-González[2#], Hasni Khelfaoui[2#], Adrianne E S Watson[3], Corinne Poilbout[2&], Anastassia Voronova[3,4], María Cecilia Angulo[2,5*]

1 Inovarion SAS, 251 rue Saint Jacques, 75005 Paris, France

2 Université Paris Cité, Institute of Psychiatry and Neuroscience of Paris (IPNP), INSERM U1266, "Team: Interactions between neurons and oligodendroglia in myelination and myelin repair", F-75014 Paris, France

3 Department of Medical Genetics, Faculty of Medicine and Dentistry, University of Alberta, Edmonton, Alberta, Canada

4 Department of Cell Biology, Faculty of Medicine and Dentistry, University of Alberta, Edmonton, Alberta, Canada

5 GHU PARIS psychiatrie & neurosciences, F-75014 Paris, France

& Present address: Inserm, UMR-S 1180, "Team: Signaling and cardiovascular pathophysiology", Université Paris-Saclay, 91400 Orsay, France.

# These authors contributed equally to this work.

[*]**Corresponding author:**

María Cecilia Angulo, Institute of Psychiatry and Neuroscience of Paris (IPNP), INSERM 1266, 102, rue de la Santé, 75014 Paris, FRANCE. Tel: 33-1-40789243. e-mail address: maria-cecilia.angulo@u-paris.fr

Abstract

Emerging evidence links oligodendrocyte (OL) lineage cells and myelin to cognitive processes, yet the role of myelination in shaping neuronal networks critical for cognitive tasks remains unknown. We demonstrate in mice that early postnatal GABAergic signaling between interneurons and oligodendrocyte precursor cells (OPCs) is crucial for myelination of parvalbumin (PV) interneurons, which facilitates in vivo low-gamma oscillations in the medial prefrontal cortex (mPFC) and supports fear extinction learning. Disruption of this signaling results in PV interneuron dysmyelination, decreases low-gamma power, and impairs tone fear extinction. These deficits are specific to PV interneuron dysmyelination, as mPFC myelination, high-gamma oscillations and contextual fear extinction are not significantly altered. Increasing PV interneuron activity or enhancing myelination do not reverse the deficits, indicating the long-term consequences of these early myelination impairments. Our findings reveal the role of OPC GABAergic signaling in PV interneuron myelination and mPFC circuit maturation, with lasting impacts on gamma rhythms and cognition.

Brief Summary

Early postnatal interneuron-oligodendroglia communication may shape neuronal networks underlying cognition. Here, the authors show that disrupting this signaling in mice impairs interneuron myelination, gamma rhythms, and fear-related learning.

Introduction

Oligodendrocytes (OLs) are specialized glial cells of the central nervous system (CNS) that produce myelin, a lipid-rich substance that envelops axons, facilitates faster impulse propagation and supplies energy. Beyond their cellular functions as myelinating glia, several lines of evidence highlight the pivotal role of OL lineage cells in the underpinnings of cognitive processes[1]. Behavioral interventions, such as fear conditioning tasks or working memory training in mice, enhance both oligodendrogenesis and myelination, while genetic models impeding OL differentiation result in compromised fear memory recall and spatial memory consolidation[2–4]. In humans, the correlation between white matter changes and cognitive functions further supports the link between OLs, myelin plasticity and cognition[1,5].

While cognitive processes rely on the orchestrated coordination of neuronal networks and the resulting brain oscillations, the contribution of glial cells to this orchestration remains underexplored. Recent advances in mathematical modeling have provided insights into the potential role of myelin in promoting temporal synchronization among neuronal ensembles[6]. This theoretical framework is consistent with in vivo experiments conducted on the auditory system of dysmyelinated mouse models characterized by abnormal myelin formation, which exhibited delayed action potentials and loss of temporal precision[7]. However, the conversion of OL-induced neuronal synchronization into brain oscillation activities is still poorly understood. For instance, cuprizone demyelinated mice exhibit an enhanced rather than a decreased cortical theta power during phases of quiet wakefulness[8] and ablation of oligodendrogenesis affects in vivo electroencephalography (EEG) recordings even in the absence of task experience[9]. Moreover, besides myelin, the generation of new OLs themselves is crucial for coupling spindle oscillations in the prefrontal cortex with sharp wave ripple oscillations in the hippocampus following learning[2]. While these initial studies offer some insight, the influence of OL lineage cells on distinct oscillation frequencies that dictate cognitive performance remains elusive.

Cortical oscillations result from the dynamic interactions of excitatory and inhibitory neurons,

with gamma oscillations being of particular importance for cognitive function[10]. Notably, gamma-band

oscillations result from the fast synchronization of excitatory neuronal activity, a phenomenon largely

modulated by parvalbumin (PV) interneurons that provide a robust synaptic inhibition[11–13].

Interestingly, PV interneurons share a unique association with OL lineage cells, arising from common

embryonic origins[14]. During postnatal development, these interneurons form bona fide GABAergic

synapses with OL precursor cells (OPCs) which disappear after the second postnatal week in mice[15–17].

Recent evidence highlights the critical role of early communication between PV interneurons and

OPCs *via* GABA receptors in the myelination and maturation of PV interneurons, as well as in shaping

cortical inhibitory circuit function and mouse behaviour[18,19]. Moreover, PV interneurons constitute the

largest myelinated population of GABAergic interneurons in the cortex[20,21]. Considering the

relationship between PV interneurons, OL lineage cells and myelin, we propose that early postnatal

interactions between PV interneurons and OPCs play a crucial role in establishing prefrontal

synchronization and oscillations in adulthood, especially in the gamma-frequency range, which could

have significant implications for cognitive function.

To address our hypothesis, we genetically disrupted $GABA_A$ receptor-mediated synapses on

OPCs during early postnatal development and subjected adult mice to a fear conditioning task while

recording oscillations in the infralimbic (IL) region of the medial prefrontal cortex (mPFC), a key area

for processing emotional and cognitive information[1]. Our results demonstrate that early postnatal

disruption of OPC GABAergic synaptic signalling results in impairments in tone fear extinction

learning which arise from a reduction in low-gamma frequency power during tone fear extinction

sessions. These functional deficits are associated with PV interneuron dysmyelination and an

excitation/inhibition (E/I) imbalance in the adult mPFC. Efforts to restore the altered phenotype in

transgenic mice by local chemogenetic-mediated enhancement of PV interneuron activity in the mPFC

or by in vivo infusion of the pro-myelinating compound fractalkine (FKN) fail to recover learning and

memory deficits. Altogether, our findings reveal the role of early interneuron-OPC synaptic signalling

in shaping mPFC circuits and associated fear extinction learning and memory, while also highlighting

the long-lasting and irreversible impact of early OL-mediated disruptions on cognitive deficits.

Results

Early GABAergic OPC synapse disruption impairs tone fear extinction

To investigate the potential impact of early postnatal interneuron-OPC interactions on cognitive functioning in adulthood, we used the *NG2cre$^{ERT2+/-}$;Gcamp3$^{f/f}$;γ2$^{f/f}$* mice (γ2$^{f/f}$ mice) in which transient γ2 subunit-mediated GABAergic synapses of OPCs are inactivated by 4-hydroxytamoxifen (4-OHT) injections administered from postnatal day 3 (P3) to P5[22] (Supplementary Fig. 1a; see Online Methods). In agreement with our previous reports which demonstrated recombination at the γ2 locus and a partial reduction in OPC evoked GABAergic postsynaptic currents (I_{GABA}) in the somatosensory cortex[18,22], our current results reveal a near-complete abolishment of I_{GABA} in layer 5 OPCs within the IL region of acute mPFC slices of γ2$^{f/f}$ mice at P9-P13 (Supplementary Fig. 1b, 1c). This finding indicates a more pronounced impairment of interneuron-OPC synaptic signaling in the mPFC compared to the somatosensory cortex.

To assess whether γ2-containing GABA-A receptor-mediated currents persist in OPCs into adulthood, we recorded I_{GABA} in layer 5 OPCs of two-month old control and γ2$^{f/f}$ mice. In contrast to the pronounced γ2-dependent currents observed in young animals, evoked I_{GABA} remained detectable at similar levels in both groups and was insensitive to zolpidem (1 μM), a positive allosteric modulator selective for GABA-A receptors containing the γ2 subunit (Supplementary Fig. 1a, 1d, 1e). This pharmacological profile as well as the presence of a GABAergic current in the γ2$^{f/f}$ mice is consistent with prior reports indicating that, while adult OPCs in the cortex and hippocampus express functional GABA-A receptors, these cells lack the γ2 subunit[23,24]. Together, these data confirm that γ2-mediated GABAergic synapses are developmentally restricted and absent in adult OPCs in the mPFC. To further validate the efficiency of OPC targeting in this region, we quantified recombination by immunolabeling Olig2 cells and enhancing GCaMP3 detection with an anti-GFP antibody in four-month old mice. We

found 65.5% of recombinant OL lineage cells in control mice and 63.4% in $\gamma2^{f/f}$ mice in the mPFC,

indicating efficient and comparable recombination for both groups (Supplementary Fig. 2a).

After confirming the early functional inactivation of OPC GABAergic synapses in this cortical

region in $\gamma2^{f/f}$ mice, we next investigated whether this developmental synaptic alteration would affect

mPFC-dependent cognitive functions in adulthood. To assess this, we first evaluated four-month-old

$\gamma2^{f/f}$ mice using a fear conditioning task, a widely used paradigm to investigate mPFC-dependent

processes such as learning, memory, emotional regulation, and perception[25].

To evaluate learning and memory processes associating environmental cues with aversive

experiences, adult mice subjected to 4-OHT injections at P3-P5, underwent a behavioral test where a

30 s conditioned tone stimulus was paired with a foot shock delivered during the last 2 s of the tone in a

specific context (Fig. 1a). In the conditioning session, adult control and $\gamma2^{f/f}$ mice exhibited similar fear

memory acquisition, albeit slightly faster in $\gamma2^{f/f}$ mice (Fig. 1b). No significant differences were found

between the two groups during the context extinction session (Fig. 1c). However, in the first tone

extinction session, although the baseline level (BL, first trial block) of fear expression were comparable

in control and $\gamma2^{f/f}$ mice (Fig. 1d), control mice showed faster and more extensive fear extinction

learning by the end of the protocol (Fig. 1d). To test whether this deficit in fear extinction learning was

persistent, we performed a second tone extinction session on the following day (extinction retrieval).

We found that, in this session, control mice exhibited significantly lower baseline values and reduced

initial freezing levels compared to the first tone extinction session, indicating successful extinction

retrieval (Fig. 1e). The tone extinction learned in the first session was therefore retained and

successfully retrieved over time in control mice. In contrast, $\gamma2^{f/f}$ mice showed persistently high

baseline values and a weak extinction, with freezing levels remaining similar to those observed in the

first tone extinction session (Fig. 1e), indicating that these mice failed to improve or consolidate fear

extinction learning between sessions. Of note, no significant sex differences were observed between

male and female mice in both control and γ2$^{f/f}$ groups (Supplementary Fig. 3a-h). To confirm that the

observed behavioral deficits specifically result from early postnatal recombination rather than to the

presence of the floxed γ2 alleles, we reproduced the same fear conditioning paradigm in γ2 mice that

were not injected with 4-OHT. These animals retain normal developmental expression of γ2 subunits in

OPCs during development, while carrying the same Cre and γ2 alleles as the injected littermates. Non-

injected γ2 mice showed no differences in fear conditioning behavior compared to controls

(Supplementary Fig. 5a–e; Fig. 1b–f), indicating that the presence of the floxed alleles does not affect

extinction behavior. We then assessed the effect of inducing recombination specifically in adulthood

(Supplementary Fig. 5a). In this condition, no significant differences in tone extinction learning or

retrieval were observed compared to non-injected γ2 mice (Supplementary Fig. 5b-e). These findings

indicate that the behavioral deficits depend on early postnatal disruption of γ2-mediated GABAergic

signaling in OPCs, and that recombination induction in adulthood, when the γ2 subunit is not expressed

in OPCs, has no effect.

Given that the deficit was specific to tone extinction and not observed in context extinction, we

considered whether this could be due to auditory impairments. To address this, we measured startle

responses to noise stimuli of varying intensities, including the one used during fear conditioning

(Supplementary Fig. 4a, 4b). Although this test does not assess potential hidden hearing loss[26], we

found that both groups exhibited comparable startle responses to random noise of varying dB levels.

This suggests that major auditory deficits are unlikely to account for the observed differences in fear

conditioning (Supplementary Fig. 4c, 4d). Building on these findings, we further analyzed the

extinction learning deficit in γ2$^{f/f}$ mice throughout different stages of the retrieval session

(Supplementary Fig. 3i). The extinction learning deficit in γ2$^{f/f}$ mice was predominant at early and

intermediate stages, but it was also evident at the late stage (Fig. 1f), showing their impaired ability to

acquire and consolidate an extinction memory.

Our results indicate that $\gamma2^{f/f}$ mice exhibit reduced fear learning and memory capabilities,

particularly in tone fear extinction learning, while no major auditory impairments were detected. These

results suggest that the disruption of OPC GABAergic synaptic signaling during the early postnatal

period leads to long-term cognitive impairments in adulthood, specifically affecting processes related

to learning and memory, without evidence of substantial auditory deficits.

Excitation-inhibition imbalance in the adult mPFC of $\gamma2^{f/f}$ mice

The long-lasting cognitive impairments observed in $\gamma2^{f/f}$ mice, particularly during tone fear

extinction, suggest that these deficits may be caused by underlying neurophysiological alterations.

Therefore, we specifically tested the local excitation-inhibition (E/I) balance in the mPFC that is

involved in regulating cognitive processes such as learning, memory, and emotional regulation[10,12].

This hypothesis is further supported by the known role of mPFC PV interneurons in regulating the E/I

balance by modulating excitatory inputs, which are necessary for higher-order cognitive functions[27].

Since fear conditioning experiments revealed specific deficits in tone fear extinction in $\gamma2^{f/f}$ mice, we

focused on the IL region of the mPFC, a critical region for fear extinction memory processes[28].

To test the E/I balance in the IL region of adult $\gamma2^{f/f}$ mice, we performed patch-clamp recordings

of layer 2/3 pyramidal cells in acute slices, while stimulating in layer V with an extracellular electrode.

The stimulation elicited both excitatory (eEPSCs) and inhibitory (eIPSCs) postsynaptic currents in

mature IL circuits (Fig. 2a). Our results revealed that, three months after inactivating the $\gamma2$ subunit-

mediated GABAergic synapses in OPCs, $\gamma2^{f/f}$ mice exhibited a significant reduction in the mean

amplitudes of eIPSCs, but not eEPSCs, in pyramidal neurons (Fig. 2a-b). Consequently, the E/I ratio

was higher in $\gamma2^{f/f}$ mice, causing a strong E/I imbalance (Fig. 2b).

To assess whether the E/I imbalance observed in the mPFC reflects a broader effect or is more

regionally restricted, we extended our analysis to two additional brain regions involved in fear

expression and extinction: the amygdala and hippocampus. First, we recorded from principal neurons

in the basolateral amygdala (BLA), while stimulating the lateral amygdala (Fig. 2c, d). Similarly, we

recorded from CA1 pyramidal neurons in the hippocampus, while stimulating Schaffer collateral inputs

from CA3 (Fig. 2e, f). Unlike the IL region, we found no significant differences in eIPSC or eEPSC

amplitudes, nor in the E/I ratio between control and $\gamma2^{f/f}$ mice in either the BLA or CA1 regions (Fig.

2d, 2f).

To further evaluate the functional specificity of these alterations, we complemented these

recordings with region-specific behavioral tests targeting the amygdala and the hippocampus: the

elevated plus maze and the novel object recognition, respectively. In both tasks, $\gamma2^{f/f}$ mice performed

similarly to controls, showing no significant differences in anxiety levels and recognition memory

(Supplementary Fig. 5f, 5g). These results are consistent with the lack of synaptic alterations in these

regions. Moreover, they align with our fear conditioning results, in which $\gamma2^{f/f}$ mice exhibited normal

fear acquisition, a process largely mediated by the amygdala, and normal contextual extinction, which

strongly depends on hippocampal function (Fig. 1b, 1c).

Since the proper regulation of E/I balance is essential for normal cognitive processing, the

observed alterations in the mPFC functioning suggest a functional impairment in this region due to

early interneuron-OPC synapse disruption. These findings indicate that the synaptic E/I imbalance is

more prominent in the mPFC than the amygdala and hippocampus, suggesting that the mPFC is

particularly sensitive to early developmental perturbations.

PV interneuron dysmyelination in the adult mPFC of $\gamma2^{f/f}$ mice

The observed cognitive and E/I ratio deficits prompted us to investigate whether early

disruption of interneuron-OPC synapses results in persistent myelination deficits, particularly in PV

interneurons known for their highly plastic myelination[29]. Previous findings in the somatosensory

cortex demonstrated that $\gamma 2^{f/f}$ mice exhibit myelination defects in PV interneurons during juvenile

stages[18]. Here, we extended this analysis to the mPFC in adulthood to determine whether these

abnormalities persist and specifically affect the myelination of PV interneurons, the primary source of

transient GABAergic synaptic input onto OPCs[16,17]. We found that $\gamma 2^{f/f}$ mice displayed normal overall

myelin coverage and intensity of MBP staining in IL cortical layers (Fig. 3a-c). We then analyzed the

length of individual myelin segments, or internodes, along PV$^+$ axons in the same region. This was

achieved by reconstructing high resolution images of axonal segments expressing MBP, PV and the

axonal marker SMI-312 (Fig. 3d). Measurements revealed a statistically significant increase in the

mean internode length of PV$^+$, but not PV$^-$, axons of $\gamma 2^{f/f}$ mice compared to controls (Fig. 3e, 3f). The

myelination defect observed in PV$^+$ axons was not accompanied by alterations in either the cell density

or distribution of PV interneurons across mPFC layers, nor by changes in the total interneuron

population identified by GAD67 expression (Supplementary Fig. 6a-e). In addition, quantification of

PV signal within SMI-312$^+$ axonal compartments—used as a proxy for PV$^+$ axon density—did not

reveal significant differences between control and $\gamma 2$ mice (Supplementary Fig. 6f, 6g). Furthermore,

we found no differences in the cell density or proportion of Olig2$^+$/CC1$^-$ OPCs and Olig2$^+$/CC1$^+$ OLs

between control and $\gamma 2^{f/f}$ mice (Supplementary Fig. 2b-d). These results suggest that the early

disruption of $\gamma 2$-mediated synapses in OPCs impacts predominantly PV interneuron myelination,

without affecting interneuron survival or the maturation of OL lineage cells in the mPFC.

Although OPCs form GABAergic synapses only transiently during postnatal development and

these synapses are absent in adulthood[15,16,23], myelination defects in PV interneurons persists into

adulthood of $\gamma 2^{f/f}$ mice. Notably, these alterations are specific to PV interneurons as we found no

significant changes in the overall mPFC myelination pattern, length of internodes of PV$^-$ axons, OL

lineage cell densities, PV interneuron survival, the distribution of interneurons within cortical layers or

of the estimated PV$^+$ axon density.

Early GABAergic OPC synapse loss irreversibly impairs fear extinction

The persistent PV interneuron dysmyelination as well as the electrophysiological and cognitive

impairments observed in $\gamma2^{f/f}$ mice raised the question of whether these deficits could be corrected

through targeted interventions. To this end, we explored two distinct strategies aimed at addressing the

underlying structural and behavioral abnormalities: enhancing PV interneuron activity and promoting

myelination.

Previous research has demonstrated that using a modified form of the human M3 muscarinic

(hM3Dq) receptor, activated by injections of the inert clozapine metabolite clozapine-N-oxide (CNO)

for 14 days, can specifically promote activity-dependent myelination of PV interneurons in the adult

mPFC[30]. In addition, acutely increasing PV interneuron activity with a chemogenetic approach should

also rebalance the observed reduced cortical inhibition, improving E/I impairments (Fig. 2a, 2b).

Building on this finding, we explored whether selectively targeting PV interneurons with a

chemogenetic approach could effectively address dysmyelination and fear learning deficits (Fig. 4a).

We performed bilateral intracranial injections into the IL region of the mPFC in $\gamma2^{f/f}$ mice using either

pAAV-S5E2-dTom-nlsdTom (dTom) or pAAV-S5E2-hM3D(Gq)-P2A-dTomato (hM3Dq-dTom) viral

vectors[31] (Fig. 4a, 4b). Approximately 80% of cells infected by these viral constructs were identified as

PV$^+$ interneurons in this region (Fig. 4b, 4c). To confirm the functional efficacy of the hM3Dq-

mediated activation of PV interneurons, we first performed whole-cell patch-clamp recordings in acute

mPFC slices from $\gamma2^{f/f}$ mice injected with pAAV-S5E2-hM3D(Gq)-P2A-dTomato. The dTom$^+$ neurons,

that express hM3Dq, exhibited an increased excitability in the presence of CNO while nearby dTom$^-$

pyramidal neurons did not, indicating that CNO effectively enhanced PV interneuron activity

(Supplementary Fig. 7a-d). This supports the specificity and efficacy of hM3Dq-mediated

chemogenetic manipulation. We then performed twice-daily in vivo injections of CNO over two weeks

in dTom-injected and hM3Dq-dTom-injected $\gamma2^{f/f}$ mice (Fig. 4a). Our results demonstrate that CNO

administration did not induce any significant changes in either the coverage or mean intensity of the

MBP fluorescence signal compared to those in dTom-injected $\gamma2^{f/f}$ mice (Fig. 4d-f). Furthermore, we

did not detect any reduction of PV interneuron internode length in hM3Dq-dTom in $\gamma2^{f/f}$ mice (Fig. 4g),

suggesting that increased PV interneuron activity in adulthood does not restore aberrant internode

length. To examine whether enhancing PV interneuron activity in vivo, independent of the effect on

interneuron dysmyelination, ameliorates cognitive deficits, we conducted the fear conditioning task in

both CNO-treated controls and $\gamma2^{f/f}$ mice. As expected, no differences were observed in the acquisition

or context sessions (Supplementary Fig. 8a, 8b). Consistent with the lack of effect on PV interneuron

internode length, we observed no significant differences during the tone extinction and extinction

retrieval sessions between the two viral injected groups, nor when compared to naive $\gamma2^{f/f}$ mice (Fig.

4h, 4i).

Our second strategy consisted in enhancing myelination in young adult mice, a developmental

period when myelination is still actively progressing in the mPFC and cognitive deficits associated with

neurodevelopmental disorders are often recognized due to the emergence of visible symptoms[1]. We

reasoned that targeting this critical period might offer greater potential for modifying myelination and

improving cognitive deficits because: (1) myelination plasticity persists well beyond early postnatal

stages in cortical regions, and (2) therapeutic interventions during active myelination phases might

rescue structural and functional deficits caused by earlier disruptions. We focused on fractalkine

(FKN), a well-studied chemokine known for its role in microglial function[32], but also shown to

promote OPC differentiation into mature OLs during cortical development through its release from

GABAergic interneurons[33]. Recent studies demonstrate that FKN can promote myelination under both

healthy and demyelinating conditions when administered exogenously[34,35]. We thus exploited the pro-

myelinating potential of FKN to examine whether its exogenous application in mice could compensate

for PV interneuron dysmyelination in the mPFC and mitigate the associated learning and memory

deficits observed during tone fear extinction. To this end, we performed in vivo FKN infusions into the

lateral ventricles of P42-46 mice over a 25-28-day period (Fig. 5a). Our results revealed that in vivo

FKN infusions in $\gamma 2^{f/f}$ mice increased both the coverage and mean intensity of MBP fluorescent signal

in the mPFC of infused mice compared to vehicle-treated mice, indicating an enhanced myelination

(Fig. 5b-d). However, measurements of internode length of $PV^+/SMI-312^+$ axons in the same animals

showed no difference between the two groups, suggesting that this intervention is insufficient to repair

PV interneuron-specific myelin abnormalities in $\gamma 2^{f/f}$ mice (Fig. 5e). To assess whether FKN-induced

myelination enhancement could still result in cognitive improvements, we conducted the fear

conditioning task in $\gamma 2^{f/f}$ mice treated with either vehicle or FKN for comparison. As expected, no

differences were observed in the acquisition or context sessions (Supplementary Fig. 8c, 8d). However,

no improvements were observed in the tone extinction or extinction retrieval sessions either (Fig. 5f,

5g), indicating that enhancing ongoing mPFC myelinationfails to rectify PV interneuron

dysmyelination or improve the associated learning and memory deficits.

Our findings show the irreversible cognitive impairments resulting from early disruption of

OPC GABAergic synapses. Despite targeted interventions aimed at increasing PV interneuron activity

or enhancing myelination, we observed no improvements in PV interneuron internode length and fear

extinction learning and memory. Once early interneuron-OPC interactions are compromised during a

critical developmental window, subsequent efforts to reverse structural and functional deficits become

particularly challenging. The early postnatal period, particularly the first two postnatal weeks[16], is

therefore essential for ensuring proper cognitive development and function.

Specific alterations in gamma oscillations during fear extinction

The E/I balance in the mPFC, which is disrupted in $\gamma2^{f/f}$ mice, plays a fundamental role in generating gamma oscillations (30-90 Hz), high-frequency neural rhythms which support prefrontal-dependent cognitive processing. These oscillations arise from the rapid synchronization of excitatory activity, with PV interneurons playing a central role by providing perisomatic inhibition onto pyramidal cell ensembles, which fine-tunes network discharges[10,12,13]. Deficits in E/I balance and PV interneuron myelination could potentially disrupt gamma oscillations, impacting cognitive functions. To investigate this possibility, we recorded local field potentials (LFPs) from the IL region of the mPFC in freely moving mice using an implanted tetrode (Fig. 6a), of which the placement was confirmed postmortem through histological analysis (Fig. 6b). After 10 days of recovery post-implantation, mice were first habituated to wearing the recording cable in their home cage for three days. Then, they were placed in an open field environment and later subjected to the fear conditioning task, during which LFP signals were recorded (Fig. 6a, 6c). We analyzed these signals using fast Fourier transformation (FFT) to generate spectrograms and power spectra allowing us to extract different oscillatory activities during the behavioral tasks (Fig. 6c-f, 6i, 6j).

In vivo LFP recordings during the open field test detected both theta (4-12 Hz) and gamma (30-90 Hz) oscillations in the mPFC of control and $\gamma2^{f/f}$ mice (Fig. 6c). Contrary to our hypothesis that disrupted myelination would impair gamma oscillations, there were no significant differences in theta or gamma oscillatory activity between the two groups, regardless of whether the mice were in the center or periphery of the arena (Fig. 6e-h). These results are consistent with the absence of significant differences in exploratory behavior between control and $\gamma2^{f/f}$ mice, including distance traveled, time spent, immobility time in the center and periphery of the arena, as well as the number of crossings or average speed (Supplementary Fig. 9a-f). These findings indicate that deficits in PV interneuron

myelination did not result in alterations in either oscillatory activity of IL region or exploratory behavior under the conditions of the open field task.

The consequences of PV interneuron dysmyelination in the mPFC may be task-specific, potentially impacting complex processes rather than general locomotor or exploratory behaviors. These same implanted mice were therefore subjected to the fear conditioning protocol which is designed to assess more cognitively demanding processes (Fig. 6a). In vivo LFP recordings were performed during the tone extinction retrieval session, when more pronounced behavioral differences between control and $\gamma2^{f/f}$ mice were observed (Fig. 1e). During this session, we optimized recordings by using a Faraday cage and minimizing external noise. Analysis of the oscillatory activity revealed no significant differences in the relative power of theta oscillations between the two groups during tone presentation and inter-trial intervals (ITI) (Fig. 6i-k). However, $\gamma2^{f/f}$ mice exhibited a significant reduction in gamma oscillations throughout both periods, suggesting a specific disruption of neural dynamics at this frequency range (Fig. 6i-j, 6l). These results indicate that early postnatal disruption of interneuron-OPC signaling leads to persistent PV interneuron dysmyelination and E/I imbalance in the mPFC, which in turn drive alterations in gamma oscillations and the subsequent impairments in fear extinction learning observed in $\gamma2^{f/f}$ mice.

The absence of significant gamma oscillation changes in the open field, in contrast to the pronounced reduction observed during fear extinction retrieval, suggests that PV interneuron dysmyelination predominantly affects higher cognitive processes, without necessarily disrupting basal neural activity or basic behaviors like exploration or locomotion.

Low-gamma oscillation disruption during fear extinction retrieval

Gamma oscillations can be subdivided into low and high frequency ranges, each contributing distinctly to cognitive processes. High-gamma frequencies have been associated with encoding

processes, whereas low-gamma frequencies support retrieval in human episodic memory[36]. To

investigate whether the differences in gamma oscillations observed during fear extinction retrieval

between control and $\gamma 2^{f/f}$ mice were frequency-specific, we conducted a detailed analysis of both low-

gamma (30-45 Hz) and high-gamma (55-90 Hz) power. Our results revealed a significant reduction in

the relative power of low-gamma oscillations in $\gamma 2^{f/f}$ mice during both tone presentation and ITI of fear

extinction retrieval, while high-gamma oscillations showed no significant differences (Fig. 7a, 7b).

These results suggest that deficits in neuronal processing in mPFC of $\gamma 2^{f/f}$ mice are primarily associated

with alterations in low-gamma oscillations.

Gamma oscillations are known to couple strongly with theta oscillations during cognitive

processes, playing a key role in coordinating network-wide communication[37]. Our analyses reveal that

the strength of the theta phase-low-gamma amplitude and theta phase-high-gamma amplitude coupling

increased during tone presentation compared to the ITI in both control and $\gamma 2^{f/f}$ mice (Supplementary

Fig. 10). This increase was observed across the entire session as well as at the different stages within

the session (Supplementary Fig. 10), indicating a robust network activation and an ability in both

groups to discriminate the tone stimuli. This strong theta–gamma coupling serves as a hallmark of

effective cognitive processing, especially since no significant changes in coupling were observed

between the center and periphery of the arena in the open field task (Supplementary Fig. 9g, 9h).

However, despite the clear increase in coupling during tone presentation compared to ITI, there were

no significant differences between control and $\gamma 2^{f/f}$ mice in the tone/ITI coupling ratio, either for low-

gamma or high-gamma oscillations (Supplementary Fig. 10c, 10g). The lack of coupling defects shows

that the coordination of broad network interactions remains intact, despite a decrease in low-gamma

power.

To further dissect the temporal dynamics of gamma power alterations, we analyzed the relative

power of low- and high-gamma oscillations at different stages of the retrieval session: early,

intermediate, and late (Fig. 7c, 7d). We found that low-gamma power was significantly reduced during

the early and intermediate stages in $\gamma2^{f/f}$ mice compared to controls (Fig. 7c) during tone presentations

and ITI, while it remained unaffected during the late stage. These early and intermediate reductions in

low-gamma power correspond with the higher freezing levels observed in $\gamma2^{f/f}$ mice at these stages

(Fig. 1f), suggesting a temporal relationship between low-gamma disruption and behavioral deficits. In

contrast, high-gamma power exhibited a trend toward reduction that did not reach statistical

significance and remained comparable between control and $\gamma2^{f/f}$ mice throughout all three stages of the

retrieval session (Fig. 7d), further supporting the conclusion that the observed alterations are more

pronounced in the low-gamma frequency range.

In summary, our findings suggest that low-gamma oscillations in the mPFC play a more critical

role than high gamma oscillations in supporting fear extinction retrieval. The reduction in low-gamma

power, particularly during the early and intermediate stages of the retrieval session in $\gamma2^{f/f}$ mice, points

to its essential role in enabling neuronal network activity required for the successful fear extinction

learning and memory. This reduction, alongside the lack of changes in theta-gamma coupling, is

consistent with an intracortical mPFC network dysfunction, resulting from PV interneuron impairments

which are key for intracortical projections.

Discussion

This study reveals that the early postnatal period is a critical window for interneuron-oligodendroglia interactions, during which disruption of OPC GABAergic synaptic signaling in $\gamma 2^{f/f}$ mice leads to persistent impairments in local neuronal network function and cognitive performance in adulthood. Specifically, deficits in learning and memory during tone fear extinction are associated with disrupted low-gamma oscillations in the IL region of the mPFC. These alterations arise from early dysfunctions in interneuron-OPC signaling, leading to impaired PV interneuron myelination and disrupted neuronal E/I balance. Our findings show the necessity of this developmental phase for oligodendroglia and PV interneuron myelination to sustain adult low-gamma oscillations that are critical for tone fear extinction learning and memory. Overall, this study emphasizes the importance of early postnatal PV interneuron myelination in the maturation of inhibitory circuits and its role in maintaining cortical low-gamma rhythms essential for specific cognitive functions. Consistent with this, adult-specific induction of $\gamma 2$ deletion had no effect on extinction behavior, suggesting that this signaling pathway is only functionally relevant during early postnatal development. This is likely due to the fact that functional $\gamma 2$-containing GABA-A receptor-mediated synaptic inputs are not present in adult OPCs[23,24].

Defects in OL lineage cells and myelination are increasingly associated with a variety of cognitive deficits[1]. For example, social isolation, whether occurring immediately after weaning or during adulthood, results in hypomyelination of the mPFC[38,39], while early maternal separation impairs emotional responses and object recognition in adult mice, linked to premature OPC differentiation in this cortical region[40]. Proper myelination is also critical for emotional regulation and memory processes, including fear conditioning and extinction, as disruptions in de novo OL production or myelin formation prolongs fear responses and hampers the consolidation of contextual fear memories[2,3]. However, while cognitive impairments are often attributed to pronounced

hypomyelination or demyelination, the impact of subtle, cell-type-specific dysmyelination, such as that in PV interneurons, remains largely unexplored. PV interneurons play a key role in regulating gamma oscillations and maintaining the E/I balance, both of which are essential for coordinated neuronal network function and cognitive performance[10,12,27]. Our findings suggest that even mild myelination defects in PV interneurons during critical developmental windows can lead to persistent alterations in inhibitory circuitry, impairing cognitive function. The alterations in low-gamma oscillations observed in our study, without significant changes in high-gamma oscillations, suggest that specific deficits in PV interneurons can selectively impact distinct gamma frequency bands. This cell-specific dysmyelination could represent a previously underestimated mechanism underlying cognitive deficits and emotional dysregulations present in neurodevelopmental and psychiatric disorders, where disruptions in PV interneuron function and gamma oscillations are commonly implicated[1]. Our findings open the possibility that subtle myelination abnormalities in PV interneurons may contribute to the E/I imbalance and network dysfunctions characteristic of conditions such as schizophrenia and autism spectrum disorders[1]. The contribution of oligodendroglia to these conditions has likely been underestimated and deserves further investigation.

Recent studies in healthy mice demonstrate that behavioral training and stimuli can induce subtle changes in myelination. Motor-skill learning induces myelin remodelling, including the elongation of nodes of Ranvier[41], highlighting the importance of a fine-tuned myelination during learning processes. Furthermore, the length of nodes of Ranvier is modulated in distinct ways by interventions like repetitive transcranial magnetic stimulation or spatial learning[42]. Despite these insights into activity-dependent delicate myelination changes, the role of oligodendroglia and myelin in modulating neuronal network activity and cortical oscillations during the execution of cognitive processes in health and disease is poorly understood. A recent study using cuprizone-treated mice has shown that prolonged impairments in social and spatial working memory, as well as in cognitive

flexibility, persist after remyelination. These impairments correlate with long-term desynchronized activity between the hippocampus and the mPFC, characterized by lasting disruptions in gamma synchronization and ripple/slow oscillation coupling, a condition that develops during the demyelination phase and is only partially restored during remyelination[43]. Furthermore, fiber photometry has revealed that animals unable to form new myelin exhibit a lack of the time-dependent modulation in mPFC neuronal activity seen in control animals, which initially show a decrease and then a progressive increase in activity during transitions from mobility to freezing[3]. Active oligodendrogenesis and myelination are therefore required for the proper evolution of prefrontal dynamics associated with fear memory recall. Moreover, the disruption of de novo production of OLs immediately before contextual fear conditioning also impairs the coupling between spindle oscillations in the prefrontal cortex and sharp wave ripple oscillations in the hippocampus, independent of myelin deficits[2]. Therefore, both OL production alterations and myelin deficiencies are linked to impaired neuronal responses in brain regions activated during cognitive tasks.

While the recognition of the involvement of oligodendroglia in cognition is growing, the mechanisms by which these cells influence specific neuronal subtypes and network dynamics during the execution of a cognitive task remain elusive. Our results demonstrate that early disruption of OPC GABAergic signaling results in a selective dysmyelination of PV interneurons, without major changes in overall oligodendrogenesis or myelination, and altered low-gamma oscillation activity during tone fear extinction. This defect does not uniformly impact neuronal networks or cognitive processes, as high-gamma oscillations show no significant change, and no impairments were observed in fear acquisition, context fear extinction, or exploratory behavior in the open field. The specific effects observed in our study may be explained by the fact that myelination characteristics such as patterns, myelin content, internode and node lengths vary across specific neuronal identities and for a specific cell type in different brain areas[21,44], which can lead to different consequences. This selective effect

may arise from differences in myelination requirements in mPFC subregions, with the IL region depending on low-gamma oscillations critical for extinction memory, while the prelimbic (PL) region, involved in fear learning, may rely on different oscillatory patterns and thus be less affected by PV interneuron dysmyelination. In any case, our findings emphasize the critical role of precise myelination patterns of specific neuronal subtypes in maintaining proper network dynamics during distinct cognitive functions. While neuronal contributions to cognitive processes are undisputable, our study positions oligodendroglia and myelination as pivotal elements driving the construction of prefrontal-dependent low-gamma oscillations necessary for a specific cognitive process. In addition to oligodendroglia, other types of glial cells, such as astrocytes, also play a significant role in regulating neuronal network dynamics. Recent findings suggest that GABAergic signaling to astrocytes, although non-synaptic in this case, also influences low-gamma oscillations without impacting high-gamma oscillations[45], further strengthening the importance of glial-neuronal interactions in maintaining cognitive functions. These findings highlight the importance of glial cells in fine-tuning network oscillations important for higher-order cognitive functions.

While slow oscillations like theta oscillations result from the coordinated activity of widely distributed neuronal ensembles, fast-frequency oscillations like gamma oscillations are considered a local phenomenon, involving smaller tissue volumes[1]. Cortical PV interneurons contribute to various frequency oscillations, but their role in gamma oscillations is crucial. This is due to their ability to discharge action potentials at high frequencies and their extensive local synaptic connectivity with surrounding excitatory neurons, which facilitates rapid and synchronized neuronal activity essential for gamma oscillations[11–13]. Our study revealed that PV interneuron dysmyelination in $\gamma2^{f/f}$ mice are associated with a reduced inhibitory drive causing an E/I imbalance and alterations of low-gamma oscillations in the IL region during tone fear extinction retrieval. A critical involvement of PV interneurons in low-gamma activity during cognitive processes is supported by a previous study

showing a rescue of low-gamma oscillation alterations and social deficits by the optogenetic

stimulation of PV interneurons at 40 Hz in a model of autism spectrum disorders characterized by a

Neuregulin-3 mutation[46]. Moreover, the optogenetic activation of PV interneurons at 30 Hz allows for

the preservation of gamma frequency oscillations in control, but not in mice with widespread

demyelination[8]. In these models as well as in $\gamma2^{f/f}$ mice, a decreased PV interneuron firing rate is also

observed[8,18,46]. Our findings uniquely highlight the role of PV interneuron myelination in sustaining

low-gamma oscillations during specific cognitive tasks, aligning with previous studies that explore the

function of these interneurons in cortical processing.

PV interneuron myelination remains particularly plastic throughout life, making these neurons

more susceptible to myelin deficiencies. For instance, monocular deprivation results in experience-

dependent adaptive myelin remodeling specifically in cortical PV interneurons, but not in glutamatergic

projection neurons[29]. Furthermore, myelination defects occur specifically in PV interneurons of the

mPFC in a rat model displaying schizophrenia-like behaviors[47], and demyelination induces selective

vulnerability of GABAergic interneurons in both progressive multiple sclerosis and a mouse model of

local cortical demyelination[48]. In a transgenic mouse, mild hypomyelination due to the loss of

oligodendrocyte ErbB receptor signaling results in a decreased PV expression and a reduction in

inhibitory but not excitatory synaptic connections, causing an E/I imbalance in L2/3 neurons of the

auditory cortex[49]. Thus, PV interneuron myelination contributes to the pathological mechanisms

underlying various neurological disorders, making it a novel and attractive pharmacological target[50].

Interestingly, it was previously shown that chemogenetic activation of hippocampal PV interneurons

adult mice subjected to a modified fear conditioning protocol with severe electric foot shocks reduced

fear generalization and restored myelination of PV interneuron axons[51]. Nevertheless, altering early

interneuron-oligodendroglia interactions, myelination, and neuronal plasticity during critical periods

can generate lasting cognitive and behavioral effects that may be challenging to reverse. In our study,

targeted interventions to increase PV interneuron activity or enhance global myelination in γ2[f/f] mice did not repair the defect in PV interneuron internode length or lead to improvements in fear extinction memory. This suggests that, once early interneuron-OPC interactions are compromised, subsequent attempts to enhance myelination or promote PV interneuron activity are ineffective in reversing the resulting cognitive deficits. The critical developmental window for these interactions, occurring transiently in the first two postnatal weeks[16], appears crucial for establishing proper cognitive development and function. Preserving the integrity of transient interneuron-OPC interactions during this period appears essential to prevent long-lasting and irreversible cognitive deficits. It should be noted that most psychiatric treatments begin after the onset of symptoms, which often occur well after these critical periods, making it challenging to reverse such early-established myelination deficits. Additional evidence suggests that early but transient dysfunctional events of interneurons can be deleterious to cortical function in adulthood. Kessaris et al. (2021)[52] demonstrated that an aberrant overproduction of PV interneurons during embryogenesis has lasting effects that persist into critical periods of cortical development, despite the normalization of cell density postnatally. This transient excess of interneurons during early development results in behavioral abnormalities in adulthood, indicating that the timing of interneuron clearance is essential for the proper maturation of neural circuits that underlie adult behavior. In humans, early life adversity correlates with increased OPC-mediated perineuronal nets production around PV interneurons in the ventromedial PFC; as perineuronal nets regulate the closure of developmental windows of cortical plasticity, this potentially limits cortical circuit neuroplasticity and promotes maladaptive behaviors[53]. Together, these findings emphasize the challenges in reversing cognitive deficits once developmental processes are disrupted. An interesting line for future research in myelin biology would be to explore how abnormalities in PV interneurons and OL lineage synergize, potentially increasing the risk of neurodevelopmental and psychiatric disorders.

 Methods

Transgenic mice and induction of Cre-mediated recombination

All experiments adhered the European Union and institutional guidelines for the care and use of

laboratory animals. Ethical approval was obtained from the French ethical committee for animal care of

university Paris Descartes (Committee N°CEEA34) and the Ministry of National Education and

Research (Project No: 202010131827131). The experimental approach involved $NG2cre^{ERT2+/-}$

$;Gcamp3^{f/f};\gamma2^{f/f}$ mice ($\gamma2^{f/f}$ mice) to inactivate $\gamma2$-GABA$_A$ receptor-mediated synapses in OPCs while

the $NG2^{creERT2+/-};Gcamp3^{f/f}$ mice were used as controls[18,22]. Genotyping was conducted by PCR using

primers specific to the different alleles, and Cre expression was induced from P3 to P5 by daily

intraperitoneal injections of 0.2 mg of 4-OHT (Tocris Bioscience) as previously described[22]. The Cre

expression in adult mice (P110-P112) was induced two weeks prior to the fear conditioning test by

administering tamoxifen (1 mg in Miglyol oil, Caesar & Loretz, Germany) intraperitonealy once daily

for three consecutive days. Animals were housed in controlled conditions with 12 h light/dark cycle,

ad libitum access to food and water, and in average ambient temperature of 21°C and 45% humidity.

Both females and males were used at different postnatal stages from P10 to P134.

Behavioral assays

Anxiety and locomotion were assessed in a 50 cm × 50 cm x 45 cm open field arena, where mice

were placed for 10 minutes under a 50 Lux illumination. An infrared camera monitored animal

movements and the ViewPoint behavioral analysis software processed the data (version 5.31.0.8,

ViewPoint, Behavior Technology, France). The fraction of time spent in an outer area of 2500 cm^2 over

an inner square area of 900 cm^2 was calculated. The open field test was used to assess locomotion by

measuring distance traveled, velocity, time spent in the inner (center) or outer (periphery) area, and number of crossings in the center area.

The fear conditioning paradigm consisted of four consecutive days of behavioral testing in a chamber of 30 cm × 25 cm × 25 cm made of two aluminum sidewalls, acrylic front and rear walls and a floor of metal bars of 3.2 mm diameter separated by 7.9 mm. An infrared camera, controlled by a computerized system interface, monitored behavior and freezing (percentage of time of immobility) using the Videofreeze software (version 2.7.2.117, MED Associates, St. Albans, VT, USA). On day-1 (fear conditioning), mice (aged P95-P103) were individually placed into a dark conditioning chamber, odorized with 1% carvone (mint odor). After a baseline period of 3 min, animals received five pairings of an auditory conditional stimulus (CS; 30 s, 5 kHz, 85 dB) with the unconditioned stimulus (US) applied at the end of the tone (2 s, 0.5 mA foot-shock) with a 2 min ITI period. On day-2 (contextual fear extinction), mice were introduced in the same conditioning chamber, odorized with carvone, and allowed to explore for 18 min without presentation of CS or US. On day-3 (tone fear extinction), mice were placed in chambers with a modified context and odor (0.5% isopentyl acetate; banana odor). After a baseline period of 3 min, the CS alone was presented 40 times (30 s), separated by 5 s ITIs. On day-4 (tone fear extinction retrieval), mice were subjected to the same protocol of day 3. The open-field arena and fear conditioning chambers box were wiped with 70% ethanol and air-dried between mice.

Freezing behavior was automatically quantified in VideoFreeze software (v2.27). Animal movement was analyzed with the default settings (motion threshold: 18 arbitrary units, sample rate: 30 frames per second, linear method, minimum freeze duration: 30 frames), with freezing defined as immobility lasting at least 1 s. Freezing responses were calculated across intertrial-intervals (ITIs) on day 1, averaged across periods of 2 minutes on day 2, and those on day 3 and day 4 were calculated using the average freezing response to five consecutive CS (5-trial blocks). The tone fear extinction

retrieval session (day 4) was divided into 3 stages (early, intermediate and late), each composed of 13

CS presentation and 13 ITIs for analysis of the behavioral and in vivo electrophysiology data. In this

case, the last CS and ITI was excluded from the analysis. Since no sex differences were observed

between male and female mice in both control and $\gamma 2^{f/f}$ groups (Supplementary Fig. 3a-h), data from

both sexes were combined for each group.

The startle response was assessed using a startle apparatus (SR-LAB Startle Response System,

v1.0.1.4, San Diego Instruments). The mice (P120) were exposed to tone stimuli of varying intensities

(measured in dB) lasting 20 ms, with an inter-trial interval (ITI) ranging from 10 to 20 s, and under a

constant background noise of 64 dB. Following a 5-minute acclimation period in this environment, the

tone stimuli were presented in random order to the animals. The startle response was quantified by

analyzing both the maximal and mean startle amplitudes in response to the randomly applied tone

stimuli.

The apparatus used for the elevated plus maze comprise two open and two perpendicular closed

arms (37x10 cm each one), elevated 50 cm above the floor (version 5.31.0.8, ViewPoint, Behavior

Technology, France). The closed arms had a plexiglass wall with a height of 15 cm. The mice (P100-

P125) were located at the center of the maze for 4 min for free exploration. We measured the

exploration time in the open arms as well as the distance traveled.

For the novel object recognition task, mice (P100-P125) were placed in the same Open Field

chamber containing two identical objects (Familiar objects), separated by 20 cm, and allowed to

explore for 5min. Following a 2-hour retention interval, the mice returned to the chamber for 5min-

exploration period, during which one of the familiar objects (always the left one) was replaced with a

novel object. Object exploration was quantified by measuring the time spent interacting with each

object, and a preference index (PI) was calculated as follows: PI = Time with Novel Object / (Time

with Novel Object + Time with Familiar Object).

Surgical implantation of the microdrive

A custom microdrive was fashioned using a 10 ml syringe cut to a length of 1.5 cm. This allows the Neuralynx EIB-16 headstage to be glued to the inside, serving as a base for a mini-Faraday cage. The cage was made by sticking a sheet of copper foil adhesive tape (noll electronic, Germany) around the syringe and grounding it with a silver-soldered wire (Ø 203.2 μm, ref. 782000, A-M systems). The conical bottom of the microdrive is made by cutting the bottom of a 10 ml Falcon® tube. This conical shape increased the surface in contact with the skull and eased the application of dental cement. The center of the conical base is drilled to glue a plastic guide for introducing the tetrode composed of 4 tungsten wires (Ø 50.8/101.6 μm) insulated with Teflon (ref. 795500, A-M systems). The tail of the tetrode along with the silver ground and reference wires were gently inserted into channels holes on the EIB-16-QC (Neuralynx), and pinned by small golden pins (Ref. 31-0603, Neuralynx). The fully assembled microdrive weighed less than 3g.

For chronic implantation of the microdrive, adult control and $\gamma 2^{f/f}$ mice (P100-P129) were deeply anaesthetized with isoflurane (4% for induction and 1.5% for maintenance, Vetflurane, Virbac). Local analgesics, including a solution of lidocaïne (Laocaïne, 7 mg/kg, Intervet-MSD), buprenorphine (BUPRECARE, 0,1 mg/kg; 2 mL/kg; Axience) and an ophtalmic gel (Ocrygel, DOMES PHARMA FR) were administered prior to or during the surgery. After fixing the animal in the stereotaxic frame (Kopf Instruments), the skin above the skull was disinfected with 70% ethanol and betadine solution (Vétédine, Vetoquinol, France) before incision. A hole was stereotaxically drilled to reach the IL region of mPFC using coordinates −0.4 mm mediolateral, +1.70 mm anterior-posterior relative to bregma. One additional hole in the interparietal region was performed to insert a screw overlying the cerebellum

where the reference wire and the soldered silver ground wire were connected together. Then, the screw

was sealed with a light-cure, cyanoacrylate-based adhesive (Loctite 4310) to ensure stability. Finally,

the microdrive was fixed to the skull using resin-modified glass ionomer cement (Ketac Cem Plus

Automix 3M) and acrylic dental cement (Pattern Resin LS, GC America Inc). The surface of the

exposed brain was covered with saline (0.9% NaCl) for the duration of the experiment. The skin was

sutured (Mersilene, EH7147H, Ethicon) and a saline solution at 37°C was administered subcutaneously

(100 to 200 µL) to rehydrate the animal. Finally, mice were recovered from anesthesia in an

environment at 37°C before they returned to their home cages.

At the end of experimental procedures, mice were deeply anesthetized using an overdose of a

solution of Ketamine (180 mg/kg, Clorketam 1000, Vetoquinol, France) and Xylazine (10 mg/kg,

Rompum, Elanco France), and electrolytic lesions were made to make the electrode implantation site

more visible. A current of 12 µA for 30 seconds was applied under gas anesthesia (isoflurane) through

the implanted tetrode *via* the EIB-16-QC interface and the NanoZ impedance measurement system

(Neuralynx, USA). Mice were then perfused with phosphate buffer saline (PBS) followed by 0.15 M

phosphate buffer (PB; pH ≈ 7.3), containing 4% paraformaldehyde (PFA). Brains were kept 2 h in

PFA, rinse three times in PB and incubated in sucrose 30% at 4°C until sinking for cryoprotection.

Then, brains were embedded with a film of OCT by gently rolling them (Microm Microtech), frozen in

dry ice and stored at -70°C. Coronal slices were cut with a cryostat (20 µm; Leica CM3050 S Cryostat),

stained with DAPI and mounted. Sections were visualized using an epifluorescence microscope (Zeiss

Axioplan 2) to confirm the recording location. Animals with the tetrode not clearly positioned in the

mPFC were excluded from the analysis.

In vivo electrophysiological recordings

Recordings were performed 7-15 days after surgery by using Neuralynx 16 channels preamplifier and

7m tether with its data acquisition software Cheetah (v6.3.2, Neuralynx, USA). LFPs were recorded at

30 kHz sampling rate (Neuralynx, USA). Animals were habituated to experimenter handling and to

recording sessions in their home cage for 15 min twice a day during 3 days; recording duration were 10

min in the open-field and 27 min on day-4 of the fear conditioning test.

The analysis of mPFC oscillations were carried out with custom-made programs written in

Python 3.9.7 (Spyder 5.1.5). All scripts used in this study are freely available at Angulo's lab gitlab

repository (https://gitlab.com/d5674/c_ibaceta/psd-volt). The mPFC LFP signals were band-pass

filtered from 1 Hz to 120 Hz (fifth-order IR Butterworth filter), 50 Hz notch filtered (Notch width 0.1

Hz) to minimize AC 50 Hz noise. For LFP analysis, the data was downsampled from 30 kHz to 1 kHz.

Then, we performed a power spectrum analysis (PSD) using the Welch method from SciPy Python and

Neurodsp library[54]. The parameters used were: 2048 FFTs, Hann window and 1024 overlapping points.

Following the PSD calculation, we normalized the data obtained during the open-field exploration by

averaging the values across all frequencies (from 1Hz to 120Hz). For fear extinction retrieval, we

normalized the data by averaging the values of frequencies ranging from 0Hz to 120Hz during the

baseline period (3 min). We defined three frequency bands, theta (4-12Hz), low gamma (30-45Hz) and

high gamma (55-90Hz), and obtained the PSD for each band range. For 3D plots depicting PSD over

time and frequency, we used Neurodsp library[54].

The PAC analysis was conducted using the Hilbert Transform, extracting the phase of Theta and

the amplitude of both low-gamma and high-gamma, with a FIT filter to isolate the three frequencies

(using SciPy Python library). For the analysis during tone fear extinction, we considered three stages:

baseline, sound and ITI. We further subdivided the sound and ITI in three phases based on trial

numbers, according to their trials: early (from 1 to 13), intermediate (from 14 to 26) and late (from 27

to 39). We calculated PSD and PAC for each of these phases.

In vivo fractalkine (FKN) infusions

In vivo FKN infusions were performed by adapting the protocol described by Watson et al. (2021)[34].
On the day of the surgery, FKN was dissolved in 0.2% BSA-PBS (75 µg/ml; Recombinant Soluble
Mouse CX3CL1/Fractalkine, R&D systems) or BSA in PBS (0.2%, Jackson ImmunoResearch). $\gamma 2^{f/f}$
mice (P42-P46) were deeply anesthetized with isoflurane (4% for induction and 1.5% for maintenance,
Vetflurane, Virbac). The surgical procedure and mouse recovery were similar to that described above.
For the local delivery of these compounds, a hole was stereotaxically drilled to implant a canula,
containing either FKN or BSA, into the right lateral ventricle using coordinates −1.000 mediolateral,
−0.300 anterior-posterior, −2.500 dorsoventral relative to bregma. Finally, the micro-osmotic pump
(model 1004, Alzet) was attached via a catheter to the cannula and placed in a subcutaneous pocket
between the shoulder blades. Infusions were administered at a flow rate of 0.11 µL per hour for 25-28
days. In total, each mouse received 200 ng of FKN per day, a dose that promoted myelination in adult
murine CNS[34,35]. At the end of the experimental procedure, mice were deeply anesthetized using an
overdose of a solution of Ketamine (180 mg/kg, Clorketam 1000, Vetoquinol, France) and Xylazine (10
mg/kg, Rompum, Elanco France) and then perfused for immunostaining analysis.

Viral injections and CNO protocol

Viral-mediated specific labeling of PV interneurons in the mPFC were performed by adapting the
protocol described by Stedehouder et al., 2018[30]. Adult $\gamma 2^{f/f}$ mice (P95-P105) were used for viral
injections, receiving either pAAV-S5E2-dTom-nlsdTom for controls (Addgene, #135630) or pAAV-
S5E2-Gq-P2A-dTomato (Addgene, #135635) for DREADD stimulation[31]. The surgical procedure and

mouse recovery were similar to that described above. Viral particles were injected into the mPFC

through a 10 µl gastight syringe (model 1701, Hamilton) using a motorized micropump (Legato 130,

KD Scientific) and silica needle (Ø 75/150 µm, PHYMEP), using coordinates −0.4 mm mediolateral,

+1.70 mm anterior-posterior, −2.9 mm dorsoventral relative to bregma. Four weeks after surgery, all

mice received intraperitoneal injections of CNO (Clozapine N-oxide (CNO) dihydrochloride, 1 mg/kg;

Hello Bio) twice daily (approximately at 9:00 h and 18:00 h) for 14 consecutive days. At the end of the

experimental procedure, mice were deeply anesthetized using an overdose of a solution of Ketamine

(180 mg/kg, Clorketam 1000, Vetoquinol, France) and Xylazine (10 mg/kg, Rompum, Elanco France)

and then perfused for immunostaining analysis.

Patch-clamp recordings on acute slices

Mice were euthanized by cervical dislocation. Patch-clamp experiments were performed in the IL

region of the mPFC using coronal slices of 300 µm-thick at P9-P13 or P59-P78 for OPCs in mPFC

(Supplementary Fig. 1), P92-P124 for layer 2/3 pyramidal neurons (Fig. 2a-b) in mPFC; P132-P146 for

layer 5 dTom$^+$ PV interneurons and layer 2/3 dTom$^-$ pyramidal neurons in mPFC (Supplementary Fig.

7), P126-P147 for principal neurons in basolateral amygdala (Fig. 2c-d) and CA1 pyramidal neurons in

the hippocampus (Fig. 2e-f). Recorded cells were visualized with an Orca Flash 4.0 LT CMOS digital

camera (Hamamatsu) and Micro-Manager software (version 1.4.21[55]) under an Olympus BX51

microscope equipped with a 40X fluorescent water-immersion objective. Excitation light to detect

fluorescent OPCs was provided by the CoolLed pE-300 ultra-system (Scientifica, UK) and 470 and

525□nm filters for excitation and emission wavelengths, respectively.

Patch-clamp recordings were performed at RT using an extracellular solution containing (in

mM): 126 NaCl, 2.5 KCl, 1.25 NaH2PO4, 26 NaHCO3, 20 glucose, 5 pyruvate, 3 CaCl2 and 1 MgCl2

(95% O2, 5% CO2). Glutamatergic and GABAergic postsynaptic currents of IL layer 2/3 pyramidal

neurons, BLA principal neurons with triangular somata, CA1 pyramidal neurons in the hippocampus

and fluorescent IL layer 5 OPCs were evoked at −70 mV and 0 mV, respectively by a monopolar

electrode (glass pipette; 100 ms pulse, 10 V; Iso-Stim 01D, npi electronic GmbH, Tamm, Germany),

with a CsMeS-based intracellular solution containing (in mM): 125 $CsCH_3SO_3H$, 5.4-aminopyridine,

10 tetraethylammonium chloride, 0.2 EGTA, 0.5 $CaCl_2$, 2 $MgCl_2$, 10 HEPES, 2 Na_2-ATP, 0.2 Na-GTP

and 10 Na_2-phosphocreatine (pH $\approx$ 7.3). The rate of stimulation was 0.2 Hz for neurons and 0.1 Hz

for OPCs. Recordings were made without series resistance (R_S) compensation. R_S was monitored

during recordings and cells showing a change of more than 30% were discarded. Potentials were

corrected for a junction potential of −10 mV. Acquisition was obtained using Multiclamp 700B and

pClamp version 10.7 software, filtered at 4 kHz and digitized at 20 kHz. Evoked postsynaptic

currents were analyzed off-line using pClamp version 10.7 software (Molecular Devices).

Immunostaining and quantification

Mice were deeply anesthetized using an overdose of a solution of Ketamine (180 mg/kg, Clorketam

1000, Vetoquinol, France) and Xylazine (10 mg/kg, Rompum, Elanco France). For PV and GAD-67

immunolabelling, mice (P115-P125) were then perfused with PBS followed by 0.15 M phosphate

buffer (PB; pH $\approx$ 7.3), containing 4% paraformaldehyde (PFA) and their brains were frozen in

isopentane at -70°C and cut into coronal slices with a cryostat (20 μm; Leica CM3050 S Cryostat). The

slices were then fixed in 4% paraformaldehyde (PFA) for 10 min with 0.1 M phosphate buffer (PB) at

4°C, washed first with PBS and then with tris buffered saline (TBS). Next, they were placed in a

blocking buffer with 50 mM TBS, 5% goat serum, 0.4% bovine serum albumin (BSA) and 0.1% Triton

x100 for 2h at RT, and incubated with primary antibodies in a 5% serum, 10% BSA, 10% Triton and 50

mM TBS solution for three nights in a cold chamber (4°C). Primary antibodies were rabbit polyclonal

anti-PV (1:500; ref. PV-27, Swant) and mouse monoclonal anti-GAD67 (1:200; ref. MAB5406, Sigma-Aldrich). Slices were washed three times in 50 mM TBS and 0.1% Triton x100 before they were incubated with secondary antibodies with 10% BSA, 10% Triton and 50 mM TBS at 4°C for two nights. We used goat anti-rabbit DyLight-405 (ref. 35551; Thermofisher Scientific) and goat anti-mouse Alexa Fluor-546 (ref. A-11030, Thermofisher Scientific) at 4°C for two nights at 1:200. Slices were washed three times with the same previous washing buffer and mounted with Vectashield mounting medium (ref. H-1000, Vectorlabs).

For other immunostainings, mice (age: P115-P125) were perfused and.brains were kept 2□h in PFA, rinsed three times in PB and incubated in sucrose 30% at 4°C until sinking for cryoprotection. Then, they were embedded with a film of OCT (Microm Microtech) by rolling them gently, frozen in dry ice and stored at -70°C. Coronal slices were cut with a cryostat (20 μm; Leica CM3050 S Cryostat), permeabilized with 0.3% triton X-100 and 4% Normal Goat Serum (NGS) overnight and incubated with primary antibodies. For internode length analysis, the slices were incubated with anti-PV and anti-SMI-312 for four nights before the incubation with the respective secondary antibodies during 2 hours at room temperature. Then, they were first incubated with anti-MBP for two nights and then with its secondary antibody for 2 hours at room temperature. For quantification of OPC and OL densities, the slices were incubated with anti-Olig2 and anti-CC1 for four nights and then with its secondary antibody for 2 hours at room temperature. To quantify recombination efficiency, we followed a similar protocol using an anti-Olig2 and anti-GFP antibody to amplify the GCaMP3 reporter signal (GCaMP3 served solely as a reporter in this analysis). Primary antibodies used for immunohistochemistry were rabbit polyclonal anti-PV (1:500; ref. PV-27, Swant), mouse anti-SMI-312 (1:200; ref. 837901, BioLegend), rat monoclonal anti-MBP (1:200; clone 26; ref. AB7349, Abcam), rabbit polyclonal anti-Olig2 (1:400; ref. AB9610, Merck Millipore), mouse monoclonal anti-APC (CC1; 1:100; clone CC-1; ref. OP80, Merck Millipore) and chicken polyclonal anti-GFP (for detection of GCaMP3 here used as a reporter;

1:1000; ref. ab13970, Abcam). Slices with primary antibodies were washed three times in PBS and

incubated in secondary antibodies coupled, respectively, to goat anti-rabbit DyLight-405 (ref. 35551,

Thermofisher Scientific), goat anti-mouse Alexa Fluor-546 (ref. A-11030, Thermofisher Scientific),

goat anti-rat Alexa Fluor-633 (ref. A-21094, Thermofisher Scientific) and goat anti-mouse DyLight-633

(ref. GTX76787; Genetex) at RT at 1:200.

For marker[+] cell analysis, confocal images were acquired using 63x oil objectives and, for

internode length and MBP analysis, confocal image stacks of 15-20 µm depth were acquired using HC

PL APO 93x/1,30 motCORR glycerol objectives with an SP8 Leica confocal microscope and the LAS

X software (version 3.5.7.23225). This objective is equipped with motorized correction for refractive

index mismatch which allows to image deeper in a sample, thereby maintaining optimal resolution and

signal intensity throughout the z-stack. With this configuration, we obtained a lateral resolution of ~200

nm, sufficient for precise measurement of internode lengths along PV+ axons in 3D. Images were then

analyzed blind using the Fiji/ImageJ2 framework (NIH, version 1.53i[56]). Cells were counted from layer

1 to 6 in the IL region of mPFC with ROI manager tool (3-4 slices per animal). Each image containing

20 z-sections of 1 µm each. Cells that were partially cut from the image were excluded in three of the

six sides of the cube to prevent border effects. PV[+], GAD-67[+], Olig2[+] and CC1[+] cell densities were

calculated as the number of cells per volume (mm^3). For the analysis of cell distributions, the x

coordinate of each cell was extracted from FIJI/ImageJ2 and its position was normalized with respect

to the total length of the layers of the mPFC, which can vary from one slice to another. Then, we

calculated the frequency of cells in the normalized mPFC length with a binning of 0.1. To quantify

MBP fluorescence coverage and intensity, we applied a mask generated from a binary (black & white)

image in FIJI. This mask was created to specifically isolate regions with MBP signal, allowing for

targeted analysis of fluorescence exclusively in these areas. To assess PV signal in axonal

compartments of deep layers of the IL region, we applied a mask generated from SMI-312[+] axon

labeling to isolate axonal regions in image stacks acquired with a 93x objective. Then, the intensity of

PV^+ pixels was quantified to estimate the PV signal in axonal compartments. Internode analyses were

systematically carried out in cortical layers V–VI of the IL region, where PV interneurons are most

abundant. For each animal, internodes were detected in 1–8 slices (mean: 2.95±0.21 slices per animal;

7.47±0.74 internodes per animal across 3-5 animals per group). Only internodes flanked by two nude

axonal regions were considered for analysis. Because such internodes are relatively sparse and difficult

to detect in thick brain sections, the sampled segments were naturally distributed across the tissue,

ensuring that quantifications reflected multiple independent axon populations rather than repeated

measures from the same axon. Internode length was measured from semi-automatic 3D reconstructions

of $SMI-312^+/PV^+/MBP^+$ segments obtained using Simple Neurite Tracer plugin in Fiji/ImageJ2

framework[18,57]. All quantifications were performed by an experimenter independent of the

immunostainings and image acquisition, and blinded to the experimental conditions, ensuring unbiased

analysis. Together, this strategy guarantees that the reported phenotype reflects consistent alterations

across multiple animals, slices, and axons, rather than a sampling artifact.

Statistics

All data are expressed as mean ± SEM. In the dot plots, individual mice (N) are represented by closed

dots while repeated measures (n) are shown as open dots. Statistical analysis were performed using

GraphPad Prism (v10.2.2; GraphPad Software Inc., USA), and JASP (version 0.16.4; JASP Team,

2022, Amsterdam, The Netherlands). Each group of data was first subjected to Shapiro-Wilk normality

test. According to the data structure and variance equality, two-group comparisons were performed

using a two-tailed Student's *t* test, a two-tailed Welch's t test or a two-tailed Mann-Whitney U test for

unpaired samples. We also used two-tailed (default output of JASP) linear mixed models with

likelihood ratio tests to account for repeated measures within each animal. Multiple group comparisons

were performed using the parametric two-way ANOVA followed by a post-hoc Sidak, Tukey's or

Fishers Least Significant Difference (LSD) multiple comparison tests, as well as mixed-effects model

followed by a Sidak's multiple comparison test for situations where data were not normality

distributed. Differences were considered significant when $P < 0.05$.

Data availability

The data generated and analyzed during this study are included in the manuscript and supplementary

data. Source data are provided with this paper. The raw datasets are available for research purposes

from the corresponding author upon request.

Code availability

Psd-volt Python scripts are hosted at https://gitlab.com/d5674/c_ibaceta/psd-volt and published under a

Free Software GNU GPLv3 license.

References

References

1. Khelfaoui, H., Ibaceta-Gonzalez, C. & Angulo, M. C. Functional myelin in cognition and
neurodevelopmental disorders. *Cell Mol Life Sci* **81**, 181 (2024).

2. Steadman, P. E. *et al.* Disruption of Oligodendrogenesis Impairs Memory Consolidation in Adult
Mice. *Neuron* **105**, 150-164.e6 (2020).

3. Pan, S., Mayoral, S. R., Choi, H. S., Chan, J. R. & Kheirbek, M. A. Preservation of a remote fear
memory requires new myelin formation. *Nat Neurosci* **23**, 487–499 (2020).

4. Shimizu, T. *et al.* Oligodendrocyte dynamics dictate cognitive performance outcomes of working
memory training in mice. *Nat Commun* **14**, 6499 (2023).

5. Kaller, M. S., Lazari, A., Blanco-Duque, C., Sampaio-Baptista, C. & Johansen-Berg, H. Myelin
plasticity and behaviour-connecting the dots. *Curr Opin Neurobiol* **47**, 86–92 (2017).

6. Pajevic, S., Plenz, D., Basser, P. J. & Fields, R. D. Oligodendrocyte-mediated myelin plasticity and
its role in neural synchronization. *Elife* **12**, e81982 (2023).

7. Moore, S. *et al.* A role of oligodendrocytes in information processing. *Nat Commun* **11**, 5497 (2020).

8. Dubey, M. *et al.* Myelination synchronizes cortical oscillations by consolidating parvalbumin-
mediated phasic inhibition. *Elife* **11**, e73827 (2022).

9. Kaller, M. S. *et al.* Ablation of oligodendrogenesis in adult mice alters brain microstructure and
activity independently of behavioral deficits. *Glia* **72**, 1728–1745 (2024).

10.   Jung, F. & Carlén, M. Neuronal oscillations and the mouse prefrontal cortex. *Int Rev Neurobiol*
158, 337–372 (2021).

11.   Bartos, M., Vida, I. & Jonas, P. Synaptic mechanisms of synchronized gamma oscillations in
inhibitory interneuron networks. *Nat Rev Neurosci* **8**, 45–56 (2007).

12.   Buzsáki, G. & Wang, X.-J. Mechanisms of gamma oscillations. *Annual review of neuroscience*
35, 203–225 (2012).

13.   Hijazi, S., Smit, A. B. & Van Kesteren, R. E. Fast-spiking parvalbumin-positive interneurons in

brain physiology and Alzheimer's disease. *Mol Psychiatry* (2023) doi:10.1038/s41380-023-02168-y.

14.   Benamer, N., Vidal, M. & Angulo, M. C. The cerebral cortex is a substrate of multiple

interactions between GABAergic interneurons and oligodendrocyte lineage cells. *Neurosci Lett* **715**,

134615 (2020).

15.   Vélez-Fort, M., Maldonado, P. P., Butt, A. M., Audinat, E. & Angulo, M. C. Postnatal Switch

from Synaptic to Extrasynaptic Transmission between Interneurons and NG2 Cells. *J Neurosci* **30**,

6921–6929 (2010).

16.   Orduz, D. *et al.* Interneurons and oligodendrocyte progenitors form a structured synaptic

network in the developing neocortex. *eLife* **4**, e06953 (2015).

17.   Orduz, D. *et al.* Developmental cell death regulates lineage-related interneuron-oligodendroglia

functional clusters and oligodendrocyte homeostasis. *Nat Commun* **10**, 4249 (2019).

18.   Benamer, N., Vidal, M., Balia, M. & Angulo, M. C. Myelination of parvalbumin interneurons

shapes the function of cortical sensory inhibitory circuits. *Nat Commun* **11**, 5151 (2020).

19.   Fang, L.-P. *et al.* Impaired bidirectional communication between interneurons and

oligodendrocyte precursor cells affects social cognitive behavior. *Nat Commun* **13**, 1394 (2022).

20.   Micheva, K. D. *et al.* A large fraction of neocortical myelin ensheathes axons of local inhibitory

neurons. *eLife* **5**, e15784 (2016).

21.   Stedehouder, J. *et al.* Fast-spiking Parvalbumin Interneurons are Frequently Myelinated in the

Cerebral Cortex of Mice and Humans. *Cereb Cortex* **27**, 5001–5013 (2017).

22.   Balia, M., Benamer, N. & Angulo, M. C. A specific GABAergic synapse onto oligodendrocyte

precursors does not regulate cortical oligodendrogenesis. *Glia* **65**, 1821–1832 (2017).

23.   Balia, M. *et al.* Postnatal Down-Regulation of the GABAA Receptor γ2 Subunit in Neocortical

NG2 Cells Accompanies Synaptic-to-Extrasynaptic Switch in the GABAergic Transmission Mode.

Cereb Cortex **25**, 1114–1123 (2015).

24.   Passlick, S. *et al.* Expression of the gamma2-subunit distinguishes synaptic and extrasynaptic

GABA(A) receptors in NG2 cells of the hippocampus. *J Neurosci* **33**, 12030–12040 (Jul 17).

25.   Maren, S., Phan, K. L. & Liberzon, I. The contextual brain: implications for fear conditioning,

extinction and psychopathology. *Nat Rev Neurosci* **14**, 417–428 (2013).

26.   Ji, L. *et al.* From hidden hearing loss to supranormal auditory processing by neurotrophin 3-

mediated modulation of inner hair cell synapse density. *PLoS Biol* **22**, e3002665 (2024).

27.   Ferguson, B. R. & Gao, W.-J. PV Interneurons: Critical Regulators of E/I Balance for Prefrontal

Cortex-Dependent Behavior and Psychiatric Disorders. *Front Neural Circuits* **12**, 37 (2018).

28.   Sung, Y. & Kaang, B.-K. The Three Musketeers in the Medial Prefrontal Cortex: Subregion-

specific Structural and Functional Plasticity Underlying Fear Memory Stages. *Exp Neurobiol* **31**,

221–231 (2022).

29.   Yang, S. M., Michel, K., Jokhi, V., Nedivi, E. & Arlotta, P. Neuron class-specific responses

govern adaptive myelin remodeling in the neocortex. *Science* **370**, eabd2109 (2020).

30.   Stedehouder, J., Brizee, D., Shpak, G. & Kushner, S. A. Activity-Dependent Myelination of

Parvalbumin Interneurons Mediated by Axonal Morphological Plasticity. *J Neurosci* **38**, 3631–3642

(2018).

31.   Vormstein-Schneider, D. *et al.* Viral manipulation of functionally distinct interneurons in mice,

non-human primates and humans. *Nat Neurosci* **23**, 1629–1636 (2020).

32.   Camacho-Hernández, N. P. & Peña-Ortega, F. Fractalkine/CX3CR1-Dependent Modulation of

Synaptic and Network Plasticity in Health and Disease. *Neural Plast* **2023**, 1–17 (2023).

33.   Voronova, A. *et al.* Migrating Interneurons Secrete Fractalkine to Promote Oligodendrocyte

Formation in the Developing Mammalian Brain. *Neuron* **94**, 500-516.e9 (2017).

34.   Watson, A. E. S. *et al.* Fractalkine signaling regulates oligodendroglial cell genesis from SVZ

precursor cells. *Stem Cell Reports* **16**, 1968–1984 (2021).

35.   De Almeida, M. M. A. *et al.* Fractalkine enhances oligodendrocyte regeneration and

remyelination in a demyelination mouse model. *Stem Cell Reports* **18**, 519–533 (2023).

36.   Griffiths, B. J. *et al.* Directional coupling of slow and fast hippocampal gamma with neocortical

alpha/beta oscillations in human episodic memory. *Proc Natl Acad Sci USA* **116**, 21834–21842

(2019).

37.   Tamura, M., Spellman, T. J., Rosen, A. M., Gogos, J. A. & Gordon, J. A. Hippocampal-

prefrontal theta-gamma coupling during performance of a spatial working memory task. *Nat*

Commun **8**, 2182 (2017).

38.   Makinodan, M., Rosen, K. M., Ito, S. & Corfas, G. A critical period for social experience-

dependent oligodendrocyte maturation and myelination. *Science* **337**, 1357–1360 (2012).

39.   Liu, J. *et al.* Impaired adult myelination in the prefrontal cortex of socially isolated mice. *Nat*

Neurosci **15**, 1621–1623 (2012).

40.   Teissier, A. *et al.* Early-life stress impairs postnatal oligodendrogenesis and adult emotional

behaviour through activity-dependent mechanisms. *Mol Psychiatry* **25**, 1159–1174 (2020).

41.   Bacmeister, C. M. *et al.* Motor learning drives dynamic patterns of intermittent myelination on

learning-activated axons. *Nat Neurosci* **25**, 1300–1313 (2022).

42.   Cullen, C. L. *et al.* Periaxonal and nodal plasticities modulate action potential conduction in the

adult mouse brain. *Cell Reports* **34**, 108641 (2021).

43.   Mercier, O. *et al.* Transient demyelination causes long□term cognitive impairment, myelin

alteration and network synchrony defects. *Glia* **72**, 960–981 (2024).

44.   Call, C. L. & Bergles, D. E. Cortical neurons exhibit diverse myelination patterns that scale

931   between mouse brain regions and regenerate after demyelination. *Nat Commun* **12**, 4767 (2021).

932   45.   Mederos, S. *et al.* GABAergic signaling to astrocytes in the prefrontal cortex sustains goal-

933   directed behaviors. *Nat Neurosci* **24**, 82–92 (2021).

934   46.   Cao, W. *et al.* Gamma Oscillation Dysfunction in mPFC Leads to Social Deficits in Neuroligin

935   3 R451C Knockin Mice. *Neuron* **97**, 1253-1260.e7 (2018).

936   47.   Maas, D. A. *et al.* Interneuron hypomyelination is associated with cognitive inflexibility in a rat

937   model of schizophrenia. *Nat Commun* **11**, 2329 (2020).

938   48.   Zoupi, L. *et al.* Selective vulnerability of inhibitory networks in multiple sclerosis. *Acta

939   Neuropathol* **141**, 415–429 (2021).

940   49.   Borges, B. C., Meng, X., Long, P., Kanold, P. O. & Corfas, G. Loss of oligodendrocyte ErbB

941   receptor signaling leads to hypomyelination, reduced density of parvalbumin☐expressing

942   interneurons, and inhibitory function in the auditory cortex. *Glia* **71**, 187–204 (2023).

943   50.   Stedehouder, J. & Kushner, S. A. Myelination of parvalbumin interneurons: a parsimonious

944   locus of pathophysiological convergence in schizophrenia. *Mol Psychiatry* **22**, 4–12 (2017).

945   51.   Yamada, J. *et al.* Altered regulation of oligodendrocytes associated with parvalbumin neurons in

946   the ventral hippocampus underlies fear generalization in male mice. *Neuropsychopharmacol* **48**,

947   1668–1679 (2023).

948   52.   Magno, L. *et al.* Transient developmental imbalance of cortical interneuron subtypes presages

949   long-term changes in behavior. *Cell Rep* **35**, 109249 (2021).

950   53.   Tanti, A. *et al.* Child abuse associates with increased recruitment of perineuronal nets in the

951   ventromedial prefrontal cortex: a possible implication of oligodendrocyte progenitor cells. *Mol

952   Psychiatry* **27**, 1552–1561 (2022).

54.   Cole, S., Donoghue, T., Gao, R. & Voytek, B. NeuroDSP: A package for neural digital signal

processing. *JOSS* **4**, 1272 (2019).

55.   D. Edelstein, A. *et al.* Advanced methods of microscope control using µManager software. *JBM*

1, 1 (2014).

56.   Schindelin, J. *et al.* Fiji: an open-source platform for biological-image analysis. *Nat Methods* **9**,

676–682 (2012).

57.   Schneider, C. A., Rasband, W. S. & Eliceiri, K. W. NIH Image to ImageJ: 25 years of image

analysis. *Nat Methods* **9**, 671–675 (2012).

Acknowledgements

We thank the NeurImag platform, Phenobrain and the animal facility of IPNP and their funding sources (Fédération pour la Recherche Médicale, Fondation Leducq). We would also thank Ludivine Therreau from Phenobrain as well as Beata Turanska and Julie Cognet in the team for their technical assistance, Steven A. Kushner and Karim Benchenane for insightful discussions on the project, Najate Benamer, Marie Vidal, Yana Kibalnyk, Philippe Faure and Elise Bousseyrol for help at the preliminary phase of the study. M.C.A. is a CNRS (Centre National de la Recherche Scientifique) investigator.

A.V and M.C.A. were supported by an Era-Net Neuron grant (CIHR and ANR under the frame of Neuron Cofund, No. 161466 and No. R19068KK for A.V. and M.C.A, respectively). A.E.S.W. was supported by CIHR CGSD. A.V. was supported by CIHR grants (PS 166120 and PJT 195940), MS Canada grant (1249541), the Scottish Rite Charitable Foundation of Canada grant as well as Canada Research Chair in Neural Stem Cell Biology and Sloan Research Fellowship in Neuroscience. M.C.A. was supported by grants from Fondation pour la Recherche Médicale (FRM, EQU202103012626 and MND202310017891), ANR (ANR CoLD: ANR-20-CE16–0001-01; ANR Myelex: ANR-21-CE37–0020-01), Fondation de France (00147199/WB-2023-50772) and Fédération pour la Recherche sur le Cerveau (1R24183PNOPES).

Author contributions

F.P. conducted the majority of the experiments and performed data analysis of the study. C.I-G. developed the Python code for in vivo electrophysiological analysis and contributed to behavioral tests and data analysis. H.K. performed patch-clamp recordings of OPCs and neurons in different brain regions as well as with virus injections. C.I-G and H.K. equally contributed to this work as second co-authors. C.P. carried out behavioral experiments and immunostainings. A.E.S.W. and A.V. performed

985   and assisted with FKN infusions. M.C.A. supervised the project, contributed to data analysis, and

986   drafted the initial version of the manuscript. All authors edited and approved the manuscript.

988   **Competing interests** All authors declare no competing interests.

 Figure Legends

Figure 1. Inactivation of GABAergic synapses in OPCs disrupts tone fear extinction

a. Mice were subjected to a four-day fear conditioning protocol consisting of four sessions: a

conditioning session which included a 180s baseline (BL) period followed by five 2s-foot shocks

delivered at the end of each 30s tone; a contextual extinction session (18 min); a tone extinction session

which included a 180s BL period followed by 40 tones of 30s each with 5s inter-trial interval (ITI); a

tone extinction retrieval session which included a 180s BL period followed by 40 tones of 30s each

with 5s ITI. **b-e.** Comparison of freezing behavior (percentage of time spent immobile) between control

(dark blue) and $\gamma 2^{f/f}$ (light blue) mice during each session. No major differences were observed between

the two groups during conditioning (**b**) and context extinction (**c**) sessions. Control mice: N=18 and

$\gamma 2^{f/f}$ mice: N=23; p= 0.9745 / 0.0748 / 0.0173 / 0.4112 / 0.1300 / 0.9752 for BL and trials 1-5 in (**b**); p=

0.9955 / 0.8627 / 0.7244 / 0.9989 / >0.9999 / >0.9999 / 0.9309 / >0.9999 / 0.9705 for 2-18 min in (**c**);

two-way ANOVA test followed by post-hoc Sidak multiple comparisons test. Data are represented as

mean±s.e.m. (**d,e**) During both tone extinction (**d**) and tone extinction retrieval (**e**) sessions, $\gamma 2^{f/f}$ mice

exhibited significantly higher freezing levels across most 5-trial blocks compared to controls. Note that

in control mice, the BL and the two first freezing levels during extinction retrieval shown in **e** were

significantly lower to BL and the two first freezing levels during tone extinction shown in **d** (p= 0.0162

for BL, p= 0.0014 for trial block 1 and p= 0.007 for trial block 2). Note also that BL freezing level was

higher in $\gamma 2^{f/f}$ mice compared to control during the extinction retrieval session (**e**). Control mice: N=18

and $\gamma 2^{f/f}$ mice: N=23; p= 0.9998 / 0.1588 / 0.0013 / 0.0109 / 0.0017 / 0.0034 / 0.0078 / <0.0001 /

0.0002 for PL and 5-trial blocks 1-8 in (**d**); p= 0.0358 / 0.0004 / <0.0001 / <0.0001 / <0.0001 / <0.0001

/ 0.0229 / 0.0086 / 0.0136 for BL and 5-trial blocks 1-8 in (**e**); two-way ANOVA test followed by post-

hoc Sidak multiple comparisons test. Data are represented as mean±s.e.m. **f.** Comparison of freezing

behavior between control and $\gamma 2^{f/f}$ mice at early, intermediate and late stages of the extinction retrieval

session (averaged freezing of 13 tones for each stage see Supplementary Fig. 3i). Control mice: N=18

and $\gamma 2^{f/f}$ mice: N=23; p<0.0001 / <0.0001/ p= 0.0018 at early-intermediate-late stages; two-way

ANOVA test followed by post-hoc Sidak multiple comparisons test. Data are represented as

mean±s.e.m. *p<0.05, **p<0.01, ***p<0.001, ****p<0.0001. Source data are provided as a Source

Data file.

1019

Figure 2. Early inactivation of GABAergic synapses of OPCs impairs inhibition in the adult

mPFC

a. eIPSCs (top trace) and eEPSCs (bottom trace) in layer 2/3 pyramidal neurons held at 0 mV and −70

mV, respectively, upon extracellular stimulation of layer 5 of the infralimbic (IL) region of acute slices

of the mPFC (*inset*, left) in 3-month-old control (dark blue) and $\gamma 2^{f/f}$ (light blue) mice. Stimulation

artifacts blanked, stimulation time indicated (black arrowheads). **b.** Dot plots of eIPSCs (left), eEPSCs

(middle) and E/I ratio (right) calculated from eEPSCs and eIPSCs of layer 2/3 pyramidal neurons

evoked by extracellular stimulation in layer 5 in control (dark blue) and $\gamma 2^{f/f}$ mice (light blue) (n=16

cells from N=9 mice and n=15 cells from N=7 mice for control and $\gamma 2^{f/f}$ mice, respectively). p= 0.0494

for eIPSCs, p= 0.3378 for eEPSCs, p= 0.0002 for E/I ratio; two-tailed Mann–Whitney U test. Data are

represented as mean±s.e.m. *p<0.05, ***p0.001, ns: not significant. **c.** eIPSCs (top trace) and eEPSCs

(bottom trace) in principal neurons of the basolateral amygdala (BLA) held at 0 mV and −70 mV,

respectively, upon extracellular stimulation of the lateral amygdala (LA) in 4-month-old control (dark

blue) and $\gamma 2^{f/f}$ (light blue) mice. Stimulation artifacts blanked, stimulation time indicated (black

arrowheads). **d.** Dot plots of eIPSCs (left) and eEPSCs (middle) and E/I ratio (right) calculated from

eEPSCs and eIPSCs of principal neurons evoked by extracellular stimulation in the LA in control (dark

blue) and $\gamma 2^{f/f}$ mice (light blue) (n=9 cells from N=5 mice and n=8 cells from N=3 mice for control and

$\gamma2^{f/f}$ mice, respectively). p= 0.9082 for eIPSCs, p= 0.8365 for eEPSCs, p= 0.4968 for E/I ratio; two-

tailed Student *t* test. Data are represented as mean±s.e.m; ns: not significant. **e.** eIPSCs (top trace) and

eEPSCs (bottom trace) in CA1 pyramidal neurons of the hippocampus held at 0 mV and −70 mV,

respectively, upon extracellular stimulation of Schaffer collaterals in 4-month-old control (dark blue)

and $\gamma2^{f/f}$ (light blue) mice. Stimulation artifacts blanked, stimulation time indicated (black arrowheads).

f. Dot plots of eIPSCs (left) and eEPSCs (middle) and E/I ratio (right) calculated from eEPSCs and

eIPSCs of pyramidal neurons evoked by extracellular stimulation in stratum radiatum in control (dark

blue) and $\gamma2^{f/f}$ mice (light blue) (n=8 cells from 5 mice and n=7 cells from 4 mice for control and $\gamma2^{f/f}$

mice, respectively). p= 0.4531 for eIPSCs, p= 0.0721 for eEPSCs, p= 0.2382 for E/I ratio; two-tailed

Student *t* test for eIPSCs and E/I ratio; two-tailed Mann–Whitney U test for eEPSCs. Data are

represented as mean±s.e.m. ns: not significant. Source data are provided as a Source Data file.

1048

Figure 3. Inactivation of GABAergic synapses of OPCs impairs PV interneuron internode length

in the mPFC

a. Confocal images of MBP in the IL region of the mPFC of control (left) and $\gamma2^{f/f}$ (right) mice. Scale

bar: 50 µm. **b, c.** Dot plots of MBP fluorescence coverage (**b**) and intensity (**c**) of control (dark blue)

and $\gamma2^{f/f}$ (light blue) mice. Control mice: n=12 technical replicates from N=3 mice (biological

replicates) and $\gamma2^{f/f}$ mice: n=16 technical replicates from N=3 mice (biological replicates). p= 0.078 for

(**b**) and p= 0.374 for (**c**); two-tailed linear mixed models with likelihood ratio tests applied to account

for repeated measures (open dots) within each animal (closed dots). Data are presented as mean±s.e.m.

ns: not significant. **d.** Confocal images of PV (red), MBP (green), and SMI-312 (blue) labeling,

illustrating myelinated internodes of PV[+] axons, identified by bare axonal regions on each side, in

control and $\gamma2^{f/f}$ mice (white arrowheads). Note that the internode of PV[+] axons in $\gamma2^{f/f}$ mice appeared

longer compared to those of control mice. Scale bar: 10 µm. **e, f.** Dot plots of internode lengths of PV[+]

(**e**) and PV$^-$ (**f**) axons in control (dark blue) and $\gamma 2^{f/f}$ (light blue) mice. Control mice: n= 21 technical

replicates from N=3 mice (biological replicates) and $\gamma 2^{f/f}$ mice: n= 21 technical replicates from N= 3

mice (biological replicates) for PV$^+$ axons; Control mice: n= 16 technical replicates from N= 3 mice

(biological replicates) and $\gamma 2^{f/f}$ mice: n= 15 technical replicates from N= 3 mice (biological replicates)

for PV$^-$ axons. p= 0.026 for (**e**) and p= 0.325 for (**f**); two-tailed linear mixed models with likelihood

ratio tests applied to account for repeated measures (open dots) within each animal (closed dots). Data

are represented as mean±s.e.m. *p<0.05, ns: not significant. Source data are provided as a Source Data

file.

1069

1070

Figure 4. Chemogenetic-induced increase in PV interneuron activity does not rescue fear

extinction deficits

a. Schematic of viral injections and clozapine-N-oxide (CNO) administration protocol for PV

interneuron activation in the mPFC. The $\gamma 2^{f/f}$ mice received bilateral injections of either pAAV-S5E2-

dTom-nlsd (dTom) or pAAV-S5E2-hM3Dq(Gq)-P2A-dTomato (hM3Dq-dTom) virus targeting PV

interneurons in the mPFC at around P90. After one month, the same mice received twice-daily

intraperitoneal injections of CNO for 14 consecutive days. The day following the final CNO injection,

they were subjected to the fear conditioning protocol (d1–d4). **b.** Confocal image of a coronal section

showing the restricted expression of hM3Dq-dTom virus on the IL region of the mPFC. Scale bar: 500

μm. **c.** Confocal images of dTom$^+$/PV$^+$ neurons in the IL region. A dTom$^+$/PV$^+$ cell is shown (Inset,

arrow); a dTom$^+$/PV$^-$ cells is indicated (arrowhead). Dot plots (right) of the percentage of dTom$^+$/PV$^+$

cells in both dTom-injected (white) and hM3Dq-dTom-injected (red) $\gamma 2^{f/f}$ mice. dTom-injected $\gamma 2^{f/f}$

mice: n= 9 technical replicates from N= 3 mice (biological replicates) and hM3Dq-dTom-injected $\gamma 2^{f/f}$

mice: n= 9 technical replicates from N= 3 mice (biological replicates). p= 0.92 for (**c**); two-tailed linear

mixed models with likelihood ratio tests applied to account for repeated measures (open dots) within

each animal (closed dots). Data are represented as mean±s.e.m. ns: not significant. Scale bar: 50 µm

and 10 µm. **d.** Confocal images of MBP in the IL region of the mPFC of dTom-injected (left) and

hM3Dq-dTom-injected (right) $\gamma 2^{f/f}$ mice. Scale bar: 50 µm. **e, f.** Dot plots of MBP fluorescence

coverage (**e**) and intensity (**f**) of the IL region of dTom-injected (white) and hM3Dq-dTom-injected

(red) $\gamma 2^{f/f}$ mice (dTom-injected $\gamma 2^{f/f}$ mice: n= 11 technical replicates from N= 3 mice (biological

replicates) and hM3Dq-dTom-injected $\gamma 2^{f/f}$ mice: n= 10 technical replicates from N= 3 mice (biological

replicates)). p= 0.486 for (**e**), p=0.158 for (**f**); two-tailed linear mixed models with likelihood ratio tests

applied to account for repeated measures (open dots) within each animal (closed dots). Data are

presented as mean±s.e.m. ns: not significant. **g.** Dot plots of internode lengths of PV$^+$ axons in dTom-

injected (white) and hM3Dq-dTom-injected (red) $\gamma 2^{f/f}$ mice (dTom-injected $\gamma 2^{f/f}$ mice: n= 19 technical

replicates from N= 3 mice (biological replicates) and hM3Dq-dTom-injected $\gamma 2^{f/f}$ mice: n= 26

technical replicates from N= 5 mice (biological replicates); p= 0.181 for (**g**); two-tailed linear mixed

models with likelihood ratio tests applied to account for repeated measures (open dots) within each

animal (closed dots). Data are represented as mean±s.e.m. ns: not significant. **h, i.** Comparison of

freezing behavior between dTom-injected (white) and hM3Dq-dTom-injected (red) $\gamma 2^{f/f}$ mice during

tone extinction (**h**) and tone extinction retrieval (**i**) sessions. No significant differences were observed

between the two groups (dTom-injected $\gamma 2^{f/f}$ mice: N=13 and hM3Dq-dTom-injected $\gamma 2^{f/f}$ mice: N=14).

p= 0.7225 / 0.5369 / 0.7494 / 0.2596 / 0.2849 / 0.8932 / >0.9999 / 0.8701 / 0.9983 for BL and 5-trial

blocks 1-8 in (**h**) and p= 0.5947 / 0.6679 / 0.4744 / 0.8512 / 0.9922 / 0.9999 / 0.9819 / 0.9242 / 0.9711

for BL and 5-trial blocks 1-8 in (**i**); two-way ANOVA test followed by post-hoc Sidak multiple

comparisons test. Data are represented as mean±s.e.m. ns: not significant. 4-OHT: 4-

hydroxytamoxifen. Source data are provided as a Source Data file.

1108

1109

Figure 5. In vivo FKN infusions increase myelination, but does not rescue fear extinction deficits

a. Schematic of the in vivo infusion protocol. Vehicle (VC) or FKN was continuously infused into the lateral ventricle of $\gamma2^{f/f}$ mice at P42-P46 for 25-28 days (D25-D28) using a micro-osmotic pump. These mice were subjected to the fear conditioning protocol (d1-d4) during the last days of infusion. **b.** Confocal images of MBP in the IL region of the mPFC of VC-infused (left) and FKN-infused (right) $\gamma2^{f/f}$ mice at P70-74. Note the increased myelination of FKN-infused $\gamma2^{f/f}$ mouse. Scale bar: 50 µm. **c, d.** Dot plots of MBP fluorescence coverage (**c**) and intensity (**d**) of the IL region of VC-infused (white) and FKN-infused (yellow) $\gamma2^{f/f}$ mice. VC-infused mice: n= 21 technical replicates from N= 4 mice (biological replicates) and FKN-infused mice: n= 23 technical replicates from N= 3 mice (biological replicates); p= 0.021 for (**c**) and p= 0.016 for (**d**); two-tailed linear mixed models with likelihood ratio tests applied to account for repeated measures (open dots) within each animal (closed dots). Data are presented as mean±s.e.m. *p<0.05 **e.** Dot plots of internode lengths of PV$^+$ axons in VC-infused (white) and FKN-infused (yellow) $\gamma2^{f/f}$ mice. VC-infused mice: n= 56 technical replicates from N= 4 mice (biological replicates) and FKN-infused mice: n= 31 technical replicates from N=3 mice (biological replicates); p= 0.722 for (**e**); two-tailed linear mixed models with likelihood ratio tests applied to account for repeated measures (open dots) within each animal (closed dots). Data are represented as mean±s.e.m. ns: not significant. **f, g.** Comparison of freezing behavior between VC-infused (white) and FKN-infused (yellow) $\gamma2^{f/f}$ mice during tone extinction (**f**) and tone extinction retrieval (**g**) sessions. No significant differences were observed between the two groups. VC-infused mice: N=8 and FKN-infused mice: N=8; p= 0.9992 / 0.5402 / 0.5641 / 0.2220 / 0.8762 / >0.9999 / 0.9109 / 0.9995 / 0.4330 for BL and 5-trial blocks 1-8 in (**f**); p= 0.9996 / 0.7776 / 0.9995 / 0.9590 / >0.9999 / 0.8786 / >0.9999 / 0.9845 / 0.9933 for BL and 5-trial blocks 1-8 in (**g**); two-way ANOVA test

1132 followed by post-hoc Sidak multiple comparisons test. Data are represented as mean±s.e.m. Source

1133 data are provided as a Source Data file.

1134

Figure 6. Disruption of gamma oscillations during fear extinction retrieval, but not exploration

a. Schematic of the protocol for conducting in vivo recordings in behaving mice. Surgery to implant the

microdrive containing the tetrode was performed at around P115. Recordings of LFPs were performed

7–15 days-post-surgery during open-field exploration and fear extinction retrieval. **b.** The tetrode was

implanted in the IL region of the mPFC (left). Brain slice of a recorded mouse (middle) after a DAPI

staining (blue) and showing the correct location of the tetrode (right, arrowhead). Scale bar, 1 mm. fmi,

forceps minor of the corpus callosum; CPu, Caudate-Putamen; PL, prelimbic cortex; IL, infralimbic

cortex. **c.** Representative raw LFP signal from the mPFC (top), used to extract the theta (4-12 Hz,

middle) and gamma (30-90 Hz, bottom) frequency components following band-pass filtering. **d.** Color-

coded spectrogram from a mouse LFP recording, showing theta (4-12 Hz) and gamma (30-90 Hz)

oscillation powers during fear extinction retrieval. **e, f, i, j.** Normalized (Norm.) power spectra of

mPFC recordings from the center (C) and periphery (P) of the open-field (OF) arena (**e,f**), as well as

during tone and inter-trial interval (ITI) phases of the fear extinction retrieval session (**i,j**) in control

(dark blue) and $\gamma2^{f/f}$ (light blue) mice. Theta (θ) and gamma (γ) frequency bands are indicated. Power

spectra are averaged of all animals, and the curves and the shaded areas indicate mean±s.e.m. **g, h, k, l.**

Dot plots showing the normalized (Norm.) theta (**g, k**) and gamma (**h, l**) oscillation powers for the

center (C) and periphery (P) of the open-field arena (**g, h**), and for tones and inter-trial intervals (ITI)

during the fear extinction retrieval session (**k, l**) in control (dark blue) and $\gamma2^{f/f}$ (light blue) mice.

Control mice: N=5 and $\gamma2^{f/f}$ mice: N=5; p= 0.8797 in center and p= 0.8966 in periphery for (**g**); p=

0.4152 in center and p= 0.4343 in periphery for (**h**); p= 0.2222 during tones (two-tailed Mann–Whitney

U test) and p= 0.1222 during ITIs for (**k**); p= 0.03 during tones and p= 0.017 during ITIs for (**l**); two-

tailed Student's *t* test. All data are represented as mean±s.e.m. *p<0.05, ns: not significant. 4-OHT: 4-

hydroxytamoxifen. Source data are provided as a Source Data file.

Figure 7. Disrupted low-Gamma oscillations during fear extinction retrieval in $\gamma2^{f/f}$ mice

a, b. Dot plots of the relative power of low-gamma and high-gamma oscillations during tone (**a**) and

inter-trial interval (ITI) (**b**) in control (dark blue) and $\gamma2^{f/f}$ (light blue) mice. Control mice: N=5 and $\gamma2^{f/f}$

mice: N=5; p=0.0033 (low-gamma) and p= 0.0978 (high-gamma) for (**a**); p=0.0013 (low-gamma) and

p= 0.0629 (high-gamma) for (**b**); two-way ANOVA test followed by post-hoc Sidak multiple

comparisons test. **p<0.01, ns: not significant **c, d.** Comparisons of the normalized (Norm.) power of

low-gamma (**c**) and high-gamma (**d**) oscillations during tone (left) or inter-trial interval (ITI; right) in

control and $\gamma2^{f/f}$ mice at early (E), intermediate (I) and late stages (L) of the extinction retrieval session

(average responses during 13 tones for each stage, see Supplementary Fig. 2i). Individual animal

responses are shown by faint traces. Control mice: N=5 and $\gamma2^{f/f}$ mice: N=5; p= 0.026 / 0.0237 / 0.0860

at early, intermediate andlate stages during tones and p= 0.0109 / 0.0061 / 0.1398 at early, intermediate

and late stages during ITIs for low-gamma (**c**); p= 0.2781 / 0.1078 / 0.1470 at early, intermediate and

late stages during tones and p= 0.2159 / 0.1005 / 0.2578 at early, intermediate and late stages during

ITIs for high-gamma (**d**); two-way ANOVA test followed by a post-hoc Sidak multiple comparisons

test. All data are represented as mean±s.e.m. *p<0.05, **p<0.01, ns: not significant. Source data are

provided as a Source Data file.

a

Conditioning — context **A**

→ 24 h →

Context extinction — context **A**

→ 24 h →

Tone extinction — context **B**   X7

→ 24 h →

Extinction retrieval — context **B**   X7

b Conditioning

○ Ctrl   ● γ2^{f/f}

Freezing (%)

BL 1 2 3 4 5
Trials

c Context extinction

○ Ctrl   ● γ2^{f/f}

Freezing (%)

2 4 6 8 10 12 14 16 18
Time (minutes)

d Tone extinction

○ Ctrl   ● γ2^{f/f}

Freezing (%)

BL 1 2 3 4 5 6 7 8
5-trial blocks

e Extinction retrieval

○ Ctrl   ● γ2^{f/f}

Freezing (%)

BL 1 2 3 4 5 6 7 8
5-trial blocks

f Extinction retrieval

○ Ctrl   ● γ2^{f/f}

Freezing (%)

early   intermediate   late

a

pAAV-S5E2-dTom-nlsdTom
or
pAAV-S5E2-hM3Dq(Gq)-P2A-dTomato

$\gamma 2^{f/f}$

$\gamma 2^{f/f}$

Age   P3 P4 P5   P90   D0   D28   14 days   D42 d1 d2 d3 d4

4-OHT
injection

D0
virus bilateral
injection

CNO injections
(I.P. twice daily)

fear
conditioning

b

hM3Dq-dTom

c

dTom
PV

dTom⁺ PV⁺ cells / dTom⁺ cells (%)

ns

dTom   hM3Dq
-dTom

d

IL

dTom   hM3Dq-dTom

MBP   MBP

e

MBP coverage (µm²)

ns

dTom   hM3Dq
-dTom

f

MBP intensity (a.u.)

ns

dTom   hM3Dq
-dTom

g

Internode length of PV⁺ axons (µm)

ns

dTom   hM3Dq
-dTom

h

Tone extinction

dTom   hM3Dq-dTom

Freezing (%)

BL 1 2 3 4 5 6 7 8
5-trial blocks

i

Extinction retrieval

dTom   hM3Dq-dTom

Freezing (%)

BL 1 2 3 4 5 6 7 8
5-trial blocks

a

Age P3 P4 P5   P42-46   d1 d2 d3 d4

↑ ↑ ↑
4-OHT
injection

D0
pump
implantation

VC or FKN infusion

D25-28
fear conditioning

b

IL

MBP   VC   FKN   MBP

c MBP coverage (μm^2) — ******** — VC, FKN

d MBP intensity (a.u.) — ******* — VC, FKN

e Internode length of PV+ axons (μm) — ns — VC, FKN

f Tone extinction — VC, FKN — Freezing (%) — BL 1 2 3 4 5 6 7 8 — 5-trial blocks

g Extinction retrieval — VC, FKN — Freezing (%) — BL 1 2 3 4 5 6 7 8 — 5-trial blocks

Plaisier et al.

Supplementary Figure 1. Elimination of γ2-GABA$_A$ receptor-mediated postsynaptic currents in OPCs of the mPFC in pups

a. Experimental protocol. Both control and γ2$^{f/f}$ mice were injected with 4-OHT from P3 to P5 followed by patch-clamp recordings of OPCs in acute mPFC slices at around P10 (P9-P13), coinciding with the peak of cortical GABAergic synaptic connectivity[15], or at around P60 (P59-P78). **b.** Representative evoked glutamatergic (I$_{Glu}$) and GABAergic (I$_{GABA}$) postsynaptic currents in OPCs of control (left) and γ2$^{f/f}$ (right) mice in response to extracellular stimulation

of layer 5 of the IL region of the mPFC at P9-P13. Note the near-complete abolishment of I_{GABA} in $\gamma 2^{f/f}$ mice. Stimulation artifacts blanked, stimulation time indicated (black arrowheads). **c.** Dot plots showing the amplitudes of I_{Glu} (left) and I_{GABA} (right) in control (dark blue; n=8 cells from N=3 mice) and $\gamma 2^{f/f}$ (light blue; n=7 cells from N=3) mice. p= 0.6107 for I_{Glu}; p= 0.0030 for I_{GABA}; two-tailed Student's t test for excitation and two-tailed Welch's t test for inhibition. Data are represented as mean±s.e.m. **p<0.01, ns: not significant. **d.** Representative evoked GABAergic (I_{GABA}) postsynaptic currents in OPCs of control (left) and $\gamma 2^{f/f}$ (right) mice in response to extracellular stimulation of layer 5 of the IL region of the mPFC at P59-P78. Recordings were obtained in normal conditions (blue traces) and in the presence of 1 µM zolpidem (ZPD; gray traces). Stimulation artifacts blanked, stimulation time indicated (black arrowheads). **e**. Dot plots showing the amplitudes of I_{GABA} before and after 4 min of bath application of ZPD in control (dark blue: n=8 cells in total and n=6 in ZPD from N=3 mice) and $\gamma 2^{f/f}$ mice (light blue: n=7 cells in total and n=4 in ZPD from N=3 mice) mice. Note that, unlike during development, evoked I_{GABA} were similar in control and $\gamma 2^{f/f}$ mice at P59-P78 and were not significantly potentiated by zolpidem (1 µM), indicating the absence of functional $\gamma 2$-containing GABA-A receptors in adult OPCs. p= 0.1996 for comparison of I_{GABA} in normal conditions, p>0.9999 for comparison of ZPD potentiation in control mice; p= 0.6250 for comparison of ZPD potentiation in $\gamma 2^{f/f}$ mice; two-tailed Mann-Whitney U test to compare I_{GABA} in control and $\gamma 2^{f/f}$ mice; Wilcoxon test for ZPD. Data are represented as mean±s.e.m. ns: not significant. 4-OHT: 4-hydroxytamoxifen. Source data are provided as a Source Data file.

Supplementary Figure 2. Recombination efficiency and no changes in the density and proportions of Olig2$^+$/CC1$^-$ OPCs and Olig2$^+$/CC1$^+$ OL of the mPFC in γ2$^{f/f}$ mice

a. Confocal image (middle) of Olig2$^+$ (white) and GFP$^+$ (green) cells in IL region of the adult mPFC (left). Note that most Olig2$^+$ cells are positive for GFP. Some Olig2$^+$/GFP$^-$ cells are

indicated (yellow arrowheads). Scale bar: 50 µm. Dot plots (right) of the recombination efficiency for control (dark blue) and $\gamma2^{f/f}$ (light blue) mice. Control mice: n= 12 technical replicates from N= 3 mice (biological replicates) and $\gamma2^{f/f}$ mice: n= 12 technical replicates from N= 3 mice (biological replicates); p= 0.598 for (**a**); two-tailed linear mixed models with likelihood ratio tests applied to account for repeated measures (open dots) within each animal (closed dots). Data are represented as mean±s.e.m. ns: not significant**. b.** Representative confocal images of Olig2$^+$ (white) and CC1$^+$ (magenta) cells in IL region of the mPFC (inset) in four-month control (top) and $\gamma2^{f/f}$ (bottom) mice. Arrowheads indicate Olig2$^+$/CC1$^+$ OL cells (white) and Olig2$^+$/CC1$^-$ OPCs (yellow). Scale bar: 50 µm. **c.** Quantification of Olig2$^+$ and Olig2$^+$/CC1$^+$ cell densities in control (dark blue) and $\gamma2^{f/f}$ mice (light blue). Control mice: n= 12 technical replicates from N= 3 mice (biological replicates) and $\gamma2^{f/f}$ mice: n= 12 technical replicates from N= 3 mice (biological replicates). p= 0.615 for Olig2$^+$ and p= 0.915 for Olig2$^+$/CC1$^+$ quantifications; two-tailed linear mixed models with likelihood ratio tests applied to account for repeated measures (open dots) within each animal (closed dots). Data are represented as mean±s.e.m. ns: not significant. **d.** Ratio of Olig2$^+$/CC1$^-$ OPCs and Olig2$^+$/CC1$^+$ OL cell densities compared to the total number of Olig2$^+$ cells in control (dark blue) and $\gamma2^{f/f}$ mice (light blue). Control mice: n= 12 technical replicates from N= 3 mice (biological replicates) and $\gamma2^{f/f}$ mice: n= 12 technical replicates from N= 3 mice (biological replicates). p= 0.624 for OPC and p= 0.624 for OL cells quantifications; two-tailed linear mixed models with likelihood ratio tests applied to account for repeated measures (open dots) within each animal (closed dots). Data are represented as mean±s.e.m. ns: not significant. Source data are provided as a Source Data file.

Supplementary Figure 3. No sex-dependent behavioral differences observed in control and γ2^{f/f} mice

a-h. Mice were subjected to a four-day fear conditioning protocol consisting of four sessions (see Fig. 1 for the protocol). Comparison of freezing behavior between males (green) and females (orange) in control (**a-d**) and γ2^{f/f} (**e-h**) mice during each session. No significant sex-related differences were observed in either group (control mice: N=9 males and N=9 males females; γ2^{f/f} mice: N=8 males and N=15 females). p= 0.9951 / >0.9999 / >0.9999 / 0.9311 / 0.7790 / 0.9555 for BL and trials 1-5 in (**a**); p= >0.9999 / 0.7735 / 0.8212 / >0.9999 / 0.9519 / >0.9999 / >0.9999 / 0.9876 / >0.9999 at 2-18 min in (**b**); p= 0.9968 / 0.3125 / 0.9905 / 0.8630 / 0.6554 / 0.7887 / 0.9998 / 0.9728 / >0.9999 for BL and 5-trial blocks 1-8 in (**c**); p= 0.9310 / 0.9032 / 0.8360 / 0.9993 / >0.9999 / 0.9680 / 0.7230 / 0.7247 / >0.9999 for BL and 5-trial blocks 1-8 in (**d**); p= >0.9999 / >0.9999 / 0.8630 / >0.9999 / >0.9999 / 0.9965 for BL and trials 1-5 in (**e**); p= 0.4722 / 0.4544 / 0.6724 / 0.9971 / >0.9999 / 0.9966 / >0.9999 / 0.9920 / >0.9999 for 2-18 min in (**f**); p= 0.9958 / >0.9999 / >0.9999 / >0.9999 / 0.9753 / 0.9993 / >0.9999 / 0.3224

/ >0.9999 for BL and 5-trial blocks 1-8 in (**g**); p= >0.9999 / >0.9999 / >0.9999 / >0.9999 / 0.9748 / >0.9999 / 0.8924 / 0.7174 / 0.9996 at BL/5-trial blocks 1-8 in (**h**); two-way ANOVA test followed by post-hoc Sidak multiple comparisons test. Data are represented as mean±s.e.m. **i.** Schematic of the early, intermediate and late stages (13 tones for each stage) of the tone extinction retrieval session. Source data are provided as a Source Data file.

Supplementary Figure 4. No auditory impairments detected in γ2^{f/f} mice

a,b. A mouse in the startle apparatus (**a**) was exposed to tone stimuli of varying intensities (**b**, measured in dB) and lasting 20 ms, with an inter-trial interval of 10 to 20 seconds. Following 5 min of acclimation period under a constant background noise of 64 dB, the stimuli were presented in random order. **c,d.** Startle response to the randomly applied tone stimuli of varying intensities. No significant differences were observed for either the maximal startle amplitude (**c**) or the mean startle amplitude (**d**). N=8 control mice and N=8 γ2^{f/f} mice. p= >0.9999 / >0.9999 / >0.9999 / >0.9999 / 0.9981 / >0.9999 / 0.9785 for NST, 70, 80, 85, 90, 100, 120 dB in (**c**); p= >0.9999 / >0.9999 / > 0.9999 / >0.9999 / 0.9993 / >0.9999 / 0.9992 for NST, 70, 80, 85, 90, 100, 120 dB in (**d**); mixed-effects model followed by a Sidak's multiple comparison test. Data are represented as mean±s.e.m. NST: No-stimulus trial. Source data are provided as a Source Data file.

Supplementary Figure 5. Comparable behavioral performance in γ2$^{f/f}$ mice following adult recombination or during elevated plus maze and novel object recognition following early postnatal recombination.

a-e. Mice received tamoxifen (TMX) injections for three consecutive days two weeks prior to be subjected to the same fear condition protocol of Fig. 1a. Comparisons of freezing behavior are shown between γ2$^{f/f}$ mice that were not injected (black) and those that received injections in adulthood (gray) for each session. No significant differences were observed between the two groups. N=15 and N=12 for not injected and injected in adulthood γ2$^{f/f}$ mice, respectively; p= 0.3115 / 0.7434 / 0.2140 / 0.4495 / 0.3692 / 0.9955 for BL and trials 1-5 in (**b**); p= 0.8014 / 0.5073 / 0.7394 / >0.9999 / 0.9991 / >0.9999 / 0.8445 / 0.5385 / 0.1911 for 2-18 min in (**c**); p= 0.5495 / >0.9999 / 0.9432 / 0.9998 / 0.9908 / 0.8998 / 0.9443 / 0.6961 / 0.9849 for BL and 5-trial blocks 1-8 in (**d**); p= 0.9978 / 0.8794 / 0.9716 / 0.9846 / 0.9463 / 0.8884 / >0.9999 / >0.9999 / 0.9997 for BL and 5-trial blocks 1-8 in (**e**); two-way ANOVA test followed by post-hoc Sidak

multiple comparisons test. Data are represented as mean±s.e.m. **f, g.** Following 4-OHT administration (P3–P5), mice were subjected to an elevated plus maze (EPM, **f**) or novel object recognition test (NOR, **g**). No significant differences were observed either for the distance and exploration time (%) during the EPM or the preference index (PI) during the NOR between control and $\gamma 2^{f/f}$ mice. EPM: N=9 for control and N=8 for $\gamma 2^{f/f}$ mice; NOR: N=9 for control and N=7 for $\gamma 2^{f/f}$ mice. p= 0.6058 for distance (**f;** two-tailed Mann-Whitney U test), p= 0.5788 for exploration time (**f**), p= 0.8533 for PI (**g**), two-tailed Student's t test. All data are represented as mean±s.e.m. ns: not significant. 4-OHT: 4-hydroxytamoxifen. Source data are provided as a Source Data file.

Supplementary Figure 6. No changes in the density and distribution of PV⁺ and GAD67⁺ interneurons of the mPFC in γ2$^{f/f}$ mice

a. Representative confocal images of PV⁺ (magenta) and GAD67⁺ interneurons of the IL region of the mPFC (inset) in four-month control (top) and γ2$^{f/f}$ (bottom) mice. Arrowheads indicate PV⁺/GAD67⁺ neurons (white) and PV⁻/GAD67⁺ neurons (yellow). Scale bar: 50 µm. **b-e.**

Quantification of PV$^+$ and GAD67$^+$ cell densities (**b, c**) and normalized distribution from layer 1 to 6 (**d, e**) in control (dark blue) and γ2$^{f/f}$ mice (light blue). Control mice: n= 11 technical replicates from N= 4 mice (biological replicates) and γ2$^{f/f}$ mice: n= 14 technical replicates from N= 4 mice (biological replicates); p= 0.719 for (**b**) and p= 0.413 for (**c**); two-tailed linear mixed models with likelihood ratio tests applied to account for repeated measures (open dots) within each animal (closed dots). For distributions, p= 0.9993 / 0.9908 / 0.9997 / 0.9950 / 0.9987 / 0.9807 / >0.9999 / >0.9999 / 0.8875 / 0.9997 for normalized bins covering layers 1-6 in (**d**) and p= >0.9999 / 0.9968 / 0.8290 / >0.9999 / >0.9999 / 0.9510 / 0.9968 / 0.9936 / 0.3770 / 0.9968 for normalized bins covering layers 1-6 in (**e**); two-way ANOVA test followed by post-hoc Sidak multiple comparisons test. Data are presented as mean±s.e.m. ns: not significant. **f.** Confocal images of SMI-312 (green) and PV (magenta) labeling, showing axonal compartments identified using a mask based on SMI-312$^+$ signal (white ROIs) in the deep layers of the IL region in a control mouse. Yellow arrowheads indicate some examples of detected PV+ axons. Scale bar: 20 µm. **g**. Dot plots of PV$^+$ fluorescence intensity within axonal compartment masks in control (dark blue) and γ2$^{f/f}$ (light blue) mice. Control mice: n= 8 technical replicates from N= 3 mice (biological replicates) and γ2$^{f/f}$ mice: n= 8 technical replicates from N= 3 mice (biological replicates); p=0.5 for (**g**); two-tailed linear mixed models with likelihood ratio tests applied to account for repeated measures (open dots) within each animal (closed dots). Data are presented as mean±s.e.m. ns: not significant. Source data are provided as a Source Data file.

Supplementary Figure 7. Electrophysiological validation of CNO-mediated activation in hM3Dq-expressing PV interneurons.

a, b. Whole-cell patch-clamp recordings from a layer 5 dTomato[+] PV interneurons (**a**) and a layer 2/3 dTomato[-] pyramidal cell (**b**) in acute mPFC slices of $\gamma2^{f/f}$ mice injected with pAAV-S5E2-hM3D(Gq)-P2A-dTomato. Responses to depolarizing current ramps are shown before (Control, top) and after 3-min bath applications of 10 µM CNO (CNO, bottom). Note the CNO selectively reduced the latency to first action potential and increased firing frequency in the PV interneuron (red), but not in the pyramidal neuron (black). **c, d**. Dot plots of the time of first action potential (**c**) and the firing frequency (**d**) of dTomato[+] PV interneurons (red) and neighboring dTomato[-] pyramidal cells (black). N=3 injected $\gamma2^{f/f}$ mice; p= 0.0467 for dTomato[+] PV interneurons and p= 0.9529 for dTomato[-] pyramidal cells in (**c**), two-tailed paired Student's

t test; p= 0.0156 for dTomato^{+} PV interneurons and p= 0.7926 for dTomato^{-} pyramidal cells in

(**d**), two-tailed Wilcoxon matched-pairs test and two-tailed paired Student's t test, respectively.

Data are represented as mean±s.e.m. *p<0.05, ns: not significant. Ctrl: control; CNO:

clozapine-N-oxide. Source data are provided as a Source Data file.

Supplementary Figure 8. Fear conditioning and context extinction sessions following chemogenetic-mediated activation of PV interneurons or in vivo FKN infusions in γ2$^{f/f}$ mice

a-d. Comparison of freezing behavior during fear conditioning and contextual extinction sessions between dTom-injected (white) and hM3Dq-dTom-injected (magenta) γ2$^{f/f}$ mice (**a, b**) and between VC-infused (white) and FKN-infused (yellow) (**c, d**). No significant differences were observed between these respective matched groups. dTom-injected γ2$^{f/f}$ mice: N=13 and hM3Dq-dTom-injected γ2$^{f/f}$ mice: N=14; VC-infused mice: N=8 and FKN-infused mice: N=8; p= 0.3668 / 0.8045 / 0.3195 / 0.2096 / 0.7111 / 0.7241 for BL and trials 1-5 in (**a**) and p= 0.4876 / 0.4341 / 0.7070 / 0.9611 / 0.6067 / 0.7291 / 0.2116 / 0.3426 / 0.9825 for 2-18 min in (**b**); p= 0.6656 / 0.9988 / 0.9988 / 0.2602 / 0.7824 / 0.8382 for BL and trials 1-5 in (**c**) and p= >0.9999 / 0.9971 / >0.9999 / >0.9999 / 0.9850 / 0.9844 / >0.9999 / >0.9999 / >0.9999 for 2-18 min in (**d**); two-way ANOVA test followed by post-hoc Sidak multiple comparisons test. Data are represented as mean±s.e.m. Source data are provided as a Source Data file.

Supplementary Figure 9. Open field behavior and in vivo recordings in control and γ2$^{f/f}$ mice

a. The open-field behavior of mice was measured at the center and periphery of the arena while they were connected *via* a cable to the amplifier. **b-d.** Dot plots of the total distance traveled (**b**), time spent (**c**) and time of immobility (**d**) in the center (C) and periphery (P) of the arena for control (dark blue) and γ2$^{f/f}$ mice (light blue). N=5 control mice and N=5 γ2$^{f/f}$ mice; p= 0.4307 in center and p= 0.9553 in periphery for (**b**); p= 0.7017 in center and p= 0.7033 in periphery for (**c**); p= 0.8587 in center and p= 0.7986 in periphery for (**d**); two-tailed Student's *t* test; All data are represented as mean±s.e.m. ns: not significant. **e, f.** Dot plots of the number of crossings and average speed during the open field test in control (dark blue) and γ2$^{f/f}$ mice (light blue). N=5 control mice and N=5 γ2$^{f/f}$ mice; p= 0.7395 for (**e**); p= 0.7886 for (**f**); two-tailed Student's. All data are represented as mean±s.e.m. ns: not significant. **g-h.** Theta phase-

low-gamma amplitude (**g**) and theta phase-high-gamma amplitude (**h**) coupling did not change in the center compared to the periphery in both control (dark blue) and $\gamma2^{f/f}$ (light blue) mice. N=5 control mice and N=5 $\gamma2^{f/f}$ mice; p= 0.204 in control mice and p= 0.6165 in $\gamma2^{f/f}$ mice for (**g**); p= 0.2273 in control mice and p= 0.7238 in $\gamma2^{f/f}$ mice for (**h**); two-tailed Student's *t* test. All data are represented as mean±s.e.m; ns: not significant. Source data are provided as a Source Data file.

Supplementary Figure 10. No disruption of theta-high-gamma coupling during tone fear extinction

a. In vivo local field potential (LFP) recordings illustrating the coupling and uncoupling of gamma and theta oscillations. **b, f.** Theta phase-low-gamma (**b**) and theta phase-high-gamma (**f**) amplitude coupling increases during tone compared to inter-trial interval (ITI) in both control (dark blue) and $\gamma 2^{f/f}$ (light blue) mice. N=5 control mice and N=5 $\gamma 2^{f/f}$ mice. p= 0.0007 in control mice and p= 0.0003 in $\gamma 2^{f/f}$ mice for (**b**); p= 0.0041 in control mice and p= 0.0001 in $\gamma 2^{f/f}$ mice for (**f**); two-way ANOVA test followed by a Fishers Least Significant Difference (LSD) test. All data are represented as mean±s.e.m. **p<0.01, ***p<0.001. **c, g.** The ratio of theta phase-low-gamma (**c**) and theta phase-high-gamma (**g**) amplitude coupling during the tone relative to ITI is not significantly different between the two groups. N=5 control mice and N=5 $\gamma 2^{f/f}$ mice. p= 0.6585 for (**c**) and p= 0.2928 for (**g**); two-tailed Student's *t* test. All data are represented as mean±s.e.m. ns: not significant. **d-e, h-i.** Theta phase-low-gamma (**d, e**) and theta phase-high-gamma (**h, i**) amplitude coupling at early (E), intermediate (I) and late (L) stages of the extinction retrieval session in control (**d, h**) and $\gamma 2^{f/f}$ (**e, i**) mice. N=5 control mice and N=5 $\gamma 2^{f/f}$

mice. p= 0.0448 / 0.4048 / 0.0073 for early, intermediate and late stages for (**d**) and p= 0.0478 / 0.0140 / 0.0381 for early, intermediate and late stages for (**e**); p= 0.0008 / 0.0003 / <0.0001 for early, intermediate and late stages for (**h**) and p= 0.002 / 0.0004 / 0.0003 for early, intermediate and late stages for (**i**); two-way ANOVA test followed by Tukey's multiple comparison test. All data are represented as mean±s.e.m. *p<0.05, **p<0.01, ***p<0.001, ****p<0.0001, ns: not significant. Source data are provided as a Source Data file.