Title: Replication stress increases de novo CNVs across the malaria parasite genome

Authors: Noah Brown[1], Aleksander Luniewski[1], Xuanxuan Yu[2,3], Michelle Warthan[1], Shiwei Liu[1,4], Julia
Zulawinska[1], Syed Ahmad[1], Molly Congdon[5], Webster Santos[5], Feifei Xiao[2], Jennifer L Guler[1*]

Affiliations:

[1]University of Virginia, Department of Biology, Charlottesville, VA, USA

[2]Unifersity of Florida, Department of Biostatistics, Gainesville, FL, USA

[3]Unifersity of Florida, Department of Surgery, College of Medicine, Gainesville, FL, USA

[4]Current affiliation: Indiana University School of Medicine, Indianapolis, IN, USA

[5]Virginia Tech, Department of Chemistry, Blacksburg, VA, USA

*Corresponding Author

Email: jlg5fw@virginia.edu

ABSTRACT

Changes in the copy number of large genomic regions, termed copy number variations (CNVs),
contribute to important phenotypes in many organisms. CNVs are readily identified using conventional
approaches when present in a large fraction of the cell population. However, CNVs that are present in
only a few genomes across a population are often overlooked but important; if beneficial under specific
conditions, a de novo CNV that arises in a single genome can expand during selection to create a larger
population of cells with novel characteristics. While the reach of single cell methods to study de novo
CNVs is increasing, we continue to lack information about CNV dynamics in rapidly evolving microbial
populations. Here, we investigated de novo CNVs in the genome of the *Plasmodium* parasite that causes
human malaria. The highly AT-rich *P. falciparum* genome readily accumulates CNVs that facilitate rapid
adaptation to new drugs and host environments. We employed a low-input genomics approach
optimized for this unique genome as well as specialized computational tools to evaluate the de novo
CNV rate both before and after the application of stress. We observed a significant increase in genome-
wide de novo CNVs following treatment with a replication inhibitor. These stress-induced de novo CNVs
encompassed genes that contribute to various cellular pathways and tended to be altered in clinical
parasite genomes. This snapshot of CNV dynamics emphasizes the connection between replication
stress, DNA repair, and CNV generation in this important microbial pathogen.

INTRODUCTION

Changes in the copy number of large genomic regions, termed copy number variations (CNVs), are a
source of phenotypic diversity for many organisms (as reviewed in [1-3]). The structure of CNVs and
their formation is complex; they can involve a few base pairs or whole genes, encompass many
structural forms (e.g. tandem arrays, inversions, translocations), and be generated by a range of DNA
repair pathways (e.g. recombination and end-joining pathways). CNVs are especially important for
rapidly evolving microbes such as bacteria, yeast, and viruses by contributing to antimicrobial resistance,
nutrient acquisition, and pathogenesis [4-6]. CNVs also contribute to cancer growth and progression ([7-
10] and reviewed in [11, 12]). Further, we are beginning to appreciate CNVs as drivers of other human
disorders and disease susceptibility including blood, metabolic, neurological, and infectious diseases [13-
21].

Increased access to genome sequencing has facilitated the identification of these important genomic
rearrangements, especially following selection. CNVs that are identified using standard "bulk" analysis
approaches (e.g.. read coverage methods) are present in a large fraction of the cell population (>50%
[2]). However, those in a minority of genomes, or even a few genomes, are "averaged" away during
analysis steps. This artifact limits our ability to assess how CNVs arise and contribute to the genomic
diversity of individual cells within a population. Such diversity is important; if beneficial under specific
conditions, a CNV that arises in a single genome can expand during selection into a larger population of
cells with novel characteristics [22, 23]. This rapid expansion is exemplified when minor bacterial
populations with higher gene copy numbers confer "heteroresistance" during clinical antibiotic selection
[24, 25, 26]. In another example, higher levels of intra-tumor heterogeneity in gene copy number
predict a poorer cancer prognosis [27-29].

In order to observe a genome's evolutionary potential in the absence of selection, we require
approaches specifically designed to detect CNVs that are not present in a predecessor "parental"
genome. Due to their rare and novel nature, these events are commonly termed "de novo" CNVs [3, 30-
34]. Early experimental progress detecting de novo CNVs involved cloning individual cells, which takes
time, is prone to contamination, and prevents detection of detrimental CNVs [30-32]. Misalignment of
short-reads to reference genomes (i.e. discordant or split reads) can also indicate the presence of de
novo CNVs, but false positives are common if matched normal samples are not available (reviewed in
[35]).

Recent methods that isolate single cells have been successful at assessing de novo CNVs during
experimental evolution, disease progression, and tissue development (as reviewed in [35-39]). De novo
CNVs have been tracked in evolving yeast genomes using flow cytometry to quantify fluorescent
reporters integrated into a specific selectable locus [33]. This approach is sensitive but provides a limited
view of CNV dynamics by focusing on a single or few specific loci. Single cell transcriptomics, which infers
gene copy number using mRNA abundance, can identify large structural changes across genomes from
heterogeneous tumor samples [40-43]. This approach averages relative expression over Mb-sized
genomic regions and thus, is not applicable to identify smaller de novo CNVs. Single cell genomics,
where individual genomes or nuclei are isolated and amplified to a level that can be sequenced, has
been used to directly quantify de novo CNV rates in brain tissue and cancer cells [27, 44-49]. A recent
approach, termed direct library preparation, is free from amplification steps, thus limiting genome
skewing; however, this method is less accessible due to the requirement for specialized cell dispensers
(microfluidics or piezoelectric) and has not been tested with CNVs smaller than 500kb [50]. While the
reach of single cell methods is expanding, we continue to lack information about de novo CNVs in
microbes and their dynamics in evolving populations.

The *Plasmodium* parasite that causes malaria readily accumulates kb-sized CNVs in its genome [51-54].
CNVs impact the survival of malaria, allowing this parasite to evade clinical detection [55], expand
beneficial gene families [56], invade new host cells [57], and develop clinical antimalarial resistance [58-
60]. We are specifically interested in the genetic diversity of one species of malaria, *P. falciparum*, since
this may explain its rapid adaptation to new drugs and host environments [61]. Due to its relatively
small, AT-rich genome (23 Mb, 19.4% GC [62]), low-input genomics is challenging in this single cell
protozoan. However, we previously optimized a single cell genomics approach and developed novel
computational tools to evaluate de novo CNVs in the *P. falciparum* genome [63] (see *Materials and*
Methods for tool details).

Here, we combined these advancements and made additional improvements to investigate de novo CNV
formation in the *P. falciparum* genome. Since various types of cellular stress can induce genetic change
(reviewed in [2, 64-66]), we also evaluated the impacts of replication stress on de novo CNV
identification. Using our low-input genomics pipeline, we observed that replication stress increased the
number of de novo CNVs across the parasite genome. This study of genome dynamics, along with
improved tools, increases our understanding of how stress can stimulate rapid microbial evolution.

RESULTS

Refined low-input genomics pipeline increased efficiency and accuracy

We developed a robust pipeline for assessing the frequency of de novo CNVs in the *Plasmodium* genome
based on our prior studies (**Fig. 1**). We adapted our single cell genomics method to improve parasite
isolation and whole genome amplification steps ([63], **Fig. 1A, Fig. S1**). In this modified protocol, we
used fluorescence-activated cell sorting (FACS) to isolate parasites to improve efficiency and modified
aspects of our whole genome amplification method to improve coverage (**Table S1**, **Fig. S1**). We also
sorted low-cell populations to increase accuracy (e.g. 2-cells per well, **Fig. S2**), and added quality control
steps that confirmed our samples were of high quality prior to short-read sequencing. The resulting low-
input genomics pipeline consisted of basic steps including parasite isolation using flow sorting, whole
genome amplification using a modified MALBAC-based approach, quality control confirmation, short-
read sequencing, and CNV analysis (**Fig. 1A**).

Figure 1. Low-input genomics approach for analysis of malaria parasite genomes under stress. Early erythrocytic stage *P.*
falciparum parasites grown in vitro were treated with a replication inhibitor (DSM1) or the solvent control (dimethylsulfoxide)
followed by recovery and reinvasion to produce a new round of ring stage parasites. **A.** For low-input genomics, parasite-
infected erythrocytes were isolated using flow sorting (10-cell control or 2-cell samples). Parasite genomes were amplified using
a modified MALBAC-based whole genome amplification approach (**Fig. S1**). Quality control steps involved DNA quantification to
assess amplification success, droplet digital PCR to assess parasite genome amplification, and PCR-high resolution melting to
assess sample cross-contamination. After Illumina short-read sequencing, reads are filtered, trimmed and used as input for
copy number variation analysis using HapCNV and LUMPY. **B.** Parasite erythrocytic life cycle including approximate timing (h),
genome number (n), and effect of treatment.

Replication stress followed by a recovery period led to isolation of healthy parasites

To explore the impact of replication stress on de novo CNV generation, we treated the parasites with an
antimalarial that inhibits DNA replication by limiting pyrimidine pools; DSM1 targets *Plasmodium* the
dihydroorotate dehydrogenase enzyme [67]. Application of DSM1 for an extended period kills parasites
by perturbing DNA synthesis (>48hrs at 10x the EC_{50}, [68]). However, short-term treatment reversibly

stalls parasite replication similar to another replication inhibitor, aphidicolin (**Fig. S3**). For low-input
genomics, we applied DSM1 to ring-stage parasites for a brief time (12hr, **Fig. S4A and B**). Similar to our
pilot experiment (**Fig. S3A**), we observed that treated parasites stalled prior to replication (**Fig. S4C and
E**) and slightly decreased their growth rate compared to untreated parasites (**Fig. S4G**). We harvested
viable parasites after a recovery period (~30hrs, **Fig. S4H**), where we allowed parasites to complete an
additional round of replication and erythrocyte invasion to produce those that have a single, haploid
genome (rings, **Fig. S4D and F**). The recovery stage was essential as it allowed parasites to repair the
DNA damage accumulated during treatment. As a control for cross-sample contamination between
isolation wells, we isolated and amplified untreated parasites with different genetic backgrounds (*FCR3*
vs *Dd2*, **Table 1, Fig. S5**).

Prior to isolation of low-cell populations, we confirmed that untreated and DSM1-treated parasites were
at a similar life cycle stage and viability using staining for mitochondrial membrane potential (**Table 1,
Fig. S5A**); we saved a portion of these samples for parasite population sequencing (i.e. bulk samples).
We then proceeded to isolate small populations of viable, ring stage parasites using FACS (**Fig. S5B**) for
whole genome amplification.

Table 1: Parasite density, staging, and health at isolation for low-input genomics.

Line/Treatment*	Mean % Parasitemia$	Mean % Rings	Mean % Viability@
Untreated *FCR3*^	0.6%	90%	91%
Untreated *Dd2*^	0.8%	87%	94%
DSM1-treated *Dd2*	0.6%	77%	95%

*Treatment conditions: 1μM DSM1 (~10x EC50) for 12hrs prior to 30.8hrs of recovery. Untreated samples were incubated with
DMSO as a solvent control for 12hrs and allowed to recover for 28-30hrs (**Fig. S4H**). $Parasitemia was determined by calculating
the number of infected erythrocytes (SYBR Green+) compared to uninfected erythrocytes (SYBR Green-) (**Fig. S4**).@Viability of
parasites was determined by measuring the mitochondrial membrane potential (Mitoprobe+) and calculating % of total SYBR
Green-based parasitemia (plots presented in **Fig. S4D, S4F,** and **Fig. S5**). ^*FCR3* (Africa) and *Dd2* (Southeast Asia) parasite lines
are from distinct geographic origins [69, 70].

Quality assessments showed effective isolation and amplification of low-input samples

We sorted low-cell populations from each parasite group into 60 wells of a 96-well plate (*FCR3*,
untreated *Dd2*, and treated *Dd2*), including 6 wells with 10-cells and 56 wells with 2-cells. Ten-cell wells
served as positive controls for the whole genome amplification step and 2-cell wells provided the
optimal balance between sorting accuracy (**Fig. S2**) and de novo CNV detection. Zero cells were sorted
into the top 2 rows of the plate ("no-cell" wells). After parasite lysis and applying *Pf*MALBAC version-2
whole genome amplification (**Fig. S1**), we assessed amplification success using three approaches. First,
we measured the resulting DNA quantity across 80% of the amplified wells (**Fig. S6**). On average,
MALBAC amplification in wells that contained sorted parasites yielded ~120ng of total DNA per reaction,
with a ~10% increase in DNA for 10- vs 2-cell samples (mean of 127ng versus 116ng total, respectively).
We did not detect position-based bias across plates or appreciable amplification from no-cell wells, but
we did observe that treated *Dd2* wells had ~3-fold lower levels of amplification than other samples

(mean of 51ng versus 151ng total, respectively). There was little difference in mean amplified DNA
amounts between the two untreated sample groups (*FCR3* at 146ng and untreated-*Dd2* at 152ng).

Second, we performed droplet digital PCR (ddPCR) for parasite-specific genes on approximately one-
third of wells post isolation to confirm the amplification of the parasite genome. DdPCR for *pfmdr1* and
pfhsp70 displayed that wells with measurable DNA contained amplified parasite DNA (**Fig. S6** and **S7**).
Additionally, we confirmed that 2-cell wells with very low total DNA amounts (**Fig. S6**) were positive for
parasite genomes while "no-cell" wells did not show evidence of parasite material (**Fig. S7**).

Finally, we employed high-resolution melting analysis to profile a drug resistance marker that differs
between *FCR3* and *Dd2* parasites. By assessing the *pfdhps* SNP profile of amplified genomes and
comparing it to the parental profile in ~10% of samples, we confirmed that there was no evidence of
cross-sample contamination during the preparation and amplification steps (**Fig. S8**). Therefore, we
proceeded to sequence the amplified bulk and low-input samples (**Table S2**).

Coverage deviation and SNP profiles exhibited expected trends in low-input samples

We sequenced 3 bulk samples and 90 low-input samples using Illumina short-read sequencing (**Table 2**
and **Table S3**). Overall, sequencing proceeded well as indicated by coverage and coverage deviation of
the bulk samples, as well as an equivalent mean mapping quality across all samples (**Table 2**). Because
we noticed that treated *Dd2* wells had lower levels of DNA following amplification (**Fig. S6**), we
sequenced higher amounts of material for this condition; this choice impacted mean coverage levels
where treated *Dd2* samples had ~4-times higher coverage than untreated *Dd2* samples (**Table 1**). The
percent of total reads that mapped to the *P. falciparum* genome was high across all samples (mean of
67%), indicating efficient amplification of the parasite genome and little contribution of environmental
contamination during sample amplification. As expected based on previous studies [63], coverage
deviation was ~3-fold higher in low-input samples when compared to bulk samples, reflecting the bias of
the whole genome amplification step to over- or under-amplify specific genomic regions.

We removed five low-input samples from further analysis based on low coverage levels; on average,
excluded samples had ~7-times lower coverage than other low-input samples (**Table S3**). Of the
remaining samples, 18 were 10-cell samples and 57 were 2-cell samples. Although mean coverage was
~2-fold higher for 2-cell samples (due to the higher coverage of treated low-input samples), mean
normalized deviation was similar between 10- and 2-cell samples (3.5 and 3.1, respectively).

To evaluate the quality of the sequencing data, we tracked SNPs in the low-input samples compared to
bulk samples. Despite some variation due to the non-clonal nature of parasite lines (see *Materials and*
Methods), low-input SNP profiles were similar to their corresponding bulk sample (**Fig. S9A**). After
normalizing total SNPs to mapped reads, we detected a lower rate of SNPs in treated samples compared
to untreated counterparts (p value of 0.0001, **Fig. S9B**). We did not detect a correlation between
normalized total SNPs and amplification quality (**Fig. S9C**, R_2: untreated *Dd2*, 0.01; treated *Dd2,* 0.01);
we did observe a positive correlation between SNP number and coverage depth in treated versus
untreated *Dd2* samples (**Fig. S9D**, R_2: untreated *Dd2*, 0.66; treated *Dd2,* 0.27), indicating that the
difference in SNP numbers is likely due to varying sensitivity at different levels of read coverage [71, 72].

202
203
204 **Table 2: Sequencing Summary for low-input samples and paired bulk samples**

		No. of samples **	Mean no. mapped reads per sample	Mean coverage per sample	Mean Coefficient of Variation (CV#)	Mean mapping quality
Bulk	Untreated *FCR3*	1	9,966,137	54.9	59.5	58.2
Low-input	Untreated *FCR3*	16	2,694,209	13.7	105.1	58.2
Bulk	Untreated *Dd2*	1	5,735,107	33.6	33.3	58.5
Low-input	Untreated *Dd2*	33	2,579,674	13.2	82.6	58.3
Bulk	Treated *Dd2*	1	72,380,215	416.3	35.2	58.5
Low-input	Treated *Dd2**	36	9,936,903	53.7	86.5	58.4

205 *On average, ~4x more material was loaded on the flow cell for the treated samples than the untreated samples due to lower
206 initial amplification yields (**Fig. S6**).
207 **Excludes samples that were removed due to low coverage. For low-input samples, includes both 10- and 2-cell samples.
208 #CV is the coefficient of variation of normalized read abundance as in [63]
209
210

Experimental and computational advances improved known CNV calls across low-input samples

For the current study, we employed two different CNV calling methods in low-input samples. HapCNV is
a novel read coverage-based CNV calling method specifically designed for low cell data from haploid
genomes [73]. In contrast to traditional methods that arbitrarily select reference samples for CNV data
normalization, HapCNV constructs a genomic location (or bin)-based pseudo-reference as a comparison
baseline. This step systematically alleviates amplification bias for the identification of de novo CNVs.
LUMPY is an established CNV calling method that exhibits high sensitivity due to the incorporation of
multiple CNV signals (i.e. split and discordant reads) generated from short-read sequencing. It is
particularly well-suited for detecting low-frequency variants in low-coverage datasets; however, for
many CNV callers, high sensitivity leads to higher false positives [74-76].

Using these two distinct CNV calling methods, combined with a recently developed CNV counting
approach, we evaluated the presence of known CNVs in our 2-cell samples (**Fig. 2A**). The identification of
known CNVs (i.e. those identified in the bulk sample, see *Materials and Methods*) within low-input
samples displays the utility of the specific CNV calling method for different size CNVs in specific genome
locations. In our previous study, we identified 2 of the 3 known CNVs in ~10% of single cell genomes
[63]. In the current study, we identified the *pfmdr1* amplicon in 100% of 2-cell samples using HapCNV
(57/57) and 79% of samples using LUMPY (45/57). We did not identify the *pf11-1* amplicon in any 2-cell
samples using HapCNV (0/57) but detected this locus in 75% of samples using LUMPY (43/57). Finally,
we identified *pf332* CNVs in 46% of 2-cell samples using HapCNV (26/57) and 100% of samples using
LUMPY (57/57). Although our two studies used different CNV calling methods and are not directly

comparable, the overall improvement in the detection of known CNVs in the current study is likely due
to advances in both the whole genome amplification method to limit amplification bias (**Table S1**) and
recently developed analyses approaches. When we evaluated the detection of three known CNVs in *Dd2*
low-input samples, we observed a somewhat higher rate of known CNV detection by either method in
treated samples (**Fig. 2B**, HapCNV: mean of 1.2 out of 3 total CNVs for untreated and 1.7 for treated
(increase of 42%), LUMPY: mean of 2.1 out of 3 total CNVs for untreated and 2.9 for treated (increase of
38%, **Table S4**).

De novo CNVs in low-input samples consisted of rare and common CNVs

We next sought to quantify de novo CNVs in low-input samples. We defined de novo CNVs as those not
present in the bulk sample and we categorized them based on their frequency in low cell samples.
"Common" CNVs were present in a larger number of genomes (≥10% of the same sample type, i.e.
untreated or treated), and "rare" CNVs were those that occurred in a small proportion of samples (<10%
of the sample type) (**Fig. 2A**). Overall, LUMPY detected more total de novo CNVs than HapCNV across all
samples (12-fold), and the proportion of rare versus common CNVs varied depending on the method
(18% vs 82% for HapCNV, 61% vs 39% for LUMPY, respectively, **Table S4**). Additionally, de novo CNVs
were more often identified as duplications than deletions for both CNV calling methods (**Fig. 2C**).

To understand the nature of common and rare CNVs, we also assessed how often their locations
overlapped across the two sample types (untreated and treated, **Fig. S10**). This analysis is useful for
tracking common/rare category utility and relevance. For utility, this step acts as a sanity check since, by
definition, we do not expect rare CNV locations to overlap as often as common CNVs. For relevance, de
novo CNVs with conserved locations across sample types are less likely to represent true CNVs newly
arising in a genome. As expected, we identified many common CNVs with conserved locations (22% for
HapCNV and 52% for LUMPY, **Fig. S10A**). The lower rate of overlapping calls across samples for HapCNV
is likely due to the bin-based normalization strategy to remove amplification artifacts [73]. Conversely,
we found that rare CNVs were predominantly called in unique genome locations (94% for HapCNV and
97% for LUMPY, **Fig. S10A**), supporting their novel nature. This pattern was consistent when we
randomly down-sampled all sequencing data to the lowest read coverage prior to CNV calling (1.3
million reads, **Fig. S10B**). This comparison not only highlights the suitability of the common and rare
categories but also the difference between the CNV calling methods. Based on these observations, for
the following analysis, we assessed common and rare CNVs using both methods to capture the broadest
view of stress effects on CNV generation.

Genome-wide de novo CNVs increased following replication stress

When we compared de novo CNVs in genomes with and without replication stress, we found that results
were consistent regardless of the CNV calling method (**Fig. 2**). While the proportion of duplication and
deletions did not change appreciably with treatment (**Fig. 2C**), we identified a highly significant
difference in de novo CNVs between treated and untreated 2-cell samples (p value of 0.0002 for
HapCNV and <0.0001 for LUMPY, **Fig. 2D**). This pattern was consistent when we down-sampled all
sequencing data (p value of 0.001 for HapCNV and <0.0001 for LUMPY, **Fig. 2E**), indicating that the
difference in de novo CNV counts between treatments was not due to read coverage. When we
assessed common and rare CNV categories, we once again observed highly significant differences
between treated and untreated 2-cell samples in common CNVs using both methods (common: p value

0.0004 for HapCNV and <0.0001 for LUMPY, **Fig. 2F;** rare: p value of 0.008 for HapCNV and <0.0001 for
LUMPY, **Fig. 2G**).

When we compared the proportion of de novo CNVs relative to total CNVs per sample, rare CNVs were
significantly increased over common CNVs (p value of 0.003 for HapCNV and 0.009 for LUMPY, **Fig. 2H**).
This difference persisted regardless of down-sampling (p value of 0.02 for HapCNV and 0.004 for LUMPY,
Fig. S11) and is in line with our assessment above that rare CNVs are more likely to be novel in nature
(**Fig. S10**). Overall, we detected a ~2-3-fold increase of de novo CNVs in treated samples, regardless of
the CNV calling method (**Table 3**). Once again, rare CNVs displayed the largest increase following
treatment (~3-4-fold, **Table 3, Fig. 2I**).

De novo CNVs represented diverse cellular pathways with clinical benefits

When we compared overlaps between the HapCNV and LUMPY (**Fig. 3A**), we detected a set of CNV
regions that was consistent within sample groups (5 for untreated and 38 for treated, **Table S5**). The
frequency of these "high-confidence" CNVs also reflected the increase in CNVs following replication
stress (with knowns excluded, ~15-fold increase in treated samples). Overall, high-confidence CNVs were
located on the majority of chromosomes (12 of 14, **Fig. 3B**) and represented both duplications and
deletions (**Fig. 3C**). Of note, approximately half of these regions were identified as "rare" by both CNV
calling methods across treated samples (17/36, 47%; **Table S5**), indicating that novel CNVs were
stimulated in parasite genomes under stress. When we searched for genes that are covered by these
regions, we identified 26 genes (across 3 de novo CNV regions) and 198 genes (across 37 de novo CNV
regions) in untreated and treated *Dd2* samples, respectively (**Table S5**). Emphasizing their random
nature, genes encompassed by the CNV regions represented diverse protein classes (**Fig. 3D**) and no
gene ontology (GO) categories were significantly enriched (using an FDR adjusted p value of 0.05, **Table
S6**).

Finally, we evaluated whether CNVs that arose under stress were important for parasite survival by
comparing genes covered by our treated CNV regions to those from the largest catalogue of clinically
relevant CNVs to date. This list of high frequency CNVs was previously called using genomes from 2855
parasite isolates from 21 malaria-endemic countries and represented genes from larger (>300bp), high-
quality, core genome variants [59]. Assuming ~5000 genes in the core *P. falciparum* genome [77], we
found that genes from the two lists were ~2-times more likely to overlap than by chance (chi-square
odds ratio of 2.5 for HapCNV-treated CNV list and 1.6 for LUMPY-treated CNV list). This significant
association between CNV locations suggests that stress-induced de novo CNVs have the potential to be
beneficial in the clinical environment (Fisher's exact p value <0.0001 for HapCNV-treated CNV list and
0.03 for LUMPY-treated CNV list).

Table 3. Mean CNV counts per 2-cell sample using two analysis methods.

CNV detection method$^\$$	Condition (2-cell only)	Rare CNVs per sample*	Common CNVs per sample*	Combined de novo CNVs per sample^

HapCNV	treated *Dd2*	7	23	15
	untreated *Dd2*	1.7	15	8
	fold change	**4.4**	**1.6**	**1.9**
LUMPY	treated *Dd2*	297	163	230
	untreated *Dd2*	99	68	84
	fold change	**3.0**	**2.4**	**2.7**

*Includes both duplications and deletions. Values are calculated by taking the mean of CNV counts per sample within each category. ^De novo CNV counts combined the subcategories of rare (<10% of samples) and common CNVs (>10% of samples, absent in bulk). $CNV analysis performed using all reads.

Figure 2. Low cell genomics displays an increase in de novo CNVs following replication stress. Number of CNVs from untreated (U-*Dd2*) and treated (T-*Dd2*) 2-cell samples from two CNV analysis methods: HapCNV and LUMPY. Statistics for all plots use an unpaired T-test with two tailed Welch's correction (****:p value <0.0001; ***: <0.001; **: <0.01; *:<0.05; no stars: not significant). Analyses include all reads, unless otherwise indicated (i.e. panel E is down-sampled). Line at mean value for

each dataset. **A**. Depiction of CNV categories used in the analysis. Known CNVs (orange) are detected in bulk samples and
present in all low cell samples. De novo CNVs are not present in bulk samples are considered common (green, <10%) or rare
(teal, <10%) depending on their frequency across the 2-cell samples. **B**. Detection of three known CNVs in 2-cell samples.
Known CNVs were identified in *Dd2* bulk sequence (either *pfmdr1, pf11-1, or pf332*). 0: no known CNVs were detected in 2-cell
sample; 1/2/3: one/two/or three known CNVs were detected in 2-cell sample (see **Table S4** for sample counts). **C.** Proportion of
total CNVs detected as duplications (Dup) or deletions (Del) in untreated (U) or treated (T) 2-cell *Dd2* samples (p values: 0.02 for
Dup and 0.03 for Del from LUMPY). **D**. Detection of de novo CNVs (common and rare combined) from all reads (p values: 0.0002
for HapCNV, <0.0001 for LUMPY). **E**. Detection of de novo CNVs (common and rare combined) from down-sampled reads (p
values: 0.0014 for HAPCNV and <0.0001 for LUMPY). **F**. Detection of common CNVs from all reads (p values: 0.0004 for HapCNV,
<0.0001 for LUMPY). **G**. Detection of rare CNVs from all reads (p values: 0.008 for HapCNV, <0.0001 for LUMPY). **H**. Proportion
of total CNVs detected as rare and common from all reads; pie charts plot the mean but statistics are calculated using all data
points from the rare CNV category (p values: 0.003 for HapCNV, 0.009 for LUMPY). Pie chart size does not represent total de
novo CNV numbers (~12x higher for LUMPY, **Table S4**) I. Mean fold change between rare and common CNVs detected by
HapCNV and LUMPY in untreated and treated *Dd2* 2-cell samples (see **Table 3**).

Figure 3: High-confidence CNVs are located across the genome and represent diverse protein classes. A. Comparison of CNV
calls showing the number of CNV regions consistent across the two CNV calling methods. High-confidence CNVs in untreated
samples (U-*Dd2*, in yellow text); high-confidence CNVs in treated samples (T-*Dd2*, in white text). Central number (grey): CNVs
consistent across all samples and calling methods (includes 2 known CNVs and 2 de novo CNVs, **Table S5**). **B**. Chromosomal
location of high-confidence CNVs identified by both HapCNV and LUMPY methods (green) from untreated (black) and treated
(red) parasites. Only core regions of the genome are included in the representation; subtelomeric regions as defined by Otto, et
al. were omitted. Each CNV region was increased by a factor of 2 to facilitate visualization relative to the rest of the genome. *,
de novo high-confidence CNVs identified in both untreated and treated samples. **C**. Summary of duplications and deletions
represented in high-confidence CNV list. U-*Dd2*, untreated; T-*Dd2*, treated. Mix: sub-regions were called as duplications and
deletions across a single CNV region by the same CNV calling method (i.e. HapCNV or LUMPY). **D**. Panther classification system
v19 protein class comparison. Top chart: protein classes from all annotated *P. falciparum* genes. Bottom charts: protein classes
represented by high-confidence CNV regions in untreated (U-*Dd2*) and treated (T-*Dd2*) samples. UC: unclassified proteins
(green). Other colors are randomly assigned by the program to represent diverse protein classes.

Discussion:

De novo CNVs are not detected when analyzing a population of parasites predominantly because their
signal (e.g. extra reads that align to that region or reads that span breakpoints) is negated by the
overwhelming signal from normal copy number at that genome location. For this reason, assessments of
fewer cells are necessary to investigate de novo CNV generation. Here, we optimized a low-input
analysis pipeline and successfully isolated, amplified, and sequenced *P. falciparum* samples. With
experimental and computational improvements, we were able to increase our rate of parental, or
"known", CNV calling over our prior study [63]. Importantly, we detected non-parental, or "de novo",
CNVs across the parasite genome and replication stress significantly increased their rate formation. By
analyzing ~45 low-input samples per condition, we are limited to observing a subset of the population.
However, our findings demonstrate that replication stress readily drives the rapid generation of de novo
CNVs. Below we cover how this study contributes to our understanding of genome evolution and
integrate it with an overarching model of *P. falciparum* adaptation.

Application of sub-lethal stress without evidence of selection to explore CNV dynamics

To evaluate the impact of replication stress on de novo CNV generation, we applied sub-lethal treatment
to parasites just prior to replication. We observed replication stall and then resume post-treatment,
which provided evidence that we successfully applied non-lethal stress (**Fig. S3 and S4**). Following this
step, we allowed the parasites to complete replication and reinvade new erythrocytes. We reasoned
that this "recovery phase" enabled the repair of the resulting DNA damage, which is likely to be
replication-dependent (reviewed in [1, 78]. Additionally, reinvasion facilitated the isolation of haploid
parasite genomes (1n, **Fig. 1B** and **S4B**), encouraging the detection of de novo CNVs due to limited
contrasting signal [36]. Because of the reinvasion step, which involves an expansion in parasite number
(~3-fold, **Fig. S4G**), there was a potential to select for beneficial DNA changes across the population of
parasites. However, we did not detect evidence of strong selection from SNP profiles (**Fig. S9A**) or high-
confidence CNV regions (**Fig. 3**, **Table S5**). Specifically, we did not observe a preference for CNVs that
encompassed the dihydroorotate dehydrogenase gene (*pfdhodh*), which contributes directly to DSM1
resistance [79], or enrichment of CNVs that include genes from DNA-related pathways (**Table S6**).

Relative comparison using multiple CNV calling methods to appreciate the impact of stress

De novo CNV estimates using single cell methods from neurons, yeast, and human cancer vary greatly
and are difficult to standardize due to the use of different experimental techniques and CNV calling
methods [33, 34, 80]. For these reasons, we are not attempting to compare the rate of *P. falciparum*
CNV formation from this study to those from other organisms. Additionally, this lack of standardization
in the field led us to use multiple CNV calling methods in our analysis. Due to the strengths and
weaknesses of HapCNV and LUMPY, we observed differences in both known (**Fig. 2B**) and de novo CNV
(**Table S4**) calling using the two methods. LUMPY identifies reads that cover breakpoint regions to
sensitively detect CNVs [74]; because we are counting regions with few reads as support, both sensitivity
and the number of false positives are high in this analysis. High known calling rates along with high
numbers of de novo CNVs in our studies exemplified this feature of LUMPY. On the other hand, HapCNV
uses a genome-specific pseudo-reference for normalization, which removes repeated patterns of over-
and under-amplification ([73], and *Materials and Methods*); because we require read coverage to span
3 consecutive 1kb bins, small CNVs in lower coverage genomic neighborhoods are excluded in HapCNV
anaylsis. This limited the detection of smaller known amplicons (*pf11-1* and *pf332*) and led to fewer de
novo CNV calls using this method. Given the high abundance of small CNVs (<300bp) in the parasite
genome [59, 81], HapCNV is likely underestimating their impact in our studies.

Ultimately, the value of our study is in the relative comparison of treated and untreated samples by
both methods. In this case, we assume that false positive CNVs occur at a similar rate across both
groups, which allowed us to confidently assess the impact of stress. Variations in known and de novo
CNV calling described above serve to remind us that no CNV calling method is perfect and combining
them can improve confidence in results [63, 76, 82, 83]. Therefore, we investigated de novo CNV
patterns using the two individual methods (**Figs. 2**, **S10**, **S11**) as well as those that overlapped between
HapCNV and LUMPY (**Fig. 3**, **Table S5**). Importantly, these "high-confidence" CNVs reflected increases
after stress previously detected using the individual tools, albeit at a greater level (3- vs >10-fold
increase). Additionally, we speculate that newly arising CNVs would have distinct locations across
samples; thus, the rare nature and unique locations of high-confidence CNVs emphasized their potential
to be novel (**Table S5**).

De novo CNV categories highlighting existing and novel genome variation

During our investigations, we identified two types of de novo CNVs; those detected in one or a few
genomes (rare) and those detected in more than a few (common). There is no precedence for these CNV
categories in the context of *Plasmodium* biology (i.e. a haploid parasite with asynchronous replication
and schizogeny [84]). However, we propose that tracking these categories helps us to understand the
biological relevance of de novo CNVs in our analysis.

Based on their frequency, common CNVs are either artifacts of low-input procedures/CNV analysis or
represent minor variants that preexist in the population or arise early in the replication cycle. For the
former, bias during the whole genome amplification step (i.e. the repeated pattern of over/under-
amplification that occurs in a reproducible pattern across the parasite genome [63]) and PCR during
library construction have the potential to skew gene copy number and increase the false positive rate
[85, 86]. However, we chose experimental and computational methods designed to limit the
contribution of amplification bias. First, MALBAC amplification itself limits the over-amplification of
genomic regions by avoiding exponential amplification at the earliest steps [85] and we used limited PCR
cycles during library preparation (3 cycles, [63]). These efforts are most clearly shown through the
reduction in CV following MALBAC optimization in both of our studies (by ~39% after modifying the
amplification primer [63] and by 43% after switching to the *Bsu* polymerase, **Table S1**). Second, LUMPY
is not dependent on read coverage and HapCNV specifically addresses amplification artifacts by
removing repeated signal present in all samples [73, 74]. Overall, we detected very few CNVs with
conserved genomic locations across low-input samples, which provides evidence that our methods limit
the effect of amplification bias on the final results; we only identified two high-confidence CNV regions
that had conserved locations across multiple *Dd2* and *FCR3* 2-cell samples (**Fig. 3B**, **Table S5**). In the
future, single-read visualization of long-reads may offer advantages in distinguishing amplification bias
from minor variants and de novo CNVs [87].

Rare CNVs, on the other hand, represent either random noise or true signal from novel CNVs arising in
the genome. We assert that most noise is removed through normalization procedures, especially with
HapCNV, and the impact of remaining false positives are minimized by the relative comparison of our
studies (see above). We identified the majority of rare CNVs in unique genome locations across sample
types, providing evidence that they are not a result of amplification bias where the same CNVs are
repeatedly detected in each sample. Additionally, the greater impact of stress on rare CNVs than
common CNVs (**Table 3** and **Figs. 2G, 2I**) supports their replication dependence. The random nature of

de novo CNVs, as well as the capacity to encompass any gene across the genome, ensures that CNVs can
alter all aspects of parasite biology in response to the host environment. Further, our finding that stress-
induced de novo CNVs tended to exhibit altered copy number in clinical isolates combined with the high
frequency of unique CNVs in previous genome-wide CNV studies [53, 59], directly illustrate the
expansive evolutionary potential of this organism.

Adaptations that encourage de novo CNV formation

The current model of CNV formation in asexual erythrocytic *P. falciparum* parasites is that AT-rich
sequences form hairpins, disrupt replication, and eventually lead to double-strand breaks that are
repaired by error-prone pathways [61]. The evolution of CNVs in this organism is especially interesting
because of its unique genome architecture and alternative repertoire of CNV-generating repair
pathways [62, 88]. Although they arise at many locations across the genome ([51, 52, 59], **Fig. 3A**), *P.*
falciparum CNVs that contribute to adaptation are commonly gene duplications with a relatively simple
structure. Many impactful duplications form in tandem head-to-tail orientation ([61, 79, 89, 90], **Fig.**
4A), which is likely due to a limited repertoire of DNA repair pathways; *P. falciparum* lacks the canonical
nonhomologous end-joining pathway that contributes to CNV formation in other organisms [1, 88].
Instead, parasites use pathways that employ varying lengths of sequencing homology (i.e. homologous
recombination, or HR, and microhomology-mediated repair, **Fig. 4B**). This repair repertoire, along with
an especially high AT-content genome that facilitates CNV formation [61, 79] and a lack of cell cycle
checkpoints that control replication forks during times of stress (reviewed in [91]), likely represent
adaptations that assist haploid *P. falciparum* parasites in accumulating CNVs across their genome (**Fig.**
4A).

Updating the model of P. falciparum genome adaptation

By combining insights from the current study with previous knowledge about CNV formation [1, 61, 65],
we propose a connection between replication stress, DNA repair, and CNV generation in *P. falciparum*
(**Fig. 4B**). Prior studies have shown that stress can either alter levels of proteins essential for HR-based
repair or increase the frequency of DNA breaks [92-94]. With a decrease in HR activity in particular, the
parasite may increase its reliance on alternative error-prone pathways to repair DNA damage.
Microhomology-mediated pathways require less homology and therefore, are more likely to interact
with diverse sequences up- and downstream of a DNA break to generate various length CNVs. So far, the
predominant evidence for this model in *P. falciparum* was the detection of microhomology-mediated
pathway signatures in CNV breakpoints [61]. Our observation of stress-induced de novo CNV formation
(**Table 3** and **Fig. 2I**) further supports this model in *P. falciparum* and is consistent with studies on
diverse organisms [30-32, 95, 96]. Interestingly, the level of de novo CNV stimulation is consistent across
organisms; treatment of mammalian cells with replication inhibitors also leads to a ~3-5-fold increase in
de novo CNVs [30-32].

Even with a change in the copy number of a single region per parasite, the genomic diversity within a
single infected human is expansive due to the sheer numbers of *P. falciparum* parasites (estimated to
reach 10^8 parasites when symptomatic and $>10^{11}$ in severe infection, [97]). This diversity becomes an
obvious advantage as a heterogeneous population prepares asexual parasites to respond to diverse
stressors (**Fig. 4C**). However, one question has been whether random CNVs constitutively arise across
the *P. falciparum* genome or only when under stress. In the former, random amplicons within individual
parasites would position the population to respond rapidly to selection (e.g. antimalarial exposure). In
the latter, specific stressors would stimulate CNV formation to increase genomic diversity. Since some

antimalarials act rapidly [68], we hypothesize that beneficial CNVs must already be present in a few parasites across the population to increase the chances of survival. Indeed, we observed a low level of de novo CNVs across the parasite genome under normal conditions (**Fig. 2D** and **2E**). However, it is also important to understand how parasites respond to stressful environments during infection, including changes in nutrient composition in different hosts, drug treatment during symptomatic infection, or attack from the human immune system. While the current study focused on replication stress, it will be important to evaluate the impact of other sources of stress on *P. falciparum* CNV formation. For example, hypoxia stimulates CNV formation in cancer cells [92] and a proteotoxic drug stimulates genetic change in yeast [98].

Figure 4: Connection between replication stress, DNA repair, and CNV generation in the malaria genome. **A**. Adaptations that encourage CNV formation in the *P. falciparum* genome (underlined). **B**. Summary of how replication stress impacts DNA repair pathways. HR, homologous recombination; NHEJ, non-homologous end-joining; MMEJ, microhomology-mediated end joining; MMBIR, microhomology-mediated break-induced repair; DSB, double-strand breaks. **C**. Benefits of a diverse parasite population for evolutionary potential. Stress elevates the frequency of de novo CNVs across the population, which leads to more rapid evolution of beneficial CNVs (blue cells).

Clinical implications & future questions

P. falciparum causes the majority of worldwide malaria deaths and readily acquires antimalarial resistance [99, 100]. Resistance-conferring CNVs that encompass multiple genes have been identified in both clinical infections [51, 59, 101-105] and laboratory selections [79, 90, 106-111]. Despite their direct contribution to resistance, CNVs may also facilitate the acquisition of point mutations in haploid *P. falciparum*; strong evidence for the close relationship comes from the observation of point mutations *within* amplifications selected in vitro [90, 107, 109, 112, 113]. Once de novo CNVs form during replication of the asexual erythrocytic stage (**Fig. 4**), meiotic recombination during the sexual phase in the mosquito can streamline beneficial CNVs to balance fitness costs [114]. Given the importance of CNVs in *P. falciparum* adaptation, it is not surprising that this organism has evolved strategies to encourage CNV formation (as described in *Adaptations that encourage de novo CNV formation*). Additionally, parasites from specific regions of the world may have an increased propensity to develop

drug resistance [115]. Evaluating whether the CNV rate correlates with the parasite background will help
to define the evolutionary potential of this successful pathogen. Despite some success with antimalarial
therapies and vaccines targeting the *Plasmodium* parasite, a strategy to impede genome evolution may
be required to control malaria infections.

Materials and Methods:

Parasite Lines, Compounds, and Treatments

We acquired *Dd2* (MRA-156) and *FCR3* (MRA-731) parasite lines from Bei Resources (ATCC, Manassas,
VA). In this study, we were interested in detecting sub-clonal levels of genomic diversity that occur
naturally in cell culture (i.e. untreated conditions); therefore, we did not re-clone parasite lines prior to
treatment. For low-input genomics, we grew parasites in complete RPMI 1640 with HEPES (Thermo
Fisher Scientific, Waltham, MA) supplemented 0.5% Albumax II Lipid-Rich BSA (MilliporeSigma,
Burlington, MA) and 50 mg/L hypoxanthine (Thermo Fisher Scientific) and donor A+ human
erythrocytes (BioIVT, Hicksville, NY). We grew all cultures at 3% hematocrit at 37°C and individually
flushed flasks with 5% oxygen, 5% carbon dioxide, and 90% nitrogen gas. We diluted cultures with
uninfected erythrocytes and changed the culture medium every other day to keep parasitemia below
2% during maintenance. We confirmed that all cultures were negative for mycoplasma contamination
approximately monthly using a LookOut Mycoplasma PCR detection kit (MilliporeSigma).

We synthesized DSM1 as in previous studies [87, 116] and purchased aphidicolin (MilliporeSigma).
DSM1 targets *P. falciparum dhodh*, which contributes to pyrimidine biosynthesis [67]. Aphidicolin
inhibits B-Family DNA polymerases, and consequently, *P. falciparum* replication [117, 118]. Both
compounds, when applied to ring-stage parasites (**Fig. 1B**), inhibit DNA replication and stall parasites at
trophozoite stage (**Fig. S3A** and **S4G**, [118]).

To assess the effects of short-term DSM1 treatment, we acquired high ring-stage cultures (>85%) by
synchronizing parasites twice with 5% sorbitol, 48hrs apart. We then applied 1μM DSM1, solvent control
(dimethylsulfoxide, DMSO), or replication inhibition control (4.4μM aphidicolin) for 12hrs. Following
treatment, we washed parasites with sterile 1x phosphate-buffered saline (PBS, Thermo Fisher
Scientific), returned them to complete RPMI, and allowed parasites to complete their life cycle and
reinvade new erythrocytes for an additional 29.5-34.5hrs (**Fig. S3B**). We tracked parasitemia and
parasite viability on an Accuri C6 flow cytometer (BD Biosciences, Franklin Lakes, NJ) as previously
performed [63, 116, 119]. We stained parasites with 1x SYBR Green (Thermo Fisher Scientific, stains the
parasite nucleus) to assess the proportion of infected erythrocytes (parasitemia) and stage of the
parasite development cycle (**Fig. 1B**) and 10nM MitoProbe DilC1 (5) (Thermo Fisher Scientific, stains
active parasite mitochondria) to indicate the proportion of the parasites that are viable over time (**Fig.
S3C**).

For parasite treatment for low-input genomics, we synchronized parasites (as above), applied 1μM
DSM1 or the DMSO control for 12hrs, and allowed recovery for 28-31hrs (**Fig. S4H**). We removed
treatments and tracked parasite number and health as described above. Following reinvasion, we
harvested viable 1n ring stage parasites for low-input genomics using flow sorting (details in *Parasite
Flow Sorting for Low-input Genomics*).

Parasite Flow Sorting for Low-input Genomics

Cell sorter calibration & accuracy assessments. We calibrated the flow sorter (SH800, Sony Biotechnology, San Jose, CA) using the manufacturer's calibration beads. We accounted for overlaps in the excitation/emission wavelengths using the integrated compensation panel matrix calculation in the SH800 software according to the manufacturer's procedure. We also manually calibrated the droplet sorting to the nearest 0.2mm, as recommended by the manufacturer, using the 96-well plate setting (Armadillo high- performance 96-well plate, Thermo Fisher Scientific). We evaluated SH800 sorting accuracy prior to low-input harvest using a colorimetric assay as previously described [120]. Briefly, we mixed SYBR Green+/MitoProbe+ parasites (see staining details in *Parasite lines, Compounds, & Treatments*) with horseradish peroxidase enzyme (Thermo Fisher Scientific) at a final concentration of 2.5mg/ml. We then sorted parasites into a 96-well plate filled with TMB-ELIZA substrate (Thermo Fisher Scientific) using the single cell (3 drops) instrument setting, in triplicate plates (**Fig. S2**). Formation of a color in the well (blue, green, or yellow) indicates the successful sorting of the enzyme, and therefore parasites, into the well with the substrate. This assessment allowed us to evaluate the accuracy of SH800 sorting (through the evaluation of success for 1- versus 2-cell wells, **Fig. S2C**), the consistency of sorting (through the evaluation of replicates), and the best plate positions for sorting (through the evaluation of performance in different plate rows/columns). Based on these evaluations, we proceeded with isolating 2- and 10-cells per well (**Fig. S2D**) and avoided sorting into the top 2 rows and the first and last column of the 96-well plate (**Figs. S2D** and **S6**).

Parasite isolation & storage. We stained parasites with SYBR Green and MitoProbe DiIC1 (5) in complete RPMI as above (see staining details in *Parasite Lines, Compounds, & Treatments*), gassed the tubes with 5% CO_2, 5% O_2, 90% N, and placed sample on ice to ensure viability prior to flow sorting within 15min (SH800, Sony Biotechnology Inc., San Jose, CA). We used a final concentration of 1 x 10^7 parasites/ml diluted in sterile 1x PBS (Thermo Fisher Scientific) as input for sorting at the "single-cell setting" (3 drop) into a 96-well plate (Armadillo high performance 96-well plate, Thermo Fisher Scientific) with each well containing 2.375µl of cell lysis buffer (0.025M Tris Ph8.8 (Roche Diagnostics, Indianapolis, IN), 0.01M NaCl (MilliporeSigma), 0.01M KCl (MilliporeSigma), 0.01M $(NH_4)_2SO_4$ (Thermo Fisher Scientific), 0.001M EDTA (Promega, Madison, WI), and 10% Triton X-100 (MilliporeSigma)). We gated viable 1n ring-stage parasites (**Fig. S5**) and sorted into the wells containing cell lysis buffer with an approximate sorting time of 10min. After sorting, we centrifuged for 30 seconds in a plate centrifuge (MPS1000, Labnet International, Madison, NJ). We immediately overlaid samples with one drop (approx. 25µl) of light mineral oil (BioReagent grade for molecular biology, MilliporeSigma) and sealed the plates with Microamp® Clear Adhesive Film (Applied Biosystems, Waltham, MA) before storage at -80°C until whole genome amplification.

MALBAC Whole Genome Amplification for Low-input Genomics

Before whole genome amplification, we thawed the plates containing sorted parasites (see *Parasite Isolation & Storage*) and added 1mg/ml Proteinase K in sterile 1x PBS (Thermo Fisher Scientific) to a final volume of 2.5µl per well. We heated the plates in a PCR cycler (C1000, Bio-Rad Laboratories, Hercules, CA) at 50°C for 3hrs, followed by 75°C for 20 min and 80°C for 5 min for proteinase k inactivation. We amplified the parasite genome using the Multiple Annealing and Looping Based Amplification Cycles (MALBAC) method essentially as previously described ([63], Version 1 in **Fig. S1**) with some modifications (Version 2, **Fig. S1**). In summary, 1) we modified the pre-amplification random primer by

adding 5 additional degenerate bases with 20% GC-content to increase annealing to AT-rich genome
sequences (5'GTGAGTGATGGTTGAGGTAGTGTGGAG<u>NNNNNNNNNNN</u>TTT 3'); 2) we performed 19 of the
21 total linear cycles with the *Bsu* DNA Polymerase (Large Fragment, New England Biosciences), which
has a lower optimal reaction temperature (37°C) to improve the amplification on AT-rich sequences
[121]; 3) we lowered the extension temperature from 40/50°C to 37°C during the linear amplification
cycles that used the *Bsu* enzyme (see full cycling parameters in **Fig. S1**); and 4) we integrated robotic
pipetting (Mosquito LV, SPT Labtech, Melbourn, UK) to increase the throughput of our assays (from 23
samples in Version 1 to 90 samples in the current Version 2) and limit contamination potential.

Overall, we performed 21 total linear cycles (19 cycles with *Bsu* polymerase and 2 cycles with *Bst*
polymerase, New England Biolabs, Ipswitch, MA) and 17 total exponential amplification cycles using
Herculase II Fusion DNA polymerase (Agilent Technologies, Santa Clara, CA). During amplification steps,
we employed standard steps to limit contamination [63]. For automated pipetting of the enzyme
solution during linear cycles, tips were changed after each round of pipetting. Post-amplification, we
purified amplified DNA with Zymo DNA Clean & Concentrator-5columns (Zymo Research, Irvine, CA)
according to the protocol and ran 2µl of all samples on 1% agarose gels to check for the presence of
DNA (generally, if >30ng/µl, samples could be visualized with a size range of 100 to >1500bp).

Assessments of Amplification Success for Low-input Genomics

DNA quantification. We quantified the MALBAC-amplified DNA using a Qubit fluorimeter (Qubit 1X
dsDNA High Sensitivity Assay Kit, Thermo Fisher Scientific).

Droplet Digital PCR. To confirm the presence of parasite DNA in MALBAC-amplified samples, we
performed droplet digital PCR (ddPCR) as described previously using the QX2000 droplet generator,
C1000 thermocycler, and QX2000 droplet reader (Bio-Rad Laboratories) [116, 122]. We used duplex
assays to evaluate two parasite genomic loci concurrently (*pfmdr1:* Forward-
TGCCCACAGAATTGCATCTA; Reverse- ACCCTGATCGAAATGGAACCT; Probe -
TCGTGTGTTCCATGTGACTG; *pfhsp70:* Forward- TGCTGTCATTACCGTTCCAG; Reverse -
AGATGCTGGTACAATTGCAGGA; Probe - AGCAGCTGCAGTAGGTTCATT (Integrated DNA
technologies, Newark, NJ). The reaction master mix contained 600nm of forward and reverse primers,
50nm probes, 10µl of ddPCR Supermix for Probes (2x, Bio-Rad Laboratories), 3µl of nuclease-free water
(QIAGEN), and 1.5ng (5µl) of template DNA per assay (total of 20µl). We used the following cycling
conditions for PCR amplification: 10 min at 95°C initial denaturation step, 1 min at 95°C second
denaturation step, and 2 min at 58°C annealing and extension step (ramp rate of 1°C per second), the
second denaturation step and the annealing/extension step repeated 60 times, and then 10 min at 98°C
to halt the reaction [122]. In addition to running amplified samples to assess amplification success, we
ran ddPCR with bulk genomic DNA as a positive control, no template controls (water replaced DNA), and
material from "no cell" wells to assess cross-well contamination. We considered the samples positive for
parasite DNA if there were more than 50 total positive droplets in target-positive clusters.

High Resolution Melting Assay. To assess potential contamination between MALBAC-amplified samples,
we performed asymmetric PCR amplification of the *pfdhps* locus followed by high-resolution melting
(HRM) as described previously [123, 124]. The *pfdhps* locus at codon 613 is distinct in *Dd2* and *FCR3*
parasite lines (*Dd2*: Ser-613 and *FCR3*: Ala-613, [125]). Each 20µL reaction contained 8µl of the 2.5x
LightScanner Master mix (BioFire^TM Defense, Salt Lake City, Utah, USA), 1/10µM of forward/reverse
primers and 8µM probes targeting the *pfdhps* gene position 613: Forward -

CTCTTACAAAATATACATGTATATGATGAGTATCCACTT;   Reverse-
CATGTAATTTTTGTTGTGTATTTATTACAACATTTTGA;   Probe - AAGATTTATTGCCCATTGCATGA/3SpC3,
(Integrated DNA technologies), 7µl of nuclease free water, and 3µl of DNA (~0.05ng total). We used the
following cycling conditions for PCR amplification with the Rotor-Gene Q instrument with a 72-well rotor
(QIAGEN): 95°C for 5 min, 45 cycles of 95°C for 10s, 55°C for 30s, and 72°C for 10s, followed by a pre-
melt at 55°C for 90s, and a HRM ramp from 65°C to 90°C, with an increase of 0.1°C every 2s. We plotted
the change in fluorescence versus temperature (dF/T) using Rotor-Gene Q software (version 2.3.5, build
1; QIAGEN) and compared HRM peaks of amplified samples to bulk genomic DNA and plasmid controls.

Bulk DNA Extraction for Short-Read Sequencing

We extracted bulk DNA for short-read sequencing as previously performed [63]. Briefly, we lysed
erythrocytes with 1.5% saponin and washed the parasite pellet 3 times with 1x PBS (Thermo Fisher
Scientific), before resuspension in a buffered solution (150mM NaCl (MilliporeSigma), 10mM EDTA
(Promega Corporation, Madison, WI), and 50mM Tris pH7.5 (Roche Diagnostics)) to a total volume of
500µl. We then lysed the parasites with 10% sarkosyl (MilliporeSigma) and 20mg/ml proteainase K
(Thermo   Fisher   Scientific)   at   37°C   overnight   before   DNA   purification   using   standard
phenol/chloroform/isoamyl alcohol extraction and chloroform washing steps (2 times each, [63]).
Finally, we precipitated DNA using 100% ethanol with 100mM of sodium acetate overnight in DNA-lo
bind tubes (Eppendorf, Enfield, CT) and then washed twice with 70% ethanol before resuspension in
50µl nuclease free water (QIAGEN). We stored bulk genomic DNA at -20°C until sequencing library
preparation.

Low-input Genomics Sample Selection & Short-Read Sequencing

Low-input sample selection. We selected 16 low-input samples from the *FCR3* plate, 36 samples from
the untreated *Dd2* plate, and 38 samples from the treated *Dd2* plate for short-read Illumina sequencing.
We based our selection on the quantity of the MALBAC amplified DNA and presence of parasite DNA
using ddPCR (**Figs. S6 and S7**). In summary, the majority of *FCR3* and untreated *Dd2* samples yielded
quantifiable parasite DNA following MALBAC amplification (53/60 *FCR3* samples and 60/60 untreated
Dd2 samples); in these conditions, we chose samples randomly to proceed with sequencing (indicated in
Fig. S6). For treated *Dd2* samples, we chose samples for sequencing if they had adequate DNA quantity
(>10ng total, 30 samples, **Fig. S6**) or had ddPCR results showing the presence of parasite DNA (an
additional 8 samples).

Short-read sequencing. Before short-read sequencing, we sheared bulk samples and low-input samples
using Covaris M220 Focused Ultrasonicator for 150s and 130s, respectively, to generate fragment sizes
of ~350bp as evaluated by an Agilent 2100 Bioanalyzer using the High Sensitivity DNA kit (Agilent, Santa
Clara). We adjusted the volume of sheared samples with nuclease-free water up to 50µl. For samples
with >100ng (**Table S2,** including bulk and MALBAC amplified samples), we diluted them to 1.2-2ng/µl;
for samples <100ng, we proceeded with no dilution. We used NEBNext Ultra II kit (Illumina Inc., San
Diego, CA) to prepare libraries for sequencing with 3 cycles of PCR amplification, as performed
previously [63]. We quantified the resulting libraries using NEBNext Library Quant Kit (Illumina Inc.)
before sequencing on the Illumina Nextseq 550 using 150 bp paired-end cycles.

Short-Read Sequence Processing & Analysis

Read processing and alignment. We performed short-read quality control steps as described previously [61, 63]. Briefly, we reordered and removed singletons and subsequently interleaved paired reads using BBMap, trimmed the MALBAC common sequence, PhiX, and Illumina adapters from the remaining reads with the BBDuk tool within BBMap, and aligned reads to the *pf3D7-62_v3* reference genome using the Speedseq genome aligner [126]. We removed reads that map to VAR regions from bam files according to previously defined genomic coordinates [77]. We filtered out reads with low mapping quality (<q30) and duplicated reads using SAMtools [127]. We employed Qualimap to report mean coverage and standard deviation across the genome [128]. Using non-overlapping 20kb size bins, we calculated the coefficient of variation of read coverage by dividing the standard deviation of coverage within a bin by the mean across a sample and multiplying by 100 [129, 130] (R version 4.4.2).

Down-sampling. For analysis that assessed down-sampled data (**Figs. 2, S10, and S11**), we first converted the processed bam files back into FASTQ files using SAMtools and then used the reformat.sh option of BBtools to select 1.3M reads from each FASTQ (represented the fewest number of reads from a sample that passed quality filtering from the final dataset). We then realigned files to the reference genome (*pf3D7-62_v3*).

Single nucleotide polymorphism analysis. We performed SNP genotyping and analysis as previously [63], based on the MalariaGen *P. falciparum* Community Project V6.0 pipeline [131-134] using the *pf3D7-62_v3* reference genome. Briefly, we applied GATK's Base Quality Score Recalibration using default settings. We detected potential SNPs using GATK's HaplotypeCaller and subsequently genotyped the SNPs using CombineGVCFs and GenotypeGVCFs. Then we employed GATK's VariantRecalibrator using previously validated SNP datasets [114]. We then applied GATK's ApplyRecalibration to assign a VQSLOD score [134]. We filtered the resulting SNPs for those with VQSLOD > 6 and for a GT quality metric >20 to ensure high-quality variant calling. We only selected variants flagged as Bi-allelic to simplify the analysis. For SNP Principle Component Analysis (PCA), we merged experiment-wide SNP data (described above) into a single file. Then we merged the VCF into a large matrix and converted the genomic data into numeric information using the 'vcfr' package in R (Version 4.2.3) (https://CRAN.R-project.org/package=vcfR). We excluded individual SNPs if >25% of the samples lacked a call in this position or if all calls were the same for each sample in that position). We scaled the remaining SNPs around the origin using the 'scale' R function. We then calculated the principal components using the 'prcomp' R function, and scored the dataset using the 'scores' function from the 'vegan' R package (https://CRAN.R-project.org/package=vegan).

Copy Number Variation Analysis

CNV calling in bulk samples. We performed CNV detection for bulk samples similar to as previously described [61, 63]. Briefly, we called CNVs independently using two methods, CNVnator (read depth based calling, [135]) and LUMPY (split and discordant read based calling, [74]). To Identify CNVs called in both methods, we used SVCROWS to define overlapping CNV regions relative to their size. Briefly, SVCROWS uses a reciprocal-overlap-based approach (i.e. two CNVs must be overlapping each other at, or greater than, a defined threshold) to determine if two CNVs are close enough in their genomic position to be called the same. The source code for SVCROWS can be accessed at https://github.com/A-Crow-Nowhere/SVCROWS.git. The program utilizes different thresholds for overlap based on the sizes of the CNVs being compared, ensuring that we account for shifts in CNV position. For known CNV calls, we

used the following SVCROWs input parameters: ExpandRORegion = FALSE, BPfactor = TRUE, DefaultSizes
= FALSE, xs = 3000, xl = 10,000, y1s = 300, y1l = 1000, y2s = 50% and y2l = 80%; based on the average size
of a *P. falciparum* gene (~2.3kb) and intergenic region (~2kb). Similar to our previous study [63], we
identified 3 known CNVs that were called by both LUMPY and CNVnator methods in the core genome of
bulk samples (untreated and treated). We determined known CNV boundaries using SVCROWS: *pfmdr1*
(*Pf3D7_05_v3*, 888001-970000, 82kb), *pf11-1* (*Pf3D7_10_v3*, 1521345-1541576, 20kb), and *pf332*
(*Pf3D7_11_v3*, 1950201-1962400, 12kb).

CNV calling in 2-cell samples. We employed two methods for CNV detection in the core genome of low-
input samples; LUMPY is a split/discordant read strategy with high sensitivity [74], and HapCNV is a read
coverage-based strategy designed for haploid genomes [73]. We ran LUMPY as part of Speedseq with
default parameters as previously described [61]. We filtered resulting structural variants to include only
duplications (DUP, >1 copy of a region) and deletions (DEL, one less copy of region than the reference).
We then filtered those calls for those GQ > 20 to ensure high-fidelity calls. In HapCNV, we used a quality
control and bias correction procedure to exclude bins of poor quality and remove bias introduced by GC
content and mappability variation. We then constructed a pseudo-reference for each *Dd2* low-input
sample using within-*Dd2* information, which enabled control of background noise while preserving CNV
signals after normalization. Finally, we used a circular binary segmentation algorithm (CBS, [136]) to
detect copy number change points followed by a Gaussian Mixture Model (GMM, [137]) for CNV
identification. The source code for HapCNV and examples of real data application can be accessed at
https://github.com/FeifeiXiao-lab/HapCNV. For statistics, we used PRISM (GraphPad Software, La Jolla,
CA), using unpaired parametric T-tests with Welch's correction.

Defining CNV regions/determination of "rarity" in CNV calling. Small differences in sequence quality
surrounding a read can lead to shifted breakpoint determination for biologically identical CNVs, which is
especially true for low-input genomics datasets [38]. To account for this, we used SVCROWS to
determine whether two CNV signals were the same within and between samples (see *CNV calling in bulk*
samples for input parameters). We assigned the categorizations of "rare" and "common" by assessing
the CNV region frequency within datasets. "Rare" CNVs were defined as occurring in <10% of the
samples within a treatment group; "common" CNVs were defined as occurring in ≥10% of samples
within a treatment group; "known" CNVs were defined by CNVs called in bulk samples (see above, *CNV*
calling in bulk samples).

High-confidence CNV region identification. To identify "high-confidence" CNV regions called by both
HapCNV and LUMPY methods, we compared the 'consensus list' generated by SVCROWS for each
detection method by combining them into a single SVCROWS input file. Because 1) HapCNV generates
imprecise breakpoints, and 2) there is a large disparity of average CNV region lengths between the two
methods (HapCNV = ~40kb, LUMPY = ~4.3kb), we relaxed the stringency of the SVCROWS parameters.
Our input parameters to generate the "high-confidence" list were as follows: xs = 3000, xl = 6000, y1s =
500, y1l = 1500, y2s = 30% and y2l = 60%. We defined "high-confidence" CNV regions as those that had
>1 match from both HapCNV and LUMPY that was the same type (i.e. either duplication or deletion or
mixed). For Venn diagram generation, we calculated overlaps using SVCROWS "Scavenge" mode (input
parameters: ExpandRORegion = FALSE, BPfactor = TRUE, DefaultSizes = FALSE, xs = 3000, xl = 6000, y1s =
500, y1l = 1500, y2s = 30% and y2l = 60%). We systematically compared lists for each overlap
comparison, and if regions had at least one match in an opposing dataset, we considered it a match. We
used the *draw.quad.venn* function in the "VennDiagram" R package (R 4.2.1) to generate the diagram.

Gene Ontology enrichment and protein class identification. We used the online Gene Ontology Resource (geneontology.org) to perform GO enrichment analysis using the PANTHER Classification System [138, 139]. Since a large portion of the *P. falciparum* genome remains unannotated (PlasmoDB, 30%) and the majority of molecular functions remain unclassified (92.8%), we used the Panther Protein class assessment (version 19.0, only 55.9% remained unclassified) with default statistics (Fisher's test with FDR adjusted p value of <0.05 for significance, which is recommended for small counts and overlaps between classes). We used the web tool to represent protein classes on pie charts.

Comparison to clinical CNV dataset. To determine genes covered by de novo CNVs from HapCNV and LUMPY, we used SVCROWS "Hunt" mode (input parameters: BPfactor = TRUE, DefaultSizes = FALSE, xs = 3000, xl = 6000, y1s = 300, y1l = 600, y2s = 30, y2l = 60), which takes a secondary input list of known genes (Pf3D7_62_v3, Plasmodb.org) against which to compare CNV regions. For the clinical dataset, we used genes covered by >300bp CNV regions present in high frequency in clinical isolates (Supplementary Table 3 from [59]), and manually reformatted to match SVCROWS input style guidelines before rerunning as above.

Acknowledgments:

We would like to thank Dr. John Campbell for use of the Mosquito LV instrument and assay reagents for the colorimetric assessment of the SH800. We would also like to thank Dr. Ali Guler for statistical consultation and Dr. Heidi Seears from the University of Virginia, Department of Biology Genomics Core Facility for sequencing assistance. This work was funded by 1R01AI150856-01A1 (to JLG).

References:

1. Hastings, P.J., J.R. Lupski, S.M. Rosenberg, and G. Ira, *Mechanisms of change in gene copy number.* Nat Rev Genet, 2009. 10(8): p. 551-64.
2. Lauer, S. and D. Gresham, *An evolving view of copy number variants.* Curr Genet, 2019.
3. Pos, O., J. Radvanszky, G. Buglyo, Z. Pos, D. Rusnakova, B. Nagy, and T. Szemes, *DNA copy number variation: Main characteristics, evolutionary significance, and pathological aspects.* Biomed J, 2021. 44(5): p. 548-559.
4. Roth, J.R., *The joys and terrors of fast adaptation: new findings elucidate antibiotic resistance and natural selection.* Mol Microbiol, 2011. 79(2): p. 279-82.
5. Payen, C., S.C. Di Rienzi, G.T. Ong, J.L. Pogachar, J.C. Sanchez, A.B. Sunshine, M.K. Raghuraman, B.J. Brewer, and M.J. Dunham, *The dynamics of diverse segmental amplifications in populations of Saccharomyces cerevisiae adapting to strong selection.* G3 (Bethesda), 2014. 4(3): p. 399-409.
6. Elde, N.C., S.J. Child, M.T. Eickbush, J.O. Kitzman, K.S. Rogers, J. Shendure, A.P. Geballe, and H.S. Malik, *Poxviruses deploy genomic accordions to adapt rapidly against host antiviral defenses.* Cell, 2012. 150(4): p. 831-41.
7. Leary, R.J., J.C. Lin, J. Cummins, S. Boca, L.D. Wood, D.W. Parsons, S. Jones, T. Sjoblom, B.H. Park, R. Parsons, J. Willis, D. Dawson, J.K. Willson, T. Nikolskaya, Y. Nikolsky, L. Kopelovich, N. Papadopoulos, L.A. Pennacchio, T.L. Wang, S.D. Markowitz, G. Parmigiani, K.W. Kinzler, B. Vogelstein, and V.E. Velculescu, *Integrated analysis of homozygous deletions, focal amplifications, and sequence alterations in breast and colorectal cancers.* Proc Natl Acad Sci U S A, 2008. 105(42): p. 16224-9.

8.   Beroukhim, R., C.H. Mermel, D. Porter, G. Wei, S. Raychaudhuri, J. Donovan, J. Barretina, J.S.
Boehm, J. Dobson, M. Urashima, K.T. Mc Henry, R.M. Pinchback, A.H. Ligon, Y.J. Cho, L. Haery, H.
Greulich, M. Reich, W. Winckler, M.S. Lawrence, B.A. Weir, K.E. Tanaka, D.Y. Chiang, A.J. Bass, A.
Loo, C. Hoffman, J. Prensner, T. Liefeld, Q. Gao, D. Yecies, S. Signoretti, E. Maher, F.J. Kaye, H.
Sasaki, J.E. Tepper, J.A. Fletcher, J. Tabernero, J. Baselga, M.S. Tsao, F. Demichelis, M.A. Rubin,
P.A. Janne, M.J. Daly, C. Nucera, R.L. Levine, B.L. Ebert, S. Gabriel, A.K. Rustgi, C.R. Antonescu, M.
Ladanyi, A. Letai, L.A. Garraway, M. Loda, D.G. Beer, L.D. True, A. Okamoto, S.L. Pomeroy, S.
Singer, T.R. Golub, E.S. Lander, G. Getz, W.R. Sellers, and M. Meyerson, *The landscape of*
somatic copy-number alteration across human cancers. Nature, 2010. 463(7283): p. 899-905.
9.   Zhang, L., Y. Yuan, K.H. Lu, and L. Zhang, *Identification of recurrent focal copy number variations*
and their putative targeted driver genes in ovarian cancer. BMC Bioinformatics, 2016. 17(1): p.
222.
10.   Peng, H., L. Lu, Z. Zhou, J. Liu, D. Zhang, K. Nan, X. Zhao, F. Li, L. Tian, H. Dong, and Y. Yao, *CNV*
Detection from Circulating Tumor DNA in Late Stage Non-Small Cell Lung Cancer Patients. Genes
(Basel), 2019. 10(11).
11.   Krijgsman, O., B. Carvalho, G.A. Meijer, R.D. Steenbergen, and B. Ylstra, *Focal chromosomal copy*
number aberrations in cancer-Needles in a genome haystack. Biochim Biophys Acta, 2014.
1843(11): p. 2698-2704.
12.   Carey-Smith, S.L., R.S. Kotecha, L.C. Cheung, and S. Malinge, *Insights into the Clinical, Biological*
and Therapeutic Impact of Copy Number Alteration in Cancer. Int J Mol Sci, 2024. 25(13).
13.   Gonzalez, E., H. Kulkarni, H. Bolivar, A. Mangano, R. Sanchez, G. Catano, R.J. Nibbs, B.I.
Freedman, M.P. Quinones, M.J. Bamshad, K.K. Murthy, B.H. Rovin, W. Bradley, R.A. Clark, S.A.
Anderson, J. O'Connell R, B.K. Agan, S.S. Ahuja, R. Bologna, L. Sen, M.J. Dolan, and S.K. Ahuja,
The influence of CCL3L1 gene-containing segmental duplications on HIV-1/AIDS susceptibility.
Science, 2005. 307(5714): p. 1434-40.
14.   Sebat, J., B. Lakshmi, D. Malhotra, J. Troge, C. Lese-Martin, T. Walsh, B. Yamrom, S. Yoon, A.
Krasnitz, J. Kendall, A. Leotta, D. Pai, R. Zhang, Y.H. Lee, J. Hicks, S.J. Spence, A.T. Lee, K. Puura, T.
Lehtimaki, D. Ledbetter, P.K. Gregersen, J. Bregman, J.S. Sutcliffe, V. Jobanputra, W. Chung, D.
Warburton, M.C. King, D. Skuse, D.H. Geschwind, T.C. Gilliam, K. Ye, and M. Wigler, *Strong*
association of de novo copy number mutations with autism. Science, 2007. 316(5823): p. 445-9.
15.   Glessner, J.T., J.P. Bradfield, K. Wang, N. Takahashi, H. Zhang, P.M. Sleiman, F.D. Mentch, C.E.
Kim, C. Hou, K.A. Thomas, M.L. Garris, S. Deliard, E.C. Frackelton, F.G. Otieno, J. Zhao, R.M.
Chiavacci, M. Li, J.D. Buxbaum, R.I. Berkowitz, H. Hakonarson, and S.F. Grant, *A genome-wide*
study reveals copy number variants exclusive to childhood obesity cases. Am J Hum Genet, 2010.
87(5): p. 661-6.
16.   Olsson, L.M. and R. Holmdahl, *Copy number variation in autoimmunity--importance hidden in*
complexity? Eur J Immunol, 2012. 42(8): p. 1969-76.
17.   Butchbach, M.E., *Copy Number Variations in the Survival Motor Neuron Genes: Implications for*
Spinal Muscular Atrophy and Other Neurodegenerative Diseases. Front Mol Biosci, 2016. 3: p. 7.
18.   Sekar, A., A.R. Bialas, H. de Rivera, A. Davis, T.R. Hammond, N. Kamitaki, K. Tooley, J. Presumey,
M. Baum, V. Van Doren, G. Genovese, S.A. Rose, R.E. Handsaker, C. Schizophrenia Working
Group of the Psychiatric Genomics, M.J. Daly, M.C. Carroll, B. Stevens, and S.A. McCarroll,
Schizophrenia risk from complex variation of complement component 4. Nature, 2016.
530(7589): p. 177-83.
19.   Zekavat, S.M., S. Ruotsalainen, R.E. Handsaker, M. Alver, J. Bloom, T. Poterba, C. Seed, J. Ernst,
M. Chaffin, J. Engreitz, G.M. Peloso, A. Manichaikul, C. Yang, K.A. Ryan, M. Fu, W.C. Johnson, M.
Tsai, M. Budoff, R.S. Vasan, L.A. Cupples, J.I. Rotter, S.S. Rich, W. Post, B.D. Mitchell, A. Correa,
A. Metspalu, J.G. Wilson, V. Salomaa, M. Kellis, M.J. Daly, B.M. Neale, S. McCarroll, I. Surakka, T.

Esko, A. Ganna, S. Ripatti, S. Kathiresan, P. Natarajan, and N.T.L.W. Group, *Deep coverage whole*
genome sequences and plasma lipoprotein(a) in individuals of European and African ancestries.
Nat Commun, 2018. 9(1): p. 2606.

20.   Farashi, S. and C.L. Harteveld, *Molecular basis of alpha-thalassemia.* Blood Cells Mol Dis, 2018.
70: p. 43-53.

21.   Wisniowiecka-Kowalnik, B. and B.A. Nowakowska, *Genetics and epigenetics of autism spectrum*
disorder-current evidence in the field. J Appl Genet, 2019. 60(1): p. 37-47.

22.   Craven, S.H. and E.L. Neidle, *Double trouble: medical implications of genetic duplication and*
amplification in bacteria. Future Microbiol, 2007. 2(3): p. 309-21.

23.   Sandegren, L. and D.I. Andersson, *Bacterial gene amplification: implications for the evolution of*
antibiotic resistance. Nat Rev Microbiol, 2009. 7(8): p. 578-88.

24.   Hjort, K., H. Nicoloff, and D.I. Andersson, *Unstable tandem gene amplification generates*
heteroresistance (variation in resistance within a population) to colistin in Salmonella enterica.
Mol Microbiol, 2016. 102(2): p. 274-289.

25.   Anderson, S.E., E.X. Sherman, D.S. Weiss, and P.N. Rather, *Aminoglycoside Heteroresistance in*
Acinetobacter baumannii AB5075. mSphere, 2018. 3(4).

26.   Nicoloff, H., K. Hjort, B.R. Levin, and D.I. Andersson, *The high prevalence of antibiotic*
heteroresistance in pathogenic bacteria is mainly caused by gene amplification. Nat Microbiol,
2019. 4(3): p. 504-514.

27.   Baslan, T., J. Kendall, K. Volyanskyy, K. McNamara, H. Cox, S. D'Italia, F. Ambrosio, M. Riggs, L.
Rodgers, A. Leotta, J. Song, Y. Mao, J. Wu, R. Shah, R. Gularte-Merida, K. Chadalavada, G.
Nanjangud, V. Varadan, A. Gordon, C. Curtis, A. Krasnitz, N. Dimitrova, L. Harris, M. Wigler, and J.
Hicks, *Novel insights into breast cancer copy number genetic heterogeneity revealed by single-*
cell genome sequencing. Elife, 2020. 9.

28.   van Dijk, E., T. van den Bosch, K.J. Lenos, K. El Makrini, L.E. Nijman, H.F.B. van Essen, N. Lansu,
M. Boekhout, J.H. Hageman, R.C. Fitzgerald, C.J.A. Punt, J.B. Tuynman, H.J.G. Snippert, G. Kops,
J.P. Medema, B. Ylstra, L. Vermeulen, and D.M. Miedema, *Chromosomal copy number*
heterogeneity predicts survival rates across cancers. Nat Commun, 2021. 12(1): p. 3188.

29.   Sobral, D., M. Martins, S. Kaplan, M. Golkaram, M. Salmans, N. Khan, R. Vijayaraghavan, S.
Casimiro, A. Fernandes, P. Borralho, C. Ferreira, R. Pinto, C. Abreu, A.L. Costa, S. Zhang, T.
Pawlowski, J. Godsey, A. Mansinho, D. Macedo, S. Lobo-Martins, P. Filipe, R. Esteves, J.
Coutinho, P.M. Costa, A. Ramires, F. Aldeia, A. Quintela, A. So, L. Liu, A.R. Grosso, and L. Costa,
Genetic and microenvironmental intra-tumor heterogeneity impacts colorectal cancer evolution
and metastatic development. Commun Biol, 2022. 5(1): p. 937.

30.   Arlt, M.F., J.G. Mulle, V.M. Schaibley, R.L. Ragland, S.G. Durkin, S.T. Warren, and T.W. Glover,
Replication stress induces genome-wide copy number changes in human cells that resemble
polymorphic and pathogenic variants. Am J Hum Genet, 2009. 84(3): p. 339-50.

31.   Arlt, M.F., A.C. Ozdemir, S.R. Birkeland, T.E. Wilson, and T.W. Glover, *Hydroxyurea induces de*
novo copy number variants in human cells. Proc Natl Acad Sci U S A, 2011. 108(42): p. 17360-5.

32.   Arlt, M.F., S. Rajendran, S.R. Birkeland, T.E. Wilson, and T.W. Glover, *De novo CNV formation in*
mouse embryonic stem cells occurs in the absence of Xrcc4-dependent nonhomologous end
joining. PLoS Genet, 2012. 8(9): p. e1002981.

33.   Lauer, S., G. Avecilla, P. Spealman, G. Sethia, N. Brandt, S.F. Levy, and D. Gresham, *Single-cell*
copy number variant detection reveals the dynamics and diversity of adaptation. PLoS Biol, 2018.
16(12): p. e3000069.

34.   Oketch, D.J.A., M. Giulietti, and F. Piva, *Copy Number Variations in Pancreatic Cancer: From*
Biological Significance to Clinical Utility. Int J Mol Sci, 2023. 25(1).

35.   Yi, D., J.W. Nam, and H. Jeong, *Toward the functional interpretation of somatic structural*
889   *variations: bulk- and single-cell approaches.* Brief Bioinform, 2023. 24(5).
890   36.   Macaulay, I.C. and T. Voet, *Single cell genomics: advances and future perspectives.* PLoS Genet,
891   2014. 10(1): p. e1004126.
892   37.   Wang, Y. and N.E. Navin, *Advances and applications of single-cell sequencing technologies.* Mol
893   Cell, 2015. 58(4): p. 598-609.
894   38.   Gawad, C., W. Koh, and S.R. Quake, *Single-cell genome sequencing: current state of the science.*
895   Nat Rev Genet, 2016. 17(3): p. 175-88.
896   39.   Wang, X., H. Chen, and N.R. Zhang, *DNA copy number profiling using single-cell sequencing.* Brief
897   Bioinform, 2018. 19(5): p. 731-736.
898   40.   Patel, A.P., I. Tirosh, J.J. Trombetta, A.K. Shalek, S.M. Gillespie, H. Wakimoto, D.P. Cahill, B.V.
899   Nahed, W.T. Curry, R.L. Martuza, D.N. Louis, O. Rozenblatt-Rosen, M.L. Suvà, A. Regev, and B.E.
900   Bernstein, *Single-cell RNA-seq highlights intratumoral heterogeneity in primary glioblastoma.*
901   Science, 2014. 344(6190): p. 1396-1401.
902   41.   Xu, Q.H., S.H. Chen, Y.B. Hu, and W. Huang, *Single-cell RNA transcriptome reveals the intra-*
903   *tumoral heterogeneity and regulators underlying tumor progression in metastatic pancreatic*
904   *ductal adenocarcinoma.* Cell Death Discovery, 2021. 7(1).
905   42.   Gao, R.L., S.S. Bai, Y.C. Henderson, Y.Y. Lin, A. Schalck, Y. Yan, T. Kumar, M. Hu, E. Sei, A. Davis, F.
906   Wang, S.F. Shaitelman, J.R. Wang, K. Chen, S. Moulder, S.Y. Lai, and N.E. Navin, *Delineating copy*
907   *number and clonal substructure in human tumors from single-cell transcriptomes.* Nature
908   Biotechnology, 2021. 39(5): p. 599-608.
909   43.   Mahdipour-Shirayeh, A., N. Erdmann, C. Leung-Hagesteijn, and R.E. Tiedemann, *sciCNV: high-*
910   *throughput paired profiling of transcriptomes and DNA copy number variations at single-cell*
911   *resolution.* Briefings in Bioinformatics, 2022. 23(1).
912   44.   Navin, N., J. Kendall, J. Troge, P. Andrews, L. Rodgers, J. McIndoo, K. Cook, A. Stepansky, D. Levy,
913   D. Esposito, L. Muthuswamy, A. Krasnitz, W.R. McCombie, J. Hicks, and M. Wigler, *Tumour*
914   *evolution inferred by single-cell sequencing.* Nature, 2011. 472(7341): p. 90-4.
915   45.   McConnell, M.J., M.R. Lindberg, K.J. Brennand, J.C. Piper, T. Voet, C. Cowing-Zitron, S. Shumilina,
916   R.S. Lasken, J.R. Vermeesch, I.M. Hall, and F.H. Gage, *Mosaic copy number variation in human*
917   *neurons.* Science, 2013. 342(6158): p. 632-7.
918   46.   Cai, X., G.D. Evrony, H.S. Lehmann, P.C. Elhosary, B.K. Mehta, A. Poduri, and C.A. Walsh, *Single-*
919   *cell, genome-wide sequencing identifies clonal somatic copy-number variation in the human*
920   *brain.* Cell Rep, 2014. 8(5): p. 1280-9.
921   47.   Francis, J.M., C.Z. Zhang, C.L. Maire, J. Jung, V.E. Manzo, V.A. Adalsteinsson, H. Homer, S. Haidar,
922   B. Blumenstiel, C.S. Pedamallu, A.H. Ligon, J.C. Love, M. Meyerson, and K.L. Ligon, *EGFR variant*
923   *heterogeneity in glioblastoma resolved through single-nucleus sequencing.* Cancer Discov, 2014.
924   4(8): p. 956-71.
925   48.   Eirew, P., A. Steif, J. Khattra, G. Ha, D. Yap, H. Farahani, K. Gelmon, S. Chia, C. Mar, A. Wan, E.
926   Laks, J. Biele, K. Shumansky, J. Rosner, A. McPherson, C. Nielsen, A.J. Roth, C. Lefebvre, A.
927   Bashashati, C. de Souza, C. Siu, R. Aniba, J. Brimhall, A. Oloumi, T. Osako, A. Bruna, J.L. Sandoval,
928   T. Algara, W. Greenwood, K. Leung, H. Cheng, H. Xue, Y. Wang, D. Lin, A.J. Mungall, R. Moore, Y.
929   Zhao, J. Lorette, L. Nguyen, D. Huntsman, C.J. Eaves, C. Hansen, M.A. Marra, C. Caldas, S.P. Shah,
930   and S. Aparicio, *Dynamics of genomic clones in breast cancer patient xenografts at single-cell*
931   *resolution.* Nature, 2015. 518(7539): p. 422-6.
932   49.   Rohrback, S., C. April, F. Kaper, R.R. Rivera, C.S. Liu, B. Siddoway, and J. Chun, *Submegabase copy*
933   *number variations arise during cerebral cortical neurogenesis as revealed by single-cell whole-*
934   *genome sequencing.* Proc Natl Acad Sci U S A, 2018. 115(42): p. 10804-10809.

50. Laks, E., A. McPherson, H. Zahn, D. Lai, A. Steif, J. Brimhall, J. Biele, B. Wang, T. Masud, J. Ting, D.
Grewal, C. Nielsen, S. Leung, V. Bojilova, M. Smith, O. Golovko, S. Poon, P. Eirew, F. Kabeer, T.
Ruiz de Algara, S.R. Lee, M.J. Taghiyar, C. Huebner, J. Ngo, T. Chan, S. Vatrt-Watts, P. Walters, N.
Abrar, S. Chan, M. Wiens, L. Martin, R.W. Scott, T.M. Underhill, E. Chavez, C. Steidl, D. Da Costa,
Y. Ma, R.J.N. Coope, R. Corbett, S. Pleasance, R. Moore, A.J. Mungall, C. Mar, F. Cafferty, K.
Gelmon, S. Chia, C.I.G.C. Team, M.A. Marra, C. Hansen, S.P. Shah, and S. Aparicio, *Clonal
Decomposition and DNA Replication States Defined by Scaled Single-Cell Genome Sequencing.*
Cell, 2019. 179(5): p. 1207-1221 e22.
51. Kidgell, C., S.K. Volkman, J. Daily, J.O. Borevitz, D. Plouffe, Y. Zhou, J.R. Johnson, K. Le Roch, O.
Sarr, O. Ndir, S. Mboup, S. Batalov, D.F. Wirth, and E.A. Winzeler, *A systematic map of genetic
variation in Plasmodium falciparum.* PLoS Pathog, 2006. 2(6): p. e57.
52. Ribacke, U., B.W. Mok, V. Wirta, J. Normark, J. Lundeberg, F. Kironde, T.G. Egwang, P. Nilsson,
and M. Wahlgren, *Genome wide gene amplifications and deletions in Plasmodium falciparum.*
Molecular and Biochemical Parasitology, 2007. 155(1): p. 33-44.
53. Cheeseman, I., N. Gomez-Escobar, C. Carret, A. Ivens, L. Stewart, K. Tetteh, and D. Conway, *Gene
copy number variation throughout the Plasmodium falciparum genome.* BMC Genomics, 2009.
10(1): p. 353.
54. Bopp, S.E., M.J. Manary, A.T. Bright, G.L. Johnston, N.V. Dharia, F.L. Luna, S. McCormack, D.
Plouffe, C.W. McNamara, J.R. Walker, D.A. Fidock, E.L. Denchi, and E.A. Winzeler, *Mitotic
evolution of Plasmodium falciparum shows a stable core genome but recombination in antigen
families.* PLoS Genet, 2013. 9(2): p. e1003293.
55. Gendrot, M., R. Fawaz, J. Dormoi, M. Madamet, and B. Pradines, *Genetic diversity and deletion
of Plasmodium falciparum histidine-rich protein 2 and 3: a threat to diagnosis of P. falciparum
malaria.* Clin Microbiol Infect, 2019. 25(5): p. 580-585.
56. Matesanz, F., M. Tellez, and A. Alcina, *The Plasmodium falciparum fatty acyl-CoA synthetase
family (PfACS) and differential stage-specific expression in infected erythrocytes.* Mol Biochem
Parasitol, 2003. 126(1): p. 109-12.
57. Dankwa, S., C. Lim, A.K. Bei, R.H. Jiang, J.R. Abshire, S.D. Patel, J.M. Goldberg, Y. Moreno, M.
Kono, J.C. Niles, and M.T. Duraisingh, *Ancient human sialic acid variant restricts an emerging
zoonotic malaria parasite.* Nat Commun, 2016. 7: p. 11187.
58. Price, R.N., A.-C. Uhlemann, A. Brockman, R. McGready, E. Ashley, L. Phaipun, R. Patel, K. Laing,
S. Looareesuwan, N.J. White, F. Nosten, and S. Krishna, *Mefloquine resistance in Plasmodium
falciparum and increased pfmdr1 gene copy number.* The Lancet, 2004. 364(9432): p. 438-447.
59. Ravenhall, M., E.D. Benavente, C.J. Sutherland, D.A. Baker, S. Campino, and T.G. Clark, *An
analysis of large structural variation in global Plasmodium falciparum isolates identifies a novel
duplication of the chloroquine resistance associated gene.* Sci Rep, 2019. 9(1): p. 8287.
60. Heinberg, A., E. Siu, C. Stern, E.A. Lawrence, M.T. Ferdig, K.W. Deitsch, and L.A. Kirkman, *Direct
evidence for the adaptive role of copy number variation on antifolate susceptibility in
Plasmodium falciparum.* Mol Microbiol, 2013. 88(4): p. 702-12.
61. Huckaby, A.C., C.S. Granum, M.A. Carey, K. Szlachta, B. Al-Barghouthi, Y.H. Wang, and J.L. Guler,
*Complex DNA structures trigger copy number variation across the Plasmodium falciparum
genome.* Nucleic Acids Res, 2019. 47(4): p. 1615-1627.
62. Gardner, M.J., N. Hall, E. Fung, O. White, M. Berriman, R.W. Hyman, J.M. Carlton, A. Pain, K.E.
Nelson, S. Bowman, I.T. Paulsen, K. James, J.A. Eisen, K. Rutherford, S.L. Salzberg, A. Craig, S.
Kyes, M.-S. Chan, V. Nene, S.J. Shallom, B. Suh, J. Peterson, S. Angiuoli, M. Pertea, J. Allen, J.
Selengut, D. Haft, M.W. Mather, A.B. Vaidya, D.M.A. Martin, A.H. Fairlamb, M.J. Fraunholz, D.S.
Roos, S.A. Ralph, G.I. McFadden, L.M. Cummings, G.M. Subramanian, C. Mungall, J.C. Venter,

D.J. Carucci, S.L. Hoffman, C. Newbold, R.W. Davis, C.M. Fraser, and B. Barrell, *Genome sequence*
of the human malaria parasite Plasmodium falciparum. Nature, 2002. 419(6906): p. 498-511.

63.   Liu, S., A.C. Huckaby, A.C. Brown, C.C. Moore, I. Burbulis, M.J. McConnell, and J.L. Guler, *Single-*
cell sequencing of the small and AT-skewed genome of malaria parasites. Genome Med, 2021.
13(1): p. 75.

64.   Galhardo, R.S., P.J. Hastings, and S.M. Rosenberg, *Mutation as a stress response and the*
regulation of evolvability. Crit Rev Biochem Mol Biol, 2007. 42(5): p. 399-435.

65.   Arlt, M.F., T.E. Wilson, and T.W. Glover, *Replication stress and mechanisms of CNV formation.*
Curr Opin Genet Dev, 2012. 22(3): p. 204-10.

66.   Glover, T.W., T.E. Wilson, and M.F. Arlt, *Fragile sites in cancer: more than meets the eye.* Nat Rev
Cancer, 2017. 17(8): p. 489-501.

67.   Phillips, M.A., R. Gujjar, N.A. Malmquist, J. White, F. El Mazouni, J. Baldwin, and P.K. Rathod,
Triazolopyrimidine-Based Dihydroorotate Dehydrogenase Inhibitors with Potent and Selective
Activity against the Malaria Parasite Plasmodium falciparum. Journal of Medicinal Chemistry,
2008. 51(12): p. 3649-3653.

68.   Sanz, L.M., B. Crespo, C. De-Cozar, X.C. Ding, J.L. Llergo, J.N. Burrows, J.F. Garcia-Bustos, and F.J.
Gamo, *P. falciparum in vitro killing rates allow to discriminate between different antimalarial*
mode-of-action. PLoS One, 2012. 7(2): p. e30949.

69.   Jensen, J.B. and W. Trager, *Plasmodium falciparum in culture: establishment of additional*
strains. Am J Trop Med Hyg, 1978. 27(4): p. 743-6.

70.   Dolan, S.A., L.H. Miller, and T.E. Wellems, *Evidence for a switching mechanism in the invasion of*
erythrocytes by Plasmodium falciparum. J Clin Invest, 1990. 86(2): p. 618-24.

71.   Tian, S., H. Yan, C. Neuhauser, and S.L. Slager, *An analytical workflow for accurate variant*
discovery in highly divergent regions. BMC Genomics, 2016. 17(1): p. 703.

72.   Sanderson, N.D., J. Swann, L. Barker, J. Kavanagh, S. Hoosdally, D. Crook, G. GonFast
Investigators, T.L. Street, and D.W. Eyre, *High precision Neisseria gonorrhoeae variant and*
antimicrobial resistance calling from metagenomic Nanopore sequencing. Genome Res, 2020.
30(9): p. 1354-1363.

73.   Yu, X., F. Qin, S. Liu, N.J. Brown, Q. Lu, G. Cai, J.L. Guler, and F. Xiao, *HapCNV: A Comprehensive*
Framework for CNV Detection in Low-input DNA Sequencing Data. bioRxiv, 2024: p.
2024.12.19.629494.

74.   Layer, R.M., C. Chiang, A.R. Quinlan, and I.M. Hall, *LUMPY: a probabilistic framework for*
structural variant discovery. Genome Biol, 2014. 15(6): p. R84.

75.   Cameron, D.L., L. Di Stefano, and A.T. Papenfuss, *Comprehensive evaluation and*
characterisation of short read general-purpose structural variant calling software. Nat Commun,
2019. 10(1): p. 3240.

76.   Gabrielaite, M., M.H. Torp, M.S. Rasmussen, S. Andreu-Sanchez, F.G. Vieira, C.B. Pedersen, S.
Kinalis, M.B. Madsen, M. Kodama, G.S. Demircan, A. Simonyan, C.W. Yde, L.R. Olsen, R.L. Marvig,
O. Ostrup, M. Rossing, F.C. Nielsen, O. Winther, and F.O. Bagger, *A Comparison of Tools for*
Copy-Number Variation Detection in Germline Whole Exome and Whole Genome Sequencing
Data. Cancers (Basel), 2021. 13(24).

77.   Otto, T.D., U. Bohme, M. Sanders, A. Reid, E.I. Bruske, C.W. Duffy, P.C. Bull, R.D. Pearson, A.
Abdi, S. Dimonte, L.B. Stewart, S. Campino, M. Kekre, W.L. Hamilton, A. Claessens, S.K. Volkman,
D. Ndiaye, A. Amambua-Ngwa, M. Diakite, R.M. Fairhurst, D.J. Conway, M. Franck, C.I. Newbold,
and M. Berriman, *Long read assemblies of geographically dispersed Plasmodium falciparum*
isolates reveal highly structured subtelomeres. Wellcome Open Res, 2018. 3: p. 52.

78.   Saxena, S. and L. Zou, *Hallmarks of DNA replication stress.* Mol Cell, 2022. 82(12): p. 2298-2314.

79.   Guler, J.L., D.L. Freeman, V. Ahyong, R. Patrapuvich, J. White, R. Gujjar, M.A. Phillips, J. DeRisi,
and P.K. Rathod, *Asexual populations of the human malaria parasite, Plasmodium falciparum,*
use a two-step genomic strategy to acquire accurate, beneficial DNA amplifications. PLoS
Pathog, 2013. 9(5): p. e1003375.
80.   Evrony, G.D., E. Lee, P.J. Park, and C.A. Walsh, *Resolving rates of mutation in the brain using*
single-neuron genomics. Elife, 2016. 5.
81.   Mills, R.E., K. Walter, C. Stewart, R.E. Handsaker, K. Chen, C. Alkan, A. Abyzov, S.C. Yoon, K. Ye,
R.K. Cheetham, A. Chinwalla, D.F. Conrad, Y. Fu, F. Grubert, I. Hajirasouliha, F. Hormozdiari, L.M.
Iakoucheva, Z. Iqbal, S. Kang, J.M. Kidd, M.K. Konkel, J. Korn, E. Khurana, D. Kural, H.Y. Lam, J.
Leng, R. Li, Y. Li, C.Y. Lin, R. Luo, X.J. Mu, J. Nemesh, H.E. Peckham, T. Rausch, A. Scally, X. Shi,
M.P. Stromberg, A.M. Stutz, A.E. Urban, J.A. Walker, J. Wu, Y. Zhang, Z.D. Zhang, M.A. Batzer, L.
Ding, G.T. Marth, G. McVean, J. Sebat, M. Snyder, J. Wang, K. Ye, E.E. Eichler, M.B. Gerstein, M.E.
Hurles, C. Lee, S.A. McCarroll, J.O. Korbel, and P. Genomes, *Mapping copy number variation by*
population-scale genome sequencing. Nature, 2011. 470(7332): p. 59-65.
82.   Zhao, M., Q. Wang, Q. Wang, P. Jia, and Z. Zhao, *Computational tools for copy number variation*
(CNV) detection using next-generation sequencing data: features and perspectives. BMC
Bioinformatics, 2013. 14 Suppl 11(Suppl 11): p. S1.
83.   Guan, P. and W.K. Sung, *Structural variation detection using next-generation sequencing data: A*
comparative technical review. Methods, 2016. 102: p. 36-49.
84.   Arnot, D.E., E. Ronander, and D.C. Bengtsson, *The progression of the intra-erythrocytic cell cycle*
of Plasmodium falciparum and the role of the centriolar plaques in asynchronous mitotic division
during schizogony. Int J Parasitol, 2011. 41(1): p. 71-80.
85.   Zong, C., S. Lu, A.R. Chapman, and X.S. Xie, *Genome-wide detection of single-nucleotide and*
copy-number variations of a single human cell. Science, 2012. 338(6114): p. 1622-6.
86.   Lasken, R.S., *Single-cell sequencing in its prime.* Nat Biotechnol, 2013. 31(3): p. 211-2.
87.   Liu, S., E.R. Ebel, A. Luniewski, J. Zulawinska, M.L. Simpson, J. Kim, N. Ene, T.W.A. Braukmann, M.
Congdon, W. Santos, E. Yeh, and J.L. Guler, *Direct long read visualization reveals metabolic*
interplay between two antimalarial drug targets. bioRxiv, 2023.
88.   Kirkman, L.A., E.A. Lawrence, and K.W. Deitsch, *Malaria parasites utilize both homologous*
recombination and alternative end joining pathways to maintain genome integrity. Nucleic Acids
Res, 2014. 42(1): p. 370-9.
89.   Triglia, T., S.J. Foote, D.J. Kemp, and A.F. Cowman, *Amplification of the multidrug resistance*
gene pfmdr1 in Plasmodium falciparum has arisen as multiple independent events. Mol. Cell.
Biol., 1991. 11(10): p. 5244-5250.
90.   Luth, M.R., K.P. Godinez-Macias, D. Chen, J. Okombo, V. Thathy, X. Cheng, S. Daggupati, H.
Davies, S.K. Dhingra, J.M. Economy, R.C.S. Edgar, M.G. Gomez-Lorenzo, E.S. Istvan, J.C. Jado,
G.M. LaMonte, B. Melillo, S. Mok, S.K. Narwal, T. Ndiaye, S. Ottilie, S. Palomo Diaz, H. Park, S.
Pena, F. Rocamora, T. Sakata-Kato, J.L. Small-Saunders, R.L. Summers, P.K. Tumwebaze, M.
Vanaerschot, G. Xia, T. Yeo, A. You, F.J. Gamo, D.E. Goldberg, M.C.S. Lee, C.W. McNamara, D.
Ndiaye, P.J. Rosenthal, S.L. Schreiber, G. Serra, J.L. De Siqueira-Neto, T.S. Skinner-Adams, A.C.
Uhlemann, N. Kato, A.K. Lukens, D.F. Wirth, D.A. Fidock, and E.A. Winzeler, *Systematic in vitro*
evolution in Plasmodium falciparum reveals key determinants of drug resistance. Science, 2024.
386(6725): p. eadk9893.
91.   Matthews, H., C.W. Duffy, and C.J. Merrick, *Checks and balances? DNA replication and the cell*
cycle in Plasmodium. Parasit Vectors, 2018. 11(1): p. 216.
92.   Bindra, R.S., P.J. Schaffer, A. Meng, J. Woo, K. Maseide, M.E. Roth, P. Lizardi, D.W. Hedley, R.G.
Bristow, and P.M. Glazer, *Down-regulation of Rad51 and decreased homologous recombination*
in hypoxic cancer cells. Molecular and Cellular Biology, 2004. 24(19): p. 8504-8518.

93.   Saleh-Gohari, N., H.E. Bryant, N. Schultz, K.M. Parker, T.N. Cassel, and T. Helleday, *Spontaneous*
homologous recombination is induced by collapsed replication forks that are caused by
endogenous DNA single-strand breaks. Mol Cell Biol, 2005. 25(16): p. 7158-69.
94.   Hastings, P.J., G. Ira, and J.R. Lupski, *A microhomology-mediated break-induced replication*
model for the origin of human copy number variation. PLoS Genet, 2009. 5(1): p. e1000327.
95.   Durkin, S.G., R.L. Ragland, M.F. Arlt, J.G. Mulle, S.T. Warren, and T.W. Glover, *Replication stress*
induces tumor-like microdeletions in FHIT/FRA3B. Proc Natl Acad Sci U S A, 2008. 105(1): p. 246-
51.
96.   Chen, L., W. Zhou, C. Zhang, J.R. Lupski, L. Jin, and F. Zhang, *CNV instability associated with DNA*
replication dynamics: evidence for replicative mechanisms in CNV mutagenesis. Hum Mol Genet,
2015. 24(6): p. 1574-83.
97.   White, N.J., *Malaria parasite clearance.* Malar J, 2017. 16(1): p. 88.
98.   Shor, E., C.A. Fox, and J.R. Broach, *The yeast environmental stress response regulates*
mutagenesis induced by proteotoxic stress. PLoS Genet, 2013. 9(8): p. e1003680.
99.   WHO, *World Malaria Report 2023.* 2023.
100.   Okombo, J. and D.A. Fidock, *Towards next-generation treatment options to combat Plasmodium*
falciparum malaria. Nat Rev Microbiol, 2024.
101.   Wilson, C., A. Serrano, A. Wasley, M. Bogenschutz, A. Shankar, and D. Wirth, *Amplification of a*
gene related to mammalian mdr genes in drug-resistant Plasmodium falciparum. Science, 1989.
244(4909): p. 1184-1186.
102.   Nair, S., D. Nash, D. Sudimack, A. Jaidee, M. Barends, A.-C. Uhlemann, S. Krishna, F. Nosten, and
T.J.C. Anderson, *Recurrent Gene Amplification and Soft Selective Sweeps during Evolution of*
Multidrug Resistance in Malaria Parasites. Mol Biol Evol, 2007. 24(2): p. 562-573.
103.   Ravenhall, M., E.D. Benavente, M. Mipando, A.T. Jensen, C.J. Sutherland, C. Roper, N. Sepulveda,
D.P. Kwiatkowski, J. Montgomery, K.S. Phiri, A. Terlouw, A. Craig, S. Campino, H. Ocholla, and
T.G. Clark, *Characterizing the impact of sustained sulfadoxine/pyrimethamine use upon the*
Plasmodium falciparum population in Malawi. Malar J, 2016. 15(1): p. 575.
104.   Leroy, D., F. Macintyre, Y. Adoke, S. Ouoba, A. Barry, G. Mombo-Ngoma, J.M. Ndong Ngomo, R.
Varo, Y. Dossou, A.K. Tshefu, T.T. Duong, B.Q. Phuc, B. Laurijssens, R. Klopper, N. Khim, E.
Legrand, and D. Menard, *African isolates show a high proportion of multiple copies of the*
Plasmodium falciparum plasmepsin-2 gene, a piperaquine resistance marker. Malar J, 2019.
18(1): p. 126.
105.   Bopp, S., P. Magistrado, W. Wong, S.F. Schaffner, A. Mukherjee, P. Lim, M. Dhorda, C.
Amaratunga, C.J. Woodrow, E.A. Ashley, N.J. White, A.M. Dondorp, R.M. Fairhurst, F. Ariey, D.
Menard, D.F. Wirth, and S.K. Volkman, *Plasmepsin II-III copy number accounts for bimodal*
piperaquine resistance among Cambodian Plasmodium falciparum. Nat Commun, 2018. 9(1): p.
1769.
106.   Dharia, N., A. Sidhu, M. Cassera, S. Westenberger, S. Bopp, R. Eastman, D. Plouffe, S. Batalov, D.
Park, S. Volkman, D. Wirth, Y. Zhou, D. Fidock, and E. Winzeler, *Use of high-density tiling*
microarrays to identify mutations globally and elucidate mechanisms of drug resistance in
Plasmodium falciparum. Genome Biology, 2009. 10(2): p. R21.
107.   Rottmann, M., C. McNamara, B.K. Yeung, M.C. Lee, B. Zou, B. Russell, P. Seitz, D.M. Plouffe, N.V.
Dharia, J. Tan, S.B. Cohen, K.R. Spencer, G.E. Gonzalez-Paez, S.B. Lakshminarayana, A. Goh, R.
Suwanarusk, T. Jegla, E.K. Schmitt, H.P. Beck, R. Brun, F. Nosten, L. Renia, V. Dartois, T.H. Keller,
D.A. Fidock, E.A. Winzeler, and T.T. Diagana, *Spiroindolones, a potent compound class for the*
treatment of malaria. Science, 2010. 329(5996): p. 1175-80.
108.   Ross, L.S., F.J. Gamo, M.J. Lafuente-Monasterio, O.M. Singh, P. Rowland, R.C. Wiegand, and D.F.
Wirth, *In Vitro Resistance Selections for Plasmodium falciparum Dihydroorotate Dehydrogenase*

Inhibitors Give Mutants with Multiple Point Mutations in the Drug-binding Site and Altered
Growth. J Biol Chem, 2014. 289(26): p. 17980-95.

109.   Phillips, M.A., J. Lotharius, K. Marsh, J. White, A. Dayan, K.L. White, J.W. Njoroge, F. El Mazouni,
Y. Lao, S. Kokkonda, D.R. Tomchick, X. Deng, T. Laird, S.N. Bhatia, S. March, C.L. Ng, D.A. Fidock,
S. Wittlin, M. Lafuente-Monasterio, F.J. Benito, L.M. Alonso, M.S. Martinez, M.B. Jimenez-Diaz,
S.F. Bazaga, I. Angulo-Barturen, J.N. Haselden, J. Louttit, Y. Cui, A. Sridhar, A.M. Zeeman, C.
Kocken, R. Sauerwein, K. Dechering, V.M. Avery, S. Duffy, M. Delves, R. Sinden, A. Ruecker, K.S.
Wickham, R. Rochford, J. Gahagen, L. Iyer, E. Riccio, J. Mirsalis, I. Bathhurst, T. Rueckle, X. Ding,
B. Campo, D. Leroy, M.J. Rogers, P.K. Rathod, J.N. Burrows, and S.A. Charman, *A long-duration*
dihydroorotate dehydrogenase inhibitor (DSM265) for prevention and treatment of malaria. Sci
Transl Med, 2015. 7(296): p. 296ra111.

110.   Cowell, A. and E. Winzeler, *Exploration of the Plasmodium falciparum Resistome and Druggable*
Genome Reveals New Mechanisms of Drug Resistance and Antimalarial Targets. Microbiol
Insights, 2018. 11: p. 1178636118808529.

111.   Rocamora, F., L. Zhu, K.Y. Liong, A. Dondorp, O. Miotto, S. Mok, and Z. Bozdech, *Oxidative stress*
and protein damage responses mediate artemisinin resistance in malaria parasites. PLoS Pathog,
2018. 14(3): p. e1006930.

112.   Thaithong, S., L.C. Ranford-Cartwright, N. Siripoon, P. Harnyuttanakorn, N.S. Kanchanakhan, A.
Seugorn, K. Rungsihirunrat, P.V. Cravo, and G.H. Beale, *Plasmodium falciparum: gene mutations*
and amplification of dihydrofolate reductase genes in parasites grown in vitro in presence of
pyrimethamine. Exp Parasitol, 2001. 98(2): p. 59-70.

113.   Istvan, E.S., N.V. Dharia, S.E. Bopp, I. Gluzman, E.A. Winzeler, and D.E. Goldberg, *Validation of*
isoleucine utilization targets in Plasmodium falciparum. Proc Natl Acad Sci U S A, 2011. 108(4): p.
1627-32.

114.   Miles, A., Z. Iqbal, P. Vauterin, R. Pearson, S. Campino, M. Theron, K. Gould, D. Mead, E. Drury, J.
O'Brien, V. Ruano Rubio, B. MacInnis, J. Mwangi, U. Samarakoon, L. Ranford-Cartwright, M.
Ferdig, K. Hayton, X.Z. Su, T. Wellems, J. Rayner, G. McVean, and D. Kwiatkowski, *Indels,*
structural variation, and recombination drive genomic diversity in Plasmodium falciparum.
Genome Res, 2016. 26(9): p. 1288-99.

115.   Rathod, P.K., T. McErlean, and P.C. Lee, *Variations in frequencies of drug resistance in*
Plasmodium falciparum. Proc Natl Acad Sci U S A, 1997. 94(17): p. 9389-93.

116.   McDaniels, J.M., A.C. Huckaby, S.A. Carter, S. Lingeman, A. Francis, M. Congdon, W. Santos, P.K.
Rathod, and J.L. Guler, *Extrachromosomal DNA amplicons in antimalarial-resistant Plasmodium*
falciparum. Mol Microbiol, 2021. 115(4): p. 574-590.

117.   Sugino, A. and K. Nakayama, *DNA polymerase alpha mutants from a Drosophila melanogaster*
cell line. Proc Natl Acad Sci U S A, 1980. 77(12): p. 7049-53.

118.   Inselburg, J. and H.S. Banyal, *Plasmodium falciparum: synchronization of asexual development*
with aphidicolin, a DNA synthesis inhibitor. Exp Parasitol, 1984. 57(1): p. 48-54.

119.   Brown, A.C., C.C. Moore, and J.L. Guler, *Cholesterol-dependent enrichment of understudied*
erythrocytic stages of human Plasmodium parasites. Sci Rep, 2020. 10(1): p. 4591.

120.   Rodrigues, O.R. and S. Monard, *A rapid method to verify single-cell deposition setup for cell*
sorters. Cytometry A, 2016. 89(6): p. 594-600.

121.   Nam, D., S. Kim, J.H. Kim, S. Lee, D. Kim, J. Son, D. Kim, B.S. Cha, E.S. Lee, and K.S. Park, *Low-*
Temperature Loop-Mediated Isothermal Amplification Operating at Physiological Temperature.
Biosensors (Basel), 2023. 13(3).

122.   Brown, N., C. da Silva, C. Webb, D. Matias, B. Dias, B. Cancio, M. Silva, R. Viegas, C. Salvador, N.
Chivale, S. Luis, P. Arnaldo, J. Zulawinska, C.C. Moore, F. Nogueira, and J.L. Guler, *Antimalarial*

resistance risk in Mozambique detected by a novel quadruplex droplet digital PCR assay.
Antimicrob Agents Chemother, 2024. 68(7): p. e0034624.

123.   Daniels, R., D. Ndiaye, M. Wall, J. McKinney, P.D. Sene, P.C. Sabeti, S.K. Volkman, S. Mboup, and
D.F. Wirth, *Rapid, field-deployable method for genotyping and discovery of single-nucleotide*
polymorphisms associated with drug resistance in Plasmodium falciparum. Antimicrob Agents
Chemother, 2012. 56(6): p. 2976-86.

124.   Kassaza, K., A.C. Long, J.M. McDaniels, M. Andre, W. Fredrickson, D. Nyehangane, P. Orikiriza,
D.J. Operario, J. Bazira, J.A. Mwanga-Amumpaire, C.C. Moore, J.L. Guler, and Y. Boum, 2nd,
Surveillance of Plasmodium falciparum pfcrt haplotypes in southwestern uganda by high-
resolution melt analysis. Malar J, 2021. 20(1): p. 114.

125.   Alker, A.P., V. Mwapasa, and S.R. Meshnick, *Rapid real-time PCR genotyping of mutations*
associated with sulfadoxine-pyrimethamine resistance in Plasmodium falciparum. Antimicrob
Agents Chemother, 2004. 48(8): p. 2924-9.

126.   Chiang, C., R.M. Layer, G.G. Faust, M.R. Lindberg, D.B. Rose, E.P. Garrison, G.T. Marth, A.R.
Quinlan, and I.M. Hall, *SpeedSeq: ultra-fast personal genome analysis and interpretation.* Nat
Methods, 2015. 12(10): p. 966-8.

127.   Li, H., B. Handsaker, A. Wysoker, T. Fennell, J. Ruan, N. Homer, G. Marth, G. Abecasis, R. Durbin,
and S. Genome Project Data Processing, *The Sequence Alignment/Map format and SAMtools.*
Bioinformatics, 2009. 25(16): p. 2078-9.

128.   Garcia-Alcalde, F., K. Okonechnikov, J. Carbonell, L.M. Cruz, S. Gotz, S. Tarazona, J. Dopazo, T.F.
Meyer, and A. Conesa, *Qualimap: evaluating next-generation sequencing alignment data.*
Bioinformatics, 2012. 28(20): p. 2678-9.

129.   Huang, L., F. Ma, A. Chapman, S. Lu, and X.S. Xie, *Single-Cell Whole-Genome Amplification and*
Sequencing: Methodology and Applications. Annu Rev Genomics Hum Genet, 2015. 16: p. 79-
102.

130.   Chen, C., D. Xing, L. Tan, H. Li, G. Zhou, L. Huang, and X.S. Xie, *Single-cell whole-genome analyses*
by Linear Amplification via Transposon Insertion (LIANTI). Science, 2017. 356(6334): p. 189-194.

131.   McKenna, A., M. Hanna, E. Banks, A. Sivachenko, K. Cibulskis, A. Kernytsky, K. Garimella, D.
Altshuler, S. Gabriel, M. Daly, and M.A. DePristo, *The Genome Analysis Toolkit: a MapReduce*
framework for analyzing next-generation DNA sequencing data. Genome Res, 2010. 20(9): p.
1297-303.

132.   DePristo, M.A., E. Banks, R. Poplin, K.V. Garimella, J.R. Maguire, C. Hartl, A.A. Philippakis, G. del
Angel, M.A. Rivas, M. Hanna, A. McKenna, T.J. Fennell, A.M. Kernytsky, A.Y. Sivachenko, K.
Cibulskis, S.B. Gabriel, D. Altshuler, and M.J. Daly, *A framework for variation discovery and*
genotyping using next-generation DNA sequencing data. Nat Genet, 2011. 43(5): p. 491-8.

133.   Van der Auwera, G.A., M.O. Carneiro, C. Hartl, R. Poplin, G. Del Angel, A. Levy-Moonshine, T.
Jordan, K. Shakir, D. Roazen, J. Thibault, E. Banks, K.V. Garimella, D. Altshuler, S. Gabriel, and
M.A. DePristo, *From FastQ data to high confidence variant calls: the Genome Analysis Toolkit*
best practices pipeline. Curr Protoc Bioinformatics, 2013. 43(1110): p. 11 10 1-11 10 33.

134.   MalariaGen, A. Ahouidi, M. Ali, J. Almagro-Garcia, A. Amambua-Ngwa, C. Amaratunga, R. Amato,
L. Amenga-Etego, B. Andagalu, T.J.C. Anderson, V. Andrianaranjaka, T. Apinjoh, C. Ariani, E.A.
Ashley, S. Auburn, G.A. Awandare, H. Ba, V. Baraka, A.E. Barry, P. Bejon, G.I. Bertin, M.F. Boni, S.
Borrmann, T. Bousema, O. Branch, P.C. Bull, G.B.J. Busby, T. Chookajorn, K. Chotivanich, A.
Claessens, D. Conway, A. Craig, U. D'Alessandro, S. Dama, N.P. Day, B. Denis, M. Diakite, A.
Djimde, C. Dolecek, A.M. Dondorp, C. Drakeley, E. Drury, P. Duffy, D.F. Echeverry, T.G. Egwang,
B. Erko, R.M. Fairhurst, A. Faiz, C.A. Fanello, M.M. Fukuda, D. Gamboa, A. Ghansah, L. Golassa, S.
Goncalves, W.L. Hamilton, G.L.A. Harrison, L. Hart, C. Henrichs, T.T. Hien, C.A. Hill, A. Hodgson, C.
Hubbart, M. Imwong, D.S. Ishengoma, S.A. Jackson, C.G. Jacob, B. Jeffery, A.E. Jeffreys, K.J.

Johnson, D. Jyothi, C. Kamaliddin, E. Kamau, M. Kekre, K. Kluczynski, T. Kochakarn, A. Konate,
D.P. Kwiatkowski, M.P. Kyaw, P. Lim, C. Lon, K.M. Loua, O. Maiga-Ascofare, C. Malangone, M.
Manske, J. Marfurt, K. Marsh, M. Mayxay, A. Miles, O. Miotto, V. Mobegi, O.A. Mokuolu, J.
Montgomery, I. Mueller, P.N. Newton, T. Nguyen, T.N. Nguyen, H. Noedl, F. Nosten, R.
Noviyanti, A. Nzila, L.I. Ochola-Oyier, H. Ocholla, A. Oduro, I. Omedo, M.A. Onyamboko, J.B.
Ouedraogo, K. Oyebola, R.D. Pearson, N. Peshu, A.P. Phyo, C.V. Plowe, R.N. Price, S.
Pukrittayakamee, M. Randrianarivelojosia, J.C. Rayner, P. Ringwald, K.A. Rockett, K. Rowlands, L.
Ruiz, D. Saunders, A. Shayo, P. Siba, V.J. Simpson, J. Stalker, X.Z. Su, C. Sutherland, S. Takala-
Harrison, L. Tavul, V. Thathy, A. Tshefu, F. Verra, J. Vinetz, T.E. Wellems, J. Wendler, N.J. White, I.
Wright, W. Yavo and H. Ye, *An open dataset of Plasmodium falciparum genome variation in*
7,000 worldwide samples. Wellcome Open Res, 2021. 6: p. 42.

135. Abyzov, A., A.E. Urban, M. Snyder, and M. Gerstein, *CNVnator: an approach to discover,*
genotype, and characterize typical and atypical CNVs from family and population genome
sequencing. Genome Res, 2011. 21(6): p. 974-84.

136. Venkatraman, E.S. and A.B. Olshen, *A faster circular binary segmentation algorithm for the*
analysis of array CGH data. Bioinformatics, 2007. 23(6): p. 657-63.

137. Xiao, F., Y. Niu, N. Hao, Y. Xu, Z. Jin, and H. Zhang, *modSaRa: a computationally efficient R*
package for CNV identification. Bioinformatics, 2017. 33(15): p. 2384-2385.

138. Mi, H., D. Ebert, A. Muruganujan, C. Mills, L.P. Albou, T. Mushayamaha, and P.D. Thomas,
PANTHER version 16: a revised family classification, tree-based classification tool, enhancer
regions and extensive API. Nucleic Acids Res, 2021. 49(D1): p. D394-D403.

139. Singh, G. and D. Gupta, *In-Silico Functional Annotation of Plasmodium falciparum Hypothetical*
Proteins to Identify Novel Drug Targets. Front Genet, 2022. 13: p. 821516.

1243