Longitudinal prevalence and co-carriage of pathogens associated with nursing home acquired pneumonia in three long-term care facilities

Ryann N. Whealy[1,2,], Alexander Roberts[1,2], Tara N. Furstenau[2], Skylar Timm[1,2], Sara Maltinsky[1], Sydney Wells[1], Kylie Drake[1], Kayla Ramirez[1], Candice Bolduc[3], Ann Ross[3], Talima Pearson[1], Viacheslav Y. Fofanov[1,2]

[1] Pathogen and Microbiome Institute, Northern Arizona University, Flagstaff, Arizona, United States of America
[2] School of Informatics, Computing, and Cyber Systems, Northern Arizona University, Flagstaff, Arizona, United States of America.
[3] Mobile Dentistry of Arizona, Mesa, Arizona, United States of America

Corresponding authors:
Viacheslav Y. Fofanov
1295 Knoles Dr, Flagstaff, AZ 86011, USA
Viacheslav.Fofanov@nau.edu

Talima Pearson
1395 Knoles Dr, Flagstaff, AZ 86011, USA
Talima.Pearson@nau.edu

Word count: 3301

Abstract

Nursing home acquired pneumonia (NHAP), and its subset – aspiration-associated pneumonia, is a leading cause of morbidity and mortality among residents in long-term care facilities (LTCFs). Understanding colonization dynamics of respiratory pathogens in LTCF residents is essential for effective infection control. This study examines the longitudinal trends in prevalence, persistence, bacterial load, and co-colonization patterns of five respiratory pathogens in three LTCFs in Phoenix, Arizona. Anterior nares and oral swabs were collected every other week and tested using qPCR for *Haemophilus influenzae*, *Pseudomonas aeruginosa*, *Streptococcus pneumoniae*, *Staphylococcus aureus*, and *Chlamydia pneumoniae*. Weekly average positivity rates were 17.75% for *H. influenzae* (0% - 39.39%), 9.95% for *P. aeruginosa* (0% - 37.74%), 31.89% for *S. pneumoniae* (1.79% - 41.67%), and for 28.00% for *S. aureus* (0% - 55.36%). *C. pneumoniae* was not detected. *H. influenzae* and *S. pneumoniae* predominantly colonized the oral cavity, while *P. aeruginosa* and *S. aureus* predominantly colonized the nasal cavity. *S. pneumoniae* and *S. aureus* colonizations were significantly more persistent than *H. influenzae* and *P. aeruginosa*, with persistence correlating with significantly higher bacterial loads. Co-colonization did occur in ~20% of positive samples, but appeared to be due to random chance. This study reveals distinct colonization patterns among respiratory pathogens in LTCF residents, highlighting differences in site-specific prevalence, persistence, and bacterial load. These findings underscore the importance of longitudinal monitoring to inform targeted infection control strategies in LTCFs.

Introduction

Nursing home acquired pneumonia (NHAP) is a leading cause of morbidity and mortality in long term care facilities (LTCFs), accounting for 40% of hospitalizations and up to 33% of deaths among residents [1]. It is estimated that 365 out of every 1000 LTCF residents are diagnosed with pneumonia annually [2], a rate 6-11 times higher than the incidence of pneumonia in elderly people living in the community [3,4]. Further, the medical care and treatment required for NHAP contribute to significant healthcare costs due to the potential for prolonged hospital stays, need for intensive care, and treatment-resistant etiologies. While pneumonia can often be treated by LTCF staff at an average cost of approximately $3,700 [5], up to 30% of patients require hospitalization which increases average treatment cost to $10,400 [6,7]. For patients who require hospitalization, mortality rates range from 13-45% [8,9]. These high costs and poor outcomes underscore the need for targeted, evidence-based infection prevention and stewardship strategies in nursing homes [10].

Micro-aspiration is the primary pathogenic mechanism of pneumonia in otherwise healthy individuals, while macro-aspiration (the cause of aspiration pneumonia) typically occurs only in those experiencing difficulty swallowing or changes in consciousness [11]. Older individuals are more likely to sustain an infection induced through aspirating large amounts of opportunistic pathogens due to decreased humoral and cell-mediated immunity, increased likelihood of underlying airway disorders, and altered levels of consciousness. Many bacterial etiological agents of pneumonia asymptomatically colonize the oropharyngeal mucosa or other areas of the body. As such, sources of many pneumonia infections are endogenous and due to autoinfection [12]. Controlling pneumonia from endogenous sources requires management and control of these pathogens at individual and institutional levels to reduce the likelihood of acquisition, carriage, and infection.

Many pneumonia-causing bacteria colonize dental plaque biofilms [13–16], providing a potential source for micro- and macro-aspiration pneumonia [14,16]. Up to 80% correlation has been found between aerobic dental plaque colonization and the causative agent of aspiration associated-pneumonia [16]. Unsurprisingly, poor oral hygiene increases the risk of pneumonia [14,17–19], highlighting its potential as an intervention target. Further, elderly LTCF residents tend to have particularly poor oral health with high plaque scores [20], indicating that improving oral health among this population may be especially impactful. While promising, oral health interventions have shown mixed results in lowering NHAP incidence. In some studies, improved oral hygiene has been shown to decrease the incidence rate of pneumonia, potentially preventing as many as 10% of NHAP deaths [16,21–24]. Other studies have found no advantage in toothbrushing and chlorhexidine rinses over standard care in preventing NHAP [25]. In short, while the relationship

95   between colonization and infection is well established, much research still needs to be done to assess the potential for oral
96   hygiene interventions in reducing NHAP.
97
Due to the varied etiological agents involved in NHAP, characterization of longitudinal oral colonization and associated
transmission dynamics are critical for guiding future intervention studies. Furthering the understanding of the transmission
dynamics of the most common etiological agents responsible for NHAP (and, simultaneously, developing better assays to
detect them) will enable future intervention studies aimed at equipping LTCF clinicians with strategies for mitigation and
prevention. Towards this end, this study characterizes the colonization dynamics of common etiological agents of NHAP
in the oral and nasal cavity in LTCF residents.
104
105   **Methods**
106
107 **Sample and data collection**
108
Across three LTCFs in the Phoenix metropolitan area, 121 residents were enrolled to participate in this study after
providing written consent (IRB: 1766728). Participants provided their pneumonia vaccination status, as well as
demographic information (age, sex, and ethnicity). Swabs of the anterior nares and oral mucosa were collected every other
week from November 2021 to November 2023 (49 events total). Oral hygiene professionals supervised the self-sampling,
which has proven to be an effective and sensitive method of detecting pathogens at these body sites [26]. Collected swabs
were stored in 1mL of Liquid Amies at -20ºC until processing.
115
116 **Sample processing, testing, and sequencing**
117
Multi-pathogen qPCR panel
Total DNA was extracted from nasal and oral swab samples using the Applied Biosystems MagMAX Viral and Pathogen
Nucleic Acid Isolation Kit with the KingFisher Flex system, according to the manufacturer's instructions. qPCR was
performed using a multi-pathogen panel (adapted from published assays) [27–31] designed to detect and quantify
Haemophilus influenzae, *Chlamydia pneumoniae*, *Pseudomonas aeruginosa*, *Streptococcus pneumoniae*, and
Staphylococcus aureus (Supplemental Table 1). We performed an in-silico PCR to evaluate the specificity of the primers
and probes across a comprehensive set of reference genomes for each pathogen. Some oligonucleotide sequences were
adjusted to maximize complementarity with a wider range of genomes. The assays were validated independently using
positive controls, and optimized to be run in duplex. There were interactions between oligos for *H. influenzae* and *S.*
pneumoniae as well as *S. aureus* and *P. aeruginosa*, but no other combinations produced cross-primer dimerization based
on in-silico PCR. Serial dilutions of quantitative genomic DNA were included in all runs to allow for relative DNA
quantification using standard curve analysis.
130
131 Each qPCR reaction was carried out in a 10 μL volume containing 5 μL of TaqMan Universal Master Mix, 2 μL of
132 template DNA, final concentrations of 0.6 μM for primers and 0.3 μM for probes, and nuclease-free water. Cycling
133 conditions were as follows: 50°C for 2 minutes, 95°C for 15 minutes, then 45 cycles of 95°C for 15 seconds and 57°C for
134 1 minute. Amplifications were completed using an Applied Biosystems QuantStudio™ 7 Pro System, and results were
135 analyzed using the QuantStudio™ Design & Analysis Software (version 2.6.0).
136
Secondary validation using multi-pathogen amplicon sequencing
To confirm that the multiplex qPCR assays accurately identified the target species, we used a species-specific targeted
amplicon sequencing approach to detect additional genomic regions unique to each pathogen in a subset of qPCR positive
samples (Supplemental Table 2). The multiplex PCR was performed using the KAPA 2G Fast Multiplex PCR Master Mix
in 25μL reaction volumes, with 5ng of template DNA, 0.2 μM concentration of each primer, and water to reach the final
volume. PCR thermocycling conditions consisted of an initial denaturation at 95°C for 3 minutes, followed by 35 cycles

of denaturation at 95°C for 15 seconds, annealing at 60°C for 30 seconds, and extension at 72°C for 1 minute and 30
seconds and a final extension at 72°C for 1 minute. PCR products were prepared for sequencing using our standard
amplicon sequencing protocol [32] and were sequenced on the Illumina MiniSeq platform. Reads were aligned to
reference genomes (Accessions in Supplemental Table 2) using BWA-MEM v0.7.17 [33].

Data analysis

All data cleaning and statistical analyses were conducted using R version 4.4.1. Statistical significance was set at a p-
value of less than 0.05 for all tests. To minimize bias from sampling inconsistency, results are only included from
participants who attended at least ⅓ of sampling events.

McNemar's test [34] was used to compare colonization rates between nasal and oral samples for each pathogen,
accounting for the paired nature of the samples collected from the same participants. Matched odds ratios with 95%
confidence intervals were calculated to quantify the magnitude of differences in colonization rates. To estimate the
probabilities of transitioning between colonization states over time, we employed a Markov chain analysis with
bootstrapping (1000 for each pathogen). The resulting transition matrices generated from bootstrapping were averaged to
produce a final transition matrix representing the mean transition probabilities. Linear mixed models were fitted using the
lme4 [35] and lmerTest [36] packages to quantify predictors of average relative DNA quantities. Full models included
transient vs. persistent colonization, age, sex, pneumococcal vaccination, and recent estimated exposure as fixed effects as
well as random intercepts for each participant to account for repeated measures. The results are reported in relative units
(RQ), indicating fold changes in bacterial abundance relative to a standardized control. Co-colonization patterns were
visualized using an upset plot created with the UpSetR [37] package, highlighting the intersections between different
pathogens. To investigate potential temporal associations between colonization with different pathogens, we performed a
correlation analysis incorporating current and lagged colonization statuses. However, no significant correlations were
found, so this analysis was exploratory in nature.

Results

Participant description
The 121 participants ranged in age from 60 to 97 (median = 83, mean = 82.97) and were a majority female (76.03%). .
Race/ethnicity was not included in this analysis due to a significantly imbalanced distribution within the sample;
approximately 96% of participants were non-Hispanic whites. The subset of participants who attended at least one-third of
sampling events (n=85) closely matched the demographic distribution of the overall study population, with a mean age of
83 and approximately ¾ being female. These individuals attended an average of 34 out of 49 sampling events,
approximately 70% were vaccinated against streptococcal pneumonia, and were evenly distributed between the three
LTCFs.

Colonization rates at nasal and oral sites
We observed wide ranges of colonization rates for the pathogens of interest and significant site-specificity towards the
nasal or oral cavity. Of the approximately 6,000 samples collected from the 85 participants in LTCFs, 9.99% of samples
were positive for *Haemophilus influenzae*, 5.78% were positive for *Pseudomonas aeruginosa*, 22.07% were positive for
Streptococcus pneumoniae, and 18.01% were positive for *Staphylococcus aureus*. Notably, *Chlamydia pneumoniae* was
not detected in any samples. Table 1 provides an overview of the colonization metrics for each pathogen. Across time
points, notable variation was observed in both sample-level and individual-level positivity rates. The cumulative incidence
values illustrate that a majority of participants were colonized by at least one of these pathogens during the study period.

Table 1: Colonization Metrics by Pathogen

Pathogen	Average weekly sample positivity (range)	Average weekly individual positivity (range)	Cumulative incidence
H. influenzae	9.81% (0–23.48)	17.75% (0–39.39)	81.18%
P. aeruginosa	5.56% (0–20)	9.95% (0–37.74)	83.53%
S. pneumoniae	21.70% (0.89–31.62)	31.89% (1.79–41.67)	78.82%
S. aureus	17.80% (0–36.13)	28.00% (0–55.36)	91.76%

Our findings indicate that these pathogens differentially colonize the oral and nasal cavities, with *H. influenzae* and *S.*
pneumoniae predominating in the oral cavity, and *P. aeruginosa* and *S. aureus* more frequently detected in the nasal
cavity (Table 2). These differences were statistically significant (McNemar's test, $p < 1 \times 10^{-6}$), suggesting that the
observed distribution patterns are not due to random variation but reflect true biological differences in site-specific
colonization. Matched odds ratios further quantify these disparities, demonstrating substantially reduced odds of nasal
colonization for *H. influenzae* and *S. pneumoniae* relative to oral colonization, but notably increased odds of nasal
colonization for *P. aeruginosa* and *S. aureus*.

Longitudinal colonization rates
There were statistically significant changes in participant positivity over time for some of the pathogens (Figure 1). *H.*
influenzae colonization decreased over time in both sample types, while *P. aeruginosa, S. aureus,* and *S. pneumoniae*
colonization decreased in nasal samples over time (p < 0.01). Oral colonization with *P. aeruginosa, S. pneumoniae,* and *S.*
aureus did not significantly change over time. Colonization at the individual level changed significantly over time for *H.*
influenzae, P. aeruginosa, and *S. aureus* (p < 0.01). These changes were not correlated with any measured variables and
there were no changes in protocol during the study.
206

Figure 1. The proportion of individuals (from oral, nasal, or both sample types), nasal samples, and oral samples testing positive for each pathogen over the course of the study. Pathogens include *Haemophilus influenzae* (Hi, red), *Pseudomonas aeruginosa* (Pa, green), *Streptococcus pneumoniae* (Sp, blue), and *Staphylococcus aureus* (Sa, purple). Each dot represents the positivity rate for a collection event, with trends highlighting temporal dynamics in colonization prevalence across anatomical sites and pathogens.

207
Duration of colonization
The distribution of time participants spent colonized shows notable variability among pathogens (Figure 2). The
proportion of time that participants spent colonized by *S. pneumoniae* and *S. aureus* exhibited a pronounced bimodal
pattern with a peak near zero, indicating never or highly transient colonization, and another near one, reflecting persistent
colonization. Interestingly, colonization by *P. aeruginosa* appears to be predominantly short-lived, with only rare cases of
persistent colonization or carriage. Colonization with *H. influenzae* was similarly skewed, though less drastically. The
observation that individuals tend to be colonized by *S. aureus* either a majority of the time or only transiently has
previously been found [38,39]. Less is known about persistence of colonization with the other pathogens of interest in
healthy individuals. To quantify this variability, participants were categorized as "never" (no colonization over the study),
"transient" (colonization in <50% of sampling events), or "persistent" (colonization in ≥50% of sampling events). For *H.*
influenzae, 67.1% of participants experienced transient colonization, 18.8% were never colonized, and 14.1% were
persistently colonized, indicating that *H. influenzae* colonization is mostly short-lived for otherwise healthy individuals.
For *P. aeruginosa*, 82.4% were transient, 16.5% were never colonized, and 1.18% were persistently colonized, indicating
that *P. aeruginosa* is not typically a long-term colonizer. In contrast, *S. pneumoniae* exhibited a more balanced pattern:
43.5% were transiently colonized, 21.2% were never colonized, and 35.3% were persistently colonized. For *S. aureus*,
only 8.24% were never colonized, with 71.8% transiently colonized and 20% persistent colonization.
224

Figure 2. The proportion of events participants spent colonized by each pathogen (if colonized at least once); *Haemophilus influenzae* (Hi, red), *Pseudomonas aeruginosa* (Pa, green), *Streptococcus pneumoniae* (Sp, blue), and *Staphylococcus aureus* (Sa, purple). The distribution indicates variability in colonization duration across pathogens, with box plots embedded to show median and quartile ranges.

The average duration of continuous colonization at any of the two body sites was approximately 4 weeks for *H. influenzae*, 2.4 weeks for *P. aeruginosa*, 9.6 weeks for *S. pneumoniae*, and 8.8 weeks for *S. aureus* (of those that were colonized at least once during the study). Recurrent colonization (recolonization after pathogen was not detected and thus assumed to be cleared from an individual) was observed in 58.8% of participants for *H. influenzae*, 61.2% for *P. aeruginosa*, 57.6% for *S. pneumoniae*, and 72.9% for *S. aureus*. These colonization patterns are illustrated below (Figure 3). *S. pneumoniae* and *S. aureus* are often persistent colonizers, while *H. influenzae* and *P. aeruginosa* typically appear transiently. Interestingly, persistent colonization with *S. aureus* tends to be in the nasal cavity while persistent colonization with *S. pneumoniae* tends to occur in the oral cavity (Figure 3).

Figure 3. Colonization patterns of *Haemophilus influenzae* (Hi, red), *Pseudomonas aeruginosa* (Pa, green), *Streptococcus pneumoniae* (Sp, blue), and *Staphylococcus aureus* (Sa, purple) shown by A) Colonization at the individual level (from oral, nasal, or both sample types) across the study period where each row represents an individual participant grouped by their LTCF; B) Nasal colonization across the study period, showing varying persistence of pathogens over time in each sampled LTCF; C) Oral colonization across the study period, showing varying persistence of pathogens over time in each sampled LTCF; and D) Duration of continuous colonization for each pathogen at the individual level and at nasal and oral sites, highlighting variability.

Markov chains were used to provide a more nuanced understanding of colonization likelihood and stability over time. For
H. influenzae, at any given time point, the probability of becoming colonized was 9.05%, with a 55.92% chance of
maintaining colonization once established, indicating moderate persistence. *P. aeruginosa* showed a similar colonization
probability of 9.05%, but only a 21.67% probability of remaining colonized, reflecting its low likelihood of sustained
colonization. In contrast, *S. pneumoniae* had a 7.56% probability of colonizing but an 83.64% chance of remaining
colonized, suggesting that *S. pneumoniae* is more persistent once established. For *S. aureus*, the colonization probability
was 10.74%, with a 72.98% likelihood of continuing colonization once established.

To identify which factors most influenced bacterial load (and thus the amount of bacterial material available for macro-
aspiration), relative DNA quantities during positive weeks were analyzed using linear mixed models for each pathogen.
Participants who were never colonized by the pathogen of interest were excluded from this analysis to focus on the
comparison between transient and persistent colonization. For all pathogens, a subjects' categorization of persistent or
transient colonization was the most significant predictor of their bacterial load (except for *P. aeruginosa* which did not
have a pronounced persistent category). For *H. influenzae*, *S. pneumoniae* and *S. aureus*, persistent colonization was
associated with significantly higher bacterial DNA levels compared to transient colonization ($p < 0.001$).

<u>Co-colonization</u>
While colonization by a single pathogen was most common, co-positivity was observed nearly ⅓ of the time that an
individual was colonized (Figure 4). However, the co-colonization rates did not appear to be more frequent than what is
expected by chance ($p > 0.4$, under assumption of independence of colonization events). Additionally, lagged colonization
status analysis suggested that there is no evidence that colonization by one pathogen predicts colonization by another
within the next month ($r < 0.2$).

Figure 4. Co-colonization observed throughout the study within individuals (i.e. considering both oral and nasal results). The bar chart at top shows the percent of instances where individuals who were positive for at least one pathogen were positive for a given combination of pathogens (filled black dots). Each row represents a pathogen, and the bar chart to the right shows the relative prevalence for each pathogen among individuals that tested positive for at least one pathogen.

259
260 **Discussion**
261

Findings from this study highlight several important colonization patterns among LTCF residents, all of which underscore
the value of longitudinal data in revealing temporal variability and individual-level dynamics that would be impossible to
capture with cross-sectional approaches. This study is strengthened by sample collection from multiple body sites,
investigation of multiple pathogens, and the utilization of qPCR for pathogen detection, which has been shown to be an
especially reliable method of detection in older populations [40]. Longitudinal data collection provides valuable
information on how prevalence, persistence, bacterial load, and co-colonization patterns are affected temporally and on an
individual and institutional basis. Additionally, this study provides valuable information about asymptomatic carriage in
otherwise healthy individuals; the very few existing longitudinal NHAP research studies have been limited to hospitalized
patients. These insights can guide the design of effective mitigation strategies for NHAP and associated etiological agents.
271

While average weekly prevalence rates of *H. influenzae* (average 17.75%), *P. aeruginosa* (9.95%), *S. pneumoniae*
(31.89%), and *S. aureus* (28.00%) offer valuable cross-sectional snapshots, the high cumulative incidences over the study
period (81.18%, 83.53%, 78.82%, and 91.76%, respectively), and the duration of individual carriage, reveal a much
broader picture of participant exposure and colonization dynamics. Without tracking these pathogens over time, the
episodic nature of colonization might obscure the extent of overall exposure. For instance, while weekly rates fluctuate
significantly (e.g., *S. pneumoniae* ranging from 0.89% to 31.62% and *S. aureus* from 0% to 36.13%), the cumulative
incidences highlight that almost all participants were colonized at least once over the course of a two-year period. These
findings emphasize that point-prevalence measures alone underestimate the true burden of colonization, potentially
leading to an incomplete understanding of pathogen carriage, transmission, and risks.
281

This study demonstrates distinct colonization patterns of these pathogens in the oral and nasal cavities. Interestingly,
Chlamydia pneumoniae was absent from all samples, possibly due to its obligate intracellular life cycle and tendency to
colonize lower respiratory tract tissue [41]. Over the two-year observation period, *H. influenzae* and *S. aureus*
colonization decreased over time in their predominant niches. In contrast, nasal colonization rates of *P. aeruginosa* and *S.*
pneumoniae increased as the study progressed. These temporal shifts occurred independently of participant demographics,
sampling methods, or changes in study protocol, suggesting that underlying biological or ecological factors, rather than

measured external variables, influenced the observed trends. While sample collection from the nasal and oral cavity
allows characterization of site-specificity in these pathogens, it is likely that different strains may exhibit distinct site-
specificity. Future studies may benefit from increased genotypic resolution and expanding the number of body sites
sampled to further understand site-specificity and potential reservoirs for colonization and spread. Studies have shown
that increasing the number of body sites sampled, increasing the number of replicates tested, and using multiple detection
methods can minimize false negatives and increase prevalence estimates [26]. Consequently, it is likely that the estimates
presented here (using only qPCR for detection and only one replicate from two body sites) are conservative and under-
represent the true prevalence of colonization and co-colonization.
296
The categorization of participants into transient and persistent colonization groups provide insight into the different
strategies utilized by these pathogens. Markov chain simulations confirmed the dominance of transient colonization for *P.
aeruginosa,* and to a lesser degree *H. influenzae*, indicating that these pathogens frequently establish and then are cleared
without long-term colonization. Analysis of bacterial loads indicated that persistent colonization is associated with a
significantly higher bacterial burden for *H. influenzae*, *S. pneumoniae*, and *S. aureus*. This finding suggests that once these
pathogens establish persistent colonization, they are able to maintain higher bacterial densities, which could have
implications for transmission potential and clinical outcomes. This should also guide potential intervention strategies - for
example, attempting to disrupt persistent *S. aureus* colonization via oral healthcare interventions (when persistent
colonization appears to be in the nasal cavity) are likely to be ineffective.
306
Polymicrobial pneumonia represents up to 40% of cases with an identified cause [42]. The involvement of multiple
pathogens can significantly affect infection dynamics, often resulting in more complex clinical outcomes [42]. In our
study, co-colonization was a relatively frequent occurrence, observed approximately ⅓ of the time that an individual was
colonized. However, these rates aligned with simulated expectations ($p > 0.4$), supporting the assumption of
independence. Consistent with this, no evidence was found that colonization by one pathogen predicts subsequent
colonization by another. Our findings suggest that effective preventions should focus on control of individual pathogen
colonization rather than a specialized approach to multiple pathogens.
314
315 This study highlights the complex dynamics of colonization in LTCFs, including variability in site-specific colonization,
316 the distinction between transient and persistent colonization, and the occurrence of co-colonization. Longitudinal data
317 collection was crucial, enabling a better understanding of colonization dynamics and persistence. These insights
318 emphasize the importance of targeted surveillance and intervention strategies that consider the temporal and specific
319 nature of colonization, particularly for high-risk pathogens like *S. aureus* and *P. aeruginosa*.
320

321   **Author Contributions**
322 This study was conceptualized by RNW, VYF, TP, and TNF. Data collection was carried out by CB, AR, TP, and TNF,
323 and formal analyses were performed by RNW and TNF. Writing was completed by RNW, TNF, TP, and AR. Wet lab
324 work and sample processing were performed by RNW, AR, ST, SM, SW, KD, and KR. All authors have reviewed the
325 manuscript.
326   **Acknowledgements**
327 This work was supported by the Centers for Disease Control and Prevention (contract 75D30121C11191 - TP) and the
328 National Institutes of Health (NIH) National Institute on Minority Health and Health Disparities (U54MD012388 - TP)
329 and National Institute of Allergy and Infectious Diseases (R15AI156771 - TP).

Appendix

337 <u>Supplemental Table 1</u>

Pathogen	Positive control	Oligo	Original sequence	Adjusted sequence	Notes
H. influenzae [27]	ATCC® 51907DQ™	Hi P6 F	CCA GCT GCT AAA GTA TTA GTA GAA	n/a	The primers amplified in 89% (85/95) of genomes (156bp amplicon).
		Hi P6 R	TTC ACC GTA AGA TAC TGT GCC	n/a	
		Hi P6 probe	CAG ATG CAG TTG AAG GTT ATT TAG	C**R**G ATG CAG TT**R** AAG GTT ATT TA**R**	The probe was not an exact match for any of the amplicons but worked for all with adjusted sequence
C. pneumoniae [28]	ATCC® VR-1360DQ™	CPTM1	CAT GGT GTC ATT CGC CAA GT	n/a	Primers amplified in 100% (12/12) of genomes (82bp) and probe was an exact match in each
		CPTM2	CGT GTC GTC CAG CCA TTT TA	n/a	
		CP probe	TCT ACG TTG CCT CTA AGA GAA AAC TTC AAG TTG GA	n/a	
P. aeruginosa [29]	ATCC® 47085DQ™	Pa23F	TCC AAG TTT AAG GTG GTA GGC TG	n/a	The primers amplified in 98% (522/532) of genomes (94bp amplicon). All of the amplicons contained an exact match to the probe. A slight change to the reverse primer improves this to 99% (527/532)
		Pa23R	ACC ACT TCG TCA TCT AAA AGA CGA C	ACC ACT TCG TCA TCT AAA AGA C**R**A C	
		Pa23P	AGG TAA ATC CGG GGT TTC AAG GCC	n/a	
S. pneumoniae [30]	Various isolates from Russakoff et al. confirmed using qPCR and WGS [26,43]	cpsA-348F	GCT GTT TTA GCA GAT AGT GAG ATC GA	GCT GTT TTA GCA GAT AGT GA**K** AT**Y** G**R**	The primers initially amplified in only 54% (107/197) of genomes (a 67bp amplicon). After adjustment, this improved
		cpsA-415R	TCC CAG TCG GTG CTG TCA	TCC CAG TCG GTG CT**R** TCA	

		cpsA probe	AAT GTT ACG CAA CTG ACG AG	**R**AT GTT A**Y**G CAA CTG A**Y**G AG	to 89% (176/197). Initially, only 41% (81/197) of the amplicons had an exact match for the probe, but adding ambiguous bases increased this to 87% (172/197).
S. aureus [31]	ATCC® 700669DQ™	SaQuant_47F	AAC TAC TAG GGG AGC CTA ATR AT	n/a	n/a
		SaQuant_99_12 2R	GGT ACT AAC CAA ATC AGG TCA TAA	n/a	n/a
		SaQuant_74_98 FP	TGG CTG AGA TGA AYT GTT CAG ACC C	n/a	n/a

369 Supplemental Table 2
370 The genomic targets included established MLST targets [44–48] as well as additional pathogen-specific targets identified
371 by selecting conserved regions from reference genome alignments within each species [49]. There were a total of 106
372 primer pairs: 11 for *C. pneumoniae,* 9 for *H. influenzae,* 36 for *P. aeruginosa,* 30 for *S. aureus,* and 20 for *S. pneumoniae.*
373

Species	Genome	Forward Primer Seq	Reverse Primer Seq
C. pneumoniae	NC_005043.1	AGCCGAATCCAGAACGT	ACCCGCGCTAGAATGAC
C. pneumoniae	NC_005043.1	AGAGAGATCCATAGCCAGAA	ACCGAAGATGTATGGGTAAC
C. pneumoniae	NC_005043.1	GCTTACATCGCGTCAGT	CCTCGAAGACAGGGATCA
C. pneumoniae	NC_005043.1	AACTTCCCCCTGAGCAT	ACGTTCAAAAATTGCTCGTTT
C. pneumoniae [45]	NC_005043.1	GCTTTAGAATTARSARAWGCT	GATCCTCCGGTATCYGATCC
C. pneumoniae [45]	NC_005043.1	ATGCGCAAGATATCAGTGGG	AAAGCTCCRSTWGMTATMGGWAG
C. pneumoniae [45]	NC_005043.1	GCTTCTARAGTACTTTTAAATG	TATTTRGAAATYTTTKCSAGYCG
C. pneumoniae [45]	NC_005043.1	GGAGTCWCTACWAAAGAAGG	TCGTAYTGYACATCRAAAGG
C. pneumoniae [45]	NC_005043.1	CCTATGATGAATCTKATCAATGG	TCTTCTTCRGCWAGMCCATCT
C. pneumoniae [45]	NC_005043.1	AGATCTTCTTCWGGRGGWAGAGA	TTCYTTCAKAACSTAGGTTTT
C. pneumoniae [45]	NC_005043.1	ATTAAAAAATGTGCTGCT	CCTTCAGGAACATTYAACCC
H. influenzae	NC_007146.2	CCGCTGTTCGCAGTTTG	CGTGCCAGTGCTGCAAT
H. influenzae	NC_007146.2	AAACTCAAGGTCGTGCAT	CGTGCGGTTTTGTACGT
H. influenzae [44]	NC_007146.2	GGTGCACCGGGTGCAGGTAA	CCTAAGATTTTATCTAACTC
H. influenzae [44]	NC_007146.2	ATGGCAGGTGCAAAAGAGAT	TTGTACAACAGGCTTTTGCG
H. influenzae [44]	NC_007146.2	CTTATCGTTGGTCTTGCCGT	TTGGCACTTTCCACTTTTCC
H. influenzae [44]	NC_007146.2	ACCACTTTCGGCGTGGATGG	AAGATTTCCCAGGTGCCAGA
H. influenzae [44]	NC_007146.2	TCATTGTATGATATTGCCCC	ACTTCTGTACCTGCATTTTG
H. influenzae [44]	NC_007146.2	GGTGAAAAAATCAATCGTAC	ATTGAAAGACCAATAGCTGA
H. influenzae [44]	NC_007146.2	ATGGCAACTCAAGAAGAAAA	TTACCAAACATCACGCCTAT
P. aeruginosa	NC_002516.2	CGCCAGCTGTTTCAGTT	ACTCGCCTATGCGATCT
P. aeruginosa	NC_002516.2	AGCGCAGGAGGTTTTCC	CGCCGTGAGGAAATCAG
P. aeruginosa	NC_002516.2	GCCACGTGCGTAATATGC	TCCACATCCGCTGCATT

P. aeruginosa	NC_002516.2	ACCAGGGTACGCTCGTT	ACGTGATCCGCGGTATC
P. aeruginosa	NC_002516.2	GTCAGCGTGACCCGMAT	CGTTGCAGCAACTGCAT
P. aeruginosa	NC_002516.2	TGTAGCCGGCTAATTTCC	CGTCAGCCTTGCGATAG
P. aeruginosa	NC_002516.2	TAGGACTCCGCGGTCAT	GAAAGCCCAGCAATTGC
P. aeruginosa	NC_002516.2	TTGCGCCGACAGACTAC	CTCGGCGTGAAGGAAGA
P. aeruginosa	NC_002516.2	TCCAGCTCGACCACSGT	ATCGTCCTCGCGGTGAT
P. aeruginosa	NC_002516.2	ACGCTGCCTCCATAGTG	TAGTGGTGCCGCTTGAG
P. aeruginosa	NC_002516.2	CAGACGCGAGATTTCGT	AATCGCGATGGGCGATA
P. aeruginosa	NC_002516.2	ATGGCGCTCTCCAAGGA	TGCAGCGGTTTCAACAG
P. aeruginosa	NC_002516.2	ACCAGGGAAAGCTGGAA	ATGCCCTGGCCGTSGTY
P. aeruginosa	NC_002516.2	ACGAAGCTGCGCAGAAC	TCGTCGGACAGTTCGAA
P. aeruginosa	NC_002516.2	GTTCCCGTTCGGCTGAA	CGCCGATGGTGTAGTTG
P. aeruginosa	NC_002516.2	TCCARCTGGAAACCGCAAT	TGGCGGACTTCCGTTTC
P. aeruginosa	NC_002516.2	ACCATCCCCTTCAAGGA	TGGCCAAAGGGATCAGA
P. aeruginosa	NC_002516.2	AGCTTGGGATCGACGTA	AATGACGGCGTGAATGC
P. aeruginosa	NC_002516.2	TCCACTTTCGCGAACAG	TTCCGCCGTCGTAAGCA
P. aeruginosa	NC_002516.2	ATGACCCCGCACGATTC	AGCCGTCATGTCCGATT
P. aeruginosa	NC_002516.2	CAACGCGGTGACCCGTG	TGGACTTGGGGAACTGC
P. aeruginosa	NC_002516.2	CGGCACCGAACTGATCA	CTCGCCCTGGATCAGTA
P. aeruginosa	NC_002516.2	ACACCTCCAGGGTGTACTC	TAGGGATGGGCGTTTCG
P. aeruginosa	NC_002516.2	CACCCTGTCGACGGTAT	ARTCCTCCAGTCCGGTC
P. aeruginosa	NC_002516.2	GAGGGTGCCGTCGAAGT	GTGCTCGGCAAGTTCCA
P. aeruginosa	NC_002516.2	CTGGTCAAGCGGGTGAT	AACTCGTCCACCAGGAT
P. aeruginosa	NC_002516.2	TGATCGATCCGCAAGCG	GGTCGAAATCGGGTGAGA
P. aeruginosa	NC_002516.2	CAGGCTGACGCAAGCAT	TGCACCTGGGCATAGAC
P. aeruginosa	NC_002516.2	ACGATGAACACGGTGGC	AACTGCCCTGTGAAACG
P. aeruginosa [46]	NC_002516.2	GCCACACCTACATCGTCTAT	AGGTTGCCGAGGTTGTCCAC
P. aeruginosa [46]	NC_002516.2	ATGTCACCGTGCCGTTCAAG	TGAAGGCAGTCGGTTCCTTG
P. aeruginosa [46]	NC_002516.2	AGGTCGGTTCCTCCAAGGTC	GACGTTGTGGTGCGACTTGA
P. aeruginosa [46]	NC_002516.2	AGAAGACCGAGTTCGACCAT	GGTGCCATAGAGGAAGTCAT

P. aeruginosa [46]	NC_002516.2	ACGGCGAGAACGAGGACTAC	TGGCGGTCGGTGAAGGTGAA
P. aeruginosa [46]	NC_002516.2	GGTGACGACGGCAAGCTGTA	GTATCGCCTTCGGCACAGGA
P. aeruginosa [46]	NC_002516.2	TTCAACTTCGGCGACTTCCA	GGTGTCCATGTTGCCGTTCC
S. pneumoniae	NZ_CP020549.1	AATGTTGCTGGACGTGC	TAGTGTTTGTRTTGTGAAATAGCA
S. pneumoniae	NZ_CP020549.1	GTCAAGACCACCAGTAACG	TCGTGAGCATGTAGACACT
S. pneumoniae	NZ_CP020549.1	GCTACAATCGGTAGCACAA	TGTCGAGACTACTCGTCAT
S. pneumoniae	NZ_CP020549.1	AACTCGGGATGACTCTCT	CCCAACGTCTGGGAGTT
S. pneumoniae	NZ_CP020549.1	AGAATGTCAACTACCTCTCTCT	AACCAAGCGAAGCTAAGG
S. pneumoniae	NZ_CP020549.1	TCTAGCCCTGCTTAAACTAGT	CAAGCGTGTCCGTTACC
S. pneumoniae	NZ_CP020549.1	TCGTAACAACTGCTAACACT	GCCCCGTAACYCTACAAC
S. pneumoniae	NZ_CP020549.1	AACAAGACATACTMCCCTACAC	CTAAAAAGTTACTAAAAATAAAAAN
S. pneumoniae	NZ_CP020549.1	AGCTAACCGTGCATTCG	AACCCRGGCTAAAGTTAGTTAG
S. pneumoniae	NZ_CP020549.1	CCGTTCACACGCGATGA	AGCCCAAGTGTAAAAGCA
S. pneumoniae	NZ_CP020549.1	ATTGTTTGGGCACCTCTT	GGGGATATAGAAAACAAGACCC
S. pneumoniae	NZ_CP020549.1	TGAGCGTGTCAAATTCCA	AGAGTACAGCACCACTGT
S. pneumoniae	NZ_CP020549.1	CAGCGCTAGCTCACAGA	CAACTGATCCAGCCCAAT
S. pneumoniae [47]	NZ_CP020549.1	GCCTTTGAGGCGACAGC	TGCAGTTCARAAACATWTTCTAA
S. pneumoniae [47]	NZ_CP020549.1	ATGGACAAACCAGCNAGYTT	GCTTGAGGTCCCATRCTNCC
S. pneumoniae [47]	NZ_CP020549.1	GGCATTGGAATGGGATCACC	TCTCCCGCAGCTGACAC
S. pneumoniae [47]	NZ_CP020549.1	GCCAACTCAGGTCATCCAGG	TGCAACCGTAGCATTGTAAC
S. pneumoniae [47]	NZ_CP020549.1	TTATTCCTCCTGATTCTGTC	GTGATTGGCCAGAAGCGGAA
S. pneumoniae [47]	NZ_CP020549.1	TTATTAGAAGAGCGCATCCT	AGATCTGCCTCCTTAAATAC
S. pneumoniae [47]	NZ_CP020549.1	TGCYCAAGTTCCTTATGTGG	CACTGGGTRAAACCWGGCAT
S. aureus	NC_007795.1	GTCCAGGTAGCATGATT	TGTCATACCAGTTAGGAATCACA
S. aureus	NC_007795.1	AATTAAGTAAGCTCCAATGCGTT	TAGTTCGCTCTCCCCTTA
S. aureus	NC_007795.1	TCCAATATCCTGGCGTGA	TTCACAACCATTACCAAG
S. aureus	NC_007795.1	TAACGATGCGACAGGTACAG	ATGATGATGCTATGCGT
S. aureus	NC_007795.1	AGGTCTCACGACATCATT	CATAATACCTGCGCCATCA

S. aureus	NC_007795.1	CGGACAAGAACATGTCAC	TGAATTAGTCCCCTGCG
S. aureus	NC_007795.1	GATGCAGAATACTCAAATGT	CGCCGTTATCATAGAAGCTATAC
S. aureus	NC_007795.1	GGAGTTATCATGCCAACAGTTATA	GCGCTTCTTCTGCACCCAT
S. aureus	NC_007795.1	CTCACCTCTACTTCTTATGGA	CACATCCATGATTTCGTGCTCT
S. aureus	NC_007795.1	TACAAGCTAGGTCGAAAGAAATAC	CTTGTAACAACTTTAACTTAGCTT
S. aureus	NC_007795.1	TCACATCATTTAAATCGCCAC	GCGCAGGTAGATTAATGC
S. aureus	NC_007795.1	GCAACTTCCATCGAACCGACAGTC	TAATAGAGGAGTGTAGCACT
S. aureus	NC_007795.1	GACCTACAGCGATTCTT	TACACAAGCTCTGAATCG
S. aureus	NC_007795.1	GTGTCGAATTTTTATGCAT	AAACTTATAACTACGGCATTCTTTG
S. aureus	NC_007795.1	ATTAGGCGTTGTCGCA	AACAGGTATCTTATTGCATAATAT
S. aureus	NC_007795.1	ATCCTGACACTTACATTCCAT	CCACTGCCACCTGTTCG
S. aureus	NC_007795.1	CACCTACACGAATCATTGACAA	AGCAGGTGGTGCACTTATC
S. aureus	NC_007795.1	ACATTGAGCCATACTTCCA	TTAGAAGGTCCTCGAGA
S. aureus	NC_007795.1	TCCTGTCCAAGTGTAATGATAA	CGTTACAGCTACAATCGCTAC
S. aureus	NC_007795.1	AAGTTATGGACTTAGACATTCTA	TGGACAAGGTTGGTAATAGCGA
S. aureus	NC_007795.1	ACCATCTTCTATCGTTTTATCA	GCTGCAAGTATATTACATGA
S. aureus	NC_007795.1	TCAGCTGTGTTTCGTTGT	AATTAGGACAATACGTATATGAACA
S. aureus	NC_007795.1	TTTTCAAGGCTGTTTTTTATGCG	CGATTGTATGTGTGCCAATAATA
S. aureus [48]	NC_007795.1	TTGATTCACCAGCGCGTATTGTC	AGGTATCTGCTTCAATCAGCG
S. aureus [48]	NC_007795.1	ATCGGAAATCCTATTTCACATTC	GGTGTTGTATTAATAACGATATC
S. aureus [48]	NC_007795.1	CTAGGAACTGCAATCTTAATCC	TGGTAAAATCGCATGTCCAATTC
S. aureus [48]	NC_007795.1	ATCGTTTTATCGGGACCATC	TCATTAACTACAACGTAATCGTA
S. aureus [48]	NC_007795.1	GTTAAAATCGTATTACCTGAAGG	GACCCTTTTGTTGAAAAGCTTAA

| S. aureus [48] | NC_007795.1 | TCGTTCATTCTGAACGTCGTGAA | TTTGCACCTTCTAACAATTGTAC |
| S. aureus [48] | NC_007795.1 | CAGCATACAGGACACCTATTGGC | CGTTGAGGAATCGATACTGGAAC |

Citations

1.   Griffith MF, Levy CR, Parikh TJ, Stevens-Lapsley JE, Eber LB, Palat S-IT, et al. Nursing Home Residents Face
Severe Functional Limitation or Death After Hospitalization for Pneumonia. J Am Med Dir Assoc. 2020;21: 1879–
1884.

2.   Muder RR. Pneumonia in residents of long-term care facilities: epidemiology, etiology, management, and
prevention. Am J Med. 1998;105: 319–330.

3.   Marrie TJ. Pneumonia in the long-term-care facility. Infect Control Hosp Epidemiol. 2002;23: 159–164.

4.   Henig O, Kaye KS. Bacterial Pneumonia in Older Adults. Infect Dis Clin North Am. 2017;31: 689–713.

5.   Kruse RL, Mehr DR, Boles KE, Lave JR, Binder EF, Madsen R, et al. Does hospitalization impact survival after
lower respiratory infection in nursing home residents? Med Care. 2004;42: 860–870.

6.   Loeb M, Carusone SC, Goeree R, Walter SD, Brazil K, Krueger P, et al. Effect of a clinical pathway to reduce
hospitalizations in nursing home residents with pneumonia: a randomized controlled trial. JAMA. 2006;295: 2503–
2510.

7.   Dempsey CL. Nursing home-acquired pneumonia: outcomes from a clinical process improvement program.
Pharmacotherapy. 1995;15: 33S–38S.

8.   Gloth FM 3rd, Burton JR. Autopsies and death certificates in the chronic care setting. J Am Geriatr Soc. 1990;38:
151–155.

9.   Lim WS, Macfarlane JT. A prospective comparison of nursing home acquired pneumonia with community acquired
pneumonia. Eur Respir J. 2001;18: 362–368.

10.   Crnich CJ, Jump R, Trautner B, Sloane PD, Mody L. Optimizing Antibiotic Stewardship in Nursing Homes: A
Narrative Review and Recommendations for Improvement. Drugs & Aging. 2015;32: 699–716.

11.   Overview of Pneumonia. Goldman's Cecil Medicine. W.B. Saunders; 2012. pp. 587–596.

12.   Pathophysiology of Pneumonia. Clinics in Chest Medicine. 2005;26: 39–46.

13.   Scannapieco FA, Stewart EM, Mylotte JM. Colonization of dental plaque by respiratory pathogens in medical
intensive care patients. Crit Care Med. 1992;20: 740–745.

14.   Russell SL, Boylan RJ, Kaslick RS, Scannapieco FA, Katz RV. Respiratory pathogen colonization of the dental
plaque of institutionalized elders. Spec Care Dentist. 1999;19: 128–134.

15.   Didilescu AC, Skaug N, Marica C, Didilescu C. Respiratory pathogens in dental plaque of hospitalized patients with
chronic lung diseases. Clin Oral Investig. 2005;9: 141–147.

16.   El-Solh AA, Pietrantoni C, Bhat A, Okada M, Zambon J, Aquilina A, et al. Colonization of dental plaques: a
reservoir of respiratory pathogens for hospital-acquired pneumonia in institutionalized elders. Chest. 2004;126:
1575–1582.

17.   Hong C, Aung MM, Kanagasabai K, Lim CA, Liang S, Tan KS. The association between oral health status and
respiratory pathogen colonization with pneumonia risk in institutionalized adults. Int J Dent Hyg. 2018;16: e96–
e102.

18.   Quagliarello V, Ginter S, Han L, Van Ness P, Allore H, Tinetti M. Modifiable risk factors for nursing home-acquired
pneumonia. Clin Infect Dis. 2005;40: 1–6.

19.   Abe S, Ishihara K, Adachi M, Okuda K. Oral hygiene evaluation for effective oral care in preventing pneumonia in
dentate elderly. Arch Gerontol Geriatr. 2006;43: 53–64.

20. Chalmers J, Pearson A. Oral hygiene care for residents with dementia: a literature review. J Adv Nurs. 2005;52: 410–
419.

21. Grap MJ, Munro CL, Hamilton VA, Elswick RK Jr, Sessler CN, Ward KR. Early, single chlorhexidine application
reduces ventilator-associated pneumonia in trauma patients. Heart Lung. 2011;40: e115–22.

22. Hua F, Xie H, Worthington HV, Furness S, Zhang Q, Li C. Oral hygiene care for critically ill patients to prevent
ventilator-associated pneumonia. Cochrane Database Syst Rev. 2016;10: CD008367.

23. Sjögren P, Nilsson E, Forsell M, Johansson O, Hoogstraate J. A systematic review of the preventive effect of oral
hygiene on pneumonia and respiratory tract infection in elderly people in hospitals and nursing homes: effect
estimates and methodological quality of randomized controlled trials. J Am Geriatr Soc. 2008;56: 2124–2130.

24. Bassim CW, Gibson G, Ward T, Paphides BM, Denucci DJ. Modification of the risk of mortality from pneumonia
with oral hygiene care. J Am Geriatr Soc. 2008;56: 1601–1607.

25. Juthani-Mehta M, Van Ness PH, McGloin J, Argraves S, Chen S, Charpentier P, et al. A cluster-randomized
controlled trial of a multicomponent intervention protocol for pneumonia prevention among nursing home elders.
Clin Infect Dis. 2015;60: 849–857.

26. Russakoff B, Wood C, Lininger MR, Barger SD, Trotter RT, Maltinsky S, et al. A Quantitative Assessment of
Staphylococcus aureus Community Carriage in Yuma, Arizona. J Infect Dis. 2023;227: 1031–1041.

27. Abdeldaim GMK, Strålin K, Korsgaard J, Blomberg J, Welinder-Olsson C, Herrmann B. Multiplex quantitative PCR
for detection of lower respiratory tract infection and meningitis caused by Streptococcus pneumoniae, Haemophilus
influenzae and Neisseria meningitidis. BMC Microbiol. 2010;10: 310.

28. Welti M, Jaton K, Altwegg M, Sahli R, Wenger A, Bille J. Development of a multiplex real-time quantitative PCR
assay to detect Chlamydia pneumoniae, Legionella pneumophila and Mycoplasma pneumoniae in respiratory tract
secretions. Diagn Microbiol Infect Dis. 2003;45: 85–95.

29. Improved early diagnosis of Pseudomonas aeruginosa by real-time PCR to prevent chronic colonisation in a
paediatric cystic fibrosis population. J Cyst Fibros. 2011;10: 21–24.

30. Park HK, Lee HJ, Kim W. Real-time PCR assays for the detection and quantification of Streptococcus pneumoniae.
FEMS Microbiol Lett. 2010;310: 48–53.

31. Wood C, Sahl J, Maltinsky S, Coyne B, Russakoff B, Yagüe DP, et al. SaQuant: a real-time PCR assay for
quantitative assessment of Staphylococcus aureus. BMC Microbiol. 2021;21: 1–7.

32. Williamson CHD, Vazquez AJ, Nunnally AE, Kyger K, Fofanov VY, Furstenau TN, et al. ColiSeq: a multiplex
amplicon assay that provides strain level resolution of directly from clinical specimens. Microbiol Spectr. 2024;12:
e0413923.

33. Li H. Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM. 2013. Available:
http://arxiv.org/abs/1303.3997

34. Website. Available: R Core Team (2024). _R: A Language and Environment for Statistical Computing_. R
Foundation for Statistical Computing, Vienna, Austria. <https://www.R-project.org/>.

35. Bates D, Mächler M, Bolker B, Walker S. Fitting linear mixed-effects models Usinglme4. J Stat Softw. 2015;67.
doi:10.18637/jss.v067.i01

36. Kuznetsova A, Brockhoff PB, Christensen RHB. LmerTest package: Tests in linear mixed effects models. J Stat
Softw. 2017;82. doi:10.18637/jss.v082.i13

37. Gehlenborg N. A More Scalable Alternative to Venn and Euler Diagrams for Visualizing Intersecting Sets [R

496 package UpSetR version 1.4.0]. 2019 [cited 1 Dec 2024]. Available: https://CRAN.R-project.org/package=UpSetR

497 38. Nouwen JL, Fieren MWJA, Snijders S, Verbrugh HA, van Belkum A. Persistent (not intermittent) nasal carriage of
498  Staphylococcus aureus is the determinant of CPD-related infections. Kidney Int. 2005;67: 1084–1092.

499 39. de Vogel CP Boelens HA Verveer J Nouwen JL Verbrugh HA Wertheim HF van BAVN. Reclassification of
500  Staphylococcus aureus nasal carriage types. J Infect Dis. 2009. doi:10.1086/599119

501 40. Sutcliffe CG, Grant LR, Cloessner E, Klugman KP, Vidal JE, Reid R, et al. Association of Laboratory Methods,
502  Colonization Density, and Age With Detection of Streptococcus pneumoniae in the Nasopharynx. Am J Epidemiol.
503  2019;188: 2110–2119.

504 41. Porritt RA, Crother TR. *Chlamydia pneumoniae* Infection and Inflammatory Diseases. OT. 2016;7.
505  doi:10.1615/ForumImmunDisTher.2017020161

506 42. Cillóniz C, Civljak R, Nicolini A, Torres A. Polymicrobial community-acquired pneumonia: An emerging entity.
507  Respirology. 2016;21: 65–75.

508 43. Pearson T, Hepp C, Trotter RT, Mbegbu M, Russakoff B, Panisello YD, et al. Genome Sequences of Community
509  Carriage Strains of Staphylococcus aureus from Yuma, Arizona. Microbiology resource announcements. 2021;10.
510  doi:10.1128/MRA.00449-21

511 44. Meats E, Feil EJ, Stringer S, Cody AJ, Goldstein R, Kroll JS, et al. Characterization of encapsulated and
512  noncapsulated Haemophilus influenzae and determination of phylogenetic relationships by multilocus sequence
513  typing. J Clin Microbiol. 2003;41: 1623–1636.

514 45. Pannekoek Y, Morelli G, Kusecek B, Morré SA, Ossewaarde JM, Langerak AA, et al. Multi locus sequence typing
515  of Chlamydiales: clonal groupings within the obligate intracellular bacteria Chlamydia trachomatis. BMC Microbiol.
516  2008;8: 42.

517 46. Curran B, Jonas D, Grundmann H, Pitt T, Dowson CG. Development of a multilocus sequence typing scheme for the
518  opportunistic pathogen Pseudomonas aeruginosa. J Clin Microbiol. 2004;42: 5644–5649.

519 47. Enright MC, Spratt BG. A multilocus sequence typing scheme for Streptococcus pneumoniae: identification of
520  clones associated with serious invasive disease. Microbiology. 1998;144 (Pt 11): 3049–3060.

521 48. Enright MC, Day NP, Davies CE, Peacock SJ, Spratt BG. Multilocus sequence typing for characterization of
522  methicillin-resistant and methicillin-susceptible clones of Staphylococcus aureus. J Clin Microbiol. 2000;38: 1008–
523  1015.

524 49. Furstenau TN, Cocking JH, Sahl JW, Fofanov VY. Variant site strain typer (VaST): efficient strain typing using a
525  minimal number of variant genomic sites. BMC Bioinformatics. 2018;19: 222.