**DUSP6 is upregulated in metastasis and influences migration and metabolism in**

**pancreatic cancer cells**

Authors: Mariana T. Ruckert[1,2,3], R. McKinnon Walsh[2], Bailey A. Bye[2], Austin E. Eades[2], Wei

Yan[4], Filip Bednar[4,5], Jiaqi Shi[5,6], Costas A. Lyssiotis[3,5,7], Michael N. VanSaun[2#], Vanessa S.

Silveira[1,8#]

[1]Department of Genetics, Ribeirao Preto Medical School, University of Sao Paulo, Ribeirao Preto, SP,

14040-901, Brazil.

[2]Department of Cancer Biology, University of Kansas Medical Center, Kansas City, KS, 66160, US.

[3]Department of Molecular & Integrative Physiology, University of Michigan, Ann Arbor, MI, 48109,

US.

[4]Department of Surgery, University of Michigan, Ann Arbor, MI, 48109, US.

[5]Rogel Cancer Center, University of Michigan, Ann Arbor, MI, 48109, US.

[6]Department of Pathology & Clinical Labs, University of Michigan, Ann Arbor, MI, 48109, US.

[7]Department of Internal Medicine, Division of Gastroenterology and Hepatology, University of

Michigan, Ann Arbor, MI, 48109, US.

[8]Massachusetts General Hospital Cancer Center, Krantz Family Center for Cancer Research, Boston,

MA; Harvard Medical School, Boston, MA, 02114, US.

Running title: DUSP6 influences migration and glycolysis in PDAC cells.

Keywords: DUSP6; Migration; Glycolysis; Pancreatic cancer; Metastasis.

[#]Corresponding authors: vdasilvasilveira@mgh.harvard.edu, mvansaun@kumc.edu

**ABSTRACT**

Pancreatic ductal adenocarcinoma (PDAC) is a highly aggressive malignancy characterized by KRAS mutations in approximately 95% of cases. Despite recent advancements with KRAS inhibitors, therapeutic resistance has emerged, and combination approaches are needed. In particular, it is important to understand how downstream signaling of KRAS supports PDAC growth. For example, DUSP6, a dual-specificity phosphatase that modulates ERK1/2 phosphorylation and RAS pathway activity, has emerged as an important regulator of KRAS-MAPK signaling. Transcriptomic analyses demonstrate that DUSP6 is markedly overexpressed in PDAC tumors compared to non-tumoral pancreatic tissue. Single-cell RNA-seq reveals its upregulation in epithelial tumor cells, with further elevation in metastatic lesions relative to primary tumors. This upregulation correlates with the quasi-mesenchymal/squamous molecular subtype, and clinically, high DUSP6 expression is associated with poorer overall survival. Gene set enrichment analyses of metastatic samples indicate that DUSP6 is linked to pathways involved in cell migration and metabolism. To elucidate DUSP6's functional roles, stable knockdown of DUSP6 in PDAC cell lines resulted in increased ERK/MAPK activation and altered migratory capacity. Metabolic profiling showed enhanced basal glycolysis following DUSP6 suppression. However, combined inhibition of glycolysis and DUSP6 downregulation did not affect the migratory phenotype, indicating that glycolytic alterations do not drive migration. These findings highlight the dual and independent roles of DUSP6 in modulating migratory capacity and glycolysis in PDAC. This study underscores the significance of DUSP6 as a potential therapeutic target and provides new insights into its contributions to PDAC progression.

**SIGNIFICANCE STATEMENT:** DUSP6 plays dual and independent roles in pancreatic

cancer, regulating both migration and glycolysis. Its upregulation in metastasis is associated

with poor prognosis and more aggressive phenotypes, highlighting its clinical relevance.

Targeting DUSP6 represents a potential therapeutic strategy to disrupt KRAS-MAPK signaling

and target key pathways driving PDAC progression.

**INTRODUCTION**

Pancreatic ductal adenocarcinoma (PDAC) accounts for over 90% of pancreatic cancer cases

and is one of the most lethal forms of invasive cancer globally, ranking as the third leading

cause of cancer-related mortality [1-3]. Despite significant advancements, the current 5-year

survival rate remains low, at 13% [3]. Late detection of PDAC primarily stems from vague

symptoms, lack of early diagnostic markers, and resistance to chemotherapy [4, 5]. Complete

resection remains the sole curative option, yet it is unattainable for the majority of patients; 80–

90% of patients present with locally advanced or metastatic disease upon diagnosis [6, 7].

Metastasis in PDAC commonly affects the liver, peritoneum, and lungs [8-12]. Studies indicate

that not all circulating tumor cells successfully initiate secondary tumors, suggesting the

requirement for genetic and/or epigenetic changes [13]. Mutations such as those occurring in

*Kras* are prevalent in both primary and metastatic tumors, with certain passenger mutations

showing a preference for metastatic sites [11, 12]. For example, loss of DPC4 (SMAD4) is

more frequent in metastatic cases, correlating with increased invasiveness [14]. Gene

expression alterations are also important considerations in this context; however, to date, most

studies have only documented changes between non-neoplastic pancreatic and tumor tissues

[15].

*Kras* mutations are found in approximately 95% of PDAC cases [16] and are the first genetic

alterations that arise in precancerous lesions. These mutations alter protein conformation and

ultimately lead to a prolonged activation state and, consequently, downstream activation of

signaling pathways such as the ERK/MAPK pathway [17]. In tumor cells, ERK

phosphorylation drives MYC activation through subsequent phosphorylation, leading to a

myriad of effects such as uncontrolled cell proliferation, increased survival, and metabolic

rewiring [18, 19]. Although it is intuitive to believe that KRAS constitutive activation is

imperative for tumors, its hyperactivation induces oxygen reactive species (ROS) production,

and cells are forced to upregulate their antioxidant systems to maintain ROS homeostasis.

However, failure to buffer intracellular ROS accumulation can result in oncogene-induced

senescence and cytotoxicity [20]. Therefore, fine-tuned KRAS-MAPK activation is required

for tumor cells to survive and concurrently evade the detrimental effects of their metabolic

adaptations.

DUSP6 is a dual-specificity phosphatase that targets serine and threonine residues, particularly

dephosphorylating ERK1/2 and downregulating the ERK/MAPK pathway [21].  While DUSP6

is reported to act in a negative feedback loop to downregulate ERK/MAPK activation, it has

also been reported that ERK1/2 phosphorylates DUSP6 to target this protein for subsequent

proteasomal degradation [22]. Its role in cancer is context dependent, acting either as an

oncogene or as a tumor suppressor [23]. DUSP6 was previously shown to be upregulated in

pancreatic intraepithelial neoplastic (PanIN) lesions (early stage of PDAC tumorigenesis) and

downregulated in invasive carcinoma, particularly in poorly differentiated subtypes [24]. This

phenomenon has been reported to occur due to hypermethylation of *DUSP6* promoter, leading

to abrogation of gene expression and increased activity of the ERK/MAPK pathway [25]. The

complex coordination of the KRAS-ERK-DUSP6 axis and the timely regulation of DUSP6

expression throughout tumorigenesis prompted us to investigate the role of this molecule in

PDAC metastasis.

In this study, we demonstrated that *in silico* gene expression and RNAscope studies in human

samples revealed upregulation of *DUSP6* in metastatic samples compared to that in primary

tumors and non-neoplastic pancreatic tissue. Gene set enrichment analysis (GSEA) of human

metastatic PDAC samples revealed enrichment of pathways related to cell migration and

metabolism. *In vitro* assessments demonstrated *DUSP6* downregulation in a KRAS mutant

cell line, which correlates with increased ERK/MAPK activation, diminished proliferation, and

migration capacity; however, this effect was absent in KRAS wild-type cells. Functionally,

*DUSP6* downregulation led to elevated basal glycolysis, indicating its involvement in glucose

metabolism. However, the combined inhibition of *DUSP6* and glycolysis did not affect the

migratory phenotype of the KRAS mutant cell line. Our findings suggest that, while DUSP6

influences migration capacity and glucose metabolism, these effects may not be directly

correlated in this context.

## MATERIALS & METHODS

### Cell lines and culture

MIA    PaCa-2    (RRID:CVCL_0428),    PANC-1    (RRID:CVCL_0480),    AsPC-1

(RRID:CVCL_0152), BxPC-3 (RRID:CVCL_0186), Capan-2 (RRID:CVCL_0026), and

SW1990 (RRID:CVCL_1723) were purchased from the ATCC (Manassas, VA, USA). K8484

(RRID:CVCL_XD12) and DT8082 (RRID:CVCL_XD11) were acquired from David

Tuveson's laboratory (Cold Spring Harbor Laboratory) [20]. P4313 was acquired from Andy

Lowy (UC San Diego) [21]. PKT62 is a murine pancreatic tumor cell line derived from a PKT

(*LSL-Kras*$^{G12D/+}$, *TGFBR2*$^{fl/fl}$, *Ptf1a*$^{Cre/+}$) mouse [26] and kindly provided by Michael VanSaun

(University of Kansas Medical Center). KPC CAFs and PKT CAFs cell lines were isolated

from the respective mouse tumors and validated by western blotting. Briefly, tumors were

collected, digested, minced and cells were sorted with EpCAM beads (BioLegend Inc., cat

#741444). EpCAM$^-$ cells were kept in culture and until full confluence, protein samples were

isolated and fibroblastic phenotype was confirmed by the presence of α-SMA and absence of

CK19 and Ras$^{G12D}$. Capan- 2, AsPC-1 and SW1990 were cultured in high glucose (4.5 g/L)

RPMI-1640 medium (ThermoFisher Scientific, Waltham, MA USA cat #A1049101)

supplemented with 10% heat-inactivated fetal bovine serum (FBS, Atlanta Biologicals/RD

Systems, Atlanta, GA USA) and antibiotic-antimycotic (Thermo Fisher Scientific cat#

15240062). The K8484 cell line was cultured in high-glucose (4.5 g/L) Dulbecco's Modified

Eagle Media (Thermo Fisher Scientific, cat# 11-995-073) supplemented with 5% heat-

inactivated FBS (R&D Systems-Atlanta Biologicals) and antibiotic-antimycotic (Thermo

Fisher Scientific). All other cell lines were maintained in high-glucose DMEM supplemented

with 10% FBS and antibiotic-antimycotic. Cells were maintained in culture conditions (37°C,

5% $CO_2$ and humid atmosphere) as recommended by the ATCC, utilized within 15 passages

from purchase and regularly tested for mycoplasma.

**Plasmids and gene knockdown**

To generate *DUSP6* knockdown cells, short hairpin RNA (shRNA) constructs targeting three

independent regions of DUSP6 mRNA (#1 5'-CTGTGGTGTCTTGGTACATTG-3'; #2 5'-

TCTAATCCAAAGGGTATATTT-3,' #3 5'-ATTCGGCATCAAGTACATCTT-3') were

obtained from Millipore (Burlington, MA, USA). The empty vector pLKO.1 puro

(RRID:Addgene_8453) was used as a control. Packaged lentiviruses were generated by

transfecting HEK293T (RRID:CVCL_0063) cells with the respective plasmids and the Trans-

Lentiviral™ Packaging Mix (Open Biosystems) to produce viral supernatants, which were

subsequently concentrated with Lenti-X™ concentrator (TakaraBio, cat #631231). Next,

$2.5 \times 10^5$ pancreatic cancer cells were seeded in 6-well plates and incubated for 24h with

lentiviral suspension. Successfully transduced cells were selected with puromycin treatment

(1-2.5µg/mL) until complete cell death was achieved in the non-transduced cells.

**RNA isolation and RT-qPCR**

Briefly, $10^6$ cells were seeded in 6-well plates and incubated overnight. The next day, cells

were washed with 1X PBS and harvested using QIAzol Lysis Reagent (Qiagen, GmbH, cat

#79306). Samples were stored at -80°C until total RNA was isolated using the Direct-zol RNA

Miniprep kit (Zymo Research Corporation), according to the manufacturer's protocols. RNA

quantification and quality analyses were performed using NanoDrop 2000 (Thermo Fisher

Scientific Inc.). For cDNA, 1.5-2µg of RNA was processed using the High-Capacity cDNA

Reverse Transcription Kit (Applied Biosystems, cat #4368814) according to the

manufacturer's instructions. The product was further diluted 1:10 before storage at -20°C.

Real-time PCR analysis was carried out with the 2X SYBR Green qPCR Master Mix (ApexBio

Technology, cat #K1070) on a ViiA 7 Real-Time PCR System (Applied Biosystems) using the

following primers: human *DUSP6* (Fwd: ATGGTAGTCCGCTGTCCAAC/Rev:

ACGTCCAAGTTGGTGGAGTC), mouse *Dusp6* (Fwd: TGTTTGAGAATGCGGGCGAGTT

/ Rev: ACAGTTTTTGCCTCGGGCTTCA), human *ACTB* (Fwd:

TCGTGATGGACTCCGGTGAC/Rev: CGTGGTGGTGAAGCTGTAG), and mouse *Actb*

(Fwd: GTGACGTTGACATCCGTAAAGA / Rev: GCCGGACTCATCGTACTCC). Each

sample was run in triplicate using the $2^{-\Delta\Delta CT}$ method (Livak & Schmittgen, 2001) and the fold-

change was evaluated relative to normal samples and determined using *ACTB/Actb* levels as a

reference.

**Western blotting**

Briefly, $10^6$ cells were plated in 6-well plates, incubated overnight, and collected the next day

for protein isolation. Cells were washed with 1X PBS, lysed in ice-cold RIPA buffer (1X RIPA

[Cell Signaling Technology]; 10mM NaF; 1mM PMSF), sonicated, and incubated on ice for

10 min. The cells were then centrifuged at $16,000 \times g$ for 10 min at 4°C. Protein quantification

was performed using the Pierce™ BCA Protein Assay Kit (Thermo Fisher Scientific),

following the manufacturer's instructions. For western blotting assays, 10 µg of total protein

was loaded in each lane of a 10% polyacrylamide gel and subjected to electrophoresis. Proteins

were transferred to nitrocellulose membranes using the semi-dry method on a Trans-Blot Turbo

Transfer System (Bio-Rad). Specific proteins were detected using the following primary

antibodies: P-ERK1/2 (Cell Signaling Technology, cat #4370, RRID:AB_2315112, link),

ERK1/2 (Cell Signaling Technology, cat #4695, RRID:AB_390779, link), E-cadherin (Cell

Signaling Technology, cat #3195, RRID:AB_2291471, link), N-cadherin (Cell Signaling

Technology, cat #13116, RRID:AB_2687616, link), and β-actin (Cell Signaling Technology,

cat #12262, RRID:AB_2566811, link) and MKP-3 (Santa Cruz Biotechnology, cat #sc-

137246, RRID:AB_2095347, link). Secondary antibodies used were Peroxidase AffiniPure

F(ab')₂ Fragment Donkey Anti-Mouse IgG (H+L) (cat #715-036-151, RRID:AB_2340774,

link) and Donkey Anti-Rabbit IgG (H+L) (cat #711-036-152, RRID:AB_2340590, link) from

Jackson ImmunoResearch Laboratories Inc. Images were obtained using FluorChem M (Bio-

Techne) or a Sapphire FL Biomolecular Imager (Azure), and analyzed using ImageJ software

(NIH, Bethesda, MD, USA).

**Immunohistochemistry staining**

Healthy pancreata, primary tumor, and metastatic samples were collected from wild-type

C57BL/6J (RRID:IMSR_JAX:000664) mice, *LSL-Kras*$^{G12D/+}$, *Ptf1a*$^{Cre/+}$ (KC), *LSL-Kras*$^{G12D/+}$,

Tp53$^{R172H/+}$, *Ptf1a*$^{Cre/+}$ mouse pancreatic tissue (KPC) maintained in a C57BL/6J background.

Male and female mice were used for experiments and mice ages varied from 5.9 to 86.6 weeks.

Samples were collected, embedded in paraffin, and sectioned into 4μm sections. The paraffin

sections were processed as follows: 15 min in xylene (2x); 5 min in 100% ethanol (2x) and 5

min in 75% ethanol. Finally, the sections were washed 3x in PBS at room temperature (RT).

Antigen retrieval was performed by boiling the sections in 10 mmol/L sodium citrate solution

for 10 min and waiting for the sections to cool down and achieve RT. Next, the sections were

washed 3x with PBS, quenched with 3% $H_2O_2$ for 15 min, and blocked with 5% BSA for 20

min. Whole-slide scanning was performed using a PANNORAMIC Scan II scanner

(3DHISTECH). Immunohistochemical staining was performed by iHisto Inc. (Salem MA;

iHisto.io) using an anti-DUSP6 antibody (Abcam, cat #ab76310, RRID:AB_1523517, link).

Ten fields per independent section from each condition were randomly selected for

quantification using the QuPath software [27]. The study weas performed under protocols

2005N000148 and 2019N000116 approved by the IACUC at MGH, and at Pharmaron Inc. and

Champions Oncology under approved protocols.

**RNA *in situ* hybridization**

RNA in situ hybridization was performed using the RNAscope® technology (Advanced Cell

Diagnostics, Inc.). Briefly, tumors were collected from the mice, sectioned, and immediately

fixed in 10% buffered formalin for 24h. Tumor sections were preserved in 70% ethanol,

embedded in paraffin blocks, cut into 5 μm sections, and placed in pre-treated slides. The

sections were freshly cut for the assay and dried at RT to preserve RNA integrity. Sections

were baked for 1h at 60°C and then deparaffinized using Histo-Clear (Electron Microscopy

Science, cat #6411004) and 100% ethanol. Next, sections were treated with RNAscope®

Hydrogen Peroxide for 10 min at RT, followed by a target retrieval step with RNAscope® 1X

Target Retrieval Reagent for 15 min, and finally, RNAscope® Protease Plus treatment for 30

min at 40°C. Finally, we used the RNAscope® 2.5 HD Duplex Reagent Kit, following the

manufacturer's protocol. Specific probes used in this assay were Mm-Dusp6 (cat #429321-C1),

Mm-Krt19-C2 (cat #402941- C2) and Mm-Pdgfrb-C2 (cat #411381-C2). The sections were

imaged at 20x magnification using an EVOS™ M5000 Imaging System (Thermo Fisher

Scientific).

**Clinical samples**

220   All the human patient samples described in this study were collected from patients undergoing

221   pancreas resection due to pancreatitis, cystic neoplasms or PDAC from 2002 to 2015 at the

222   University of Michigan Health System. Collection of these samples was approved by the

223   Institutional Review Board at the University of Michigan (IRB number: HUM00098128).

**RNA *in situ* hybridization combined with immunofluorescence multiplex staining**

Tissue microarray (TMA) sections were obtained, set up and processed as described in [28].

Fluorometric in-situ hybridization was performed according to the manufacturer's (Advanced

Cell Diagnostics, Inc.) instructions. Briefly, FFPE preserved TMAs were cut in 5 µm sections,

and placed in pre-treated slides. Sections were freshly cut for the assay and dried at RT to

preserve RNA integrity. Sections were baked for 1h at 60°C, deparaffinized in xylene three

times for 15 mins, washed in 100% ethanol twice for 3 mins and air dried. Slides were treated

with pretreatment 1, 2 and 3 buffers, rinsed in double-distilled water (ddH$_2$O) and incubated

with the DUSP6 (cat #1032061-C1) or control probes for 2h at 40°C in a humidity chamber.

Slides were then treated with Amp solutions 1-6 before the fluorogenic substrate Opal 690 TSA

(Akoya Biosciences, Inc.) was applied to the slide. After a 10 min incubation, the slides were

washed and then subjected to dual immunofluorescent staining on a Ventana Discovery Ultra

stainer (Roche Diagnostics, North America) with smooth muscle actin (BioCare Medical, cat

#001, RRID:AB_2910606, link, 1:100, 60 mins, Opal 570) and pan-cytokeratin (Abcam, cat

#ab9377, RRID:AB_307222, link, 1:200, 20 mins, Opal 520). Slides were mounted using

Prolong Gold with DAPI and scanned using a Vectra Polaris™ (Akoya Biosciences, Inc.)

whole slide scanner. Images were analyzed using the de-array TMA tool in QuPath and DAPI

was used to identify individual cells in each core.

**Proliferation assay**

Cell proliferation was assessed by ethynyldeoxyuridine (EdU) incorporation. Briefly, 2.5-

$5 \times 10^4$ cells were seeded in 24-well plates in complete media and incubated overnight. The next

day, the media was replaced with the appropriate treatments and incubated for 18 h. At this

point, EdU was added to the wells at a final concentration of 10μM and cells were incubated

for an additional 6h. At the end of this period, the cells were trypsinized, harvested, and fixed

with 10% buffered formalin overnight at 4°C with gentle agitation. The next day, the cells were

washed with PEB-T (1x PBS, 2mM EDTA, 1% BSA, 0.5% Triton-X) and subsequently with

PB (1x PBS, 1% BSA). Next, the cells were resuspended in Click-It Reaction ($H_2O$, 100mM

Tris pH 8.5, 1mM $CuSO_4$, 1μM azide-dye) and 100mM ascorbic acid. Cells were incubated

with 0.5μg/mL propidium iodide (PI), washed twice, and resuspended in PEB (1x PBS, 2mM

EDTA, 1% BSA) for flow cytometry analysis. The cells were analyzed for 530 nm and 695 nm

emission using an Attune NxT Flow Cytometer (Thermo Fisher Scientific). FCS files were

analyzed with FlowJo™ v10.8 Software (BD Biosciences) and the percentage of EdU-positive

cells was taken from the PI-positive single cells.

**Migration assay**

First, $2 \text{-} 4 \times 10^5$ cells were seeded into 24-well plates and cultured until they reached full

confluence. Then, a scratch was made at the center of the plate using a 200μL pipette tip. Wells

were washed twice with 1X PBS to discard all debris, and media with appropriate treatments

were added. The plates were photographed at 0 and 24h using an EVOS™ M5000 Imaging

System (Thermo Fisher Scientific). Scratch area was calculated using the Wound Healing

Coherency Tool (Montpellier Ressources Imagerie) on ImageJ (NIH, Bethesda, MD,

RRID:SCR_003070, link) and results are expressed as healing percentage – the difference in

scratch area between time 0 and time 24.

**In silico analysis**

*DUSP6* expression analysis was performed using publicly available datasets obtained from The

Cancer Genome Atlas (TGCA_PAAD), Genotype-Tissue Expression (GTEx, normal

pancreatic tissue) and Gene Expression Omnibus (GEO, RRID:SCR_005012, link; GSE62452

[29], GSE28735 [30], GSE15471 [31], GSE71729 [32] and GSE224564 [33]).   Overall

survival analysis was performed by Gene Expression Profiling Interactive Analysis (GEPIA,

RRID:SCR_018294, link) using the UCSC Xena project (http://xena.ucsc.edu) datasets.

Differential expression of *DUSP6* in the epithelial and stromal compartments was performed

using GSE93326 [34] with the R2 Genomics Analysis and Visualization Platform. The latter

was also utilized for accession of Bailey *et al.* [35] dataset since ICGC Data Portal Repository

was discontinued. Finally, *DUSP6* expression was correlated with Metastatic Potential

(MetMap 500 [36]) at the DepMap Portal using the Cancer Cell Line Encyclopedia (CCLE).

We specifically filtered the cell lines for Exocrine Adenocarcinoma, which corresponds to

PDAC cell lines. Gene set enrichment analysis was conducted in a cohort of metastatic samples

from the publicly available dataset GSE71729 [32] using the hallmark gene set collection

(version 2024) available at GSEA 4.3.3. software (RRID:SCR_003199, link) [37]. Single-cell

analysis was performed using a previously published [38] pancreatic tissue single-cell atlas

(link).

**Glycolysis stress test**

Glycolysis rates were determined using the Seahorse XF Glycolysis Stress Test Kit (Agilent,

Santa Clara, CA, USA), following the manufacturer's protocol. Briefly, 2-4x10$^4$ cells were

seeded into a 96-well plate and incubated overnight. The next day, cells were washed twice

with Seahorse XF DMEM (pH 7.4 (2mM glutamine) and a final volume of 180μL of the same

was added to each well. The extracellular acidification rate (ECAR) was measured (mpH/min)

at 11 time points with sequential injections of 10mM Glucose, 1μM Oligomycin and 100mM

2-Deoxy-D-glucose (Thermo Scientific, cat. #AC111980010). For nuclear staining, 5μg/mL

Hoechst was added to the oligomycin solution. The ECAR values were calculated and

normalized to the cell count number using an automated imaging and cell counting workflow.

For normalization, the ECAR value per $10^3$ cells was set, and the Agilent Seahorse XF system

automatically calculated and generated normalized results.

**Lactate assay**

For lactate measurement, $2.5 \times 10^4$ cells were seeded in 96-well plates and allowed to sit

overnight in the incubator. The next day, the complete media in the plate was replaced with

new media containing 0.5% FBS, and the cells were incubated for 24h. At the end of the

experiment, culture plates were centrifuged at $450 \times g$ for 5 minutes and 10μL of media was

collected from each well for the subsequent assay. For lactate quantification, we utilized the

Glycolysis Cell-Based Assay Kit (Cayman Chemical Company, Ann Arbor, MI, USA, cat

#600450), following the manufacturer's protocol. The absorbance was read at 490 nm using a

Synergy H1 Multimode microplate reader (Bio-Tek/Agilent Technologies).

**Statistical analysis**

Sample sizes were chosen to achieve a minimum of triplicate samples for all the experiments.

For the assessment of statistical significance between two groups, Welch's t-test was used for

parametric data, while Mann-Whitne test was used for non-parametric data. For comparison

across multiple groups, Brown-Forsythe and Welch ANOVA test or mixed-effects analysis

with Šídák's multiple comparisons test were used when appropriate for parametric data; for

non-parametric data, Kruskal-Wallis test were performed. Data normality was determined

based on Q-Q plots distribution. All statistical tests were performed using GraphPad Prism 9

software (RRID:SCR_002798, link), and the results were considered significant at α = 0.05. *

P < 0.05, ** P <0.01, *** P < 0.001, and **** P < 0.0001.

**RESULTS**

*DUSP6* **overexpression is correlated with advanced stage PDAC.**

To investigate the role of DUSP6 in metastasis, we first confirmed its expression in tumor

samples. To evaluate *DUSP6* expression in PDAC samples, we used three distinct in silico

datasets available on public platforms (GSE62452 [29], GSE28735 [30] and TCGA/GTEx).

Our analysis revealed that *DUSP6* was upregulated in primary tumor samples compared to non-

tumoral pancreatic tissue across all analyzed datasets (Figure 1A-C). Based on these findings,

we aimed to assess *DUSP6* expression in metastatic cases. Utilizing the *in silico* dataset

GSE71729 [32], we compared *DUSP6* expression across a cohort comprising 134 non-tumor

tissue samples, 145 primary tumor samples, and 62 metastatic samples derived from various

secondary sites. Notably, we observed that *DUSP6* was significantly overexpressed in

metastatic samples compared to both primary tumor samples (P < 0.0001) and non-tumorous

pancreatic tissue (Figure 1D). Considering that *DUSP6* is overexpressed in primary tumors and

metastatic samples, we hypothesized that there is a possible correlation between DUSP6

expression and patient prognosis. To answer this question, we first assessed its expression in

the PDAC molecular subtypes. We performed *in silico* analysis utilizing PDAC molecular

subtyping [35, 39]. We took advantage of the datasets from Badea *et al.* (GSE15471) and

Bailey *et al.* [35] and analyzed *DUSP6* expression in distinct PDAC subtypes. In the Badea *et*

*al.* cohort (Figure 1E), we observed that *DUSP6* was overexpressed in the quasi-mesenchymal

subtype compared to non-tumoral pancreatic tissue (P < 0.0001) and the exocrine-like subtype

(P = 0.0211). Likewise, in the Bailey *et al.* cohort (Figure 1G), we observed DUSP6

overexpression in the squamous subtype compared to that in the pancreatic progenitor (P =

0.0065) and immunogenic (P = 0.0197) subtypes. Next, we performed the same analysis using

another subtyping paradigm [32]; however, no differences were observed (Figure 1F). The

same subtyping was used to stratify samples in the GSE224564 [33] (Figure 1H) dataset, and

we observed overexpression of *DUSP6* in the basal-like subtype compared to the classical

subtype (P = 0.0292). The quasi-mesenchymal, squamous, and basal-like subtypes share

similar molecular signatures and worse prognoses than the other PDAC subtypes. Collectively,

these results suggest that DUSP6 is enriched in this tumor phenotype. To establish a link

between DUSP6 expression and ERK1/2 activation, we performed a correlation analysis

between *DUSP6* expression and *MAPK1* expression in quasi-mesenchymal samples from

GSE15471 [31]. We observed a moderate positive correlation in this context (R = 0.6787, P =

0.02), suggesting that DUSP6 overexpression may be required to counteract MAPK signaling

(Figure S1A).

Supporting the latter, our analysis of TCGA dataset revealed that high *DUSP6* expression in

PDAC patients correlated with worse overall survival (P = 0.0058; Figure 1I). Despite the

counterintuitive nature of *DUSP6* overexpression in metastatic samples, given the prevailing

dogma suggesting its negative regulation of the ERK/MAPK pathway, our findings were

substantiated by further correlation analysis conducted with 30 PDAC cell lines from the

Cancer Cell Line Encyclopedia (CCLE) and the Metastasis Map (MetMap 500 [36]). This

analysis revealed a moderate positive correlation between *DUSP6* expression and metastatic

potential in these cell lines (R = 0.611; P = 0.0003; Figure 1J). Furthermore, this correlation

analysis indicated that *DUSP6* gene expression ranked as the eighth strongest positive

correlation for metastatic potential within this PDAC cell line cohort among a list of 1000 genes

(Figure 1K). Collectively, these observations suggest that *DUSP6* overexpression in PDAC

samples is associated with the poorest prognostic outcomes and correlates with progression to

metastasis.

To validate these findings, we assessed DUSP6 expression in lesions across different stages of

the disease using healthy murine pancreatic tissue, primary tumors, and liver metastasis

samples. We observed a trend of temporal control of DUSP6 expression with disease

progression, although the results were not statistically significant (Figure 2A). DUSP6

expression is upregulated during the acinar-to-ductal metaplasia (ADM) process and

subsequently reduced in pancreatic intraepithelial neoplastic (PanIN) lesions. DUSP6 was

again increased in primary tumor lesions and, finally, in liver metastasis samples. These results

suggest that temporal control of DUSP6 expression is differentially utilized during PDAC

tumorigenesis and progression, perhaps to fine-tune MAPK signaling.

PDAC has a highly complex microenvironment and is mainly composed of dense

fibroinflammatory stroma consisting predominantly of activated stromal fibroblasts and

immunosuppressive immune cell types. Our previous analyses were primarily performed on

bulk tumors, where gene expression represents a composite of myriad cell types in the tumor

microenvironment. Accordingly, we assessed whether DUSP6 overexpression was localized to

specific cell types within the tumor. To answer this question, we examined *DUSP6* expression

in different tumor compartments using GSE93326 [34]. Indeed, our analysis revealed that

*DUSP6* was significantly upregulated in tumor cells compared to stromal cells (P < 0.0001;

Figure S1B). We validated this analysis by assessing DUSP6 protein levels in four mouse

PDAC cell lines and two mouse cancer-associated fibroblast (CAF) cell lines. The mouse cell

lines exhibited varying levels of the protein, with DT8082 and K8484 demonstrating the

highest DUSP6 levels, P4313 showing intermediate levels, and PKT62 displaying the lowest

levels (Figure S1C). Subsequently, we investigated *Dusp6* expression in mouse tumor tissue

using RNA in situ hybridization (RNA-ISH) technology. We examined *Dusp6* mRNA levels

in tumor sections obtained from *LSL-Kras*[G12D/+], *Ptf1a*[Cre/+] (KC), *LSL-Kras*[G12D/+], Tp53[R172H/+],

*Ptf1a*[Cre/+] (KPC) mice, co-stained for either *Krt19* (a tumor epithelial cell marker) or *Pdgfrb* (a

stromal marker). As depicted in Figure 1SC, *Dusp6* (green dots) exhibited strong co-expression

with *Krt19* (red dots, upper images), but not with *Pdgfrb* (red dots, lower images), in both

genetic backgrounds. Consistent with the western blot analysis of protein levels, both KC and

KPC pancreata demonstrated *Dusp6* overexpression in malignant epithelial cells, whereas

stromal compartment expression remained low (Figure S1D). To further explore the link

between DUSP6 expression and MAPK signaling, we leverage GSE93326 [34] to perform a

correlation analysis between *DUSP6* and *MAPK1* in the epithelial and stromal compartments.

Interestingly, we observed a moderate positive correlation in gene expression levels in the

epithelial compartment ($R = 0.6105$, $P < 0.0001$) but not in the stromal compartment ($P =$

$0.0321$, $P = 0.789$) in this cohort (Figure S1E-F). Altogether, these data indicate that DUSP6

is upregulated in PDAC epithelial cells in the context of hyperactivated MAPK signaling.

To confirm whether these observations could be translated into human disease, we used a panel

of human samples from two TMAs containing normal pancreatic tissue/pancreatitis, intraductal

papillary mucinous neoplasms (IPMN), mucinous cystic pancreatic neoplasms (MCN), PanIN,

PDAC, and PDAC metastasis. We differentially stained tissue compartments with duplex

RNA-ISH specific markers using panCK for epithelial cells and αSMA for CAFs (Figure 2C).

A general assessment of *DUSP6* positive cells throughout the cores revealed a trend similar to

that observed in mouse tissue. Although not statistically significant, we observed a slight

decrease in expression in PDAC primary tumors compared to that in normal pancreatic tissue

and a subsequent increase in metastatic samples (Figure 2D). Furthermore, we observed a

significant increase in *DUSP6* expression in epithelial cells throughout tumor progression, and

once again, a notable decrease in *DUSP6* expression in PDAC primary tumors, followed by

upregulation in metastatic samples (Figure 2E). These findings suggest that *DUSP6* timely

upregulation is required for metastatic tumor cells in the foreign environment.

The observation that *DUSP6* expression is significantly lower in PDAC primary tumor cells is

counterintuitive, considering what we previously observed in bulk RNA-seq datasets.

Considering these previous observations and the complexity of the PDAC microenvironment,

we decided to dig further into a publicly available pancreatic tissue single-cell atlas [38] to

characterize *DUSP6* expression patterns in different cell types within the healthy donor

pancreas, adjacent normal pancreatic tissue, and primary tumor microenvironment. Figure 2F

depicts the UMAP plot of all the cell types identified in the bulk of human patient samples.

Interestingly, we observed that regardless of the context, DUSP6 was majorly expressed in

myeloid cells, with increased expression in macrophages (Figures 2G-I, Figure S2A).

Nevertheless, when we look specifically at the epithelial cells, in these three contexts, we

observe a 2.4-fold increase in the percentage of *DUSP6* expressing epithelial cells in PDAC

compared to healthy pancreata (Figure S2A). Lastly, we performed co-expression analysis of

*DUSP6* and *KRT19* and confirmed a 1.8 × increase in *DUSP6* positive cells in PDAC *KRT19*

positive cells compared to healthy pancreata (Figure 2J). These results support *DUSP6*

upregulation in epithelial cells within the tumor microenvironment. Interestingly, *in silico*

analysis utilizing a bulk RNAseq dataset from Bojmar *et al.* [40] reveals that *DUSP6* is

upregulated in liver biopsy samples from patients who eventually developed PDAC liver

metastasis in comparison to those who did not in a 3-year follow up study (Figure S2B). Our

results also indicate a role for DUSP6 in the immune cell compartment, which may explain the

variations in DUSP6 expression observed in different PDAC human samples when evaluated

through bulk RNA sequencing. Additionally, DUSP6 expression may influence the

establishment of the pre-metastatic niche.

**433**  ***DUSP6* expression in metastatic samples correlates with decreased oxidative**

**434**  **phosphorylation and increased glycolysis.**

**435**  To gain further insight into the functional significance of *DUSP6* overexpression in metastatic

**436**  samples, we aimed to elucidate the pathways involved. To accomplish this, we revisited the

**437**  GSE71729 [32] dataset and conducted gene set enrichment analysis using 62 metastatic

**438**  samples. Intriguingly, our analyses revealed enrichment of several pathways associated with

**439**  cell motility, epithelial-to-mesenchymal transition, and metabolism. Notably, these

**440**  enrichments revealed a positive correlation between *DUSP6* expression and cell migration

**441**  (Figure 3B), suggesting an upregulation of this process, along with a negative correlation

**442**  between its expression and oxidative phosphorylation (Figure 3A), suggesting a

**443**  downregulation of this pathway. Supporting these observations, Panther Pathway enrichment

**444**  analyses conducted across two independent datasets (TCGA_PAAD and CPTAC_PDAC)

**445**  highlighted Glycolysis (P00024) as one of the pathways most strongly correlated with *DUSP6*

**446**  expression in PDAC across all three datasets (Table S1). Moreover, other enrichment analyses

**447**  utilizing the CPTAC Protein dataset indicated a correlation between DUSP6 protein levels and

**448**  the Pentose phosphate pathway (P02762), Glucose 6-phosphate metabolic process

**449**  (GO:0051156), and Carbon metabolism (hsa01200) (Table S1). These findings prompted us to

**450**  hypothesize that *DUSP6* overexpression may play a pivotal role in metabolic reprogramming

**451**  of tumor cells during metastasis.

**452**  ***DUSP6* downregulation reduces proliferation and migration in PDAC cell lines.**

**453**  Considering *DUSP6* upregulation in metastatic samples, we performed *DUSP6* knockdown in

**454**  PDAC cell lines exhibiting the highest levels of the protein, one of which was derived from

**455**  metastatic ascites (AsPC-1, Figure S3A). *DUSP6* knockdown was confirmed through gene

**456**  expression and protein level analyses (Figure 3D, S3). As expected, *DUSP6* knockdown led to

increased activation of ERK1/2, as confirmed by enhanced phosphorylation on western blotting

(Figure 3D). Subsequently, considering that ERK1/2 activation elicits a proliferative

phenotype, we evaluated cell proliferation using ethynyldeoxyuridine (EdU) incorporation

assays under conditions of full serum (5%) and serum deprivation, designed to either stimulate

or inhibit proliferation. Surprisingly, both AsPC-1 and BxPC-3 cell lines exhibited impaired

proliferation after *DUSP6* knockdown. Notably, AsPC-1 cells displayed a more pronounced

decrease, particularly under serum-starved conditions (P < 0.0001; Figure S4A). While BxPC-

3 cells also showed significant proliferation impairment (P < 0.0001), they were less affected

by serum starvation compared to AsPC-1 cells (Figure S4B).

To confirm our prior observations in human samples, we investigated the impact of *DUSP6*

knockdown on cell migratory capacity. *In vitro* migration assays conducted within 24h of

scratching (time 0) revealed distinct phenotypes among the cell lines. AsPC-1 cells exhibited

a significant reduction in migratory capacity upon DUSP6 knockdown (P < 0.0001; Figure

S4C), whereas BxPC-3 cells showed no significant effect (Figure S4D).

In summary, our findings suggest that DUSP6 plays a role in regulating both proliferation and

migration capacity of PDAC cells, albeit with potential context-dependent differences.

***DUSP6* downregulation increases glycolytic flux in PDAC cell lines.**

To ascertain DUSP6's involvement in tumor cell metabolism, we first evaluated basal lactate

secretion in extracellular media following *DUSP6* knockdown using a colorimetric assay. In

this context, distinct phenotypes were observed in both cell lines: BxPC-3 cells exhibited a

significant increase in lactate secretion compared to the control (P < 0.0001), whereas AsPC-1

cells showed no significant difference compared to the control (Figure 3F).

Considering that glucose is the primary source of energy for tumor cells *in vitro*, we sought to

delineate how DUSP6 affects glucose metabolism in each cell line and conducted a metabolic

assay to measure glycolysis activity in the absence of *DUSP6* (Figure 3E). Upon *DUSP6*

knockdown and glucose injection, both AsPC-1 and BxPC-3 cells demonstrated a statistically

significant increase in basal glycolysis (Figure 3G), consistent with observations from the

lactate secretion assay. Following treatment with oligomycin, an inhibitor of ATP synthase

(complex V), which shuts down mitochondrial respiration and makes cells dependent on

glycolysis for ATP production, we observed a significant increase in glycolytic capacity in

AsPC-1 cells upon *DUSP6* knockdown compared to the control ($P < 0.05$; Figure 3H), with no

significant differences in glycolytic reserve (Figure 3I). This indicates that the absence of

*DUSP6* alone leading to pronounced ERK1/2 activation pushes these cells towards their

maximum glycolytic potential. Upon the addition of 2-DG, glycolysis was completely

abolished under all conditions. These results demonstrate that *DUSP6* inhibition affects

glycolysis in PDAC under basal metabolic conditions, but subsequent rescue experiments are

required to determine the observed changes in glycolytic reserve.

**Glycolysis inhibition does not provide an additional impact cell migratory capacity in**

***DUSP6* depleted cells.**

To gain deeper insights into how metabolic changes induced by *DUSP6* knockdown affect cell

behavior, we conducted a cell proliferation assay. By reducing fetal bovine serum in the media

to minimize its metabolic influence, we assessed cell proliferation with and without 2-DG

addition. Consistent with our previous findings, both AsPC-1 and BxPC-3 cells exhibited a

significant reduction in proliferation upon *DUSP6* knockdown compared to their respective

controls. Following 2-DG treatment, BxPC-3 cells displayed a notable decrease in proliferation

under both conditions, with the reduction being more pronounced in the control cells ($P <$

0.0001; Figure S5B). In contrast, AsPC-1 showed no discernible effect on proliferation

compared to the same conditions without 2-DG (Figure S5A).

Subsequently, to establish a link between the metabolic phenotype and metastatic potential of

these cells, we conducted a migratory capacity assay combined with 2-DG treatment. *DUSP6*

knockdown significantly reduced the migratory capacity of AsPC-1 cells, whereas no change

was observed in BxPC-3 cells (Figure S4). Unexpectedly, treatment with 2-DG did not

significantly alter the migratory phenotype observed in AsPC-1 cells, but significantly reduced

BxPC-3 migratory capacity upon DUSP6 knockdown (Figure 3J-K). Interestingly, we noted

that in the presence of an external source of pyruvate in the media, the cells maintained their

glycolytic potential, even when treated with electron transport chain (ETC) inhibitors

(Rotenone and Antimycin A) and 2-DG (Figure S5C). This suggests that pyruvate availability

plays a critical role in the decision to sustain aerobic glycolysis or to feed the TCA cycle. These

findings suggest that, under nutrient-rich conditions, both AsPC-1 and BxPC-3 utilize different

energy sources for migration.

**DISCUSSION**

DUSP6 is differentially expressed during PDAC tumorigenesis and progression [24]. However,

little is known about the role of phosphatases in the development and establishment of

metastasis. Our group observed that *DUSP6* was overexpressed in primary tumor samples

compared to non-tumoral pancreatic tissues by analyzing *in silico* datasets and spatial transcript

profiling. To further investigate the complex feedback loops involving DUSP6 and the

ERK/MAPK signaling pathway, we evaluated *in silico* data to assess *DUSP6* expression in

late-stage PDAC samples. We observed that *DUSP6* was overexpressed in metastatic samples

compared to primary tumor samples by leveraging *in silico,* immunohistochemistry, and

multiplex RNAscope analysis. Translational relevance for *DUSP6* expression was observed in

survival data; TCGA analysis showed that high DUSP6 expression correlates with lower

overall survival in PDAC patients. Finally, when we analyzed *DUSP6* expression among

different PDAC molecular subtypes [35, 39], *DUSP6* overexpression was pronounced in the

quasi-mesenchymal/squamous subtype, which is associated with a worse prognosis [41]. These

data support the hypothesis that *DUSP6* overexpression confers a more aggressive phenotype,

further impacting metastasis and worsening overall survival in PDAC.

Spatial localization is critical for investigating the functionality. Through *in silico* analysis of

pancreatic tumor samples, we determined that *DUSP6* expression was upregulated in the

epithelial compartment of the tumor rather than in the stromal compartment. The same pattern

was observed in mouse models assessing protein presence by immunohistochemistry and RNA

*in situ* hybridization in mouse pancreatic tumors. In human samples, we observed a significant

decrease in *DUSP6* expression in PDAC epithelial cells versus normal pancreatic tissue, but

increased expression in metastatic epithelial cells. By leveraging a single-cell RNA-seq Human

Pancreatic Cancer Atlas, we were able to confirm that *DUSP6* expression is increased in PDAC

epithelial cells. However, the results also highlighted macrophages as a subset of cells with

important changes in *DUSP6* expression throughout tumorigenesis. Kidger *et al.* recently

reported similar RNA *in situ* hybridization results using KC mice, demonstrating that *DUSP6*

expression is found in ductal cells in PanINs, but not in stromal cells [42]. Our results

confirmed that PDAC tumor cells overexpress *DUSP6*, suggesting the need for further

investigation of its role in tumor development.

Previous studies on multiple tumor types have linked high DUSP6 expression to worse

prognosis and metastasis [43-45]. *DUSP6* has alternatively been described as a tumor

suppressor gene in PDAC. Furukawa *et al.* reported that downregulation of *DUSP6* coordinated

PDAC progression via hypermethylation of its promoter [24, 25]. Kidger *et al.* also proposed

a tumor suppressor role for *Dusp6*, observing that KCD6$^{-/-}$ animals develop a higher number

of poorly differentiated tumors and liver metastases [42]. In their model, *Dusp6* was

constitutively absent, leading to a permanent state of Erk1/2 activation, which would

predictably result in more aggressive tumors and more metastatic foci. Our current hypothesis

is that *DUSP6* suppression might allow tumor initiation, while its expression is required for

metastatic cells to re-establish in a new environment. Reinforcing this idea, *DUSP6*

overexpression appears to correlate with the more invasive quasi-mesenchymal subtype.

To investigate this, we subjected PDAC cell lines to stable *DUSP6* knockdown and evaluated

their proliferation and migration capacities. Surprisingly, we did not find consensus regarding

the observed phenotypes. All statistically significant changes in proliferation indicated an

impairment in this capacity. This observation is counterintuitive because it is widely known as

a dogma in the field that ERK1/2 activation leads to increased cell proliferation. However, it is

also known that tumor cells require fine-tuning of ERK1/2 activation, in which too much

MAPK signaling can bypass the fitness advantage and become detrimental to cell growth [46].

Similarly, Gutierrez-Prat *et al.* recently reported that depletion of DUSP4, another ERK1/2

negative regulator, reduces proliferation by driving "oncogene overdose" in BRAF- and

NRAS-mutant melanoma cell lines [47]. The migration phenotype varied among the cell lines,

complicating any definitive conclusions on this topic. We hypothesized that this variability

might be due to natural compensatory mechanisms within the cells, resulting from dependence

on this signaling pathway. AsPC-1 cells carry two alleles of the $KRAS^{G12D}$ mutation, which is

the most commonly found among PDAC patients and, therefore, more clinically relevant.

BxPC-3 cells present with wild-type *KRAS* which might explain the differences observed

between these two cell lines. A recent study by Bulle *et al.* showed that upon MAPK

pharmacological inhibition, *DUSP6* is one of the top downregulated genes, demonstrating that

*DUSP6* expression correlates with ERK/MAPK activation. Supporting this evidence, we

observed that *DUSP6* was significantly upregulated in tumor cell lines harboring a *KRAS*

mutation compared to tumor cells driven by other mutations in the CCLE dataset (Figure S6).

Bulle *et al.* also demonstrated that upon MAPK inhibition, DUSP6 was targeted for

proteasomal degradation, leading to an increase in HER2 activation [48]. Interestingly, this is

the first time that DUSP6 has been described as a negative regulator of HER2, since it was

previously thought to be ERK-specific. This discovery shifts the paradigm of DUSP6's

functional role and raises the possibility that it may be involved in the negative feedback for

other signaling pathways. Additionally, it is important to consider that DUSP6 is not the only

phosphatase acting as a regulator of ERK phosphorylation. Therefore, its downregulation may

elicit compensatory responses from other molecules, ultimately offsetting its absence. Ito and

colleagues recently demonstrated a co-dependence of DUSP4 and DUSP6 in MAPK-pathway-

driven tumors as a mechanism of drug resistance [49]. In this regard, we highlight that many

DUSPs, among other proteins, are involved in ERK1/2 negative feedback control and have

been shown to act synergistically to prevent oncogene-induced senescence [23]. Similarly, we

found a positive correlation between DUSP6 and MAPK1 expression in tumor cells, but not in

stromal cells, in a human cohort, suggesting that these cells are also dependent on DUSP6 for

ERK1/2 modulation; we cannot, however, rule out the involvement of other DUSPs in the

process. Moreover, the BxPC-3 cell line does not harbor a *KRAS* mutation, and thus, its

tumorigenic behavior does not rely on this condition. The observed phenotype in this cell line

might be driven by mechanisms different from those initially hypothesized. Conversely, the

AsPC-1 cell line, a metastatic PDAC line harboring a *KRAS* mutation, more closely mimics the

genetic background that we aimed to investigate. In this cell line, we observed a significant

impairment in migration capacity upon *DUSP6* knockdown and increased ERK1/2 activation.

To understand the mechanisms by which *DUSP6* affects the metastatic potential of PDAC cells,

we performed gene set enrichment analysis using the metastatic cohort of samples from the

GSE71729 [32] dataset. This analysis indicated a positive correlation with cell migration and

a consistent negative correlation with oxidative phosphorylation. Consequently, we conducted

a glycolysis stress test and observed that the absence of *DUSP6* led to increased glycolytic

function in PDAC cell lines. This upregulation of glycolysis was confirmed by the increased

extracellular acidification rate upon glucose supplementation (in the absence of pyruvate), which could be completely reversed by the addition of 2-DG. This shift towards a more glycolytic phenotype was expected based on previous literature showing that $KRAS^{G12D}$ increases glucose uptake and glycolytic flux through MAPK activation in PDAC cells: Ying and colleagues demonstrated that increased glycolytic flux induces the non-oxidative pentose phosphate pathway (PPP), which is involved in processes such as nucleic acid synthesis and is closely related to cell proliferative capacity [18]. However, in this context, we observed that although glycolysis was induced upon *DUSP6* knockdown, the cells showed a decreased proliferative capacity, a phenotype that is consistent among the cell lines. This behavior might be linked to inherent differences in cell metabolism among PDAC cell lines, as it has been previously reported that cells can present distinct metabolic profiles despite their growth rates [50]. Interestingly, Hsu *et al.* have demonstrated that DUSP6 is required for T-cell receptor-mediated glycolysis, and that in the absence of DUSP6, T cells show higher dependence on glutamine and fatty acids to fuel the TCA cycle [51]. These findings reinforce DUSP6's involvement in metabolic reprogramming but highlight the context-dependency of the observed phenotypes.

Finally, we hypothesized that the induction of glycolysis through *DUSP6* inhibition would decrease the migratory potential of these cells, which could be rescued by the addition of 2-DG, leading to irreversible glycolysis inhibition. We treated the cells with 2-DG and evaluated their migratory capacity but did not observe significant changes with the combination of *DUSP6* and glycolysis inhibition. Previous studies have demonstrated that the induction of glycolysis leads to increased metastatic potential due to the promotion of epithelial-to-mesenchymal transition (EMT) [52, 53]. In the present study, we demonstrated that while cells acquire a more glycolytic phenotype upon *DUSP6* inhibition, their migratory capacity is abrogated. However, inhibition of glucose flux failed to rescue the observed phenotype. We

hypothesized that, in a context where pyruvate is available in the culture media, cells can bypass 2-DG-mediated glycolysis inhibition. To estimate the effect of glycolysis induction on the metastatic behavior of *DUSP6* knockdown cells, it is recommended to closely control nutrient availability in the media. While it is tempting to extrapolate our observations to the decreased metastatic potential of the cells, we must reiterate that metastasis encompasses a variety of complex processes involving both intrinsic and extrinsic signals [54]. To further understand DUSP6's role in metastatic potential, it will be necessary to explore other mechanisms involved in metastasis, such as invasive potential and the ability of these cells to adapt to a new environment, which is highly correlated with their metabolic reprogramming capacity.

**CONCLUSIONS**

This study provides new insights into the role of DUSP6 in PDAC metastasis and metabolism. We demonstrated that DUSP6 inhibition elicits cell-specific effects on migratory potential, while inducing glycolysis in PDAC cells. However, we were unable to establish a direct correlation between the two processes. Future studies should include *in vivo* metastatic assays upon DUSP6 inhibition, as well as metabolomic analysis, to better understand how PDAC cells reprogram their metabolism in the absence of DUSP6 to thrive and survive in a secondary niche.

**ACKNOWLEDGEMENTS**

We would like to acknowledge Dr. David Tuveson (CRUK) for supplying the murine K8484 cell line used in these studies: Dr. Andrew Lowy (UCSD) for supplying the P4313 cell line; Dr. Dafydd Thomas, at the Research Histology and Immunohistochemistry Core (University of Michigan), for conducting the RNA-ISH multiplex assay; the Merchant Lab (University of Miami) for use of the PKT62 cell line; and the Flow Cytometry Core Laboratory at KUMC,

which is sponsored, in part, by the NIH/NIGMS COBRE grant P30 GM103326 and the
NIH/NCI Cancer Center Grant P30 CA168524.

**FINANCIAL SUPPORT**

This study was supported by São Paulo Research Foundation (FAPESP 2015/10694-5) granted to VSS. MTR was supported by CAPES Foundation (Finance Code 001) and Fulbright Association. JS is supported in part by the National Cancer Institute of the National Institutes of Health under award number R37CA262209. MVS received funds from the University of Kansas Cancer Center. CAL was supported by the NCI (R37CA237421, R01CA248160, and R01CA244931).

**AUTHOR'S CONTRIBUTIONS**

MTR: methodology, data collection, formal analysis, visualization, paper writing, and editing; RMW: data collection, formal analysis, and editing; BBB, AEE, WY, FB, JS: data collection and analysis; CAL: supervision, methodology, data analysis, editing; MNV: supervision, methodology, data analysis, paper writing, and editing; VSS: funding acquisition, project administration, conceptualization, supervision, methodology, data analysis, paper writing, and editing.

**COMPLIANCE WITH ETHICAL STANDARDS**

All experiments involving mice were approved by the Ethics Committee on Animal Research at Ribeirao Preto Medical School, University of São Paulo (protocol# 016/2018), and were conducted in accordance with the Guidelines of the Brazilian College of Animal Experimentation.

**CONFLICTS OF INTEREST**

In the past three years, CAL has consulted for Astellas Pharmaceuticals, Odyssey Therapeutics,

Third Rock Ventures, and T-Knife Therapeutics, and is an inventor on patents pertaining to

Kras regulated metabolic pathways, redox control pathways in pancreatic cancer, and targeting

the GOT1-ME1 pathway as a therapeutic approach (US Patent No: 2015126580-A1,

05/07/2015; US Patent No: 20190136238, 05/09/2019; International Patent No:

WO2013177426-A2, 04/23/2015).

**REFERENCES**

1.  Wood, L.D. and R.H. Hruban, *Pathology and Molecular Genetics of Pancreatic*
*Neoplasms.* Cancer journal (Sudbury, Mass.), Nov-Dec 2012. **18**(6).

2.  Muñoz, A.R., et al., *Pancreatic cancer: Current status and Challenges.* Current
pharmacology reports, 2017/12. **3**(6).

3.  Siegel, R.L., A.N. Giaquinto, and A. Jemal, *Cancer statistics, 2024.* CA: A Cancer
Journal for Clinicians, 2024/01/01. **74**(1).

4.  Bardeesy, N., et al., *Pancreatic cancer biology and genetics.* Nature Reviews Cancer
2002 2:12, 2002/12. **2**(12).

5.  McGuigan, A., et al., *Pancreatic cancer: A review of clinical diagnosis, epidemiology,*
*treatment and outcomes.* World Journal of Gastroenterology, 2018/11/11. **24**(43).

6.  Vincent, A., et al., *Pancreatic cancer.* Lancet (London, England), 2011/08/08.
**378**(9791).

7.  Rawla, P., T. Sunkara, and V. Gaduputi, *Epidemiology of Pancreatic Cancer: Global*
*Trends, Etiology and Risk Factors.* World Journal of Oncology, 2019/02. **10**(1).

8.  Disibio, G. and S.W. French, *Metastatic patterns of cancers: results from a large*
*autopsy study.* Arch Pathol Lab Med, 2008. **132**(6): p. 931-9.

9.  Kamisawa, T., et al., *Hematogenous metastases of pancreatic ductal carcinoma.*
Pancreas, 1995. **11**(4): p. 345-9.

10. Mao, C., et al., *Observations on the Developmental Patterns and the Consequences of*
*Pancreatic Exocrine Adenocarcinoma.* Archives of Surgery, 1995/02/01. **130**(2).

11. Embuscado, E.E., et al., *Immortalizing the complexity of cancer metastasis: genetic*
*features of lethal metastatic pancreatic cancer obtained from rapid autopsy.* Cancer
Biol Ther, 2005. **4**(5): p. 548-54.

12. Yachida, S. and C.A. Iacobuzio-Donahue, *The pathology and genetics of metastatic*
*pancreatic cancer.* Arch Pathol Lab Med, 2009. **133**(3): p. 413-22.

13. Haeno, H., et al., *Computational modeling of pancreatic cancer reveals kinetics of*
*metastasis suggesting optimum treatment strategies.* Cell, 2012. **148**(1-2): p. 362-75.

14. Wilentz, R.E., et al., *Loss of expression of Dpc4 in pancreatic intraepithelial neoplasia:*
*evidence that DPC4 inactivation occurs late in neoplastic progression.* Cancer Res,
2000. **60**(7): p. 2002-6.

15. CA, I.-D., *Genetic evolution of pancreatic cancer: lessons learnt from the pancreatic*
*cancer genome sequencing project - PubMed.* Gut, 2012 Jul. **61**(7).

16. Hayashi, A., et al., *The pancreatic cancer genome revisited.* Nature Reviews
Gastroenterology & Hepatology 2021 18:7, 2021-06-04. **18**(7).

17. EA, C., et al., *A central role for RAF→MEK→ERK signaling in the genesis of*
*pancreatic ductal adenocarcinoma - PubMed.* Cancer discovery, 2012 Aug. **2**(8).

18. Ying, H., et al., *Oncogenic Kras Maintains Pancreatic Tumors through Regulation of*
*Anabolic Glucose Metabolism.* Cell, 2012/04/04. **149**(3).

19. Than, M.T., et al., *KRAS-Driven Tumorigenesis and KRAS-Driven Therapy in*
*Pancreatic Adenocarcinoma.* Molecular Cancer Therapeutics, 2024/10/01. **23**(10).

20. N, D. and S. P, *Targeting reactive oxygen species in development and progression of*
*pancreatic cancer - PubMed.* Expert review of anticancer therapy, 2017 Jan. **17**(1).

21. Alonso, A., et al., *Protein tyrosine phosphatases in the human genome.* Cell, 2004.
**117**(6): p. 699-711.

22. S, M., et al., *Extracellular signal-regulated kinases phosphorylate mitogen-activated*
*protein kinase phosphatase 3/DUSP6 at serines 159 and 197, two sites critical for its*
*proteasomal degradation - PubMed.* Molecular and cellular biology, 2005 Jan. **25**(2).

23. Ruckert, M.T., et al., *Protein tyrosine phosphatases: promising targets in pancreatic*
*ductal adenocarcinoma.* Cell Mol Life Sci, 2019. **76**(13): p. 2571-2592.

24. Furukawa, T., et al., *Potential tumor suppressive pathway involving DUSP6/MKP-3 in*
*pancreatic cancer.* Am J Pathol, 2003. **162**(6): p. 1807-15.

25. Xu, S., et al., *Abrogation of DUSP6 by hypermethylation in human pancreatic cancer.*
J Hum Genet, 2005. **50**(4): p. 159-167.

26.  Nagathihalli, N.S., et al., *STAT3 Mediated Remodeling of the Tumor Microenvironment*
*Results in Enhanced Tumor Drug Delivery in a Mouse Model of Pancreatic Cancer.*
Gastroenterology, 2015 Aug 7. **149**(7).
27.  Bankhead, P., et al., *QuPath: Open source software for digital pathology image*
*analysis.* Scientific Reports 2017 7:1, 2017-12-04. **7**(1).
28.  Yi, Z., et al., *KDM6A Regulates Cell Plasticity and Pancreatic Cancer Progression by*
*Noncanonical Activin Pathway.* Cellular and Molecular Gastroenterology and
Hepatology, 2022/01/01. **13**(2).
29.  S, Y., et al., *A Novel MIF Signaling Pathway Drives the Malignant Character of*
*Pancreatic Cancer by Targeting NR3C2 - PubMed.* Cancer research, 07/01/2016.
**76**(13).
30.  G, Z., et al., *DPEP1 inhibits tumor cell invasiveness, enhances chemosensitivity and*
*predicts clinical outcome in pancreatic ductal adenocarcinoma - PubMed.* PloS one,
2012. **7**(2).
31.  Badea, L., et al., *Combined gene expression analysis of whole-tissue and*
*microdissected pancreatic ductal adenocarcinoma identifies genes specifically*
*overexpressed in tumor epithelia.* Hepatogastroenterology, 2008. **55**(Nov-Dec): p. 11.
32.  Moffitt, R.A., et al., *Virtual microdissection identifies distinct tumor- and stroma-*
*specific subtypes of pancreatic ductal adenocarcinoma.* Nature Genetics 2015 47:10,
2015-09-07. **47**(10).
33.  Y, O., et al., *SERPINB3-MYC axis induces the basal-like/squamous subtype and*
*enhances disease progression in pancreatic cancer - PubMed.* Cell reports, 12/26/2023.
**42**(12).
34.  C, M., et al., *Experimental microdissection enables functional harmonisation of*
*pancreatic cancer subtypes - PubMed.* Gut, 2019 Jun. **68**(6).
35.  Bailey, P., et al., *Genomic analyses identify molecular subtypes of pancreatic cancer.*
Nature, 2016. **531**(7592): p. 47-52.
36.  Jin, X., et al., *A metastasis map of human cancer cell lines.* Nature, 2020 Dec 9.
**588**(7837).
37.  A, S., et al., *Gene set enrichment analysis: a knowledge-based approach for*
*interpreting genome-wide expression profiles - PubMed.* Proceedings of the National
Academy of Sciences of the United States of America, 10/25/2005. **102**(43).

38.    Carpenter, E.S., et al., *Analysis of Donor Pancreata Defines the Transcriptomic*
770           *Signature and Microenvironment of Early Neoplastic Lesions.* Cancer Discovery,
771           2023/06/06. **13**(6).

772    39.    Collisson, E.A., et al., *Subtypes of pancreatic ductal adenocarcinoma and their*
773           *differing responses to therapy.* Nature Medicine 2011 17:4, 2011-04-03. **17**(4).

774    40.    Bojmar, L., et al., *Multi-parametric atlas of the pre-metastatic liver for prediction of*
775           *metastatic outcome in early-stage pancreatic cancer.* Nature Medicine 2024 30:8,
776           2024-06-28. **30**(8).

777    41.    Birnbaum, D.J., et al., *Validation and comparison of the molecular classifications of*
778           *pancreatic carcinomas.* Mol Cancer, 2017. **16**(1): p. 168.

779    42.    Kidger, A.M., et al., *Suppression of mutant Kirsten-RAS (KRAS(G12D))-driven*
780           *pancreatic carcinogenesis by dual-specificity MAP kinase phosphatases 5 and 6.*
781           Oncogene, 2022. **41**(20): p. 2811-2823.

782    43.    Kato, M., et al., *Dual-specificity phosphatase 6 plays a critical role in the maintenance*
783           *of a cancer stem-like cell phenotype in human endometrial cancer.* Int J Cancer, 2020.
784           **147**(7): p. 1987-1999.

785    44.    Wu, F., et al., *Nuclear-Biased DUSP6 Expression is Associated with Cancer Spreading*
786           *Including Brain Metastasis in Triple-Negative Breast Cancer.* International Journal of
787           Molecular Sciences, 2019/06. **20**(12).

788    45.    Wu, Q.N., et al., *Pharmacological inhibition of DUSP6 suppresses gastric cancer*
789           *growth and metastasis and overcomes cisplatin resistance.* Cancer Lett, 2018. **412**: p.
790           243-255.

791    46.    Timofeev, O., et al., *ERK pathway agonism for cancer therapy: evidence, insights, and*
792           *a target discovery framework.* npj Precision Oncology 2024 8:1, 2024-03-14. **8**(1).

793    47.    Gutierrez-Prat, N., et al., *DUSP4 protects BRAF- and NRAS-mutant melanoma from*
794           *oncogene overdose through modulation of MITF.* Life Science Alliance, 2022 May 17.
795           **5**(9).

796    48.    Bulle, A., et al., *Combined KRAS-MAPK pathway inhibitors and HER2-directed drug*
797           *conjugate is efficacious in pancreatic cancer.* Nature Communications 2024 15:1,
798           2024-03-20. **15**(1).

799    49.    T, I., et al., *Paralog knockout profiling identifies DUSP4 and DUSP6 as a digenic*
800           *dependence in MAPK pathway-driven cancers - PubMed.* Nature genetics, 2021 Dec.
801           **53**(12).

50.    Daemen, A., et al., *Metabolite profiling stratifies pancreatic ductal adenocarcinomas*
*into subtypes with distinct sensitivities to metabolic inhibitors.* Proceedings of the
National Academy of Sciences of the United States of America, 2015/08/08. **112**(32).

51.    Hsu, W.-C., et al., *DUSP6 mediates T cell receptor-engaged glycolysis and restrains*
*TFH cell differentiation.* Proceedings of the National Academy of Sciences, 2018-8-21.
**115**(34).

52.    Li, M., et al., *Biological role of metabolic reprogramming of cancer cells during*
*epithelial-mesenchymal transition (Review).* Oncology Reports, 2019-02-01. **41**(2).

53.    Kang, H., et al., *Role of Metabolic Reprogramming in Epithelial–Mesenchymal*
*Transition (EMT).* International Journal of Molecular Sciences, 2019/04. **20**(8).

54.    Joshi, V.B., O.L.G. Ruiz, and G.L. Razidlo, *The Cell Biology of Metastatic Invasion in*
*Pancreatic Cancer: Updates and Mechanistic Insights.* Cancers, 2023/04. **15**(7).

# Figure 1

# Figure 2

# Figure 3

**Figure 1 – *DUSP6* overexpression is correlated to tumor metastatic potential and worse prognosis.** *In silico* datasets from GSE62452 [29] **(A)**, GSE28735 [30] **(B)** and GTex/TCGA **(C)** assessing *DUSP6*'s differential expression between non-tumoral pancreatic tissue and primary tumor tissue; **(D)** GSE71729 [32] assessing *DUSP6* expression between non-tumoral pancreatic tissue, primary and metastatic tumor tissue; *In silico* gene expression analysis utilizing **(E)** Collisson [39], **(F)** Moffitt [32], **(G)** Bailey [35] and **(H)** GSE224564 [33] datasets to assess *DUSP6* expression among PDAC molecular subtypes; **(I)** Kaplan-Meier (Log-rank) plot comparing *DUSP6* expression and patients' overall survival using the TCGA dataset. Padj. = 0.0339, N = 67 for each group; **(J & K)** Pearson correlation analysis between *DUSP6* expression in CCLE PDAC cell lines and their metastatic potential (MetMap 500 [36]). R = 0.6105, P = 0.0003. ADEX: aberrantly differentiated endocrine–exocrine; QM: quasi-mesenchymal. (A, H) Mann-Whitney unpaired test. (B, C, F) Welch's unpaired t-test. (D, G) Brown-Forsythe and Welch unpaired ANOVA test. (E) Kruskal-Wallis unpaired test. * P < 0.05; ** P < 0.01; *** P < 0.001; **** P < 0.0001.

**Figure 2 – *DUSP6* is upregulated in PDAC epithelial cells in human and mouse tissue.** **(A)** DUSP6 immunohistochemistry staining (20x magnification) in wild type (WT) pancreatic tissue (N=3), KC (N=3) and KPC (N=6) mice. Yellow arrows indicate acinar-to-ductal metaplasia (ADM) lesions and red arrows indicate Pancreatic Intraepithelial Neoplasia (PanIN) lesions. Quantification (B) shows DAB OD mean ± SD in DUSP6 positive cells from 10 randomly selected areas in each slide stratified by lesion grade; **C)** Multiplex *in situ* hybridization with *DUSP6* (red, RNAscope), panCK (green), αSMA (orange) and DAPI (blue). Images are representative of normal pancreatic tissue/pancreatitis, Intraductal papillary mucinous neoplasms (IPMN), Mucinous Cystic Pancreatic Neoplasms (MCN), Pancreatic Intraepithelial Neoplasia (PanIN), Pancreatic Ductal Adenocarcinoma (PDAC) and PDAC metastasis samples composed from 2 human TMA sections; **D)** Quantification of the percentage *DUSP6* positive cells normalized by total number of cells in each TMA core; **E)** Quantification of the percentage of cells that are positive for both *DUSP6* and panCK, normalized by total number of cells in each TMA core. Welch's unpaired t-test; **F)** UMAP plot of all cells captured from single-cell RNA sequencing of eleven human tumor samples [28]. Cell types are identified by color. Following UMAPs show DUSP6 expression among cell types in healthy pancreatic tissue **(G)**, adjacent normal pancreatic tissue **(H)** and

PDAC samples **(I)**; **J)** Percentage of cells co-expressing *DUSP6* and *KRT19* (epithelial marker) in healthy pancreatic tissue, adjacent normal pancreatic tissue and PDAC samples assessed by single-cell RNA-seq [28]. * $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$; **** $P < 0.0001$.

**Figure 3 – DUSP6 downregulation increases glycolysis in PDAC cell lines, but glycolysis inhibition does not rescue migratory phenotype.** GSEA analysis of GSE71729 dataset showing pathways significantly negatively **(A & C)** and positively **(B & C)** correlated to *DUSP6* expression in metastatic samples; **D)** Western blot analysis of DUSP6 expression and ERK1/2 phosphorylation upon DUSP6 knockdown. Quantification shows DUSP6 expression normalized to β-actin, and P-ERK1/2 normalized to total ERK1/2 levels; **E)** Representative curves of extracellular acidification rates (ECAR) measured in AsPC-1 and BxPC-3 with sequential injections of glucose (10 mM), oligomycin (1 µM) and 2-DG (100 mM), respectively in a Seahorse assay; **F)** Basal extracellular lactate secretion levels upon DUSP6 knockdown in complete media; **G)** Basal glycolysis corresponds to glycolytic levels under basal conditions, no stimulation; **H)** Glycolytic capacity is measured after inhibition of ATP synthase (complex V of the mitochondrial respiratory chain); **I)** Glycolytic reserve corresponds to the difference between glycolytic capacity and basal glycolysis. Assessment of migratory capacity upon 2-DG treatment in AsPC-1 **(J)** and BxPC-3 **(K)** upon DUSP6 knockdown. Seahorse Glycolysis Stress Test was performed in 9-12 replicates in 2 independent experiments. Results represent mean ± SD of one representative experiment. L-Lactate assay was performed in 2 independent experiments with 6 replicates each. Results represent mean ± SEM. Welch's unpaired t-test; Migration graphs show 3 independent experiments with 4-6 replicates each. Results represent mean ± SEM. Mixed-effects analysis with Šídák's multiple comparisons test. *ns*: not significant; * $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$; **** $P < 0.0001$.