1 **Global diversity of putative functional viral auxiliary metabolic genes and their**
2 **relationship with soil organic carbon stocks**
3

4 Dominik Merges*
5
6 *Corresponding author:
7 Dominik Merges
8 Mergesd01@gmail.com

9
10 **Running title:** Functional AMG diversity in soils
11
12 **Key words:** bacteria, bacteriophages, ecosystem functioning, ecosystem processes, nutrient
13 cycling, carbon, nitrogen, species interactions
14

Abstract

Global biodiversity decline has prompted extensive research and debate on how biodiversity
loss affects ecosystem functioning. Microbial taxonomic and functional diversity loss can
impair critical processes such as plant biomass degradation or nitrogen fixation. Biotic
interactions, especially those involving microbes and their viruses, are increasingly
recognized as drivers of such processes. Microbial viruses carry auxiliary metabolic genes
(AMGs), functional genes acquired from bacterial hosts, that mediate processes like carbon
and nitrogen cycling, positioning viruses as potential reservoirs of essential ecosystem
functions. To investigate the putative contribution of viral AMGs to microbial functional
diversity and ecosystem functioning in soils, the global dataset of soil viral genomes,
comprising 616,935 uncultivated viral genomes and 38,508 unique viral operational
taxonomic units from 2,953 soil metagenomes, was mined for carbon (C) and nitrogen (N)
cycling genes. For C cycling, 301 viral genomes with AMGs from glycoside hydrolase (GH)
families (n cellulase = 182; n hemicellulase = 119) were detected. The viruses containing
cellulase and hemicellulase AMGs were distributed across most major ecosystem types,
including agricultural, forest, grassland, and permafrost soils, but were absent from deserts,
salt marshes, and shrublands. Forest ecosystems harbored the highest abundance of
cellulase and hemicellulase AMGs. The ratio of cellulase to hemicellulase AMGs varied by
ecosystem, showing near parity in forests but cellulase dominance in agricultural, grassland,
and permafrost soils. Relating AMGs to ecosystem functioning, sites with cellulase and
hemicellulase AMGs showed significantly lower soil organic carbon (SOC) content in
agricultural soils, whereas no significant differences were observed in forest, grassland, or
permafrost soils. These results suggest that viral AMGs may influence SOC dynamics,
particularly in agricultural systems, and highlight the need for deeper investigation into their
functional roles in soil ecosystems. This study underscores the importance of viral
contributions to ecosystem functioning, emphasizing the potential of microbial viruses to
serve as mutualistic agents in global biogeochemical cycles.

Global biodiversity decline has prompted extensive research and debate on how biodiversity
loss affects ecosystem functioning (WWF 2022). Microbial taxonomic and functional diversity
loss can impair critical processes such as plant biomass degradation or nitrogen fixation, and
denitrification. The loss of microbial taxonomic and functional diversity can impair processes
such as plant biomass decomposition (Wagg et al., 2019), nitrogen fixation, and
denitrification (Castaño et al., 2023; Philippot et al., 2013). However, in certain instances,
diversity loss appears to have little to no impact on these processes, highlighting the
complexity of interactions within microbial communities (Wertz et al., 2006). These
interactions may be dependent on attributes of communities differing in their structure and
species composition that may result in different patterns of ecosystem functional response to
diversity loss (Tilman et al., 1997; Philippot et al., 2013). Currently, it remains uncertain
which specific community attributes are most relevant to an ecosystem's functional response
to the loss of diversity.

Among the community attributes relevant to ecosystem processes and functions, biotic
interactions, particularly those involving microbes and their viruses, are increasingly
recognized as relevant (Braga et al., 2018). While significant attention has been given to
viruses' antagonistic roles, their potential mutualistic effects on ecosystem functioning remain
relatively unexplored. Recent findings indicate that microbial viruses can carry auxiliary
metabolic genes (AMGs), functional genes acquired from bacterial hosts, that play critical
roles in processes like carbon and nitrogen cycling, stress response, and energy production,
positioning viruses as potential reservoirs of essential ecosystem functions (Roux et al.,
2016).

Although most AMG-related evidence originates from marine systems (Roux et al., 2016),
soil studies have also demonstrated virus-encoded AMGs supporting bacterial and archaeal
functions, including pesticide degradation, carbon and methane cycling, and nitrification
(Emerson et al., 2018; Zheng et al., 2022). Moreover, specific functional AMGs have been
experimentally verified for roles in plant biomass decomposition (Emerson et al., 2018) and
organochlorine pesticide degradation (Zheng et al., 2022). These findings suggest that
microbial viruses may function as mutualists, equipping microbial communities with additional
functional genes and thereby significantly influencing ecosystem functions (Jurburg et al.,
2023).

The aim of this study was to investigate 1) the prevalence of putative functional AMGs across
different ecosystem types and 2) their potential contribution to microbial functional diversity
and ecosystem functioning in soils.

To quantify the prevalence of viral AMGs in soils, functionally annotated viral genomes from
the global dataset of soil viral genomes (Graham et al. 2024) were screened, to collect viral
genomes with AMGs linked to ecosystem functioning. The global dataset of soil viral
genomes contained 616935 uncultivated viral genomes and 38508 unique viral operational
taxonomic units assembled from 2953 previously sequenced soil metagenomes (Graham et
al. 2024). From this dataset, a database of viral genomes with functional AMGs, relevant for
C (i.e. degradation of plant matter) and N (nitrogen fixation) cycles (Figure 1a) was build, by
identifying AMGs linked to C and N cycles, leveraging functional annotations from CAZy
database (Drula et al. 2022 and the KEGG database modules to phenotype annotations
(Kanehisa et al. 2016). For C cycling, the viral genomes were screened for AMGs associated
with the degradation of plant biomass, i.e. microbial GH families targeting plant cell wall
degradation, specifically for cellulose (CAZyme identifier: GH5, GH6, GH7, GH9, GH16,
GH17, GH44, GH45, GH48, GH74; Auer et al. 2023) and hemicellulose (CAZyme identifier:
GH8, GH10, GH11, GH12, GH26, GH28, GH30, GH53; Auer et al. 2023). In total, 301 viral
genomes with AMGs from the GH families (n cellulase = 182; n hemicellulase = 119) were
found (Fig. 1a). For N cycling, I found one viral genome with a nifH AMG. The NifH gene
encodes for the Nitrogenase iron enzyme responsible for nitrogen fixation of nitrogen-fixing
bacteria and archaea (nifH, KEGG ortholog K02588; Fig 1a).

The cellulase and hemicellulase AMGs were present within most major ecosystem types (i.e.
agricultural, forests, grasslands, permafrost), but absent in deserts, salt marshes and
shrublands (Fig. 1b). Cellulase and hemicellulase AMGs were most common in forest
ecosystems, which typically have rather stable environmental conditions and a highly diverse
plant biomass, including leaves, wood and understorey vegetation which contain a mix of
cellulose and hemicellulose, and accordingly, both cellulase and hemicellulase AMGs are
abundant. The two degradation types (i.e. cellulase and hemicellulase) show different ratios
across ecosystems, with an almost balanced ratio in forests, but dominance of cellulase over
hemicellulose in agricultural, grassland and permafrost ecosystems (Fig. 1c). Cellulase
dominance may be attributed to several factors, such as that microorganisms may prioritize
cellulase production 1) based on the lower diversity of plant biomass in agricultural,
grassland, and permafrost soils and 2) under less stable environmental conditions (e.g.,
tillage, moisture variability, and temperature extremes), as cellulose provides a more
recalcitrant and consisted carbon source compared to hemicellulose.

Relating the presence of cellulase and hemicellulase AMGs to ecosystem functioning, a
comparison of soil organic matter (SOC) content in sites lacking degradation genes (grey
boxes) versus those containing degradation genes (blue boxes) reveals significantly lower
SOC content in agricultural samples harboring putative viral degradation genes (t-test $p <$
0.001; Fig. 1d). This observation aligns with findings that agricultural practices accelerate
SOC breakdown through microbial activity (Lal, 2004; Six et al., 2002). The presence of viral
degradation genes may indicate active viral and microbial communities contributing to SOC
decomposition, consistent with evidence linking viral auxiliary metabolic genes (AMGs) to
ecosystem-level carbon cycling (Roux et al., 2016). In agricultural systems, intensive land-
use practices (e.g., tillage, crop harvest) expose organic matter to microbial degradation,
potentially enhancing SOC breakdown by active viral and microbial communities (Paustian et
al., 1997). Conversely, the stable microclimate and continuous organic matter input in
forests, low temperatures and frozen soils in permafrost, as well as extensive root biomass
and well-structured soils in grasslands, may contribute to maintaining stable SOC levels
regardless of the presence of viral AMGs (Jobbágy & Jackson, 2000; Schuur et al., 2015;
Jackson et al., 1996).

The observed association of putative viral AMG presence on SOC in agricultural systems
underscores the need for a more comprehensive evaluation of their ecological roles.
Controlled experiments offer a promising approach to elucidate the contribution of viral
AMGs to microbial-mediated SOC degradation. Such investigations will advance our
understanding of the interplay between viral and microbial communities in shaping
ecosystem carbon dynamics.

References

Burke, C., P. Steinberg, D. Rusch, S. Kjelleberg, T. Thomas, Bacterial community assembly based on functional genes rather than species, Proc. Natl. Acad. Sci. U.S.A. 108 (34) 14288-14293.

Braga, L. P., Soucy, S. M., Amgarten, D. E., Da Silva, A. M., & Setubal, J. C. (2018). Bacterial diversification in the light of the interactions with phages: the genetic symbionts and their role in ecological speciation. Frontiers in Ecology and Evolution, 6, 6.

Castaño, C., Hallin, S., Egelkraut, D., Lindahl, B.D., Olofsson, J. and Clemmensen, K.E. (2023), Contrasting plant–soil–microbial feedbacks stabilize vegetation types and uncouple topsoil C and N stocks across a subarctic–alpine landscape. New Phytol, 238: 2621-2633.

Emerson, J. B., Roux, S., Brum, J. R., Bolduc, B., Woodcroft, B. J., Jang, H. B., ... & Sullivan, M. B. (2018). Host-linked soil viral ecology along a permafrost thaw gradient. Nature microbiology, 3(8), 870-880.

Graham, E.B., Camargo, A.P., Wu, R. et al. A global atlas of soil viruses reveals unexplored biodiversity and potential biogeochemical impacts. Nat Microbiol 9, 1873–1883 (2024).

Grime, J. Competitive Exclusion in Herbaceous Vegetation. Nature 242, 344–347 (1973.

Jackson, R. B., Canadell, J., Ehleringer, J. R., Mooney, H. A., Sala, O. E., & Schulze, E. D. (1996). A global analysis of root distributions for terrestrial biomes. Oecologia, 108(3), 389–411. Jobbágy, E. G., & Jackson, R. B. (2000). The vertical distribution of soil organic carbon and its relation to climate and vegetation. Ecological Applications, 10(2), 423–436. Jurburg SD, Hom EFY, Chatzinotas A (2023) Beyond pathogenesis: Detecting the full spectrum of ecological interactions in the virosphere. PLOS Biology 21(5): e3002109.

Kuzyakov, Y., & Mason-Jones, K. (2018). Viruses in soil: Nano-scale undead drivers of microbial life, biogeochemical turnover and ecosystem functions. Soil Biology and Biochemistry, 127, 305-317.

Lal, R. (2004). Soil carbon sequestration impacts on global climate change and food security. Science, 304(5677), 1623–1627. McNaughton S J. Biodiversity and function of grazing ecosystems. In: Schulze E D, Mooney H A, editors. Biodiversity and ecosystem functioning. New York, N.Y: Springer Verlag; 1993. pp. 361–384.

McNaughton S J. Biodiversity and function of grazing ecosystems. In: Schulze E D, Mooney H A, editors. Biodiversity and ecosystem functioning. New York, N.Y: Springer Verlag; 1993. pp. 361–384.

Merges, D., Dal Grande, F., Greve, C., Otte, J., & Schmitt, I. (2021). Virus diversity in metagenomes of a lichen symbiosis (Umbilicaria phaea): complete viral genomes, putative hosts and elevational distributions. Environmental Microbiology, 23(11), 6637-6650.

Merges, D., Schmidt, A., Schmitt, I., Neuschulz, E. L., Dal Grande, F., & Bálint, M. (2023). Metatranscriptomics reveals contrasting effects of elevation on the activity of bacteria and bacterial viruses in soil. Molecular Ecology, 32(23), 6552-6563.

Naeem, S., Thompson, L. J., Lawler, S. P., Lawton, J. H., & Woodfin, R. M. (1994). Declining biodiversity can alter the performance of ecosystems. Nature, 368(6473), 734-737. Naeem S. (1998). Species redundancy and ecosystem reliability. Conserv Biol 12: 39–45

Le Provost G., Thiele J., Zaitsev A. S., Manning P. (2021): Contrasting responses of above- and belowground diversity to multiple components of land-use intensity. Nature Communications 12:3918.

Roux, S. et al. Ecogenomics and potential biogeochemical impacts of globally abundant ocean viruses. Nature 537, 689–693 (2016).Tilman, D., Reich, P. B. & Knops, J. M. H. Biodiversity and ecosystem stability in a decade-long grassland experiment. Nature 441, 629–632 (2006).

Tilman D, Lehman C, Thomson K. (1997). Plant-diversity and ecosystem productivity: theoretical considerations. Proc Natl Acad Sci USA 94: 1857–1861.

Paustian, K., Collins, H. P., & Paul, E. A. (1996). Management controls on soil carbon. In E. A. Paul et al. (Eds.), Soil organic matter in temperate agroecosystems (pp. 15–49). CRC Press.

Philippot, L. et al. Loss in microbial diversity affects nitrogen cycling in soil. ISME J. 7, 1609–1619 (2013).

Santos-Medellín, C., Estera-Molina, K., Yuan, M., Pett-Ridge, J., Firestone, M. K., &
Emerson, J. B. (2022). Spatial turnover of soil viral populations and genotypes overlain by
cohesive responses to moisture in grasslands. Proceedings of the National Academy of
Sciences, 119(45), e2209132119.
Schuur, E. A. G., McGuire, A. D., Schädel, C., Grosse, G., Harden, J. W., Hayes, D. J., et al.
(2015). Climate change and the permafrost carbon feedback. Nature, 520(7546), 171–179.
Six, J., Conant, R. T., Paul, E. A., & Paustian, K. (2002). Stabilization mechanisms of soil
organic matter: Implications for C-saturation of soils. Plant and Soil, 241(2), 155–176.
ter Horst, A. M., Jane D Fudyma, Jacqueline L Sones, Joanne B Emerson, Dispersal, habitat
filtering, and eco-evolutionary dynamics as drivers of local and global wetland viral
biogeography, The ISME Journal, Volume 17, Issue 11, November 2023, Pages 2079–2089.
Trubl, G., Kimbrel, J.A., Liquet-Gonzalez, J. et al. Active virus-host interactions at sub-
freezing temperatures in Arctic peat soil. Microbiome 9, 208 (2021).
van der Plas, F. (2019). Biodiversity and ecosystem functioning in naturally assembled
communities. Biological Reviews of the Cambridge Philosophical Society, 94(4), 1220–1245.
Wagg, C., Schlaeppi, K., Banerjee, S. et al. Fungal-bacterial diversity and microbiome
complexity predict ecosystem functioning. Nat Commun 10, 4841 (2019.
Wertz S, Degrange V, Prosser JI, Poly F, Commeaux C, Freitag T et al. (2006). Maintenance
of soil functioning following erosion of microbial diversity. Environ Microbiol 8: 2162–2169.
WWF. (2022). Living planet report 2022—building a nature-positive society. M. Grooten, D.
Bignoli Juffe, & T. Petersen (Eds.). WWF.
Zheng, X., Jahn, M.T., Sun, M. et al. (2022) Organochlorine contamination enriches virus-
encoded metabolism and pesticide degradation associated auxiliary genes in soil
microbiomes. ISME J 16, 1397–1408
212

Figure 1: A) Number of found viral genomes containing AMGs for the degradation of plant
biomass and nitrogen fixation. B) Prevalence of putative cellulase and hemicellulase AMGs.
C) Putative cellulase and hemicellulase AMGs distributions differ across ecosystems D)
Agricultural sites with cellulase and hemicellulase AMGs hold significantly less SOC than
agricultural without. In forest, grassland and permafrost ecosystems sites with and without
cellulase and hemicellulase AMGs do not differ in SOC content.

222