To Peck or Not To Peck: The influence of early-life social environment

on response inhibition and impulsive aggression in Japanese quails

Alizée Vernouillet[1*], Kathryn Willcox[1], Reinoud Allaert[1], Anneleen Dewulf[1], Wen Zhang[1,2],

Camille A. Troisi[1], Sophia Knoch[1], An Martel[2], Luc Lens[1], Frederick Verbruggen[1]

1. Centre for Research on Ecology, Cognition and Behaviour of Birds, Ghent University

2. Department of Veterinary Sciences, Ghent University

Alizée Vernouillet ORCID: https://orcid.org/0000-0003-2525-2936

Kathryn Willcox ORCID: https://orcid.org/0000-0002-3222-2850

Reinoud Allaert ORCID: https://orcid.org/0000-0002-6122-496X

Anneleen Dewulf ORCID: https://orcid.org/0000-0003-4215-0421

Wen Zhang ORCID: https://orcid.org/0000-0001-9120-9763

Camille A. Troisi ORCID: https://orcid.org/0000-0002-4036-3848

Sophia Knoch ORCID: https://orcid.org/0000-0001-9156-5725

An Martel ORCID: https://orcid.org/0000-0001-7609-5649

Luc Lens ORCID: https://orcid.org/0000-0002-0241-2215

Frederick Verbruggen ORCID: https://orcid.org/0000-0002-7958-0719

Correspondence concerning this article should be addressed to Alizée Vernouillet. Email:

alizee.vernouillet@gmail.com

1. Summary

Deficits in response inhibition (i.e., the ability to suppress inappropriate responses) may contribute to increased impulsive aggression (i.e., unplanned behaviors that harm others). Since early-life environment might influence the development of response inhibition, it could also indirectly affect impulsive aggression. However, this relationship has rarely been directly examined. Here, we investigated whether response inhibition is associated with impulsive aggression and whether this relationship explains the effects of early-life social environments on impulsive aggression in Japanese quails (*Coturnix japonica*). Quails ($n = 120$) were raised in two social conditions: Small groups of five birds or Large groups of 15 birds. Response inhibition was assessed using the Barrier and Cylinder tasks, while impulsive aggression was measured in two contexts - within a group of five familiar individuals and in a dyad with an unfamiliar individual. We found that some aspects of response inhibition were related to impulsive aggression. Furthermore, quails reared in small groups showed significantly poorer response inhibition than those reared in large groups. Yet, impulsive aggression did not significantly differ between the two conditions. These findings suggest that, while impulsive aggression is partly related to response inhibition, other factors mitigate the effects of early social environments on its expression.

Keywords: aggression, early-life social environment, inhibitory control, impulsive behaviour, personality, response inhibition

2. Introduction

Group living brings many benefits, such as reducing predation pressure and increasing foraging efficiency, but can also create conflicts between members due to increased competition (1). These agonistic interactions between individuals may result in *impulsive aggression* (i.e., the unplanned behavioural response of inflicting damage to self or others, (2)), that in turn may sometimes have negative fitness consequences for one or multiple individuals involved (e.g., reduced access to resources, injuries, or even death). While a lot of research has focused on aggression as a personality trait in animals (as reviewed by e.g., (3)), little is known about the cognitive mechanisms underlying aggression.

Some researchers suggest that impulsive aggression results from deficits in *response inhibition* (also referred to in the animal cognition field as *inhibitory control*; (4–6); but see (7)), the suppression of inappropriate pre-potent behavioural responses or actions (8–10). Indeed, inhibition deficits are correlated with increased aggression in rats (11), dogs (12), long-tailed macaques (13), and humans (14–16). Additionally, both increased levels of aggression and decreased levels of response inhibition have been correlated with lower levels of central serotonin (5-HT), a well-conserved neuromodulator of nervous systems in both invertebrates and vertebrates (9,11,17,18). However, most studies investigating the link between response inhibition and impulsive aggression are correlational. This has two important consequences. First, response inhibition deficits could cause or contribute to impulsive aggression, but it is also possible that a common underlying factor simply influences both (independently) (as suggested by (19)). Second, it has been proposed that environmental factors during early life contribute to response inhibition deficits and impulsive aggression later in life. Yet again, causal relationships have been hard to establish, and systematic research is lacking.

Early-life environmental conditions, both physical and social, can shape an individual's cognitive abilities and personality (e.g., (20–24)). For example, response inhibition is key to

maintain spatial and temporal cohesion between group members (25). Indeed, individuals within a group benefit from being surrounded by others and performing the same activity together, so they need to adjust their activity to that of other group members (e.g., (26,27)). Additionally, within a group, subordinate individuals might inhibit feeding or mating in the presence of dominant individuals to avoid potential aggressive interactions or confrontations (28). Given that social complexity is multi-faceted and highly variable, both within and between species (29), in the present study we focused on one key aspect of the social organization – social group size (30). An increase in group size means an increase in potential interactions with other group members, and therefore an increase in possibilities to learn how to interact with other individuals. Social group size has been shown to correlate with higher levels of response inhibition in Australian magpies (*Cracticus tibicen dorsalis*, (31)), in hyenas (*Crocuta crocuta*, (32)), and in primates (33), but not in guppies (*Poecilia reticulata*, (24,34)). Hence, we expected that a larger group size required individuals to develop better response inhibition due to greater social requirements.

Many studies have found aggression is usually suppressed in larger groups (e.g., calves – (35), chickens – (36,37), fish – (38), pigs – (39)). Based on the same reasoning described in the previous paragraph, with larger groups come a higher number of potential competitors for resources, and individuals might thus benefit to switch to a more socially tolerant strategy (e.g., (37,40), but see e.g., (41)). Yet, methodological differences between studies make it difficult to determine whether, and if so in what direction, early-life social environment shapes an individual's aggressive tendencies (42). Based on previous research and on the theoretical link between inhibition and aggression, and because we predicted that individuals raised in larger groups would have better response inhibition, we expected levels of impulsive aggression to decrease with increasing group size.

In this study, we investigated the link(s) between early-life social environment, response inhibition and aggression in Japanese quails (*Coturnix japonica*). Japanese quails are a precocial social bird species that is easy to raise in captivity, which enabled us to rear them without the presence of

adult quails. This controlled raising environment allowed us to manipulate their postnatal social

conditions, while controlling all other environmental parameters. Additionally, Japanese quails have

long been recognized as an ideal model for studying aggression, as hierarchies within their social

group are established and maintained through frequent aggressive interactions (e.g., (26,43,44)). As

a result, their typical behaviors, including aggressive behaviors towards conspecifics, have been

extensively documented (e.g., (44–46)). These combined factors were crucial in establishing causal

relationships between early-life social environment and an individual's response inhibition and

aggression.

We evaluated response inhibition in Japanese quails with two commonly used tasks to

evaluate an animal's response inhibition: the *Barrier task*, during which individuals must detour

around a vertical transparent barrier to reach a reward, and the *Cylinder task*, during which

individuals must reach through the openings of a transparent cylinder to obtain a reward. Both

tasks are purporting to measure the same aspect of response inhibition and have been referred to

as "detouring" tasks or "motoric self-regulation" tasks, as, in both tasks individuals must inhibit a

prepotent motor action (i.e., going towards the visible reward) and instead change it into a different

one (i.e., detouring around a physical barrier) (47) (but see (48)). Aggression was evaluated in two

different contexts: once in a group of familiar individuals (e.g., (44)), and once in presence of an

unfamiliar individual (e.g., (49)).

The main goal of the study was to experimentally determine the influence of early-life social

environment on response inhibition and impulsive aggression in Japanese quails. We focused on

three research questions: (A) do deficits in response inhibition cause impulsive aggression at an

individual-level, and (B) does social group size during early life shape an individual's response

inhibition and (C) impulsive aggression? Based on previous correlational research and on

theoretical models, we predicted that, if indeed higher impulsive aggression can be attributed to a

deficit in response inhibition, (A) response inhibition and impulsive aggression would be inversely

related at an individual-level, and that (B and C) individuals raised in larger groups would have a

higher level of response inhibition and be less aggressive than individuals raised in smaller groups.

3. Material and methods

Subjects

We used 120 Japanese quails ($n = 60$ females, $n = 60$ males) in this study. Birds were hatched

simultaneously over two days and kept together at the Opvangcentrum Vogels en Wilde Dieren,

Ostend, Belgium. Birds were housed in four identical heated lofts, each loft containing four 1m² (1

m length x 1 m width x 2 m height) enclosures. Each enclosure included an electric hen heating

plate (25 cm length x 25 cm width, Comfort Chicks), which was replaced with a plastic cover when

birds were 16 days old. Outside of the experimental food restriction procedure, birds were provided

with *ad lib* grain food mix (Aveve), presented in a single 50-cm long linear chick feeder (Voss

Framing). Birds had free access to water. Each loft followed a 12:12 day-night cycle, with light

onset at 07:00 am. All birds were visually checked at least twice a day (once in the morning and

once in the evening). Measurements (weight and tarsus length) were taken when the birds were 20

days old (before the start of the study) and at 44 days old (at the end of the study). Quail care was

performed by AV, KW, RA, AD, WZ, and CT, under AV and RA's supervision.

Within each loft, one of the four enclosures housed a large group of 15 individuals (Large),

and the three other enclosures housed a small group of five individuals (Small). On day 2 post-

hatch, birds were randomly assigned to either a Small or a Large enclosure. Hence, 60 birds were

assigned to each experimental condition.

General Experimental Procedures

Quails were habituated to being handled and to the testing environment during their second

and third week. Testing started when birds were 24 days old. All 120 individuals underwent the

Barrier task, the Cylinder task, the Familiar Group task, and the Unfamiliar Individual task, given

in that order for all individuals. Birds were tested individually during the Detour and the Cylinder

tasks (i.e., response inhibition tasks), and with other individuals during the Familiar Group and the

Unfamiliar Individual tasks (i.e., impulsive aggression tasks), and tasks were performed on separate

days. A schematic overview of the study timeline can be found in the Supplementary Material (Fig.

S1).

Testing took place between 08:30 and 14:00 (local time). Food was removed the evening

prior testing to ensure motivation to participate in the experiment. On the morning of testing, birds

were individually placed in a portable cage with multiple compartments (Ducatillon) and

transported to a waiting room, where they were kept before and after testing until all individuals

had been tested.

All testing occurred in a test box (86 cm length x 146 cm width x 91 cm height). The testing

box was lit with LED lights to control for lighting. All trials were recorded using a camera

(BASCOM) mounted to the ceiling of the test box. A 'start box' (29 cm length x 45 cm width x 29

cm height) was attached to the front of the test box. The start box had two doors – one transparent

and one opaque – which slid vertically on rails between the two boxes, and a separate light. Inside

the start box was another square piece of wood that served as the back of the box attached to a

handle, which could slide forward and was used to gently push the birds into the test box when

required.

Every testing trial followed the same general procedure. First, the bird(s) to be tested was

(were) placed in the start box with both start box doors closed. Next, the opaque door was opened,

allowing the bird(s) to see within the test box but not enter the test box. After 15 seconds the

transparent door was opened, indicating the start of the trial. Birds that did not leave the start box

within 30 seconds were gently pushed into the test box. Once the bird(s) either reached the reward

location, or the time limit for the trial was reached, the test box lights were turned off, indicating the end of the trial. The bird(s) was (were) given 15 seconds to return to the start box; if it(they) did not, it was (they were) caught by the experimenter and placed back in the start box. Once all birds completed all required trials for a session, they were returned to their home enclosures. Data collection was performed by AV, KW, RA, AD, WZ, CT, and LL, under AV and KW's supervision.

Videos were coded using BORIS v.7.13 (Behavioral Observation Research Interactive Software (50)). All the videos of the response inhibition tasks were coded by one experimenter (KW), who was naïve to the conditions. All the videos of the impulsive aggression tasks were coded by another coder (SK), who was naïve to the conditions and the experimental hypotheses. To assess inter-observer reliability, 20 % of the videos of the response inhibition tasks were coded by one coder (SK), who was naïve to the conditions and the experimental hypotheses, and 20 % of the videos of the impulsive aggression tasks were coded by one experimenter (AV). Inter-rater reliability was calculated using BORIS, and was high, with average Cohen's kappa coefficients of 0.840 and 0.805, for all response inhibition videos and all impulsive aggression videos respectively (51).

Habituation

Birds were gradually habituated to the portable cage from 20-days-old onwards and to the test box at 22-23 days old. For the habituation in the test box, a feeder was placed 50cm from the test box entrance. The feeder used was filled with grains and was the same design as those used in the birds' enclosures, but smaller (30cm length). A petri dish with two live mealworms was also placed in front of the feeder. Both rewards were familiar to the birds and were used to encite them to participate in testing. The total trial time was 5 minutes; each bird received one habituation trial.

Response inhibition tasks

<u>Barrier task</u>

Apparatus. The apparatus for the Barrier task consisted of a square wooden frame (40 cm

length x 41.2 cm height) with two horizontal supports (20 cm width) (Fig. 1). The frame was placed

55 cm from the entrance of the test box, centrally such that there was 22 cm between it and the

test box wall on each side. A reward feeder filled with grains was placed directly behind the barrier,

and a petri dish containing two mealworms was placed directly behind the feeder. An opaque

barrier was used during the training trials and a transparent barrier was used during the testing trials,

so the reward food was visible to the birds through the barrier during the test session (Fig. 1). Both

barriers had a strip of green tape on the left side and a strip of yellow tape on the right side (the

same colours as the enclosure and reward feeders).

Procedures. This task included a training session and a testing session on two separate,

consecutive days.

During the training session, the opaque barrier was attached to the frame. During the first

trial, the bird was given two minutes to detour around the barrier to reach the food; from the

second trial onwards, it had one minute and a half. Trials were repeated until the bird successfully

detoured on three consecutive trials, with a maximum of seven trials per bird. If the bird did not

detour once within three consecutive trials, the session ended. All birds passed the training session.

The testing session on the following day consisted of one training trial and one test trial per

bird. The test trial followed the same procedure as during the training trials, but the transparent

barrier was used instead of the opaque barrier.

'Successful' trials were defined as when the bird detoured around the barrier (opaque or

transparent) without interacting (i.e., pecking or pushing) with it.

Cylinder task

     Apparatus. The cylinder apparatus consisted of a transparent Plexiglas tube with open ends (5.4 cm diameter x 13.2 cm length) mounted on a small wooden platform (13 cm length x 3 cm width x 2 cm height; Fig. 1). The wooden base was covered with horizontal strips of yellow and green tape. The apparatus was placed 40 cm from the entrance of the test box, so that the entrances were perpendicular to the start box and secured to the floor with tape. One freshly killed mealworm and 1tsp of grain were placed centrally inside the cylinder. Quails had to put their head within the cylinder to retrieve the food.

     Procedures. The Cylinder task occurred during a single test day. During each trial, the bird had two minutes to reach the food inside the cylinder. Trials were repeated until the bird completed ten trials, which was the maximum number of trials given, or until the bird failed to reach the food for three consecutive trials.

     'Successful' trials were defined as when the bird reached the food through one of the two open sides of the cylinder without first pecking at the outside of the cylinder. For the purpose of this study, we will only focus on the first trial, as we believe it best measure the stopping behaviour of an individual (52). Indeed, the subsequent trials might reflect the inhibitory learning of the individual and is investigated in a different study (53).

Behavioural measures of response inhibition tasks

     For each quail, the dependent measures evaluated were (1) trial performance (i.e., whether the trial was a success), and (2) time spent 'interacting' with the barrier/cylinder (i.e., time the bird spends actively and repeatedly pecking/pushing the barrier).

Figure 1. Apparatuses used for the *Barrier task* (left) and for the *Cylinder task* (right) with Japanese quails (*n* = 120).

Impulsive aggression tasks

<u>Familiar Group</u>

Apparatus. A feeder, identical to the one present in the home enclosure of individuals and filled with grains, was placed in the middle of the text box. A sheet of semi-transparent plastic was placed inside the feeder to control access to the food. The feeder was present through the duration of the trial, but access to the food was adjusted during the trial (see Procedures).

Procedures. Five quails from the same enclosure were tested together for two trials on two consecutive days. The five individuals were placed in the start box and, following the same procedure as during the previous tasks, were let into the test box after 15 seconds after the opaque door of the start box was removed. Trials lasted 10 minutes. During the first trial, access to the feeder was completely thwarted for the first five minutes of the trial (No Food phase), then partially thwarted so that only one individual could access the food for the last five minutes of the trial

(Limited Access phase). During the second trial, the procedure was reversed (i.e., Limited Access

phase during the first five minutes, No Food phase during the last five minutes).

The same set of five birds were tested in the task. Repeated trials and different experimental

phases allowed us to investigate whether quails' aggressive behaviour was repeatable across

contexts differing in food availability. This aspect is being investigated in a different project and

not analysed in this study (Vernouillet et al. in prep.). Here, we focused on the behaviour of

individuals during the first trial in this study.

<u>Unfamiliar Individual</u>

280
Apparatus. Similar to the Familiar Group task, a feeder, identical to the one present in the

home enclosure of individuals and filled with grains, was placed in the middle of the test box. A

sheet of semi-transparent plastic was placed inside the feeder to control access to the food. The

feeder was placed inside the test box during the trial (see Procedures).

Procedures. Two quails from a different enclosure and loft (but of the same experimental

condition) were tested together for three trials on three consecutive days. Two individuals in the

start box, and following the same procedure as during the precedent tasks, were let into the test

box after 15 seconds from removing the opaque door of the start box. Trials lasted three minutes.

The feeder was placed in the test box after one minute. During the first trial, only one bird had

access to food when the feeder was placed in the test box. During the second trial, both birds could

freely access the food when the feeder was placed in the test box. During the third trial, none of

the birds had access to the food, as access was thwarted.

The same set of two birds were tested for each trial in the task. Repeated trials and different

experimental phases allowed us to investigate whether quails' aggressive behaviour was repeatable

across contexts differing in food availability. This aspect is being investigated in a different project

and not analysed in this study (Vernouillet et al. in prep.). Here, we focused on the behaviour of

individuals during the first trial in this study.

<u>Behavioural measures of impulsive aggression tasks</u>

For each quail, the dependent measures evaluated were (1) number of pecks given, (2)

number of pecks received, (3) time spent chasing, (4) time spent escaping, and (5) time spent

pushing. Descriptions of each behaviour can be found in Table S1.

Statistical analyses

We first wanted to perform a structural equation modelling approach (SEM) to determine

whether response inhibition, as estimated by the measures assessed during the Barrier and the

Cylinder tasks, could predict impulsive aggression, as estimated by the behavioral measures

assessed during the Familiar Group and the Unfamiliar Individual tasks. However, a preliminary

Confirmatory Factor Analysis indicated a lack of correlations among measures assessing the latent

factors of interest (i.e., response inhibition and impulsive aggression), and none of the models

tested had a decent fit (see Supplementary Material, Figures S2 and S3, Table S2 for more details

on factors analyses and SEM).

We thus used a different approach for our analyses. Importantly, unlike the SEM approach,

this approach does not allow us to infer the direction of the potential link between response

inhibition and impulsive aggression. First, to determine whether both the Barrier task and the

Cylinder task were assessing the same aspect of response inhibition, we correlated (1) first trial

performance using a tetrachoric correlation (for binary data), and (2) time spent 'interacting' with

the barrier/cylinder using a Pearson's correlation. Second, to determine whether both impulsive

aggression tasks were assessing the same aspect of aggression, we conducted a PCA with (1)

number of pecks given, (2) number of pecks received, (3) time spent chasing, (4) time spent

escaping, and (5) time spent pushing for both tasks. PCA was based on a correlation matrix. We

only considered components whose Eigenvalues were higher than 1 and for each component, we

only considered loadings whose contribution was higher than what was expected if all loadings

were of equal weight. Third, as we did not find any significant correlation between the different

response inhibition variables (see Results), we assessed whether higher levels of impulsive

aggression were linked to response inhibition deficits by evaluating whether aggression scores (as

calculated during the PCA) significantly differ between individuals that were successful during the

first trial of each response inhibition task using a Student t-test and whether they correlated with

time spent 'interacting' with the barrier/cylinder using a Pearson's correlation.

To determine whether early-life social environment influenced an individual's response

inhibition and/or impulsive aggression, we used a generalized linear model (GLM) approach, with

(1) trial performance (i.e., whether the barrier/cylinder testing trial was successful) for each

response inhibition task separately, and (2) time spent 'interacting' with the barrier/cylinder (i.e.,

time the bird spends actively and repeatedly pecking/pushing the barrier) for each response

inhibition task separately, as dependent variables associated with response inhibition, and (3) using

the PCA scores obtained for the first two axes, as aggressive behaviours during the Unfamiliar

Individual task mostly loaded on PC1 and aggressive behaviours during the Familiar Group task

mostly loaded on PC2 (see Results) as dependent variables associated with impulsive aggression.

Full models included social group condition, as well as morphological factors such as body weight

and sex.

To assess whether specific factors influenced our dependent variables, we compared full

models to a reduced model that did not include the factor that was evaluated (following the

procedures described by (54,55)). Parameter estimation was achieved using the residual maximum

likelihood ratio test.

For trial performance during the Barrier and the Cylinder tasks, we used a binomial

distribution. For time spent interacting with the barrier/cylinder, we rounded the values to the

nearest second and used a Poisson distribution. We first checked the assumptions of the GLMs

were met using the *DHARMa* package (56). Models were checked for dispersion and zero-inflation

using residual plots. In case residuals were over-dispersed, we added a dispersion parameter (this

was the case for the GLM on the second impulsive aggression score). In case there were too many

zero in our dataset, we added a zero-inflated model with a constant at the origin, and in the case

that the assumptions of the models were still not met, we changed the family to a negative binomial

family (this was the case for the GLM on time spent pecking for both response inhibition tasks).

Analyses were conducted using R version 4.3.2 (57) with the *glmmTMB* (58), *psych* (59),

FactoMineR (60), and *factoextra* (61). Descriptive statistics were obtained using the *Rmisc* package

(62). Plots were created using the *ggplot2* (63), *ggsignif* (64), *ggpubr* (65), *patchwork* (66), *wesanderson* (67)

packages. Alpha was set at 0.05 for all statistical analyses. All statistical analyses were performed by

AV, verified by KW and CAT.

4. Results

Overview of response inhibition tasks

Overview of Barrier task

During the testing trial, 19 individuals detoured successfully (i.e., without touching the

transparent barrier), 93 individuals detoured unsuccessfully (i.e., after touching the transparent

barrier), seven individuals touched the barrier but did not detour during the entire duration of the

trial, and one did not interact with the apparatus nor detoured around the transparent barrier (view

Table S3 for an overview of the response outcomes of the quails based on their social group and

on sex). Among individuals who pecked the barrier ($n = 100$), average time spent pecking the

barrier was 17.5 ($\pm SE$: 1.4) seconds.

Overview of Cylinder task

During the first testing trial, 14 individuals detoured successfully (i.e., without touching the

transparent cylinder), 46 individuals detoured unsuccessfully (i.e., after touching the transparent

cylinder), 36 individuals touched the cylinder but did not detour, and 24 did not interact with nor

detoured around the cylinder (view Table S3 for an overview of the response outcomes of the

quails based on their social group and on sex). Among individuals who pecked the cylinder ($n =$

82), average time spent pecking the cylinder was 15.0 ($\pm SE$: 1.6) seconds.

Repeatability of response inhibition

There were 88 individuals who were not successful during either response inhibition task,

18 individuals who were successful during the Barrier task only, 13 individuals who were successful

during the Cylinder task only, and only one individual who was successful during both response

inhibition tasks.

For this analysis, only individuals who interacted (either by detouring around and/or by

pecking the transparent barrier) during both tasks were included (*n* = 95). Results from the

correlations indicated that both tasks were measuring different aspects of RI, as neither

performance between tasks (tetrachoric correlation coefficient: -0.250) nor time spent interacting

with the transparent barrier/cylinder (Pearson's correlation: *t* = -1.129, *df* = 93, *p* = 0.262, Fig. 2)

correlated between tasks.

Figure 2. Time spent pecking at the transparent barrier during the Barrier task did not significantly

correlate with the time spent pecking the cylinder during the Cylinder task in Japanese quails

(Pearson's correlation: *t* = -1.129, *df* = 93, *p* = 0.262).

Overview of impulsive aggression tasks

<u>Overview of Familiar Group task</u>

All 120 individuals participated during the Familiar Group task. Quails gave on average 8.2

$\pm$ 16.7 pecks, received on average 8.2 $\pm$ 9.0 pecks, chased another individual for 1.0 $\pm$ 5.0 seconds,

escaped for 1.7 $\pm$ 4.4 seconds, and pushed others for 2.3 $\pm$ 6.0 seconds.

<u>Overview of Unfamiliar Individual task</u>

All 120 individuals participated during the Unfamiliar Individual task. Quails gave on

average 2.5 $\pm$ 5.3 pecks, received on average 2.5 $\pm$ 5.3 pecks, chased another individual for 1.8 $\pm$

5.7 seconds, escaped for 3.3 $\pm$ 7.5 seconds, and pushed the other for 0.3 $\pm$ 0.9 seconds.

<u>Repeatability of impulsive aggression</u>

All individuals were included in the analysis, as they all participated in both tasks ($n = 120$).

Results of the PCA indicated that there was not a unique impulsive aggression aspect measured

during the tasks.

The first five components had an Eigenvalue higher than 1 and together explained 78.1 %

of the variance in the data. The first component explained 21.7 % of the variance, the second

component explained 18.6 % of the variance, the third component explained 14.8 % of the variance,

the fourth component explained 12.5 % of the variance, and the fifth component explained an

additional 10.5 % of the variance.

Additionally, we found that both tasks investigated a different impulsive aggression aspect,

as measures related to each task loaded on different components (Table 1). Interestingly, number

of pecks given, amount of time spent chasing, and amount of time spent pushing loaded on the

opposite of number of pecks received and amount of time escaping on the two first components,

suggesting that individuals engaging in aggressive behaviors were receiving aggressive behaviors

from other individuals.

Table 1. PCA loadings for the measures on impulsive aggression in Japanese quails. Loadings

considered important (i.e., whose contribution was higher than what was expected if all loadings

were of equal weight) are indicated in bold.

		Dimensions				
		1	2	3	4	5
Familiar Group	Number of pecks given	0.044	**-0.737**	**0.440**	0.176	0.186
	Time spent chasing	0.148	**-0.644**	0.191	0.300	**0.530**
	Time spent pushing	-0.036	-0.264	**0.606**	0.070	**-0.626**
	Number of pecks received	0.254	**0.563**	**0.488**	0.020	0.310
	Time spent escaping	**0.483**	**0.509**	0.128	0.082	**0.346**
Unfamiliar Individual	Number of pecks given	**0.716**	-0.036	-0.271	**0.457**	-0.148
	Time spent chasing	**0.493**	-0.183	**-0.553**	**0.382**	-0.170
	Time spent pushing	**0.515**	0.310	**0.491**	0.279	-0.264

| | Number of pecks received | **-0.624** | 0.280 | 0.058 | **0.599** | -0.010 |
| | Time spent escaping | **-0.662** | 0.211 | -0.059 | **0.574** | 0.053 |

For the subsequent analyses, we used the individual scores obtained on the first two

components, as they best represent the aggression scores during each task (PC1 for Unfamiliar

Individual, PC2 for Familiar Group).

Relationship between response inhibition and impulsive aggression

We evaluated whether the different response inhibition behavioural measures were related

to the aggression scores of the individuals. For these analyses, we used reduced datasets that

included quails that interacted with the apparatuses (either by detouring successfully or by pecking

at the apparatus) either during Barrier task ($n = 119$) for measures associated to that task or during

the Cylinder task ($n = 96$) for measures associated to that task.

Aggression scores for the Unfamiliar Individual task (PC1) significantly correlated with the

amount of time spent pecking on the cylinder during the Cylinder task, with individuals that spent

more time pecking at the cylinder having a higher aggression score than individuals that spent less

time pecking at the cylinder ($t = 2.199$, $d.f. = 94$, $p = 0.030$; Fig. 3) and tended to be higher in

unsuccessful individuals in comparison to successful individuals ($t = 2.072$, $d.f. = 18.08$, $p = 0.053$).

Response inhibition measures assessed during the Barrier task did not significantly explain the

aggression scores for that task (Performance during the Barrier task: $t = -0.090$, $d.f. = 38.55$, $p =$

0.929; Time spent pecking the transparent barrier: $t = -0.197$, $d.f. = 117$, $p = 0.844$).

Figure 3. Time spent pecking on the cylinder during the Cylinder task significantly predicted the

aggression scores obtained during the Unfamiliar Individual task.

Aggression scores for the Familiar Group task (PC2) were significantly linked to the

performance during the Barrier task, with individuals that successfully detoured during the Barrier

task being more aggressive than individuals that were unsuccessful (t = 2.757, $d.f.$ = 20.79, p =

0.012; Fig. 4). No other response inhibition measures significantly influenced the aggression scores

for this task (Performance during the Cylinder task: t = 1.588, $d.f.$ = 20.54, p = 0.128; Time spent

pecking the transparent barrier: t = -1.757, $d.f.$ = 117, p = 0.082; Time spent pecking the transparent

cylinder: t = -1.645, $d.f.$ = 94, p = 0.103).

Figure 4. Performance during the Barrier task significantly predicted the aggression scores obtained

during the Familiar Group task. Note that for this aggression score, negative scores indicate more

aggressive behaviours than positive scores.

Influence of group size on response inhibition and impulsive aggression

Influence of Group size on response inhibition

Given the lack of contextual repeatability, we evaluated the effects of Group Size, Sex, and Body

Weight on performance and on the time spent interacting with the transparent barrier/cylinder for

both Response Inhibition tasks separately.

Only individuals who interacted (either by detouring around and/or by pecking the

transparent barrier) were included for the analyses on performance (n = 119 for the Barrier task, n

= 96 for the Cylinder task; see Table S3 for overview of number). Performance during the Barrier

484   task and performance during the Cylinder task were not explained by Group Size, Sex, nor Body

485   Weight (Table 2).

487   Table 2. Factors evaluated to explain performance during the Barrier task (n = 119 quails) and

488   during the Cylinder task (n = 96 quails).

	Performance Barrier task			Performance Cylinder task		
	χ^2	d.f.	p	χ^2	d.f.	p
Group Size	0.732	1	0.392	0.421	1	0.517
Sex	3.725	1	0.054	1.410	1	0.235
Body Weight	1.298	1	0.255	0.002	1	0.965

490
491

Only individuals who pecked at the transparent barrier were included for the analyses on

time spent pecking (n = 100 for the Barrier task, n = 82 for the Cylinder task; see Table S3 for

overview of number). Time spent pecking at the transparent barrier during the Barrier task was not

explained by Group Size (χ^2 = 1.608, d.f. = 1, p = 0.205; Fig. 5) nor by Sex (χ^2 = 0.063, d.f. = 1, p

= 0.802). However, Body Weight significantly influenced time spent pecking at the transparent

barrier, with individuals with a lower weight spending more time interacting with the barrier than

individuals with a higher weight (χ^2 = 5.835, d.f. = 1, p = 0.016; Fig. S4).

Figure 5. Left. Time spent pecking at the transparent barrier during the Barrier task did not
significantly differ between social groups (χ^2 = 1.608, $d.f.$ = 1, p = 0.205). Right. Quails in the Large
group spent significantly less time pecking at the transparent cylinder than quails in the Small group
(χ^2 = 6.256, df = 1, p = 0.012).

Group Size significantly influenced the time spent pecking at the transparent cylinder
during the Cylinder task, with individuals in the Large group spending less time pecking at the
transparent cylinder than individuals in the Small group (Time spent pecking at the transparent
cylinder $M \pm SE$: Large group: 11.64 ± 1.92 s, Small group: 18.70 ± 2.50 s; χ^2 = 6.256, df = 1, p =
0.012; Fig. 5). Body Weight also significantly influenced time spent pecking at the cylinder, with
individuals with a higher weight spending more time interacting with the cylinder than individuals
with a lower weight (χ^2 = 14.268, $d.f.$ = 1, p < 0.001; Figure S4). There was no significant difference
between males and females (χ^2 = 3.719, $d.f.$ = 1, p = 0.054).

Influence of Group size on impulsive aggression

517   Given the lack of repeatability in impulsive aggression between contexts, we evaluated the

518 effects of Group Size, Sex, Body Weight on the aggressive scores for both impulsive aggression

519 tasks separately (n = 120 quails).

520   Aggression scores for the Unfamiliar Individual task (PC1) were not explained by Group

521 Size (χ^2 = 0.001, $d.f.$ = 1, p = 0.973; Fig. 6) nor by Body Weight (χ^2 = 1.178, $d.f.$ = 1, p = 0.278).

522 However, male quails had a significantly higher aggression score than female quails (Females: -0.28

523 ± 1.54, Males: 0.28 ± 1.38; χ^2 = 4.318, $d.f.$ = 1, p = 0.038; Figure S5).

524   Aggression scores for the Familiar Group task (PC2) were not explained by Group Size (χ^2

525 = 0.183, $d.f.$ = 1, p = 0.670; Fig. 6). However, males were significantly more aggressive than females

526 (Females: 0.35 ± 0.97, Males: -0.35 ± 1.61; χ^2 = 8.554, $d.f.$ = 1, p = 0.004; Figure S5), and quails

527 with a higher weight were significantly more aggressive than quails with a lower weight (χ^2 = 6.002,

528 $d.f.$ = 1, p = 0.016; Figure S6).

531 Figure 6. There were no significant differences in aggression scores between social groups a) during

532 the Unfamiliar Individual (χ^2 = 0.001, $d.f.$ = 1, p = 0.973), and b) during the Familiar Group task

533 $(\chi^2 = 0.183, d.f. = 1, p = 0.670)$. *Note: for the axis PC2, more aggressive individuals have a negative

534 score.

5. Discussion

The main goals of this study were threefold: (A) to examine the link between response inhibition and impulsive aggression at an individual level, and (B) to assess whether group size during early-life environment influenced the development of response inhibition and (C) impulsive aggression. We found (A) some significant correlations between measures of response inhibition and impulsive aggression, providing some limited support to the theoretical model linking both concepts. We also found (B) evidence that early-life social environment indeed shapes the development of response inhibition, as quails raised in smaller social groups spent more time repeating an inefficient action (i.e., pecking at the cylinder) than those raised in larger social groups. However, contrary to what we expected, (C) we found no evidence that early-life social environment influenced the development of impulsive aggression.

Relationship between response inhibition and impulsive aggression

In Japanese quails, we found some evidence that response inhibition was significantly linked to impulsive aggression at an individual-level. More specifically, increased perseveration on an unrewarded action towards an unfamiliar object (i.e., time spent pecking at the transparent cylinder) was significantly correlated to increased aggression towards unfamiliar individuals. This result may suggest that individuals tend to interact aggressively towards novelty, be it an object or an individual. The link between higher aggression and lower inhibition has also been found in previous observational studies in mammals (12,13,68,69), as suggested by theoretical models (14). In primates, aggressive individuals with low response inhibition are thought to have an advantage to access and retain food resources (70). A similar observation was found in Australian magpies, where individuals involved in aggressive interactions (both initiating and receiving) performed worse in an associative learning task, possibly due to lower time and/or energy to attend the task (71). While

the authors did not test inhibition directly, previous work in that study population has found a

significant correlation between inhibition (as assessed with a cylinder) and associative learning (31).

Together, these results highlight how the social environment, its dynamics, and cognitive

performance are tightly linked with one another.

Interestingly, we also found that individuals that successfully fully cancelled a discrete action

(i.e., by detouring without pecking at the transparent barrier) had significantly increased levels of

aggression towards familiar individuals. This result goes in the opposite direction to what we

predicted, and contrast with previous studies that investigated the link between inhibition and

aggression. One potential explanation for this seemingly contradictory result could be by viewing

the adaptive significance of aggression (3). Indeed, aggressive individuals are more likely to initiate

contests and to subsequently win them (72), which potentially gives them higher fitness payoffs.

Hence, individuals with better response inhibition may use the most advantageous coping strategy

in response to the social environment demands.

These contradictory results may suggest three things: a) that the mechanisms involved during

the cancellation of an action may depend on whether the individuals are already engaged in the

action they need to inhibit, and/or b) that the tasks are not assessing response inhibition or

aggression, but rather pecking tendency (whether it is towards an object or an individual), and/or

c) that response inhibition and aggression depends highly on the social context.

Indeed, when examining in detail the requirements of both the Detour and the Cylinder tasks,

we observe that individuals may have to inhibit two actions: first, the prepotent response of going

directly at the food, thus pecking at the transparent barrier, and second, the repetitive action of

pecking along the barrier if they failed to inhibit the first action (73). Not only the nature of the

action to be stopped, but the overall task context in which actions need to be stopped could thus

involve different mechanisms.

The lack of correlations between the different measures of the two response inhibition tasks

we observed is consistent with results from previous studies who have used the same tasks (e.g., in

dogs: (52,74,75); in pheasants: (76)). This occurred despite both tasks having similar task demands

in general (i.e., detouring tasks requiring individuals to detour around a transparent barrier, (47))

and despite both tasks having a similar "go" stimulus (i.e., the visible food reward) and a similar

"stop" stimulus (i.e., the transparent barrier) present at the same time (for a recent in-depth

discussion about the importance of the "go" and 'stop" stimuli in response inhibition tasks, please

refer to (73)). Rather, the characteristics of the stop stimulus itself, such as the shape (e.g., (77))

and the motor action required to successfully detour, might influence the individual's response: the

transparent barrier in the Barrier task is large and requires individuals to move their entire body

around it, whereas the transparent barrier in the Cylinder task is small and requires individuals to

place their head through the openings (52).

Alternatively, the lack of repeatability in performance might be due to training differences.

Indeed, for the Barrier task, individuals were trained to go around the opaque barrier and might

have learned the detouring solution (78), explaining the high success of individuals in this task

during the testing trial. In comparison, individuals were not trained during the Cylinder task. Hence,

during the Cylinder task, individuals must not only learn to inhibit pecking at the transparent

cylinder, but also to successfully "solve" the task (52). This may also explain why the number of

individuals who successfully detoured during the Cylinder task was lower than the ones who did

during the Barrier task.

Interestingly, the lower detouring performance during the Cylinder task, i.e., the second

testing trial involving detouring around a transparent barrier, suggest that individuals were unable

to use their previously experience of detouring around a transparent barrier in a different context.

Unlike what was found with pheasants by van Horik et al. (76), quails did not learn to inhibit

interacting with a transparent surface between the two tasks, as the duration of pecking at the

transparent surface did not differ between tasks. However, we found evidence that Japanese quails

did learn to inhibit pecking with repeated trials during the Cylinder task (53).

Even when carefully choosing tasks purporting to assess the same aspect of response

inhibition, there is a lack of repeatability in an individual's behavioural response. Understanding

how and why exactly there is such a low reliability between tasks is beyond the scope of the current

study but warrants further investigation (79). Yet, together, these results suggest that, despite using

similar tasks, response inhibition is multi-faceted and highly dependent on the action that needs to

be stopped and/or on the context in which stopping need to occur.

Similarly, the contradictory results we obtained regarding the link between aggression and

response inhibition may be due to the contextual differences between both aggression tasks. The

lack of repeatability in individual aggression scores is surprising as it goes against the idea of

aggression as a personality trait (e.g., (80–82)). Instead, this result suggests that aggression is subject

to high intra-individual variation (83). Plasticity in aggressive responses could depend on the

outcome of past antagonistic interactions (84) and on the traits of the other individual, such as

status signals or signals of fighting ability (reviewed by (85)) or use social information to assess the

costs of engaging in an antagonistic behaviour towards another individual (86). Individuals'

behaviour may also differ when they are in the presence of familiar and of unfamiliar individuals

(e.g., (87), "winner and loser effects" – (88), "dear enemy" hypothesis – as reviewed by (89); but

see (44) for a counter-example with male Japanese quails). Indeed, because agonistic interactions

between individuals can be costly, familiarity is thought to play a key role in the formation of

hierarchies, both within species (90) and between species (91). For this reason, aggression is usually

reduced within the group once hierarchies are established (3). Regardless of the lack of correlations

in aggressive behaviours between tasks, we found that behaviours associated with individuals that

were the target of aggressive behaviours (as assessed by the time spent escaping and the number

of pecks received) were less likely to engage in aggressive behaviours (as assessed by the time spent

chasing and pushing, and the number of pecks given) in each task context.

Finally, another methodological aspect that may explain this duality in our results linking

response inhibition and aggression is the use of food rewards in a group context during the

aggression tasks and in a solitary context during the response inhibition tasks. Indeed, the social

context may exert different inhibitory pressures on individuals, despite having similar motivation

levels in both cases. For instance, in the presence of a more dominant or more aggressive group

member, an individual might inhibit going towards the food reward to reduce the risk of receiving

directed aggression (92). However, when alone, that same individual may or may not have lower

response inhibition levels towards the food rewards. This possibility suggests, once again, the

importance of accounting for the social context when evaluating impulsivity.

In sum, we found evidence that response inhibition is linked to impulsive aggression. But

given that the relationship is not as clear cut as we expected and that we were unable to assess the

direction of that relationship due to the lack of correlations between tasks, our conclusions

regarding the link between the two are not as definitive as we had hoped them to be. For instance,

we cannot exclude the possibility that other factors might mediate the relationship between the

two (19) nor can we definitely say that aggression results from deficits in response inhibition. Next,

we explored whether social environment during early life might contribute to deficits in response

inhibition and/or to impulsive aggression.

Influence of early-life social environment on response inhibition and impulsive aggression

The importance of the social environment on the development of cognition has been

suggested in many species (20,21,23,30,71,93). More specifically, social group size, a key

component of the social environment, usually correlates with higher levels of inhibition in hyenas

(32) and in Australian magpies (31), but not in guppies (34,94) nor in horses (95). Here, we found

that quails raised in larger groups inhibited a repetitive unrewarded action (i.e., pecking at the

transparent cylinder) faster, on average, than quails raised in smaller groups. This result aligns with

those from previous studies evaluating intraspecific variability in response inhibition using the

Cylinder task on Australian magpies and on hyenas (31,32), and support what we predicted.

However, we found no significant effect of social group size on the overall performance for both

tasks, nor on the time spent pecking on the transparent barrier during the Barrier task. As we

previously mentioned, both tasks may not measure the same aspect of response inhibition or may

measure pecking tendency in one or the other. Therefore, the influence of the social environment

might be mediated differently depending on which aspect of response inhibition is considered.

Early-life social environmental conditions have strong effects on aggressive behaviour in

many species, including birds (96), farm animals (97), and humans (e.g., (42)), with usually a

decrease in aggression as group size increases. For quails specifically, they are more aggressive when

they are raised in social isolation than when they have been raised with others (98). Unlike previous

research, we found no significant difference in levels of aggression between groups during our

aggression tasks. Hence, we found no evidence that quails become more socially tolerant with

increasing group size (e.g., (37,40,49)), even if admittedly we did not assess the aggression

experienced by individuals outside of the experimental context. However, this lack of significant

results might be caused by the increase in density associated with the increase in group size, as

individuals of both conditions were given the same size enclosure. Higher population density has

been associated with higher aggression levels (e.g., lizards – (99)). Hence, in our study, the potential

reduction in aggression due to group size might be confounded with increase in aggression due to

increase in density and might have switched to a resource defense strategy instead. A next step is

to disentangle the effects of group size and group density on aggression as well as on response

inhibition. Alternatively, differences in social group size can have affected social motivation (i.e.,

the motivation to seek proximity with other group members) and social bonding in quails. Indeed,

Japanese quails raised in larger social group show less positive social interactions (100), potentially

because their ability to recognize familiar individuals and remember past interactions with them

decrease as the number of group members increase (30).

Social group size is not the only aspect of social complexity. Group dynamics and stability,

as well as the nature of the relationships between group members, are other important aspects that

contribute to the variation in social complexity within and between groups (29). Further

investigation is needed to determine which aspects of the social life influence the development of

aggression and of response inhibition.

Intra- and inter-individual variation in response inhibition and in impulsive aggression

Research on non-human animals had, until recently, tried to minimize the importance of

inter-individual variation in cognition (101,102). Now, there is a general effort in pinpointing which

factors may underlie individual differences in cognition generally (103,104). Indeed, despite

controlling the early-life physical and social environment of our quails, as well as the acclimatation,

habituation and testing procedures, we still observed pronounced individual differences in

performance and in the time spent performing a repetitive unrewarded action during the response

inhibition tasks and in the aggressive behaviours during the impulsive aggression tasks that were

not explained by the difference in group size. Here, we can only speculate about the causes of these

differences in our sample.

Inter-individual differences in response inhibition and impulsive aggression, as well as the

lack of repeatability across tasks, may be due to differences in motivation and attention (105,106).

Once again, despite controlling the food deprivation procedure to ensure similar motivation to

participate in each task, some individuals might have greater motivation than others, due to higher

basal energetic demands (107) or lower availability of food in their home enclosure (108). This

might also explain why we found an effect of body weight on time spent pecking at the transparent

barrier (also found in New Zealand robins, (109)). However, given that the direction of that effect

was opposite for both response inhibition tasks, we cannot confirm this explanation as to why or

how it might be the case. Additionally, in group-living domestic species, more aggressive individuals

often outcompete less aggressive individuals for limited resources and have a higher weight as a

result (e.g., (110)), which support our findings on heavier individuals being more aggressive towards

familiar individuals. Concurrently, being aggressive might be less costly and more beneficial for

larger individuals, as they are more likely to physically outcompete smaller ones.

Cognitive abilities and personality traits have also been known to differ between females and

males (111). Previous work on mosquitofish (112), baboons (113), dogs (114), and guppies (115),

has shown that females are significantly better than males at inhibiting their responses. Yet this was

not the case in our quails nor in other species tested, such as Australian skinks (107), spotted

hyaenas (32), gulls (73), and red junglefowls (116). Despite not finding any differences in response

inhibition between males and females, we found that males were significantly more aggressive than

females. Indeed, early studies suggest that most aggressive behaviors are initiated by sexually mature

males towards other males and females (117). For this reason, sexually experienced females tend

to avoid relatively aggressive males (118), as cohabitation with aggressive males lead to increased

baseline cortisol levels (119), lower egg quality (120), and lower plumage quality (44), amongst

others. Aggressive behaviors can also be initiated by sexually mature females when paired with an

unfamiliar individual (both males and females), but their aggression rates decreased quickly over

time, which was not the case for males (121). Here, interestingly, we found such a sex difference in

aggression when the quails were between 5- and 6-weeks old, at the onset of sexual maturity in

Japanese quails (between 6- to 8-weeks old). Little is known regarding aggression in immature, non-

sexually active or castrated quails, but it is usually assumed they have lower aggressive tendencies

than sexually mature ones (43,122). Hence, inter-individual variation in aggression that we observed

might also be due to inter-individual differences in reaching sexual maturity, despite keeping all

other parameters constant for all individuals (photoperiod, temperature, food).

Personality traits, especially neophobia (i.e., the fear to interact with novelty) and exploration

(i.e., the willingness to interact with novelty, (123)) have also been suggested to explain differences

in cognitive abilities ((124), also reviewed in (125)), or even the motivation to participate in the

tasks or interact with the apparatuses (e.g., self-selection by participation of less neophobic

individuals in the population, (126,127)). To a lesser extent, neophobia and exploration could also

explain some inter-individual behavioural differences we observed during the aggression tasks, as

less neophobic and/or more exploratory individuals might approach unfamiliar individuals and

interact with them. For instance, more exploratory great tits (*Parus major*) are more likely to initiate

fights (128). Yet, whether personality traits derive from cognitive abilities, or *vice versa*, remains

debated in the field of animal cognition (and even between the authors of the current paper).

Finally, a large inter-individual variation might indicate a lack of strong selection on a given

trait. For instance, there was no evidence that response inhibition contributed to social rank,

foraging efficiency, or risk taking in red junglefowl females (116), suggesting that there might be a

lack of strong selection on response inhibition, potentially explaining why there is such strong

variation in response inhibition between individuals. Alternatively, a large inter-individual variation

in response inhibition within the population might indicate that inhibiting a behavioral response

might be favored in certain environments or contexts, but not in others (129,130). For example,

lower-ranked individuals, especially in large social groups, may benefit from having a higher level

of response inhibition, possibly to reduce conflicts with higher-ranked individuals (32). Similar

arguments can be made to explain intra-individual variation in response inhibition and in aggression.

Indeed, the lack of repeatability in response inhibition at an individual-level may suggest that

individuals are flexible in their cognitive ability and in their social behavior, depending on the

context. For instance, impulsive actions might be more adaptive in an unstable or unpredictable

environment or when travel time between resources increase (131). Understanding in which

contexts extreme values of response inhibition and if might be favored and why such variation in

cognitive traits and in personality exists within the population is a topic that warrants further

investigation.

6. Conclusion

In summary, our study provides one of the first experimental evidence that early-life social

environment can influence the development of response inhibition. We also experimentally

showed that certain aspects of aggression are linked to deficits in response inhibition, especially

regarding perseverating in performing an unrewarded action towards novelty (objects and/or

individual). Hence, our study shed some light regarding the mechanisms underlying impulsive

aggression using Japanese quails. Yet, contrary to what was expected based on previous literature,

group size, a key aspect of early-life social environment, did not influence aggression in our quails,

despite it influencing certain aspects of response inhibition (the same ones that were linked to

aggression). Aggression is a multifaceted and complex behaviour that involves multiple processes

(132,133). Thus, despite controlling the physical and social environment, intrinsic factors such as

genetics, motivation, non-social personality traits (i.e., neophobia, exploration, activity), and the

ability to get social information might explain inter-individual variability in aggression that we

witnessed and mediate the link between response inhibition and aggression. Regarding the

influence of sociality on the emergence of aggressive behaviors, other social factors, such as group

dynamics and stability, and further research is warranted to fully understand how sociality, response

inhibition, and aggression relate to one another.

Acknowledgements

We thank the staff of the Wildlife Rescue Centre Ostend for general project support and Harry Suter for insightful discussions and comments on the draft of the manuscript.

Ethical Statement

The experiment was performed in accordance with the Association for the Study of Animal Behaviour ethical guidelines under permission of the ethical committee of animal experimentation (VIB Site Ghent, Universiteit Gent): EC2022-005.

Funding Statement

Research support was provided by a BOF postdoc fellowship (#BOF.PDO.2021.0035.01) to AV, a FWO (Flemish Research Foundation) PhD Fellowship grant (No. 11F0823N) to AD, a CSC (China Scholarship Council) PhD Scholarship (No. 202106180023) to WZ, a Marie Skłodowska-Curie Action fellowship ('UrbanCog', project number: 101062662) under the European Union's Horizon Europe Programme to CAT, a FWO (Flemish Research Foundation) PhD Fellowship grant (No. 11P3G24N) to SK, an ERC Consolidator grant (European Union's Horizon 2020 research and innovation programme, grant agreement No 769595) to FV, and Methusalem Project 01M00221 (Ghent University) to FV, LL, and AM.

Data Accessibility

All videos of the experiment can be found on Zenodo (https://zenodo.org/records/13361318).

R codes and extracted data from BORIS can be found on OSF (https://osf.io/2b5uy/).

Competing Interests

None to declare.

Author contributions

Based on the CRediT system, the contributions of the authors are as follow:

'Conceptualization' – AV, LL, FV; 'Data curation' – AV, KW; 'Formal analysis' – AV; 'Funding

acquisition' – AV, AD, WZ, CAT, AM, LL, FV; 'Investigation' – AV, KW, RA, AD, WZ, CAT,

SK, LL, FV, 'Methodology' – AV, KW, RA, AD, LL, FV; 'Project administration' – AV, FV;

'Resources' – FV, 'Supervision' – AV, LL, FV; 'Validation' – AV, KW, SK, CAT; 'Visualization' –

AV, 'Writing—original draft' – AV, and 'Writing—review & editing' – AV, KW, RA, AD, WZ,

CAT, AM, LL, FV. In the Methods section, we followed the MERIT system (Nakagawa et al. 2023).

References

1.   Hamilton WD. Geometry for the selfish herd. J Theor Biol. 1971 May 1;31(2):295–311.
2.   Barratt ES, Stanford MS, Dowdy L, Liebman MJ, Kent TA. Impulsive and premeditated aggression: a factor analysis of self-reported acts. Psychiatry Res. 1999 May 31;86(2):163–73.
3.   Holekamp KE, Strauss ED. Aggression and dominance: an interdisciplinary overview. Curr Opin Behav Sci. 2016 Dec;12:44–51.
4.   Breuer J, Scharkow M, Quandt T. Sore losers? A reexamination of the frustration–aggression hypothesis for colocated video game play. Psychol Pop Media Cult. 2015;4(2):126–37.
5.   Osumi T, Nakao T, Kasuya Y, Shinoda J, Yamada J, Ohira H. Amygdala dysfunction attenuates frustration-induced aggression in psychopathic individuals in a non-criminal population. J Affect Disord. 2012 Dec 15;142(1):331–8.
6.   Tremblay RE. Developmental origins of chronic physical aggression: An international perspective on using singletons, twins and epigenetics. Eur J Criminol. 2015 Sep;12(5):551–61.
7.   Chester DS. Aggression as successful self-control. Soc Personal Psychol Compass. 2024 Feb;18(2):e12832.
8.   Verbruggen F, Logan GD. Response inhibition in the stop-signal paradigm. Trends Cogn Sci. 2008 Nov 1;12(11):418–24.
9.   Bari A, Robbins TW. Inhibition and impulsivity: Behavioral and neural basis of response control. Prog Neurobiol. 2013 Sep;108:44–79.
10.   Diamond A. Executive Functions. Annu Rev Psychol. 2013 Jan 3;64(Volume 64, 2013):135–68.
11.   Homberg JR, Pattij T, Janssen MCW, Ronken E, De Boer SF, Schoffelmeer ANM, et al. Serotonin transporter deficiency in rats improves inhibitory control but not behavioural flexibility: Serotonin transporter knockout and impulse control. Eur J Neurosci. 2007 Sep 25;26(7):2066–73.
12.   Gobbo E, Zupan Šemrov M. Dogs Exhibiting High Levels of Aggressive Reactivity Show Impaired Self-Control Abilities. Front Vet Sci. 2022 Mar 24;9:869068.

13.   Overduin-de Vries AM, Vermande MM, Hessen DJ, Sterck EHM. The ability to inhibit
impulses is related to social behavior in long-tailed macaques. Am J Primatol. 2024
Feb;86(2):e23587.

14.   Vigil-Colet A, Codorniu-Raga MJ. Aggression and inhibition deficits, the role of functional
and dysfunctional impulsivity. Personal Individ Differ. 2004 Nov;37(7):1431–40.

15.   Hoaken PNS, Shaughnessy VK, Pihl R o. Executive cognitive functioning and aggression:
Is it an issue of impulsivity? Aggress Behav. 2003;29(1):15–30.

16.   Van Den Bergh F, Spronk M, Ferreira L, Bloemarts E, Groenink L, Olivier B, et al.
Relationship of delay aversion and response inhibition to extinction learning, aggression, and sexual
behaviour. Behav Brain Res. 2006 Nov;175(1):75–81.

17.   Bacqué-Cazenave J, Bharatiya R, Barrière G, Delbecque JP, Bouguiyoud N, Di Giovanni
G, et al. Serotonin in Animal Cognition and Behavior. Int J Mol Sci. 2020 Feb 28;21(5):1649.

18.   Miyazaki K, Miyazaki KW, Doya K. The Role of Serotonin in the Regulation of Patience
and Impulsivity. Mol Neurobiol. 2012 Apr 1;45(2):213–24.

19.   Puiu AA, Wudarczyk O, Goerlich KS, Votinov M, Herpertz-Dahlmann B, Turetsky B, et
al. Impulsive aggression and response inhibition in attention-deficit/hyperactivity disorder and
disruptive behavioral disorders: Findings from a systematic review. Neurosci Biobehav Rev. 2018
Jul 1;90:231–46.

20.   Foraita M, Howell T, Bennett P. Environmental influences on development of executive
functions in dogs. Anim Cogn. 2021 Jul;24(4):655–75.

21.   Johnson D, Policelli J, Li M, Dharamsi A, Hu Q, Sheridan MA, et al. Associations of Early-
Life Threat and Deprivation With Executive Functioning in Childhood and Adolescence: A
Systematic Review and Meta-analysis. JAMA Pediatr. 2021 Nov 1;175(11):e212511.

22.   Kotrschal A, Taborsky B. Environmental Change Enhances Cognitive Abilities in Fish.
PLOS Biol. 2010 Apr 6;8(4):e1000351.

23.   Nettle D, Bateson M. Adaptive developmental plasticity: what is it, how can we recognize
it and when can it evolve? Proc R Soc B Biol Sci. 2015 Aug 7;282(1812):20151005.

24.   Triki Z, Zhou T, Argyriou E, Sousa De Novais E, Servant O, Kolm N. Social complexity
affects cognitive abilities but not brain structure in a Poecilid fish [Internet]. Animal Behavior and
Cognition;   2023   Aug   [cited   2024   Feb   7].   Available   from:
http://biorxiv.org/lookup/doi/10.1101/2023.08.20.554009

25.   Dunbar RIM, Shultz S. The Social Role of Inhibition in Primates and Other Animals
[Internet]. bioRxiv; 2022 [cited 2024 Dec 19]. p. 2020.10.26.354852. Available   from:
https://www.biorxiv.org/content/10.1101/2020.10.26.354852v3

26.   Alcala RS, Caliva JM, Flesia AG, Marin RH, Kembro JM. Aggressive dominance can
decrease behavioral complexity on subordinates through synchronization of locomotor activities.
Commun Biol. 2019 Dec;2(1):467.

27.   Papageorgiou D, Farine DR. Group size and composition influence collective movement
in a highly social terrestrial bird. Díaz-Muñoz SL, Rutz C, Díaz-Muñoz SL, editors. eLife. 2020
Nov 10;9:e59902.

28.   Byrne RW, Whiten A. Cognitive Evolution in Primates: Evidence from Tactical Deception.
Man. 1992;27(3):609–27.

29.   Kappeler PM. A framework for studying social complexity. Behav Ecol Sociobiol. 2019
Jan;73(1):13.

30.   Croney CC, Newberry RC. Group size and cognitive processes. Appl Anim Behav Sci. 2007

Mar;103(3–4):215–28.

31.   Ashton BJ, Ridley AR, Edwards EK, Thornton A. Cognitive performance is linked to group size and affects fitness in Australian magpies. Nature. 2018 Feb;554(7692):364–7.

32.   Johnson-Ulrich L, Holekamp KE. Group size and social rank predict inhibitory control in spotted hyaenas. Anim Behav. 2020 Feb;160:157–68.

33.   Amici F, Aureli F, Call J. Fission-Fusion Dynamics, Behavioral Flexibility, and Inhibitory Control in Primates. Curr Biol. 2008 Sep 23;18(18):1415–9.

34.   Lucon-Xiccato T, Montalbano G, Reddon AR, Bertolucci C. Social environment affects inhibitory control via developmental plasticity in a fish. Anim Behav. 2022 Jan;183:69–76.

35.   Færevik G, Andersen IL, Jensen MB, Bøe KE. Increased group size reduces conflicts and strengthens the preference for familiar group mates after regrouping of weaned dairy calves (Bos taurus). Appl Anim Behav Sci. 2007 Dec;108(3–4):215–28.

36.   Estevez I, Newberry RC, Keeling LJ. Dynamics of aggression in the domestic fowl. Appl Anim Behav Sci. 2002 Apr 2;76(4):307–25.

37.   Estevez I, Keeling LJ, Newberry RC. Decreasing aggression with increasing group size in young domestic fowl. Appl Anim Behav Sci. 2003 Dec;84(3):213–8.

38.   Syarifuddin S, Kramer DL. The effect of group size on space use and aggression at a concentrated food source in blue gouramis, Trichogaster trichopterus (Pisces: Belontiidae). Environ Biol Fishes. 1996 Jul 1;46(3):289–96.

39.   Turner SP, Horgan GW, Edwards SA. Effect of social group size on aggressive behaviour between unacquainted domestic pigs. Appl Anim Behav Sci. 2001 Nov;74(3):203–15.

40.   Pagel M, Dawkins MS. Peck orders and group size in laying hens: `futures contracts' for non-aggression. Behav Processes. 1997 Apr 1;40(1):13–25.

41.   Van Loo PLP, Mol JA, Koolhaas JM, Van Zutphen BFM, Baumans V. Modulation of aggression in male mice: influence of group size and cage size. Physiol Behav. 2001 Apr 1;72(5):675–83.

42.   Haller J, Harold G, Sandi C, Neumann ID. Effects of Adverse Early-Life Events on Aggression and Anti-Social Behaviours in Animals and Humans. J Neuroendocrinol. 2014 Oct;26(10):724–38.

43.   Caliva JM, Falkenburger Melleu F, Marino-Neto J, Marin RH, Kembro JM. Expression of aggressiveness modulates mesencephalic c-fos activation during a social interaction test in Japanese quail (Coturnix japonica). Behav Brain Res. 2019 Jul;367:221–9.

44.   Pellegrini S, Condat L, Caliva JM, Marin RH, Guzman DA. Can Japanese quail male aggressions toward a female cagemate predict aggressiveness toward unknown conspecifics? Livest Sci. 2019 Apr;222:65–70.

45.   Mills AD, Crawford LL, Domjan M, Faure JM. The behavior of the japanese or domestic quail Coturnix japonica. Neurosci Biobehav Rev. 1997;21(3):261–81.

46.   Millman ST, Duncan IJ, Widowski TM. Male broiler breeder fowl display high levels of aggression toward females. Poult Sci. 2000 Sep;79(9):1233–41.

47.   Kabadayi C, Bobrowicz K, Osvath M. The detour paradigm in animal cognition. Anim Cogn. 2018 Jan;21(1):21–35.

48.   van Horik JO, Langley EJG, Whiteside MA, Laker PR, Beardsworth CE, Madden JR. Do detour tasks provide accurate assays of inhibitory control? Proc R Soc B Biol Sci. 2018 Mar 28;285(1875):20180150.

49.   Samarakone TS, Gonyou HW. Domestic pigs alter their social strategy in response to social

group size. Appl Anim Behav Sci. 2009 Oct;121(1):8–15.

50. Friard O, Gamba M. BORIS: a free, versatile open-source event-logging software for video/audio coding and live observations. Methods Ecol Evol. 2016;7(11):1325–30.

51. McHugh ML. Interrater reliability: the kappa statistic. Biochem Medica. 2012 Oct 15;22(3):276–82.

52. Vernouillet AAA, Stiles LR, Andrew McCausland J, Kelly DM. Individual performance across motoric self-regulation tasks are not correlated for pet dogs. Learn Behav. 2018 Dec;46(4):522–36.

53. Willcox K, Vernouillet A, Lens L, Verbruggen F. Early-life group size influences response inhibition, but not the learning of it, in Japanese quails. Learn Behav [Internet]. 2024 Sep 19 [cited 2024 Nov 21]; Available from: https://link.springer.com/10.3758/s13420-024-00643-2

54. Harrison XA, Donaldson L, Correa-Cano ME, Evans J, Fisher DN, Goodwin CED, et al. A brief introduction to mixed effects modelling and multi-model inference in ecology. PeerJ. 2018 May 23;6:e4794.

55. Tredennick AT, Hooker G, Ellner SP, Adler PB. A practical guide to selecting models for exploration, inference, and prediction in ecology. Ecology [Internet]. 2021 Jun [cited 2022 May 18];102(6). Available from: https://onlinelibrary.wiley.com/doi/10.1002/ecy.3336

56. Hartig F, Lohse L, leite M de S. DHARMa: Residual Diagnostics for Hierarchical (Multi-Level / Mixed) Regression Models [Internet]. 2024 [cited 2024 Dec 19]. Available from: https://cran.r-project.org/web/packages/DHARMa/index.html

57. R Core Team. R: The R Project for Statistical Computing [Internet]. [cited 2024 Dec 19]. Available from: https://www.r-project.org/

58. Brooks M, Bolker B, Kristensen K, Maechler M, Magnusson A, McGillycuddy M, et al. glmmTMB: Generalized Linear Mixed Models using Template Model Builder [Internet]. 2024 [cited 2024 Dec 19]. Available from: https://cran.r-project.org/web/packages/glmmTMB/index.html

59. Revelle W. psych: Procedures for Psychological, Psychometric, and Personality Research [Internet]. 2024 [cited 2024 Dec 19]. Available from: https://cran.r-project.org/web/packages/psych/index.html

60. Husson F, Josse J, Le S, Mazet J. FactoMineR: Multivariate Exploratory Data Analysis and Data Mining [Internet]. 2024 [cited 2024 Dec 19]. Available from: https://cran.r-project.org/web/packages/FactoMineR/index.html

61. Kassambara A, Mundt F. factoextra: Extract and Visualize the Results of Multivariate Data Analyses [Internet]. 2020 [cited 2024 Dec 19]. Available from: https://cran.r-project.org/web/packages/factoextra/index.html

62. Hope RM. Rmisc: Ryan Miscellaneous [Internet]. 2022 [cited 2024 Dec 19]. Available from: https://cran.r-project.org/web/packages/Rmisc/index.html

63. Wickham H, Chang W, Henry L, Pedersen TL, Takahashi K, Wilke C, et al. ggplot2: Create Elegant Data Visualisations Using the Grammar of Graphics [Internet]. 2024 [cited 2024 Dec 19]. Available from: https://cran.r-project.org/web/packages/ggplot2/index.html

64. Ahlmann-Eltze [aut C, Patil (@patilindrajeets) I. ggsignif: Significance Brackets for "ggplot2" [Internet]. 2022 [cited 2024 Dec 19]. Available from: https://cran.r-project.org/web/packages/ggsignif/index.html

65. Kassambara A. ggpubr: "ggplot2" Based Publication Ready Plots [Internet]. 2023 [cited 2024 Dec 19]. Available from: https://cran.r-project.org/web/packages/ggpubr/index.html

66.   Pedersen TL. patchwork: The Composer of Plots [Internet]. 2024 [cited 2024 Dec 19].
Available from: https://cran.r-project.org/web/packages/patchwork/index.html

67.   Ram [aut K, cre, Wickham H, Richards C, Baggett A. wesanderson: A Wes Anderson
Palette Generator [Internet]. 2023 [cited 2024 Dec 19]. Available from: https://cran.r-
project.org/web/packages/wesanderson/index.html

68.   García-Forero C, Gallardo-Pujol D, Maydeu-Olivares A, Andrés-Pueyo A. Disentangling
impulsiveness, aggressiveness and impulsive aggression: An empirical approach using self-report
measures. Psychiatry Res. 2009 Jun 30;168(1):40–9.

69.   Rudebeck PH, Walton ME, Millette BHP, Shirley E, Rushworth MFS, Bannerman DM.
Distinct contributions of frontal areas to emotion and social behaviour in the rat. Eur J Neurosci.
2007;26(8):2315–26.

70.   Overduin-de Vries AM, Vries H de, Vermande MM, Reijntjes AHA, Sterck EHM. Both
aggressive and affiliative behaviour facilitate resource access in high-ranking female long-tailed
macaques (Macaca fascicularis). Behaviour [Internet]. 2020 Mar 20 [cited 2024 Dec 19];157.
Available from: https://brill.com/view/journals/beh/157/3-4/article-p267_3.xml

71.   Speechley EM, Ashton BJ, Thornton A, King SL, Simmons LW, Woodiss-Field SL, et al.
Aggressive interactions influence cognitive performance in Western Australian magpies. Proc R
Soc B Biol Sci. 2024 Jun 5;291(2024):20240435.

72.   Garnham LC, Porthén SA, Child S, Forslind S, Løvlie H. The role of personality, cognition,
and affective state in same-sex contests in the red junglefowl. Behav Ecol Sociobiol. 2019 Nov
19;73(11):149.

73.   Troisi CA, Vernouillet A, Allaert R, Knoch S, Martel A, Lens L, et al. Beyond a unitary
construct: Dissecting stopping behaviour in two bird species. bioRxiv;

74.   Brucks D, Marshall-Pescini S, Wallis LJ, Huber L, Range F. Measures of Dogs' Inhibitory
Control Abilities Do Not Correlate across Tasks. Front Psychol. 2017 May 24;8:849.

75.   Fagnani J, Barrera G, Carballo F, Bentosela M. Is previous experience important for
inhibitory control? A comparison between shelter and pet dogs in A-not-B and cylinder tasks.
Anim Cogn. 2016 Nov 1;19(6):1165–72.

76.   Van Horik JO, Langley EJG, Whiteside MA, Laker PR, Madden JR. Intra-individual
variation in performance on novel variants of similar tasks influences single factor explanations of
general cognitive processes. R Soc Open Sci. 2018 Jul;5(7):171919.

77.   Zucca P, Antonelli F, Vallortigara G. Detour behaviour in three species of birds: quails
(Coturnix sp.), herring gulls (Larus cachinnans) and canaries (Serinus canaria). Anim Cogn. 2005
Apr;8(2):122–8.

78.   Marshall-Pescini S, Virányi Z, Range F. The Effect of Domestication on Inhibitory Control:
Wolves and Dogs Compared. PLOS ONE. 2015 Feb 25;10(2):e0118469.

79.   Cauchoix M, Chow PKY, van Horik JO, Atance CM, Barbeau EJ, Barragan-Jason G, et al.
The repeatability of cognitive performance: a meta-analysis. Philos Trans R Soc B Biol Sci. 2018
Sep 26;373(1756):20170281.

80.   Sih A, Bell A, Johnson JC. Behavioral syndromes: an ecological and evolutionary overview.
Trends Ecol Evol. 2004 Jul 1;19(7):372–8.

81.   Bell AM. Future directions in behavioural syndromes research. Proc R Soc B Biol Sci. 2006
Dec 21;274(1611):755–61.

82.   Réale D, Reader SM, Sol D, McDougall PT, Dingemanse NJ. Integrating animal
temperament within ecology and evolution. Biol Rev Camb Philos Soc. 2007 May;82(2):291–318.

83.   Briffa M, Sneddon LU, Wilson AJ. Animal personality as a cause and consequence of contest behaviour. Biol Lett. 2015 Mar;11(3):20141007.

84.   Hsu Y, Earley RL, Wolf LL. Modulation of aggressive behaviour by fighting experience: mechanisms and contest outcomes. Biol Rev Camb Philos Soc. 2006 Feb;81(1):33–74.

85.   Tibbetts EA, Pardo-Sanchez J, Weise C. The establishment and maintenance of dominance hierarchies. Philos Trans R Soc B Biol Sci. 2022 Feb 28;377(1845):20200450.

86.   Hobson EA. Differences in social information are critical to understanding aggressive behavior in animal dominance hierarchies. Curr Opin Psychol. 2020 Jun;33:209–15.

87.   Doran C, Bierbach D, Laskowski KL. Familiarity increases aggressiveness among clonal fish. Anim Behav. 2019 Feb;148:153–9.

88.   Dugatkin LA, Earley RL. Individual recognition, dominance hierarchies and winner and loser effects. Proc R Soc Lond B Biol Sci. 2004 Jul 22;271(1547):1537–40.

89.   Werba JA, Stuckert AM, Edwards M, McCoy MW. Stranger danger: A meta-analysis of the dear enemy hypothesis. Behav Processes. 2022 Jan;194:104542.

90.   Marler P. On animal aggression: The roles of strangeness and familiarity. Am Psychol. 1976;31(3):239–46.

91.   Leighton GM, Drury JP, Small J, Miller ET. Unfamiliarity generates costly aggression in interspecific avian dominance hierarchies. Nat Commun. 2024 Jan 6;15(1):335.

92.   Hector AK, Raleigh MJ. The effects of temporary removal of the alpha male on the behavior of subordinate male vervet monkeys. Am J Primatol. 1992;26(2):77–87.

93.   Lambert CT, Guillette LM. The impact of environmental and social factors on learning abilities: a meta-analysis. Biol Rev. 2021 Dec;96(6):2871–89.

94.   Triki Z, Zhou T, Argyriou E, Sousa De Novais E, Servant O, Kolm N. Social complexity affects cognitive abilities but not brain structure in a Poeciliid fish. Le Roux A, editor. Behav Ecol. 2024 May 1;35(3):arae026.

95.   Brucks D, Härterich A, König Von Borstel U. Horses wait for more and better rewards in a delay of gratification paradigm. Front Psychol. 2022 Jul 22;13:954472.

96.   Ruploh T, Henning M, Bischof H, Von Engelhardt N. Effects of social conditions during adolescence on courtship and aggressive behavior are not abolished by adult social experience. Dev Psychobiol. 2015 Jan;57(1):73–82.

97.   Rodenburg TB, Koene P. The impact of group size on damaging behaviours, aggression, fear and stress in farm animals. Appl Anim Behav Sci. 2007 Mar;103(3–4):205–14.

98.   Middlebush CW, Mountjoy PT, Riley V. Unconditioned aggression in the Japanese quail, Coturnix coturnix japonica. Psychon Sci. 1968 Mar;11(3):113–4.

99.   Cooper Jr WE, Dimopoulos I, Pafilis P. Sex, Age, and Population Density Affect Aggressive Behaviors in Island Lizards Promoting Cannibalism. Ethology. 2015;121(3):260–9.

100.   Schweitzer C, Lévy F, Arnould C. Increasing group size decreases social bonding in young Japanese quail, Coturnix japonica. Anim Behav. 2011 Mar;81(3):535–42.

101.   Shaw RC, Schmelz M. Cognitive test batteries in animal cognition research: evaluating the past, present and future of comparative psychometrics. Anim Cogn. 2017 Nov 1;20(6):1003–18.

102.   Boogert NJ, Madden JR, Morand-Ferron J, Thornton A. Measuring and understanding individual differences in cognition. Philos Trans R Soc B Biol Sci. 2018 Sep 26;373(1756):20170280.

103.   Thornton A, Lukas D. Individual variation in cognitive performance: developmental and evolutionary perspectives. Philos Trans R Soc B Biol Sci. 2012 Oct 5;367(1603):2773–83.

104.   Völter CJ, Tinklenberg B, Call J, Seed AM. Comparative psychometrics: establishing what

differs is central to understanding what evolves. Philos Trans R Soc B Biol Sci. 2018 Aug
13;373(1756):20170283.

105.   Brucks D, Marshall-Pescini S, Range F. Dogs and wolves do not differ in their inhibitory
control abilities in a non-social test battery. Anim Cogn. 2019 Jan;22(1):1–15.

106.   Rowe C, Healy SD. Measuring variation in cognition. Behav Ecol. 2014 Jan 1;25(6):1287–
92.

107.   Szabo B, Hoefer S, Whiting MJ. Are lizards capable of inhibitory control? Performance on
a semi-transparent version of the cylinder task in five species of Australian skinks. Behav Ecol
Sociobiol. 2020 Oct;74(10):118.

108.   Mayack C, Naug D. Starving honeybees lose self-control. Biol Lett. 2015
Jan;11(1):20140820.

109.   Shaw RC. Testing cognition in the wild: factors affecting performance and individual
consistency in two measures of avian cognition. Behav Processes. 2017 Jan;134:31–6.

110.   Hanas AK, Guigueno MF, Fernie KJ, Letcher RJ, Ste-Marie Chamberland F, Head JA.
Assessment of the effects of early life exposure to triphenyl phosphate on fear, boldness, aggression,
and activity in Japanese quail (Coturnix japonica) chicks. Environ Pollut. 2020 Mar;258:113695.

111.   Lucon-Xiccato T. The contribution of executive functions to sex differences in animal
cognition. Neurosci Biobehav Rev. 2022 Jul;138:104705.

112.   Vinogradov IM, Jennions MD, Van Veen E, Fichtel C, Kappeler PeterM, Fox RJ. The
effect of sex, age and boldness on inhibitory control. Anim Behav. 2022 Nov;193:133–43.

113.   Lacreuse A, Gullstrand J, Fagot J. Sex differences in inhibitory control in socially-housed
baboons (Papio papio). Behav Brain Res. 2016 Oct;312:231–7.

114.   Junttila S, Huohvanainen S, Tiira K. Effect of Sex and Reproductive Status on Inhibitory
Control and Social Cognition in the Domestic Dog (Canis familiaris). Animals. 2021 Aug
20;11(8):2448.

115.   Lucon-Xiccato T, Bisazza A. Discrimination reversal learning reveals greater female
behavioural flexibility in guppies. Biol Lett. 2014 Jun 30;10(6):20140206.

116.   Garnham LC, Boddington R, Løvlie H. Variation in inhibitory control does not influence
social rank, foraging efficiency, or risk taking, in red junglefowl females. Anim Cogn [Internet].
2022 Feb 4 [cited 2022 May 18]; Available from: https://link.springer.com/10.1007/s10071-022-
01598-5

117.   Schlinger BA, Palter B, Callard GV. A method to quantify aggressiveness in Japanese quail
(Coturnix c. japonica). Physiol Behav. 1987 Jan;40(3):343–8.

118.   Ophir AG, Persaud KN, Galef BG. Avoidance of Relatively Aggressive Male Japanese
Quail (Coturnix japonica) by Sexually Experienced Conspecific Females. J Comp Psychol.
2005;119(1):3–7.

119.   Rutkowska J, Place NJ, Vincent S, Adkins-Regan E. Adrenocortical response to mating,
social interaction and restraint in the female Japanese quail. Physiol Behav. 2011 Oct;104(5):1037–
40.

120.   Abuoghaba AAK, Ali F, Selim DAF, Abdelwahab AAM, Abdelfattah MG. Impact of male-
female cohabitation period on behavioral aspects, fertility, hatchability, and hormonal estimates of
Japanese quail. Poult Sci. 2022 Jan;101(1):101530.

121.   Pellegrini S, Orso GA, Marin RH, Guzman DA. The insemination status and social context
influences quail laying and social behavior: A novel experimental set up. Poult Sci. 2023
Feb;102(2):102328.

122. Georgelin M, Ferreira VHB, Cornilleau F, Meurisse M, Poissenot K, Beltramo M, et al.
Short photoperiod modulates behavior, cognition and hippocampal neurogenesis in male Japanese
quail. Sci Rep. 2023 Jan 18;13(1):951.

123. Greenberg R, Mettke-hofmann C. Ecological Aspects of Neophobia and Neophilia in Birds.
In: Nolan V, Thompson CF, editors. Current Ornithology [Internet]. Boston, MA: Springer US;
2001 [cited 2024 Dec 20]. p. 119–78. Available from: https://doi.org/10.1007/978-1-4615-1211-
0_3

124. Lucon-Xiccato T, Montalbano G, Bertolucci C. Personality traits covary with individual
differences in inhibitory abilities in 2 species of fish. Carere C, editor. Curr Zool. 2020 Apr
1;66(2):187–95.

125. Dougherty LR, Guillette LM. Linking personality and cognition: a meta-analysis. Philos
Trans R Soc B Biol Sci. 2018 Aug 13;373(1756):20170282.

126. Webster MM, Rutz C. How STRANGE are your study animals? Nature. 2020
Jun;582(7812):337–40.

127. Ferreira VHB, Reiter L, Germain K, Calandreau L, Guesdon V. Uninhibited chickens:
ranging behaviour impacts motor self-regulation in free-range broiler chickens (*Gallus gallus*
domesticus). Biol Lett. 2020 Jan;16(1):20190721.

128. Verbeek MEM, Boon A, Drent PJ. Exploration, Aggressive Behaviour and Dominance in
Pair-Wise Confrontations of Juvenile Male Great Tits. Behaviour. 1996;133(11–12):945–63.

129. Madden JR, Langley EJG, Whiteside MA, Beardsworth CE, van Horik JO. The quick are
the dead: pheasants that are slow to reverse a learned association survive for longer in the wild.
Philos Trans R Soc Lond B Biol Sci. 2018 Sep 26;373(1756):20170297.

130. Suter H, Houston AI, Radford AN. Self-control: waiting longer is not necessarily better.
ISBE 2024; 2024 Oct 3; Melbourne, Australia.

131. Stevens JR, Stephens DW. The adaptive nature of impulsivity. In: Madden GJ, Bickel WK,
editors. Impulsivity: The behavioral and neurological science of discounting [Internet].
Washington: American Psychological Association; 2010 [cited 2022 May 18]. p. 361–87. Available
from: http://content.apa.org/books/12069-013

132. Kelly AM, Wilson LC. Aggression: Perspectives from social and systems neuroscience.
Horm Behav. 2020 Jul;123:104523.

133. Flanigan ME, Russo SJ. Recent advances in the study of aggression.
Neuropsychopharmacology. 2019 Jan;44(2):241–4.

1149